

RESIDUAL INTERSECTIONS AND THE ANNIHILATOR OF KOSZUL HOMOLOGIES

S. H. HASSANZADEH¹ AND J. NAÉLITON²

ABSTRACT. Cohen-Macaulayness, unmixedness, the structure of the canonical module and the stability of the Hilbert function of algebraic residual intersections are studied in this paper. Some conjectures about these properties are established for large classes of residual intersections without restricting local number of generators of the ideals involved. A family of approximation complexes for residual intersections is constructed to determine the above properties. Moreover some general properties of the symmetric powers of quotient ideals are determined which were not known even for special ideals with a small number of generators. Acyclicity of a prime case of these complexes is shown to be equivalent to find a common annihilator for higher Koszul homologies. So that, a tight relation between residual intersections and the uniform annihilator of positive Koszul homologies is unveiled that sheds some light on their structure.

1. INTRODUCTION

Understanding the residual of the intersection of two algebraic varieties is an intriguing concept in algebraic geometry. Similar to many other notions in intersection theory, the concept of residual intersections need the proper interpretation to encompass the desired algebraic and geometric properties. In the sense of Artin and Nagata [AN], which is our point of view, the notion is a vast generalization of linkage (or liaison) which is more ubiquitous, but also harder to understand. Let X and Y be irreducible closed subschemes of a Noetherian scheme Z with $\text{codim}_Z(X) \leq \text{codim}_Z(Y) = s$ and $Y \not\subset X$, Then Y is called a residual intersection of X if the number of equations needed to define $X \cup Y$ as a subscheme of Z is the smallest possible that is s . Precisely, if R is a Noetherian ring, I an ideal of height g and $s \geq g$ an integer, then

- An *(algebraic) s-residual intersection* of I is a proper ideal J of R such that $\text{ht}(J) \geq s$ and $J = (\mathfrak{a} :_R I)$ for some ideal $\mathfrak{a} \subset I$ generated by s elements.
- A *geometric s-residual intersection* of I is an algebraic s -residual intersection J of I such that $\text{ht}(I + J) \geq s + 1$.

Based on a construction of Laksov for residual intersection, Fulton [Fu, Definition 9.2.2] presents a formulation for residual intersection that, locally, can be expressed as follows: Suppose that $X = \text{Spec}(R)$

Date: October 16, 2018.

⁰Mathematics Subject Classification 2010 (MSC2010). Primary 13C40, 13H10, 13D02, Secondary 14C17, 14M06.

⁰Keywords: Residual Intersection, Canonical Module, Type, Sliding depth, Approximation Complex, Koszul annihilator

¹Corresponding author, Partially supported by a CNPq grant.

²Under a CNPq Doctoral scholarship.

and Y and S are closed subschemes of X defined by the ideals \mathfrak{a} and I respectively. Let $\tilde{X} = \text{Proj}(\mathcal{R}_R(I))$ be the blow-up of X along S . Consider the natural map $\pi : \tilde{X} \rightarrow X$. Let $\tilde{Y} = \pi^{-1}(Y)$ and $\tilde{S} = \pi^{-1}(S)$. Then $\tilde{Y} = \text{Proj}(\mathcal{R}_R(I)/\mathfrak{a}\mathcal{R}_R(I))$ and $\tilde{S} = \text{Proj}(\mathcal{R}_R(I)/I\mathcal{R}_R(I))$ are closed subschemes of \tilde{X} with ideal sheaves $\mathcal{I}_{\tilde{Y}}$ and $\mathcal{I}_{\tilde{S}}$. $\mathcal{I}_{\tilde{S}}$ is an invertible sheaf. Let $Z' \subseteq \tilde{X}$ be the closed subscheme defined by the ideal sheaf $\mathcal{I}_{Z'} = \mathcal{I}_{\tilde{Y}} \cdot \mathcal{I}_{\tilde{S}}^{-1}$. Z' is called the *residual scheme to \tilde{S} in \tilde{Y}* . Precisely $Z' = \text{Proj}(\mathcal{R}_R(I)/\gamma\mathcal{R}_R(I))$ in which $\gamma = \mathfrak{a} \subseteq (\mathcal{R}_R(I))_{[1]}$. Finally, the residual intersection to S in Y [Fu, 9.2.2] is the direct image of $\mathcal{O}_{Z'}$ i.e. $\pi_*(\mathcal{O}_{Z'})$. Since $\mathcal{O}_{Z'}$ is a coherent sheaf, by [H, III, Proposition 8.5],

$$\pi_*(\mathcal{O}_{Z'}) = H^0(\widetilde{\tilde{X}}, \widetilde{\mathcal{O}_{Z'}}) = \Gamma(\widetilde{\tilde{X}}, \widetilde{\mathcal{O}_{Z'}}).$$

Note that $\Gamma(\widetilde{\tilde{X}}, \widetilde{\mathcal{O}_{Z'}}) = \Gamma_*(\tilde{X}, \mathcal{O}_{Z'}(0))$ which is equal to the ideal transform $D_{\mathcal{R}_R(I)_+}(\mathcal{R}_R(I)/\gamma\mathcal{R}_R(I))_{[0]}$, by an application of Čech complex. The later is closely related, and in many cases determined, by the ideal

$$H_{\mathcal{R}_R(I)_+}^0(\mathcal{R}_R(I)/\gamma\mathcal{R}_R(I))_{[0]} = \bigcup (\mathfrak{a}I^i :_R I^{i+1}).$$

In [Ha] and in the current work, we consider the symmetric algebra $\text{Sym}_R(I)$ instead of the Rees algebra $\mathcal{R}_R(I)$. This generalization has already proved its usefulness in studying multiple-point formulas by Kleiman c.f. [Fu, 17.6.2]. We then find a kind of *Arithmetical* residual intersection to be $\bigcup (\gamma\text{Sym}_R^i(I) :_R \text{Sym}_R^{i+1}(I))$ where $\gamma = \mathfrak{a} \subseteq \text{Sym}_R^i(I)_{[1]}$. Comparing the above three definitions for residual intersection, we have

$$J = (\mathfrak{a} : I) \subseteq \bigcup (\gamma\text{Sym}_R^i(I) :_R \text{Sym}_R^{i+1}(I)) \subseteq \bigcup (\mathfrak{a}I^i :_R I^{i+1}).$$

Interestingly, these ideals coincide if the algebraic residual intersection J does not share any associated primes with I , e.g if J is unmixed and the residual is geometric.

Determining the cases where the first inclusion above is an equality leads us to define a third variation of algebraic residual intersection.

Definition 1.1. An *arithmetic s -residual intersection* $J = (\mathfrak{a} : I)$ is an algebraic s -residual intersection such that $\mu_{R_{\mathfrak{p}}}((I/\mathfrak{a})_{\mathfrak{p}}) \leq 1$ for all prime ideal $\mathfrak{p} \supseteq (I + J)$ with $\text{ht}(\mathfrak{p}) \leq s$. (μ denotes the minimum number of generators.)

Clearly any geometric s -residual intersection is arithmetic. Moreover for any algebraic s -residual intersection $J = (\mathfrak{a} : (f_1, \dots, f_r))$ which is not geometric, all of the colon ideals $(\mathfrak{a} : f_i)$ are arithmetic s -residual intersection and at least one of them is not geometric.

In this paper we introduce a family of complexes, denoted by $\{\mathcal{Z}_{\bullet}^+\}_{i=0}^{\infty}$, to approximate i th symmetric power of I/\mathfrak{a} , $\text{Sym}_R^i(I/\mathfrak{a})$ for $i > 1$. The idea to define this family is inspired by [Ha] in which the single complex ${}_0\mathcal{Z}_{\bullet}^+$ is treated. $H_0({}_0\mathcal{Z}_{\bullet}^+)$ is a cyclic module of the form R/K where K is called the *Disguised s -residual intersection of I w.r.t. \mathfrak{a}* , see Definition 2.1. The study of the other members of the above family of complexes shed some more light on the structure of residual intersections. A flavor of our main results in Section 2 is the following,

Let (R, \mathfrak{m}) be a CM local ring of dimension d and I be an ideal with $\text{ht}(I) = g > 0$. Let $s \geq g$, $1 \leq k \leq s - g + 2$ and $J = (\mathfrak{a} : I)$ be any (algebraic) s -residual intersection. Suppose that I is Strongly Cohen-Macaulay. Then

- (1) Corollary 2.14, the canonical module of R/J is $\text{Sym}_R^{s-g+1}(I/\mathfrak{a})$, provided the residual is arithmetic and R is Gorenstein;
- (2) Corollary 2.8, $\text{depth}(R/\mathfrak{a}) = d - s$;
- (3) Corollary 2.8, J is unmixed of codimension s ;
- (4) Theorem 2.6 and Proposition 3.1, ${}_k\mathcal{Z}_\bullet^+$ is acyclic, $H_0({}_k\mathcal{Z}_\bullet^+) = \text{Sym}_R^k(I/\mathfrak{a})$ and the latter is CM of dimension $d - s$; (the acyclicity of ${}_k\mathcal{Z}_\bullet^+$ implies conjecture(5) below in the arithmetic case.)

These results address the following (implicit) conjectures made during the development of the theory of algebraic residual intersections.

Let R be a Cohen-Macaulay(CM) local ring and I be Strongly Cohen-Macaulay (SCM) or even just satisfies Sliding depth (SD). Then

- (1) R/J is Cohen-Macaulay.
- (2) The canonical module of R/J is the $(s - g + 1)$ th symmetric power of I/\mathfrak{a} , if R is Gorenstein.
- (3) \mathfrak{a} is minimally generated by s elements.
- (4) J is unmixed.
- (5) The Hilbert series of R/J depends only on I and the degrees of the generators of \mathfrak{a} .

The first conjecture essentially goes back to [AN], it has been asked as an open question in [HU]. However in the Sundance conference in 1990 [U1], Ulrich mentioned the above conjectures (1) – (4) as desirable facts to be proved. The property of the Hilbert function is rather recent and has been analyzed by Chardin, Eisenbud and Ulrich in [CEU].

It should be mentioned that these conjectures are proved if one supposes in addition that the ideal I has locally few number of generators, a condition which is called G_s , or if it has a deformation with G_s property. Over time the G_s condition became a 'standard' assumption in the theory of residual intersection which is not avoidable in some cases. However, the desire is to prove the above assertions without restricting the local number of generators of I .

In comparison, obtaining the structure of the canonical module in the absence of the G_s condition is more challenging. To achieve this, we show that under the above mentioned hypotheses, $\text{Sym}_R^{s-g+1}(I/\mathfrak{a})$ is a faithful maximal Cohen-Macaulay R/J -module of type 1. Concerning the type of modules, we prove even more. We show in Theorem 2.12 that the following inequality holds for any $1 \leq k \leq s - g + 1$,

$$r_R(\text{Sym}_R^k(I/\mathfrak{a})) \leq \binom{r+s-g-k}{r-1} r_R(R),$$

where for a finitely generated R -module M , $r_R(M) := \dim_{R/\mathfrak{m}} \text{Ext}_R^{\text{depth}(M)}(R/\mathfrak{m}, M)$ is the Cohen-Macaulay type of M .

In section 3, we present several applications of the theorems and constructions so far. We state how much the Hilbert functions of R/J and R/\mathfrak{a} depend on the generators and/or degrees of I and \mathfrak{a} . If I

satisfies SCM, then the Hilbert function of the disguised residual intersection, and that of $Sym_R^k(I/\mathfrak{a})$, if $1 \leq k \leq s - g + 2$, depends only on the **degrees** of the generators of \mathfrak{a} and the Koszul homologies of I . In particular, $k = 1$ implies that the Hilbert function of R/\mathfrak{a} is constant on the open set of ideals \mathfrak{a} generated by s forms of the given degrees such that $\text{ht}(\mathfrak{a} : I) \geq s$. This is comparable with results in [CEU] where the same assertion is concluded under some G_s hypotheses.

The graded structure of ${}_k\mathcal{Z}_\bullet^+$ shows that if I satisfies the SD_1 condition, then

$$\text{reg}(Sym_R^k(I/\mathfrak{a})) \leq \text{reg}(R) + \dim(R_0) + \sigma(\mathfrak{a}) - (s - g + 1 - k) \text{indeg}(I/\mathfrak{a}) - s.$$

for $k \geq 1$. These applications were not known even if ideal I satisfies the G_s condition. Finally in this section, Proposition 3.4, by a combination of older and newer facts, we show that for any algebraic s -residual intersection $J = \mathfrak{a} : I$, if I is SCM and evenly linked to a G_s ideal (or has a deformation with these properties) then the disguised residual intersection and the algebraic residual intersection coincide. Based on this fact, we conjecture that, Conjecture 5.9, *in the presence of the sliding depth condition the disguised residual intersection is the same as the algebraic residual intersection.*

In section 4, we try to understand better the structure of complex ${}_0\mathcal{Z}_\bullet^+$ in the case where I/\mathfrak{a} is principal, say $I = (\mathfrak{a}, b)$. We find in Theorem 4.4 that $H_i({}_0\mathcal{Z}_\bullet^+(\mathfrak{a}, \mathbf{f})) \simeq bH_i(a_1, \dots, a_s)$ for all $i \geq 1$ and $H_0({}_0\mathcal{Z}_\bullet^+(\mathfrak{a}, \mathbf{f})) \simeq R/(\mathfrak{a} : b)$. This fact shows how much the homologies of \mathcal{Z}_\bullet^+ complexes may depend on the generating sets, as well it shows a tight relation between uniform annihilator of Koszul homologies and acyclicity of \mathcal{Z}_\bullet^+ . As a byproduct, Corollary 4.6 suggests that in the way of studying the properties of colon ideals, instead of assuming $\text{ht}((a_1, \dots, a_s) : I) \geq s$, one may only need to suppose that $IH_i(a_1, \dots, a_s) = 0$ locally at codimension $s - 1$.

Motivated by the facts in section 4, we investigate on the uniform annihilator of positive Koszul homologies in Section 5. Not much is known about the annihilator of Koszul homologies. In Corollary 5.6, we show that for a residual intersection $J = \mathfrak{a} : I$ where I satisfies SD, and $\text{depth}(R/I) \geq d - s$,

$$I \subseteq \bigcap_{j \geq 1} \text{Ann}(H_j(\mathfrak{a})).$$

Surprisingly, this result contradicts one of the unpublished, but well-known results of G. Levin [V2, Theorem 5.26] which yielded in [CHKV] that $\text{Supp}(H_1(\mathfrak{a})) = \text{Supp}(H_0(\mathfrak{a}))$. Simple examples of residual intersection disprove this last claim. Moreover in this section it is shown in Theorem 5.4 that for an s -residual intersection $J = \mathfrak{a} : I$, if I satisfies SD and $\text{depth}(R/I) \geq d - s$, so does \mathfrak{a} . This is an interesting result since for a long time it was known that the residual intersections of the ideal \mathfrak{a} are Cohen-Macaulay although no one was aware of the SD property of \mathfrak{a} .

2. RESIDUAL APPROXIMATION COMPLEXES

In this section we introduce a family of complexes which approximate the residual intersection and some of its related symmetric powers. We denote this family by $\{\mathcal{Z}_\bullet^+\}_{i=0}^\infty$. The complex ${}_0\mathcal{Z}_\bullet^+$ was already defined in [Ha] and used to prove the CM-ness of arithmetic residual intersection of ideals with sliding depth.

Throughout this section, R is a Noetherian ring of dimension d , $I = (\mathbf{f}) = (f_1, \dots, f_r)$ is an ideal of grade $g \geq 1$. Although by adding one variable we may as well treat the case $g = 0$; but, for simplicity, we keep the assumption $g \geq 1$. $\mathfrak{a} = (a_1, \dots, a_s)$ is an ideal contained in I , $s \geq g$, $J = \mathfrak{a} :_R I$, and $S = R[T_1, \dots, T_r]$ is a polynomial extension of R with indeterminates T_i 's. We denote the symmetric algebra of I over R by \mathcal{S}_I or in general the symmetric algebra of an R -module M by $\text{Sym}_R(M)$ and the k th symmetric power of M by $\text{Sym}_R^k(M)$. We consider \mathcal{S}_I as an S -algebra via the ring homomorphism $S \rightarrow \mathcal{S}_I$ sending T_i to f_i as an element of $(\mathcal{S}_I)_1 = I$ then $\mathcal{S}_I = S/\mathcal{L}$. Let $a_i = \sum_{j=1}^r c_{ji} f_j$, $\gamma_i = \sum_{j=1}^r c_{ji} T_j$, $\gamma = (\gamma_1, \dots, \gamma_s)$ and $\mathbf{g} := (T_1, \dots, T_r)$.

For a sequence of elements \mathfrak{r} in a commutative ring A and an A -module M , we denote the Koszul complex by $K_\bullet(\mathfrak{r}; M)$, its cycles by $Z_i(\mathfrak{r}; M)$ and homologies by $H_i(\mathfrak{r}; M)$. For a graded module M , $\text{indeg}(M) := \inf\{i : M_i \neq 0\}$ and $\text{end}(M) := \sup\{i : M_i \neq 0\}$. Setting $\text{deg}(T_i) = 1$ for all i , S is a standard graded ring over $S_0 = R$.

To set one more convention, when we draw the picture of a double complex obtained from a tensor product of two finite complexes (in the sense of [W, 2.7.1]), say $\mathcal{A} \otimes \mathcal{B}$; we always put \mathcal{A} in the vertical direction and \mathcal{B} in the horizontal one. We also label the module which is in the up-right corner by $(0, 0)$ and consider the labels for the rest, as the points in the third-quadrant.

2.1. ${}_k\mathcal{Z}_\bullet^+$ complexes. The first object in the construction of the family of ${}_k\mathcal{Z}_\bullet^+$ complexes is one of the approximation complexes- the \mathcal{Z} -complex [HSV]. We consider the approximation complex $\mathcal{Z}_\bullet(\mathbf{f})$:

$$0 \rightarrow Z_{r-1} \otimes_R S(1-r) \rightarrow \dots \rightarrow Z_1 \otimes_R S(-1) \rightarrow Z_0 \otimes_R S \rightarrow 0$$

where $Z_i = Z_i(\mathbf{f})$ is the i th cycle of the Koszul complex $K_\bullet(\mathbf{f}, R)$.

The second object is the Koszul complex $K_\bullet(\gamma, S)$:

$$0 \rightarrow K_s(\gamma_1, \dots, \gamma_s)(-s) \rightarrow \dots \rightarrow K_1(\gamma_1, \dots, \gamma_s)(-1) \rightarrow K_0(\gamma_1, \dots, \gamma_s) \rightarrow 0.$$

Let $\mathcal{D}_\bullet = \text{Tot}(K_\bullet(\gamma, S) \otimes_S \mathcal{Z}_\bullet(\mathbf{f}))$. Then

$$\mathcal{D}_i = \bigoplus_{j=i-s}^{\min\{i, r-1\}} [Z_j \otimes_R S]^{(i-s)}(-i). \quad (2.1)$$

For a graded S -module M the k th graded component of M is denoted by $M_{[k]}$. Let $(\mathcal{D}_\bullet)_{[k]}$ for $k \in \mathbb{Z}$ be the k th graded strand of \mathcal{D}_\bullet . We have $(\mathcal{D}_i)_{[k]} = 0$, for all $k < i$, in particular

$$H_k((\mathcal{D}_\bullet)_{[k]}) = \text{Ker}(D_k \rightarrow D_{k-1})_{[k]}. \quad (2.2)$$

Now, let $C_\mathbf{g}^\bullet = C_\mathbf{g}^\bullet(S)$ be the Čech complex of S with respect to the sequence $\mathbf{g} = (T_1, \dots, T_r)$,

$$C_\mathbf{g}^\bullet : 0 \rightarrow C_\mathbf{g}^0(= S) \rightarrow C_\mathbf{g}^1 \rightarrow \dots \rightarrow C_\mathbf{g}^r \rightarrow 0.$$

We then consider the bi-complex $C_\mathbf{g}^\bullet \otimes_S \mathcal{D}_\bullet$ with $C_\mathbf{g}^0 \otimes_S \mathcal{D}_0$ in the corner. This bi-complex gives rise to two spectral sequences for which the second terms of the horizontal spectral are

$${}^2 E_{hor}^{-i,-j} = H_{\mathfrak{g}}^j(H_i(D_{\bullet})), \quad (2.3)$$

and the first terms of the vertical spectral are

$$E_{ver}^{-i,-j} = \begin{cases} 0 \rightarrow H_{\mathfrak{g}}^r(D_{r+s-1}) \rightarrow \cdots \rightarrow H_{\mathfrak{g}}^r(D_1) \rightarrow H_{\mathfrak{g}}^r(D_0) \rightarrow 0 & \text{if } j = r \\ 0 & \text{otherwise} \end{cases} \quad (2.4)$$

Since $H_{\mathfrak{g}}^r(D_i) = H_{\mathfrak{g}}^r(\bigoplus_j [Z_j \otimes S](-i)) = \bigoplus_j Z_j \otimes H_{\mathfrak{g}}^r(S)(-i)$ and $\text{end}(H_{\mathfrak{g}}^r(S)) = -r$, it follows that $\text{end}(H_{\mathfrak{g}}^r(D_i)) = i - r$, thus $H_{\mathfrak{g}}^r(D_i)_{[i-r+j]} = 0$, for all $j \geq 1$. It then leads to define the following sequence of complexes indexed by $k \geq 0$:

$$0 \longrightarrow H_{\mathfrak{g}}^r(D_{r+s-1})_{[k]} \longrightarrow \cdots \longrightarrow H_{\mathfrak{g}}^r(D_{r+k+1})_{[k]} \xrightarrow{\phi_k} H_{\mathfrak{g}}^r(D_{r+k})_{[k]} \longrightarrow 0 \quad (2.5)$$

Since the vertical spectral (2.4) collapses at the second step, the horizontal spectral converges to the homologies of $H_{\mathfrak{g}}^r(D_{\bullet})$. Since all of the homomorphisms are homogeneous of degree 0, the convergence inherits to any graded component. Therefore for any $k \geq 0$ there exists a filtration $\cdots \subseteq \mathcal{F}_{2k} \subseteq \mathcal{F}_{1k} \subseteq \text{Coker}(\phi_k)$ such that

$$\frac{\text{Coker}(\phi_k)}{\mathcal{F}_{1k}} \simeq ({}^{\infty} E_{hor}^{-k,0})_{[k]}. \quad (2.6)$$

Observing that $H_{\mathfrak{g}}^{-t}(H_h(D_{\bullet})) = 0$, for all $(t, h) \in \mathbb{N} \times \mathbb{N}_0$, one has ${}^l E_{hor}^{-k,0} \subseteq H_{\mathfrak{g}}^0(H_k(D_{\bullet}))$, for all $l \geq 2$. Hence we have the following chain of maps for which, except the isomorphism in the middle, all maps are canonical and the map on the right is given by (2.2):

$$\begin{array}{ccccccc} H_{\mathfrak{g}}^r(D_{r+k})_{[k]} & \longrightarrow & \text{Coker}(\phi_k) & \longrightarrow & \frac{\text{Coker}(\phi_k)}{\mathcal{F}_{1k}} & & \\ & & \simeq & & & & \\ {}^{\infty} (E_{hor}^{-k,0})_{[k]} & \xleftarrow{1-1} & H_{\mathfrak{g}}^0(H_k(D))_{[k]} & \xrightarrow{1-1} & H_k(D)_{[k]} & \xrightarrow{1-1} & (D_k)_{[k]}. \end{array} \quad (2.7)$$

We denote the composition of the above chain of R -homomorphisms by τ_k .

Finally, we define the promised family of complexes as follows: For any integer $k \geq 0$, ${}_k \mathcal{Z}_{\bullet}^+$ is a complex of length s consists of two parts: the right part is $(D_{\bullet})_{[k]}$ and the left part is $({}^1 E_{ver})_{[k]}$. These parts are joined via τ_k . More precisely,

$${}_k \mathcal{Z}_{\bullet}^+ : \quad 0 \longrightarrow {}_k \mathcal{Z}_s^+ \longrightarrow \cdots \xrightarrow{\phi_k} {}_k \mathcal{Z}_{k+1}^+ \xrightarrow{\tau_k} {}_k \mathcal{Z}_k^+ \longrightarrow \cdots \longrightarrow {}_k \mathcal{Z}_0^+ \longrightarrow 0 \quad (2.8)$$

wherein

$${}_k \mathcal{Z}_i^+ = \begin{cases} (D_i)_{[k]} & i \leq \min\{k, s\}, \\ H_{\mathfrak{g}}^r(D_{i+r-1})_{[k]} & i > k. \end{cases} \quad (2.9)$$

The structure of ${}_k\mathcal{Z}_\bullet^+$ depends in two ways on the generating sets. Namely, it depends on the generating set of I , \mathbf{f} , and the expression of the generators of \mathfrak{a} in terms of the generators of I , c_{ij} . However, for $k \geq 1$, we have

$$H_0({}_k\mathcal{Z}_\bullet^+) = H_0(D_\bullet)_{[k]} = (\mathcal{S}_I/(\gamma)\mathcal{S}_I)_{[k]} = \text{Sym}_R^k(I/\mathfrak{a}).$$

The case where $k = 0$ is also very interesting. However the structure of $H_0({}_0\mathcal{Z}_\bullet^+)$ is not as clear as the cases where $k > 0$.

Definition 2.1. Let R be a Noetherian ring and $\mathfrak{a} \subseteq I$ be two ideals of R . The *Disguised s -residual intersection of I w.r.t. \mathfrak{a}* is the unique ideal K such that $H_0({}_0\mathcal{Z}_\bullet^+) = R/K$. The reasons for choosing the attribute *Disguised* is that, K is contained in $J = (\mathfrak{a} : I)$, it has the same radical as J has; and in the case where R is CM, J is an algebraic residual intersection and I satisfies some sliding depth condition, K is Cohen-Macaulay; moreover K coincides with J in the case where the residual is arithmetic, by [Ha, Theorem 2.11]. We conjecture, 5.9, that under the above assumptions, K is always the same as J despite it does not appear so.

2.2. Acyclicity and Cohen-Macaulayness. One more occasion where the properties of ${}_k\mathcal{Z}_\bullet^+$ are independent of the generating sets of I and \mathfrak{a} is the following lemma, which is crucial to prove the acyclicity of ${}_k\mathcal{Z}_\bullet^+$.

Lemma 2.2. *Let R be a Noetherian ring and $\mathfrak{a} \subseteq I$ be two ideals of R . If $I = \mathfrak{a}$, then every complex ${}_k\mathcal{Z}_\bullet^+$ defined in (2.8) is exact.*

Proof. Since the approximation complex $\mathcal{Z}_\bullet(\mathbf{f})$ (resp. $K_\bullet(\gamma)$) is a differential graded algebra, $H_i(\mathcal{Z}_\bullet)$ (resp. $H_i(K_\bullet(\gamma))$) is a $\mathcal{S}_I = S/\mathcal{L}$ -module (resp. $S/(\gamma)$ -module) for all i . The bi-complex $K_\bullet(\gamma) \otimes_S \mathcal{Z}_\bullet$, gives rise to the horizontal spectral sequence with second terms ${}^2E_{hor}^{-i,-j} = H_j(K_\bullet(\gamma); H_i(\mathcal{Z}_\bullet))$. It follows that $(\mathcal{L} + (\gamma))$ annihilates ${}^2E_{hor}^{-i,-j}$ and consequently annihilates ${}^\infty E_{hor}^{-i,-j}$ which is a sub-quotient of ${}^2E_{hor}^{-i,-j}$. By the convergence of the spectral sequence to the homologies of the total complex $H_\bullet(D_\bullet)$, it is straightforward to deduce that $H_i(D_\bullet)$ is $(\mathcal{L} + (\gamma))$ -torsion modules for all i , i.e. $(\mathcal{L} + (\gamma))^N H_{i+j}(D_\bullet) = 0$ for all i, j and for some N . Considering the equation $I = \mathfrak{a}$ in $(\mathcal{S}_I)_1$, we have $\mathcal{L} + (\gamma) = \mathcal{L} + \mathfrak{g}$. Therefore

$$H_{\mathfrak{g}}^j(H_i(D_\bullet)) = H_{\mathfrak{g}+\mathcal{L}}^j(H_i(D_\bullet)) = H_{(\gamma)+\mathcal{L}}^j(H_i(D_\bullet)) = \begin{cases} H_i(D_\bullet), & j = 0 \\ 0 & j > 0. \end{cases} \quad (2.10)$$

Once more we study the spectral sequences arising from $C_{\mathfrak{g}}^\bullet \otimes D_\bullet$ as it was already mentioned in (2.8). Applying (2.10), we draw both horizontal and vertical spectral sequence simultaneously as follows:

$$\begin{array}{ccccccc}
{}^\infty E_{hor} : & 0 & \cdots & 0 & H_k(D_\bullet)_{[k]} & \cdots & H_0(D_\bullet)_{[k]} \\
{}^1 E_{ver} : & 0 & \cdots & & & & 0 \\
H_{\mathfrak{g}}^r(D_{r+s-1})_{[k]} & \longrightarrow & \cdots & \xrightarrow{\phi_k} & H_{\mathfrak{g}}^r(D_{r+k})_{[k]} & \longrightarrow & 0 & 0 & \cdots & 0
\end{array}$$

Since $H_j(D_\bullet)$ is a sub-quotient of D_j , $H_j(D_\bullet)_{[k]} = 0$, for all $j > k$. Also by (2.10) there is only one non-zero row in the horizontal spectral. On the other hand \mathfrak{g} is a regular sequence on the D_j 's which in turn shows that the vertical spectral is just one line.

Consequently $H_j({}_k\mathcal{Z}_\bullet^+) := H_{j+r-1}(H_{\mathfrak{g}}^r(D_\bullet))_{[k]} = H_{j-1}(D_\bullet)_{[k]} = 0$ whenever $j > k + 1$. On the other hand, if $j < k$, $H_j({}_k\mathcal{Z}_\bullet^+) := H_j(D_\bullet)_{[k]} = H_{j+r}(H_{\mathfrak{g}}^r(D_\bullet)_{[k]})$. However the latter is zero since $\text{end}(H_{\mathfrak{g}}^r(D_{r+j}) \leq j$. This shows that ${}_k\mathcal{Z}_\bullet^+$ is exact on the left and also on the right hand of τ_k .

It remains to prove the exactness in the joint points k and $k + 1$. For this, notice that, $({}^\infty E_{hor}^{-i,-j})_{[k]} = 0$ for $j \geq 1$ hence $\mathcal{F}_{jk}/\mathcal{F}_{(j+1)k} = ({}^\infty E_{hor}^{-k-j,-j})_{[k]} = 0$ for $j \geq 1$. Consequently, $\mathcal{F}_{1k} = 0$ in (2.6). Therefore the map τ_k defined in (2.7) is exactly the canonical map $H_{\mathfrak{g}}^r(D_{r+k})_{[k]} \xrightarrow{Can.} \text{Coker}(\phi_k)_{[k]}$ as required. \square

As already mentioned in the introduction, some sliding depth conditions are needed to prove the acyclicity of the ${}_k\mathcal{Z}_\bullet^+$ complexes.

Definition 2.3. Let (R, \mathfrak{m}) be a Noetherian local ring of dimension d and $I = (f_1, \dots, f_r)$ an ideal of grade $g \geq 1$. Let k and t be two integers, we say that the ideal I satisfies SD_k at level t if $\text{depth}(H_i(\mathbf{f}; R)) \geq \min\{d - g, d - r + i + k\}$ for all $i \geq r - g - t$ (whenever $t = r - g$ we simply say that I satisfies SD_k , also SD stands for SD_0).

I is strongly Cohen-Macaulay, SCM, if $H_i(\mathbf{f}; R)$ is CM for all i . Clearly SCM is equivalent to SD_{r-g} .

Similarly, we say that I satisfies the sliding depth condition on cycles, SDC_k , at level t , if $\text{depth}(Z_i(\mathbf{f}, R)) \geq \min\{d - r + i + k, d - g + 2, d\}$ for all $i \geq r - g - t$. Again if $t = r - g$, we simply say that I satisfies SDC_k and we use SDC instead of SDC_0 .

Some of the basic properties and relations between conditions SD_k and SDC_k are explained in the following proposition.

Proposition 2.4. *Let (R, \mathfrak{m}) be a CM local ring of dimension d and $I = (f_1, \dots, f_r)$ be an ideal of grade $g \geq 1$. Let k and t be two integers. Then*

- (1) *The properties SDC_k and SD_k at level t localizes; they only depend on I and not the generating set, if $t = r - g$.*
- (2) *SD_k implies SDC_{k+1} .*

- (3) SD_k at level $t < r - g$ implies SDC_a at level t for $a \leq g + t + 2 - d$.
- (4) SDC_{k+1} at level t implies SD_k at level t for any t , if $g \geq 2$. This implication is also the case if $g = 1$ and $k = 0$.
- (5) SD_0 at level $t \geq 1$ implies SDC_0 at level t .

Proof. (1) is essentially proved in [V]. (2) was already proved in [Ha, 2.5]. (3) follows after analyzing the spectral sequences derived from the tensor product of the truncated Koszul complex

$$0 \longrightarrow Z_i(\mathbf{f}) \longrightarrow K_i \longrightarrow K_{i-1} \longrightarrow \cdots \longrightarrow K_0 \longrightarrow 0.$$

and the Čech complex, $C_m^\bullet(R)$. To prove (4), we consider the depth inequalities derived from the short exact sequences $0 \longrightarrow Z_{i+1}(\mathbf{f}) \longrightarrow K_{i+1} \longrightarrow B_i(\mathbf{f}) \longrightarrow 0$, and $0 \longrightarrow B_i(\mathbf{f}) \longrightarrow Z_i(\mathbf{f}) \longrightarrow H_i(\mathbf{f}) \longrightarrow 0$. To show (5) we use the fact that SDC_k holds for any k at level -1 then a recursive induction applying $\text{depth}(Z_i) \geq \min\{\text{depth}(H_i), d, \text{depth}(Z_{i+1}) - 1\}$ proves the assertion. \square

In the next Theorem we present sufficient conditions for the acyclicity of ${}_k\mathcal{Z}_\bullet^+$. The strategy taken here to prove the acyclicity is the same as the one applied in [Ha] (**however the reader should notice that the complex \mathcal{C}_\bullet defined in [Ha] is a little bit different from ${}_0\mathcal{Z}_\bullet^+$ here, in the former the tail is substituted by a free complex but still \mathcal{C}_\bullet remains quasi-isomorphic to ${}_0\mathcal{Z}_\bullet^+$**). Since the complex is finite we avail ourselves of "lemme d'aciclicité" of Peskine-Szpiro; so that we assume some sliding depth conditions and prove the acyclicity in height $s - 1$ wherein $I = \mathfrak{a}$. By Lemma 2.2, ${}_k\mathcal{Z}_\bullet^+$ is exact, if $I = \mathfrak{a}$. Then an induction will show the acyclicity globally.

Although the proof here is more involved than [Ha, 2.8], we prefer to omit it to go faster to newer theorems. The Cohen-Macaulay hypothesis in this theorem is needed to show that if for an R -module M , $\text{depth}(M) \geq d - t$ then for any prime \mathfrak{p} , $\text{depth}(M_{\mathfrak{p}}) \geq \text{ht}(\mathfrak{p}) - t$, [V, Section 3.3].

Proposition 2.5. *Let (R, \mathfrak{m}) be a CM local ring of dimension d and $J = (\mathfrak{a} : I)$ be an s -residual intersection. Assume that $I = (f_1, \dots, f_r)$ and $\text{ht}(I) = g \geq 1$. Fix $0 \leq k \leq \max\{s, s - g + 2\}$. Then the complex ${}_k\mathcal{Z}_\bullet^+$ is acyclic, if any of the following hypotheses holds*

- (1) $1 \leq s \leq 2$ and $s = k$, or
- (2) $r + k \geq s + 1$, $k \leq 2$ and I satisfies SDC_1 at level $s - g - k$, or
- (3) $r + k \leq s$ and I satisfies SD , or
- (4) $r + k \geq s + 1$, $k \geq 3$, I satisfies SDC_1 at level $s - g - k$, and $\text{depth}(Z_i(\mathbf{f})) \geq d - s + k$, for $0 \leq i \leq k$.

Consequently, having a finite acyclic complex whose components have sufficient depth, one can estimate the depth of the zeroth homology.

Theorem 2.6. *Let (R, \mathfrak{m}) be a CM local ring of dimension d , $I = (f_1, \dots, f_r)$ be an ideal with $\text{ht}(I) = g \geq 1$. Let $s \geq g$ and fix $0 \leq k \leq \min\{s, s - g + 2\}$. Suppose that one the following hypotheses holds*

- (i) $r + k \leq s$ and I satisfies SD condition, or
- (ii) $r + k \geq s + 1$, I satisfies SDC_1 at level $s - g - k$ and $\text{depth}(Z_i(\mathbf{f})) \geq d - s + k$ for $0 \leq i \leq k$, or
- (iii) $k \leq s - r + 2$ and I satisfies SD , or
- (iv) $\text{depth}(H_i(\mathbf{f})) \geq \min\{d - s + k - 2, d - g\}$, for $0 \leq i \leq k - 1$ and I satisfies SD , or
- (v) I is strongly Cohen-Macaulay.

Then for any s -residual intersection $J = (\mathfrak{a} : I)$, the complex ${}_k Z_{\bullet}^+$ is acyclic. Furthermore, $\text{Sym}_R^k(I/\mathfrak{a})$ if $1 \leq k \leq s - g + 2$, or the disguised residual intersection if $k = 0$, is CM of codimension s .

In the above theorem, the cases where $k = 0, 1$ and $s - g + 1$ are the most important ones. For, $k = 0$ is related to the disguised residual intersection, $k = 1$ is connected to R/\mathfrak{a} and $k = s - g + 1$ is about the canonical module of R/J .

Remark 2.7. Any of the conditions (iii)-(v) are stronger than (i) and (ii). Condition (i) in Theorem 2.6 implies that $\text{depth}(R/I) \geq d - s + k$ and condition (v) implies that $\text{depth}(R/I) \geq d - g \geq d - s$. The other conditions (ii), (iii) and (iv) yield $\text{depth}(R/I) \geq d - s + k - 2$. Therefore to have a better consequence of the theorem it may not be too much to ask $\text{depth}(R/I) \geq d - s$.

The first two items in the next corollary settle conjectures (3) and (4) mentioned in the introduction.

Corollary 2.8. Let (R, \mathfrak{m}) be a CM local ring of dimension d , $I = (f_1, \dots, f_r)$ be an ideal with $\text{ht}(I) = g \geq 1$. Suppose that $\text{depth}(R/I) \geq d - s$ and that I satisfies any of the conditions in Theorem 2.6 for $k = 1$, for instance if I satisfies SD . Then for any algebraic s -residual intersection $J = (\mathfrak{a} : I)$,

- (1) $\text{depth}(R/\mathfrak{a}) = d - s$ and thus \mathfrak{a} is minimally generated by s elements.
- (2) J is unmixed of height s .
- (3) $\text{Ass}(R/\mathfrak{a}) \subseteq \text{Ass}(R/I) \cup \text{Ass}(R/J)$. Furthermore if I is unmixed then the equality holds.
- (4) If the residual is arithmetic then $\text{Sym}_R^i(I/\mathfrak{a})$ is a faithful R/J -module for all i .

Proof. (1). Applying Theorem 2.6 for $k = 1$, we have $\text{depth}(I/\mathfrak{a}) = d - s$. Therefore the result follows from the exact sequence $0 \rightarrow I/\mathfrak{a} \rightarrow R/\mathfrak{a} \rightarrow R/I \rightarrow 0$ and the fact that $\text{depth}(R/I) \geq d - s$ by hypothesis.

(2). J has codimension at least s , thus for any $\mathfrak{p} \in \text{Ass}(R/J)$, $\text{ht}(\mathfrak{p}) \geq s$. On the other hand, $\text{Ass}(R/J) \subseteq \text{Ass}(R/\mathfrak{a})$ and any prime ideal in the latter has codimension less than s by (1) (see [BH, 1.2.13]), this yields the unmixedness of J .

(3). Let $\mathfrak{p} \in \text{Ass}(R/\mathfrak{a})$, then $\text{ht}(\mathfrak{p}) \leq s$, by (1). If $\text{ht}(\mathfrak{p}) < s$ or $\mathfrak{p} \not\supseteq J$, then $\mathfrak{a}_{\mathfrak{p}} = I_{\mathfrak{p}}$ and thus $\mathfrak{p} \in \text{Ass}(R/I)$ otherwise $\text{ht}(\mathfrak{p}) = s$ and $\mathfrak{p} \supseteq J$ which clearly means $\mathfrak{p} \in \text{Ass}(R/J)$. Verification of the last statement is straightforward.

(4) To see this, we refer to the proof of [Ha, Theorem 2.11] wherein it is shown that in the spectral sequence defined in (2.3), $K := (\infty E_{hor}^{0,0})_{[0]} \subseteq J \subseteq ({}^2 E_{hor}^{0,0})_{[0]}$. Moreover in the case where the residual is arithmetic this spectral sequence, locally in height s , collapses in the second page, hence all of these inclusions are equality. We just notice that $({}^2 E_{hor}^{0,0})_{[0]} = H_{\mathfrak{g}}^0(H_0(D_{\bullet}))_{[0]} = \bigcup_{i=0}^{\infty} (\gamma \cdot \text{Sym}_R^i(I) :_R \text{Sym}_R^{i+1}(I))$. Thus once the equality in the above line happens, $J = \gamma \cdot \text{Sym}_R^i(I) :_R \text{Sym}_R^{i+1}(I)$ for all i . Therefore $\text{Sym}_R^i(I/\mathfrak{a}) = \text{Sym}_R^{i+1}(I)/\gamma \cdot \text{Sym}_R^i(I)$ is a faithful R/J -module.

□

The following examples show the accuracy of the conditions in Theorem 2.6 and Corollary 2.8(4).

Example 2.9. Let $R = \mathbb{Q}[x_1, \dots, x_7]$, $M = \begin{pmatrix} 0 & x_1 & x_2 & x_3 \\ x_4 & x_5 & x_6 & x_7 \end{pmatrix}$ and $I = I_2(M)$. Set $f_0 = -x_4x_1$, $f_1 = -x_2x_4$, $f_2 = x_1x_6 - x_2x_5$, $f_3 = -x_3x_4$, $f_4 = x_1x_7 - x_3x_5$ and $f_5 = x_2x_7 - x_3x_6$. *Macaulay2* computations show that $\text{depth}(Z_1(I)) = 6$, $\text{depth}(Z_2(I)) = 2$ and $\text{depth}(Z_3(I)) = 6$ hence I satisfies SDC_1 at level 0 but not at level 1. However, Z_1 has enough depth, thence condition (ii) in Theorem 2.6 is fulfilled whenever $2 \leq s \leq 4$. The non-trivial case will be $s = 4$. Also $\text{depth}(R/I) = 4 > d - s = 3$. It then follows from Corollary 2.8 that for any 4-residual intersection $J = \mathfrak{a} : I$, $\text{ht}(J) = 4 = \mu(\mathfrak{a})$. It is easy to see that I satisfies G_∞ ; hence [EHU, 3.7bis] implies that $\ell(I) \geq 5$. On the other hand the Plücker relations show that $f_2f_3 - f_1f_4 + f_0f_5 = 0$. Hence I is not generated by analytically independent elements- that is $\ell(I) = 5$. Therefore the same proposition in *loc. cit.* proves that there must exist a 5-residual intersection $J = \mathfrak{a} : I$ such that $\text{ht}(J) > 5$. By the latter, Corollary 2.8 guarantees that for that ideal \mathfrak{a} , the module I/\mathfrak{a} cannot be CM and thus Theorem 2.6 does not work because SDC_1 at level 1 is not satisfied.

Another point about this example is that $\text{depth}(R/I^2) = 2$, according to *Macaulay2*; therefore $\text{Ext}^5(R/I^2, R) \neq 0$. The vanishing of this Ext module would be a sufficient to prove that I is 4-residually S_2 in [CEU, Theorem 4.1] whereas we saw in the above corollary that I is 4-residually unmixed.

Example 2.10. The arithmetic hypothesis in Corollary 2.8(4) cannot be dropped. Let $R = \mathbb{Q}[x, y]$, $I = (x, y)$ and $\mathfrak{a} = (x^2, y^2)$. Then $J = (\mathfrak{a}, xy)$ and $\text{Ann}(\text{Sym}_R^2(I/\mathfrak{a})) = (\mathfrak{a}I : I^2) = (x, y)$. The latter is not J , since the residual(linkage) is not arithmetic.

In the case where the residual intersection is geometric we have stronger corollaries.

Corollary 2.11. *Let (R, \mathfrak{m}) be a CM local ring of dimension d , $I = (f_1, \dots, f_r)$ an ideal with $\text{ht}(I) = g \geq 1$. Suppose that $\text{depth}(R/I) \geq d - s$ and that I satisfies any of the conditions in Theorem 2.6 for some $1 \leq k \leq s - g + 2$. Then for any geometric s -residual intersection $J = (\mathfrak{a} : I)$ and for that k ,*

- (1) $I^k/\mathfrak{a}I^{k-1} \simeq \text{Sym}_R^k(I/\mathfrak{a})$ and it is a faithful maximal Cohen-Macaulay R/J -module.
- (2) $\mathfrak{a}I^{k-1} = I^k \cap J$.
- (3) $I^k + J$ is a CM ideal of height $s + 1$ and $\text{ht}(I^k + J/J) = 1$.

Proof. (1). Consider the natural map $\varphi : \frac{\text{Sym}_R^k(I)}{\gamma \text{Sym}_R^{k-1}(I)} \rightarrow \frac{I^k}{\mathfrak{a}I^{k-1}}$ that sends T_i to f_i . φ is onto. Let $K = \text{Ker}(\varphi)$. We show that $\text{Ass}(K) = \emptyset$, hence $K = 0$. $\text{Ass}(K) \subseteq \text{Ass}(\frac{\text{Sym}_R^k(I)}{\gamma \text{Sym}_R^{k-1}(I)})$. By Theorem 2.6, $\text{depth}(\frac{\text{Sym}_R^k(I)}{\gamma \text{Sym}_R^{k-1}(I)}) = d - s$, hence for any associated prime \mathfrak{p} in the latter $\text{ht}(\mathfrak{p}) \leq s$. However, since J is a geometric residual, for any \mathfrak{p} of codimension $\leq s$, either $I_{\mathfrak{p}} = \mathfrak{a}_{\mathfrak{p}}$ or $I_{\mathfrak{p}} = (1)$. In the former case $\frac{\text{Sym}_R^k(I_{\mathfrak{p}})}{\gamma_{\mathfrak{p}} \text{Sym}_R^{k-1}(I_{\mathfrak{p}})} = 0$, for $k \geq 1$, which implies that $\mathfrak{p} \notin \text{Ass}(K)$. In the latter case, $\text{Sym}_R^k(I_{\mathfrak{p}}) = R_{\mathfrak{p}}$, hence $\varphi_{\mathfrak{p}}$ is an isomorphism, which means $K_{\mathfrak{p}} = 0$ in particular $\mathfrak{p} \notin \text{Ass}(K)$. Therefore $\text{Ass}(K) = \emptyset$, hence $K = 0$. The CM ness of $\text{Sym}_R^k(I/\mathfrak{a})$ was already shown in Theorem 2.6. To see that $I^k/\mathfrak{a}I^{k-1}$ is faithful one may

use Corollary 2.8 and part (1), or, just notice that $J \subseteq \mathfrak{a}I^{k-1} : I^k$ and that the equality holds locally at all associated primes of J which are of height s .

(2). Consider the canonical map $\psi : \frac{I^k}{\mathfrak{a}I^{k-1}} \rightarrow \frac{I^k+J}{J}$. By (2), $I^k/\mathfrak{a}I^{k-1}$ is CM of dimension $d-s$ which in conjunction with the fact that the residual is geometric shows that ψ is an isomorphism for all $0 \leq k \leq s-g+2$. Notice that $\frac{I^k+J}{J} \simeq \frac{I^k}{J \cap I^k}$. Thus the canonical map imply $J \cap I^k = \mathfrak{a}I^{k-1}$.

(3). By (1) and (2), $(I^k + J)/J$ is CM of dimension $d-s$. Also by Theorem 2.6, R/J is CM of dimension $d-s$. Thus the exact sequence $0 \rightarrow (I^k + J)/J \rightarrow R/J \rightarrow R/(I^k + J) \rightarrow 0$ implies $\text{depth}(R/I^k + J) \geq d-s-1$. On the other hand, since the residual is geometric, $\text{ht}(I^k + J) \geq s+1$ which in turn completes the proof. To see that $\text{ht}(I^k + J/J) = 1$, it is just enough to notice that R/J is CM. \square

We now proceed to determine the structure of the canonical module of R/J . This achievement is via a study of the type of an appropriate candidate for the canonical module. Recall that in a Noetherian local ring (R, \mathfrak{m}) the type of a finitely generated module M is the dimension of the R/\mathfrak{m} -vector space $\text{Ext}_R^{\text{depth}(M)}(R/\mathfrak{m}, M)$ and it is denoted by $r_R(M)$ or just $r(M)$.

Theorem 2.12. *Let (R, \mathfrak{m}) be a CM local ring of dimension d and $I = (f_1, \dots, f_r)$ be an ideal with $\text{ht}(I) = g \geq 2$. Let $J = (\mathfrak{a} : I)$ be an s -residual intersection of I and $1 \leq k \leq s-g+1$. Suppose that I is SCM. Then*

$$r_R(\text{Sym}_R^k(I/\mathfrak{a})) \leq \binom{r+s-g-k}{r-1} r_R(R).$$

Proof. First we deal with the case where $g = 2$. Consider the complex ${}_k\mathcal{Z}_\bullet^+$

$$0 \longrightarrow H_{\mathfrak{g}}^r(D_{r+s-1})_{[k]} \longrightarrow \cdots \xrightarrow{\phi_k} H_{\mathfrak{g}}^r(D_{r+k})_{[k]} \xrightarrow{\tau_k} (D_k)_{[k]} \longrightarrow \cdots \longrightarrow (D_0)_{[k]} \longrightarrow 0 \quad (2.11)$$

We study the depth of the components of this complex. $Z_{r-1} = R$ hence $\text{depth}({}_k\mathcal{Z}_{r+s-1}^+) = d$. For $k \leq j \leq s-2$, the first cycle which appears in D_{r+j} is Z_{r-s+j} and it has depth equal to $d = d-g+2$, by SCM condition, which is at least $d-s+(j+1)+1$. As for the right part of the complex we have $\text{depth}(Z_i) = d-g+2 \geq d-s+k+1$ for all $0 \leq i \leq k$, since by assumption $k \leq s-g+1$.

Summing up we have

$$\text{depth}({}_k\mathcal{Z}_i^+) \geq d-s+i+1 \text{ for all } i = 0, \dots, s-1. \quad (2.12)$$

Now, let $F_\bullet \rightarrow R/\mathfrak{m}$ be a free resolution of R/\mathfrak{m} . We put the double complex $\text{Hom}_R(F_\bullet, {}_k\mathcal{Z}_\bullet^+)$ in the third quadrant with $\text{Hom}_R(F_0, {}_k\mathcal{Z}_0^+)$ in the center. The two arisen spectral sequences are:

$${}^\infty E_{hor}^{-i,-j} = \begin{cases} \text{Ext}_R^j(R/\mathfrak{m}, \text{Sym}_R^k(I/\mathfrak{a})), & i = 0 \\ 0 & i \neq 0, \end{cases}$$

and ${}^1 E_{ver}^{-i,-j} = \text{Ext}_R^j(R/\mathfrak{m}, {}_k\mathcal{Z}_i^+)$. The latter vanishes for all (i, j) where $j-i \leq d-s$ and $i \neq s$, according to (2.12). The convergence of both complexes to the homology module of the total complex

implies that $\text{Ext}_R^{d-s}(R/\mathfrak{m}, \text{Sym}_R^k(I/\mathfrak{a})) \simeq {}^\infty E_{\text{ver}}^{-s,-d}$. The latter is a submodule of $\text{Ext}_R^d(R/\mathfrak{m}, {}_k\mathcal{Z}_s^+)$. Therefore $r_R(\text{Sym}_R^k(I/\mathfrak{a})) \leq \dim_{R/\mathfrak{m}}(\text{Ext}_R^d(R/\mathfrak{m}, {}_k\mathcal{Z}_s^+))$.

Following from the construction of \mathcal{D}_\bullet , $D_{r+s-1} = S(-r+1-s)$. Hence

$${}_k\mathcal{Z}_s^+ = H_{\mathfrak{g}}^r(S)_{[-r+1-s+k]}.$$

To calculate the dimension of the above Ext module we just need to know how many copies of R appear in the inverse polynomial structure of the above local cohomology module. The answer is simply the positive solutions of the numerical equation $\alpha_1 + \cdots + \alpha_r = r-1+s-k$ which is $\binom{r-k+s-2}{r-1}$. Therefore $r_R(\text{Sym}_R^k(I/\mathfrak{a})) \leq \binom{r+s-2-k}{r-1} r(R)$ in the case where $g=2$.

Now, let $\text{ht}(I) = g \geq 2$. We choose a regular sequence \mathfrak{a} of length $g-2$ inside \mathfrak{a} which is a part of a minimal generating set of \mathfrak{a} . I/\mathfrak{a} is still SCM by [H, Corollary 1.5], moreover $J/\mathfrak{a} = \mathfrak{a}/\mathfrak{a} : I/\mathfrak{a}$. Recall that $\mu(\mathfrak{a}) = s$ by Corollary 2.8, therefore J/\mathfrak{a} is an $(s-g+2)$ -residual intersection of I/\mathfrak{a} which is of height 2. Hence we return to the case $g=2$, so that

$$r_R(\text{Sym}_R^k(I/\mathfrak{a})) = r_{R/\mathfrak{a}}(\text{Sym}_{R/\mathfrak{a}}^k(I/\mathfrak{a})) \leq \binom{r+(s-g+2)-2-k}{r-1} r_{R/\mathfrak{a}}(R/\mathfrak{a}) = \binom{r+s-g-k}{r-1} r_R(R).$$

(see [BH, Exercise 1.2.26]).

□

Remark 2.13. Clearly the above proof works as well for $k=0$, hence if I is SCM of $\text{ht}(I) \geq 2$ and K is a disguised s -residual intersection of I then

$$r_R(R/K) \leq \binom{r+s-g}{r-1} r_R(R).$$

Everything is now ready to explain the structure of the canonical module.

Theorem 2.14. *Suppose that (R, \mathfrak{m}) is a Gorenstein local ring of dimension d . Let I be a SCM ideal with $\text{ht}(I) = g$ and $J = (\mathfrak{a} : I)$ be an arithmetic s -residual intersection of I . Then $\omega_{R/J} \simeq \text{Sym}_R^{s-g+1}(I/\mathfrak{a})$. It is isomorphic to $I^{s-g+1} + J/J$ in the geometric case; Furthermore R/J is generically Gorenstein and, R/J is Gorenstein if and only if I/\mathfrak{a} is principal.*

Proof. Without loss of generality, we may suppose that $g \geq 2$, otherwise we may add a new variable x to R and consider ideals $\mathfrak{a} + (x)$ and $I + (x)$ in the local ring $R[x]_{\mathfrak{m}+(x)}$ wherein we have $J + (x) = (\mathfrak{a} + (x)) : (I + (x))$.

R/J is CM by Theorem 2.6. Respectively, according to Theorem 2.6, Corollary 2.8 and Theorem 2.12, $\text{Sym}_R^{s-g+1}(I/\mathfrak{a})$ is a maximal CM, faithful R/J -module and of type 1. Therefore it is the canonical module of R/J by [BH, 3.3.13]. In the geometric case $\text{Sym}_R^{s-g+1}(I/\mathfrak{a}) \simeq I^{s-g+1} + J/J$ by Corollary 2.11.

To see that R/J is generically Gorenstein, notice that for any prime ideal $\mathfrak{p} \supseteq J$ of height s . $r((R/J)_{\mathfrak{p}}) = \mu((\omega_{R/J})_{\mathfrak{p}}) = \mu(\text{Sym}_{R_{\mathfrak{p}}}^{s-g+1}((I/\mathfrak{a})_{\mathfrak{p}}))$ which is 1 since the residual is arithmetic by the assumption. Finally let $t = \mu(I/\mathfrak{a})$ then $r(R/J) = \mu(\omega_{R/J}) = \binom{s-g+t}{t-1}$ which is one if and only if $t=1$. □

3. HILBERT FUNCTION, CASTELNUOVO-MUMFORD REGULARITY

Although the family $\{ {}_k\mathcal{Z}_\bullet^+ \}$ does not consist of free complexes, however it can approximate R/J and $Sym_R^k(I/\mathfrak{a})$ very closely. Several numerical invariants or functions such as Castelnuovo-Mumford regularity, Projective dimension and Hilbert function can be estimated via this family. In [CEU], Chardin, Eisenbud and Ulrich restated an old question ([S]) asking for which open sets of ideals \mathfrak{a} the Hilbert function of R/\mathfrak{a} depends only on the degrees of the generators of \mathfrak{a} . In [CEU], the authors consider the following two conditions

- (A1) If the Hilbert function of R/\mathfrak{a} is constant on the open set of ideals \mathfrak{a} generated by s forms of the given degrees such that $\text{codim}(\mathfrak{a} : I) \geq s$; and
- (A2) if the Hilbert function of $R/(\mathfrak{a} : I)$ is constant on this set.

It is shown in [CEU, Theorem 2.1] that ideals with some slight depth conditions in conjunction with G_{s-1} or G_s satisfy these two conditions. In this direction we have the following result which in addition to show the validity of the above conditions (A1) and (A2) under some sliding depth condition, provides a method to compute the desired Hilbert functions.

Proposition 3.1. *Let R be a CM graded ring over an Artinian local ring R_0 , I and \mathfrak{a} be two homogenous ideals of R . Let $J = (\mathfrak{a} : I)$ be an s -residual intersection. Let $0 \leq k \leq s - g + 2$ and suppose that I satisfies any of the following hypotheses:*

- (1) $1 \leq s \leq 2$ and $s = k$, or
- (2) $r + k \geq s + 1$, $k \leq 2$ and I satisfies SDC_1 at level $s - g - k$, or
- (3) $r + k \leq s$ and I satisfies SD , or
- (4) $r + k \geq s + 1$, $k \geq 3$, I satisfies SDC_1 at level $s - g - k$, and $\text{depth}(Z_i(\mathfrak{f})) \geq d - s + k$, for $0 \leq i \leq k$.

Then the Hilbert function of the disguised s -residual intersection, if $k = 0$, and that of $Sym_R^k(I/\mathfrak{a})$, if $1 \leq k \leq s - g + 2$, depends only on the **degrees** of the generators of \mathfrak{a} and the Koszul homologies of I . In particular if I satisfies any of the above conditions for $k = 1$ then the Hilbert function of R/\mathfrak{a} satisfies (A1).

Proof. By Proposition 2.5, the standing assumptions on I imply the acyclicity of ${}_k\mathcal{Z}_\bullet^+$. Hence the Hilbert function of $H_0({}_k\mathcal{Z}_\bullet^+)$ is derived from the Hilbert functions of the components of ${}_k\mathcal{Z}_\bullet^+$ which, according to (2.1) and (2.9), are just some direct sums of the Koszul's cycles of I shifted by the twists appearing in the Koszul complex $K_\bullet(\gamma, S)$. Since the Hilbert functions of Koszul's cycles are inductively calculated in terms of those of the Koszul's homologies, the Hilbert function of $H_0({}_k\mathcal{Z}_\bullet^+)$ depends on the Koszul's homologies of I and just the degrees of the generators of \mathfrak{a} . In the case where $k = 1$ we know the Hilbert function of $I/\mathfrak{a} = Sym_R^1(I/\mathfrak{a})$ so that we know the Hilbert function of R/\mathfrak{a} by $0 \rightarrow I/\mathfrak{a} \rightarrow R/\mathfrak{a} \rightarrow R/I \rightarrow 0$. \square

The next important numerical invariant associated to an algebraic or geometric object is the Castelnuovo-Mumford regularity. Here we present an upper bound for the regularity of the disguised s -residual intersection K and the regularity of $Sym_R^k(I/\mathfrak{a})$ (if $1 \leq k \leq s - g + 1$).

Assume that $R = \bigoplus_{n \geq 0} R_n$ is a positively graded \ast local Noetherian ring of dimension d over a Noetherian local ring (R_0, \mathfrak{m}_0) set $\mathfrak{m} = \mathfrak{m}_0 + R_+$. Suppose that I and \mathfrak{a} are homogeneous ideals of R generated by homogeneous elements f_1, \dots, f_r and a_1, \dots, a_s , respectively. Let $\deg f_t = i_t$ for all $1 \leq t \leq r$ with $i_1 \geq \dots \geq i_r$ and $\deg a_t = l_t$ for $1 \leq t \leq s$. For a graded ideal \mathfrak{b} , the sum of the degrees of a minimal generating set of \mathfrak{b} is denoted by $\sigma(\mathfrak{b})$.

Proposition 3.2. *With the same notations just introduced above, let (R, \mathfrak{m}) be a CM \ast local ring, I an ideal with $\text{ht}(I) = g \geq 2$ and $s \geq g$. Suppose that $J = (\mathfrak{a} : I)$ is an s -residual intersection of I and K is the disguised s -residual intersection of I w.r.t \mathfrak{a} . Suppose that any of the following conditions hold for some $0 \leq k \leq s - g + 1$,*

- (i) $r + k \leq s$ and I satisfies SD_1 condition;
- (ii) $r + k \geq s + 1$, I satisfies SDC_2 at level $s - g - k$ and $\text{depth}(Z_i(\mathfrak{f})) \geq d - s + k + 1$ for $0 \leq i \leq k$.

Then if $k \geq 1$,

$$\text{reg}(\text{Sym}_R^k(I/\mathfrak{a})) \leq \text{reg}(R) + \dim(R_0) + \sigma(\mathfrak{a}) - (s - g + 1 - k) \text{indeg}(I/\mathfrak{a}) - s.$$

And if $k = 0$, $\text{reg}(R/K) \leq \text{reg}(R) + \dim(R_0) + \sigma(\mathfrak{a}) - (s - g + 1) \text{indeg}(I/\mathfrak{a}) - s$.

Proof. The proof of this fact is essentially the same as that of [Ha, Theorem 3.6] wherein the case of $k = 0$ is verified. The crucial point to repeat that proof is that we have to change the structure of the families $\{ {}_k \mathcal{Z}_\bullet^+ \}$ from the beginning by substituting the tails with a free complex. Another way to prove this result in the case where I is SCM, may be by applying the " $g = 2$ " trick in Theorem 2.12. \square

The inequality in the above proposition becomes more interesting in particular cases, for example if R is a polynomial ring over a field and $k = 1$ this inequality reads as

$$\text{reg}_R(I/\mathfrak{a}) + (s - g) \text{indeg}(I/\mathfrak{a}) \leq \sigma(\mathfrak{a}) - s.$$

Hence we have a numerical obstruction for $J = (\mathfrak{a} : I)$ to be an s -residual intersection.

At the other end of the spectrum we have $k = s - g + 1$, this case corresponds to the canonical module of R/J .

The next proposition improves [Ha, Proposition 3.5] by removing the G_s condition.

Proposition 3.3. *Suppose that (R, \mathfrak{m}) is a positively graded Gorenstein \ast local ring, over a Noetherian local ring (R_0, \mathfrak{m}_0) , with canonical module $\omega_R = R(b)$ for some integer b . Let I be a homogeneous SCM ideal with $\text{ht}(I) = g$, $\mathfrak{a} \subset I$ homogeneous and $J = (\mathfrak{a} : I)$ be an arithmetic s -residual intersection of I . Then $\omega_{R/J} = \text{Sym}_R^{s-g+1}(I/\mathfrak{a})(b + \sigma(\mathfrak{a}))$.*

In particular, if $\dim(R_0) = 0$ then $\text{reg}(R/J) = \text{reg}(R) + \sigma(\mathfrak{a}) - \text{indeg}(\text{Sym}_R^{s-g+1}(I/\mathfrak{a})) - s$.

Proof. We may and do assume that $g \geq 2$. First suppose that $g = 2$. We keep the notations defined at the beginning of the section 2 to define ${}_k \mathcal{Z}_\bullet^+$ complex. Considering R as a subalgebra of $S = R[T_1, \dots, T_r]$, we write the degrees of an element x of R as the 2-tuple $(\deg x, 0)$ with the second entry zero. So that $\deg f_t = (i_t, 0)$ for all $1 \leq t \leq r$, $\deg a_t = (l_t, 0)$ for all $1 \leq t \leq s$, $\deg T_t = (i_t, 1)$ for all $1 \leq t \leq r$, and thus $\deg \gamma_t = (l_t, 1)$ for all $1 \leq t \leq s$. With these notations the \mathcal{Z} -complex has the following shape

$$\mathcal{Z}_\bullet : 0 \rightarrow Z_{r-1} \otimes_R S(0, -r+1) \rightarrow \cdots \rightarrow Z_1 \otimes_R S(0, -1) \rightarrow Z_0 \otimes_R S \rightarrow 0.$$

Consequently,

$$Z_{r-1} = R(-\sum_{t=1}^r i_t, 0) \otimes_R S(0, -r+1),$$

and, taking into account that a_1, \dots, a_s is a minimal generating set of \mathfrak{a} by Corollary 2.8,

$$D_{r+s-1} = S(-\sum_{t=1}^r i_t - \sigma(\mathfrak{a}), -s-r+1).$$

It then follows that ${}_{(s-1)}\mathcal{Z}_s^+ = H_{\mathfrak{g}}^r(D_{r+s-1})_{[*], [s-1]}$ is isomorphic to

$$H_{\mathfrak{g}}^r(S)(-\sum_{t=1}^r i_t - \sigma(\mathfrak{a}), 0)_{[*], [-r]} \simeq (RT_1^{-1} \cdots T_r^{-1}(-\sum_{t=1}^r i_t))(-\sigma(\mathfrak{a})) \simeq R(-\sigma(\mathfrak{a})). \quad (3.1)$$

Notice that $\deg(T_j^{-1}) = (-i_j, -1)$.

We next turn to the proof of Theorem 2.12 and consider the spectral sequences therein for $k = s-g+1 = s-1$. We have ${}^1E_{ver}^{-s, -d} = \text{Ext}_R^d(R/\mathfrak{m}, {}_{(s-1)}\mathcal{Z}_s^+)$ which is isomorphic to $\text{Ext}_R^d(R/\mathfrak{m}, R(b))(-b - \sigma(\mathfrak{a}))$ by (3.1). The latter is in turn homogeneously isomorphic to $R/\mathfrak{m}(-b - \sigma(\mathfrak{a}))$, since R is Gorenstein. The inclusion $\text{Ext}_R^{d-s}(R/\mathfrak{m}, \text{Sym}_R^{s-g+1}(I/\mathfrak{a})) \simeq {}^\infty E_{hor}^{0, -(d-s)} \hookrightarrow {}^1E_{ver}^{-s, -d} = \text{Ext}_R^d(R/\mathfrak{m}, {}_{s-1}\mathcal{Z}_s^+)$ is indeed an isomorphism, since the latter is a vector space of dimension one. In the graded case all of the homomorphisms in the proof of Theorem 2.12 are homogeneous, therefore the above isomorphism implies that $\text{Ext}_R^{d-s}(R/\mathfrak{m}, \text{Sym}_R^{s-g+1}(I/\mathfrak{a})) \simeq R/\mathfrak{m}(-b - \sigma(\mathfrak{a}))$. Since we already know that in the local case $\omega_{R/J} = \text{Sym}_R^{s-2+1}(I/\mathfrak{a})$ by Corollary 2.14, hence in the graded case we have $\omega_{R/J} = \text{Sym}_R^{s-2+1}(I/\mathfrak{a})(b + \sigma(\mathfrak{a}))$.

Now suppose that $g \geq 2$ and let $\mathfrak{a} = a_1, \dots, a_{g-2}$ be a regular sequence in \mathfrak{a} which is a part of its minimal generating set. Then $\omega_{R/\mathfrak{a}} = \frac{R}{\mathfrak{a}}(b + \sigma(\mathfrak{a}))$ and moreover we go back to the case where $g = 2$. We then have

$$\omega_{R/J} = \text{Sym}_{R/\mathfrak{a}}^{(s-g+2)-2+1} \left(\frac{I/\mathfrak{a}}{\mathfrak{a}/\mathfrak{a}} \right) ((b + \sigma(\mathfrak{a})) + \sigma(\mathfrak{a}/\mathfrak{a})) = \text{Sym}_R^{s-g+1}(I/\mathfrak{a})(b + \sigma(\mathfrak{a})).$$

□

Finally, one may ask whether under the G_s condition our methods can contribute more information about the structure of residual intersections than what is known so far. The next result shows that in the presence of the G_s condition the complex ${}_0\mathcal{Z}_\bullet^+$ approximates R/J for any (algebraic) s -residual intersection J . In particular the above facts about the regularity and the Hilbert function hold for R/J .

Proposition 3.4. *Let (R, \mathfrak{m}) be a Gorenstein local ring and suppose that I is SCM evenly linked to an ideal which satisfies G_s . Then the disguised s -residual intersections of I with respect to an ideal \mathfrak{a} , K , is the same as the algebraic residual intersection $J = \mathfrak{a} : I$.*

Proof. It is shown in [HU, Theorem 5.3] that under the above assumptions on I there exists a deformation (R', I') of (R, I) such that I' is SCM and G_s . Let $I = (f_1, \dots, f_r)$, $\mathfrak{a} = (a_1, \dots, a_s)$ and suppose that $a_i = \sum_{j=1}^r c_{ij} f_j$. Let s_{ij} be a lifting of c_{ij} to R' and f'_i be a lifting of f_i to R' . Set $X = (x_{ij})$ to be an $s \times r$ matrix of invariants, $\alpha := (\alpha_1, \dots, \alpha_s) = X \cdot (f'_1, \dots, f'_r)^t$. Let $\tilde{R} = R'[x_{ij}]_{(\mathfrak{m} + (x_{ij} - s_{ij}))}$ and $\tilde{J} :=$

$(\alpha_1, \dots, \alpha_s) :_{\tilde{R}} I' \tilde{R}$. By [HU, Lemma 3.2] \tilde{J} is a geometric s -residual intersection of $I' \tilde{R}$. We construct the complex ${}_0\mathcal{Z}_{\bullet}^+(\alpha_1, \dots, \alpha_s; f'_1, \dots, f'_s)$ in \tilde{R} and set $\tilde{R}/\tilde{K} = H_0({}_0\mathcal{Z}_{\bullet}^+(\alpha_1, \dots, \alpha_s; f'_1, \dots, f'_s))$. Since \tilde{J} is a geometric residual, $\tilde{J} = \tilde{K}$.

Assume that π is the projection from \tilde{R} to R which sends x_{ij} to c_{ij} . By [Ha, Proposition 2.13], $R/\pi(\tilde{K}) = H_0({}_0\mathcal{Z}_{\bullet}^+)$ and by [HU, Theorem 4.7], $\pi(\tilde{J}) = J$. Therefore $R/K = R/\pi(\tilde{K}) = R/\pi(\tilde{J}) = R/J$. \square

4. ARITHMETIC RESIDUAL INTERSECTIONS

In this section we scrutinize the structure of the ${}_0\mathcal{Z}_{\bullet}^+$ complex in the case where I/\mathfrak{a} is cyclic. We find that the homologies of ${}_0\mathcal{Z}_{\bullet}^+$ determines the uniform annihilator of non-zero Koszul homologies. The importance of this fact is that although the Koszul complex is an old object, not much is known about the annihilator of higher homologies. Besides the rigidity and differential graded algebra structure of the homologies, there is an interesting paper of Corso, Huneke, Katz and Vasconcelos [CHKV] wherein the authors make efforts to find out "whether the annihilators of non-zero Koszul homology modules of an unmixed ideal is contained in the integral closure of that ideal". However, it seems that the situation of mixed ideals is more involved, for example for $\mathfrak{a} = (x^2 - xy, y^2 - xy, z^2 - zw, w^2 - zw)$, (taken from *loc. cit.*), $\text{Ann}(H_1) = \bar{\mathfrak{a}}$, the integral closure, and $\text{Ann}(H_2) = \sqrt{\bar{\mathfrak{a}}}$. This example shows that the intersection of the annihilators has a better structure than the union. This is an interesting example for us since $(\mathfrak{a} : \bar{\mathfrak{a}})$ is a 4-residual intersection with $\bar{\mathfrak{a}}/\mathfrak{a}$ cyclic and $\bar{\mathfrak{a}}$ satisfies SD.

Since a part of our results works well with arithmetic s -residual intersections, it is worth mentioning some general properties of these ideals.

Proposition 4.1. *Let R be a Noetherian ring, $\mathfrak{a} = (a_1, \dots, a_s) \subset I$ be ideals of R and $J = \mathfrak{a} : I$ be an s -residual intersection. Then*

- (i) *J is an arithmetic s -residual intersection if and only if $\text{ht}(0 : \bigwedge^2(I/\mathfrak{a})) \geq s + 1$.*
- (ii) *If 2 is unit in R then $I + J \subseteq (0 : \bigwedge^2(I/\mathfrak{a}))$.*
- (iii) *The followings statements are equivalent*
 - (1) *For all $i \leq s$, an i -residual intersection of I exists.*
 - (2) *For any prime ideal $\mathfrak{p} \supset I$ such that $\text{ht}(\mathfrak{p}) \leq s - 1$, $\mu(I_{\mathfrak{p}}) \leq \text{ht}(\mathfrak{p}) + 1$.*
 - (3) *There exist $a_1, \dots, a_s \in I$ such that for $i \leq s - 1$, $J_i = (a_1, \dots, a_i) : I$ is an arithmetic i -residual intersection and $J = (a_1, \dots, a_s) : I$ is an (algebraic) s -residual intersection.*
- (iv) *J is an arithmetic s -residual intersection if and only if there exists an element $b \in I$ such that $\text{ht}(\mathfrak{a}, b) : I \geq s + 1$.*

Proof. (i). $J = \mathfrak{a} : I$ is an arithmetic s -residual if and only if $\text{ht}(\text{Fitt}_j(I/\mathfrak{a})) \geq s + j$ for $j = 0, 1$ [Eis, Proposition 20.6]. Moreover $\text{Fitt}_j(I/\mathfrak{a})$ and the annihilator of $\bigwedge^{j+1}(I/\mathfrak{a})$ have the same radical [Eis, Exercise 20.10] which yields the assertion. (ii) is straightforward. (iii) goes along the same line as the proof of [U, Lemma 1.4]. (iv) follows from [U, Lemma 1.3]. \square

The concept of s -parsimonious ideals first appeared in [CEU]. It is shown that weakly s -residually S_2 ideals which satisfy G_s are s -parsimonious [CEU, 3.1]. However if I is an ideal which satisfies the condition of Corollary 2.8 then it follows from [CEU, 3.3 (a)] that I is s -parsimonious. So that we have

Remark 4.2. Let (R, \mathfrak{m}) be a CM local ring of dimension d , I an ideal such that $\text{depth}(R/I) \geq d - s$ and that I satisfies any of the conditions in Theorem 2.6 for $k = 1$. Then for any arithmetic s -residual intersection $J = (\mathfrak{a} : I)$, there exists an element $b \in I$ such that $J = (\mathfrak{a} : b)$.

This remark is another evidence to study the cases where I/\mathfrak{a} is cyclic.

To exploit the structure of $H_\bullet({}_0\mathcal{Z}_\bullet^+(\mathfrak{a}; \mathfrak{a}, b))$, the following lemma is crucial.

Lemma 4.3. Let $\mathfrak{a} = (a_1, \dots, a_s)$, $I = (\mathfrak{f}) = (b, a_1, \dots, a_s)$ and consider the complex \mathcal{D}_\bullet defined in (2.1). Let Z'_\bullet , B'_\bullet and H'_\bullet be the cycles, boundaries and homologies of the Koszul complex $K_\bullet(a_1, \dots, a_s)$. Put $\widetilde{B}_i = (B'_i :_{Z'_i} b)$ and $S = R[T_0, T_1, \dots, T_s]$ with standard grading. Then

(i)

$$H_i(\mathcal{D}_\bullet) \cong \frac{(Z'_i \otimes_R R[T_0])(-i)}{(\widetilde{B}_i \otimes_R T_0 R[T_0])(-i)},$$

in particular

$$(H_i(\mathcal{D}_\bullet)_{T_0})_{[0]} \cong bH_i(a_1, \dots, a_s) \text{ for all } i.$$

(ii) If R is a graded ring and b, a_1, \dots, a_s are homogeneous then we have a homogeneous isomorphism

$$(H_i(\mathcal{D}_\bullet)_{T_0})_{[0]} \cong bH_i(a_1, \dots, a_s)((i+1) \deg(b)) \text{ for all } i.$$

Proof. To compute the homology modules of \mathcal{D}_\bullet , we consider the two spectral sequences arising from its double complex structure (2.1). We put this double complex in the second quadrant and locate $K_0(\gamma) \otimes_S \mathcal{Z}_0(\mathfrak{f})$ in the corner.

Set $S = R[T_0, T_1, \dots, T_s]$ where T_0 corresponds to b and T_i to a_i . Then $(\gamma_1 \cdots \gamma_s) = (T_1 \cdots T_s)$ is a regular sequence on $\mathcal{Z}_i = Z_i[T_0, T_1, \dots, T_s](-i)$ with $Z_i = Z_i(b, a_1, \dots, a_s)$. We then have ${}^1E_{ver}^{-i, j} = H_j(K_\bullet(\gamma) \otimes \mathcal{Z}_i) = 0$ for all $j \geq 1$ and all i . Also $(S/(T_1, T_2, \dots, T_s)) \otimes_S \mathcal{Z}_i = R[T_0] \otimes_R Z_i$ in which we consider $R[T_0]$ as an S -module by trivial multiplication $T_i R[T_0] = 0$, $1 \leq i \leq s$. It then follows that ${}^1E_{ver}^{0, \bullet}$ is the complex

$$0 \rightarrow Z_{s+1} \otimes_R R[T_0](-s-1) \xrightarrow{\partial_{s+1}} \cdots \rightarrow Z_1 \otimes_R R[T_0](-1) \xrightarrow{\partial_1} Z_0 \otimes_R R[T_0] \rightarrow 0 \quad (4.1)$$

wherein the differentials are induced by those in $\mathcal{Z}_\bullet(\mathfrak{f})$. As well,

$$H_i({}^1E_{ver}^{0, \bullet}) = H_i(\mathcal{Z}_\bullet \otimes_S (S/(T_1, T_2, \dots, T_s)))(-i) \quad (4.2)$$

Considering the convergence of the spectral sequences to the homology module of the total complex \mathcal{D}_\bullet , we have $H_i(\mathcal{D}_\bullet) = H_i({}^1E_{ver}^{0, \bullet}) = \text{Ker}(\partial_i) / \text{Im}(\partial_{i+1})$. We compute this homology in two steps.

Step 1 $\text{Ker}(\partial_i)$: Let $z_i = (\sum_j r_j e_{j_1} \wedge e_{j_2} \wedge \dots \wedge e_{j_i})\omega(T_0) \in Z_i[T_0](-i)$. We study two cases: First, if $j_1 \neq 0$ for every j which appears in the above presentation of z_i , in this case $z_i \in Z'_i[T_0]$ and moreover

$$\partial_i(z_i) = \sum_j r_j \left(\sum_{l=1}^i e_{j_1} \wedge \dots \wedge e_{j_{l-1}} \wedge e_{j_{l+1}} \wedge \dots \wedge e_{j_i} T_l \omega(T_0) \right) = 0$$

where the last vanishing is due to the fact that $T_i R[T_0] = 0$ for all $1 \leq i \leq s$. It follows that $z_i \in \text{Ker}(\partial_i)$.

Second, we show that if $j_1 = 0$ for some j that appears in z_i then $z_i \notin \text{Ker}(\partial_i)$. Writing $z_i = z'_i + z''_i \wedge e_0$, where z'_i and z''_i do not involve e_0 , one has

$$\partial_i(z_i) = \partial_i(z'_i) + \partial_{i-1}(z''_i) \wedge e_0 + (-1)^{i+1} T_0 z''_i$$

by the DG-algebra structure of $(\mathcal{Z}_\bullet, \partial')$. Since z'_i and z''_i do not involve e_0 , it follows that $\partial_i(z'_i) = 0$ and $\partial_{i-1}(z''_i) = 0$ by the multiplication on $R[T_0]$, thus $\partial_i(z_i) = (-1)^{i+1} T_0 z''_i \neq 0$. Therefore $\text{ker}(\partial_i) = Z_i(a_1, \dots, a_s)[T_0](-i) = Z'_i[T_0](-i)$.

Step 2 $\text{Im}(\partial_{i+1})$: To compute the image, we consider an arbitrary element

$$z_{i+1}\omega(T_0) = \left(\sum_j r_j e_{j_1} \wedge e_{j_2} \wedge \dots \wedge e_{j_{i+1}} \right) \omega(T_0) \in Z_{i+1}[T_0](-i-1)$$

and write $z_{i+1} = z'_{i+1} + z''_{i+1} \wedge e_0$, where z'_{i+1} and z''_{i+1} do not involve e_0 . Recall that $Z_{i+1} = Z_{i+1}(\mathbf{f})$. Consider the Koszul complex $(K_\bullet(\mathbf{f}), d_\bullet)$. We have

$$0 = d(z_{i+1}) = d(z'_{i+1}) + d(z''_{i+1}) \wedge e_0 + (-1)^i b z''_{i+1}.$$

Since z'_{i+1} and z''_{i+1} do not involve e_0 , the above equation implies that $d(z''_{i+1}) = 0$ and $b z''_{i+1} = d((-1)^{i+1} z'_{i+1})$; so that $z''_{i+1} \in (B'_i :_{Z'_i} b)$.

Conversely, for any $z''_{i+1} \in (B'_i :_{Z'_i} b)$, $b z''_{i+1} = d((-1)^{i+1} z'_{i+1})$ for some $z'_{i+1} \in K'_{i+1}$, hence $z_{i+1} = (z'_{i+1} + z''_{i+1} \wedge e_0) \in Z_{i+1}$.

On the other hand $\partial(z_{i+1}) = \partial(z'_{i+1}) + \partial(z''_{i+1} \wedge e_0) = 0 + \partial(z''_{i+1} \wedge e_0) = (-1)^i T_0 z''_{i+1}$.

It follows that $\text{Im}(\partial_{i+1}) = (T_0 \widetilde{B}_i[T_0])(-i)$. Finally the above two steps show that

$$H_i(\mathcal{D}_\bullet) = \frac{(Z'_i \otimes_R R[T_0])(-i)}{(\widetilde{B}_i \otimes_R T_0 R[T_0])(-i)}.$$

To see the last assertion, just notice that

$$((Z'_i[T_0]_{T_0})(-i))_{[0]} = Z'_i[T_0, T_0^{-1}]_{[-i]} = T_0^{-i} Z'_i \quad \text{and} \quad ((\widetilde{B}_i T_0[T_0]_{T_0})(-i))_{[0]} = T_0^{-i} \widetilde{B}_i.$$

Therefore

$$(H_i(\mathcal{D}_\bullet)_{T_0})_{[0]} \simeq Z'_i / \widetilde{B}_i = Z'_i / (B'_i :_{Z'_i} b) = H'_i / (0 :_{H'_i} b) \simeq b H_i(a_1, \dots, a_s). \quad (4.3)$$

(ii). In the graded case every homomorphism in the above argument is homogenous except the first and the last isomorphism in (4.3). To see the desired shift, notice that the bi-degree of T_0 is $(\deg(b), 1)$. Hence considering $(H_i(\mathcal{D}_\bullet)_{T_0})$ in bidegree $(0, 0)$, we have

$$(H_i(\mathcal{D}_\bullet)_{T_0})_{[(0,0)]} \simeq \frac{T_0^{-i}((Z'_i)_{[i \deg(b)])}}{T_0^{-i}((\widetilde{B}_i)_{[i \deg(b)])}} \simeq \left(\frac{Z'_i}{\widetilde{B}_i} \right)_{[i \deg(b)]}.$$

Moreover there exists the following homogeneous exact sequence which yields the assertion

$$0 \rightarrow \widetilde{B}_i \rightarrow Z'_i \rightarrow bH_i(a_1, \dots, a_s)(\deg(b)) \rightarrow 0.$$

□

Now, we are ready to explain our main Theorem in this section. The dependence of the homology modules of ${}_0\mathcal{Z}_\bullet^+(\mathbf{a}, \mathbf{f})$ to the generating sets becomes clear by this theorem, furthermore it attracts attention to study the uniform annihilator of the Koszul homology modules.

Theorem 4.4. *Let R be a (Noetherian) ring, $\mathbf{a} = (a_1, \dots, a_s)$ and $I = (\mathbf{f}) = (b, a_1, \dots, a_s)$. Then $H_i({}_0\mathcal{Z}_\bullet^+(\mathbf{a}, \mathbf{f})) \simeq bH_i(a_1, \dots, a_s)$ for all $i \geq 1$ and $H_0({}_0\mathcal{Z}_\bullet^+(\mathbf{a}, \mathbf{f})) \simeq R/(\mathbf{a} : b)$; furthermore in the graded case $H_i({}_0\mathcal{Z}_\bullet^+(\mathbf{a}, \mathbf{f})) \simeq bH_i(a_1, \dots, a_s)((i+1)\deg(b))$ for $i \geq 1$.*

Proof. We study the spectral sequences arising from the third quadrant double complex $C_\mathfrak{g}^\bullet \otimes \mathcal{D}_\bullet$ with $C_\mathfrak{g}^0 \otimes D_0$ in the center. Since T_0, T_1, \dots, T_s is a regular sequence on D_i 's, $({}^1E_{ver})_{[0]} = H_\mathfrak{g}^{s+1}(\mathcal{D}_\bullet)_{[0]}$. The latter is \mathcal{Z}_\bullet^+ by definition (2.5) which is the truncated complex ${}_0\mathcal{Z}_\bullet^+(\mathbf{a}, \mathbf{f})_{\geq 1}$.

On the other hand, in the horizontal spectral sequence we deal with $H_\mathfrak{g}^j(H_i(\mathcal{D}_\bullet))$. $H_i(\mathcal{D}_\bullet)$ is annihilated by (T_1, \dots, T_s) by (4.2) which implies that $H_\mathfrak{g}^j(H_i(\mathcal{D}_\bullet)) = H_{(T_0)}^j(H_i(\mathcal{D}_\bullet))$ for all i and j , also notice that $H_{(T_0)}^0(H_i(\mathcal{D}_\bullet))_{[0]} = 0$ for $i \geq 1$ since $\text{indeg}(D_i) = i$. Hence

$$({}^2E_{hor}^{-i, -j})_{[0]} = \begin{cases} H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_{[0]} & i = 0 \text{ and } j = 0, \\ H_{(T_0)}^1(H_i(\mathcal{D}_\bullet))_{[0]} = (H_i(\mathcal{D}_\bullet)_{T_0})_{[0]} & i \geq 1 \text{ and } j = 1, \\ 0 & \text{otherwise.} \end{cases} \quad (4.4)$$

Considering both spectrals at a same time, we have

$$\begin{array}{cccc} ({}^2E_{hor})_{[0]} : & 0 & 0 & H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_{[0]} \\ \dots & H_{(T_0)}^1(H_2(\mathcal{D}_\bullet))_{[0]} & H_{(T_0)}^1(H_1(\mathcal{D}_\bullet))_{[0]} & H_{(T_0)}^1(H_0(\mathcal{D}_\bullet))_{[0]} \\ \dots & 0 & 0 & 0 \\ \mathcal{Z}_\bullet^+ = ({}^1E_{ver})_{[0]} : & \vdots & \vdots & \vdots \\ & \swarrow & \swarrow & \swarrow \\ H_\mathfrak{g}^{s+1}(\mathcal{D}_{s+1})_{[0]} & \longrightarrow & H_\mathfrak{g}^{s+1}(\mathcal{D}_s)_{[0]} & \longrightarrow & H_\mathfrak{g}^{s+1}(\mathcal{D}_{s-1})_{[0]} & \longrightarrow & \dots \end{array}$$

By the convergence of the spectral sequences, $H_{i+1}({}_0\mathcal{Z}_\bullet^+) = H_i(\mathcal{Z}_\bullet^+) = H_{(T_0)}^1(H_{i+1}(\mathcal{D}_\bullet))_{[0]}$ for $i \geq 1$. Quite generally the latter is isomorphic to $(H_{i+1}(\mathcal{D}_\bullet)_{T_0})_{[0]}$ which in turn is isomorphic to $bH_{i+1}(a_1, \dots, a_s)$ by

Lemma 4.3. In the case where $i = 0$ the convergence of the spectral sequences provides the following exact sequence

$$0 \longrightarrow H_{(T_0)}^1(H_1(\mathcal{D}_\bullet))_0 \longrightarrow H_0(\mathcal{Z}_\bullet^+) \xrightarrow{\psi} H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_0 \longrightarrow 0. \quad (4.5)$$

According to (2.5), $H_0(\mathcal{Z}_\bullet^+) = \text{Coker}(\phi_0)$ and ψ is used to define τ_0 in (2.7). Therefore $\text{Ker}(\psi) = H_1({}_0\mathcal{Z}_\bullet^+)$ and its image determines $H_0({}_0\mathcal{Z}_\bullet^+)$ that we will compute below.

Since the horizontal spectral stabilizes in the second step, ${}^\infty E_{hor}^{0,0} = {}^2 E_{hor}^{0,0} = H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))$. Moreover $\text{end}(H_{\mathfrak{g}}^{s+1}(\mathcal{D}_{s+1})) = 0$ therefore $({}^\infty E_{hor}^{0,0})_{[i]} = 0$ for $i \geq 1$ which in particular implies that $H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_{[1]} = 0$. Specially we then have $T_0 H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_{[0]} = 0$. Recall that $H_0(\mathcal{D}_\bullet) = \text{Sym}(I)/\gamma \text{Sym}(I)$ and that we consider $\mathcal{S}_I := \text{Sym}_R(I)$ as an $S = R[T_0, \dots, T_s]$ -module via the ring homomorphism $S \rightarrow \mathcal{S}_I$ sending T_0 to b and T_i to a_i as an element of $(\mathcal{S}_I)_{[1]} = I$. Hence $T_0 H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_{[0]} = 0$ as an element in $(\text{Sym}(I)/\gamma \text{Sym}(I))_{[1]} = I/\mathfrak{a}$ is equivalent to say that $H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_{[0]} \subseteq (\mathfrak{a} : b)$.

On the other hand, $H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_{[0]} = (\mathfrak{a} : b) \cup (\cup_{i=1}^\infty (\mathfrak{a} \text{Sym}_R^i(I) :_R \text{Sym}_R^{i+1}(I)))$, hence $H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_0 \supseteq (\mathfrak{a} : b)$, which yields $H_{(T_0)}^0(H_0(\mathcal{D}_\bullet))_{[0]} = (\mathfrak{a} : b)$.

Further, by Lemma 4.3, $H_{(T_0)}^1(H_1(\mathcal{D}_\bullet))_{[0]} = (H_1(\mathcal{D}_\bullet)_{T_0})_{[0]} = bH_1(f_1, f_2, \dots, f_s)$. Substituting these facts in the short exact sequence (4.5) the assertion follows. \square

As a summary we have the following corollary

Corollary 4.5. *Let R be a (Noetherian) ring, $\mathfrak{a} = (\mathbf{a}) = (a_1, \dots, a_s)$ and $I = (\mathbf{f}) = (b, a_1, \dots, a_s)$. If $bH_i(a_1, \dots, a_s) = 0$ for all $i \geq 1$, then there exists an exact complex*

$$0 \longrightarrow \mathcal{Z}_s^+ \longrightarrow \mathcal{Z}_{s-1}^+ \longrightarrow \dots \longrightarrow \mathcal{Z}_0^+ \longrightarrow (\mathfrak{a} :_R b) \longrightarrow 0$$

wherein $\mathcal{Z}_i^+ = \bigoplus_{j=i+1}^r \mathcal{Z}_j(\mathbf{f})^{\oplus n_j}$ for some positive integers n_j and r . In particular the index of the lowest cycle which appears in the components increases along the complex.

Theorem 4.4 in conjunction with other conditions which imply the acyclicity of the complex ${}_0\mathcal{Z}_\bullet^+$ yields to non trivial facts about the uniform annihilators of non-zero koszul homologies.

Corollary 4.6. *Let (R, \mathfrak{m}) be a CM local ring of dimension d , $I = (b, a_1, \dots, a_s)$ satisfies SD. If $\dim(bH_i(a_1, \dots, a_s)) \leq d - s$ for all $i \geq 1$ then*

- $bH_i(a_1, \dots, a_s) = 0$ for all $i \geq 1$ and
- $\text{depth}(R/((a_1, \dots, a_s) : I)) \geq d - s$.

Proof. We consider the complex ${}_0\mathcal{Z}_\bullet^+(\mathfrak{a}; (b, \mathfrak{a}))$. By Theorem 4.4 the hypothesis on the dimensions implies that this complex is acyclic locally at codimension $s - 1$. Then appealing the acyclicity lemma, the sliding depth hypothesis shows that the complex is acyclic on the punctured spectrum and finally it is acyclic (the same technique which applied to prove Proposition 2.5). Once more applying Theorem 4.4, we conclude the first assertion. The depth inequality follows from the proof of Theorem 2.6. \square

As a final remark in this section, we notice that if $(\mathfrak{a} : I)$ is an s -residual intersection then $\dim(IH_i(\mathfrak{a})) \leq d - s$ for all i . So that the latter property may be considered as a generalization of algebraic residual Intersection.

5. UNIFORM ANNIHILATOR OF KOSZUL'S HOMOLOGIES

Motivated by the results in the previous section, we concentrate on the uniform annihilator of Koszul homologies in this section. In the main theorem of this section we see that in any residual intersection $J = \mathfrak{a} : I$ the sliding depth condition passes from I to \mathfrak{a} . This fact was known to experts only in the presence of the G_∞ condition [U, 1.8(3)+1.12] which is not that surprising since I is then generated by a d -sequence. However this was unknown even for perfect ideals of height 2 which are not G_∞ , see [EU] for such an example.

The next interesting result in this section is Corollary 5.6, in which we show that for any residual intersection $J = \mathfrak{a} : I$ with I satisfies sliding depth, I is contained in the uniform annihilator of the non-zero Koszul homologies of \mathfrak{a} , hence a kind of universal properties for such ideals.

Next proposition is fundamental to prove the aforementioned results though it is also interesting by itself.

Proposition 5.1. *Let (R, \mathfrak{m}) be a CM local ring of dimension d , $\mathfrak{a} = (f_1, f_2, \dots, f_s)$, $I = (f_1, f_2, \dots, f_r)$ be ideals with $s \leq r$, $J = (\mathfrak{a} : I)$ and $g = \text{ht}(I) = \text{ht}(\mathfrak{a}) \geq 1$. Let n be an integer and suppose that $\text{ht}(J) \geq n \geq 1$ and that $\text{depth}(H_l(f_1, f_2, \dots, f_r)) \geq d - r + j$ for $l \geq r - n + 2$. Then for any k with $r \geq k \geq s$ and any $j \geq k - n + 2$,*

- i) $\text{depth}(H_j(f_1, f_2, \dots, f_k)) \geq d - k + j$; and
- ii) $I \subseteq \text{Ann}(H_j(f_1, f_2, \dots, f_k))$.

Supplement, both assertions hold for sequences of the form $(f_1, \dots, f_s, f_{i_1}, \dots, f_{i_{k-s}})$ where $\{f_{i_1}, \dots, f_{i_{k-s}}\} \subseteq \{f_{s+1}, \dots, f_r\}$.

Proof. The proof is by a **recursive** induction on k . By assumption, the result is true for $k = r$. Suppose that it is true for $k \leq r$. We now apply a recursive induction on $j \geq (k - 1) - n + 2$ to show the following claims

- 1) $f_k H_j(f_1, f_2, \dots, f_{k-1}) = 0$;
- 2) $\text{depth}(H_j(f_1, f_2, \dots, f_{k-1})) \geq d - (k - 1) + j$;
- 3) $\text{depth}(Z_j(f_1, f_2, \dots, f_{k-1})) \geq d - (k - 1) + j$;
- 4) $\text{depth}(B_j(f_1, f_2, \dots, f_{k-1})) \geq d - (k - 1) + j$.

Since these claims are clear for $j > k - g$, we suppose that $j \geq (k - 1) - n + 3$ and that these claims hold for j . A simple depth chasing between cycles, boundaries and homologies shows that (2) – (4), for $j - 1$, follows from (1) for $j - 1$ and the induction hypotheses of (2) – (4) for j . We just notice that condition (1) provides the following exact sequence

$$0 \longrightarrow H_j(f_1, \dots, f_{k-1}) \longrightarrow H_j(f_1, \dots, f_k) \longrightarrow H_{j-1}(f_1, \dots, f_{k-1}) \longrightarrow 0.$$

We then proceed to prove (1) for $j - 1$ while assuming the four claims hold for j . To do this, we show that $f_k H_{j-1}(f_1, f_2, \dots, f_{k-1})_{\mathfrak{p}} = 0$ by an induction on $\text{ht}(\mathfrak{p})$. For $\text{ht}(\mathfrak{p}) < n$ it is clear, since $I_{\mathfrak{p}} = a_{\mathfrak{p}}$. Now let \mathfrak{q} be a prime ideal with $\text{ht}(\mathfrak{q}) \geq n$ and suppose that (1) to (4) hold locally for all \mathfrak{p} such that $\mathfrak{p} \subset \mathfrak{q}$. We may, and will, replace (R, \mathfrak{m}) by $(R_{\mathfrak{q}}, \mathfrak{q}R_{\mathfrak{q}})$. Considering the following exact sequence, out of the Koszul complex,

$$0 \longrightarrow Z_j(f_1, \dots, f_{k-1}) \longrightarrow \bigwedge^j (R^{k-1}) \longrightarrow B_{j-1}(f_1, \dots, f_{k-1}) \longrightarrow 0,$$

we obtain $\text{depth}(B_{j-1}) \geq \min\{\text{depth}(\bigwedge^j (R^{k-1})), \text{depth}(Z_j) - 1\} \geq d - (k - 1) + j - 1$. By assumption, $j \geq k - n + 2$, which implies that $\text{depth}(B_{j-1}(f_1, \dots, f_{k-1})) \geq d - n + 2 \geq 2$. This time

$$0 \longrightarrow B_{j-1}(f_1, \dots, f_{k-1}) \longrightarrow Z_{j-1}(f_1, \dots, f_{k-1}) \longrightarrow H_{j-1}(f_1, \dots, f_{k-1}) \longrightarrow 0$$

implies that $\text{depth}(H_{j-1}) \geq 1$, since $\text{depth}(Z_{j-1}) \geq 1$, as $\dim(R_{\mathfrak{q}}) \geq n \geq 1$.

We now consider the exact sequence

$$0 \longrightarrow H_j(f_1, \dots, f_{k-1}) \longrightarrow H_j(f_1, \dots, f_{k-1}, f_k) \longrightarrow \Gamma_{j-1} \longrightarrow 0$$

where $\Gamma_{j-1} = (0 :_{H_{j-1}(f_1, \dots, f_{k-1})} f_k)$.

The induction hypotheses imply that

$$\begin{aligned} \text{depth}(\Gamma_{j-1}) &\geq \min\{\text{depth}(H_j(f_1, \dots, f_{k-1}, f_k)), \text{depth}(H_j(f_1, \dots, f_{k-1})) - 1\} \\ &\geq \min\{d - k + j, d - (k - 1) + j - 1\} \geq d - k + j \geq d - k + (k - n + 2) = d - n + 2 \geq 2. \end{aligned}$$

At last, we have the inclusion $\Gamma_{j-1} \subseteq H_{j-1}$ where the former has depth at least 2 and the latter has depth at least 1 while the equality holds on the punctured spectrum, by induction hypothesis, so that this inclusion must be an equality, which proves the claim(1) for $j - 1$.

Applying the above argument for $k = r$ we have $\text{depth}(H_j(f_1, \dots, f_{r-1})) \geq d - (r - 1) + j$ for all $j \geq (r-1) - n + 2$ and that $f_r H_j(f_1, f_2, \dots, f_{r-1}) = 0$. However, as far as we fix f_1, \dots, f_s we may change the role of f_r with any one of $\{f_{s+1}, \dots, f_r\}$. In other words, we have $\text{depth}(H_j(f_1, \dots, f_s, \dots, \hat{f}_i, \dots, f_r)) \geq d - (r - 1) + j$ for all $j \geq (r - 1) - n + 2$ and that $f_i H_j(f_1, \dots, f_s, \dots, \hat{f}_i, \dots, f_r) = 0$. Also notice that for any ideal $\mathfrak{a} \subseteq I' \subseteq I$, $\text{ht}(\mathfrak{a} : I') \geq \text{ht}(\mathfrak{a} : I) \geq n$. Therefore if the assertion (i) holds for $(f_1, \dots, f_s, f_{i_1}, \dots, f_{i_{k-s}})$ where $\{f_{i_1}, \dots, f_{i_{k-s}}\} \subseteq \{f_{s+1}, \dots, f_r\}$, then the above argument shows that (i) holds for $(f_1, \dots, f_s, f_{i_1}, \dots, f_{i_{k-s-1}})$ and moreover $f_{i_{k-s}} H_j(f_1, \dots, f_s, f_{i_1}, \dots, f_{i_{k-s-1}}) = 0$ for $j \geq k - 1 - n + 2$. Since $f_{i_{k-s}}$ varies over the set $\{f_{s+1}, \dots, f_r\}$ we must have $I H_j(f_1, \dots, f_s, f_{i_1}, \dots, f_{i_{k-s-1}}) = 0$ for $j \geq k - 1 - n + 2$ which completes the proof. \square

We next want to show, in Lemma 5.3, that under the sliding depth condition, there exists a similar depth inequality for cycles of the Koszul complex. For this, we need the following lemma which is the reminiscence of the inductive Koszul's long exact sequence.

Lemma 5.2. *Let R be a commutative ring, $\{f_0, f_1, \dots, f_r\} \subseteq R$, Z_\bullet, B_\bullet the Koszul's cycles and boundaries with respect to sequence $\{f_1, f_2, \dots, f_r\}$, Z'_\bullet the Koszul's cycles with respect to sequence (f_0, f_1, \dots, f_r) and $\Gamma_\bullet = (B_\bullet :_{Z_\bullet} f_0)$. Then for any $j \geq 0$, there exists an exact sequence*

$$0 \longrightarrow Z_j \longrightarrow Z'_j \longrightarrow \Gamma_{j-1} \longrightarrow 0$$

Proof. The Koszul complex $K_\bullet(f_1, f_2, \dots, f_r)$ is canonically a sub complex of $K_\bullet(f_0, f_1, \dots, f_r)$. We will denote the differential of both complexes by d . Let $\{e_0, e_1, \dots, e_r\}$ be the canonical basis of R^{r+1} . Every $\theta \in Z'_j$ can be written uniquely as a sum $\theta = e_0 \wedge w + v$ where e_0 does not appear in terms w and v . Since $\theta \in Z'_j$

$$0 = d(\theta) = d(e_0 \wedge w + v) = f_0 w - e_0 \wedge d(w) + d(v).$$

It follows that

$$\begin{cases} e_0 \wedge d(w) = 0 \\ f_0 w = -d(v) \end{cases} \text{ hence } \begin{cases} d(w) = 0 \\ f_0 w = -d(v) \end{cases} \text{ therefore } w \in \Gamma_{j-1}.$$

We define $\varphi : Z'_j \rightarrow \Gamma_{j-1}$ to be $\varphi(\theta) = \varphi(e_0 \wedge w + v) = w$. The homomorphism φ is well defined, as the expression is unique. It is onto, since for any $w \in \Gamma_{j-1}$ there is $v \in K_j(f_1, f_2, \dots, f_r)$ such that $f_0 w = -d(v)$. Thence $\theta = e_0 \wedge w + v \in K_j(f_0, f_1, \dots, f_r)$ is the preimage of w .

Moreover, $Z_j = \ker(\varphi)$. On one hand, if $\theta \in Z'_j$ belongs to Z_j , then there is no e_0 in its decomposition in terms of the canonical basis of R^{r+1} , consequently $\theta = e_0 \wedge w + v = v$ and $w = 0$. On the other hand, if $0 = d(\theta) = d(e_0 \wedge w + v) = w$, then $\theta = v$. Therefore $\theta \in Z'_j \cap K_j(f_1, f_2, \dots, f_r) = Z_j$. \square

The next Lemma is needed in the proof of the main theorem 5.4. Notice that the sliding depth condition does not pass automatically from homologies to cycles because there is a level restriction, see Proposition 2.4.

Lemma 5.3. *Let (R, \mathfrak{m}) be a CM local ring of dimension d , $\mathfrak{a} = (f_1, f_2, \dots, f_s)$ and $I = (f_1, f_2, \dots, f_r)$ be ideals with $s \leq r$. Let $J = (\mathfrak{a} : I)$ and $g = \text{ht}(I) = \text{ht}(\mathfrak{a}) \geq 1$. Suppose that $\text{ht}(J) \geq n \geq 1$ and that I satisfies SD, then for $k \geq s$ and $j \geq k - n + 2$,*

$$\text{depth}(Z_j(f_1, f_2, \dots, f_k)) \geq d - k + j + 1.$$

Proof. By Proposition 5.1, $IH_j(f_1, f_2, \dots, f_k) = 0$ whenever $s \leq k \leq r$ and $j \geq k - n + 2$. Therefore by Lemma 5.2, we have the following exact sequences for $k - 1 \geq s$ and $j - 1 \geq (k - 1) - n + 2$

$$0 \longrightarrow Z_j(f_1, f_2, \dots, f_{k-1}) \longrightarrow Z_j(f_1, f_2, \dots, f_k) \longrightarrow Z_{j-1}(f_1, f_2, \dots, f_{k-1}) \longrightarrow 0.$$

We will prove the result by a recursive induction on k . If $k = r$, then it follows from Proposition 2.4. Suppose that the result holds for $k \geq s + 1$. Since $Z_j(f_1, f_2, \dots, f_{k-1})$ satisfies SDC_1 for $j \geq (k - 1) - g + 1$, we can use a new induction on j to conclude that:

$$\text{depth}(Z_{j-1}(f_1, f_2, \dots, f_{k-1})) \geq d - (k - 1) + (j - 1) + 1 = d - k + j + 1.$$

\square

We are now ready to prove the main Theorem of this section. As we mentioned at the beginning of the section, this fact was known just in the presence of the G_∞ condition, under which the problem reduced to study the properties of ideals generated by d -sequences. Here the only assumption is the sliding depth and that $\text{depth}(R/I) \geq d - s$.

Theorem 5.4. *Let (R, \mathfrak{m}) be a CM local ring of dimension d and $J = (\mathfrak{a} : I)$ be an s -residual intersection with $s \geq \text{ht}(I) \geq 1$. If I satisfies SD and $\text{depth}(R/I) \geq d - s$, then \mathfrak{a} satisfies SD as well.*

Proof. We apply Lemma 5.3 for $n = s$, it then follows that

$$\text{depth}(Z_j(f_1, f_2, \dots, f_s)) \geq d - s + j + 1$$

for $j \geq 2$. The hypothesis $\text{depth}(R/I) \geq d - s$ enables us to apply Corollary 2.8 which in turn implies that $\text{depth}(R/\mathfrak{a}) \geq d - s$; so that $\text{depth}(\mathfrak{a}) \geq d - s + 1$ and hence $\text{depth}(Z_1(\mathfrak{a})) \geq d - s + 2$ that is \mathfrak{a} satisfies the SDC_1 condition which is equivalent to SD by Proposition 2.4. \square

In the next proposition we prove a partial converse of Theorem 5.4.

Proposition 5.5. *Let (R, \mathfrak{m}) be a CM local ring of dimension d , $\mathfrak{a} = (f_1, f_2, \dots, f_s)$ and $I = (f_1, f_2, \dots, f_r)$ with $s \leq r$. Let $J = (\mathfrak{a} : I)$ be an s -residual intersection. If \mathfrak{a} satisfies SD, then for any $s \leq k \leq r$ and $j \geq k - s + 1$,*

- (i) $\text{depth}(H_j(f_1, f_2, \dots, f_k)) \geq d - k + j$ and
- (ii) $\text{depth}(Z_j(f_1, f_2, \dots, f_k)) \geq d - k + j + 1$.

Proof. We will use induction on k . If $k = s$, then (ii) is clear by Proposition 2.4. Suppose that the result holds for $k \geq s + 1$. Then by induction hypothesis, we have

$$\text{depth}(H_j(f_1, f_2, \dots, f_k)) \geq d - k + j \geq d - s + 1$$

By a similar argument as in the proof of Proposition 5.1, $IH_j(f_1, f_2, \dots, f_k) = 0$ for all $j \geq k - s + 1$. Then we have the short exact sequences below provided by Lemma 5.2 for any $j \geq k + 1 - s + 1$

$$0 \longrightarrow H_j(f_1, f_2, \dots, f_k) \longrightarrow H_j(f_1, f_2, \dots, f_{k+1}) \longrightarrow H_{j-1}(f_1, f_2, \dots, f_k) \longrightarrow 0$$

and

$$0 \longrightarrow Z_j(f_1, f_2, \dots, f_k) \longrightarrow Z_j(f_1, f_2, \dots, f_{k+1}) \longrightarrow Z_{j-1}(f_1, f_2, \dots, f_k) \longrightarrow 0.$$

A depth chasing then completes the proof. \square

The next corollary of Theorem 5.4 shows a tight relation between the uniform annihilator of non-zero Koszul homology modules and residual intersections.

Corollary 5.6. *Let (R, \mathfrak{m}) be a CM local ring of dimension d , I satisfies SD and $\text{depth}(R/I) \geq d - s$ with $s \geq \text{ht}(I) \geq 1$. Let $J = (\mathfrak{a} : I)$ be an s -residual intersection and use $H_j(\mathfrak{a})$ to denote the j 'th Koszul homology module with respect to a minimal generating set of \mathfrak{a} . Then*

$$I \subseteq \bigcap_{j \geq 1} \text{Ann}(H_j(\mathfrak{a})). \quad (\star)$$

Furthermore, the equality happens in the following cases, if $\text{depth}(R/I) \geq d - s + 1$ (hence $s \geq g + 1$).

- If $\mu(I_{\mathfrak{p}}) \leq s - i$ for a fix positive integer i and for all $\mathfrak{p} \in \text{Ass}(R/I)$, then $I = \text{Ann}(H_j(\mathfrak{a}))$ for all $1 \leq j \leq i$.
- If R is Gorenstein and $\text{Ass}(R/I) = \text{min}(R/I)$ then $I = \bigcap_{j \geq 1} \text{Ann}(H_j(\mathfrak{a}))$.

Proof. By Corollary 2.8, \mathfrak{a} is minimally generated by s elements. According to Theorem 5.4, for any j , $\text{depth}(H_j(\mathfrak{a})) \geq d - s + j$, in particular $\text{ht}(\mathfrak{p}) \leq s - 1$ for any $\mathfrak{p} \in \text{Ass}(H_j(\mathfrak{a}))$ with $j \geq 1$ c.f. [BH, 1.2.13].

To show that $IH_j(\mathfrak{a}) = 0$, we show that it has no associated prime. Notice that $IH_j(\mathfrak{a}) \subseteq H_j(\mathfrak{a})$, hence for any $\mathfrak{p} \in \text{Ass}(IH_j(\mathfrak{a}))$, $\text{ht}(\mathfrak{p}) \leq s - 1$ for which $I_{\mathfrak{p}} = \mathfrak{a}_{\mathfrak{p}}$. Therefore $(IH_j(\mathfrak{a}))_{\mathfrak{p}} = 0$ as desired.

To see the equality, we localize at associated primes of I which have height at most $s - 1$ so that $I_{\mathfrak{p}} = \mathfrak{a}_{\mathfrak{p}}$ for them. Now let (a_1, \dots, a_s) be a minimal generating set of \mathfrak{a} . If $\mu(I_{\mathfrak{p}}) \leq s - i$ then $H_j(a_1, \dots, a_s)_{\mathfrak{p}}$ has a direct summand of $(R/\mathfrak{a})_{\mathfrak{p}}$ for all $1 \leq j \leq i$ and thus its annihilator is $\mathfrak{a}_{\mathfrak{p}} = I_{\mathfrak{p}}$. In the case where R is Gorenstein and $\text{Ass}(R/I) = \text{min}(R/I)$, we see that locally at the associated primes of R/I , $I_{\mathfrak{p}}$ is unmixed and moreover $(a_1, \dots, a_s)_{\mathfrak{p}}$ is not a regular sequence since $s > \text{ht}(\mathfrak{p})$. Hence for $t = \text{ht}(I_{\mathfrak{p}})$, $\text{Ann}(H_{s-t}(\mathfrak{a}_{\mathfrak{p}}))$ is the unmixed part of $\mathfrak{a}_{\mathfrak{p}} = I_{\mathfrak{p}}$ which is $I_{\mathfrak{p}}$. \square

The next amazing example shows a benefit of relaxing the G_s condition in our Theorems. However we took this example from [EU] wherein Eisenbud and Ulrich used it to show the necessity of the G_s condition in their theorems.

Example 5.7. Let $(R, \mathfrak{m}) = \mathbb{Q}[x_1, \dots, x_5]_{(x_1, \dots, x_5)}$, $I = I_4(N)$ where

$$N = \begin{pmatrix} x_2 & x_3 & x_4 & x_5 \\ x_1 & x_2 & x_3 & x_4 \\ 0 & x_1 & x_2 & 0 \\ 0 & 0 & x_1 & x_2 \\ 0 & 0 & 0 & x_1 \end{pmatrix}.$$

Let Δ_i be the minors obtained by omitting i 'th row of N , we then consider \mathfrak{a} to be the ideal generated by entries of

$$\left(\begin{array}{ccccc} \Delta_1 & -\Delta_2 & \Delta_3 & -\Delta_4 & \Delta_5 \end{array} \right) \begin{pmatrix} x_1 & 0 & 0 & 0 & 0 \\ 0 & x_5 & 0 & 0 & 0 \\ x_3 & x_4 & x_5 & x_3 & 0 \\ 0 & x_3 & x_4 & x_5 & 0 \\ 0 & x_2 & x_3 & x_4 & x_5 \end{pmatrix}.$$

$J = (\mathfrak{a} : I)$ is a 5-residual intersection of I . I is a perfect ideal of height 2 hence it is SCM. Easily one can see that any prime that contains I must contain $\mathfrak{p} = (x_1, x_2)$ that is I is \mathfrak{p} -primary, since it is unmixed. The minimal presentation of I is given by N ; so that $\text{Fitt}_4(I) = \mathfrak{m}$ as well $\text{Fitt}_3(I) \not\subseteq \mathfrak{p}$ because $x_4^2 - x_3x_5 \notin \mathfrak{p}$, $\text{Fitt}_2(I) \subseteq \mathfrak{p}$ because any combination of 3 "diagonal" elements contains either 0 or x_1 or x_2 . Thus $\mu(I_{\mathfrak{p}}) = 3$, in particular I does not satisfy G_3 . However it completely suits in our Corollary 5.6 for $s = 5$ and $i = 2$, hence $\text{Ann}(H_1(\mathfrak{a})) = \text{Ann}(H_2(\mathfrak{a})) = I$. Also, since $\text{ht}(J) = 5$, any prime ideal of height 2 which contains \mathfrak{a} must contain I that means $I = \mathfrak{a}^{unm}$. Since the ring is Gorenstein, we then have $I = \mathfrak{a}^{unm} = \text{Ann}(H_3(\mathfrak{a}))$ (our Macaulay2 system was not able to calculate $\text{Ann}(H_1(\mathfrak{a}))$ directly!).

We notice that Corollary 5.6 presents a universal property for any ideal \mathfrak{a} such that $J = (\mathfrak{a} : I)$ is an s -residual intersection of I . For instance in the above example for any other \mathfrak{a} such that $J = (\mathfrak{a} : I)$ is a 5-residual intersection of I , we have $\text{Ann}(H_1(\mathfrak{a})) = \text{Ann}(H_2(\mathfrak{a})) = \text{Ann}(H_3(\mathfrak{a})) = \mathfrak{a}^{unm} = I$.

An unpublished result of G.Levin, [V2, Theorem 5.26] yields in [CHKV, Corollary 2.7] that *If R is a Noetherian ring and \mathfrak{a} is an ideal of finite projective dimension n , then $(\text{Ann}(H_1(\mathfrak{a})))^{n+1} \subseteq \mathfrak{a}$.* Combining this result with the above Corollary 5.6, one faces the strange fact that \mathfrak{a} and I have a same radical if one adopts the conditions of Corollary 5.6. However the following simple example (taken from [Hu]) shows that [CHKV, Corollary 2.7] is not correct, and thus the Levin's result is not true.

Example 5.8. Let $X = \begin{pmatrix} x_1 & x_3 & x_5 & x_7 \\ x_2 & x_4 & x_6 & x_8 \end{pmatrix}$ be a generic 2×4 matrix, $J = I_2(X)$, $\mathfrak{a} = (x_1x_4 - x_2x_3, x_1x_6 - x_2x_5, x_1x_8 - x_2x_7)$ and $I = (x_1, x_2)$. Then $J = \mathfrak{a} : I$ is a geometric 3-residual intersection of I , J is prime and $\mathfrak{a} = I \cap J$ hence \mathfrak{a} is radical. Furthermore $\text{Ann}(H_1(\mathfrak{a})) = \mathfrak{a}^{unm} = I$ which is not contained in \mathfrak{a}

A simpler example would be the following degeneration of the above example. Let $\mathfrak{a} = (y_1^2 - y_2^2, y_1y_3, y_2y_3) \in k[y_1, \dots, y_4]$. Then $\text{Ann}(H_1(\mathfrak{a})) = (y_1, y_2) \neq \text{rad}(\mathfrak{a})$.

At the end, we make a conjecture and ask a question. The conjecture will imply in turn the Cohen-Macaulayness of algebraic residual intersections for ideals with sliding depth condition.

Conjecture 5.9. If (R, \mathfrak{m}) is a CM local ring of dimension d and I satisfies SD with $\text{depth}(R/I) \geq d - s$. Then the disguised s -residual intersection of I coincides with the algebraic s -residual intersection.

Here is a peculiar question arises from Corollary 5.6.

Question 5.10. Does the equality hold in (\star) , if $s > \text{ht}(I)$?

Finally we present, without proof (a proof is based on Proposition 5.1), the following proposition which is an effort toward Question 5.10.

Proposition 5.11. *Let (R, \mathfrak{m}) be a CM local ring of dimension d , $\mathfrak{a} = (f_1, f_2, \dots, f_s)$, $I = (f_1, f_2, \dots, f_r)$ ideals with $s \leq r$ and $\text{ht}(I) = \text{ht}(\mathfrak{a}) \geq 1$. Suppose that $\text{ht}(\mathfrak{a} : I) \geq n \geq 1$.*

- *If $s + 1 \leq r$ and $\text{depth}(R/I) \geq d - n + 1$, then $\text{Ann}(H_j(f_1, f_2, \dots, f_r)) = I$ for $0 \leq j \leq r - s$.*

- If I satisfies SD , then for $\mu \geq s$ and $\mu - g \geq j \geq \mu - n + 2$,

$$\text{Ann}(H_j(f_1, f_2, \dots, f_\mu)) = \bigcap_{i=0}^{\mu-s} \text{Ann}(H_{j-i}(f_1, f_2, \dots, f_s))$$

Acknowledgment. The authors would like to thank Bernd Ulrich for encouragement and illuminating questions during his stays in Brazil in 2013 and 2014. A part of this work was done while the first author visiting the university of Utah supported by the PDE-fellowship, 233035/2014-1, from CNPq-Brazil, he expresses his gratitude to CNPq and thanks the university of Utah for the hospitality. Thanks to E. Tavanfar, A. Boocher and S. Iyengar for reading the draft and correcting some important typos. We also thank the referee for his/her careful reading of the manuscript and useful comments.

REFERENCES

- [AN] M. Artin and M. Nagata, *Residual intersection in Cohen-Macaulay rings*, J.Math. Kyoto Univ. (1972), 307–323.
- [BH] W. Bruns, J. Herzog, *Cohen–Macaulay Rings*, revised version, Cambridge University Press, Cambridge, 1998.
- [CEU] M. Chardin, D.Eisenbud and B. Ulrich, *Hilbert Functions, Residual Intersections, and Residually S_2 Ideals*, Composito Mathematica **125** (2001), 193–219.
- [CEU1] M. Chardin, D.Eisenbud and B. Ulrich, *Hilbert Series of Residual Intersections*, Compositio Mathematica, **151** (2015), no. 9, 1663 –1687
- [CHKV] A.Corso, C.Huneke, D. Katz, W. Vasconcelos, *Integral closure of ideals and annihilators of homology*, Commutative algebra, 3348, Lect. Notes Pure Appl. Math., 244, Chapman and Hall/CRC, Boca Raton, FL, 2006.
- [Eis] D. Eisenbud, *Commutative Algebra with a view toward algebraic geometry*, Graduate Texts in Math. **150**, Springer, New York, 1995.
- [EHU] D. Eisenbud, C. Huneke, B. Ulrich, *order ideals and a generalized height theorem*, Math. Annalen **330** (2004), 417–439.
- [EU] D.Eisenbud and B. Ulrich, *Residual intersections and duality*, arXiv:1309.2050.
- [Fu] W. Fulton, *Intersection Theory*, second edition, Springer, 1998.
- [H] R. Hartshorne, *Algebraic Geometry*, Springer, 1977.
- [Ha] S. H. Hassanzadeh, *Cohen-Macaulay residual intersections and their Castelnuovo-Mumford regularity*, Trans. Amer. Math. Soc. **364**(2012), 6371–6394.
- [HSV] J. Herzog, A. Simis, and W. Vasconcelos, *Koszul homology and blowing-up rings*, Commutative Algebra (Trento, 1981), Lecture Notes in Pure and Appl. Math., vol. 84, Marcel Dekker, NY,(1983), pp. 79–169.
- [Hu] C. Huneke, *Strongly Cohen-Macaulay schemes and residual intersections*, Trans. Amer. Math. Soc. **277**(1983), 739–763.
- [HU] C. Huneke, B. Ulrich, *Residual Intersection*, J. reine angew. Math.**390**(1988), 1–20.
- [M] H. Matsumura, *Commutative ring theory*, Cambridge University Press, Cambridge, 1986.
- [S] R. Stanley, *Weyl groups, the Hard Lefschetz theorem and the Sperner property*, SIAMJ. Algebra Discrete Math. **1** (1980), 168–184.
- [U] B. Ulrich, *Artin-Nagata properties and reduction of ideals*, Contemp. Math. **159**(1994), 373–400.
- [U1] B. Ulrich, *Remarks on Residual Intersections*, Free resolutions in Commutative Algebra and Algebraic Geometry, Eds. D. Eisenbud and C. Huneke, Research Notes in Mathematics **2** (1992), 133–138.
- [V] W. V. Vasconcelos, *Arithmetic of Blowup Algebras*, **195**, Cambridge University Press, 1994.
- [V2] W. V. Vasconcelos, *Integral Closure Rees Algebras, Multiplicities, Algorithms*, Springer, 2005.
- [W] C. A. Weibel, *An introduction to homological algebra*, **38**, Cambridge University Press, 1994.

¹ INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, BRAZIL.

E-mail address: `hamid@im.ufrj.br` and `hassanzadeh.ufrj@gmail.com`

² INSTITUTO DE MATEMÁTICA PURA E APLICADA, IMPA, RIO DE JANEIRO, BRAZIL.

E-mail address: `jnaeliton@yahoo.com.br`