

Periodic points in the intersection of immediate attracting basins boundaries

Bastien Rossetti

Abstract : We give sufficient conditions under which the set of eventually periodic points in the intersection of immediate attracting basins boundaries is non-empty. We give other conditions under which this set is dense in the intersection.

For a rational map f of degree $D \geq 2$ with two distinct attracting periodic points a_1 and a_2 , we wonder if $\partial B_1 \cap \partial B_2$ contains a periodic point, where B_i is the connected component of the complementary of the Julia set \mathcal{J} of f containing a_i . In this note we give conditions to a positive answer.

To state the theorem, let $\mathcal{P} := \{f^n(c) : f'(c) = 0 ; n \geq 1\}$ be the postcritical set of f and let \mathcal{P}_b be the set of $x \in \mathcal{P}$ such that there is no component of $\hat{\mathbb{C}} - \mathcal{J}$ whose boundary contains x .

Theorem Suppose that the B_i are simply connected, the boundaries ∂B_i are locally connected and that $\partial B_1 \cap \partial B_2 \neq \emptyset$.

1. If $x \in \partial B_1 \cap \partial B_2$ is multiply accessible from B_1 and if the orbit of x does not contain any critical point, then x is eventually periodic.
2. If $\partial B_1 \cap \partial B_2$ contains no critical point with infinite orbit and is disjoint from the ω -limit set of every recurrent critical point, then $\partial B_1 \cap \partial B_2$ contains a periodic point.
3. Assume furthermore that \mathcal{J} is connected. If $\partial B_1 \cap \partial B_2$ contains no accumulation point of \mathcal{P}_b and $\mathcal{P} \cap (\partial B_1 \cup \partial B_2)$, then the subset of eventually periodic points in $\partial B_1 \cap \partial B_2$ is non-empty and dense in $\partial B_1 \cap \partial B_2$.

As a particular case if f is postcritically finite then the set of eventually periodic points in $\partial B_1 \cap \partial B_2$ is non-empty (and dense in $\partial B_1 \cap \partial B_2$). Nevertheless the theorem does not require \mathcal{P} to be finite.

The hypothesis of part 2 are not useless for the existence of a periodic point in $\partial B_1 \cap \partial B_2$. Pick $\theta \in \mathbb{R}/\mathbb{Q}$ and consider $F_\theta(z) = \rho_\theta z^2(z-3)/(1-3z)$ with $\rho_\theta \in S^1$ such that F_θ has rotation number θ on S^1 . Such a map is studied in [Pet]. The map F_θ has two attracting points $a_1 = 0$ and $a_2 = \infty$, with $1 \in \partial B_1 \cap \partial B_2 \subset S^1$. Although the B_i are simply connected and the ∂B_i are locally connected there is no periodic point in $\partial B_1 \cap \partial B_2$, since $F_\theta|_{S^1}$ is topologically conjugate to $z \mapsto e^{2i\pi\theta}z$. Note that for these examples the point 1 is a critical point in $\partial B_1 \cap \partial B_2$ with an infinite orbit.

For the proof of the theorem we assume that the B_i are fixed, up to consider an iterate of f , and that a_i is the only critical point of f in B_i , up to make a quasi-conformal deformation (see [CG], theorem VI 5.1). Let $\phi_i : \mathbb{D} \rightarrow B_i$ conjugate f to z^{d_i} (where d_i is the local degree of f at a_i), and for $t \in \mathbb{R}$ let $R_i(t) := \phi_i(\{re^{2i\pi t} : 0 \leq r < 1\})$. Since ∂B_i is locally connected, ϕ_i extends continuously to $\overline{\phi_i} : \overline{\mathbb{D}} \rightarrow \overline{B_i}$. A point $y \in \partial B_i$ is *multiply accessible from B_i* if $\#((\overline{\phi_i})^{-1}(\{y\})) \geq 2$.

Proof of part 1. Let $\theta, \theta' \in (\overline{\phi_1})^{-1}(\{x\})$ be distinct. For any $n \geq 0$, $R_1(d_1^n \theta)$ and $R_1(d_1^n \theta')$ are distinct, for otherwise the orbit of x contains a critical point. The union $\overline{R_1(d_1^n \theta)} \cup \overline{R_1(d_1^n \theta')}$ is a

Jordan curve whose complementary has one connected component V_n such that $\overline{V_n} \cap \overline{B_2} = \{f^n(x)\}$. Let $I_n := (\overline{\phi_1})^{-1}(\partial B_1 \cap \overline{V_n})$. If there exist $M \neq N$ with $\{d_1^N \theta, d_1^N \theta'\} \cap I_M \neq \emptyset$, then $f^M(x) = f^N(x)$. Otherwise the I_k are disjoint intervals. This is impossible since $t \mapsto t^{d_1}$ is expanding on the unit circle, so that there are infinitely many I_n with $|I_n| \geq 1/d_1^2$. ■

For the proof of part 2 we show (under hypothesis) that $f|_{\partial B_1 \cap \partial B_2}$ is distance-expanding using a theorem of Mañé and we conclude with a result about periodic points for distance-expanding maps.

Theorem 0.1 ([Ma]) *Let g be a rational map of degree at least 2, and Λ be a compact forward invariant set in the Julia set containing no critical point nor parabolic point. If Λ is disjoint from the ω -limit set of every recurrent critical point, then there exists $N \geq 0$ such that $\min_{z \in \Lambda} \|(g^n)'(z)\| > 1$ for every $n \geq N$.*

Let (X, ρ) be a metric space. A continuous map $T : X \rightarrow X$ is *distance-expanding* if there exist $\lambda > 1$, $\eta > 0$ and $N \geq 0$ such that for any $x, y \in X$, if $\rho(x, y) \leq \eta$ then $\rho(T^N(x), T^N(y)) \geq \lambda \rho(x, y)$.

Theorem 0.2 ([PrUr], chapter 4) *Let (X, ρ) be a compact metric space. If $T : X \rightarrow X$ is continuous, open and distance-expanding, then there exists $\alpha > 0$ such that if $x \in X$ and $\rho(x, T^L(x)) \leq \alpha$ for some $L \geq 1$, then X contains a periodic point.*

Proof of part 2. Assume that $\partial B_1 \cap \partial B_2$ contains no critical point with finite orbit nor parabolic point, nor multiple point in ∂B_i (for otherwise $\partial B_1 \cap \partial B_2$ contains a periodic point). By theorem 0.1 there exists $N \geq 0$ such that $\min_{x \in \partial B_1 \cap \partial B_2} \|(f^N)'(x)\| > 1$. We show now that $f|_{\partial B_1 \cap \partial B_2}$ is distance-expanding and open, in order to apply theorem 0.2.

Lemma 0.3 *The restriction $f|_{\partial B_1 \cap \partial B_2}$ is distance-expanding with respect to the spherical metric.*

Proof. By continuity of $x \mapsto \|(f^N)'(x)\|$, there exist $\lambda > 1$ and a neighborhood U of $\partial B_1 \cap \partial B_2$ such that $\min_{x \in U} \|(f^N)'(x)\| \geq \lambda$. By compactness of $\partial B_1 \cap \partial B_2$ there exists $\eta > 0$ such that if $d(x, y) \leq \eta$ then the geodesic Γ between $f^N(x)$ and $f^N(y)$ lifts to a path γ from x to y with $\gamma \subset U$. Thus we get $d(f^N(x), f^N(y)) = \text{length}(\Gamma) \geq \lambda \cdot \text{length}(\gamma) \geq \lambda d(x, y)$. □

Lemma 0.4 *The restriction $f|_{\partial B_1 \cap \partial B_2}$ is open.*

Proof. Let $O \subset \partial B_1 \cap \partial B_2$ and suppose $f(O)$ is not open. There exists a sequence (y_n) in $(\partial B_1 \cap \partial B_2) - f(O)$ tending to some $y \in f(O)$. Let $x \in O$ be such that $f(x) = y$. There exist a neighborhood U of x and a neighborhood V of y such that $f : U \rightarrow V$ is a homeomorphism. Thus for n large enough the point $x_n = f^{-1}(y_n) \cap U$ is well defined, and $x_n \rightarrow x$. We show now that $x_n \in \partial B_1 \cap \partial B_2 - O$ and get that O is not open. It is clear that $x_n \notin O$ since $f(x_n) \notin f(O)$. For any n there exists a Fatou component B_i^n such that $f(B_i^n) = B_i$ and $x_n \in \partial B_1^n \cap \partial B_2^n$. For n large enough $B_i^n = B_i$. Otherwise, for some $i_0 \in \{1, 2\}$ there exists a Fatou component B such that $B \neq B_{i_0}$, $f(B) = B_{i_0}$ and $x \in \partial B$. The boundary ∂B has finitely many connected components, thus each one of them is locally connected. Let \tilde{B} be either B or B_{i_0} . There exists a connected component $U_{\tilde{B}}$ of $U \cap \tilde{B}$ such that $x \in \partial U_{\tilde{B}}$. Since $f(x)$ is simple in ∂B_{i_0} , there exists a unique connected component $V_{B_{i_0}}$ of $V \cap B_{i_0}$ such that $f(x) \in \partial V_{B_{i_0}}$. Hence $f(U_{\tilde{B}}) = V_{B_{i_0}}$. Since $B \neq B_{i_0}$, we have $U_{B_{i_0}} \cap U_B = \emptyset$ and $f(U_{B_{i_0}}) = f(U_B)$ which contradicts the injectivity of $f|_U$. □

Now we apply theorem 0.2. Let w be an accumulation point of the orbit of some point z in $\partial B_1 \cap \partial B_2$. There exist $P > Q \geq 0$ such that $f^P(z)$ and $f^Q(z)$ are both in $B(w; \alpha/2)$, hence $d(f^Q(z), f^P(z)) = d(f^Q(z), f^{P-Q}(f^Q(z))) \leq \alpha$, where α is the constant in theorem 0.2. Thus we get a periodic point in $\partial B_1 \cap \partial B_2$. ■

The proof of part 3 uses ideas and techniques developed by Kevin Pilgrim in his thesis ([Pil]). In case where f is postcritically finite and hyperbolic, the part 3 is a corollary of his work. Up to make a quasi-conformal deformation, we assume that all the critical points in the basin of a_i have a finite orbit. We also assume that $\#\mathcal{P} > 2$ otherwise f is conjugate to $z \mapsto z^{\pm D}$ and the conclusion follows.

For any $t, t' \in \mathbb{R}$, if $\overline{R_1(t)} \cap \overline{R_2(t')} \neq \emptyset$ then $\overline{R_1(t)} \cup \overline{R_2(t')}$ is called a *chord*. Denote χ the set of chords. If $\alpha \in \chi$ is periodic then the point $\alpha \cap \mathcal{J} \in \partial B_1 \cap \partial B_2$ is periodic. We equip χ with the Hausdorff distance d_H , which turns it into a compact metric space. The construction of a periodic chord will use the theory of isotopies. For any chord α and any set $X \subset \hat{\mathbb{C}}$, $[\alpha]_X$ will denote the isotopy class of α rel X . For any $\alpha, \beta \in \chi$ distinct the complement $\hat{\mathbb{C}} - (\alpha \cup \beta)$ has at least two connected components and at most three (with points of \mathcal{J} in each one of them). Thus for any $m \geq 0$, $[\alpha]_{f^{-m}(\mathcal{P})} = [\beta]_{f^{-m}(\mathcal{P})}$ if and only if one connected component of $\hat{\mathbb{C}} - (\alpha \cup \beta)$ contains all the points of $f^{-m}(\mathcal{P}) - \{a_1, a_2\}$.

(*) Let $GO(\mathcal{P}) := \{f^m(c) : f'(c) = 0, m \in \mathbb{Z}\}$. From the hypothesis the set $GO(\mathcal{P}) \cap (\partial B_1 \cap \partial B_2)$ is finite. We deduce that if a chord α satisfies $\alpha \cap \mathcal{J} \in GO(\mathcal{P})$ then $\alpha \cap \mathcal{J}$ is eventually periodic. We denote $\chi(S^2, GO(\mathcal{P})) := \{\alpha \in \chi : \alpha \cap \mathcal{J} \notin GO(\mathcal{P})\}$. We study the dynamics of the chords in $\chi(S^2, GO(\mathcal{P}))$.

The proof is as follows. For $\alpha \in \chi(S^2, GO(\mathcal{P}))$ if the sequence $([f^j(\alpha)]_{f^{-1}(\mathcal{P})})_{j \geq 0}$ is eventually cyclical then α is eventually periodic (lemma 0.7). Otherwise, remarking that $([f^j(\alpha)]_{\mathcal{P}})_{j \geq 0}$ contains twice the same element (lemma 0.8) we will build a sequence $(\beta_n) \subset \chi(S^2, GO(\mathcal{P}))$ by a series of adjustments (lemma 0.5) such that β_n converges (lemma 0.6) to a chord β with $\beta \cap \mathcal{J}$ eventually periodic. The density part will follow from the fact that we can build β as close as we want from α .

Let $\alpha \in \chi(S^2, GO(\mathcal{P}))$. A *lift* of α is the closure of a connected component of $f^{-1}(\alpha - \{a_1, a_2\})$. If a lift of α is a chord then it belongs to $\chi(S^2, GO(\mathcal{P}))$.

Lemma 0.5 *Let $N \geq 1$ and $\alpha, \beta_N \in \chi(S^2, GO(\mathcal{P}))$ be such that $\beta_N \in [f^N(\alpha)]_{\mathcal{P}}$. There exists a unique $\beta_0 \in \chi(S^2, GO(\mathcal{P})) \cap [\alpha]_{f^{-1}(\mathcal{P})}$, such that $f^N(\beta_0) = \beta_N$ and $[f^i(\beta_0)]_{f^{-1}(\mathcal{P})} = [f^i(\alpha)]_{f^{-1}(\mathcal{P})}$ for every $0 \leq i \leq N-1$. More precisely, $\beta_0 \in [\alpha]_{f^{-N}(\mathcal{P})}$.*

Proof. Since $f^{N-1}(\alpha)$ is a lift of α , there exists a unique lift β_{N-1} of β_N with $\beta_{N-1} \in [f^{N-1}(\alpha)]_{f^{-1}(\mathcal{P})}$. Moreover $\beta_{N-1} \in \chi(S^2, GO(\mathcal{P}))$. In particular, $\beta_{N-1} \in [f^{N-1}(\alpha)]_{\mathcal{P}}$. For each $1 \leq i \leq N$ we construct inductively a unique $\beta_{N-i} \in [f^{N-i}(\alpha)]_{f^{-1}(\mathcal{P})}$ with $f^i(\beta_{N-i}) = \beta_N$ and $f^k(\beta_{N-i}) \in [f^k(f^{N-i}(\alpha))]_{f^{-1}(\mathcal{P})}$ for any $0 \leq k \leq i-1$ (note that more precisely $\beta_{N-i} \in [f^{N-i}(\alpha)]_{f^{-i}(\mathcal{P})}$). Moreover $\beta_{N-i} \in \chi(S^2, GO(\mathcal{P}))$. ■

Lemma 0.6 $\forall \varepsilon > 0 \exists N \geq 0 : \forall \alpha, \beta \in \chi(S^2, GO(\mathcal{P}))$, if $[\alpha]_{f^{-N}(\mathcal{P})} = [\beta]_{f^{-N}(\mathcal{P})}$ then $d_H(\alpha, \beta) < \varepsilon$. As a consequence, if $[\alpha]_{f^{-n}(\mathcal{P})} = [\beta]_{f^{-n}(\mathcal{P})}$ for every $n \geq 0$ then $\alpha = \beta$.

Proof. Let $\varepsilon > 0$. There exists a constant $\eta > 0$ such that : for any $\alpha, \beta \in \chi(S^2, GO(\mathcal{P}))$, if $d_H(\alpha, \beta) \geq \varepsilon$ then in two of the connected components of $\hat{\mathbb{C}} - (\alpha \cup \beta)$ lie an open ball centered at a point of \mathcal{J} and with radius η (for otherwise we can construct a sequence $((\delta_n, \gamma_n)) \subset \chi(S^2, GO(\mathcal{P}))^2$ such that $d_H(\delta_n, \gamma_n) \geq \varepsilon$, and such that there are no two connected components of $\hat{\mathbb{C}} - (\delta_n \cup \gamma_n)$ in which lie an open ball centered at a point of \mathcal{J} with radius η_n , with $\eta_n \rightarrow 0$; taking an accumulation point (δ, γ) of $((\delta_n, \gamma_n))$, we have $d_H(\delta, \gamma) \geq \varepsilon$ and in two of the connected components of $\hat{\mathbb{C}} - (\delta \cup \gamma)$ lie an open ball centered at a point of \mathcal{J} and with fixed radius $\rho > 0$; we see that for N large enough in two of the connected components of $\hat{\mathbb{C}} - (\delta_N \cup \gamma_N)$ lie an open ball centered at a point of \mathcal{J} and with radius $\rho/2$; take N large enough so that $\eta_N < \rho/2$, this is a contradiction). Since $\#\mathcal{P} > 2$, there exists $N \geq 0$ (depending only on η) such that each one of these balls contains a point of $f^{-N}(\mathcal{P})$. Hence there is a point of $f^{-N}(\mathcal{P})$ in two connected components of $\hat{\mathbb{C}} - (\alpha \cup \beta)$, thus $[\alpha]_{f^{-N}(\mathcal{P})} \neq [\beta]_{f^{-N}(\mathcal{P})}$. ■

Lemma 0.7 For $\alpha \in \chi(S^2, GO(\mathcal{P}))$, if the sequence $([f^n(\alpha)]_{f^{-1}(\mathcal{P})})_{n=0}^\infty$ is cyclical then α is periodic.

Proof. Suppose that $[f^{n+Q}(\alpha)]_{f^{-1}(\mathcal{P})} = [f^n(\alpha)]_{f^{-1}(\mathcal{P})}$ for any $n \geq 0$. We apply lemma 0.5 to $f^{n+Q}(\alpha) \in [f^n(\alpha)]_{\mathcal{P}}$: there exists a chord $\beta_n \in [\alpha]_{f^{-1}(\mathcal{P})}$ such that $f^n(\beta_n) = f^{n+Q}(\alpha)$ and for all $0 \leq i \leq n-1$, $[f^i(\beta_n)]_{f^{-1}(\mathcal{P})} = [f^i(\alpha)]_{f^{-1}(\mathcal{P})}$. This chord is $f^Q(\alpha)$. Thanks to lemma 0.5 we also have $f^Q(\alpha) = \beta_n \in [\alpha]_{f^{-n}(\mathcal{P})}$. Since this is true for any $n \geq 0$, we conclude by lemma 0.6 that $f^Q(\alpha) = \alpha$. ■

Lemma 0.8 For any $\alpha \in \chi(S^2, GO(\mathcal{P}))$ there exist $M \neq N$ such that $[f^M(\alpha)]_{\mathcal{P}} = [f^N(\alpha)]_{\mathcal{P}}$.

Proof. Suppose that for any $m, n \geq 0$ distinct we have $[f^m(\alpha)]_{\mathcal{P}} \neq [f^n(\alpha)]_{\mathcal{P}}$. Since (χ, d_H) is compact, there exists a chord $\beta \in \chi$ on which accumulate a sub-sequence $(f^{n(k)}(\alpha))_{k=0}^\infty$. Since for any k, k' distinct at least two connected components of $\hat{\mathcal{C}} - (f^{n(k)}(\alpha) \cup f^{n(k')}(\alpha))$ contain a point of \mathcal{P} , one can construct a non-stationary sequence $(z_n) \subset \mathcal{P}$ which accumulates a point $z \in \beta$. Up to extraction, we have $(z_n) \subset \mathcal{P}_b$ or $(z_n) \subset \mathcal{P} \cap (\partial B_1 \cup \partial B_2)$. Indeed, assume that $(z_n) \cap \mathcal{P} - \mathcal{P}_b$ is infinite. Up to extraction assume $(z_n) \subset \mathcal{P} - \mathcal{P}_b$. By construction, there is an infinite subset of (z_n) whose elements are pairwise separated by chords. We deduce that $(z_n) \subset \mathcal{P} \cap (\overline{B_1} \cup \overline{B_2})$ since there exist finitely many Fatou components U_1, \dots, U_N such that $\mathcal{P} - \mathcal{P}_b \subset \overline{U_1} \cup \dots \cup \overline{U_N}$. Since we assume that a_i is the only point of \mathcal{P} in B_i , we conclude that $(z_n) \subset \mathcal{P} \cap (\partial B_1 \cup \partial B_2)$. In particular, $(z_n) \subset \mathcal{J}$ and $z = \lim_{n \rightarrow \infty} z_n = \beta \cap \mathcal{J} \in \partial B_1 \cap \partial B_2$. This contradicts the hypothesis of part 3. ■

Proof of part 3. Let α be a chord. We have three cases.

Case 1 : $\alpha \cap \mathcal{J} \in GO(\mathcal{P})$. Thus $\alpha \cap \mathcal{J}$ is eventually periodic, as explained in (*).

Case 2 : $\alpha \in \chi(S^2, GO(\mathcal{P}))$ and $([f^n(\alpha)]_{f^{-1}(\mathcal{P})})_{n=0}^\infty$ is eventually cyclical. Then α is eventually periodic by lemma 0.7, hence the point $\alpha \cap \mathcal{J}$ is eventually periodic.

Case 3 : $\alpha \in \chi(S^2, GO(\mathcal{P}))$ and $([f^n(\alpha)]_{f^{-1}(\mathcal{P})})_{n=0}^\infty$ is not eventually cyclical. We built from α a chord β fitting case 1 or 2, as follows.

By lemma 0.8 there exist $N \geq 0$ and $Q \geq 1$ such that $[f^{N+Q}(\alpha)]_{\mathcal{P}} = [f^N(\alpha)]_{\mathcal{P}}$. Set $\beta_0 := f^N(\alpha)$. We apply lemma 0.5 to $\beta_0 \in [f^Q(\beta_0)]_{\mathcal{P}}$: there exists a chord $\beta_1 \in [\beta_0]_{f^{-Q}(\mathcal{P})}$ such that

$$[f^i(\beta_1)]_{f^{-1}(\mathcal{P})} = [f^i(\beta_0)]_{f^{-1}(\mathcal{P})}$$

for any $0 \leq i \leq Q-1$, and $f^Q(\beta_1) = \beta_0$. Thus $[f^{kQ+i}(\beta_1)]_{f^{-1}(\mathcal{P})} = [f^i(\beta_0)]_{f^{-1}(\mathcal{P})}$ for any $k \in \{0, 1\}$ and any $0 \leq i \leq Q-1$, and $\beta_1 \in [f^{2Q}(\beta_1)]_{\mathcal{P}}$.

We build inductively a sequence of chords $(\beta_q)_{q=0}^\infty$ such that :

(i) $\beta_{q+n} \in [\beta_q]_{f^{-2^q Q}(\mathcal{P})}$ for any $n \geq 0$, and

(ii) $[f^{kQ+i}(\beta_q)]_{f^{-1}(\mathcal{P})} = [f^i(\beta_0)]_{f^{-1}(\mathcal{P})}$ for any $k \in \{0, \dots, 2^q - 1\}$ and $0 \leq i \leq Q-1$.

The sequence $(\beta_q)_{q=0}^\infty$ converges to a chord β whose point $\beta \cap \mathcal{J}$ is eventually periodic. Indeed, the convergence follows from (i) and lemma 0.6. The limit β is a chord since χ is compact. If β is not in $\chi(S^2, GO(\mathcal{P}))$ then β fits case 1. If $\beta \in \chi(S^2, GO(\mathcal{P}))$, then we get for the limit $[f^{kQ+i}(\beta)]_{f^{-1}(\mathcal{P})} = [f^i(\beta_0)]_{f^{-1}(\mathcal{P})}$ for every $k \in \mathbb{N}$ and $0 \leq i \leq Q-1$. Hence $([f^i(\beta)]_{f^{-1}(\mathcal{P})})_{i=0}^\infty$ is cyclical, and β fits case 2.

We have obtained a chord $\beta \in [f^N(\alpha)]_{f^{-Q}(\mathcal{P})}$ such that the point $\beta \cap \mathcal{J}$ is eventually periodic. We apply lemma 0.5 to $\beta \in [f^N(\alpha)]_{f^{-Q}(\mathcal{P})}$: there exists a chord $\gamma \in [\alpha]_{f^{-(N+Q)}(\mathcal{P})}$ such that $\gamma \cap \mathcal{J}$ is eventually periodic. Now we make N tend to ∞ by applying lemma 0.8 to $f^m(\alpha)$ for m as large as wished. Then we get a sequence of chords $(\gamma_n) \rightarrow \alpha$ with $\gamma_n \cap \mathcal{J}$ eventually periodic. By local connectivity of the ∂B_i we get that $\gamma_n \cap \mathcal{J}$ converges to $\alpha \cap \mathcal{J}$. This proves the density. ■

References

- [CG] L. CARLESON AND T. W. GAMELIN, *Complex Dynamics*. Second edition, Springer, 1995.
- [Ma] R. MAÑÉ , *On a theorem of Fatou*, Bol. Soc. Bras. Mat. vol. 24, p. 1-11 (1993)
- [Pet] C.L. PETERSEN , *Local connectivity of some Julia sets containing a circle with an irrational rotation*. Acta Math., 177 (1996), 163-224
- [Pil] K. PILGRIM , *Cylinders for iterated rational maps*, Thesis (1994)
- [PrUr] F. PRZYTYCKI AND M. URBAŃSKI , *Conformal Fractals : Ergodic Theory Methods*, The London Mathematical Society, Lecture Note Series 371 (2010)