

**FACTORIZATION IN $SL_n(R)$ WITH ELEMENTARY
MATRICES WHEN R IS THE DISK ALGEBRA AND THE
WIENER ALGEBRA**

AMOL SASANE

ABSTRACT. Let R be the polydisc algebra or the Wiener algebra. It is shown that the group $SL_n(R)$ is generated by the subgroup of elementary matrices with all diagonal entries 1 and at most one nonzero off-diagonal entry. The result an easy consequence of the deep result due to Ivarsson and Kutzschebauch [4].

1. INTRODUCTION

Let R be a commutative unital ring. Let I_n denote the $n \times n$ identity matrix, that is the square matrix with all diagonal entries equal to $1 \in R$ and off-diagonal entries equal to $0 \in R$. Recall that an elementary matrix $E_{ij}(\alpha)$ over R is a matrix of the form $I_n + \alpha \mathbf{e}_{ij}$, where $i \neq j$, $\alpha \in R$, and \mathbf{e}_{ij} is the $n \times n$ matrix whose entry in the i th row and j th column is 1 and all other entries are zeros. Let $SL_n(R)$ be the group of all $n \times n$ matrices whose entries are elements of R and whose determinant is 1. Let $E_n(R)$ be the subgroup of $SL_n(R)$ generated by the elementary matrices.

A classical question in commutative algebra is the following:

Question 1.1. Is $SL_n(R)$ equal to $E_n(R)$?

The answer to this question depends on the ring R , and here is a list of a few known results.

- (1) If $R = \mathbb{C}$, then the answer is “Yes”, and this is standard exercise in linear algebra; see for example [1, Exercise 18.(c), page 71].
- (2) Let R be the polynomial ring $\mathbb{C}[z_1, \dots, z_n]$ in the indeterminates z_1, \dots, z_n with complex coefficients.

If $n = 1$, then the answer is “Yes”, and this follows from the Euclidean Division Algorithm in $\mathbb{C}[z]$.

If $n = 2$, then the answer is “No”, and [2] gave the following counterexample:

$$\begin{bmatrix} 1 + z_1 z_2 & z_1^2 \\ -z_2^2 & 1 - z_1 z_2 \end{bmatrix} \in SL_2(\mathbb{C}[z_1, z_2]) \setminus E_2(\mathbb{C}[z_1, z_2]).$$

1991 *Mathematics Subject Classification.* Primary 46J10; Secondary 15A23, 15A54.

Key words and phrases. Gromov’s Vasertein Problem, Banach algebras, polydisc algebra, Wiener algebra.

For $n \geq 3$, the answer is “Yes”, and this is the K_1 -analogue of Serre’s Conjecture, which is the Suslin Stability Theorem [5].

- (3) The case of R being a ring of continuous functions was considered in [6]. Let $C(X; \mathbb{C})$ be the ring of continuous complex-valued functions on the finite-dimensional normal topological space X with pointwise operations. $C_b(X; \mathbb{C})$ denotes the subring of $C(X; \mathbb{C})$ consisting of *bounded* functions. It was shown in [6] that for $R = C(X; \mathbb{C})$ or $C_b(X; \mathbb{C})$, the answer is “Yes” if there is no homotopy obstruction. Indeed, if E is an elementary matrix, then $(X \ni) x \mapsto E(x) \in SL_n(\mathbb{C})$ is null-homotopic (to the constant map $x \mapsto I_n : X \rightarrow SL_n(\mathbb{C})$). So it follows that if $\pi(F)$ denotes the homotopy class of the map $x \mapsto F(x) : X \rightarrow SL_n(\mathbb{C})$ corresponding to $F \in SL_n(R)$, then a necessary condition for $F \in E_n(C(X; \mathbb{C}))$ is that $\pi(F) = 0$. It turns out that this condition is also sufficient, and this is the content of [6, Theorem 4].
- (4) Based on the above result, it is natural to consider the question also for the ring $\mathcal{O}(X)$ of holomorphic functions on Stein spaces in \mathbb{C}^n . This was posed as an explicit open problem by Gromov in [3], and was recently solved by Ivarsson and Kutzschebauch [4]. The main result in [4] is the following:

Theorem 1.2 ([4]). *If X is a finite-dimensional reduced Stein space and $F : X \rightarrow SL_n(\mathbb{C})$ is a holomorphic mapping that is null-homotopic, then there exists a natural number K and holomorphic mappings $G_1, \dots, G_K : X \rightarrow \mathbb{C}^{m(m-1)/2}$ such that F can be written as a product of upper and lower diagonal unipotent matrices*

$$F(x) = M_1(G_1(x)) \cdots M_K(G_K(x)), \quad x \in X,$$

where the matrices $M_j(G_j(x))$ are defined by

$$M_j(G_j(x)) := \begin{bmatrix} 1 & & 0 \\ & \ddots & \\ G_j(x) & & 1 \end{bmatrix} \quad \text{if } j \text{ is odd,}$$

while

$$M_j(G_j(x)) := \begin{bmatrix} 1 & & G_j(x) \\ & \ddots & \\ 0 & & 1 \end{bmatrix} \quad \text{if } j \text{ is even.}$$

In particular, the assumption of null-homotopy is always satisfied if X is contractible.

We wish to consider Question 1.1 for commutative, semisimple, unital complex Banach algebras R . A special case is when $R = C_b(X; \mathbb{R})$, where X is a compact Hausdorff topological space, and item (3) above describes the answer in this special case. Motivated by this, we formulate the following question/conjecture, but first we introduce some convenient notation.

Let R be a commutative, semisimple, unital complex Banach algebra with maximal ideal space denoted by X_R , equipped with the weak-* topology induced from the dual space $R^* := \mathcal{L}(R; \mathbb{C})$ of R .

Let $\widehat{\cdot} : R \rightarrow C(X_R; \mathbb{C})$ denote the Gelfand transform. For $F \in SL_n(R)$, let \widehat{F} be the matrix with elements in $C(X_R; \mathbb{C})$ obtained by taking the Gelfand transform of the entries of F , and $\pi(\widehat{F})$ denotes the homotopy class of $\varphi \mapsto \widehat{F}(\varphi) : X_R \rightarrow SL_n(\mathbb{C})$.

Conjecture 1.3. Let R be a commutative, semisimple, unital complex Banach algebra. $F \in SL_n(R)$ belongs to $E_n(R)$ if and only if $\pi(\widehat{F}) = 0$.

We consider Question 1.1 for two important Banach algebras of holomorphic functions: the polydisc algebra $A(\overline{\mathbb{D}}^n)$ and the Wiener algebra $W^+(\overline{\mathbb{D}}^n)$.

Let $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ and $\overline{\mathbb{D}} := \{z \in \mathbb{C} : |z| \leq 1\}$. Let $d \in \mathbb{N}$. The Wiener algebra $W^+(\overline{\mathbb{D}}^n)$ is the Banach algebra defined by

$$W^+(\overline{\mathbb{D}}^d) = \left\{ \sum_{k_1=0}^{\infty} \cdots \sum_{k_d=0}^{\infty} a_{(k_1, \dots, k_d)} z_1^{k_1} \cdots z_d^{k_d} : \sum_{k_1=0}^{\infty} \cdots \sum_{k_d=0}^{\infty} |a_{(k_1, \dots, k_d)}| < \infty \right\},$$

with pointwise addition and multiplication, and the $\|\cdot\|_1$ -norm given by

$$\|f\|_1 = \sum_{k_1=0}^{\infty} \cdots \sum_{k_d=0}^{\infty} |a_{(k_1, \dots, k_d)}|, \quad f = \sum_{k_1=0}^{\infty} \cdots \sum_{k_d=0}^{\infty} a_{(k_1, \dots, k_d)} z_1^{k_1} \cdots z_d^{k_d}.$$

The polydisc algebra $A(\overline{\mathbb{D}}^d)$ is the Banach algebra of all continuous functions $f : \overline{\mathbb{D}}^d \rightarrow \mathbb{C}$ which are holomorphic in \mathbb{D}^d , with pointwise addition and multiplication, and the supremum norm $\|\cdot\|_{\infty}$ given by

$$\|f\|_{\infty} := \sup_{(z_1, \dots, z_d) \in \overline{\mathbb{D}}^d} |f(z_1, \dots, z_d)|, \quad f \in A(\overline{\mathbb{D}}^d).$$

The ball algebra $A(\overline{\mathbb{B}}_d)$ is defined similarly, with the polydisc $\overline{\mathbb{D}}^d$ replaced by the ball

$$\overline{\mathbb{B}}_d := \{(z_1, \dots, z_d) \in \mathbb{C}^d : |z_1|^2 + \cdots + |z_d|^2 \leq 1\}.$$

For a $n \times n$ matrix F with entries in $A(\overline{\mathbb{D}}^d)$, $A(\overline{\mathbb{B}}_d)$ or $W^+(\overline{\mathbb{D}}^d)$, we define

$$\|F\| := \sum_{i,j=1}^n \|F_{ij}\|_{\infty},$$

where F_{ij} denotes the entry in the i th row and j th column of F . Then $\|FG\| \leq \|F\| \|G\|$, for $n \times n$ matrices F, G with entries from any of the Banach algebras $A(\overline{\mathbb{D}}^d)$, $A(\overline{\mathbb{B}}_d)$ or $W^+(\overline{\mathbb{D}}^d)$.

Our main result is the following.

Theorem 1.4. *If $R = A(\overline{\mathbb{D}}^d)$, $A(\overline{\mathbb{B}}_d)$ or $W^+(\overline{\mathbb{D}}^d)$, then $SL_n(R) = E_n(R)$.*

If $R = A(\overline{\mathbb{D}^d})$ or $W^+(\overline{\mathbb{D}^d})$, then in both cases, the maximal ideal space X_R can be identified with $\overline{\mathbb{D}^d}$ as a topological space. Similarly, $X_{A(\overline{\mathbb{B}_d})} = \overline{\mathbb{B}_d}$. If Conjecture 1.3 is true, then Theorem 1.4 follows from the observation that $\overline{\mathbb{D}^d}, \overline{\mathbb{B}_d}$ are contractible (since then $\pi(\widehat{F})$ is always trivial).

We will derive our main result as a consequence of the result from [4] quoted above, and [6, Lemma 9] reproduced below.

Lemma 1.5 ([6]). *Let R be a commutative topological unital ring such that the set of invertible elements of R is open in R . If $F \in SL_n(R)$ is sufficiently close to I_n , then F belongs to $E_n(R)$.*

2. PROOF OF THEOREM 1.4

Proof. We will simply prove the result in the case of the disc algebra $A(\overline{\mathbb{D}^d})$; the proofs in the cases of the ball algebra $A(\overline{\mathbb{B}_d})$ and the Wiener algebra being analogous.

Let $F \in SL_n(A(\overline{\mathbb{D}^d}))$. Let $r \in (0, 1)$ (to be determined later). Define

$$F_r(z_1, \dots, z_d) := F(rz_1, \dots, rz_d), \quad (z_1, \dots, z_d) \in \mathbb{D}^d.$$

As $F_r \in \mathcal{O}(\frac{1}{r}\mathbb{D}^d)$, and $\det F_r \equiv 1$, it follows from Theorem 1.2 (since $\frac{1}{r}\mathbb{D}^d$ is a contractible Stein domain) that there are elementary matrices G_1, \dots, G_K belonging to $E_n(\mathcal{O}(\frac{1}{r}\mathbb{D}^d))$ such that

$$F_r = E_1 \cdots E_K \in E_n(\mathcal{O}(\frac{1}{r}\mathbb{D}^d)) \subset E_n(A(\overline{\mathbb{D}^d})).$$

Thus $F(I_n + F^{-1}(F_r - F)) = F_r \in E_n(A(\overline{\mathbb{D}^d}))$. As $\det F = \det F_r = 1$, it follows that also $\det(I_n + F^{-1}(F_r - F)) = 1$. We will be done if we manage to show that $I_n + F^{-1}(F_r - F) \in E_n(A(\overline{\mathbb{D}^d}))$ too. But this is clear by Lemma 1.5, since

$$\left\| \left(I_n + F^{-1}(F_r - F) \right) - I_n \right\| = \|F^{-1}(F_r - F)\| \leq \|F^{-1}\| \|F_r - F\|,$$

and we can make $\|F_r - F\|$ as small as we like by choosing r close enough to 1. \square

Remark 2.1. The above proof also works for some other Banach algebras of smooth functions contained in the polydisc algebra, for example, if $N \in \mathbb{N}$, the Banach algebra $\partial^{-N}A(\overline{\mathbb{D}^d})$ of all functions $f \in A(\overline{\mathbb{D}^d})$ whose complex partial derivatives of all orders up to N belong to $A(\overline{\mathbb{D}^d})$, with the norm

$$\|f\|_{\partial^{-N}A(\overline{\mathbb{D}^d})} := \sum_{\alpha_1 + \dots + \alpha_d \leq N} \frac{1}{\alpha_1! \cdots \alpha_d!} \sup_{(z_1, \dots, z_d) \in \mathbb{D}^d} \left| \frac{\partial^{\alpha_1 + \dots + \alpha_d} f}{\partial z_1^{\alpha_1} \cdots \partial z_d^{\alpha_d}}(z_1, \dots, z_d) \right|.$$

In light of Theorem 1.4, it is natural to ask the analogous question also for the Hardy algebra. Recall that if U is an open set in \mathbb{C}^d , then the Hardy algebra $H^\infty(U)$ is the Banach algebra of all complex-valued functions on U that are bounded and holomorphic in U .

Conjecture 2.2. $SL_n(H^\infty(U)) = E_n(H^\infty(U))$ if U is the polydisc \mathbb{D}^d or open unit ball $U = \mathbb{B}_d := \{(z_1 \cdots, z_d) \in \mathbb{C}^d : |z_1|^2 + \cdots + |z_d|^2 < 1\}$.

REFERENCES

- [1] Michael Artin. *Algebra*. Prentice Hall, Englewood Cliffs, NJ, 1991.
- [2] Paul M. Cohn. On the structure of the GL_2 of a ring. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, No. 30, 5-53, 1966.
- [3] Michael Gromov. Oka's principle for holomorphic sections of elliptic bundles. *Journal of the American Mathematical Society*, 2:851-897, no. 4, 1989.
- [4] Björn Ivarsson and Frank Kutzschebauch. Holomorphic factorization of mappings into $SL_n(\mathbb{C})$. *Annals of Mathematics* (2), 175:45-69, no. 1, 2012.
- [5] Andrei A. Suslin. The structure of the special linear group over rings of polynomials. *Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya*, 41:235-252, no. 2, 1977.
- [6] Leonid N. Vaserstein. Reduction of a matrix depending on parameters to a diagonal form by addition operations. *Proceedings of the American Mathematical Society*, 103:741-746, no. 3, 1988.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, LUND UNIVERSITY, SWEDEN.
E-mail address: amol.sasane@math.lu.se