

BV-differential on Hochschild cohomology of Frobenius algebras.

Y. V. Volkov*

Abstract

For a finite-dimensional Frobenius k -algebra R with the Nakayama automorphism ν we define an algebra $\mathrm{HH}^*(R)^{\nu\uparrow}$. If the order of ν is not divisible by the characteristic of k , this algebra is isomorphic to the Hochschild cohomology algebra of R . We prove that this algebra is a BV-algebra. We use this fact to calculate the Gerstenhaber algebra structure and BV-structure on the Hochschild cohomology algebras of a family of self-injective algebras of tree type D_n .

1 Introduction

Hochschild cohomology is a subtle invariant of an associative algebra which carries a lot of information about its structure. The cohomology theory of associative algebras was introduced by Hochschild. It was later shown in [1], that the Hochschild cohomology algebra is a Gerstenhaber algebra. Sometimes we can define a BV-differential on the Hochschild cohomology algebra in such way that the structure of the Gerstenhaber algebra can be defined using this differential. In [2] Tradler shows that such differential can be defined for symmetric algebras. It is proved in [3] that a BV-differential can be constructed if the given algebra is a Frobenius algebra with finite stable Calabi-Yau dimension and periodic Hochschild cohomology. Moreover it is shown in [4] that a BV-differential exists in the case of Calabi-Yau algebra.

Let R be a Frobenius algebra over an algebraically closed field. Let ν be its Nakayama automorphism. In this paper we define an algebra $\mathrm{HH}^*(R)^{\nu\uparrow}$. It turns out that $\mathrm{HH}^*(R)^{\nu\uparrow} \simeq \mathrm{HH}^*(R)$ in many cases. The Lie bracket on the algebra $\mathrm{HH}^*(R)$ induces a bracket on $\mathrm{HH}^*(R)^{\nu\uparrow}$. We slightly modify the Tradler's proof for symmetric algebras to construct the BV-differential on the algebra $\mathrm{HH}^*(R)^{\nu\uparrow}$.

In section 5 we apply the obtained results to describe the Gerstenhaber algebra structure and the BV-differential for a family of self-injective algebras of tree type D_n . The Hochschild cohomology ring for these algebras was described in terms of generators and relations in [5]. Note that the calculation of these structures is a difficult task. There are only a few examples of such calculations. The structure of Gerstenhaber algebra was described for serial self-injective algebras in [6]. The calculus structure which includes the Gerstenhaber algebra structure was calculated for preprojective algebras of type A_n , D_n , E_n and L_n in [7] and [8]. Furthermore the Gerstenhaber algebra structure and the BV-differential for the group algebra of the quaternion group of order 8 over a field of characteristic 2 were calculated in [9].

*The author was supported by RFBR (13-01-00902 A and 14-01-31084 mol_a).

2 Basic definitions and constructions

Throughout the paper we suppose that k is an algebraically closed field, R is a finite-dimensional k -algebra, $\Lambda = R \otimes R^{\text{op}}$ is the enveloping algebra of R (we write \otimes instead of \otimes_k). The bar-resolution

$$\text{Bar}_*(R) : (R \xleftarrow{\mu} R^{\otimes 2} \xleftarrow{d_0} R^{\otimes 3} \leftarrow \dots \leftarrow R^{\otimes(n+2)} \xleftarrow{d_n} R^{\otimes(n+3)} \leftarrow \dots$$

of the algebra R is defined in the following way: $\text{Bar}_n(R) = R^{\otimes(n+2)}$ ($n \geq 0$), $\mu : R \otimes R \rightarrow R$ is the multiplication of R and d_n ($n \geq 0$) is defined by the formula

$$d_n(a_0 \otimes \dots \otimes a_{n+2}) = \sum_{i=0}^{n+1} (-1)^i a_0 \otimes \dots \otimes a_{i-1} \otimes a_i a_{i+1} \otimes a_{i+2} \otimes \dots \otimes a_{n+2},$$

where $a_i \in R$ ($0 \leq i \leq n+2$). The homology of the complex $C^*(R) = \text{Hom}_\Lambda(\text{Bar}_*(R), R)$ is called Hochschild cohomology of the algebra R . Note that $C^0(R) \simeq R$ and $C^n(R) = \text{Hom}_\Lambda(R^{\otimes(n+2)}, R) \simeq \text{Hom}_k(R^{\otimes n}, R)$. Let introduce notation

$$\delta_n^i(f)(a_1 \otimes \dots \otimes a_{n+1}) := \begin{cases} a_1 f(a_2 \otimes \dots \otimes a_{n+1}), & \text{if } i = 0, \\ (-1)^i f(a_1 \otimes \dots \otimes a_i a_{i+1} \otimes \dots \otimes a_{n+1}), & \text{if } 1 \leq i \leq n, \\ (-1)^{n+1} f(a_1 \otimes \dots \otimes a_n) a_{n+1}, & \text{if } i = n+1. \end{cases}$$

for $f \in C^n(R)$. Then $C^*(R)$ has the form

$$R \xrightarrow{\delta_0} \text{Hom}_k(R, R) \rightarrow \dots \rightarrow \text{Hom}_k(R^{\otimes n}, R) \xrightarrow{\delta_n} \text{Hom}_k(R^{\otimes(n+1)}, R) \rightarrow \dots,$$

where $\delta_n = \sum_{i=0}^{n+1} \delta_n^i$.

The cup product $f \smile g \in C^{n+m}(R) = \text{Hom}_k(R^{\otimes(n+m)}, R)$ for $f \in C^n(R)$ and $g \in C^m(R)$ is given by

$$(f \smile g)(a_1 \otimes \dots \otimes a_{n+m}) := f(a_1 \otimes \dots \otimes a_n) \cdot g(a_{n+1} \otimes \dots \otimes a_{n+m}).$$

This cup product induces a well-defined product on Hochschild cohomology

$$\smile : \text{HH}^n(R) \times \text{HH}^m(R) \longrightarrow \text{HH}^{n+m}(R)$$

which turns the graded k -vector space $\text{HH}^*(R) = \bigoplus_{n \geq 0} \text{HH}^n(R)$ into a graded commutative algebra ([1, Corollary 1]).

Let now define the Lie bracket. Let $f \in C^n(R)$, $g \in C^m(R)$. If $n, m \geq 1$, then for $1 \leq i \leq n$, we set

$$(f \circ_i g)(a_1 \otimes \dots \otimes a_{n+m-1}) := f(a_1 \otimes \dots \otimes a_{i-1} \otimes g(a_i \otimes \dots \otimes a_{i+m-1}) \otimes a_{i+m} \otimes \dots \otimes a_{n+m-1});$$

if $n \geq 1$ and $m = 0$, then $g \in R$ and for $1 \leq i \leq n$, we set

$$(f \circ_i g)(a_1 \otimes \dots \otimes a_{n-1}) := f(a_1 \otimes \dots \otimes a_{i-1} \otimes g \otimes a_i \otimes \dots \otimes a_{n-1}).$$

Now define

$$f \circ g := \sum_{i=1}^n (-1)^{(m-1)(i-1)} f \circ_i g$$

and

$$[f, g] := f \circ g - (-1)^{(n-1)(m-1)} g \circ f.$$

Note that $[f, g] \in C^{n+m-1}(R)$. Then $[,]$ induces a well-defined Lie bracket on Hochschild cohomology

$$[,] : \mathrm{HH}^n(R) \times \mathrm{HH}^m(R) \longrightarrow \mathrm{HH}^{n+m-1}(R)$$

such that $(\mathrm{HH}^*(R), \smile, [,])$ is a Gerstenhaber algebra ([1]).

A Batalin–Vilkovisky algebra (BV-algebra for short) is a Gerstenhaber algebra $(A^\bullet, \smile, [,])$ together with an operator $\Delta : A^\bullet \rightarrow A^{\bullet-1}$ of degree -1 such that $\Delta \circ \Delta = 0$ and

$$[a, b] = -(-1)^{(|a|-1)|b|} (\Delta(a \smile b) - \Delta(a) \smile b - (-1)^{|a|} a \smile \Delta(b)) \quad (2.1)$$

for homogeneous elements $a, b \in A^\bullet$. The following Theorem is proved in [2].

Theorem 1. [2, Theorem 1] *Let R be a symmetric algebra, i.e. an algebra with a nondegenerate associative symmetric bilinear form $\langle , \rangle : R \times R \rightarrow k$. For $f \in C^n(R) = \mathrm{Hom}_k(R^{\otimes n}, R)$ define $\Delta(f) \in C^{n-1}(R) = \mathrm{Hom}_k(R^{\otimes(n-1)}, R)$ by the formula*

$$\begin{aligned} & \langle \Delta(f)(a_1 \otimes \cdots \otimes a_{n-1}), a_n \rangle \\ &= \sum_{i=1}^n (-1)^{i(n-1)} \langle f(a_i \otimes \cdots \otimes a_n \otimes a_1 \otimes \cdots \otimes a_{i-1}), 1 \rangle, \end{aligned}$$

where $a_i \in R$ ($1 \leq i \leq n$). The map Δ induces the differential $\Delta : \mathrm{HH}^n(R) \rightarrow \mathrm{HH}^{n-1}(R)$. Then $(\mathrm{HH}^*(R), \smile, [,], \Delta)$ is a BV-algebra.

If $\sigma : R \rightarrow R$ is an automorphism of the algebra R , then we can define a map $\phi_\sigma : C^n(R) \rightarrow C^n(R)$ by the formula

$$(\phi_\sigma(f))(a_1 \otimes \cdots \otimes a_n) = \sigma^{-1}(f(\sigma(a_1) \otimes \cdots \otimes \sigma(a_n))),$$

where $f \in C^n(R) = \mathrm{Hom}_k(R^{\otimes n}, R)$, $a_i \in R$ ($1 \leq i \leq n$). From here on we write f^σ instead of $\phi_\sigma(f)$. It is easy to show that $(\delta_n f)^\sigma = \delta_n f^\sigma$ for $f \in C^n(R)$. Consequently, the map ϕ induces a map on Hochschild cohomology $()^\sigma : \mathrm{HH}^n(R) \rightarrow \mathrm{HH}^n(R)$. For a map $\sigma : X \rightarrow X$ we denote by X^σ the set $\{x \in X \mid \sigma(x) = x\}$. Then $\delta_n : C^n(R) \rightarrow C^{n+1}(R)$ induces the map $\delta_n^\sigma : C^n(R)^\sigma \rightarrow C^{n+1}(R)^\sigma$. We denote by $\mathrm{HH}^n(R)^{\sigma\uparrow}$ the homology of the complex $(C^n(R)^\sigma, \delta_n^\sigma)$ and define

$$\mathrm{HH}^*(R)^{\sigma\uparrow} := \bigoplus_{n \geq 0} \mathrm{HH}^n(R)^{\sigma\uparrow} = \bigoplus_{n \geq 0} \mathrm{Ker}(\delta_n^\sigma) / \mathrm{Im}(\delta_{n-1}^\sigma).$$

It is proved in [10] that $\smile : C^n(R) \times C^m(R) \longrightarrow C^{n+m}(R)$ defined above determines an algebra structure on $\mathrm{HH}^*(R)^{\sigma\uparrow}$ and the inclusion of $C^n(R)^\sigma$ into $C^n(R)$ induces an algebra homomorphism

$$\Theta_R^\sigma : \mathrm{HH}^*(R)^{\sigma\uparrow} \rightarrow \mathrm{HH}^*(R)^\sigma.$$

Moreover, Θ_R^σ is bijective if $\text{ord}(\sigma) < \infty$ and $\text{char } k \nmid \text{ord}(\sigma)$. The fact that $[\cdot, \cdot] : C^n(R) \times C^m(R) \longrightarrow C^{n+m-1}(R)$ induces a Gerstenhaber algebra structure on $\text{HH}^*(R)^{\sigma^\uparrow}$ can be proved analogously to the fact that $[\cdot, \cdot]$ induces a Gerstenhaber algebra structure on $\text{HH}^*(R)$ (cf. [1]). It is easy to see that Θ_R^σ is a homomorphism of Gerstenhaber algebras.

If $d : M \rightarrow N$ is a morphism of R -bimodules and σ_1, σ_2 are automorphisms of the algebra R , then we denote by $d_{(\sigma_1, \sigma_2)} : {}_{\sigma_1}M_{\sigma_2} \rightarrow {}_{\sigma_1}N_{\sigma_2}$ the morphism of R -bimodules, which is determined by the formula $d_{(\sigma_1, \sigma_2)}(m) = d(m)$ for $m \in M$ (we denote by ${}_{\sigma_1}M_{\sigma_2}$ the bimodule which is equal to M as k -linear space with multiplication $*$ defined by the equality $a * m * b = \sigma_1(a)m\sigma_2(b)$ for $m \in M, a, b \in R$).

Algebra R is called a Frobenius algebra if there is a linear map $\epsilon : R \rightarrow k$ such that the bilinear form $\langle a, b \rangle = \epsilon(ab)$ is nondegenerated. The Nakayama automorphism $\nu : R \rightarrow R$ is the automorphism which satisfies the equation $\langle a, b \rangle = \langle b, \nu(a) \rangle$ for all $a, b \in R$. From here on we assume that R is a Frobenius algebra, $\langle \cdot, \cdot \rangle$ is the corresponding bilinear form and ν is the Nakayama automorphism defined by it.

3 The generalization of Tradler's Theorem

Let $f \in C^n(R)$, $n \geq 1$. Define $\Delta_i f \in C^{n-1}(R)$ by the equation

$$\begin{aligned} & \langle \Delta_i f(a_1 \otimes \cdots \otimes a_{n-1}), a_n \rangle \\ &= \langle f(a_i \otimes \cdots \otimes a_{n-1} \otimes a_n \otimes \nu a_1 \otimes \cdots \otimes \nu a_{i-1}), 1 \rangle, \end{aligned}$$

where $a_i \in R$ ($1 \leq i \leq n$). Further define

$$\Delta := \sum_{i=1}^n (-1)^{i(n-1)} \Delta_i : C^n(R) \rightarrow C^{n-1}(R). \quad (3.1)$$

Lemma 1. $\delta_{n-1}(\Delta f) + \Delta \delta_n(f) = f^\nu - f \quad \forall f \in C^n(R)$.

Proof. Let $f \in C^n(R)$, $a_i \in R$ ($1 \leq i \leq n+1$). Set

$$A_i := a_i \otimes \cdots \otimes a_{n+1} \otimes \nu a_1 \otimes \cdots \otimes \nu a_{i-1} \in R^{\otimes(n+1)}.$$

Direct calculations show that

$$\begin{aligned} & \langle (-1)^{i(n-1)} \delta_{n-1}^j(\Delta_i f)(a_1 \otimes \cdots \otimes a_n), a_{n+1} \rangle \\ &= \begin{cases} \langle (-1)^{(i+1)n+1} \delta_n^{n-i+j+1}(f)(A_{i+1}), 1 \rangle, & \text{if } 0 \leq j \leq i-1, \\ \langle (-1)^{in+1} \delta_n^{j-i+1}(f)(A_i), 1 \rangle, & \text{if } i \leq j \leq n. \end{cases} \end{aligned}$$

Adding these equalities for $0 \leq j \leq n$ and $1 \leq i \leq n$, we obtain

$$\begin{aligned} & \langle \delta_{n-1}(\Delta f)(a_1 \otimes \cdots \otimes a_n), a_{n+1} \rangle + \langle \Delta \delta_n(f)(a_1 \otimes \cdots \otimes a_n), a_{n+1} \rangle \\ &= \left\langle \sum_{i=1}^{n+1} (-1)^{in+1} \sum_{j=1}^n \delta_n^j(f)(A_i), 1 \right\rangle + \left\langle \sum_{i=1}^{n+1} (-1)^{in} \delta_n(f)(A_i), 1 \right\rangle \\ &= \left\langle \sum_{i=1}^{n+1} (-1)^{in} (\delta_n^0(f)(A_i) + \delta_n^{n+1}(f)(A_i)), 1 \right\rangle. \end{aligned}$$

Since $\langle \delta_n^0(f)(A_i) + (-1)^n \delta_n^{n+1}(f)(A_{i+1}), 1 \rangle = 0$ for $1 \leq i \leq n$, we have

$$\begin{aligned} & \langle \delta_{n-1}(\Delta f)(a_1 \otimes \cdots \otimes a_n), a_{n+1} \rangle + \langle \Delta \delta_n(f)(a_1 \otimes \cdots \otimes a_n), a_{n+1} \rangle \\ &= \langle (-1)^n \delta_n^{n+1}(f)(A_1) + \delta_n^0(f)(A_{n+1}), 1 \rangle \\ &= \langle a_{n+1} f(\nu a_1 \otimes \cdots \otimes \nu a_n), 1 \rangle - \langle f(a_1 \otimes \cdots \otimes a_n) a_{n+1}, 1 \rangle \\ &= \langle (f^\nu - f)(a_1 \otimes \cdots \otimes a_n), a_{n+1} \rangle. \end{aligned}$$

□

The next two corollaries follow directly from Lemma 1.

Corollary 1. *The map Δ defined by the formula (3.1) induces a map*

$$\Delta : \mathrm{HH}^n(R)^{\nu\uparrow} \rightarrow \mathrm{HH}^{n-1}(R)^{\nu\uparrow}.$$

Corollary 2. $\mathrm{HH}^*(R)^\nu = \mathrm{HH}^*(R)$.

For a special case, the second corollary is proved in [11]. Now we are able to formulate the following generalization of Theorem 1.

Theorem 2. *Let R be a Frobenius algebra with bilinear form $\langle \cdot, \cdot \rangle$ and Nakayama automorphism ν . Then Δ defined by (3.1) induces a BV-algebra structure on the Gerstenhaber algebra $(\mathrm{HH}^*(R)^{\nu\uparrow}, \smile, [\cdot, \cdot])$.*

Proof. Let us prove that $\Delta \circ \Delta = 0$ in $\mathrm{HH}^*(R)^{\nu\uparrow}$. It is enough to prove that any element of $\mathrm{HH}^n(R)^{\nu\uparrow}$ can be represented by an element $f \in \mathrm{Ker} \delta_n^\nu$ such that $f(a_1 \otimes \cdots \otimes a_n) = 0$ if $a_i = 1$ for some $1 \leq i \leq n$. Let us define the maps $s_n^i : C^{n+1}(R) \rightarrow C^n(R)$ ($0 \leq i \leq n$) by the formula

$$(s_n^i(g))(a_1 \otimes \cdots \otimes a_n) = (-1)^i g(a_1 \otimes \cdots \otimes a_i \otimes 1 \otimes a_{i+1} \cdots \otimes a_n)$$

for $g \in C^{n+1}(R)$. It is clear that s_n^i induces a map $s_n^{i,\nu} : C^{n+1}(R)^\nu \rightarrow C^n(R)^\nu$. Now we can prove by induction on N that any element $f' \in \mathrm{HH}^n(R)^{\nu\uparrow}$ can be represented by an element $f \in \mathrm{Ker} \delta_n^\nu$ such that $f(a_1 \otimes \cdots \otimes a_n) = 0$ if $a_i = 1$ for some $1 \leq i \leq N$. Indeed, if $N = 0$, then the required assertion is obvious. Now assume that the cohomology class of f is equal to f' and $f(a_1 \otimes \cdots \otimes a_n) = 0$ if $a_i = 1$ for some $1 \leq i \leq N - 1$. Then direct calculations show that

$$(f - \delta_{n-1}^\nu s_{n-1}^{N-1,\nu}(f))(a_1 \otimes \cdots \otimes a_n) = (s_n^{N-1,\nu} \delta_n^\nu(f))(a_1 \otimes \cdots \otimes a_n) = 0$$

if $a_i = 1$ for some $1 \leq i \leq N$.

Now assume that $f \in C^n(R)^\nu$, $g \in C^m(R)^\nu$. Let us introduce the following notation

$$\Delta'(f \otimes g) := \sum_{i=1}^m (-1)^{i(n+m-1)} \Delta_i(f \smile g).$$

Then $\Delta(f \smile g) = \Delta'(f \otimes g) + (-1)^{nm} \Delta'(g \otimes f)$.

The remaining part of the proof is analogous to the proof of Theorem 1 (cf. [2]). □

Note that a Frobenius algebra is symmetric if and only if its Nakayama automorphism equals Id_R for some bilinear form. So Theorem 1 is a special case of Theorem 2.

Corollary 3. *If $\mathrm{ord}(\nu) < \infty$ and $\mathrm{char} k \nmid \mathrm{ord}(\nu)$, then the map defined by the formula (3.1) induces a BV-differential on the Gerstenhaber algebra $(\mathrm{HH}^*(R), \smile, [\cdot, \cdot])$.*

Proof. This follows from Corollary 2 of Lemma 1 and the fact that $\Theta_R^\nu : \mathrm{HH}^*(R)^{\nu\uparrow} \rightarrow \mathrm{HH}^*(R)^\nu$ is an isomorphism of Gerstenhaber algebras if $\mathrm{ord}(\nu) < \infty$ and $\mathrm{char} k \nmid \mathrm{ord}(\nu)$. □

4 On a family of self-injective algebras of tree type D_n

Let $n \geq 4$, $r \geq 1$. Let us define a k -algebra $R = R(n, r)$. Consider a quiver with relations (\mathcal{Q}, I) . Its set of vertices is $\mathcal{Q}_0 = \mathbb{Z}_r \times \{j \in \mathbb{N} \mid 1 \leq j \leq n\}$. The set of arrows \mathcal{Q}_1 of the quiver \mathcal{Q} consists of the following elements:

$$\begin{aligned} \gamma_{i,p} &: (i, n-2) \rightarrow (i, p), \quad \beta_{i,p} : (i, p) \rightarrow (i+1, 1), \\ \alpha_{i,j} &: (i, j) \rightarrow (i, j+1) \quad (1 \leq i \leq r, p \in \{n-1, n\}, 1 \leq j \leq n-3). \end{aligned}$$


In addition we use the following auxiliary notation:

$$\tau_i = \beta_{i,n} \gamma_{i,n}, \quad \omega_{i,j_2,j_1} = \alpha_{i,j_2} \cdots \alpha_{i,j_1}, \quad \mu_{i,j} = \omega_{i,j,1}, \quad \eta_{i,j} = \omega_{i,n-3,j}.$$

From here on for the uniformity of the notation we additionally suppose that the empty product of the arrows of the quiver is identified with an appropriate idempotent of the algebra R ; for example, $\mu_{i,0} = e_{i,1}$, $\omega_{i,j-1,j} = e_{i,j}$, $\eta_{i,n-2} = e_{i,n-2}$. Denote by $\phi : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ the map such that $\phi(j) = j$ for $1 \leq j \leq n-2$, $\phi(n-1) = n$ and $\phi(n) = n-1$.

The ideal I is generated by the elements

$$\begin{aligned} &\gamma_{i,\phi(p)} \eta_{i,1} \beta_{i-1,p}, \quad \beta_{i,n-1} \gamma_{i,n-1} - \tau_i, \quad \mu_{i+1,j} \tau_i \eta_{i,j} \\ &(1 \leq i \leq r, 1 \leq j \leq n-3, p \in \{n-1, n\}). \end{aligned}$$



Set $R = R(n, r) = k\mathcal{Q}/I$. It is easily verified that R is a Frobenius algebra. For two paths w_1 and w_2 of the quiver \mathcal{Q} the bilinear form $\langle \cdot, \cdot \rangle$ is defined in the following way:

$$\langle w_1, w_2 \rangle = \epsilon(w_1 w_2) := \begin{cases} 1, & \text{if } w_1 w_2 \text{ is a nonzero path of length } n-1, \\ 0 & \text{otherwise.} \end{cases}$$

This bilinear form determines the Nakayama automorphism ν . This automorphism is defined on idempotents and arrows by the formulas

$$\begin{aligned} \nu(e_{i,t}) &= e_{i-1,t}, \quad \nu(\alpha_{i,j}) = \alpha_{i-1,j}, \quad \nu(\gamma_{i,p}) = \gamma_{i-1,p}, \quad \nu(\beta_{i,p}) = \beta_{i-1,p} \\ &(1 \leq i \leq r, 1 \leq t \leq n, 1 \leq j \leq n-2, p \in \{n-1, n\}). \end{aligned}$$

It is clear that the set

$$\begin{aligned} B_R &= \{\mu_{i+1,t-1} \tau_i \eta_{i,j} \mid 1 \leq i \leq r, 1 \leq t \leq j \leq n-2\} \\ &\cup \{\omega_{i,t-1,j} \mid 1 \leq i \leq r, 1 \leq j \leq t \leq n-2\} \\ &\cup \{\gamma_{i,p} \eta_{i,j} \mid 1 \leq i \leq r, 1 \leq j \leq n-2, p \in \{n-1, n\}\} \\ &\cup \{\mu_{i+1,j-1} \beta_{i,p} \mid 1 \leq i \leq r, 1 \leq j \leq n-2, p \in \{n-1, n\}\} \\ &\cup \{\gamma_{i,p} \mu_{i,n-3} \gamma_{i-1,p} \mid 1 \leq i \leq r, p \in \{n-1, n\}\} \end{aligned}$$

is a k -basis of R . We define for $b \in B_R$ the element $\bar{b} \in B_R$ by the equalities

$$\begin{aligned}\overline{\mu_{i+1,t-1}\tau_i\eta_{i,j}} &= \omega_{i+1,j-1,t} \quad (1 \leq i \leq r, 1 \leq t \leq j \leq n-2) \\ \overline{\omega_{i,t-1,j}} &= \mu_{i+1,j-1}\tau_i\eta_{i,t} \quad (1 \leq i \leq r, 1 \leq j \leq t \leq n-2) \\ \overline{\gamma_{i,p}\eta_{i,j}} &= \mu_{i+1,j-1}\beta_{i,p} \quad (1 \leq i \leq r, 1 \leq j \leq n-2, p \in \{n-1, n\}) \\ \overline{\mu_{i+1,j-1}\beta_{i,n-1}} &= \gamma_{i+1,p}\eta_{i+1,j} \quad (1 \leq i \leq r, 1 \leq j \leq n-2, p \in \{n-1, n\}) \\ \overline{\gamma_{i,p}\mu_{i,n-3}\gamma_{i-1,p}} &= e_{i,p} \quad (1 \leq i \leq r, p \in \{n-1, n\}).\end{aligned}$$

It is easy to show that $\langle a, b \rangle = \begin{cases} 1, & \text{if } b = \bar{a}, \\ 0, & \text{if } b \neq \bar{a}, \end{cases}$ for $a, b \in B_R$.

The Hochschild cohomology algebra of $R(n, r)$ is described in [12] and [5]. Let us recall some results of these works. Denote by e_x the idempotent of the algebra R corresponding to a vertex x of the quiver \mathcal{Q} . Then $\{e_x \otimes e_y\}_{x,y}$ is a full set of orthogonal primitive idempotents for the algebra Λ . Denote by $P_{[x][y]} = \Lambda e_x \otimes e_y$ the projective Λ -module, which corresponds to idempotent $e_x \otimes e_y$. Let $\sigma : R \rightarrow R$ be the automorphism of R , which is defined on the idempotents and arrows by the formulas

$$\begin{aligned}\sigma(e_{i,j}) &= e_{i+n-1, \phi^n(j)}, \sigma(\alpha_{i,j}) = \alpha_{i+n-1,j}, \sigma(\gamma_{i,p}) = -\gamma_{i+n-1, \phi^n(p)}, \\ \sigma(\beta_{i,p}) &= \beta_{i+n-1, \phi^n(p)} \quad (1 \leq i \leq r, 1 \leq j \leq n-2, p \in \{n-1, n\}).\end{aligned}$$

Denote by Q_t the t -th module in the minimal projective bimodule resolution of R . Then

$$\begin{aligned}Q_{2m} &= \bigoplus_{i=1}^r \left(\left(\bigoplus_{j=1}^{n-2-m} P_{[i+m, j+m][i, j]} \right) \right. \\ &\quad \oplus \left(\bigoplus_{j=n-1-m}^{n-2} P_{[i+m, j+m-(n-2)][i, j]} \right) \\ &\quad \left. \oplus P_{[i+m, \phi^m(n-1)][i, n-1]} \oplus P_{[i+m, \phi^m(n)][i, n]} \right) \quad (0 \leq m \leq n-2), \\ Q_{2m+1} &= \bigoplus_{i=1}^r \left(\left(\bigoplus_{j=1}^{n-3-m} P_{[i+m, j+m+1][i, j]} \right) \oplus P_{[i+m, n-1][i, n-2-m]} \right. \\ &\quad \oplus P_{[i+m, n][i, n-2-m]} \oplus \left(\bigoplus_{j=n-1-m}^{n-2} P_{[i+m+1, j+m-(n-2)][i, j]} \right) \\ &\quad \left. \oplus P_{[i+m+1, m+1][i, n-1]} \oplus P_{[i+m+1, m+1][i, n]} \right) \quad (0 \leq m \leq n-3).\end{aligned}$$

Moreover, $Q_{t+l(2n-3)} = {}_{\sigma^l}(Q_t)_1$ for $0 \leq t \leq 2n-4$, $l > 0$. The definitions of the differentials $d_t^Q : Q_{t+1} \rightarrow Q_t$ can be found in [12]. The augmentation map $\mu : Q_0 \rightarrow R$ is defined by the formula $\mu(a \otimes b) = ab \in R$ for $a \otimes b \in P_{[x][x]}$ ($x \in \mathcal{Q}_0$). It is easy to check that $\nu(P_{[i_1, j_1][i_2, j_2]})_\nu \simeq P_{[i_1+1, j_1][i_2+1, j_2]}$. These isomorphisms give isomorphisms $\theta_t : Q_t \rightarrow {}_\nu(Q_t)_\nu$ for $t \geq 0$. Using the description of the differentials d_t^Q it is easy to verify that θ_* is a chain map, which lifts the isomorphism $\phi_\nu : R \rightarrow {}_\nu R_\nu$ defined by the formula $\phi_\nu(a) = \nu a$ for $a \in R$. Let $f \in \text{Hom}_\Lambda(Q_t, R)$, $f d_t^Q = 0$ and f is defined on the direct summands of Q_t by the

equalities $f(e_{[i_1, j_1][i_2, j_2]}) = f_{[i_1, j_1][i_2, j_2]} \in R$. It follows from [10, Lemma 2] that f^ν is defined by the equalities

$$f^\nu(e_{[i_1, j_1][i_2, j_2]}) = \nu^{-1} f_{[i_1-1, j_1][i_2-1, j_2]}.$$

To calculate the Lie bracket we will partially use the algorithm described in [9]. To apply this algorithm we need the homomorphism of left R -modules $D_t : Q_t \rightarrow Q_{t+1}$ ($t \geq 0$), $D_{-1} : R \rightarrow Q_0$, which satisfy the equations

$$\begin{aligned} D_{t+1}D_t &= 0 \quad (t \geq 0), \quad D_{t-1}d_{t-1}^Q + d_t^Q D_t = \text{Id}_{Q_t} \quad (t \geq 1), \\ D_{-1}\mu + d_0^Q D_0 &= \text{Id}_{Q_0}, \quad \mu D_{-1} = \text{Id}_R. \end{aligned} \quad (4.1)$$

Let $D_{-1} : R \rightarrow Q_0$ be a homomorphism of left modules such that $D_{-1}(e_x) = e_{[x][x]}$ for all $x \in Q_0$.

Now let us define the homomorphisms D_t ($t \geq 0$). We will define them on the elements of the form $e_x \otimes b$ ($x \in Q_0, b \in B_R$). Let $0 \leq m \leq n-3$. Then $D_{2m}(e_{i+m, j+m} \otimes e_{i, j} b)$ ($1 \leq i \leq r, 1 \leq j \leq n-2-m$) is equal to

$$\sum_{s=q}^{j-1} \omega_{i+m, j+m-1, s+m+1} \otimes \omega_{i, s-1, q}$$

if $b = \omega_{i, j-1, q}$, $1 \leq q \leq j$,

$$\sum_{s=1}^{j-1} \omega_{i+m, j+m-1, s+m+1} \otimes \mu_{i, s-1} \beta_{i-1, p} + \omega_{i+m, j+m-1, m+1} \otimes e_{i-1, p}$$

if $b = \mu_{i, j-1} \beta_{i-1, p}$, $p \in \{n-1, n\}$,

$$\begin{aligned} &\sum_{s=1}^{j-1} \omega_{i+m, j+m-1, s+m+1} \otimes \mu_{i, s-1} \tau_{i-1} \eta_{i-1, q} \\ &+ \omega_{i+m, j+m-1, m+1} \otimes \gamma_{i-1, n-1} \eta_{i-1, q} - \omega_{i+m, j+m-1, q+m-(n-2)} \otimes e_{i-1, q} \end{aligned}$$

if $b = \mu_{i, j-1} \tau_{i-1} \eta_{i-1, q}$, $n-1-m \leq q \leq n-2$,

$$\begin{aligned} &\sum_{s=1}^{j-1} \omega_{i+m, j+m-1, s+m+1} \otimes \mu_{i, s-1} \tau_{i-1} \eta_{i-1, q} \\ &+ \omega_{i+m, j+m-1, m+1} \otimes \gamma_{i-1, n-1} \eta_{i-1, q} \\ &+ \mu_{i+m, j+m-1} \beta_{i+m-1, \phi^m(n-1)} \otimes \omega_{i-1, n-3-m, q} \\ &+ \sum_{s=q}^{n-3-m} \mu_{i+m, j+m-1} \tau_{i+m-1} \eta_{i+m-1, s+m+1} \otimes \omega_{i-1, s-1, q} \end{aligned}$$

if $b = \mu_{i, j-1} \tau_{i-1} \eta_{i-1, q}$, $j \leq q \leq n-2-m$;

$D_{2m}(e_{i+m, j+m-(n-2)} \otimes e_{i, j} b)$ ($1 \leq i \leq r, n-1-m \leq j \leq n-2$) is equal to 0 if $b \neq \mu_{i, j-1} \tau_{i-1} \eta_{i-1, j}$, and is equal to $-e_{i+m, j+m-(n-2)} \otimes e_{i-1, j}$ if $b = \mu_{i, j-1} \tau_{i-1} \eta_{i-1, j}$;

$D_{2m}(e_{i+m, \phi^m(p)} \otimes e_{i, p} b)$ ($1 \leq i \leq r, p \in \{n-1, n\}$) is equal to 0

if $b = e_{i,p}$ or $b = \gamma_{i,p}\eta_{i,q}$, $n-1-m \leq q \leq n-2$,

$$e_{i+m,\phi^m(p)} \otimes \omega_{i,n-3-m,q} + \sum_{s=q}^{n-3-m} \gamma_{i+m,\phi^m(p)}\eta_{i+m,s+m+1} \otimes \omega_{i,s-1,q}$$

if $b = \gamma_{i,p}\eta_{i,q}$, $1 \leq q \leq n-2-m$,

$$\begin{aligned} & e_{i+m,\phi^m(p)} \otimes \mu_{i,n-3-m}\beta_{i-1,p} \\ & + \sum_{s=q}^{n-3-m} \gamma_{i+m,\phi^m(p)}\eta_{i+m,s+m+1} \otimes \mu_{i,s-1,q}\beta_{i-1,p} \\ & + \gamma_{i+m,\phi^m(p)}\eta_{i+m,m+1} \otimes e_{i-1,p} \end{aligned}$$

if $b = \gamma_{i,p}\eta_{i,1}\beta_{i-1,p}$.

Let $0 \leq m \leq n-3$. Then

$D_{2m+1}(e_{i+m,j+m+1} \otimes e_{i,j}b)$ ($1 \leq i \leq r, 1 \leq j \leq n-3-m$) is equal to 0 if $b \neq \mu_{i,j-1}\tau_{i-1}\eta_{i-1,j}$, and is equal to $e_{i+m,j+m+1} \otimes e_{i-1,j}$ if $b = \mu_{i,j-1}\tau_{i-1}\eta_{i-1,j}$;

$D_{2m+1}(e_{i+m+1,j+m-(n-2)} \otimes e_{i,j}b)$ ($1 \leq i \leq r, n-1-m \leq j \leq n-2$) is equal to

$$\sum_{s=q}^{j-1} \omega_{i+m+1,j+m-(n-1),s+m+(n-3)} \otimes \omega_{i,s-1,q}$$

if $b = \omega_{i,j-1,q}$, $n-1-m \leq q \leq j$,

$$\sum_{s=n-2-m}^{j-1} \omega_{i+m+1,j+m-(n-1),s+m+(n-3)} \otimes \omega_{i,s-1,q}$$

if $b = \omega_{i,j-1,q}$, $1 \leq q \leq n-2-m$,

$$\begin{aligned} & \sum_{s=n-2-m}^{j-1} \omega_{i+m+1,j+m-(n-1),s+m+(n-3)} \otimes \mu_{i,s-1}\beta_{i-1,n-1} \\ & + \mu_{i+m+1,j+m-(n-1)}\beta_{i+m,\phi^{m+1}(n-1)} \otimes e_{i-1,n-1} \end{aligned}$$

if $b = \mu_{i,j-1}\beta_{i-1,n-1}$,

$$\begin{aligned} & \sum_{s=n-2-m}^{j-1} \omega_{i+m+1,j+m-(n-1),s+m+(n-3)} \otimes \mu_{i,s-1}\beta_{i-1,n} \\ & - \mu_{i+m+1,j+m-(n-1)}\beta_{i+m,\phi^{m+1}(n)} \otimes e_{i-1,n} \end{aligned}$$

if $b = \mu_{i,j-1}\beta_{i-1,n}$,

$$\begin{aligned} & \sum_{s=n-2-m}^{j-1} \omega_{i+m+1,j+m-(n-1),s+m+(n-3)} \otimes \mu_{i,s-1}\tau_{i-1}\eta_{i-1,q} \\ & - \mu_{i+m+1,j+m-(n-1)}\beta_{i+m,\phi^{m+1}(n)} \otimes \gamma_{i-1,n}\eta_{i-1,q} \\ & + \mu_{i+m+1,j+m-(n-1)}\beta_{i+m,\phi^{m+1}(n-1)} \otimes \gamma_{i-1,n-1}\eta_{i-1,q} \\ & - \sum_{s=q}^{n-2} \mu_{i+m+1,j+m-(n-1)}\tau_{i+m}\eta_{i+m,s+m-(n-3)} \otimes \omega_{i-1,s-1,q} \end{aligned}$$

if $b = \mu_{i,j-1}\tau_{i-1}\eta_{i,q}$ ($j \leq q \leq n-2$);
 $D_{2m+1}(e_{i+m+1,m+1} \otimes e_{i,n-1}b)$ ($1 \leq i \leq r$) is equal to 0
if $b \neq \gamma_{i,n-1}\eta_{i,1}\beta_{i-1,n-1}$, and is equal to

$$\sum_{s=n-2-m}^{n-2} \omega_{i+m+1,m,s+m-(n-1)} \otimes \mu_{i,s-1}\beta_{i-1,n-1} \\ + \mu_{i+m+1,m}\beta_{i+m,\phi^{m+1}(n-1)} \otimes e_{i-1,n-1}$$

if $b = \gamma_{i,n-1}\eta_{i,1}\beta_{i-1,n-1}$;
 $D_{2m+1}(e_{i+m+1,m+1} \otimes e_{i,n}b)$ ($1 \leq i \leq r$) is equal to 0
if $b = e_{i,n}$,

$$- \sum_{s=q}^{n-2} \omega_{i+m+1,m,s+m+(n-3)} \otimes \omega_{i,s-1,q}$$

if $b = \gamma_{i,n}\eta_{i,q}$, $n-1-m \leq q \leq n-2$,

$$- \sum_{s=n-2-m}^{n-2} \omega_{i+m+1,m,s+m+(n-3)} \otimes \omega_{i,s-1,q}$$

if $b = \gamma_{i,n}\eta_{i,q}$, $1 \leq q \leq n-2-m$,

$$- \sum_{s=n-2-m}^{n-2} \omega_{i+m+1,m,s+m-(n-1)} \otimes \mu_{i,s-1}\beta_{i-1,n-1} \\ + \mu_{i+m+1,m}\beta_{i+m,\phi^{m+1}(n)} \otimes e_{i-1,n}$$

if $b = \gamma_{i,n}\eta_{i,1}\beta_{i-1,n}$;
 $D_{2m+1}(e_{i+m,\phi^m(p)} \otimes e_{i,n-2-m}b)$ ($1 \leq i \leq r, p \in \{n-1, n\}$) is equal to 0
if $b = \omega_{i,n-3-m,q}$, $1 \leq q \leq n-2-m$ or $b = \mu_{i,n-3-m}\beta_{i-1,p}$,

$$e_{i+m\phi^m(p)} \otimes e_{i-1,\phi(p)}$$

if $b = \mu_{i,n-3-m}\beta_{i-1,\phi(p)}$,

$$e_{i+m\phi^m(n)} \otimes \gamma_{i-1,n-1}\eta_{i,q}$$

if $p = n$, $b = \mu_{i,n-3-m}\tau_{i-1}\eta_{i-1,q}$ ($n-2-m \leq q \leq n-2$),

$$e_{i+m\phi^m(n-1)} \otimes \gamma_{i-1,n}\eta_{i,q} + \sum_{s=q}^{n-2} \gamma_{i+m\phi^m(n-1)}\eta_{i+m,s+m-(n-3)} \otimes \omega_{i-1,s-1,q}$$

if $p = n-1$, $b = \mu_{i,n-3-m}\tau_{i-1}\eta_{i-1,q}$ ($n-2-m \leq q \leq n-2$).

We define the homomorphism D_{2n-4} in the following way.

$D_{2n-4}(e_{i+n-2,j} \otimes e_{i,j}b)$ is equal to 0 if $b \neq \mu_{i,j-1}\tau_{i-1}\eta_{i,j}$, and is equal to $-e_{i+n-2,j} \otimes e_{i-1,j}$ if $b = \mu_{i,j-1}\tau_{i-1}\eta_{i,j}$;

$D_{2n-4}(e_{i+n-2,\phi^{n-2}(n-1)} \otimes e_{i,n-1}b)$ is equal to 0 if $b \neq \gamma_{i,n-1}\eta_{i-1,1}\beta_{i-1,n-1}$, and is equal to $-e_{i+n-2,\phi^{n-2}(n-1)} \otimes e_{i-1,n-1}$ if $b = \gamma_{i,n-1}\eta_{i-1,1}\beta_{i-1,n-1}$;

$D_{2n-4}(e_{i+n-2, \phi^{n-2}(n)} \otimes e_{i,n} b)$ is equal to 0 if $b \neq \gamma_{i,n} \eta_{i-1,1} \beta_{i-1,n}$, and is equal to $e_{i+n-2, \phi^{n-2}(n)} \otimes e_{i-1,n}$ if $b = \gamma_{i,n} \eta_{i-1,1} \beta_{i-1,n}$.

Moreover, $D_{t+l(2n-3)} = (D_t)_{(\sigma^l, 1)}$ for $0 \leq t \leq 2n-4$, $l > 0$. Direct calculations show that the maps D_t ($t \geq -1$) satisfy the equalities (4.1).

Let $\theta_{\text{Bar}, t} : \text{Bar}_n(R) \rightarrow {}_\nu(\text{Bar}_n(R))_\nu$ be defined by the formula $\theta_{\text{Bar}, t}(a_0 \otimes \cdots \otimes a_{t+1}) = \nu a_0 \otimes \cdots \otimes \nu a_{t+1}$. It is easy to show that the chain maps $\Phi_* : Q_* \rightarrow \text{Bar}_*(R)$ and $\Psi_* : \text{Bar}_*(R) \rightarrow Q_*$ constructed using the maps d_t^Q and D_t ($t \geq 0$) by the algorithm from [9] satisfy the equalities $\theta_{\text{Bar}, t} \Phi_t = (\Phi_t)_{(\nu, \nu)} \theta_t$ and $\theta_t \Psi_t = (\Psi_t)_{(\nu, \nu)} \theta_{\text{Bar}, t}$. Then it follows from [10, Lemma 2] that t -cocycle lies in $\text{Im } \Theta_R^\nu$ if and only if it can be represented by a homomorphism $f \in \text{Hom}_\Lambda(Q_t, R)$ such that

$$f(e_{[i_1, j_1][i_2, j_2]}) = \nu^{-1} f_{[i_1-1, j_1][i_2-1, j_2]}. \quad (4.2)$$

5 The description of the Lie bracket and the BV-structure

In this section we suppose that $R = R(n, r)$. Let w be a path from a vertex x to a vertex y . Denote by w^* the element of $\text{Hom}_\Lambda(P_{[y][x]}, R)$ such that $w^*(e_y \otimes e_x) = w$. For $1 \leq i \leq r$ consider the following auxiliary homomorphisms:

$$\begin{aligned} w_{i,m,j} &= (\omega_{i,j+m-1,j})^* \quad (0 \leq m \leq n-3, 1 \leq j \leq n-2-m); \\ t_{i,m,j} &= (\mu_{i+1,j+m-(n-1)} \tau_i \eta_{i,j})^* \\ &\quad (1 \leq m \leq n-2, n-1-m \leq j \leq n-2); \\ u_{i,m,q} &= (\gamma_{i,q} \eta_{i,n-2-m})^* \quad (0 \leq m \leq n-3, q \in \{n-1, n\}); \\ v_{i,m,q} &= (\mu_{i+1,m} \beta_{i,q})^* \quad (0 \leq m \leq n-3, q \in \{n-1, n\}); \\ u_{i,q} &= e_{i,q}^* \quad (q \in \{n-1, n\}); \\ v_{i,q} &= (\gamma_{i+1,q} \eta_{i+1,1} \beta_{i,q})^* \quad (q \in \{n-1, n\}). \end{aligned}$$

Let now define some elements of the algebra $\text{HH}^*(R)$.

a) Define 1-cocycle $\varepsilon_1 \in \text{Hom}_\Lambda(Q_1, R)$ by the formula

$$\varepsilon_1 = u_{1,0,n-1} + u_{1,0,n}.$$

b) Let $s = 2m + l(2n-3)$, $0 \leq m \leq n-2$, $r | m + l(n-1)$, $2 | m + ln$ and one of the following conditions is satisfied: $\text{char } k = 2$ or $2 | l$. Define s -cocycle $f_s \in \text{Hom}_\Lambda(Q_s, R)$ by the formula

$$f_s = \sum_{i=1}^r \left(\sum_{j=1}^{n-2-m} w_{i,m,j} + u_{i,n-1} + u_{i,n} \right).$$

c) Let $s = 2m + 1 + l(2n-3)$, $0 \leq m \leq n-3$, $r | m + l(n-1)$, $2 \nmid m + ln$ and one of the following conditions is satisfied: $\text{char } k = 2$ or $2 \nmid l$. Define s -cocycle $g_s \in \text{Hom}_\Lambda(Q_s, R)$ by the formula

$$g_s = \sum_{i=1}^r \left(\sum_{j=n-1-m}^{n-2} t_{i,m,j} + u_{i,m,n-1} + v_{i,m,n-1} \right).$$

d) Let $s = (l + 1)(2n - 3) - 1$, $r|(l + 1)(n - 1) - 1$, $2 \nmid (l + 1)n$. Define s -cocycle $h_s \in \text{Hom}_\Lambda(Q_s, R)$ by the formula

$$h_s = \sum_{i=1}^r \sum_{j=1}^{n-2} (-1)^j w_{i,0,j}.$$

e) Let $s = (l + 1)(2n - 3) - 1$, $r|(l + 1)(n - 1) - 1$, $2|n$ and one of the following conditions is satisfied: $\text{char } k = 2$ or $2|l$. Define s -cocycle $p_s \in \text{Hom}_\Lambda(Q_s, R)$ by the formula

$$p_s = \sum_{i=1}^r \left(\sum_{j=1}^{n-2} (-1)^j w_{i,0,j} + u_{i,n-1} \right).$$

f) Let $s = l(2n - 3)$, $l \geq 1$, $r|l(n - 1) - 1$, $2|ln$ and either $\text{char } k = 2$ or $2 \nmid l$. Denote by $\chi_s \in \text{Hom}_\Lambda(Q_s, R)$ the s -cocycle, which is equal to $v_{r,n}$ on $P_{[1,n][r,n]}$ and is equal to 0 on other direct summands of Q_s .

g) Let $s = l(2n - 3)$, $r|l(n - 1) - 1$, $2 \nmid ln$. Denote by $\xi_s \in \text{Hom}_\Lambda(Q_s, R)$ the s -cocycle, which is equal to $t_{r,n-2,1}$ on $P_{[1,1][r,1]}$ and is equal to 0 on other direct summands of Q_s .

h) Let $r = 1$. For $1 \leq j \leq n - 2$ denote by $\varepsilon_0^{(j)}$ the 0-cocycle, which is equal to $t_{1,n-2,j}$ on $P_{[1,j][1,j]}$ and is equal to 0 on other direct summands of Q_0 . For $q \in \{n - 1, n\}$ denote by $\varepsilon_0^{(q)}$ the 0-cocycle, which is equal to $v_{1,q}$ on $P_{[1,q][1,q]}$ and is equal to 0 on other direct summands of Q_0 .

It was shown in [5] that the elements defined in a)–h) are cocycles for the corresponding values of s and that they generate $\text{HH}^*(R)$ as a k -algebra. In addition, ξ_s can be excluded from the set of generators in the case where $\text{char } k \nmid \frac{n-1}{2}$ and χ_s can be excluded from the set of generators in the case where $2|n$, $\text{char } k \nmid n - 1$. Moreover, it is proved in the same work that the elements of the form $f_s, g_s, h_s, p_s, \varepsilon_1 f_s, \varepsilon_1 g_s, \chi_s, \xi_s$ and $\varepsilon_0^{(q)}$ generate $\text{HH}^*(R)$ as a k -linear space.

If $\text{char } k \nmid r$, then $\text{HH}^*(R)$ is a BV-algebra by the Corollary of Theorem 2. Since f_s, g_s, h_s and p_s satisfy the condition (4.2), they lie in the image of Θ_R^ν (even if $\text{char } k|r$).

Let us introduce the following notation

$$F(x) = \begin{cases} 0, & \text{if } x = \varepsilon_1, \\ m + l(n - 1), & \text{if } x = f_{2m+l(2n-3)} \text{ or } x = g_{2m+1+l(2n-3)}, \\ l(n - 1) - 1, & \text{if } x = p_{l(2n-3)-1}, x = h_{l(2n-3)-1}, \\ & x = \xi_{l(2n-3)} \text{ or } x = \chi_{l(2n-3)}. \end{cases}$$

Note that $r|F(x)$ in all the cases.

Proposition 1. *Let $x \in \{\varepsilon_1, f_s, g_s, h_s, p_s\}$. Then*

$$[x, \varepsilon_1] = \frac{F(x)}{r} x \quad (5.1)$$

in $\text{HH}^*(R)$.

Proof. Recall the construction of the chain maps $\Phi_t : Q_t \rightarrow \text{Bar}_t(R)$ and $\Psi_t : \text{Bar}_t(R) \rightarrow Q_t$ from [9].

Firstly, define Φ_0 and Ψ_0 by the equalities $\Phi_0(e_x \otimes e_x) = e_x \otimes e_x$, $\Psi_0(e_x \otimes e_x) = e_x \otimes e_x$, $\Psi_0(e_x \otimes e_y) = 0$ for $x, y \in \mathcal{Q}_0$, $x \neq y$. For $t > 0$ the map Ψ_t is defined by the formula

$$\Psi_t(1 \otimes a_1 \otimes \cdots \otimes a_n \otimes 1) = D_{t-1}(\Psi_{t-1}(1 \otimes a_1 \otimes \cdots \otimes a_{n-1} \otimes 1)a_n). \quad (5.2)$$

From now on we assume that $Q_t = \bigoplus_{p \in X_t} P_{[x_p][y_p]}$ for all $t \geq 0$, where $\{X_t\}_{t \geq 0}$ is a set of disjoint sets. Let $\pi_p : Q_t \rightarrow P_{[x_p][y_p]}$ be the canonical projection and

$$\pi_{p'} d_{t-1}^Q(e_{x_p} \otimes e_{y_p}) = a_{p,p'} \otimes e_{y_{p'}} + b_{p,p'} \otimes c_{p,p'}$$

for $p' \in X_{t-1}$, $p \in X_t$, where $a_{p,p'}, b_{p,p'} \in R$ and $c_{p,p'}$ are in Jacobson radical of R . Then

$$\Phi_t(e_{x_p} \otimes e_{y_p}) = \sum_{p' \in X_{t-1}} b_{p,p'} \Phi_{t-1}(e_{x_{p'}} \otimes e_{y_{p'}}) c_{p,p'} \otimes 1. \quad (5.3)$$

Consequently,

$$\Psi_t \Phi_t(e_{x_p} \otimes e_{y_p}) = \sum_{p' \in X_{t-1}} b_{p,p'} D_{t-1}(\Psi_{t-1} \Phi_{t-1}(e_{x_{p'}} \otimes e_{y_{p'}}) c_{p,p'}).$$

It follows from these formulas and induction on t that for all $t \geq 0$, $p \in X_t$ we have

$$\Phi_t(e_{x_p} \otimes e_{y_p}) = e_{x_p} \otimes a_{p,1} \otimes \cdots \otimes a_{p,t} \otimes e_{y_p} + \sum_{z \in Y_p} a_{z,0} \otimes \cdots \otimes a_{z,t+1}, \quad (5.4)$$

where

$$\Psi_t(e_{x_p} \otimes a_{p,1} \otimes \cdots \otimes a_{p,t} \otimes e_{y_p}) = e_{x_p} \otimes e_{y_p}$$

and $\Psi_t(a_{z,0} \otimes \cdots \otimes a_{z,t+1}) = 0$ for all $z \in Y_p$.

The equality $[\varepsilon_1, \varepsilon_1] = 0$ follows from the definition of the Lie bracket and the fact that ε_1 is an element of odd degree. For other elements we use the formula

$$[f, \varepsilon_1] = ((f\Psi_t) \circ (\varepsilon_1\Psi_1) - (\varepsilon_1\Psi_1) \circ (f\Psi_t))\Phi_t.$$

Consider a \mathbb{Z} -grading on the algebra R such that the idempotents and arrows, except $\gamma_{1,n-1}$ and $\gamma_{1,n}$, are of degree 0 and the arrows $\gamma_{1,n-1}$ and $\gamma_{1,n}$ are of degree 1. This grading induces a grading on Λ . We can define a grading on the direct summands of Q_t ($t \geq 0$) in such a way that Q_* is a graded resolution of the module R . Let $t = t' + l(2n - 3)$ ($0 \leq t' \leq 2n - 4$), $p \in X_t$ and $P_{[x_p][y_p]} \simeq \sigma^l(P)_1$, where the module P appears in the formula for $Q_{t'}$ as the module $P_{[i_2, j_2][i_1, j_1]}$. We define the degree of the element $e_{x_p} \otimes e_{y_p} \in Q_t$ in the following way:

1) if $j_1, j_2 \notin \{n - 1, n\}$, then

$$\deg(e_{x_p} \otimes e_{y_p}) = \left\lceil \frac{i_2 + l(n - 1) - 1}{r} \right\rceil - \left\lceil \frac{i_1 - 1}{r} \right\rceil;$$

2) if $j_1 \notin \{n - 1, n\}$, $j_2 \in \{n - 1, n\}$, then

$$\deg(e_{x_p} \otimes e_{y_p}) = \left\lceil \frac{i_2 + l(n - 1)}{r} \right\rceil - \left\lceil \frac{i_1 - 1}{r} \right\rceil;$$

3) if $j_1 \in \{n-1, n\}$, $j_2 \notin \{n-1, n\}$, then

$$\deg(e_{x_p} \otimes e_{y_p}) = \left\lceil \frac{i_2 + l(n-1) - 1}{r} \right\rceil - \left\lceil \frac{i_1}{r} \right\rceil;$$

4) if $j_1, j_2 \in \{n-1, n\}$, then

$$\deg(e_{x_p} \otimes e_{y_p}) = \left\lceil \frac{i_2 + l(n-1)}{r} \right\rceil - \left\lceil \frac{i_1}{r} \right\rceil.$$

Here we denote by $\lceil a \rceil$ the smallest integer which is greater or equal to a . It is easy to show that the differentials d_t^Q are actually of degree 0 for the grading introduced in 1)–4). Moreover, it is easy to check that, if we introduce the grading on the modules $\text{Bar}_t(R)$ in such a way that $\deg(a_0 \otimes \cdots \otimes a_{t+1}) = \sum_{i=0}^{t+1} \deg(a_i)$ for homogeneous elements $a_i \in R$ ($0 \leq i \leq t+1$), then $\text{Bar}_*(R)$ becomes a graded resolution of R . In addition Φ_t ($t \geq 0$) is a homomorphism of graded modules. It is easy to check that $\varepsilon_1 \Psi_1(b) = \deg(b)b$ for $b \in B_R$.

It is clear that for elements $f \in C^1(R) = \text{Hom}_k(R, R)$ and $g \in C^t(R) = \text{Hom}_k(R^{\otimes t}, R)$ the composition product $f \circ g$ is just a composition of f and g . Then $((\varepsilon_1 \Psi_1) \circ (f \Psi_t)) \Phi_t = \varepsilon_1 \Psi_1 f$ for $f \in \text{Hom}_\Lambda(Q_t, R)$, $f d_t^Q = 0$. Then it is easy to show that

$$\begin{aligned} & ((\varepsilon_1 \Psi_1) \circ (f \Psi_t)) \Phi_t \\ &= \begin{cases} 0, & \text{if } f \in \{f_s, h_s, p_s\}, \\ \sum_{j=n-1-m}^{n-2} t_{1,m,j} + u_{1,m,n-1}, & \text{if } f = g_s. \end{cases} \end{aligned} \quad (5.5)$$

It follows from the formula (5.4) and the formula $(\varepsilon_1 \Psi_1)(b) = \deg(b)b$ ($b \in B_R$) that

$$((f \circ \Psi_t) \circ (\varepsilon_1 \Psi_1)) \Phi_t(e_{x_p} \otimes e_{y_p}) = \deg(e_{x_p} \otimes e_{y_p}) f(e_{x_p} \otimes e_{y_p})$$

for $p \in X_t$, $f \in \text{Hom}_\Lambda(Q_t, R)$, $f d_t^Q = 0$. The assertion of proposition follows from this formula and (5.5). \square

Now we prove a theorem which combined with Proposition 1 and the results of [5] gives a full description of the algebra $\text{HH}^*(R)$ as a Gerstenhaber algebra in all cases and as a BV-algebra in the case $\text{char } k \nmid r$.

Theorem 3. 1) *If $\text{char } k \nmid r$, then $\text{HH}^*(R)$ is a BV-algebra. In this case the BV-differential Δ is defined by the following equalities:*

$$\begin{aligned} \Delta(\varepsilon_1) &= \frac{1}{r}, \quad \Delta(f_s) = \Delta(g_s) = \Delta(h_s) = \Delta(p_s) = 0, \\ \Delta(\varepsilon_1 f_s) &= \frac{f_s}{r} + [f_s, \varepsilon_1], \quad \Delta(\varepsilon_1 g_s) = \frac{g_s}{r} + [g_s, \varepsilon_1], \\ \Delta(\chi_s) &= \frac{l}{r} \left(\frac{n}{2} f_{s-1} - p_{s-1} \right) \quad (2|n, s = l(2n-3)), \\ \Delta(\chi_s) &= 0 \quad (2 \nmid n), \quad \Delta(\xi_s) = \frac{2l}{r} h_{s-1} \quad (s = l(2n-3)), \quad \Delta(\varepsilon_0^{(q)}) = 0. \end{aligned}$$

2) Suppose that $\text{char } k | r$. Then χ_s and ξ_s can be excluded from the set of generators and the Lie bracket is defined on the generators of $\text{HH}^*(R)$ by the equalities (5.1) and the equalities

$$\begin{aligned} [f_{s_1}, f_{s_2}] &= [f_{s_1}, g_{s_2}] = [f_{s_1}, h_{s_2}] = [f_{s_1}, p_{s_2}] = [g_{s_1}, g_{s_2}] = [g_{s_1}, h_{s_2}] \\ &= [g_{s_1}, p_{s_2}] = [h_{s_1}, h_{s_2}] = [p_{s_1}, p_{s_2}] = 0. \end{aligned}$$

Proof. 1) In this case $\text{HH}^*(R)$ is a BV-algebra. Let us consider $\tilde{\varepsilon}_1 \in \text{Hom}_\Lambda(Q_1, R)$ defined by the equality

$$\tilde{\varepsilon}_1 = \frac{\sum_{i=1}^r (u_{i,0,n-1} + u_{i,0,n})}{r} = \frac{\sum_{i=1}^r \nu^{-i} \varepsilon_1 \theta^i}{r}.$$

Then $\tilde{\varepsilon}_1 = \varepsilon_1$ in $\text{HH}^*(R)$ and it is easy to show that $\Delta(\tilde{\varepsilon}_1 \Psi_1) = \frac{1}{r}$.

Let us consider a \mathbb{Z} -grading on the algebra R , which is induced by length. This grading induces a grading on Λ . We can define a grading on the direct summands of Q_t ($t \geq 0$) in such a way that Q_* is a graded resolution of the module R . Let $t = t' + l(2n - 3)$ ($0 \leq t' \leq 2n - 4$), $p \in X_t$ and $P_{[x_p][y_p]} \simeq \sigma^l(P)_1$, where the module P appears in the formula for $Q_{t'}$ as the module $P_{[i_2, j_2][i_1, j_1]}$. Then we define the degree of the element $e_{x_p} \otimes e_{y_p} \in Q_t$ by the formula

$$\deg(e_{x_p} \otimes e_{y_p}) = l(n - 1)^2 + a(n - 1) + \min(j_2, n - 1) - \min(j_1, n - 1).$$

It is easy to show that the differentials d_t^Q are actually of degree 0 for this grading. Moreover, it is easy to check that, if we introduce a grading on the modules $\text{Bar}_t(R)$ in such a way that $\deg(a_0 \otimes \cdots \otimes a_{t+1}) = \sum_{i=0}^{t+1} \deg(a_i)$ for homogeneous elements $a_i \in R$ ($0 \leq i \leq t + 1$), then $\text{Bar}_*(R)$ becomes a graded resolution of R . In addition Φ_t and Ψ_t ($t \geq 0$) are homomorphisms of graded modules.

Let M, N be \mathbb{Z} -graded spaces. We say that a linear map $\varphi : M \rightarrow N$ is of degree q and write $\deg \varphi = q$ if $\deg \varphi(m) = \deg m - q$ for any homogeneous element $m \in M$. Thus it is easy to show that a grading on R induces a grading on $\text{HH}^*(R)$. Then direct inspection shows that $\deg x = (n - 1)F(x)$ for $x \in \{\varepsilon_1, f_s, g_s, h_s, p_s, \xi_s, \chi_s\}$. Note that any $b \in B_R$ is homogeneous and satisfies the equalities $\deg \bar{b} = n - 1 - \deg b$ and $\deg \nu b = \deg b$. In addition we have $\langle a, 1 \rangle = 0$ for homogeneous $a \in R$ such that $\deg a \neq n - 1$. Suppose that $x \in \text{Ker } \delta_s^y$ and $\deg x = q$. Let $a_i \in B_R$ ($1 \leq i \leq n - 1$), $A = a_1 \otimes \cdots \otimes a_{n-1}$. Then

$$\Delta_i x(A) = \sum_{a \in B_R} \langle f(a_i \otimes \cdots \otimes a_{n-1} \otimes \bar{a} \otimes \nu a_1 \otimes \cdots \otimes \nu a_{i-1}), 1 \rangle a.$$

Since

$$\deg f(a_i \otimes \cdots \otimes a_{n-1} \otimes \bar{a} \otimes \nu a_1 \otimes \cdots \otimes \nu a_{i-1}) = \deg A + \deg \bar{a} - q,$$

nonzero coefficients can appear only for $a \in B_R$ such that $\deg a = \deg A - q$, i.e. $\deg(\Delta_i x) = q$. So $\deg(\Delta x) = \deg x$.

If $x \in \{f_s, g_s, h_s, p_s\}$, then $\deg x = (n - 1)F(x)$ and there are no nonzero elements of such degree in $\text{HH}^t(R)$ for $t < s$. So $\Delta(f_s) = \Delta(g_s) = \Delta(h_s) = \Delta(p_s) = 0$. Then the equalities for $\Delta(\varepsilon_1 f_s)$ and $\Delta(\varepsilon_1 g_s)$ follow from (2.1). We can calculate $\Delta(\varepsilon_1 p_s)$ and $\Delta(\varepsilon_1 h_s)$ using the same formula. The formula for $\Delta(\chi_s)$ in the case where $2|n$, $\text{char } k \nmid n - 1$ and the formula for $\Delta(\xi_s)$ in the case where $2 \nmid n$, $\text{char } k \nmid \frac{n-1}{2}$ follow from [5, Lemma 1].

Let now $\tilde{\xi}_s = \sum_{i=1}^r \nu^{-i} \xi_s \theta^i$, $\tilde{\chi}_s = \sum_{i=1}^r \nu^{-i} \chi_s \theta^i$. Then $\xi_s = \frac{\tilde{\xi}_s}{r}$ and $\chi_s = \frac{\tilde{\chi}_s}{r}$ in $\text{HH}^*(R)$. Note that the elements $\tilde{\xi}_s \Psi_s$ and $\tilde{\chi}_s \Psi_s$ belong to $C^s(R)^\nu$. In addition, if the elements $\tilde{\xi}_s \Psi_s$ and $\tilde{\chi}_s \Psi_s$ are defined for some field k , then they are defined for any field (for the same quiver). As it was said before the formulas for $\Delta(\xi_s) = \frac{\Delta(\tilde{\xi}_s \Psi_s) \Phi_{s-1}}{r}$ and $\Delta(\chi_s) = \frac{\Delta(\tilde{\chi}_s \Psi_s) \Phi_{s-1}}{r}$ are valid for a field k with zero characteristic.

Let us introduce the notion of the standard basis for some modules. The standard basis for Λ is the set $B_\Lambda = \{a \otimes b\}_{a,b \in B_R}$. If $x, y \in \mathcal{Q}_0$, then the standard basis for $P_{[x][y]}$ is the set $B_\Lambda \cap P_{[x][y]}$. And the standard basis for $R^{\otimes t}$ is the set $\{a_1 \otimes \cdots \otimes a_t\}_{a_1, \dots, a_t \in B_R}$. Thus we define the standard basis for Q_t ($t \geq 0$) and $\text{Bar}_t(R)$. Note that in all cases the definition of the standard basis does not depend on the field. If the standard basis is defined for a module M , we denote it by B_M . We denote by $\mathbb{Z}B_M$ the set of linear combinations with integer coefficients of elements from B_M . Note that

- a) if $b \in \mathbb{Z}B_M$ and $a \in B_R$, then $ab, ba \in \mathbb{Z}B_M$ and the coefficients do not depend on the field;
- b) if $a \in \mathbb{Z}B_{Q_t}$, then $d_{t-1}^Q(a) \in \mathbb{Z}B_{Q_{t-1}}$, $D_t(a) \in \mathbb{Z}B_{Q_{t+1}}$ and the coefficients do not depend on the field;
- c) if $a \in B_R$, then $\bar{a} \in B_R$ and $\nu a \in B_R$ and they do not depend on the field;
- d) if $a, b \in \mathbb{Z}B_R$, then $\langle a, b \rangle \in \mathbb{Z}$ and it does not depend on the field.

It follows from a) and b) that the elements of the matrices of Ψ_t and Φ_t in the standard basis are integer and do not depend on the field. It follows from c) and d) that if the matrix of $x : R^{\otimes(n+2)} \rightarrow R$ written in the standard basis consists of integer numbers, then the matrix of $\Delta(x) : R^{\otimes(n+1)} \rightarrow R$ written in the standard basis consists of integer numbers which do not depend on the field. It follows from our arguments that the matrices of $\Delta(\tilde{\xi}_s \Psi_s) \Phi_{s-1} : Q_{s-1} \rightarrow R$ and $\Delta(\tilde{\chi}_s \Psi_s) \Phi_{s-1} : Q_{s-1} \rightarrow R$ written in the standard bases consist of integer numbers which do not depend on the field. If $x \in \{\xi_s, \chi_s\}$, then it follows from [5, Remark 5] that if the set of elements of degree $(n-1)F(x)$ is linear independent in $\text{Ker}(\text{Hom}_\Lambda(d_{s-1}^Q, R))$, then it is linear independent in $\text{HH}^{s-1}(R)$. The formulas for $\Delta(\tilde{\xi}_s \Psi_s) \Phi_{s-1}$ and $\Delta(\tilde{\chi}_s \Psi_s) \Phi_{s-1}$ are written in terms of elements of $\text{Ker}(\text{Hom}_\Lambda(d_{s-1}^Q, R))$ whose definitions do not depend on the field. So these formulas remain true for any field.

2) Since the elements χ_s and ξ_s appear only in the case $\text{CMD}(n-1, r) = 1$, they can be excluded from the set of generators in this case. Let $f, g \in \{f_s, g_s, h_s, p_s\}$. Then there are such $x, y \in \text{HH}^*(R)^{\nu\uparrow}$ that $\Theta_R^\nu(x) = f$ and $\Theta_R^\nu(y) = g$. Since Θ_R^ν is a homomorphism of Gerstenhaber algebras we have $[f, g] = \Theta_R^\nu([x, y])$. Since $\text{char } k | r$ we have $\text{char } k \neq 2$ or $2|r$. It is easy to check that in both cases elements f and g have even degree. In addition it follows from the proof of [12, Lemma 3] and the formula (4.2) that if $\text{char } k \neq 2$ or $2|r$, then for any $a \in \text{Ker}(\text{Hom}_\Lambda(d_{2s+1}^Q, R))$ such that $a = \nu^{-1} a \theta_{2s+1}$ there is $\bar{a} \in \text{Ker}(\text{Hom}_\Lambda(d_{2s+1}^Q, R))$ such that $a = \sum_{i=0}^{r-1} \nu^{-i} \bar{a} \theta_{2s+1}^i$. Since $\bar{a}^\nu = \bar{a}$ in $\text{HH}^*(R)$ by Corollary 2 of Lemma 1 we have

$$a = \sum_{i=0}^{r-1} \nu^{-i} \bar{a} \theta_{2s+1}^i = \sum_{i=0}^{r-1} \bar{a}^{\nu^i} = r\bar{a} = 0$$

in $\text{HH}^{2s+1}(R)$. Consequently, $\Theta_R^\nu(\text{HH}^s(R)^{\nu\uparrow}) = 0$ for odd s . The element $[x, y]$ has odd degree because elements x and y have even degree. Then $[f, g] = \Theta_R^\nu([x, y]) = 0$ and 2) is proved. \square

Remark 1. It is easy to show that we can introduce the BV-structure on $\mathrm{HH}^*(R)$ in the case where $\mathrm{char} k|r$. For example, we can set Δ equal to 0 on all generators of $\mathrm{HH}^*(R)$.

References

- [1] M. Gerstenhaber, *The cohomology structure of an associative ring.* — Ann. Math. (2), 1963, 78, 267–288.
- [2] T. Tradler, *The Batalin–Vilkovisky algebra on Hochschild cohomology induced by infinity inner products.* — Ann. Inst. Fourier, 2008, 58, no. 7, 2351–2379.
- [3] C. H. Eu, T. Schedler, *Calabi–Yau Frobenius algebras.* — J. Algebra, 2009, 321, no. 3, 774–815.
- [4] V. Ginzburg, *Calabi–Yau algebras.* — arXiv:math/0612139.
- [5] Yu. V. Volkov, *The Hochschild cohomology algebra for a family of self-injective algebras of the tree class D_n .* — Algebra i Analiz, 2011, 23, no. 5, 99–139.
- [6] Y. G. Xu, C. Zhang, *Gerstenhaber brackets for truncated quiver algebras.* — Scientia Sinica Mathematica, 2011, 41, no. 1, 17–32.
- [7] C. H. Eu, *The calculus structure of the Hochschild homology/cohomology of preprojective algebras of Dynkin quivers.* — Journal of Pure and Applied Algebra, 2010, 214, 28–46.
- [8] C. H. Eu, *Hochschild and cyclic (co)homology of preprojective algebras of quivers of type T .* — arXiv:0710.4176.
- [9] A. Ivanov, S. Ivanov, Y. Volkov, G. Zhou, *On the Hochschild cohomology ring of the quaternion group of order eight in characteristic two.* — to appear.
- [10] Yu. V. Volkov, *Hochschild cohomology for self-injective algebras of tree class D_n .* VI. — Zap. Nauchn. Semin. POMI, 2014, preprint.
- [11] J. A. Guccione, J. J. Guccione, *Hochschild cohomology of Frobenius algebras.* — Proc. Amer. Math. Soc., 2004, 132, no. 5, 1241–1250.
- [12] Yu. V. Volkov, A. I. Generalov, *Hochschild cohomology for self-injective algebras of tree class D_n .* I. — Zap. Nauchn. Semin. POMI, 2007, 343, 121–182.