

SPECTRA OF SOME WEIGHTED COMPOSITION OPERATORS ON H^2

CARL C. COWEN, EUNGIL KO, DEREK THOMPSON, AND FENG TIAN

ABSTRACT. We completely characterize the spectrum of a weighted composition operator $W_{\psi,\varphi}$ on $H^2(\mathbb{D})$ when φ has Denjoy-Wolff point a with $0 < |\varphi'(a)| < 1$, the iterates, φ_n , converge uniformly to a , and ψ is in H^∞ and continuous at a . We also give bounds and some computations when $|a| = 1$ and $\varphi'(a) = 1$ and, in addition, show that these symbols include all linear fractional φ that are hyperbolic and parabolic non-automorphisms. Finally, we use these results to eliminate possible weights ψ so that $W_{\psi,\varphi}$ is seminormal.

1. INTRODUCTION

If ψ is in H^∞ and φ is analytic map of the unit disk into itself, the *weighted composition operator on H^2 with symbols ψ and φ* is the operator $W_{\psi,\varphi}$, where T_ψ is the analytic Toeplitz operator given by $T_\psi(h) = \psi h$ for h in H^2 , C_φ is the composition operator on H^2 given by $C_\varphi(h) = h \circ \varphi$. Clearly, if ψ is bounded on the disk, then $W_{\psi,\varphi}$ is bounded on H^2 and $\|W_{\psi,\varphi}\| \leq \|\psi\|_\infty \|C_\varphi\|$. Although it will have little impact on our work, it is not necessary for ψ to be bounded for $W_{\psi,\varphi}$ to be bounded. To avoid trivialities and special cases, we will assume ψ is not identically zero and φ is not a constant mapping.

Weighted composition operators have been studied occasionally over the past few decades, but have usually arisen in answering other questions related to operators on spaces of analytic functions, such as questions about multiplication operators or composition operators. For example, Forelli [11] showed that the only isometries of H^p for $p \neq 2$ are weighted composition operators and that the isometries for H^p with $p \neq 2$ have analogues that are isometries of H^2 (but there are also many other isometries of H^2). Weighted composition operators also arise in the description of commutants of analytic Toeplitz operators (see for example [4, 5] and in the adjoints of composition operators (see for example [6, 7, 8]).

Recently, work has begun on studying the spectrum of weighted composition operators on H^2 more carefully. Gunatillake [12] characterized the spectrum when φ has an interior fixed point and $W_{\psi,\varphi}$ is compact. The first two authors [9] characterized the spectrum when $W_{\psi,\varphi}$ is a self-adjoint operator. Bourdon and Narayan extended their work [2] to characterize the spectrum when $W_{\psi,\varphi}$ is unitary and when $W_{\psi,\varphi}$ is normal with interior fixed point. Gunatillake [13] defined invertible weighted composition operators on H^2 and identified their spectrum. Very recently, Hyvärinen, Lindström, Nieminen, and Saukko [14] extended his work to when φ is an automorphism but $W_{\psi,\varphi}$ is not necessarily invertible.

Our work finds the spectra of $W_{\psi,\varphi}$ with relatively weak conditions on ψ , but a rather strong one on φ , which is that the iterates of φ converge uniformly on *all* of \mathbb{D} to the Denjoy-Wolff point a , rather than just on compact subsets of \mathbb{D} . In Section 2, we identify situations when φ satisfies

Date: 16 May 2014.

2010 Mathematics Subject Classification. Primary: 47B33, 47B35; Secondary: 47A10, 47B20, 47B38.

Key words and phrases. weighted composition operator, spectrum of an operator, hyponormal operator.

The second author was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) grant funded by the Ministry of Education, Science and Technology (2012R1A2A2A02008590).

this uniformity condition on the convergence of its iterates, and show that this class of symbols is non-trivial. In Section 3, we give general bounds for $\sigma(W_{\psi,\varphi})$ that define the spectrum when $\sigma(C_\varphi)$ is given by the closure of $\sigma_p(C_\varphi)$. In Section 4, we are much more specific about $\sigma_p(W_{\psi,\varphi})$ when $\varphi'(a) < 1$ and give some examples. In Section 5, we eliminate some possibilities where $W_{\psi,\varphi}$ could be seminormal. Finally, we suggest further areas of study in Section 6.

2. WHEN ARE THE ITERATES OF φ UNIFORMLY CONVERGENT?

To accomplish the work of this paper, we make a rather strong assumption that φ converges uniformly on all of \mathbb{D} to the Denjoy-Wolff point a . Our work in this section will further explain when this phenomenon occurs. To facilitate reference to the property of uniform convergence of the iterates of φ , we make the following definition.

Definition We say *UCI holds for φ* or *φ satisfies UCI* if φ is an analytic map of the unit disk \mathbb{D} into itself with Denjoy-Wolff point a and the iterates φ_n of φ converge uniformly, on all of \mathbb{D} , to a .

We begin by showing that this condition is not particularly helpful when the Denjoy-Wolff point a belongs to \mathbb{D} .

Theorem 1. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic and continuous on $\partial\mathbb{D}$. If the Denjoy-Wolff point a of φ is in \mathbb{D} , then $\varphi_n \rightarrow a$ uniformly if and only if there is $N > 0$ such that $\varphi_N(\overline{\mathbb{D}}) \subseteq \mathbb{D}$.*

Proof. Suppose there is $N > 0$ such that $\varphi_N(\overline{\mathbb{D}}) \subseteq \mathbb{D}$. Since φ_n always converges uniformly on compact subsets of \mathbb{D} to a by the Denjoy-Wolff Theorem [7] and $\varphi_N(\overline{\mathbb{D}})$ is a compact subset of \mathbb{D} , we have that $\varphi_n \rightarrow a$ uniformly on \mathbb{D} .

To prove the other direction, let M be the minimum distance between a and the unit circle. Since $\varphi_n \rightarrow a$ uniformly on \mathbb{D} , for $\epsilon = M/2$, there exists $N > 0$ such that $|\varphi_N(z) - a| < \epsilon, \forall z \in \mathbb{D}$. Suppose $\varphi_N(b_1) = b_2, |b_1| = |b_2| = 1$. Then for our given ϵ , since φ_N is continuous on the unit circle, there exists $\delta > 0$ so that $|b_1 - z| < \delta \Rightarrow |b_2 - \varphi_N(z)| < \epsilon$. However, for z such that $|b_1 - z| < \delta$,

$$\begin{aligned} M &\leq |b_2 - a| = |b_2 - \varphi_N(z) + \varphi_N(z) - a| \\ &\leq |b_2 - \varphi_N(z)| + |\varphi_N(z) - a| \\ &< 2\epsilon = M \end{aligned}$$

which is a contradiction, so $\varphi_N(\overline{\mathbb{D}}) \subseteq \mathbb{D}$. □

The following corollary shows that this is of interest.

Corollary 2. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic and continuous on $\partial\mathbb{D}$. If the Denjoy-Wolff point a of φ is in \mathbb{D} and $\varphi_n \rightarrow a$ uniformly, then C_φ is power-compact. Furthermore, any associated weighted composition operator $W_{\psi,\varphi}$ with $\psi \in H^\infty$ is power-compact.*

Proof. Since $\varphi(\overline{\mathbb{D}}) \subseteq \mathbb{D}$ is a sufficient condition for C_φ to be compact [7], we see that by Theorem 1 $C_{\varphi_N} = C_\varphi^N$ is compact for some $N > 0$ and C_φ is power-compact. Since compact operators are an ideal in $\mathcal{B}(H^2)$, $(W_{\psi,\varphi})^N = T_\zeta C_{\varphi_N}$ is compact, where $\zeta = \psi(\psi \circ \varphi) \dots (\psi \circ \varphi_{N-1})$. □

Since Gunatillake [12] and others have already characterized the spectrum of compact weighted composition operators (and therefore power compact weighted composition operators) when φ has an interior fixed point, we will instead turn our work to when the Denjoy-Wolff point is on $\partial\mathbb{D}$, although our results will include the interior fixed point case. Next, we indicate some conditions on φ when the Denjoy-Wolff point is on $\partial\mathbb{D}$, and give some examples.

Theorem 3. *If $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic in \mathbb{D} and continuous on $\partial\mathbb{D}$, has Denjoy-Wolff point a with $|a| = 1$ and $\varphi_n \rightarrow a$ uniformly, then a is the only fixed point of φ in the closed unit disk.*

Proof. Suppose $\varphi(b) = b$, $b \neq a$. Since the Denjoy-Wolff point is on the boundary, we must have $|b| = 1$, or else b would be the Denjoy-Wolff point. Since $\varphi_n \rightarrow a$ uniformly, given $\epsilon > 0$, there is an N such that $|\varphi_N(z) - a| < \epsilon$, $\forall z \in \mathbb{D}$. Note that φ_N is continuous at b . For the same ϵ , there is δ such that $|b - z| < \delta \Rightarrow |\varphi_N(b) - \varphi_N(z)| = |b - \varphi_N(z)| < \epsilon$. Let z be such that $|b - z| < \delta$. Then

$$|b - a| = |b - \varphi_N(z) + \varphi_N(z) - a| \leq |b - \varphi_N(z)| + |\varphi_N(z) - a| < 2\epsilon$$

However, if we take $\epsilon < |b - a|/2$, we have a contradiction. \square

Although our work so far indicates that the class of weighted composition operators where φ satisfies UCI may be small, we now give sufficient conditions for φ to satisfy UCI and follow with some examples. Much of the following proof is owed to [1].

Theorem 4. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic in \mathbb{D} and continuous on $\partial\mathbb{D}$ and has Denjoy-Wolff point a with $|a| = 1$, $\varphi'(a) < 1$. If $\varphi_N(\overline{\mathbb{D}}) \subseteq \mathbb{D} \cup \{a\}$ for some N , then $\varphi_n \rightarrow a$ uniformly in \mathbb{D} .*

Proof. Without loss of generality, we will assume $\varphi(\overline{\mathbb{D}}) \subseteq \mathbb{D} \cup \{a\}$. Since $\varphi(\overline{\mathbb{D}}) \subseteq \mathbb{D} \cup \{a\}$ and $\varphi(\overline{\mathbb{D}})$ is connected, it fits within the disk

$$H(a, \lambda) := \left\{ z \in \mathbb{C} : |a - z|^2 \leq \lambda(1 - |z|^2) \right\}$$

for some fixed $\lambda > 0$. Disks of this type are Euclidean subdiscs of \mathbb{D} centered at $a/(1 + \lambda)$ with radius $\lambda/(1 + \lambda)$, and are tangent to the unit circle at a . Julia's Lemma [7] shows that $\varphi(H(a, \lambda)) \subseteq H(a, \varphi'(a)\lambda)$. Applying φ iteratively, we see that for any z in this set, we have

$$|a - \varphi_n(z)|^2 \leq \varphi'(a)^n \lambda (1 - |\varphi_n(z)|^2)$$

and therefore

$$|a - \varphi_n(z)| \leq \sqrt{\lambda} \varphi'(a)^{n/2} (1 - |\varphi_n(z)|) \leq \sqrt{\lambda} \varphi'(a)^{n/2}$$

Thus, for any $\epsilon > 0$, there is $N > 0$ such that for $n > N$, $\sqrt{\lambda} \varphi'(a)^{n/2} < \epsilon$ (since $\varphi'(a) < 1$). Then $|\varphi_n(z) - a| \leq \sqrt{\lambda} \varphi'(a)^{n/2} < \epsilon$ for $n > N$. \square

Although this does not completely characterize UCI holding for φ when $|a| = 1$ and $\varphi'(a) < 1$, we see from this sufficient condition that this class includes, at least, φ that are linear fractional non-automorphisms, such as $\varphi(z) = \frac{1}{2}z + \frac{1}{2}$. When $\varphi'(a) = 1$, the situation is even more delicate because the conditions above are not sufficient, as can be seen when φ is a parabolic automorphism. However, if φ is a linear fractional non-automorphism with $\varphi'(a) = 1$, we see that φ actually satisfies UCI:

Example 5. *Let φ be a linear fractional map, not an automorphism, with Denjoy Wolff point a such that $|a| = 1$ and $\varphi'(a) = 1$. Without loss of generality, assume $a = 1$. Such symbols form a*

semigroup $\varphi_t(z) = \frac{t+(2-t)z}{(2+t)-tz}$. Then we have

$$\begin{aligned} |\varphi_t(z) - 1| &= \left| \frac{t + (2-t)z}{(2+t) - tz} - \frac{(2+t) - tz}{(2+t) - tz} \right| \\ &= \left| \frac{2z - 2}{(2+t) - tz} \right| \\ &= \left| \frac{2(z-1)}{2+t(1-z)} \right| \\ &= \left| \frac{2(1-z)}{2+t(1-z)} \right| \\ &= \left| \frac{2}{\frac{2}{1-z} + t} \right| \\ &\leq \left| \frac{2}{t} \right| = \frac{2}{t} \rightarrow 0, \text{ as } t \rightarrow \infty \end{aligned}$$

since $\operatorname{Re} \left\{ \frac{2}{1-z} \right\} > 1$ for $z \in \mathbb{D}$. Thus if φ is a linear fractional non-automorphism with Denjoy-Wolff point a and $\varphi'(a) = 1$, then $\varphi_n \rightarrow a$ uniformly in \mathbb{D} .

Now we see that UCI holding for φ can arise when $\varphi'(a) < 1$ or $\varphi'(a) = 1$. Next, we show general bounds for the spectra of a weighted composition operator with UCI holding for the compositional symbol, and later we discuss the differences more specifically between the two cases.

3. SPECTRAL BOUNDS FOR $W_{\psi, \varphi}$

Throughout the remainder of the paper, we will assume that ψ is in H^∞ , continuous at the Denjoy-Wolff point a of φ , and that $\psi(a) \neq 0$.

We now offer some lemmas which will give us an inequality between the spectra of $W_{\psi, \varphi}$ and $\psi(a)C_\varphi$.

Lemma 6. *If A and B are bounded operators on a Hilbert space \mathcal{H} , then:*

- (1) *If $ABv = \lambda v$ and $Bv \neq 0$, then Bv is an eigenvector for BA with eigenvalue λ .*
- (2) $\sigma(AB) \cup \{0\} = \sigma(BA) \cup \{0\}$.

Proof. (1) If $ABv = \lambda v$ for some λ , then $BA(Bv) = B(ABv) = B(\lambda v) = \lambda(Bv)$, but we require $Bv \neq 0$ since eigenvectors need to be non-zero.

(2) It can be seen by direct computation that if $\lambda \neq 0$ and $(\lambda I - AB)$ is invertible, then $(\lambda I - BA)$ is invertible with inverse $\frac{1}{\lambda}(I + B(\lambda I - AB)^{-1}A)$. Clearly this argument also works if A and B are reversed, so $(\lambda I - AB)$ is invertible if and only if $(\lambda I - BA)$ is. Since we required $\lambda \neq 0$, we have $\sigma(AB) \cup \{0\} = \sigma(BA) \cup \{0\}$. \square

Although part (1) of Lemma 6 requires that $Bv \neq 0$, we will only be using analytic Toeplitz operators and composition operators with trivial kernels when we apply the lemma.

Lemma 7. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $\varphi_n \rightarrow a$ uniformly in \mathbb{D} , and $\psi \in H^\infty$ is continuous at $z = a$. Then $\|T_{\psi(a)}C_\varphi - T_{\psi \circ \varphi_n}C_\varphi\| \rightarrow 0$ in $\mathcal{B}(H^2)$ as $n \rightarrow \infty$.*

Proof. If $\varphi_n \rightarrow a$ uniformly in \mathbb{D} , and ψ is continuous at a , then $\psi \circ \varphi_n \rightarrow \psi(a)$ uniformly in D , which implies that $\|\psi(a) - \psi \circ \varphi_n\|_\infty \rightarrow 0$ as $n \rightarrow \infty$. Then

$$\begin{aligned} \|T_{\psi(a)}C_\varphi - T_{\psi \circ \varphi_n}C_\varphi\| &\leq \|T_{\psi(a) - \psi \circ \varphi_n}\| \|C_\varphi\| \\ &= \|\psi(a) - \psi \circ \varphi_n\|_\infty \|C_\varphi\| \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$, since C_φ is bounded. \square

Theorem 8. *If $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $\varphi_n \rightarrow a$ uniformly in \mathbb{D} , and $\psi \in H^\infty$ is continuous at $z = a$, then $\sigma(T_\psi C_\varphi) \subseteq \sigma(\psi(a)C_\varphi)$.*

Proof. Note that $(T_\psi C_\varphi - \lambda I)$ is invertible if and only if $(C_\varphi T_\psi - \lambda I) = (T_{\psi \circ \varphi} C_\varphi - \lambda I)$ is invertible by Lemma 6. Applying this iteratively, we see that $(T_\psi C_\varphi - \lambda I)$ if and only if $(T_{\psi \circ \varphi_n} C_\varphi - \lambda I)$ is invertible for all n .

Let $\lambda \in \sigma(T_\psi C_\varphi)$. Then $\lambda \in \sigma(T_{\psi \circ \varphi_n} C_\varphi)$ for all n by above. By Lemma 7, the operators $(T_{\psi \circ \varphi_n} C_\varphi - \lambda I)$ converge to $(T_{\psi(a)} C_\varphi - \lambda I)$ in H^2 norm. Since the invertible operators in $\mathcal{B}(H^2)$ are an open set and each operator in the sequence is not invertible, we know that $(T_{\psi(a)} C_\varphi - \lambda I)$ is also not invertible, so $\lambda \in \sigma(\psi(a)C_\varphi)$. \square

Given the theorem above, it is seen that we assume $\psi(a) \neq 0$ simply to avoid trivial cases where $\sigma(W_{\psi, \varphi}) = \{0\}$. Our next goal is to find a lower bound on the spectra of $W_{\psi, \varphi}$ and use a squeeze-type argument. The following theorems will accomplish that.

Theorem 9. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $\varphi_n \rightarrow a$ uniformly in \mathbb{D} , and $\psi \in H^\infty$ is continuous at $z = a$ with $\psi(a) \neq 0$. Then $\psi(a)$ is in $\sigma_{ap}(T_\psi C_\varphi)$.*

Proof. Let

$$h_m(z) = \prod_{n=0}^m \frac{\psi \circ \varphi_n}{\psi(a)}$$

Note that $T_\psi C_\varphi h_m(z) = \psi(a)h_{m+1}(z)$. These vectors are finite products of H^∞ functions, so they belong in H^∞ and therefore to H^2 as well. The unit vectors we will use will be $H_m = \frac{h_m}{\|h_m\|_2}$. We wish to show $\|(T_\psi C_\varphi - \psi(a)I)H_m\|_2 \rightarrow 0$ as $m \rightarrow \infty$.

$$\begin{aligned} \|(T_\psi C_\varphi - \psi(a)I)H_m\|_2 &= \frac{1}{\|h_m\|_2} \|(T_\psi C_\varphi - \psi(a)I)h_m\|_2 \\ &= \frac{1}{\|h_m\|_2} \|\psi(a)h_{m+1} - \psi(a)h_m\|_2 \\ &= \frac{1}{\|h_m\|_2} \|h_m \psi \circ \varphi_{m+1}(z) - \psi(a)h_m\|_2 \\ &= \frac{1}{\|h_m\|_2} \|h_m (\psi \circ \varphi_{m+1}(z) - \psi(a))\|_2 \\ &\leq \frac{1}{\|h_m\|_2} \|h_m\|_2 \|\psi \circ \varphi_{m+1}(z) - \psi(a)\|_\infty \\ &= \|\psi \circ \varphi_{m+1}(z) - \psi(a)\|_\infty \rightarrow 0; \end{aligned}$$

and the last line is by UCI holding for φ and continuity of ψ . \square

Theorem 10. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $\varphi_n \rightarrow a$ uniformly in D , and $\psi \in H^\infty$ is continuous at $z = a$ with $\psi(a) \neq 0$. Then for any eigenvalue λ of C_φ , $\psi(a)\lambda \in \sigma_{ap}(T_\psi C_\varphi)$. In particular, $\sigma_p(\psi(a)C_\varphi) \subseteq \sigma_{ap}(T_\psi C_\varphi)$.*

Proof. Let h_m be defined as in Theorem 9, and let g be an eigenvector for C_φ with eigenvalue λ . Then since the vectors h_m are all in H^∞ , $g_m = gh_m$ are all in H^2 . As before, we have

$$\begin{aligned} T_\psi C_\varphi g_m &= T_\psi C_\varphi h_m g \\ &= (\psi h_m \circ \varphi) (g \circ \varphi) \\ &= (\psi(a) h_{m+1}) (\lambda g) \\ &= \psi(a) \lambda h_{m+1} g \\ &= \psi(a) \lambda g_{m+1}. \end{aligned}$$

Let $G_m = \frac{g_m}{\|g_m\|_2}$. Then we have

$$\begin{aligned} \|(T_\psi C_\varphi - \psi(a)\lambda I) G_m\|_2 &= \frac{1}{\|g_m\|_2} \|(T_\psi C_\varphi - \psi(a)\lambda I) g_m\|_2 \\ &= \frac{1}{\|g_m\|_2} \|T_\psi C_\varphi g_m - \psi(a)\lambda g_m\|_2 \\ &= \frac{1}{\|g_m\|_2} \|\psi(a)\lambda g_{m+1} - \psi(a)\lambda g_m\|_2 \\ &= \frac{1}{\|g_m\|_2} \left\| \psi(a)\lambda g_m \left(\frac{\psi \circ \varphi_{m+1}}{\psi(a)} \right) - \psi(a)\lambda g_m \right\|_2 \\ &= \frac{1}{\|g_m\|_2} \|\lambda g_m (\psi \circ \varphi_{m+1} - \psi(a))\|_2 \\ &\leq \frac{1}{\|g_m\|_2} \|\lambda g_m\|_2 \|\psi \circ \varphi_{m+1} - \psi(a)\|_\infty \\ &= |\lambda| \|\psi \circ \varphi_{m+1} - \psi(a)\|_\infty \rightarrow 0. \end{aligned}$$

The last line is again by UCI holding for φ and continuity of ψ . Since this is true for any eigenvalue λ of C_φ , we have $\sigma_p(\psi(a)C_\varphi) \subseteq \sigma_{ap}(T_\psi C_\varphi)$. \square

Taking this together with Theorem 8, we get the following string of inequalities:

Corollary 11. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $\varphi_n \rightarrow a$ uniformly in \mathbb{D} , and $\psi \in H^\infty$ is continuous at $z = a$ with $\psi(a) \neq 0$. Then we have*

$$\overline{\sigma_p(\psi(a)C_\varphi)} \subseteq \overline{\sigma_{ap}(T_\psi C_\varphi)} \subseteq \sigma(T_\psi C_\varphi) \subseteq \sigma(\psi(a)C_\varphi)$$

In particular, if $\overline{\sigma_p(C_\varphi)} = \sigma(C_\varphi)$, then $\sigma(T_\psi C_\varphi) = \sigma(\psi(a)C_\varphi)$.

Proof. The first containment is given by Theorem 10; the second containment is trivial; the third containment is given by Theorem 8. \square

As a consequence of this corollary, we can define the spectrum in the case where $\varphi'(a) < 1$, and give some examples where $\varphi'(a) = 1$.

Corollary 12. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $\varphi_n \rightarrow a$ uniformly in \mathbb{D} , and $\varphi'(a) < 1$. Then for any $\psi \in H^\infty$ continuous at $z = a$ with $\psi(a) \neq 0$, $\sigma(W_{\psi, \varphi}) = \sigma(\psi(a)C_\varphi)$.*

Proof. When the Denjoy-Wolff point of φ is on the boundary with $\varphi'(a) < 1$ and a is the only fixed point of φ , then every point in the spectrum except for 0 and the peripheral spectrum is an eigenvalue of infinite multiplicity [7]. Thus $\overline{\sigma_p(C_\varphi)} = \sigma(C_\varphi)$ and $\sigma(W_{\psi, \varphi}) = \sigma(\psi(a)C_\varphi)$ by Corollary 11. \square

Example 13. If $\varphi(z) = \frac{1}{2-z}$ so that $\varphi(1) = 1, \varphi'(1) = 1$, then it is known that C_φ has spectrum $[0, 1]$ and point spectrum $(0, 1)$ [7]. Since $\overline{\sigma_p(C_\varphi)} = \sigma(C_\varphi)$, we have $\sigma(W_{\psi, \varphi}) = \sigma(\psi(a)C_\varphi)$ by Corollary 11, for any $\psi \in H^\infty$ continuous at $z = 1$ with $\psi(1) \neq 0$.

So far, our work in this section has not taken the value of $\varphi'(a)$ into account until the corollary above. When $\varphi'(a) < 1$, we can actually be much more specific about the point spectrum, which we will do in the next section.

4. POINT SPECTRA OF $W_{\psi, \varphi}$ WHEN $\varphi'(a) < 1$

For this section, our goal is to show that except for 0 and the peripheral spectrum, $\sigma(W_{\psi, \varphi})$ otherwise consists entirely of eigenvalues when $\varphi'(a) < 1$. We accomplish this by extending the vector in the proof of Theorem 9 to an infinite series bounded by $\varphi'(a)^n$.

Theorem 14. Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $0 < |\varphi'(a)| < 1$, and $\varphi_n \rightarrow a$ uniformly in \mathbb{D} . Then for any ψ in H^∞ that is bounded away from zero on \mathbb{D} and continuous at a , there is an eigenvector h for $T_\psi C_\varphi$ with eigenvalue $\psi(a)$ and h invertible in H^∞ .

Proof. Since ψ is a bounded, analytic, and non-vanishing map on \mathbb{D} , we may assume that there exists a bounded analytic map η so that $\psi = e^\eta$. Since η is analytic and bounded on \mathbb{D} , it has bounded derivative there, so η is Lipschitz on \mathbb{D} , i.e. $|\eta(z_1) - \eta(z_2)| \leq K|z_1 - z_2|$ for $z_1, z_2 \in \mathbb{D}$ and some constant \tilde{K} independent of z_1, z_2 . Since η is continuous at a , it can be seen that the above inequality holds on $\mathbb{D} \cup \{a\}$. Additionally, since φ_n converges uniformly on \mathbb{D} , $|\varphi_n(z) - a| \leq K\varphi'(a)^n$ for some constant K independent of z , as seen in the proof of Theorem 4 above. Since

$$\lim_{n \rightarrow \infty} \eta \circ \varphi_n = \eta(a),$$

we want to show that $\sum_{n=0}^{\infty} (\eta \circ \varphi_n - \eta(a))$ converges in H^∞ . Since

$$|\eta(\varphi_n(z)) - \eta(a)| \leq \tilde{K}|\varphi_n(z) - a| \leq K\tilde{K}\varphi'(a)^n$$

and $|\varphi'(a)| < 1$, the series converges. Set

$$g = \sum_{n=0}^{\infty} (\eta \circ \varphi_n - \eta(a)).$$

Then $h(z) = e^{g(z)}$ is an eigenfunction for $W_{\psi, \varphi}$ with eigenvalue $\psi(a)$. Since ψ is bounded below, so is η , and now $g(z)$ is bounded above and below, so $\frac{1}{h} = e^{-g(z)}$ is also in H^∞ . \square

The next theorem shows that the special eigenvector above completely identifies the point spectrum with that of $\psi(a)C_\varphi$.

Theorem 15. If $T_\psi C_\varphi$ has eigenvalue α with an eigenvector $g \in H^\infty$ for α , and λ is any eigenvalue of C_φ with eigenvector f , then $\alpha\lambda$ is an eigenvalue of $T_\psi C_\varphi$ with eigenvector gf . Furthermore, if $\frac{1}{g} \in H^\infty$ as well, then $\sigma_p(\alpha C_\varphi) = \sigma_p(T_\psi C_\varphi)$. In particular, if $\alpha = \psi(a)$, then $\sigma_p(\psi(a)C_\varphi) = \sigma_p(T_\psi C_\varphi)$.

Proof. We have $T_\psi C_\varphi g = \alpha g$ and $C_\varphi f = \lambda f$. Note since $g \in H^\infty, gf \in H^2$. Then

$$T_\psi C_\varphi(gf) = \psi(gf) \circ \varphi = (\psi g \circ \varphi)(f \circ \varphi) = (\alpha g)(\lambda f) = \alpha\lambda(gf)$$

so gf is an eigenfunction for $T_\psi C_\varphi$ with eigenvalue $\alpha\lambda$. So $\sigma_p(\alpha C_\varphi) \subseteq \sigma_p(T_\psi C_\varphi)$.

Now, if $\frac{1}{g} \in H^\infty$ as well, then for any eigenvalue $\mu \in \sigma_p(T_\psi C_\varphi)$ with eigenvector h , we can write $v = \frac{h}{g}$ which is in H^2 , so $gv = h$. Then

$$\mu gv = \mu h = T_\psi C_\varphi h = T_\psi C_\varphi gv = (\psi g \circ \varphi)(v \circ \varphi) = \alpha gv \circ \varphi$$

Dividing the far sides by g , we see that $\mu v = \alpha v \circ \varphi$, so $\mu \in \sigma_p(\alpha C_\varphi)$. Thus $\sigma_p(T_\psi C_\varphi) \subseteq \sigma_p(\alpha C_\varphi)$, so now $\sigma_p(\alpha C_\varphi) = \sigma_p(T_\psi C_\varphi)$. \square

Putting these theorems together, we have the following corollary.

Corollary 16. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $0 < |\varphi'(a)| < 1$, and $\varphi_n \rightarrow a$ uniformly in \mathbb{D} . Then for any $\psi \in H^\infty$ that is bounded away from zero on \mathbb{D} and continuous at a , $\sigma_p(W_{\psi, \varphi}) = \sigma_p(\psi(a)C_\varphi)$.*

Although we required stricter conditions on ψ to achieve the above corollary, we can in fact use UCI holding for φ to relax those conditions:

Corollary 17. *Suppose $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is analytic with Denjoy-Wolff point a , $0 < |\varphi'(a)| < 1$, and $\varphi_n \rightarrow a$ uniformly in \mathbb{D} . Then for any $\psi \in H^\infty$ that is continuous at a with $\psi(a) \neq 0$, $\sigma_p(W_{\psi, \varphi}) = \sigma_p(\psi(a)C_\varphi)$.*

Proof. By Lemma 6, $\sigma_p(T_\psi C_\varphi) = \sigma_p(T_{\psi \circ \varphi_n} C_\varphi)$ for all n . Since ψ is continuous at a and $\psi(a) \neq 0$, there is $\epsilon > 0$ so that $\psi(z)$ is bounded away from zero on the set $\{z : |z - a| < \epsilon\}$. Since $\varphi_n \rightarrow a$ uniformly, there is N such that for $n \geq N$, $|\varphi_n(z) - a| < \epsilon$, for all $z \in \mathbb{D}$. Then $T_{\psi \circ \varphi_n} C_\varphi$ satisfies the conditions of Corollary 16, so $\sigma_p(T_\psi C_\varphi) = \sigma_p(T_{\psi \circ \varphi_n} C_\varphi) = \sigma_p(\psi(a)C_\varphi)$. \square

Since we can now entirely characterize the spectrum and point spectrum when $\varphi'(a) < 1$, we illustrate this with an example below.

Example 18. *Let $\varphi(z) = \frac{1}{2}z + \frac{1}{2}$ and $\psi(z) = e^{(2-z)}$. Note $\varphi_n(z) = \frac{1}{2^n}z + 1 - \frac{1}{2^n}$. Then for η as in the proof of Theorem 14, we have $\eta(z) = 2 - z$ and we can compute $g(z)$ as in Theorem 14:*

$$\begin{aligned} \sum_{n=0}^{\infty} \eta \circ \varphi_n - \eta(1) &= \sum_{n=0}^{\infty} 2 - \left(\frac{1}{2^n}z + 1 - \frac{1}{2^n}\right) - 1 \\ &= \sum_{n=0}^{\infty} \frac{1}{2^n}(1 - z) \\ &= 2 - 2z \end{aligned}$$

Then $h(z) = e^{(2-2z)}$ is an H^∞ eigenvector for $W_{\psi, \varphi}$ with eigenvalue $\psi(1) = e$, as is seen below:

$$\psi h \circ \varphi = e^{(2-z)} e^{(2-2(\frac{1}{2}z + \frac{1}{2}))} = e^{(2-z)} e^{(1-z)} = e^{(1+2-2z)} = e^1 e^{(2-2z)} = eh$$

Note that $\frac{1}{h} = e^{(2z-2)}$ is also in H^∞ . It is known that the functions $(1-z)^\lambda$ are eigenvectors of C_φ with eigenvalue $(\frac{1}{2})^\lambda$, that these belong to H^2 when $\text{Re}(\lambda) > -\frac{1}{2}$, and that $\sigma_p(C_\varphi) = \{\lambda : 0 < \lambda < \sqrt{2}\}$ [7]. Then $\sigma_p(W_{\psi, \varphi}) = \{\lambda : 0 < \lambda < \sqrt{2}e\}$ and $e^{(2-2z)}(1-z)^\lambda$ is an eigenvector for eigenvalue $\frac{e}{2^\lambda}$.

Our work here depended on the fact that $\varphi'(a) < 1$. The following two examples show that an analogous statement cannot be made when $\varphi'(a) = 1$.

Example 19. Let $\varphi(z) = \frac{1}{2-z}$ and $\psi(z) = 2 - z$. Then we see that $h(z) = 1 - z$ is an eigenvector for $W_{\psi,\varphi}$ with eigenvalue 1. It is known that C_φ has spectrum $[0, 1]$ with point spectrum $(0, 1)$ [7]. Since h is in H^∞ , any eigenvector g for an eigenvalue λ of C_φ corresponds to an eigenvector gh of $W_{\psi,\varphi}$ with eigenvalue λ . Thus $W_{\psi,\varphi}$ has spectrum $[0, 1]$ and every element $0 < \lambda \leq 1$ is an eigenvalue.

Example 20. Let $\varphi(z) = \frac{1}{2-z}$ and $\psi(z) = \frac{1}{1-\frac{1}{2}z}$. The first two authors [9] showed that the operator $W_{\psi,\varphi}$ is self-adjoint and has no eigenvalues, but rather consists entirely of approximate point spectrum.

5. SEMINORMALITY OF $W_{\psi,\varphi}$

In [9], the first two authors showed that the semigroup of parabolic non-automorphisms studied in this paper have a companion weight so that $W_{\psi,\varphi}$ is self-adjoint. The form of the companion weight associated with most known self-adjoint [9], normal [2], and cohyponormal [10] weighted composition operators is $\psi = pK_{\sigma(0)}$, where p is a constant and σ is the Cowen auxillary function of φ (which is linear fractional in these situations). As a result of our work above, we eliminate possibilities for ψ when φ is a linear fractional non-automorphism with Denjoy-Wolff point on ∂D and $W_{\psi,\varphi}$ is seminormal.

First, we show that when φ is a parabolic non-automorphism, there are no other weight functions ψ continuous at the Denjoy-Wolff point a so that $W_{\psi,\varphi}$ is (co)hyponormal.

Theorem 21. Let $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ be a parabolic non-automorphism with Denjoy-Wolff point a and let $\psi \in H^\infty$ be continuous at $z = a$. If $W_{\psi,\varphi}$ is (co)hyponormal, then it is normal and ψ is a multiple of $K_{\sigma(0)}$, where σ is the Cowen auxillary function of φ . Furthermore, if $\psi(a)$ is real, then $W_{\psi,\varphi}$ is self-adjoint.

Proof. Without loss of generality, assume $a = 1$ since composition with a rotation is unitary. For now, assume $\psi(a)$ is real. Any (co)hyponormal operator whose spectrum has zero area is normal [15]. Since C_φ has spectrum $[0, 1]$ and point spectrum $(0, 1)$, $W_{\psi,\varphi}$ (and therefore also $W_{\psi,\varphi}^*$) has spectrum equal to the line segment $[0, \psi(a)]$ by Corollary 11. Since line segments have zero area, $W_{\psi,\varphi}$ is normal. Since $W_{\psi,\varphi}$ is (sub)normal and $\sigma(W_{\psi,\varphi}) \subseteq \mathbb{R}$, it is self-adjoint [3]. The self-adjoint weighted composition operators on H^2 have been completely characterized in [9] and ψ must therefore be a real multiple of $K_{\sigma(0)}$.

If $\psi(a)$ is not real, we get the same result for the weight $\lambda\psi$, where λ is a non-zero constant so that $\lambda\psi(a)$ is real. Then we see that ψ must be a (non-real) multiple of $K_{\sigma(0)}$ and that $W_{\psi,\varphi}$ is normal. \square

Next, let φ be a hyperbolic non-automorphism. Here, $W_{\psi,\varphi}$ can be cohyponormal and in fact cosubnormal. For example, if $\varphi(z) = sz + 1 - s, 0 < s < 1$, then $\sigma(0) = 0, K_0 = 1$ and C_φ is a “weighted” composition operator which is cosubnormal. In [2], it is shown that if ψ is in C^1 on $\overline{\mathbb{D}}$ then $W_{\psi,\varphi}$ cannot be essentially normal. Due to our understanding of the spectrum from Section 4 above, we can show that no weight ψ in H^∞ continuous at the Denjoy-Wolff point (but with no other conditions on ψ at the boundary) creates a hyponormal weighted composition operator when φ is a hyperbolic non-automorphism. However, first we need a lemma.

Lemma 22. Let g be a vector in H^2 such that $\langle g, z^n g \rangle = \langle g, g \rangle$ for all integers $n \geq 1$. Then g is the zero vector.

Proof. Suppose that g is not the zero vector. Writing g as $g = \sum_{k=0}^{\infty} a_k z^k$, since g is not the zero vector, not all a_k are zero. Therefore, there is an integer n such that the vector $g_n = \sum_{k=n}^{\infty} a_k z^k$ satisfies $\|g_n\| < \|g\|/2$. Then

$$\|g\|^2 = |\langle g, g \rangle| = |\langle g, z^n g \rangle| = |\langle g_n, g \rangle| \leq \|g_n\| \|g\| < \|g\|^2 / 2$$

which is impossible. Therefore g is the zero vector. \square

Theorem 23. *Let $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ be a hyperbolic non-automorphism. There is no $\psi \in H^\infty$ continuous at $z = a$ such that $W_{\psi, \varphi}$ is hyponormal.*

Proof. Without loss of generality, assume $\varphi(z) = sz + 1 - s$ for some $0 < s < 1$. (Otherwise, conjugate $W_{\psi, \varphi}$ by the unitary weighted composition operator $T_g C_\zeta$ where $g = K_{\zeta(0)}$ and ζ is an automorphism so that $\zeta \circ \varphi \circ \zeta$ is in this form. This will change the weight function ψ , but it will still be continuous at a and it is otherwise arbitrary.) Now assume $W_{\psi, \varphi}$ is hyponormal.

It is known that $(1-z)^n$ is an eigenvector for C_φ with eigenvalue s^n . By Theorem 14, there is an eigenfunction $h \in H^\infty$ for $W_{\psi, \varphi}$ with eigenvalue $\psi(a)$, and thus $h(1-z)^n$ is an eigenfunction for $W_{\psi, \varphi}$ with eigenvalue $\psi(a)s^n$ by Theorem 15. Since $W_{\psi, \varphi}$ is hyponormal, eigenvectors corresponding to different eigenvalues must be perpendicular [3]. Then

$$0 = \langle h, (1-z)h \rangle = \langle h, h \rangle - \langle h, zh \rangle \Rightarrow \langle h, h \rangle = \langle h, zh \rangle$$

Keeping this result in mind, we now consider the vectors h and $(1-z)^2 h$:

$$0 = \langle h, (1-z)^2 h \rangle = \langle h, h \rangle - 2\langle h, zh \rangle + \langle h, z^2 h \rangle \Rightarrow \langle h, h \rangle = \langle h, z^2 h \rangle$$

Continuing inductively, we have $\langle h, h \rangle = \langle h, z^n h \rangle$ for all integers $n > 0$. Therefore, by Lemma 22, h is the 0 vector, which is a contradiction since eigenvectors are non-zero. Therefore $W_{\psi, \varphi}$ cannot be hyponormal. \square

6. FURTHER QUESTIONS

Below is a list of questions that would extend our work:

- (1) Characterize exactly when the iterates φ_n converge uniformly to a on all of \mathbb{D} .
- (2) Completely characterize the point spectrum of $W_{\psi, \varphi}$ when $|a| = 1$, $\varphi'(a) = 1$ and the iterates φ_n converge uniformly to a in all of \mathbb{D} .
- (3) Completely characterize (co)(hypo)normal weighted composition operators on H^2 . (For example, it has *not* been shown that if $W_{\psi, \varphi}$ is normal, φ must be linear fractional.)
- (4) In our work and many of our referenced papers, it seems that when φ has exactly one fixed point a in $\overline{\mathbb{D}}$, that $\sigma(W_{\psi, \varphi}) = \sigma(\psi(a)C_\varphi)$. How often is this true?

REFERENCES

- [1] P. S. Bourdon, V. Matache, and J. H. Shapiro, On convergence to the Denjoy-Wolff point, *Illinois J. Math.* **49**(2005), 405–430.
- [2] P. S. Bourdon and S. Narayan, Normal weighted composition operators on the Hardy space $H^2(U)$, *J. Math. Anal. App.* **367**(2010), 278–286.
- [3] J. B. Conway, *The Theory of Subnormal Operators*, Math. Surveys and Monographs, vol. 36, American Mathematical Society, Providence, RI, 2000.
- [4] C. C. Cowen, The commutant of an analytic Toeplitz operator, *Trans. Amer. Math. Soc.* **239**(1978), 1–31.

- [5] C. C. Cowen, An analytic Toeplitz operator that commutes with a compact operator, *J. Functional Analysis* **36**(1980), 169–184.
- [6] C. C. Cowen, Linear fractional composition operators on H^2 , *Integral Equations Operator Theory* **11**(1988), 151–160.
- [7] C. C. Cowen and B. D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, CRC Press, Boca Raton, 1995.
- [8] C. C. Cowen and E. A. Gallardo Gutierrez, A new class of operators and a description of adjoints of composition operators, *J. Functional Analysis* **238**(2006), 447–462.
- [9] C. C. COWEN and E. KO, Hermitian weighted composition operators on H^2 , *Trans. Amer. Math. Soc.* **362**(2010), 5771–5801.
- [10] C. C. COWEN, S. JUNG, and E. KO, Normal and cohyponormal weighted composition operators on H^2 , to appear: *Operator Theory: Advances and Applications*.
- [11] F. Forelli, The isometries of H^p , *Canadian J. Math.* **16**(1964), 721–728.
- [12] G. Gunatillake, *Weighted Composition Operators*, Thesis, Purdue University, 2005.
- [13] G. Gunatillake, Invertible weighted composition operators, *J. Functional Analysis* **261**(2011), 831–860.
- [14] O. Hyvärinen, M. Lindström, I. Nieminen, and E. Saukko, Spectra of weighted composition operators with automorphic symbols, *J. Functional Analysis* **261**(2011), 831–860.
- [15] M. Martin and M. Putinar, *Lectures on Hyponormal Operators*, Operator Theory: Advances and Applications, vol. 39. Birkhauser Verlag, Basel, 1989.

IUPUI (INDIANA UNIVERSITY – PURDUE UNIVERSITY, INDIANAPOLIS), INDIANAPOLIS, INDIANA 46202-3216
E-mail address: ccowen@math.iupui.edu

EWHA WOMANS UNIVERSITY, SEOUL 120-750, S. KOREA
E-mail address: eiko@ewha.ac.kr

TRINE UNIVERSITY, ANGOLA, INDIANA 46703
E-mail address: theycallmedt@gmail.com
E-mail address: tianf@trine.edu