

STABILITY CONDITIONS ON CY_N CATEGORIES ASSOCIATED TO A_n -QUIVERS AND PERIOD MAPS

AKISHI IKEDA

ABSTRACT. In this paper, we study the space of stability conditions on a certain N -Calabi-Yau (CY_N) category associated to an A_n -quiver. Recently, Bridgeland and Smith constructed stability conditions on some CY_3 categories from meromorphic quadratic differentials with simple zeros. Generalizing their results to higher dimensional Calabi-Yau categories, we describe the space of stability conditions as the universal cover of the space of polynomials of degree $n+1$ with simple zeros. In particular, central charges of stability conditions on CY_N categories are constructed as the periods of quadratic differentials with zeros of order $N-2$ which are associated to polynomials.

1. INTRODUCTION

The notion of a stability condition on a triangulated category \mathcal{D} was introduced by Bridgeland [Bri07] following Douglas's work on Π -stability [Dou02]. Bridgeland also showed that the space of stability conditions $\text{Stab}(\mathcal{D})$ has the structure of a complex manifold.

In this paper, we study the space of stability conditions on a certain N -Calabi-Yau (CY_N) category \mathcal{D}_n^N associated to an A_n -quiver [ST01, Tho06]. The category \mathcal{D}_n^N is defined by the derived category of finite dimensional dg modules over the Ginzburg CY_N dg algebra [Gin] of the A_n -quiver.

We show that the distinguished connected component $\text{Stab}^\circ(\mathcal{D}_n^N) \subset \text{Stab}(\mathcal{D}_n^N)$ is isomorphic to the universal cover of the space of polynomials $p_n(z) = z^{n+1} + u_1 z^{n-1} + \cdots + u_n$ ($u_i \in \mathbb{C}$) with simple zeros. The central charges of stability conditions in $\text{Stab}^\circ(\mathcal{D}_n^N)$ are constructed as the periods of quadratic differentials of the form $p_n(z)^{N-2} dz^{\otimes 2}$ on the Riemann sphere \mathbb{P}^1 .

To prove the result, we use the recent results of Bridgeland and Smith [BS15]. Motivated by the work of physicists Gaiotto-Moore-Neitzke [GMN], they constructed stability conditions on a class of CY_3 categories associated to triangulations of marked bordered surfaces from quadratic differentials with simple zeros.

By considering quadratic differentials with zeros of order $N-2$ instead of simple zeros, we can generalize the framework of [BS15] for CY_3 categories to CY_N categories. Here we only treat the simplest case that the surface is a disk with some marked points on the boundary. In the future, we expect that the framework of [BS15] can be generalized to a class of CY_N categories associated to N -angulations of marked bordered surfaces, though these categories are not (well) defined yet.

1.1. Background geometry. Let $N \geq 2$ be an integer. Here we review the geometric background for this paper and explain why we consider the periods of the differential form $p_n(z)^{\frac{N-2}{2}} dz$. Section 2 in [Tho06] is an excellent survey of related topics.

The category \mathcal{D}_n^N was first introduced as the derived Fukaya category [KS02, Tho06] of Lagrangian N -spheres which form the basis of vanishing cycles in the universal deformation of the A_n -singularity

$$X_p^N := \{x_1^2 + x_2^2 + \cdots + x_N^2 = p_n(z)\} \subset \mathbb{C}^N \times \mathbb{C}$$

where $p_n(z) = z^{n+1} + u_1 z^{n-1} + \cdots + u_n$ ($u_i \in \mathbb{C}$) with simple zeros. We consider a symplectic structure on X_p^N coming from the standard Kähler form on $\mathbb{C}^N \times \mathbb{C}$, and a nowhere vanishing holomorphic N -form Ω_p on X_p^N given by taking Poincaré residue of the standard form $dx_1 \cdots dx_N dz$ on \mathbb{C}^{N+1} .

Consider the projection

$$\pi: X_p^N \rightarrow \mathbb{C}, \quad (x_1, \dots, x_N, z) \mapsto z.$$

If $z \in \mathbb{C}$ is not a root of p_n , then the fiber $\pi^{-1}(z)$ is an affine quadric of complex dimension $N-1$ with a real slice $S^{N-1} \subset \pi^{-1}(z)$. Thus, each Lagrangian N -sphere is constructed by the total space of the S^{N-1} -fibration over a path $\gamma: [0, 1] \rightarrow \mathbb{C}$ which connects distinct roots of p_n (see [TY02, Section 6] or [Tho06, Section 2]).

As in [Dou02], a central charge “in the physics literature” of a Lagrangian $L \subset X_p^N$ is given by the period

$$\int_L \Omega_p.$$

Assume that L is a S^{N-1} -fibration over the path $\gamma: [0, 1] \rightarrow \mathbb{C}$. Then, up to constant multiplication, the above integration is reduced to the integration of the differential form $p_n(z)^{\frac{N-2}{2}} dz$ along the path γ (see [TY02, Section 6]):

$$\int_\gamma p_n(z)^{\frac{N-2}{2}} dz.$$

From this observation, we expect that the central charges of Bridgeland stability conditions on \mathcal{D}_n^N can be constructed by using the period of the differential form $p_n(z)^{\frac{N-2}{2}} dz$.

1.2. Summary of results. First we explain the space of polynomials and the associated period map (Section 2). Let M_n be the space of polynomials $p_n(z) = z^{n+1} + u_1 z^{n-1} + \cdots + u_n$ ($u_i \in \mathbb{C}$) with $n+1$ simple zeros. Note that the fundamental group of M_n is the Artin group B_{n+1} . The action of \mathbb{C}^* on M_n is given by $(u_1, \dots, u_n) \mapsto (ku_1, \dots, k^n u_n)$ for $k \in \mathbb{C}^*$.

We regard $p_n(z)$ as a meromorphic function on the Riemann sphere $\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$. For $p_n(z) \in M_n$, we define the quadratic differential on \mathbb{P}^1 by $\phi(z) := p_n(z)^{N-2} dz^{\otimes 2}$ and denote by $Q(N, n)$ the space of such differentials.

To consider the period of $\sqrt{\phi} = p_n(z)^{\frac{N-2}{2}} dz$, we need to introduce homology groups $H_+(\phi)$ and $H_-(\phi)$ which provide appropriate cycles for the integration of $\sqrt{\phi}$. If N is even, the homology group is defined by the relative homology group $H_+(\phi) = H_1(\mathbb{C}, \text{Zero}(\phi); \mathbb{Z})$ where $\text{Zero}(\phi) \subset \mathbb{C}$ is the set of zeros of ϕ . If N is odd, the homology group is defined by $H_-(\phi) := H_1(S \setminus \pi^{-1}(\infty); \mathbb{Z})$ where $\pi: S \rightarrow \mathbb{P}^1$ is the double cover of the hyperelliptic curve $S = \{y^2 = p_n(z)\}$.

A linear map $Z_\phi: H_\pm(\phi) \rightarrow \mathbb{C}$, called the period of ϕ , is defined by

$$Z_\phi(\gamma) := \int_\gamma \sqrt{\phi}$$

for $\gamma \in H_\pm(\phi)$.

Let Γ be a free abelian group of rank n . An isomorphism of abelian groups $\theta: \Gamma \xrightarrow{\sim} H_{\pm}(\phi)$ is called a Γ -framing of $\phi \in Q(N, n)$. Denote by $Q(N, n)^{\Gamma}$ the set of framed differentials. For a framed differential $(\phi, \theta) \in Q(N, n)^{\Gamma}$, the composition of $\theta: \Gamma \rightarrow H_{\pm}(\phi)$ and $Z_{\phi}: H_{\pm}(\phi) \rightarrow \mathbb{C}$ gives a linear map $Z_{\phi} \circ \theta: \Gamma \rightarrow \mathbb{C}$. We define the period map by

$$\mathcal{W}_N: Q(N, n)^{\Gamma} \rightarrow \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}), \quad (\phi, \theta) \mapsto Z_{\phi} \circ \theta.$$

Fix a point $* = (\phi_0, \theta_0) \in Q(N, n)^{\Gamma}$ and let $Q(N, n)_{*}^{\Gamma} \subset Q(N, n)^{\Gamma}$ be the connected component containing $*$. Then the universal cover of $Q(N, n)_{*}^{\Gamma}$ is isomorphic to the universal cover \widetilde{M}_n of M_n . Thus the period map \mathcal{W}_N is extended on \widetilde{M}_n :

$$\mathcal{W}_N: \widetilde{M}_n \longrightarrow \text{Hom}(\Gamma, \mathbb{C}).$$

Next we consider the category \mathcal{D}_n^N (Section 4) and stability conditions on it (Section 5). For a quiver Q , we can define a dg algebra $\Gamma_N Q$, called the Ginzburg CY_N dg algebra [Gin]. Let $\mathcal{D}_{\text{fd}}(\Gamma_N Q)$ be the derived category of dg modules over $\Gamma_N Q$ with finite total dimension. It was proved by Keller [Kel11] that the category $\mathcal{D}_{\text{fd}}(\Gamma_N Q)$ is a CY_N triangulated category. Let $Q = \overrightarrow{A}_n$ be the A_n -quiver and set $\mathcal{D}_n^N := \mathcal{D}_{\text{fd}}(\Gamma_N \overrightarrow{A}_n)$. An object $S \in \mathcal{D}_n^N$ is called N -spherical if

$$\text{Hom}_{\mathcal{D}_n^N}(S, S[i]) = \begin{cases} k & \text{if } i = 0, N \\ 0 & \text{otherwise.} \end{cases}$$

There is a full subcategory \mathcal{H}_0 , called the standard heart of \mathcal{D}_n^N , and there are N -spherical objects $S_1, \dots, S_n \in \mathcal{H}_0$ such that \mathcal{H}_0 is generated by these N -spherical objects.

From such objects, Seidel and Thomas defined autoequivalences $\Phi_{S_1}, \dots, \Phi_{S_n} \in \text{Aut}(\mathcal{D}_n^N)$, called spherical twists [ST01]. They also proved that the subgroup $\text{Br}(\mathcal{D}_n^N) \subset \text{Aut}(\mathcal{D}_n^N)$ generated by $\Phi_{S_1}, \dots, \Phi_{S_n}$ is isomorphic to the braid group B_{n+1} .

A stability condition $\sigma = (Z, \mathcal{H})$ on a triangulated category \mathcal{D} consists of a heart $\mathcal{H} \subset \mathcal{D}$ of a bounded t-structure and a linear map $Z: K(\mathcal{H}) \rightarrow \mathbb{C}$ called the central charge, which satisfy some axioms. In [Bri07], Bridgeland showed that the set of stability conditions $\text{Stab}(\mathcal{D})$ has the structure of a complex manifold, and the projection map

$$\mathcal{Z}: \text{Stab}(\mathcal{D}) \longrightarrow \text{Hom}_{\mathbb{Z}}(K(\mathcal{D}), \mathbb{C}), \quad (Z, \mathcal{H}) \mapsto Z$$

is a local isomorphism. There are two group actions on $\text{Stab}(\mathcal{D})$; one is the action of \mathbb{C} and the other is the action of $\text{Aut}(\mathcal{D})$.

Consider the category \mathcal{D}_n^N . There is a distinguished connected component $\text{Stab}^{\circ}(\mathcal{D}_n^N) \subset \text{Stab}(\mathcal{D}_n^N)$ which contains stability conditions with the standard heart \mathcal{H}_0 . The action of the group $\text{Br}(\mathcal{D}_n^N)$ preserves the connected component $\text{Stab}^{\circ}(\mathcal{D}_n^N)$. Since $\text{Br}(\mathcal{D}_n^N) \cong B_{n+1}$, the space $\text{Stab}^{\circ}(\mathcal{D}_n^N)$ has the action of the braid group B_{n+1} .

Our main result is the following.

Theorem 1.1 (Theorem 7.12). *Assume that $N \geq 2$. There is a B_{n+1} -equivariant and \mathbb{C} -equivariant isomorphism of complex manifolds \widetilde{K}_N such that the diagram*

$$\begin{array}{ccc} \widetilde{M}_n & \xrightarrow{\widetilde{K}_N} & \text{Stab}^\circ(\mathcal{D}_n^N) \\ & \searrow \mathcal{W}_N & \swarrow \mathcal{Z} \\ & \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}) & \end{array}$$

commutes.

When $N = 2$, the result was proved by Thomas [Tho06]. When $N = 3$ and $n = 2$, the result was proved by Sutherland [Sut]. The result for $N = 3$ and arbitrary n follows from the result of Bridgeland-Smith [BS15] with some modifications (see Example 12.1 in [BS15]). After submitting this paper on arXiv, I received a preliminary version of the paper [BQS] from Bridgeland, which also proves the result for the case $n = 2$ and $2 \leq N \leq \infty$ by using a different method. Their approach is based on [Sut], while my approach is based on [BS15].

The result is compatible with the observation in [Bri06] that if there is a Frobenius structure on the space of stability conditions on a CY_N category, then the central charge corresponds to the twisted period of the Frobenius structure with the twist parameter $\nu = (N - 2)/2$. The twisted periods are characterized as the horizontal sections of the second structure connection of the Frobenius structure with some parameter ν (see [Dub04, Section 3]). By K. Saito's theory of primitive forms [Sai83], there is a Frobenius structure on M_n and the map \mathcal{W}_N naturally arises as the period map associated to the primitive form. As in [Dub04, Section 5], the map \mathcal{W}_N coincides with the twisted period map with the parameter $\nu = (N - 2)/2$. Therefore, the result supports the Bridgeland's conjecture in [Bri06].

1.3. Basic ideas. Many arguments of this paper are parallel to those of [BS15] with appropriate generalizations. We replace CY_3 categories with CY_N categories, quivers with $(N - 2)$ -colored quivers, simple zeros with zeros of order $N - 2$ and triangulations with N -angulations.

However as in Section 6, there are some points where parallel arguments can not be applied. In the case of N -angulations, it is not known what objects correspond to quivers with potential and their mutation in the case of triangulations. Hence we do not have natural dg algebras associated to N -angulations of surfaces and the categorical equivalence of Keller-Yang [KY11] is not established yet.

Fortunately, King-Qiu's result [KQ] on the hearts of \mathcal{D}_n^N together with the geometric realization of the $(N - 2)$ -cluster category of type A_n by Baur-Marsh [BM08] allows us to apply the framework of [BS15] to our cases.

The key idea of constructing stability conditions on \mathcal{D}_n^N from quadratic differentials is the fact that saddle-free quadratic differentials determine N -angulations of surfaces as follows. The differential ϕ determines a foliation on \mathbb{P}^1 by

$$\text{Im} \int p_n(z)^{\frac{N-2}{2}} dz = \text{constant}.$$

A leaf which connects zeros of ϕ is called a saddle trajectory. Near a zero of order $N - 2$, the foliation behaves as in Figure 2. As a result, the foliation of a saddle-free differential ϕ determines the N -angulation Δ_ϕ of the $((N - 2)(n + 1) + 2)$ -gon (Lemma 6.11) similar to the WKB triangulations in [BS15, GMN]. This is the reason why

we consider zeros of order $N - 2$. Using the results of Baur-Marsh and King-Qiu, we have a natural heart $\mathcal{H}(\Delta_\phi) \subset \mathcal{D}_n^N$ with simple objects S_1, \dots, S_n (Corollary 6.4). Corresponding to these simple objects, there are cycles $\gamma_1, \dots, \gamma_n \in H_\pm(\phi)$, called standard saddle classes (Corollary 6.13). The central charge of S_i is defined by the period of the corresponding saddle class γ_i :

$$Z(S_i) := Z_\phi(\gamma_i) = \int_{\gamma_i} p_n(z)^{\frac{N-2}{2}} dz.$$

Thus, we can construct the stability condition $(Z, \mathcal{H}(\Delta_\phi))$ on \mathcal{D}_n^N (Lemma 7.4).

1.4. Homotopy type problem. Ishii-Ueda-Uehara [IUU10] proved that the space $\text{Stab}(\mathcal{D}_n^2)$ is connected, so $\text{Stab}^\circ(\mathcal{D}_n^2) = \text{Stab}(\mathcal{D}_n^2)$. Together with the result of Thomas in [Tho06], we have $\text{Stab}(\mathcal{D}_n^2) \cong \widetilde{M}_n$. In particular, $\text{Stab}(\mathcal{D}_n^2)$ is contractible. For $N \geq 3$, Qiu proved that $\text{Stab}^\circ(\mathcal{D}_n^N)$ is simply connected (Corollary 5.5 in [Qiu15]). Our Theorem 1.1 implies the following stronger result (Conjecture 5.8 in [Qiu15]).

Corollary 1.2 (Corollary 7.13). *The distinguished connected component $\text{Stab}^\circ(\mathcal{D}_n^N)$ is contractible.*

The remaining problem is the connectedness of the space $\text{Stab}(\mathcal{D}_n^N)$. As in the case $N = 2$, we naively expect the following.

Conjecture 1.3. *The space $\text{Stab}(\mathcal{D}_n^N)$ is connected.*

Acknowledgements. First of all, I am deeply grateful to my advisor Akishi Kato for many valuable discussions, comments, correction of errors and constant encouragement. I would like to thank Kyoji Saito and Atsushi Takahashi for valuable comments on primitive forms. I also would like to thank Fumihiko Sanda for helpful discussions about Fukaya categories.

This work is supported by World Premier International Research Center Initiative (WPI initiative), MEXT, Japan.

This paper is dedicated to the memory of Kentaro Nagao, who explained many basics of this research area and gave me helpful comments at the starting point of this work.

Notations. We put $d_n^N := (N - 2)(n + 1) + 2$. We work over an algebraically closed field k . We use homological gradings $d: M_i \rightarrow M_{i-1}$ for dg algebras and dg modules. Assume that all triangulated categories in this paper are k -linear and their Grothendieck groups are free of finite rank ($K(\mathcal{D}) \cong \mathbb{Z}^n$ for some n).

2. QUADRATIC DIFFERENTIALS ASSOCIATED TO POLYNOMIALS

In the following, we assume that $N \geq 3$ and $n \geq 1$.

2.1. Braid groups and their representations. In this section, we introduce the braid groups and their representations on some lattices associated to A_n -quivers.

Definition 2.1. The Artin braid group B_{n+1} is the group generated by generators $\sigma_1, \sigma_2, \dots, \sigma_n$ with the following relations:

$$\begin{aligned} \sigma_i \sigma_j \sigma_i &= \sigma_j \sigma_i \sigma_j & \text{if } |i - j| = 1 \\ \sigma_i \sigma_j &= \sigma_i \sigma_j & \text{if } |i - j| \geq 2. \end{aligned}$$

It is known that the element

$$C := (\sigma_1 \sigma_2 \cdots \sigma_n)^{n+1}$$

generates the center of B_{n+1} .

We consider two representations ρ_+ and ρ_- of B_{n+1} on some lattices associated to the A_n -quivers. Such representations are equivalent to the reduced Burau representations with the parameters ± 1 (see [KT08, Section 3]).

An A_n -quiver \overrightarrow{A}_n is a quiver obtained by giving orientations on edges of the Dynkin diagram of type A_n . We set vertices of \overrightarrow{A}_n by $\{1, \dots, n\}$ and assume that vertices i and $i+1$ are connected by some arrow. Let q_{ij} be the number of arrows from i to j .

Let L be a free abelian group of rank n with generators $\alpha_1, \dots, \alpha_n$:

$$L := \bigoplus_{i=1}^n \mathbb{Z}\alpha_i.$$

We define a symmetric bilinear form $\langle \cdot, \cdot \rangle_+$ and a skew-symmetric bilinear form $\langle \cdot, \cdot \rangle_-$ on L by

$$\begin{aligned} \langle \alpha_i, \alpha_j \rangle_+ &:= \delta_{ij} + \delta_{ji} - (q_{ij} + q_{ji}) \\ \langle \alpha_i, \alpha_j \rangle_- &:= \delta_{ij} - \delta_{ji} - (q_{ij} - q_{ji}) \end{aligned}$$

where δ_{ij} is the Kronecker delta.

Reflections (which depend on the sign \pm) $r_1^\pm, \dots, r_n^\pm: L \rightarrow L$ via the bilinear forms $\langle \cdot, \cdot \rangle_\pm$ is defined by

$$r_i^\pm(\alpha_j) := \alpha_j - \langle \alpha_i, \alpha_j \rangle_\pm \alpha_i.$$

Let $W_\pm := \langle r_1^\pm, \dots, r_n^\pm \rangle$ be the groups generated by these reflections.

Two representations $\rho_\pm: B_{n+1} \rightarrow \mathrm{GL}(L, \mathbb{Z})$ are defined by

$$\rho_\pm(\sigma_i) := r_i^\pm.$$

We put the kernel of ρ_\pm by P_\pm . Then we have a short exact sequence

$$1 \longrightarrow P_\pm \longrightarrow B_{n+1} \longrightarrow W_\pm \longrightarrow 1.$$

Note that the group W_+ is a symmetric group of degree $n+1$ and the group P_+ is a pure braid group.

2.2. Spaces of polynomials. Let $p_n(z)$ be a polynomial of the following form

$$p_n(z) = z^{n+1} + u_1 z^{n-1} + u_2 z^{n-2} + \cdots + u_n, \quad u_1, \dots, u_n \in \mathbb{C}.$$

We denote by $\Delta(p_n)$ the discriminant of p_n . (If $p_n(z) = \prod_{i=1}^{n+1} (z - a_i)$, then $\Delta(p_n) = \prod_{i < j} (a_i - a_j)^2$.)

Denote by M_n the space of such polynomials with simple zeros:

$$M_n := \{p_n(z) \mid \Delta(p_n) \neq 0\}.$$

Define the free \mathbb{C}^* -action on M_n by

$$(k \cdot p_n)(z) := z^{n+1} + k u_1 z^{n-1} + k^2 u_2 z^{n-2} + \cdots + k^n u_n \quad (k \in \mathbb{C}^*).$$

Note that for the above action, we have the equation

$$(k \cdot p_n)(kz) = k^{n+1} p_n(z).$$

Consider the description of M_n as the configuration space of \mathbb{C} . Let $C_{n+1}(\mathbb{C})$ be the configuration space of $(n+1)$ distinct points in \mathbb{C} and $C_{n+1}^0(\mathbb{C}) \subset C_{n+1}(\mathbb{C})$ be the subspace consists of configurations with the center of mass zero:

$$C_{n+1}(\mathbb{C}) := \{(a_1, \dots, a_{n+1}) \in \mathbb{C}^{n+1} \mid a_i \neq a_j \text{ for } i \neq j\} / S_n$$

$$C_{n+1}^0(\mathbb{C}) := \{(a_1, \dots, a_{n+1}) \in \mathbb{C}^{n+1} \mid \sum_{i=1}^{n+1} a_i = 0, a_i \neq a_j \text{ for } i \neq j\} / S_n.$$

A complex number $c \in \mathbb{C}$ acts on $C_{n+1}(\mathbb{C})$ as a parallel transportation $(a_1, \dots, a_{n+1}) \mapsto (a_1 + c, \dots, a_{n+1} + c)$. Hence we have $C_{n+1}^0(\mathbb{C}) = C_{n+1}(\mathbb{C}) / \mathbb{C}$.

As a result, we have an isomorphism $C_{n+1}^0(\mathbb{C}) \xrightarrow{\sim} M_n$ given by

$$(a_1, \dots, a_{n+1}) \mapsto \prod_{i=1}^{n+1} (z - a_i).$$

Next we consider the description of a braid group B_{n+1} as the fundamental group of M_n . Take a configuration $\{a_1, \dots, a_{n+1}\} \in C_{n+1}(\mathbb{C})$. We fix some order (a_1, \dots, a_{n+1}) for these $n+1$ points. Consider the collection of non-crossing paths $\gamma_1, \dots, \gamma_n$ in \mathbb{C} such that γ_i connects a_i and a_{i+1} . We call it the A_n -chain of paths. Let τ_i be a closed path in $C_{n+1}(\mathbb{C})$ which corresponds to the half-twist along γ_i in the clockwise direction (see Figure 1), and consider its image in M_n via the projection $C_{n+1}(\mathbb{C}) \rightarrow M_n$ (we also write it by τ_i).




FIGURE 1. Half-twists along the A_n -chain.

It is well-known that there is an isomorphism of groups

$$\pi_1(M_n) \xrightarrow{\sim} B_{n+1}, \quad [\tau_i] \mapsto \sigma_i.$$

For more details, we refer to [KT08, Section 1.6].

2.3. Quadratic differentials associated to polynomials. Let \mathbb{P}^1 be the Riemann sphere, and we fix the coordinate $z \in \mathbb{C} \cup \{\infty\} \cong \mathbb{P}^1$. For a polynomial $p_n \in M_n$, define the meromorphic quadratic differential ϕ on \mathbb{P}^1 by

$$\phi(z) := p_n(z)^{N-2} dz^{\otimes 2}.$$

ϕ has zeros of order $(N-2)$ at distinct $n+1$ points on $\mathbb{C} \subset \mathbb{P}^1$, and a unique pole of order $(d_n^N + 2)$ at $\infty \in \mathbb{P}^1$.

We denote by $Q(N, n)$ the space of such quadratic differentials associated to $p_n \in M_n$:

$$Q(N, n) := \{p_n(z)^{N-2} dz^{\otimes 2} \mid p_n \in M_n\}.$$

Independent of N , the space $Q(N, n)$ is isomorphic to M_n as a complex manifold, but we give this space various structures which depend on N .

We introduce the action of \mathbb{C}^* on $Q(N, n)$. For $\phi(z) = p_n(z)^{N-2} dz^{\otimes 2}$, the action of \mathbb{C}^* is defined by

$$(k \cdot \phi)(z) := \{(k \cdot p_n)(z)\}^{N-2} dz^{\otimes 2} \quad (k \in \mathbb{C}^*)$$

where $k \cdot p_n$ is the action defined in Section 2.2. By using the equation $(k \cdot p_n)(kz) = k^{n+1} p_n(z)$, we have the following formula.

Lemma 2.2. *For $\phi \in Q(N, n)$, the equation*

$$(k \cdot \phi)(kz) = k^{d_n^N} \phi(z) \quad (k \in \mathbb{C}^*)$$

holds.

For $\phi \in Q(N, n)$, denote by $\text{Zero}(\phi) \subset \mathbb{P}^1$ the set of zeros of ϕ and by $\text{Crit}(\phi) \subset \mathbb{P}^1$ the set of critical points of ϕ . Note that $\text{Crit}(\phi) = \text{Zero}(\phi) \cup \{\infty\}$.

The action $z \mapsto kz$ of $k \in \mathbb{C}^*$ on \mathbb{P}^1 maps the critical points of ϕ to those of $k \cdot \phi$:

$$k(\text{Crit}(\phi)) = \text{Crit}(k \cdot \phi).$$

2.4. Homology groups and periods.

Definition 2.3. Depending on the parity of N , we introduce homology groups $H_+(\phi)$ and $H_-(\phi)$ in the following.

- If N is even, $H_+(\phi)$ is defined by the relative homology group

$$H_+(\phi) := H_1(\mathbb{C}, \text{Zero}(\phi); \mathbb{Z}).$$

- If N is odd, $H_-(\phi)$ is given as follows. Let S be a hyperelliptic curve defined by the equation $y^2 = p_n(z)$ and $\pi: S \rightarrow \mathbb{P}^1$ be an associated branched covering map of degree 2. Then the homology group $H_-(\phi)$ is defined by

$$H_-(\phi) := H_1(S \setminus \pi^{-1}(\infty); \mathbb{Z}).$$

Note that if $N = 3$, the homology group $H_-(\phi)$ corresponds to the hat-homology group in [BS15] which is written as $\hat{H}(\phi)$.

Lemma 2.4. *Both $H_+(\phi)$ and $H_-(\phi)$ are free abelian groups of rank n .*

Proof. If N is even, $|\text{Zero}(\phi)| = n + 1$ implies $\text{rank } H_+(\phi) = n$.

If N is odd, there are two cases, n is even or odd. If n is even, the genus of the hyperelliptic curve S is $n/2$ and the fiber $\pi^{-1}(\infty)$ consists of one point. Hence we have $\text{rank } H_-(\phi) = 2(n/2) = n$. When n is odd, the genus of S is $(n-1)/2$ and the fiber $\pi^{-1}(\infty)$ consists of two points. Hence we have $\text{rank } H_-(\phi) = 2\{(n-1)/2\} + 1 = n$. \square

Let $\psi := \sqrt{\phi} = p_n(z)^{\frac{N-2}{2}} dz$ be the square root of ϕ . (If N is odd, ψ is determined up to the sign.) If N is even, then ψ is a holomorphic 1-form on $\mathbb{P}^1 \setminus \{\infty\}$, and if N is odd, then ψ is a holomorphic 1-form on $S \setminus \pi^{-1}(\infty)$.

Therefore, for any cycle $\alpha \in H_{\pm}(\phi)$ the integration of 1-form ψ via the cycle α

$$Z_{\phi}(\alpha) := \int_{\alpha} \psi \in \mathbb{C}$$

is well-defined.

Thus we have a group homomorphism $Z_{\phi}: H_{\pm}(\phi) \rightarrow \mathbb{C}$, called the period of ϕ .

2.5. Framings. Following [BS15, Section 2.6], we introduce framings on $Q(N, n)$. Let Γ be a free abelian group of rank n . A Γ -framing of $\phi \in Q(N, n)$ is an isomorphism of abelian groups

$$\theta: \Gamma \xrightarrow{\sim} H_{\pm}(\phi).$$

Denote by $Q(N, n)^{\Gamma}$ the space of pairs (ϕ, θ) consisting of a quadratic differential $\phi \in Q(N, n)$ and a Γ -framing θ . The projection

$$Q(N, n)^{\Gamma} \longrightarrow Q(N, n), \quad (\phi, \theta) \mapsto \phi$$

defines a principal $GL(n, \mathbb{Z})$ -bundle on $Q(N, n)$.

For $(\phi, \theta) \in Q(N, n)^{\Gamma}$, the composition of a framing θ and a period Z_{ϕ} gives a group homomorphism $Z_{\phi} \circ \theta: \Gamma \rightarrow \mathbb{C}$. Thus, we have the map

$$\mathcal{W}_N: Q(N, n)^{\Gamma} \longrightarrow \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$$

defined by $\mathcal{W}_N((\phi, \theta)) := Z_{\phi} \circ \theta \in \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$. We call \mathcal{W}_N a period map.

Proposition 2.5. *The period map*

$$\mathcal{W}_N: Q(N, n)^{\Gamma} \longrightarrow \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$$

is a local isomorphism of complex manifolds.

Proof. Theorem 4.12 in [BS15] can be easily extended to our cases. In our cases, the result also follows from K. Saito's theory on primitive forms [Sai83, Section 5] since \mathcal{W}_N is interpreted as the period map associated to the primitive form of the A_n -singularity. \square

2.6. Intersection forms. In this section, we introduce intersection forms on $H_{\pm}(\phi)$.

First we consider the case N is even. The exact sequence of relative homology groups gives the isomorphism of homology groups

$$H_+(\phi) = H_1(\mathbb{C}, \text{Zero}(\phi); \mathbb{Z}) \cong \tilde{H}_0(\text{Zero}(\phi); \mathbb{Z})$$

where $\tilde{H}_0(\text{Zero}(\phi); \mathbb{Z})$ is the reduced homology group of $H_0(\text{Zero}(\phi); \mathbb{Z})$, and this isomorphism is induced by the boundary map. There is a trivial intersection form on $H_0(\text{Zero}(\phi); \mathbb{Z})$, and it induces the intersection form on $\tilde{H}_0(\text{Zero}(\phi); \mathbb{Z})$. Through the above isomorphism of homology groups, we have the intersection form

$$I_+: H_+(\phi) \times H_+(\phi) \longrightarrow \mathbb{Z}.$$

Note that I_+ is symmetric and non-degenerate.

Next consider the case N is odd. Let $\pi: S \rightarrow \mathbb{P}^1$ be the hyperelliptic curve introduced in Section 2.4, and $\iota: S \setminus \pi^{-1}(\infty) \rightarrow S$ be the inclusion. This inclusion induces the surjective group homomorphism

$$\iota_*: H_-(\phi) = H_1(S \setminus \pi^{-1}(\infty); \mathbb{Z}) \longrightarrow H_1(S; \mathbb{Z}).$$

Since there is a skew-symmetric non-degenerate intersection form on $H_1(S; \mathbb{Z})$, through the inclusion ι_* , we have the intersection form

$$I_-: H_-(\phi) \times H_-(\phi) \longrightarrow \mathbb{Z}.$$

The map ι_* is an isomorphism if n is even and has the kernel of rank one if n is odd. Therefore, I_- is non-degenerate if n is even or has the kernel of rank one if n is odd.

For $\phi \in Q(N, n)$, fix the order of $\text{Zero}(\phi)$ and consider the A_n -chain of path $\gamma_1, \dots, \gamma_n$. If N is even, the path γ_i determines the homology class $\hat{\gamma}_i \in H_+(\phi)$ since

γ_i connects zeros of ϕ . If N is odd, since the lift $\pi^{-1}(\gamma_i)$ on the hyperelliptic curve S is a closed path in $S \setminus \pi^{-1}(\infty)$, we have the homology class $\hat{\gamma}_i := [\pi^{-1}(\gamma_i)] \in H_-(\phi)$.

The pair of the homology group $H_{\pm}(\phi)$ and the intersection form I_{\pm} gives a geometric realization of the lattice $(L, \langle, \rangle_{\pm})$ which is introduced in Section 2.1.

Lemma 2.6. *There is an isomorphism $H_{\pm}(\phi) \cong L$ given by $\hat{\gamma}_i \mapsto \alpha_i$. Further, by choosing suitable orientations of $\hat{\gamma}_1, \dots, \hat{\gamma}_n$, this isomorphism takes the intersection form I_{\pm} to the bilinear form \langle, \rangle_{\pm} .*

Proof. The result follows from the direct calculation. \square

2.7. Local system. For each $\phi \in Q(N, n)$, by corresponding to the homology group $H_{\pm}(\phi)$, we have a local system

$$\bigcup_{\phi \in Q(N, n)} H_{\pm}(\phi) \longrightarrow Q(N, n).$$

For a continuous path $c: [0, 1] \rightarrow Q(N, n)$ with $\phi_0 = c(0)$, $\phi_1 = c(1)$, a parallel transportation along c gives the identification

$$\text{GM}_c: H_{\pm}(\phi_0) \xrightarrow{\sim} H_{\pm}(\phi_1),$$

and this identification determines a flat connection on the local system. We call it the Gauss-Manin connection.

In the following, we fix a point $(\phi_0, \theta_0) \in Q(N, n)^{\Gamma}$ and denote by $Q(N, n)^{\Gamma}_*$ the connected component containing (ϕ_0, θ_0) . Any point $(\phi, \theta) \in Q(N, n)^{\Gamma}$ is contained in $Q(N, n)^{\Gamma}_*$ if and only if there is a continuous path $c: [0, 1] \rightarrow Q(N, n)$ connecting ϕ_0 and ϕ such that the diagram

$$\begin{array}{ccc} & \Gamma & \\ \theta_0 \swarrow & & \searrow \theta \\ H_{\pm}(\phi_0) & \xrightarrow{\text{GM}_c} & H_{\pm}(\phi) \end{array}$$

commutes.

Note that different connected components of $Q(N, n)^{\Gamma}$ are all isomorphic since these are related by the action of $\text{GL}(n, \mathbb{Z})$.

We introduce the \mathbb{C} -action on $Q(N, n)^{\Gamma}_*$, which is compatible with the \mathbb{C} -action on the space of stability conditions given in Section 5.3.

Definition 2.7. For a framed differential $(\phi, \theta) \in Q(N, n)^{\Gamma}_*$, the new framed differential $(\phi', \theta') := t \cdot (\phi, \theta)$ for $t \in \mathbb{C}$ is defined by

$$\phi' := e^{-(2\pi i/d_n^N)t} \cdot \phi, \quad \theta' := \text{GM}_c \circ \theta$$

where GM_c is the Gauss-Manin connection along the path $c(s) := e^{-(2\pi i/d_n^N)ts} \cdot \phi$ for $s \in [0, 1]$ which connects ϕ and ϕ' .

2.8. Monodromy representations. By using the intersection form I_{\pm} on $H_{\pm}(\phi)$ defined in Section 2.6, we can compute how cycles change by the Gauss-Manin connection along a closed path in $Q(N, n)$. The result is known as the Picard-Lefschetz formula.

Let $\phi_0 = \left[\prod_{i=1}^{n+1} (z - a_i) \right]^{N-2} dz^{\otimes 2}$ be the fixed differential in the previous section. Take the A_n -chain of paths $\gamma_1, \dots, \gamma_n$ for ordered zeros $(a_1, a_2, \dots, a_{n+1})$ as in Section 2.2. Under the identification $M_n \cong Q(N, n)$, we consider the half-twists

τ_1, \dots, τ_n as in Section 2.2 and take the orientations of $\hat{\gamma}_1, \dots, \hat{\gamma}_n \in H_{\pm}(\phi_0)$ as in Lemma 2.6.

Proposition 2.8. *The monodromy representation of $[\tau_i] \in \pi_1(Q(N, n), \phi_0)$ on $H_{\pm}(\phi_0)$ is given by*

$$\text{GM}_{\tau_i}(\beta) = \beta - I_{\pm}(\hat{\gamma}_i, \beta) \hat{\gamma}_i \quad (\beta \in H_{\pm}(\phi_0)).$$

Lemma 2.6 and Proposition 2.8 imply that the monodromy representation of the braid group $\pi_1(M_n, *) \cong B_{n+1}$ on $H_{\pm}(\phi_0)$ gives a geometric realization of two representations ρ_+ and ρ_- introduced in Section 2.1. Further recall from Section 2.1 that the subgroup $P_{\pm} \subset B_{n+1}$ is the kernel of ρ_{\pm} and the group W_{\pm} is the image of ρ_{\pm} on $L \cong H_{\pm}(\phi_0)$.

Let \widetilde{M}_n be a universal cover of M_n . Note that B_{n+1} acts on \widetilde{M}_n as the group of deck transformations.

Corollary 2.9. *The connected component $Q(N, n)_*^{\Gamma}$ is isomorphic to the space \widetilde{M}_n/P_{\pm} and the projection map $Q(N, n)_*^{\Gamma} \rightarrow Q(N, n)$ gives a principal W_{\pm} -bundle.*

Proof. It follows from the above remarks. \square

3. TRAJECTORIES ON \mathbb{P}^1

In Section 3, in order to fix notations, we collect basic facts and definitions for trajectories on \mathbb{P}^1 from [BS15, Section 3]. Since our Riemann surface is only \mathbb{P}^1 , many results are simplified.

3.1. Foliations. Let $\phi \in Q(N, n)$. For any neighborhood of a point in $\mathbb{P}^1 \setminus \text{Crit}(\phi)$, the differential $\phi(z) = \varphi(z)dz^{\otimes 2}$ defines the distinguished local coordinate w by

$$w := \int \sqrt{\varphi(z)} dz,$$

up to the transformation $w \mapsto \pm w + c$ ($c \in \mathbb{C}$). The coordinate w also characterized by

$$\phi(w) = dw \otimes dw.$$

The coordinate w determines a foliation of the phase θ on $\mathbb{P}^1 \setminus \text{Crit}(\phi)$ by

$$\text{Im}(w/e^{i\pi\theta}) = \text{constant}.$$

A straight arc $\gamma: I \rightarrow \mathbb{P}^1 \setminus \text{Crit}(\phi)$ of the phase θ is an integral curve along the foliation $\text{Im}(w/e^{i\pi\theta}) = \text{constant}$, defined on an open interval $I \subset \mathbb{R}$. Note that a maximal straight arc of the phase θ is a leaf of the foliation of the phase θ .

A foliation of the phase $\theta = 0$ is called a horizontal foliation, and a maximal straight arc of the phase $\theta = 0$ is called a trajectory.

Lemma 3.1. *Let $\phi \in Q(N, n)$. For $k = re^{i\pi\theta} \in \mathbb{C}^*$, set $\phi' := k \cdot \phi$. Then the distinguished coordinates w and w' which come from ϕ and ϕ' have the relation*

$$w' = k^{\frac{dN}{2}} w = r^{\frac{dN}{2}} e^{i\pi \frac{dN}{2} \theta} w.$$

In addition, the transformation $z \mapsto kz$ on \mathbb{P}^1 maps the horizontal straight arcs for ϕ to the straight arcs of the phase $(d_n^N \theta)/2$ for ϕ' .

Proof. Let $\gamma: I \rightarrow \mathbb{P}^1 \setminus \text{Crit}(\phi)$ be a horizontal straight arc of ϕ . Then the following computation gives the result:

$$\begin{aligned} \int^{\gamma(t)} \sqrt{\phi'(z')} dz' &= \int^{\gamma(t)} \sqrt{(k \cdot \phi)(z')} dz' \\ &= \int^{\gamma(t)} \sqrt{(k \cdot \phi)(kz)} d(kz) \quad (\text{put } z' = kz.) \\ &= k^{\frac{dN}{2}} \int^{\gamma(t)} \phi(z) dz \quad (\text{by Lemma 2.2}). \end{aligned}$$

□

A maximal straight arc γ of the phase θ defined on a finite open interval (a, b) can be extended to a continuous path $\gamma: [a, b] \rightarrow \mathbb{P}^1$ with $\gamma(a), \gamma(b) \in \text{Zero}(\phi)$ since \mathbb{P}^1 is compact. According to $\theta = 0$ or $\theta \neq 0$, we call γ a saddle trajectory or a saddle connection respectively.

3.2. Foliations near critical points. Here we see the local behavior of the horizontal foliation near a critical point. In details, we refer to [BS15, Section 3.3]. First we consider the behavior near a zero. Let $\phi \in Q(N, n)$ and $p \in \text{Zero}(\phi)$ be a zero of order $N-2$. Then there is a local coordinate t of a neighborhood $p \in U \subset \mathbb{P}^1$ ($t = 0$ corresponds to p) such that

$$\phi = c^2 t^{N-2} dt^{\otimes 2}, \quad c = \frac{1}{2} N.$$

Hence on $U \setminus \{p\}$, the distinguished local coordinate w takes the form $w = t^{N/2}$. Figure 2 illustrates the local behavior of the horizontal foliation in the case $N = 3, 4, 5$.




FIGURE 2. Trajectories near zeros of order 1, 2, 3.

Note that there are N horizontal straight arcs departing from p . This fact is the reason why the differential ϕ determines N -angulations.

Next we consider the behavior near a pole. Recall from Section 2.3 that ϕ has a unique pole of order $d_n^N + 2$ at $\infty \in \mathbb{P}^1$. Therefore it is sufficient to consider a pole of order > 2 . In this case, there is a neighborhood $\infty \in U \subset \mathbb{P}^1$ and a collection of d_n^N distinguished tangent directions v_i ($i = 1, \dots, d_n^N$) at ∞ , such that any trajectory entering U tends to ∞ and becomes asymptotic to one of the v_i . For more details, we refer to [BS15, Section 3.3].

3.3. Classification of trajectories. In this section, we summarize the global structure of the horizontal foliation on \mathbb{P}^1 following [BS15, Section 3.4], and [Str84, Section 9-11 and 15].

We first note that ϕ is holomorphic on the contractible domain $\mathbb{P}^1 \setminus \{\infty\}$. Then Theorem 14.2.2 in [Str84] implies that there are no closed trajectories and Theorem 15.2 in [Str84] implies that there are no recurrent trajectories. Hence in our setting, every trajectory of ϕ is classified to exactly one of the following classes.

- (1) A saddle trajectory approaches distinct zeros of ϕ at both ends.
- (2) A separating trajectory approaches a zero of ϕ at one end and a pole $\infty \in \mathbb{P}^1$ at the other end.
- (3) A generic trajectory approaches a pole $\infty \in \mathbb{P}^1$ at both ends.

As explained in the previous section, there are only finite horizontal straight arcs departing from zeros of ϕ . Therefore, the number of saddle trajectories and separating trajectories is finite. For a quadratic differential ϕ , denote by s_ϕ the number of saddle trajectories, and by t_ϕ the number of separating trajectories. Then the fact that there are N horizontal straight arcs departing from each zero of ϕ implies the equation

$$(3.1) \quad 2s_\phi + t_\phi = N(n + 1).$$

By removing these two type of trajectories and $\text{Crit}(\phi)$ from \mathbb{P}^1 , the remaining open surface splits into a disjoint union of connected components. In our setting, each connected component coincides with one of the following two type of surfaces.

- (1) A half-plane is the upper half-plane

$$\{w \in \mathbb{C} \mid \text{Im } w > 0\}$$

equipped with the differential $dw^{\otimes 2}$. It is swept out by generic trajectories with the both end points at $\infty \in \mathbb{P}^1$. The boundary $\{\text{Im } w = 0\}$ consists of saddle trajectories and two separating trajectories.

- (2) A horizontal strip is the strip domain

$$\{w \in \mathbb{C} \mid a < \text{Im } w < b\}$$

equipped with the differential $dw^{\otimes 2}$. It is also swept out by generic trajectories with the both end points at $\infty \in \mathbb{P}^1$. Each of the two boundaries $\{\text{Im } w = a\}$ and $\{\text{Im } w = b\}$ consists of saddle trajectories and two separating trajectories.

Note that every saddle trajectory or separating trajectory appears just two times as a boundary element of these open surfaces.




FIGURE 3. Examples of a half plane and a horizontal strip.

Let k be the number of half planes and l be the number of horizontal strips appearing in such a decomposition of \mathbb{P}^1 . Since every half plane has 2 separating trajectories on the boundary, and every horizontal strip has 4 separating trajectories on the two boundaries, we have the equation

$$(3.2) \quad 2k + 4l = 2t_\phi.$$

We also note that near a pole of order $m > 2$, just $m - 2$ half planes appear. Since ϕ has a unique pole of order $(d_n^N + 2)$ at $\infty \in \mathbb{P}^1$, we have

$$(3.3) \quad k = d_n^N = (N - 2)(n + 1) + 2.$$

Lemma 3.2. *For $\phi \in Q(N, n)$, the number of saddle trajectories s_ϕ and the number of horizontal strips l satisfy the equation*

$$s_\phi + l = n.$$

Proof. This immediately follows from equations (3.1), (3.2) and (3.3). \square

Definition 3.3. A differential $\phi \in Q(N, n)$ is called saddle-free if ϕ has no saddle trajectories (i.e. $s_\phi = 0$).

For a saddle-free differential $\phi \in Q(N, n)$, Lemma 3.2 implies that exactly n horizontal strips appear in the decomposition of \mathbb{P}^1 by ϕ . In this situation, any boundary component of a half plane or a horizontal strip consists of two separating trajectories.

3.4. Hat-homology classes and standard saddle connections. In this section, we collect some constructions of homology classes from [BS15, Section 3.2 and 3.4].

Take $\phi \in Q(N, n)$. Recall from Section 2.4 that the square root of ϕ defines a holomorphic 1-form $\psi = \sqrt{\phi}$ on $\mathbb{P}^1 \setminus \{\infty\}$ or $S \setminus \pi^{-1}(\infty)$ depending on the parity of N . We set

$$\hat{S}^\circ := \begin{cases} \mathbb{P}^1 \setminus \{\infty\} & \text{if } N \text{ is even} \\ S \setminus \pi^{-1}(\infty) & \text{if } N \text{ is odd.} \end{cases}$$

We first note that the 1-form ψ defines the distinguished local coordinate \hat{w} on $\hat{S}^\circ \setminus \text{Zero}(\psi)$ by $\psi = d\hat{w}$, up to the transformation $\hat{w} \mapsto \hat{w} + c$ ($c \in \mathbb{C}$). As a result, ψ determines the horizontal foliation on $\hat{S}^\circ \setminus \text{Zero}(\psi)$ by $\text{Im } \hat{w} = \text{constant}$. This foliation has the canonical orientation to make the tangent vectors lie in $\mathbb{R}_{>0}$. We call it the oriented horizontal foliation of ψ .

For a saddle trajectory $\gamma: [a, b] \rightarrow \mathbb{P}^1$, we can define a homology class $\hat{\gamma} \in H_\pm(\phi)$, called the hat-homology class of γ by the following.

If N is even, we can orient γ to be compatible with the oriented horizontal foliation of ψ . Since $\gamma(a), \gamma(b) \in \text{Zero}(\phi)$, the oriented curve γ defines a homology class

$$\hat{\gamma} := [\gamma] \in H_+(\phi) = H_1(\mathbb{C}, \text{Zero}(\phi); \mathbb{Z}).$$

. By definition, it satisfies $Z_\phi(\hat{\gamma}) \in \mathbb{R}_{>0}$.

If N is odd, since zeros of ϕ are branched points of the double cover $\pi: S \rightarrow \mathbb{P}^1$, the inverse image $\pi^{-1}(\gamma)$ becomes a closed curve in \hat{S}° . We can orient $\pi^{-1}(\gamma)$ to be compatible with the oriented horizontal orientation of ψ . Then the oriented closed curve $\pi^{-1}(\gamma)$ defines a homology class

$$\hat{\gamma} := [\pi^{-1}(\gamma)] \in H_-(\phi) = H_1(S \setminus \pi^{-1}(\infty); \mathbb{Z}).$$

In the case that N is even, it satisfies $Z_\phi(\hat{\gamma}) \in \mathbb{R}_{>0}$.

A similar construction works for a saddle connection $\gamma: [a, b] \rightarrow \mathbb{P}^1$ with the phase $\theta \in (0, \pi)$. In this case, the orientation is determined to satisfy that $Z_\phi(\hat{\gamma}) \in \{re^{i\pi\theta} \mid r > 0\} \subset \mathbb{H}$.

Let $h = \{a < \text{Im } w < b\}$ be a horizontal strip appearing in the domain decomposition of \mathbb{P}^1 by a saddle-free differential ϕ . Then each of two boundary components $\{\text{Im } w = a\}$ and $\{\text{Im } w = b\}$ contains exactly one zero of ϕ and consists of two separating trajectories intersecting at the zero. Hence, there is a unique saddle connection l_h with the phase $\theta \in (0, \pi)$ which connects two zeros appearing in two boundary components. We denote by $\gamma_h := \hat{l}_h \in H_\pm(\phi)$ the hat-homology class associated to the saddle connection l_h .




FIGURE 4. Standard saddle connection.

Since ϕ is saddle-free, Lemma 3.2 implies that there are precisely n horizontal strips in the decomposition of \mathbb{P}^1 by ϕ . Thus, we obtain n saddle connections l_{h_1}, \dots, l_{h_n} and n hat-homology classes $\gamma_{h_1}, \dots, \gamma_{h_n}$. l_{h_i} is called a standard saddle connection, and γ_{h_i} is called a standard saddle class.

Lemma 3.4 ([BS15], Lemma 3.2). *Standard saddle classes $\gamma_{h_1}, \dots, \gamma_{h_n}$ form a basis of $H_\pm(\phi)$.*

Proof. The argument of Lemma 3.2 in [BS15] for $N = 3$ also works for $N \geq 3$. \square

4. CY_N CATEGORIES ASSOCIATED TO ACYCLIC QUIVERS

For objects E, F in a triangulated category \mathcal{D} , we set

$$\text{Hom}^i(E, F) := \text{Hom}_{\mathcal{D}}(E, F[i])$$

$$\text{hom}^i(E, F) := \dim_k \text{Hom}_{\mathcal{D}}(E, F[i])$$

where $[i]$ is the i -th shift functor.

4.1. Ginzburg dg algebras. Let $Q = (Q_0, Q_1)$ be a finite acyclic quiver with vertices $Q_0 = \{1, \dots, n\}$ and arrows Q_1 . The Ginzburg dg algebra [Gin] $\Gamma_N Q := (k\overline{Q}, d)$ is defined as follows. Define a graded quiver \overline{Q} with vertices $\overline{Q}_0 = \{1, \dots, n\}$ and following arrows:

- an original arrow $a: i \rightarrow j \in Q_1$ (degree 0);
- an opposite arrow $a^*: j \rightarrow i$ for the original arrow $a: i \rightarrow j \in Q_1$ (degree $N - 2$);
- a loop t_i for each vertex $i \in Q_0$ (degree $N - 1$).

Let $k\overline{Q}$ be a graded path algebra of \overline{Q} , and define a differential $d: k\overline{Q} \rightarrow k\overline{Q}$ of degree -1 by

- $da = da^* = 0$ for $a \in Q_1$;
- $dt_i = e_i \left(\sum_{a \in Q_1} (aa^* - a^*a) \right) e_i$;

where e_i is the idempotent at $i \in Q_0$. Thus we have the dg algebra $\Gamma_N Q = (k\overline{Q}, d)$. Note that the 0-th homology is given by $\mathbf{H}_0(\Gamma_N Q) \cong kQ$.

Let $\mathcal{D}(\Gamma_N Q)$ be the derived category of right dg-modules over $\Gamma_N Q$, and \mathcal{D}_Q^N be the full subcategory of $\mathcal{D}(\Gamma_N Q)$ consists of dg modules M whose homology is of finite total dimension:

$$\mathcal{D}_Q^N := \{ M \in \mathcal{D}(\Gamma_N Q) \mid \sum_{i \in \mathbb{Z}} \dim_k \mathbf{H}_i(M) < \infty \}.$$

A triangulated category \mathcal{D} is called N -Calabi-Yau (CY_N) if for any objects $E, F \in \mathcal{D}$, there is a natural isomorphism of k -vector spaces

$$\nu: \text{Hom}(E, F) \xrightarrow{\sim} \text{Hom}^N(F, E)^*.$$

(For a vector space V , we denote by V^* the dual vector space.)

Theorem 4.1 ([Kel11], Theorem 6.3). *The category \mathcal{D}_Q^N is a CY_N triangulated category.*

Let $K(\mathcal{D}_Q^N)$ be the Grothendieck group of \mathcal{D}_Q^N . A bilinear form $\chi: K(\mathcal{D}_Q^N) \times K(\mathcal{D}_Q^N) \rightarrow \mathbb{Z}$, called the Euler form, is defined to be

$$\chi(E, F) := \sum_{i \in \mathbb{Z}} (-1)^i \text{hom}^i(E, F), \quad \text{for } E, F \in \mathcal{D}_Q^N.$$

The CY_N property of \mathcal{D}_Q^N implies that χ is symmetric if N is even, and skew-symmetric if N is odd.

For a quiver \overrightarrow{A}_n , we set $\mathcal{D}_n^N := \mathcal{D}_{\overrightarrow{A}_n}^N$.

4.2. Hearts of bounded t-structures.

Definition 4.2 ([BBD82]). A t-structure on \mathcal{D} is a full subcategory $\mathcal{P} \subset \mathcal{D}$ satisfying the following conditions:

- $\mathcal{P}[1] \subset \mathcal{P}$,
- define $\mathcal{P}^\perp := \{G \in \mathcal{D} \mid \text{Hom}(F, G) = 0 \text{ for all } F \in \mathcal{P}\}$, then for every object $E \in \mathcal{D}$ there is an exact triangle $F \rightarrow E \rightarrow G \rightarrow F[1]$ in \mathcal{D} with $F \in \mathcal{P}$ and $G \in \mathcal{P}^\perp$.

In addition, the t-structure $\mathcal{P} \subset \mathcal{D}$ is said to be bounded if \mathcal{P} satisfies the condition:

$$\mathcal{D} = \bigcup_{i, j \in \mathbb{Z}} \mathcal{P}^\perp[i] \cap \mathcal{P}[j].$$

For a t-structure $\mathcal{P} \subset \mathcal{D}$, we define its heart \mathcal{H} by

$$\mathcal{H} := \mathcal{P}^\perp[1] \cap \mathcal{P}.$$

It was proved in [BBD82] that \mathcal{H} becomes an abelian category.

Remark 4.3. *For a heart $\mathcal{H} \subset \mathcal{D}$ of a bounded t-structure, we have a canonical isomorphism of Grothendieck groups*

$$K(\mathcal{H}) \cong K(\mathcal{D}).$$

In the following, we only treat bounded t-structures and their hearts. The word heart always refers to the heart of some bounded t-structure.

Proposition 4.4 ([Ami09], Section 2). *There is a canonical bounded t -structure $\mathcal{P}_0 \subset \mathcal{D}_Q^N$ with the heart \mathcal{H}_0 such that the functor $\mathbf{H}_0: \mathcal{D}_Q^N \rightarrow \text{mod-}kQ$ induces an equivalence of abelian categories:*

$$\mathbf{H}_0: \mathcal{H}_0 \xrightarrow{\sim} \text{mod-}kQ.$$

We call \mathcal{H}_0 the standard heart of \mathcal{D}_Q^N . \mathcal{H}_0 is generated by simple $\Gamma_N Q$ -modules R_1, \dots, R_n which correspond to vertices $1, \dots, n$ of Q .

4.3. Spherical twists. We define some autoequivalences of \mathcal{D}_Q^N , called spherical twists, introduced by P. Seidel and R. Thomas [ST01].

An object $S \in \mathcal{D}_Q^N$ is called N -spherical if

$$\text{Hom}^i(S, S) = \begin{cases} k & \text{if } i = 0, N \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 4.5 ([ST01], Proposition 2.10). *For a spherical object $S \in \mathcal{D}_Q^N$, there is an exact autoequivalence $\Phi_S \in \text{Aut}(\mathcal{D}_Q^N)$ defined by the exact triangle*

$$\text{Hom}^\bullet(S, E) \otimes S \longrightarrow E \longrightarrow \Phi_S(E)$$

for any object $E \in \mathcal{D}_Q^N$. The inverse functor $\Phi_S^{-1} \in \text{Aut}(\mathcal{D}_Q^N)$ is given by

$$\Phi_S^{-1}(E) \longrightarrow E \longrightarrow S \otimes \text{Hom}^\bullet(E, S)^*.$$

Let $R_1, \dots, R_n \in \mathcal{H}_0$ be simple $\Gamma_N Q$ -modules. Then R_1, \dots, R_n are N -spherical in \mathcal{D}_Q^N (see Lemma 4.4 in [Kel11]). Hence they define spherical twists $\Phi_{R_1}, \dots, \Phi_{R_n} \in \text{Aut}(\mathcal{D}_Q^N)$. The Seidel-Thomas braid group $\text{Br}(\mathcal{D}_Q^N)$ is defined to be the subgroup of $\text{Aut}(\mathcal{D}_Q^N)$ generated by these spherical twists:

$$\text{Br}(\mathcal{D}_Q^N) := \langle \Phi_{R_1}, \dots, \Phi_{R_n} \rangle.$$

In the rest of this section, we assume that $Q = \overrightarrow{A}_n$.

Theorem 4.6 ([ST01], Theorem 1.2 and Theorem 1.3). *The correspondence of generators $\sigma_i \mapsto \Phi_{R_i}$ gives an isomorphism of groups*

$$B_{n+1} \cong \text{Br}(\mathcal{D}_n^N).$$

On $K(\mathcal{D}_n^N)$, a spherical twist Φ_{R_i} induces the reflection $[\Phi_{R_i}]: K(\mathcal{D}_n^N) \rightarrow K(\mathcal{D}_n^N)$ given by

$$[\Phi_{R_i}](E) = [E] - \chi(R_i, E)[R_i].$$

Recall from Section 2.1 that for a quiver \overrightarrow{A}_n , we define the lattice $(L, \langle, \rangle_\pm)$. By the map $[R_i] \mapsto \alpha_i$, we have an isomorphism of abelian groups

$$K(\mathcal{D}_n^N) \cong L.$$

Further, this map takes the Euler form χ to the bilinear form \langle, \rangle_\pm . Therefore, under the above isomorphism, spherical twists $\Phi_{R_1}, \dots, \Phi_{R_n}$ act on $K(\mathcal{D}_n^N)$ as reflections r_1^\pm, \dots, r_n^\pm in Section 2.1 :

$$[\Phi_{R_1}] = r_1^\pm, \dots, [\Phi_{R_n}] = r_n^\pm.$$

4.4. Koszul duality and spherical twists. As in Section 4.1, let $\Gamma_N Q$ be the Ginzburg dg CY $_N$ algebra of an acyclic quiver Q , $\mathcal{D}(\Gamma_N Q)$ be the derived category of right dg $\Gamma_N Q$ -modules and $\mathcal{D}_Q^N \subset \mathcal{D}(\Gamma_N Q)$ be the full subcategory consisting of right dg $\Gamma_N Q$ -modules with finite dimensional total cohomology.

In the rest of this section, we write $A := \Gamma_N Q$.

Consider the standard heart $\mathcal{H}_0 \subset \mathcal{D}_Q^N$ and its simple objects $R_1, R_2, \dots, R_n \in \mathcal{H}_0$. Set $R := \bigoplus_{i=1}^n R_i$ and define the endomorphism dg algebra B by

$$B := \mathbb{R} \operatorname{Hom}_A(R, R).$$

(We can easily check that the dg algebra B is quasi-isomorphic to the underlying graded algebra, i.e. the differential of B is zero.)

Denote by $\mathcal{D}(B)$ the derived category of right dg B -modules. The perfect derived category $\operatorname{Per}(B)$ is defined to be the smallest triangulated full subcategory of $\mathcal{D}(B)$ containing B and closed under taking direct summands.

The derived functor

$$F := \mathbb{R} \operatorname{Hom}_A(R, -): \mathcal{D}(A) \rightarrow \mathcal{D}(B)$$

has a right adjoint functor

$$G := - \otimes_B^{\mathbb{L}} R: \mathcal{D}(B) \rightarrow \mathcal{D}(A).$$

By restricting F on $\mathcal{D}_Q^N \subset \mathcal{D}(A)$ and G on $\operatorname{Per}(B) \subset \mathcal{D}(B)$, we have the following (see Section 9 in [KN13]).

Proposition 4.7. *The restriction of F on $\mathcal{D}_Q^N \subset \mathcal{D}(A)$ gives an equivalence of triangulated categories:*

$$F: \mathcal{D}_Q^N \xrightarrow{\sim} \operatorname{Per}(B).$$

Its inverse is given by the restriction of G on $\operatorname{Per}(B) \subset \mathcal{D}(B)$:

$$G: \operatorname{Per}(B) \xrightarrow{\sim} \mathcal{D}_Q^N.$$

Remark 4.8. *If $Q = \overrightarrow{A_n}$, then the graded algebra B coincides with the graded algebra A_n^N in [Tho06, Section 3] and the category $\operatorname{Per}(B)$ coincides with the category D_n^N in [Tho06, Definition 3.2].*

We set $P_i := F(R_i) = \mathbb{R} \operatorname{Hom}_A(R, R_i) \in \operatorname{Per}(B)$. Then we have $B = \bigoplus_{i=1}^n P_i$.

For each P_i , define the object $Q_i \in \operatorname{Per}(B^{op} \otimes B)$ by the cone

$$Q_i := \operatorname{Cone}(\mathbb{R} \operatorname{Hom}_B(P_i, B) \otimes P_i \rightarrow B).$$

Since R_i is N -spherical in \mathcal{D}_Q^N , the object P_i is also N -spherical in $\operatorname{Per}(B)$. We denote by $\Psi_{P_i} \in \operatorname{Aut}(\operatorname{Per}(B))$ the spherical twist along P_i .

Lemma 4.9. *The spherical twist Ψ_{P_i} is given by the derived tensor functor $- \otimes_B^{\mathbb{L}} Q_i$:*

$$\Psi_{P_i} \cong - \otimes_B^{\mathbb{L}} Q_i.$$

Proof. Since P_i is perfect, there is an isomorphism

$$N \otimes_B^{\mathbb{L}} \mathbb{R} \operatorname{Hom}_B(P_i, B) \cong \mathbb{R} \operatorname{Hom}_B(P_i, N)$$

for $N \in \operatorname{Per}(B)$. Therefore we have

$$\begin{aligned} N \otimes_B^{\mathbb{L}} Q_i &= \operatorname{Cone}(N \otimes_B^{\mathbb{L}} \mathbb{R} \operatorname{Hom}_B(P_i, B) \otimes P_i \rightarrow N \otimes_B^{\mathbb{L}} B) \\ &\cong \operatorname{Cone}(\mathbb{R} \operatorname{Hom}_B(P_i, N) \otimes P_i \rightarrow N) \\ &= \Psi_{P_i}(N). \end{aligned}$$

Since the above computation is functorial for N , the result follows. \square

Define the Seidel-Thomas braid group of $\text{Per}(B)$ by

$$\text{Br}(\text{Per}(B)) := \langle \Psi_{P_1}, \dots, \Psi_{P_n} \rangle \subset \text{Aut}(\text{Per}(B)).$$

Proposition 4.10. *If $\Psi \in \text{Br}(\text{Per}(B))$ preserves P_1, \dots, P_n , i.e. there is an isomorphism $\Psi(P_i) \cong P_i$ for each P_i , then Ψ is isomorphic to the identity functor : $\Psi \cong \text{id}$.*

Proof. Assume that

$$\Psi = \Psi_{P_{i_1}} \circ \dots \circ \Psi_{P_{i_s}}.$$

Then by Lemma 4.9, Ψ is equivalent to the derived tensor functor $-\otimes_B^{\mathbb{L}} Q$ where the object $Q \in \text{Per}(B^{op} \otimes B)$ is given by

$$Q := Q_{i_s} \otimes_B^{\mathbb{L}} \dots \otimes_B^{\mathbb{L}} Q_{i_1}.$$

Since $\Psi(P_i) \cong P_i$, we have an isomorphism $P_i \otimes_B^{\mathbb{L}} Q \cong P_i$ for each P_i . Recall that $B = \bigoplus_{i=1}^n P_i$. By taking a direct sum, we have

$$Q \cong B \otimes_B^{\mathbb{L}} Q = (\bigoplus_{i=1}^n P_i) \otimes_B^{\mathbb{L}} Q \cong (\bigoplus_{i=1}^n P_i) = B.$$

Thus, $\Psi \cong - \otimes_B^{\mathbb{L}} B$ since $Q \cong B$. For each $N \in \text{Per}(B)$,

$$N \rightarrow N \otimes_B^{\mathbb{L}} B, \quad n \mapsto n \otimes 1$$

gives a natural isomorphism $\text{id} \cong - \otimes_B^{\mathbb{L}} B$. \square

Corollary 4.11. *If $\Phi \in \text{Br}(\mathcal{D}_Q^N)$ preserves R_1, \dots, R_n , i.e. there is an isomorphism $\Phi(R_i) \cong R_i$ for each R_i , then Φ is isomorphic to the identity functor : $\Phi \cong \text{id}$.*

4.5. Simple tilted hearts. Let $\mathcal{H} \subset \mathcal{D}$ be a heart in a triangulated category \mathcal{D} . We denote by $\text{Sim } \mathcal{H}$ the set of simple objects in \mathcal{H} .

Definition 4.12. \mathcal{H} is called finite if $\text{Sim } \mathcal{H}$ is a finite set and any object of \mathcal{H} is generated by finitely many extensions of $\text{Sim } \mathcal{H}$.

For a given heart, one can construct a new heart by using the technique of tilting in [HRS96]. In particular for a finite heart $\mathcal{H} \subset \mathcal{D}$ and a simple object $S \in \mathcal{H}$, there is a new heart \mathcal{H}_S^\sharp (respectively \mathcal{H}_S^b), called the forward (respectively backward) simple tilt of \mathcal{H} by S . For the precise definition of \mathcal{H}_S^\sharp and \mathcal{H}_S^b , we refer to [KQ, Section 3]. (Our notation is based on [KQ].) In the present paper, we do not use the explicit construction of simple tilted hearts.

We say an object $S \in \mathcal{H}$ is rigid if $\text{Hom}^1(S, S) = 0$.

Proposition 4.13 ([KQ], Proposition 5.4). *Let \mathcal{D} be a triangulated category and $\mathcal{H} \subset \mathcal{D}$ be a finite heart. For any rigid simple object $S \in \mathcal{H}$, the sets of simple objects in the simple tilted hearts \mathcal{H}_S^\sharp and \mathcal{H}_S^b are given by*

$$\begin{aligned} \text{Sim } \mathcal{H}_S^\sharp &= \{S[1]\} \cup \{\psi_S^\sharp(X) \mid X \in \text{Sim } \mathcal{H}, X \neq S\} \\ \text{Sim } \mathcal{H}_S^b &= \{S[-1]\} \cup \{\psi_S^b(X) \mid X \in \text{Sim } \mathcal{H}, X \neq S\} \end{aligned}$$

where

$$\begin{aligned} \psi_S^\sharp(X) &= \text{Cone}(X \rightarrow S[1] \otimes \text{Hom}^1(X, S)^*)[-1] \\ \psi_S^b(X) &= \text{Cone}(S[-1] \otimes \text{Hom}^1(S, X) \rightarrow X). \end{aligned}$$

In addition, \mathcal{H}_S^\sharp and \mathcal{H}_S^b are also finite.

4.6. Exchange graphs. In this section, we summarize basic properties of finite hearts in \mathcal{D}_Q^N showed in [KQ].

Definition 4.14. The oriented graph with labeled edges $\text{EG}(\mathcal{D}_Q^N)$, called the exchange graph of \mathcal{D}_Q^N , is defined as follows. The set of vertices in $\text{EG}(\mathcal{D}_Q^N)$ consists of all finite hearts of \mathcal{D}_Q^N , and the set of labeled edges consists of all forward tiltings, i.e. two hearts \mathcal{H} and \mathcal{H}' are connected by the arrow $\mathcal{H} \xrightarrow{S} \mathcal{H}'$ if there is some rigid simple object $S \in \mathcal{H}$ such that $\mathcal{H}_S^\sharp = \mathcal{H}'$. The connected subgraph $\text{EG}^\circ(\mathcal{D}_Q^N) \subset \text{EG}(\mathcal{D}_Q^N)$ is defined to be the connected component which contains the standard heart \mathcal{H}_0 .

For $\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_Q^N)$ and a rigid simple object $S \in \mathcal{H}$, we define inductively

$$\mathcal{H}_S^{m\sharp} := (\mathcal{H}_S^{(m-1)\sharp})_{S[m-1]}^\sharp, \quad \mathcal{H}_S^{mb} := (\mathcal{H}_S^{(m-1)b})_{S[-m+1]}^b$$

for $m \geq 1$.

Definition 4.15. We say that a heart \mathcal{H} is

- spherical, if all its simples are spherical,
- monochromatic, if for any simples $S \neq T \in \mathcal{H}$, $\text{Hom}^\bullet(S, T)$ is concentrated in a single positive degree.

Proposition 4.16 ([KQ], Corollary 8.5). *Every heart in $\text{EG}^\circ(\mathcal{D}_Q^N)$ is finite, spherical and monochromatic. Let $\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_Q^N)$ a finite heart and consider simple objects $\text{Sim } \mathcal{H} = \{T_1, \dots, T_n\}$. Then spherical twists $\Phi_{T_1}, \dots, \Phi_{T_n}$ generate the Seidel-Thomas braid group $\text{Br}(\mathcal{D}_Q^N)$. In addition, we have*

$$\mathcal{H}_{T_i}^{(N-1)\sharp} = \Phi_{T_i}^{-1}(\mathcal{H}), \quad \mathcal{H}_{T_i}^{(N-1)b} = \Phi_{T_i}(\mathcal{H}).$$

The result implies that the action of $\text{Br}(\mathcal{D}_Q^N)$ on $\text{EG}(\mathcal{D}_Q^N)$ preserves the connected component $\text{EG}^\circ(\mathcal{D}_Q^N)$. By using the result in Section 4.4, we can show that this action is free.

Proposition 4.17. *The action of $\text{Br}(\mathcal{D}_Q^N)$ on $\text{EG}^\circ(\mathcal{D}_Q^N)$ is free.*

Proof. If $\Phi \in \text{Br}(\mathcal{D}_Q^N)$ acts trivially on $\text{EG}^\circ(\mathcal{D}_Q^N)$, then Φ preserves the vertex \mathcal{H}_0 and labeled edges connected to \mathcal{H}_0 . This implies that Φ preserves simple objects R_1, \dots, R_n in \mathcal{H}_0 . Hence by Corollary 4.11, it follows that $\Phi \cong \text{id}$. \square

Let \mathcal{H}_i ($i = 1, 2$) be the hearts of bounded t-structures \mathcal{P}_i ($i = 1, 2$). We denote by

$$\mathcal{H}_1 \leq \mathcal{H}_2$$

if they satisfy the condition $\mathcal{P}_2 \subset \mathcal{P}_1$.

We define the full subgraph of $\text{EG}^\circ(\mathcal{D}_Q^N)$ by

$$\text{EG}_N^\circ(\mathcal{H}_0) := \{ \mathcal{H} \in \text{EG}^\circ(\mathcal{D}_Q^N) \mid \mathcal{H}_0 \leq \mathcal{H} \leq \mathcal{H}_0[N-2] \}.$$

Let $\mathcal{H} \in \text{EG}(\mathcal{D}_Q^N)$ be a finite heart and $S \in \mathcal{H}$ be a simple object. Since

$$\Phi(\mathcal{H}_S^\sharp) = (\Phi(\mathcal{H}))_{\Phi(S)}^\sharp$$

for any autoequivalence $\Phi \in \text{Aut}(\mathcal{D}_Q^N)$, the element of $\text{Aut}(\mathcal{D}_Q^N)$ acts on $\text{EG}(\mathcal{D}_Q^N)$ as an automorphism of the oriented graph. Denote by $\text{Aut}^\circ(\mathcal{D}_Q^N)$ the subgroup consisting of autoequivalences which preserve the connected component $\text{EG}^\circ(\mathcal{D}_Q^N)$.

The following implies that $\text{EG}_N^\circ(\mathcal{H}_0)$ is a fundamental domain of the action of $\text{Br}(\mathcal{D}_Q^N)$ on $\text{EG}^\circ(\mathcal{D}_Q^N)$.

Theorem 4.18 ([KQ], Theorem 8.6). *There is a map of oriented graphs*

$$p_0: \text{EG}_N^\circ(\mathcal{H}_0) \longrightarrow \text{EG}^\circ(\mathcal{D}_Q^N)/\text{Br}(\mathcal{D}_Q^N)$$

such that p_0 is embedding and bijective on vertices.

4.7. Cluster categories and cluster tilting objects. In this section, we introduce the notion of an $(N-2)$ -cluster category, an $(N-2)$ -cluster tilting object and its mutation. In particular, $(N-2)$ -cluster tilting objects will play the role of intermediate between N -angulations and hearts of the derived category of the Ginzburg CY_N dga. For details of cluster categories, we refer to [KQ, Section 4].

Let Q be an acyclic quiver and $D^b(kQ)$ be the bounded derived category of finite dimensional modules over the path algebra kQ . We denote by τ the Auslander-Reiten translation of $D^b(kQ)$. The $(N-2)$ -cluster category $\mathcal{C}_{N-2}(Q)$ is defined to be the orbit category

$$\mathcal{C}_{N-2}(Q) := D^b(kQ)/\langle F^m | m \in \mathbb{Z} \rangle$$

where $F := \tau^{-1} \circ [N-2] \in \text{Aut}(D^b(kQ))$. Assume that Q has n vertices. An object $P \in \mathcal{C}_{N-2}(Q)$ is called the $(N-2)$ -cluster tilting object if

- $\text{Hom}_{\mathcal{C}_{N-2}(Q)}(P, P[i]) = 0$ for $i = 1, \dots, N-2$,
- P has n indecomposable direct summands, up to isomorphism.

Assume that an $(N-2)$ -cluster tilting object P decomposes as $P = \bigoplus_{i=1}^n P_i$. Then, one can construct a new $(N-2)$ -cluster tilting object $\mu_i^\sharp P$ (respectively $\mu_i^\flat P$), called the forward (respectively backward) mutation of P at the i -th object P_i . For the construction of $\mu_i^\sharp P$ and $\mu_i^\flat P$, we refer to [KQ, Section 4].

Definition 4.19. The cluster exchange graph $\text{CEG}_{N-2}(Q)$ is an oriented graph whose vertices are $(N-2)$ -cluster tilting objects and whose labeled edges are forward mutations, i.e. two tilting objects P and P' are connected by the arrow $P \xrightarrow{P_v} P'$ if $P' = \mu_v^\sharp P$ where P_v is an indecomposable direct summand of $P = \bigoplus_{i=1}^n P_i$.

Theorem 4.20 ([KQ], Theorem 8.6). *There is an isomorphism of oriented graphs with labeled edges*

$$\text{EG}^\circ(\mathcal{D}_Q^N)/\text{Br}(\mathcal{D}_Q^N) \cong \text{CEG}_{N-2}(Q).$$

Remark 4.21. Let $\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_Q^N)$ be a finite heart and $P \in \text{CEG}_{N-2}(Q)$ be the corresponding $(N-2)$ -cluster tilting object given by the above isomorphism. Since both $\text{EG}^\circ(\mathcal{D}_Q^N)/\text{Br}(\mathcal{D}_Q^N)$ and $\text{CEG}_{N-2}(Q)$ are n -regular graphs, the above isomorphism also gives the bijection between all simple objects S_1, \dots, S_n in \mathcal{H} and all indecomposable direct summands P_1, \dots, P_n of P :

$$\{S_1, \dots, S_n\} \cong \{P_1, \dots, P_n\}, \quad S_i \mapsto P_i.$$

4.8. Colored quivers and mutation. Following [BT09, Section 2], we introduce the notion of a colored quiver and its mutation.

An $(N - 2)$ -colored quiver \mathcal{Q} consists of vertices $\{1, \dots, n\}$ and colored arrows $i \xrightarrow{(c)} j$, where $c \in \{0, 1, \dots, N - 2\}$. Write by $q_{ij}^{(c)}$ the number of arrows from i to j of color (c) .

In addition, we assume the following conditions:

- (1) No loops, $q_{ii}^{(c)} = 0$ for all c .
- (2) Monochromaticity, if $q_{ij}^{(c)} \neq 0$, then $q^{(c')} = 0$ for $c \neq c'$.
- (3) Skew-symmetry, $q_{ij}^{(c)} = q_{ji}^{(N-2-c)}$.

Let \mathcal{Q} be a $(N - 2)$ -colored quiver satisfying additional conditions and fix a vertex v in \mathcal{Q} . Then a new $(N - 2)$ -colored quiver $\mu_v^\# \mathcal{Q} = (\tilde{q}_{ij}^{(c)})$ is defined by

$$\tilde{q}_{ij}^{(c)} = \begin{cases} q_{ij}^{(c-1)} & \text{if } v = i \\ q_{ij}^{(c+1)} & \text{if } v = j \\ \max\{0, q_{ij}^{(c)} - \sum_{t \neq c} q_{ij}^{(t)} + (q_{iv}^{(c)} - q_{iv}^{(c+1)})q_{vj}^{(0)} \\ \quad + q_{iv}^{(N-2)}(q_{vj}^{(c)} - q_{vj}^{(c-1)})\} & \text{otherwise.} \end{cases}$$

We call this operation the forward colored quiver mutation of \mathcal{Q} at a vertex v . The backward mutated colored quiver $\mu_v^b \mathcal{Q}$ can be similarly defined. (Our notation is different to the notation in [BT09]. The backward mutation $\mu_v^b \mathcal{Q}$ corresponds to the colored quiver mutation $\mu_v \mathcal{Q}$ in [BT09, Section 2].)

For each $(N - 2)$ -cluster tilting object P , we can associate an $(N - 2)$ -colored quiver $\mathcal{Q}(P)$ whose vertices $\{1, \dots, n\}$ correspond to indecomposable direct summands of $P = \bigoplus_{i=1}^n P_i$. The definition of $\mathcal{Q}(P)$ can be found in [BT09, Section 2] or [KQ, Section 4], though the precise definition will not be used later.

Theorem 4.22 ([BT09], Theorem 2.1). *Let P be an $(N - 2)$ -cluster tilting object and v be a vertex of $\mathcal{Q}(P)$ corresponding to an indecomposable direct summand P_v of P . Then we have*

$$\mu_v^\# \mathcal{Q}(P) = \mathcal{Q}(\mu_v^\# P), \quad \mu_v^b \mathcal{Q}(P) = \mathcal{Q}(\mu_v^b P).$$

Next we consider colored quivers associated to finite hearts in $\text{EG}^\circ(\mathcal{D}_Q^N)$.

Definition 4.23. For a finite heart $\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_Q^N)$, we define the colored quiver $\mathcal{Q}(\mathcal{H})$ whose vertices $1, \dots, n$ correspond to simple objects $S_1, \dots, S_n \in \text{Sim } \mathcal{H}$ and the number of arrows $q_{ij}^{(c)}$ from i to j of color (c) is given by $q_{ij}^{(c)} := \text{hom}^{c+1}(S_i, S_j)$ for $i \neq j$ and $q_{ii}^{(c)} := 0$.

By definition, $\mathcal{Q}(\mathcal{H})$ has no loops. Further, the CY_N property of \mathcal{D}_Q^N implies that $\mathcal{Q}(\mathcal{H})$ is skew-symmetric and Proposition 4.16 implies the monochromaticity of $\mathcal{Q}(\mathcal{H})$. Therefore, the colored quiver $\mathcal{Q}(\mathcal{H})$ satisfies additional conditions.

Remark 4.24. *For a finite heart \mathcal{H} , we can associate a graded quiver $\mathcal{Q}^\epsilon(\mathcal{H})$, called the Ext quiver of \mathcal{H} (Definition 6.2 in [KQ]), whose vertices are simple objects of \mathcal{H} and whose degree k arrows from S_i to S_j correspond to a basis of $\text{Hom}_{\mathcal{D}_Q^N}(S_i, S_j[k])$. On the other hand, if we consider the augmented graded quiver $\mathcal{Q}^+(\mathcal{H})$ (Definition 6.1 in [KQ]) of the colored quiver $\mathcal{Q}(\mathcal{H})$, then we have $\mathcal{Q}^\epsilon(\mathcal{H}) = \mathcal{Q}^+(\mathcal{H})$.*

The following result is stated in terms of the Ext quiver and the augmented quiver in the original paper [KQ], however here we state in terms of the colored quiver.

Theorem 4.25 ([KQ], Theorem 8.10). *Let $\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_Q^N)$ be a finite heart and $P \in \text{CEG}_{N-2}(Q)$ be the corresponding $(N-2)$ -cluster tilting object given by Theorem 4.20. Then we have*

$$\mathcal{Q}(\mathcal{H}) = \mathcal{Q}(P).$$

By Theorem 4.20, Theorem 4.22 and Theorem 4.25, we have the following corollary.

Corollary 4.26. *Let $\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_Q^N)$ be a finite heart and $\mathcal{Q}(\mathcal{H})$ be the corresponding colored quiver. Then for any simple object $S_v \in \mathcal{H}$, we have*

$$\mu_v^\sharp \mathcal{Q}(\mathcal{H}) = \mathcal{Q}(\mathcal{H}_{S_v}^\sharp), \quad \mu_v^\flat \mathcal{Q}(\mathcal{H}) = \mathcal{Q}(\mathcal{H}_{S_v}^\flat).$$

Proof. Let P be the $(N-2)$ -cluster tilting object corresponding to \mathcal{H} given by Theorem 4.20. Again by Theorem 4.20, the $(N-2)$ -cluster tilting object corresponding to $\mathcal{H}_{S_v}^\sharp$ is the forward mutation $\mu_v^\sharp P$. Then, Theorem 4.22 and Theorem 4.25 imply that

$$\mu_v^\sharp \mathcal{Q}(\mathcal{H}) = \mu_v^\sharp \mathcal{Q}(P) = \mathcal{Q}(\mu_v^\sharp P) = \mathcal{Q}(\mathcal{H}_{S_v}^\sharp).$$

□

4.9. Colored quiver lattices. In this section, we define a lattice associated to a colored quiver.

For a $(N-2)$ -colored quiver \mathcal{Q} with vertices $\{1, \dots, n\}$, we introduce a free abelian group $L_{\mathcal{Q}}$ generated by vertices of \mathcal{Q} . Denote by $\alpha_1, \dots, \alpha_n$ generators corresponding to vertices $1, \dots, n$. Then we have

$$L_{\mathcal{Q}} = \bigoplus_{i=1}^n \mathbb{Z} \alpha_i.$$

Define the bilinear form $\langle \cdot, \cdot \rangle : L_{\mathcal{Q}} \times L_{\mathcal{Q}} \rightarrow \mathbb{Z}$ by

$$\langle \alpha_i, \alpha_j \rangle := \delta_{ij} + (-1)^N \delta_{ij} - \sum_{c=0}^{N-2} (-1)^c q_{ij}^{(c)}.$$

Since $q_{ij}^{(c)} = q_{ji}^{(N-2-c)}$, the bilinear form $\langle \cdot, \cdot \rangle$ is symmetric if N is even, and skew-symmetric if N is odd.

Let $\mathcal{Q}(\mathcal{H})$ be the colored quiver of a heart $\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_Q^N)$ with simple objects S_1, \dots, S_n . We write by α_i the basis of colored quiver lattice $L_{\mathcal{Q}(\mathcal{H})}$ which corresponds to a vertex i of $\mathcal{Q}(\mathcal{H})$. Then we have the following obvious result.

Lemma 4.27. *There is an isomorphism of abelian groups*

$$\lambda_{\mathcal{H}} : L_{\mathcal{Q}(\mathcal{H})} \rightarrow K(\mathcal{H}) \cong K(\mathcal{D}_Q^N)$$

defined by $\alpha_i \mapsto [S_i]$. The map λ takes the bilinear form $\langle \cdot, \cdot \rangle$ on $L_{\mathcal{Q}(\mathcal{H})}$ to the Euler form χ on $K(\mathcal{D}_Q^N)$.

Proof. Since

$$K(\mathcal{H}) \cong \bigoplus_{i=1}^n \mathbb{Z}[S_i],$$

$\lambda_{\mathcal{H}}$ is an isomorphism. Recall from Definition 4.23 that $q_{ij}^{(c)} = \text{hom}^{c+1}(S_i, S_j)$. In addition, since \mathcal{H} is spherical, $\text{hom}^0(S_i, S_j) = \text{hom}^N(S_i, S_j) = \delta_{ij}$. Hence the result follows. \square

Let $\mathcal{H}_{S_v}^{\sharp}$ be the forward simple tilted heart and S'_1, \dots, S'_1 be its simple objects. By Proposition 4.13, we have

$$(4.1) \quad S'_j = \begin{cases} S_v[1] & \text{if } j = v \\ \psi_{S_v}^{\sharp}(S_j) & \text{if } j \neq v \end{cases}$$

where

$$\psi_{S_v}^{\sharp}(S_j) = \text{Cone}(S_j \rightarrow S_v[1] \otimes \text{Hom}^1(S_v, S_j)^*)[-1].$$

Let $\mu_v^{\sharp}\mathcal{Q}(\mathcal{H})$ be the forward colored quiver mutation of $\mathcal{Q}(\mathcal{H})$ at v . Consider two colored quiver lattices

$$L_{\mathcal{Q}(\mathcal{H})} = \bigoplus_{j=1}^n \mathbb{Z}\alpha_j, \quad L_{\mu_v^{\sharp}\mathcal{Q}(\mathcal{H})} = \bigoplus_{j=1}^n \mathbb{Z}\alpha'_j.$$

We define a linear map $F_v : L_{\mu_v^{\sharp}\mathcal{Q}(\mathcal{H})} \rightarrow L_{\mathcal{Q}(\mathcal{H})}$ by

$$F_v(\alpha'_j) := \begin{cases} -\alpha_v & \text{if } j = v \\ \alpha_j + q_{vj}^{(0)}\alpha_v & \text{if } j \neq v. \end{cases}$$

Since $\mu_v^{\sharp}\mathcal{Q}(\mathcal{H}) = \mathcal{Q}(\mathcal{H}_{S_v}^{\sharp})$ by Corollary 4.26, we have $L_{\mu_v^{\sharp}\mathcal{Q}(\mathcal{H})} \cong L_{\mathcal{Q}(\mathcal{H}_{S_v}^{\sharp})}$. Hence from Lemma 4.27, we have an isomorphism

$$\lambda_{\mathcal{H}_{S_v}^{\sharp}} : L_{\mu_v^{\sharp}\mathcal{Q}(\mathcal{H})} \longrightarrow K(\mathcal{H}_{S_v}^{\sharp}) \cong K(\mathcal{D}_Q^N)$$

defined by $\alpha'_j \mapsto [S'_j]$.

With the above notations, we have the following diagram.

Lemma 4.28. *Let $\lambda_{\mathcal{H}} : L_{\mathcal{Q}(\mathcal{H})} \rightarrow K(\mathcal{D}_n^N)$ and $\lambda_{\mathcal{H}_{S_v}^{\sharp}} : L_{\mu_v^{\sharp}\mathcal{Q}(\mathcal{H})} \rightarrow K(\mathcal{D}_n^N)$ be the isomorphisms given by Lemma 4.27. Then the diagram*

$$\begin{array}{ccc} L_{\mu_v^{\sharp}\mathcal{Q}(\mathcal{H})} & \xrightarrow{F_v} & L_{\mathcal{Q}(\mathcal{H})} \\ & \searrow \lambda_{\mathcal{H}_{S_v}^{\sharp}} & \swarrow \lambda_{\mathcal{H}} \\ & & K(\mathcal{D}_Q^N) \end{array}$$

commutes.

Proof. First note that by the equation (4.1), we have

$$[S'_j] = \begin{cases} -[S_v] & \text{if } j = v \\ [S_j] + \text{hom}^1(S_v, S_j)[S_v] & \text{if } j \neq v \end{cases}$$

in $K(\mathcal{D}_Q^N)$. Recall from Definition 4.23 that $q_{vj}^{(0)} = \text{hom}^1(S_v, S_j)$. Since $\lambda_{\mathcal{H}}(\alpha_j) = [S_j]$ and $\lambda_{\mathcal{H}_{S_v}^{\sharp}}(\alpha'_j) = [S'_j]$, we have the result. \square

5. BRIDGELAND STABILITY CONDITIONS

5.1. Spaces of stability conditions. In this section, we recall the notion of stability conditions on triangulated categories introduced by T. Bridgeland [Bri07] and collect some results for spaces of stability conditions.

Definition 5.1. Let \mathcal{D} be a triangulated category and $K(\mathcal{D})$ be its K -group. A stability condition $\sigma = (Z, \mathcal{P})$ on \mathcal{D} consists of a group homomorphism $Z: K(\mathcal{D}) \rightarrow \mathbb{C}$ called central charge and a family of full additive subcategories $\mathcal{P}(\phi) \subset \mathcal{D}$ for $\phi \in \mathbb{R}$ satisfying the following conditions:

- (a) if $0 \neq E \in \mathcal{P}(\phi)$, then $Z(E) = m(E) \exp(i\pi\phi)$ for some $m(E) \in \mathbb{R}_{>0}$,
- (b) for all $\phi \in \mathbb{R}$, $\mathcal{P}(\phi + 1) = \mathcal{P}(\phi)[1]$,
- (c) if $\phi_1 > \phi_2$ and $A_i \in \mathcal{P}(\phi_i)$ ($i = 1, 2$), then $\text{Hom}_{\mathcal{D}}(A_1, A_2) = 0$,
- (d) for $0 \neq E \in \mathcal{D}$, there is a finite sequence of real numbers

$$\phi_1 > \phi_2 > \cdots > \phi_m$$

and a collection of exact triangles

$$0 = E_0 \longrightarrow E_1 \longrightarrow E_2 \longrightarrow \cdots \longrightarrow E_{m-1} \longrightarrow E_m = E$$

with $A_i \in \mathcal{P}(\phi_i)$ for all i .

It follows from the definition that the subcategory $\mathcal{P}(\phi) \subset \mathcal{D}$ is an abelian category (see Lemma 5.2 in [Bri07]). Nonzero objects in $\mathcal{P}(\phi)$ are called semistable of phase ϕ in σ and simple objects in $\mathcal{P}(\phi)$ are called stable of phase ϕ in σ .

For a stability condition $\sigma = (Z, \mathcal{P})$, we introduce the set of semistable classes $\mathcal{C}^{\text{ss}}(\sigma) \subset K(\mathcal{D})$ by

$$\mathcal{C}^{\text{ss}}(\sigma) := \{ \alpha \in K(\mathcal{D}) \mid \text{there exists a semistable object } E \in \mathcal{D} \text{ in } \sigma \text{ such that } [E] = \alpha \}.$$

We always assume that our stability conditions satisfy the additional assumption, called the support property in [KS].

Definition 5.2. Let $\|\cdot\|$ be some norm on $K(\mathcal{D}) \otimes \mathbb{R}$. A stability condition $\sigma = (Z, \mathcal{P})$ satisfies the support property if there is a some constant $C > 0$ such that

$$C \cdot |Z(\alpha)| > \|\alpha\|$$

for all $\alpha \in \mathcal{C}^{\text{ss}}(\sigma)$.

Let $\text{Stab}(\mathcal{D})$ be the set of all stability conditions on \mathcal{D} with the support property.

In [Bri07], Bridgeland introduced a natural topology on the space $\text{Stab}(\mathcal{D})$ characterized by the following theorem.

Theorem 5.3 ([Bri07], Theorem 1.2). *The space $\text{Stab}(\mathcal{D})$ has the structure of a complex manifold and the projection map of central charges*

$$\mathcal{Z}: \text{Stab}(\mathcal{D}) \longrightarrow \text{Hom}_{\mathbb{Z}}(K(\mathcal{D}), \mathbb{C})$$

defined by $(Z, \mathcal{P}) \mapsto Z$ is a local isomorphism of complex manifolds onto an open subset of $\text{Hom}_{\mathbb{Z}}(K(\mathcal{D}), \mathbb{C})$.

In [Bri07], Bridgeland gave the alternative description of a stability condition on \mathcal{D} as a pair of a bounded t-structure and a central charge on its heart.

Definition 5.4. Let \mathcal{H} be an abelian category and let $K(\mathcal{H})$ be its Grothendieck group. A central charge on \mathcal{H} is a group homomorphism $Z: K(\mathcal{H}) \rightarrow \mathbb{C}$ such that for any nonzero object $0 \neq E \in \mathcal{H}$, the complex number $Z(E)$ lies in semi-closed upper half-plane $H = \{re^{i\pi\phi} \in \mathbb{C} \mid r \in \mathbb{R}_{>0}, \phi \in (0, 1]\}$.

The real number

$$\phi(E) := \frac{1}{\pi} \arg Z(E) \in (0, 1]$$

for $0 \neq E \in \mathcal{H}$ is called the phase of E .

A nonzero object $0 \neq E \in \mathcal{H}$ is said to be Z -(semi)stable if for any nonzero proper subobject $0 \neq A \subsetneq E$ satisfies $\phi(A) < (\leq)\phi(E)$.

For a central charge $Z: K(\mathcal{H}) \rightarrow \mathbb{C}$, the set of semistable classes can be defined similarly.

Proposition 5.5 ([Bri07], Proposition 5.3). *Let \mathcal{D} be a triangulated category. To give a stability condition on \mathcal{D} is equivalent to giving the heart $\mathcal{H} \subset \mathcal{D}$ of a bounded structure on \mathcal{D} and a central charge with the Harder-Narasimhan property on \mathcal{H} .*

For details of the Harder-Narasimhan property (HN property), we refer to [Bri07, Section 2].

We denote by $\text{Stab}(\mathcal{H})$ the set of central charges on the heart $\mathcal{H} \subset \mathcal{D}$ with the HN property and the support property.

5.2. Simple tilting and gluing. Here by using Proposition 5.5, we construct stability conditions on a finite heart.

Let $\mathcal{H} \subset \mathcal{D}$ be a finite heart with simple objects $\{S_1, \dots, S_n\}$. Then $K(\mathcal{H}) \cong \bigoplus_{i=1}^n \mathbb{Z}[S_i]$. For any point $(z_1, \dots, z_n) \in H^n$, the central charge $Z: K(\mathcal{H}) \rightarrow \mathbb{C}$ is constructed by $Z(S_i) := z_i$. Conversely, for a given central charge $Z: K(\mathcal{H}) \rightarrow \mathbb{C}$, the complex number $Z(S_i)$ lies in H for all i . Hence Z determines a point $(z_1, \dots, z_n) \in H^n$ where $z_i := Z(S_i)$. As a result, the set of central charges on \mathcal{H} is isomorphic to H^n .

Lemma 5.6 ([Bri09], Lemma 5.2). *Let $Z: K(\mathcal{H}) \rightarrow \mathbb{C}$ be a central charge given by the above construction. Then Z has a Harder-Narasimhan property. In particular, we have*

$$\text{Stab}(\mathcal{H}) \cong H^n.$$

Let $\mathcal{H} \subset \mathcal{D}$ be a finite rigid heart with simple objects $\{S_1, \dots, S_n\}$. From the above results, there is a natural inclusion

$$\text{Stab}(\mathcal{H}) \subset \text{Stab}(\mathcal{D}).$$

From such a heart, we can construct the another hearts $\mathcal{H}_{S_i}^\sharp$ (or $\mathcal{H}_{S_i}^b$) by simple tilting as in Section 4.5. The relationship between $\text{Stab}(\mathcal{H})$ and $\text{Stab}(\mathcal{H}_{S_i}^\sharp)$ (or $\text{Stab}(\mathcal{H}_{S_i}^b)$) in $\text{Stab}(\mathcal{D})$ is given by the following lemma.

Lemma 5.7 ([BS15], Lemma 7.9). *Let $\sigma = (Z, \mathcal{P}) \in \text{Stab}(\mathcal{D})$ lies on a unique codimension one boundary of the region $\text{Stab}(\mathcal{H})$ so that $\text{Im } Z(S_i) = 0$ for a unique simple object S_i . Then there is a neighborhood $U \subset \text{Stab}(\mathcal{D})$ such that one of the following holds:*

- (1) $Z(S_i) \in \mathbb{R}_{>0}$ and $U \subset \text{Stab}(\mathcal{H}) \cup \text{Stab}(\mathcal{H}_{S_i}^\sharp)$,
- (2) $Z(S_i) \in \mathbb{R}_{<0}$ and $U \subset \text{Stab}(\mathcal{H}) \cup \text{Stab}(\mathcal{H}_{S_i}^b)$.

Consider $\mathcal{D} = \mathcal{D}_n^N$, which is the derived category of finite total dimensional dg modules over the Ginzburg dg algebra $\Gamma_N \overrightarrow{A}_n$ associated to the A_n -quiver (see section 4.1). By Proposition 4.4, there is a special heart $\mathcal{H}_0 \subset \mathcal{D}_n^N$, called the standard heart. Since \mathcal{H}_0 is finite, we have $\text{Stab}(\mathcal{H}_0) \cong H^n$.

Definition 5.8. The distinguished connected component $\text{Stab}^\circ(\mathcal{D}_n^N) \subset \text{Stab}(\mathcal{D}_n^N)$ is defined to be the connected component which contains the connected subset $\text{Stab}(\mathcal{H}_0)$.

By using the result in [Woo10], Qiu described the space $\text{Stab}^\circ(\mathcal{D}_n^N)$ as follows.

Proposition 5.9 ([Qiu15], Corollary 5.3). *The distinguished connected component $\text{Stab}^\circ(\mathcal{D}_n^N)$ is given by*

$$\text{Stab}^\circ(\mathcal{D}_n^N) = \bigcup_{\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_n^N)} \text{Stab}(\mathcal{H}).$$

5.3. Group actions. For the space $\text{Stab}(\mathcal{D})$, we introduce two commuting group actions.

First we consider the action of $\text{Aut}(\mathcal{D})$ on $\text{Stab}(\mathcal{D})$. Let $\Phi \in \text{Aut}(\mathcal{D})$ be an autoequivalence of \mathcal{D} and $(Z, \mathcal{P}) \in \text{Stab}(\mathcal{D})$ be a stability condition on \mathcal{D} . Then the element $\Phi \cdot (Z, \mathcal{P}) = (Z', \mathcal{P}')$ is defined by

$$Z'(E) := Z(\Phi^{-1}(E)), \quad \mathcal{P}'(\phi) := \Phi(\mathcal{P}(\phi)),$$

where $E \in \mathcal{D}$ and $\phi \in \mathbb{R}$.

Second we define the \mathbb{C} -action on $\text{Stab}(\mathcal{D})$. For $t \in \mathbb{C}$ and $(Z, \mathcal{P}) \in \text{Stab}(\mathcal{D})$, the element $t \cdot (Z, \mathcal{P}) = (Z', \mathcal{P}')$ is defined by

$$Z'(E) := e^{-i\pi t} \cdot Z(E), \quad \mathcal{P}'(\phi) := \mathcal{P}(\phi + \text{Re}(t))$$

where $E \in \mathcal{D}$ and $\phi \in \mathbb{R}$. Clearly, this action is free.

Consider the category $\mathcal{D} = \mathcal{D}_n^N$. Recall from Section 4.3 that there is a subgroup $\text{Br}(\mathcal{D}_n^N) \subset \text{Aut}(\mathcal{D}_n^N)$ consisting of spherical twists. By Proposition 4.17 and Proposition 5.9, we have the following.

Lemma 5.10. *The action of $\text{Br}(\mathcal{D}_n^N)$ on $\text{Stab}^\circ(\mathcal{D}_n^N)$ is free and properly discontinuous.*

Proof. Proposition 4.17 and Proposition 5.9 directly imply that the action of $\text{Br}(\mathcal{D}_n^N)$ on $\text{Stab}^\circ(\mathcal{D}_n^N)$ is free.

We show that the action is properly discontinuous. Again by Proposition 5.9, it is sufficient to prove that for a stability condition $\sigma \in \text{Stab}(\mathcal{H})$ with $\mathcal{H} \in \text{EG}^\circ(\mathcal{D}_n^N)$, there is an open neighborhood $\sigma \in U \subset \text{Stab}^\circ(\mathcal{D}_n^N)$ such that $U \cap \Phi(U) = \emptyset$ for $\text{id} \neq \Phi \in \text{Br}(\mathcal{D}_n^N)$. Let $\sigma \in \text{Int}(\text{Stab}(\mathcal{H}))$ be a stability condition in the interior of $\text{Stab}(\mathcal{H})$ and take an open neighborhood $\sigma \in U \subset \text{Int}(\text{Stab}(\mathcal{H}))$. Since $\mathcal{H} \neq \Phi(\mathcal{H})$ for $\text{id} \neq \Phi \in \text{Br}(\mathcal{D}_n^N)$, $\text{Stab}(\mathcal{H}) \cap \text{Stab}(\Phi(\mathcal{H})) = \emptyset$. Hence $U \cap \Phi(U) = \emptyset$. For a general element $\sigma \in \text{Stab}(\mathcal{H})$, there is some complex number $t \in \mathbb{C}$ such that $t \cdot \sigma \in \text{Int}(\text{Stab}(\mathcal{H}))$. Since the action of \mathbb{C} commutes with the action of $\text{Br}(\mathcal{D}_n^N)$, we can apply the same argument as for σ an element in the interior of $\text{Stab}(\mathcal{H})$. \square

6. N -ANGULATIONS AND HEARTS

6.1. **N -angulations of polygons.** In [BM08], they give a geometric description of the $(N - 2)$ -cluster category of type A_n . By using their model, we give a geometric realization of the cluster exchange graph of an $(N - 2)$ -cluster category of type A_n .

Let $\Pi_{d_n^N}$ be a d_n^N -gon. An $(N - 2)$ -diagonal of $\Pi_{d_n^N}$ is a diagonal which divides $\Pi_{d_n^N}$ into an $((N - 2)i + 2)$ -gon and an $((N - 2)(n + 1 - i) + 2)$ -gon ($i = 1, \dots, n$). A collection of $(N - 2)$ -diagonals is called non-crossing if they intersect only at the boundary of $\Pi_{d_n^N}$.

Definition 6.1. An N -angulation Δ of $\Pi_{d_n^N}$ is the maximal set of non-crossing $(N - 2)$ -diagonals of $\Pi_{d_n^N}$.

Note that the number of $(N - 2)$ -diagonals in the N -angulation Δ is just n and they divide $\Pi_{d_n^N}$ into $(n + 1)$ N -gons.

For any $(N - 2)$ -diagonal $\delta \in \Delta$, there are just two N -gons whose common edge is δ . By removing δ , we have a $(2N - 2)$ -gon. We call δ a diameter of the $(2N - 2)$ -gon.

For an N -angulation Δ and an $(N - 2)$ -diagonal $\delta \in \Delta$, we introduce the flip operation to make new N -angulations $\mu_\delta^\sharp \Delta$ (called the forward flip w.r.t. δ) and $\mu_\delta^\flat \Delta$ (called the backward flip w.r.t. δ) by the following.

For an $(N - 2)$ -diagonal δ , there is a unique $(2N - 2)$ -gon which has δ as a diameter. By rotating two boundary points of δ one step clockwise (respectively counterclockwise), we have a new $(N - 2)$ -diagonal δ^\sharp (respectively δ^\flat). Thus, we obtain a new N -angulation

$$\mu_\delta^\sharp \Delta := (\Delta \setminus \{\delta\}) \cup \{\delta^\sharp\} \quad (\mu_\delta^\flat \Delta := (\Delta \setminus \{\delta\}) \cup \{\delta^\flat\}).$$




FIGURE 5. Example of an N -angulation and its forward flip in the case $N = 4$ and $n = 3$.

Definition 6.2. Define the graph $A(\Pi_{d_n^N}, N)$ to be an oriented graph whose vertices are all N -angulations of $\Pi_{d_n^N}$, and for any two N -angulations Δ and Δ' , the arrow $\Delta \xrightarrow{\delta} \Delta'$ is given if there is some $(N - 2)$ -diagonal $\delta \in \Delta$ such that $\Delta' = \mu_\delta^\sharp \Delta$.

The geometric description of the $(N - 2)$ -cluster category of type A_n in [BM08] implies the following correspondence.

Theorem 6.3 ([BM08], Theorem 5.6). *Let Q be an A_n -quiver. Then, there is an isomorphism of oriented graphs with labeled arrows*

$$\mathbb{A}(\Pi_{d_n^N}, N) \xrightarrow{\sim} \text{CEG}_{N-2}(Q).$$

6.2. Correspondence between hearts and N -angulations. Combining the geometric description of the $(N-2)$ -cluster category of type A_n ([BM08]) with the correspondence between the cluster exchange graph and the heart exchange graph ([KQ]), we have the geometric description of the heart exchange graph for an A_n -quiver. From Theorem 4.20 and Theorem 6.3, we have the following.

Corollary 6.4. *There is an isomorphism of oriented graphs with labeled arrows*

$$\mathbb{A}(\Pi_{d_n^N}, N) \xrightarrow{\sim} \text{EG}^\circ(\mathcal{D}_n^N) / \text{Br}(\mathcal{D}_n^N).$$

For $\Delta \in \mathbb{A}(\Pi_{d_n^N}, N)$ and the corresponding heart $\mathcal{H}(\Delta) \in \text{EG}^\circ(\mathcal{D}_n^N)$ (which is determined up to modulo $\text{Br}(\mathcal{D}_n^N)$), there is a canonical bijection between $(N-2)$ -diagonals $\Delta = \{\delta_1, \dots, \delta_n\}$ and simple objects $\text{Sim } \mathcal{H}(\Delta) = \{S_1, \dots, S_n\}$:

$$(6.1) \quad \{\delta_1, \dots, \delta_n\} \xrightarrow{\sim} \{S_1, \dots, S_n\}, \quad \delta_i \mapsto S_i.$$

In addition, we have

$$\mathcal{H}(\mu_{\delta_v}^\# \Delta) \equiv \mathcal{H}(\Delta)_{S_v}^\# \pmod{\text{Br}(\mathcal{D}_n^N)}$$

where $\delta_v \in \Delta$ is an $(N-2)$ -diagonal and $S_v \in \text{Sim } \mathcal{H}(\Delta)$ is the corresponding simple object.

We fix the ambiguity of $\mathcal{H}(\Delta)$ as follows.

Remark 6.5. *By Theorem 4.18, for an N -angulation Δ we can take the corresponding heart $\mathcal{H}(\Delta)$ uniquely in the fundamental domain $\text{EG}_N^\circ(\mathcal{H}_0)$. In the following, we always assume that $\mathcal{H}(\Delta)$ is in $\text{EG}_N^\circ(\mathcal{H}_0)$.*

Lemma 6.6. *Let $\Delta \in \mathbb{A}(\Pi_{d_n^N}, N)$ be an N -angulation and $\mathcal{H}(\Delta) \in \text{EG}_N^\circ(\mathcal{H}_0)$ be the corresponding heart. Take a $(N-2)$ -diagonal $\delta_v \in \Delta$ and consider the new N -angulation $\mu_{\delta_v}^\# \Delta$. Then there is a unique spherical twist $\Phi \in \text{Br}(\mathcal{D}_n^N)$ such that*

$$\Phi \cdot \mathcal{H}(\mu_{\delta_v}^\# \Delta) = \mathcal{H}(\Delta)_{S_v}^\#.$$

Proof. By Corollary 6.4, such a spherical twist exists. In addition, since the action of $\text{Br}(\mathcal{D}_n^N)$ on $\text{EG}^\circ(\mathcal{D}_n^N)$ is free, the result follows. \square

6.3. Colored quivers from N -angulations. Following [BT09, Section 11], we associate the colored quiver $\mathcal{Q}(\Delta)$ to an N -angulation Δ .

Definition 6.7. For an N -angulation $\Delta \in \mathbb{A}(\Pi_{d_n^N}, N)$, define the $(N-2)$ -colored quiver $\mathcal{Q}(\Delta)$ whose vertices are $(N-2)$ -diagonals in Δ and the colored arrow $\delta_i \xrightarrow{(c)} \delta_j$ is given if both δ_i and δ_j lie on the same N -gon in the N -angulated d_n^N -gon and there are just c edges forming the segment of the boundary of the N -gon from δ_i to δ_j in the counterclockwise direction.

Proposition 6.8. *Let $\Delta \in \mathbb{A}(\Pi_{d_n^N}, N)$ be an N -angulation and $\mathcal{H}(\Delta) \in \text{EG}^\circ(\mathcal{D}_n^N)$ be the corresponding heart. Then there is a canonical isomorphism of colored quivers*

$$\mathcal{Q}(\Delta) = \mathcal{Q}(\mathcal{H}(\Delta)).$$

In addition, the correspondence between vertices of two colored quivers is given by the correspondence between $(N - 2)$ -diagonals and simple objects of $\mathcal{H}(\Delta)$ in Corollary 6.4.

Proof. By Proposition 11.1 in [BT09], the colored quiver $\mathcal{Q}(\Delta)$ is equal to the colored quiver $\mathcal{Q}(P(\Delta))$ associated to the cluster tilting object $P(\Delta)$ given by Theorem 6.3. On the other hand, Theorem 4.25 implies that the colored quiver $\mathcal{Q}(P(\Delta))$ coincides with the colored quiver $\mathcal{Q}(\mathcal{H}(\Delta))$. \square

The definition of the colored quiver $\mathcal{Q}(\Delta)$ is compatible with the mutation and the flip.

Lemma 6.9. *Let $\delta \in \Delta$ be an N -angulation. Then we have*

$$\mu_{\delta}^{\sharp} \mathcal{Q}(\Delta) = \mathcal{Q}(\mu_{\delta}^{\sharp} \Delta).$$

Let $\Delta = \{\delta_1, \dots, \delta_n\}$ be an N -angulation and $\mu_{\delta_v}^{\sharp} \Delta$ be the forward flip w.r.t. δ_v . By Lemma 6.8 and Lemma 6.9, we have canonical isomorphisms of colored quiver lattices

$$\begin{aligned} L_{\mathcal{Q}(\Delta)} &\cong L_{\mathcal{Q}(\mathcal{H}(\Delta))} \\ L_{\mu_{\delta_v}^{\sharp} \mathcal{Q}(\Delta)} &\cong L_{\mathcal{Q}(\mu_{\delta_v}^{\sharp} \Delta)} \cong L_{\mathcal{Q}(\mathcal{H}(\mu_{\delta_v}^{\sharp} \Delta))}. \end{aligned}$$

Let S_1, \dots, S_n and $S_1^{\sharp}, \dots, S_n^{\sharp}$ be simple objects in $\mathcal{H}(\Delta)$ and $\mathcal{H}(\mu_{\delta_v}^{\sharp} \Delta)$.

By using Lemma 4.27, we have isomorphisms of lattices (for simplicity, we write $\lambda := \lambda_{\mathcal{H}(\Delta)}$ and $\lambda^{\sharp} := \lambda_{\mathcal{H}(\mu_{\delta_v}^{\sharp} \Delta)}$)

$$\begin{aligned} \lambda: L_{\mathcal{Q}(\Delta)} &\rightarrow K(\mathcal{H}(\Delta)) \cong K(\mathcal{D}_n^N) \\ \lambda^{\sharp}: L_{\mu_{\delta_v}^{\sharp} \mathcal{Q}(\Delta)} &\rightarrow K(\mathcal{H}(\mu_{\delta_v}^{\sharp} \Delta)) \cong K(\mathcal{D}_n^N) \end{aligned}$$

given by $\lambda(\alpha_i) := [S_i]$ and $\lambda^{\sharp}(\alpha'_i) := [S_i^{\sharp}]$.

Then we have the following diagram.

Lemma 6.10. *Let $F_v: L_{\mu_{\delta_v}^{\sharp} \mathcal{Q}(\Delta)} \rightarrow L_{\mathcal{Q}(\Delta)}$ be the linear map defined in Section 4.9 and $\Phi \in \text{Br}(\mathcal{D}_n^N)$ be a unique spherical twist satisfying $\Phi \cdot \mathcal{H}(\mu_{\delta_v}^{\sharp} \Delta) = \mathcal{H}(\Delta)_{S_v}^{\sharp}$ (see Lemma 6.6). Then the diagram*

$$\begin{array}{ccc} L_{\mu_{\delta_v}^{\sharp} \mathcal{Q}(\Delta)} & \xrightarrow{F_v} & L_{\mathcal{Q}(\Delta)} \\ \lambda^{\sharp} \downarrow & & \downarrow \lambda \\ K(\mathcal{D}_n^N) & \xrightarrow{[\Phi]} & K(\mathcal{D}_n^N) \end{array}$$

commutes.

Proof. As in Section 4.9, define the isomorphism

$$\lambda' := \lambda_{\mathcal{H}(\Delta)_{S_v}^{\sharp}}: L_{\mu_{\delta_v}^{\sharp} \mathcal{Q}(\Delta)} \rightarrow K(\mathcal{H}(\Delta)_{S_v}^{\sharp}) \cong K(\mathcal{D}_n^N)$$

by $\alpha'_i \mapsto [S'_i]$ where S'_1, \dots, S'_n are simple objects in $\mathcal{H}(\Delta)_{S'_v}^\sharp$. Then, since $[\Phi] \cdot [S_i^\sharp] = [S'_i]$ by Lemma 6.6, we have a commutative diagram

$$\begin{array}{ccc} & L_{\mu_v^\sharp} \mathcal{Q}(\Delta) & \\ \lambda^\sharp \swarrow & & \searrow \lambda' \\ K(\mathcal{D}_n^N) & \xrightarrow{[\Phi]} & K(\mathcal{D}_n^N). \end{array}$$

Combining the above diagram with the diagram in Lemma 4.28, we have the result. \square

6.4. N -angulations from saddle-free differentials. Let $\phi \in Q(N, n)$ be a quadratic differential. Recall from Section 3.2 that ϕ defines d_n^N distinguished tangent directions at $\infty \in \mathbb{P}^1$ since ϕ has a pole of order $(d_n^N + 2)$ at ∞ . Consider the real oriented blow-up at $\infty \in \mathbb{P}^1$ (we replace ∞ with S^1 which corresponds to tangent directions at ∞), then we have a disk Π . By adding d_n^N points on the boundary of Π , which correspond to d_n^N distinguished tangent directions, we have a d_n^N -gon $\Pi_{d_n^N}$.

In [GMN, Section 6], they constructed ideal triangulations of marked bordered surfaces from saddle-free differentials with simple zeros (see also [BS15, Section 10.1]). Motivated by their works, we consider the following construction.

Lemma 6.11. *Let $\phi \in Q(N, n)$ be a saddle-free differential. By taking one generic trajectory from each horizontal strip in the decomposition of \mathbb{P}^1 by ϕ , we have an N -angulation Δ_ϕ of $\Pi_{d_n^N}$.*

Proof. By Lemma 3.2, there are just n horizontal strips in the decomposition, and generic trajectories taken from different horizontal strips are non-crossing. Hence, it is sufficient to prove that these generic trajectories are $(N - 2)$ -diagonals. Assume that some generic trajectory divides $\Pi_{d_n^N}$ into two disks Π_1 and Π_2 , and Π_1 contain i zeros. Then the restriction of ϕ on the interior of Π_1 is also saddle-free with i zeros. Therefore, the foliation in Π_1 has $(i - 1)$ horizontal strips and Ni separating trajectories. By the equation (3.2), the number of half planes in Π_1 is $((N - 2)i + 2)$, and this implies that Π_1 is $((N - 2)i + 2)$ -gon. \square




FIGURE 6. An example of an N -angulation from a saddle-free differential in the case $N = 4$ and $n = 3$

We write by Δ_ϕ the N -angulation determined by a saddle-free differential $\phi \in Q(N, n)$. We also write by $h(\delta)$ the horizontal strip which contains a generic trajectory δ .

Set $\Delta_\phi = \{\delta_1, \dots, \delta_n\}$ and consider the colored quiver lattice

$$L_{Q(\Delta_\phi)} = \bigoplus_{i=1}^n \mathbb{Z}\alpha_i$$

where $\alpha_1, \dots, \alpha_n$ are the basis corresponding to $\delta_1, \dots, \delta_n$. Recall from Section 3.4 that we can associate standard saddle classes $\gamma_1, \dots, \gamma_n \in H_\pm(\phi)$ to horizontal strips $h(\delta_1), \dots, h(\delta_n)$.

Lemma 6.12. *There is an isomorphism of abelian groups*

$$\mu_\phi: L_{Q(\Delta_\phi)} \rightarrow H_\pm(\phi)$$

defined by $\alpha_i \mapsto \gamma_i$. The map μ_ϕ takes the bilinear form $\langle \cdot, \cdot \rangle$ on $L_{Q(\Delta_\phi)}$ to the intersection form I_\pm on $H_\pm(\phi)$.

Proof. Since standard saddle classes $\gamma_1, \dots, \gamma_n$ form the basis of $H_\pm(\phi)$ by Lemma 3.4, the map μ_ϕ is an isomorphism.

The remaining part is to prove that $\langle \alpha_i, \alpha_j \rangle = I_\pm(\gamma_i, \gamma_j)$. First note that $\langle \alpha_i, \alpha_i \rangle = I_\pm(\gamma_i, \gamma_i) = 1 + (-1)^N$ is clear by the definitions of $\langle \cdot, \cdot \rangle$ and I_\pm . Next consider the case $q_{ij}^{(c)} = 1$ for some color c and $i \neq j$. If $c = 2m$, there are $2m+1$ separating trajectories counterclockwise from γ_i to γ_j as in the left of Figure 7. Hence the oriented foliation of the 1-form $\sqrt{\phi}$ is locally given as in the left of Figure 7. Therefore, we have $\langle \alpha_i, \alpha_j \rangle = (-1)^{2m+1} q_{ij}^{(2m)} = -1 = I_\pm(\gamma_i, \gamma_j)$.

Similarly, if $c = 2m - 1$, the oriented foliation is given as in the right of Figure 7, and we have $\langle \alpha_i, \alpha_j \rangle = (-1)^{2m} q_{ij}^{(2m-1)} = 1 = I_\pm(\gamma_i, \gamma_j)$.




FIGURE 7. Oriented foliations near a zero.

□

Let $\phi \in Q(N, n)$ be a saddle-free differential and $\Delta_\phi = \{\delta_1, \dots, \delta_n\}$ be the N -angulation associated to ϕ . By Corollary 6.4, we have simple objects $S_1, \dots, S_n \in \mathcal{H}(\Delta_\phi)$ which correspond to $\delta_1, \dots, \delta_n$.

The next result is the analogue of Lemma 10.5 in [BS15].

Corollary 6.13. *There is an isomorphism of abelian groups*

$$\nu_\phi: K(\mathcal{D}_n^N) \cong K(\mathcal{H}(\Delta_\phi)) \rightarrow H_\pm(\phi)$$

defined by $[S_i] \mapsto \gamma_i$. The map ν_ϕ takes the Euler form χ on $K(\mathcal{D}_n^N)$ to the intersection form I_\pm on $H_\pm(\phi)$.

Proof. First note that by Proposition 6.8, we have a canonical isomorphism $L_{Q(\Delta_\phi)} \cong L_{Q(\mathcal{H}(\Delta_\phi))}$. Then combining Lemma 4.27 and Lemma 6.12, the result immediately follows. \square

7. FROM QUADRATIC DIFFERENTIALS TO STABILITY CONDITIONS

7.1. Stratifications. Following [BS15, Section 5.2], we introduce the stratification on $Q(N, n)$. Many results of [BS15] for the stratification also work in our cases. For $\phi \in Q(N, n)$, denote by s_ϕ the number of saddle trajectories. Define the subsets of $Q(N, n)$ by

$$D_p := \{ \phi \in Q(N, n) \mid s_\phi \leq p \}.$$

(Note that our notation differs from that of [BS15]. Our D_p corresponds to B_{2p} in [BS15].) Since Lemma 3.2 implies that $s_\phi \leq n$, we have $D_n = Q(N, n)$. Note that the stratum B_0 consists of all saddle-free differentials. These subsets have the following property.

Lemma 7.1 ([BS15], Lemma 5.2). *The subsets $D_p \subset Q(N, n)$ form an increasing chain of dense open subsets*

$$D_0 \subset D_1 \subset \cdots \subset D_n = Q(N, n).$$

Let $F_0 := D_0$ and $F_p := D_p \setminus D_{p-1}$ for $p \geq 1$. Then we have a finite stratification

$$Q(N, n) = \bigcup_{p=0}^n F_p$$

by locally closed subset F_p .

Proposition 7.2 ([BS15], Proposition 5.5 and Proposition 5.7). *Let $p \geq 1$ and assume that $\phi \in F_p$. Then there is a neighborhood $\phi \in U \subset D_p$ and a class $0 \neq \alpha \in H_\pm(\phi)$ such that*

$$\phi \in U \cap F_p \implies Z_\phi(\alpha) \in \mathbb{R}.$$

Further, if $p \geq 2$ we can take a neighborhood $\phi \in U \subset D_p$ which satisfies that $U \cap D_{p-1}$ is connected.

Following in [BS15, Section 5.2], we define generic differentials. $\phi \in Q(N, n)$ is called generic if for any $\gamma_1, \gamma_2 \in H_\pm(\phi)$,

$$\mathbb{R}Z_\phi(\gamma_1) = \mathbb{R}Z_\phi(\gamma_2) \implies \mathbb{Z}\gamma_1 = \mathbb{Z}\gamma_2.$$

It is easy to show that generic differentials are dense in $Q(N, n)$.

A differential $\phi \in Q(N, n)$ defines the length $l_\phi(\gamma)$ of a smooth path $\gamma: [0, 1] \rightarrow \mathbb{P}^1 \setminus \{\infty\}$ by

$$l_\phi(\gamma) := \int_0^1 |\varphi(\gamma(t))|^{1/2} \left| \frac{d\gamma(t)}{dt} \right| dt$$

where $\varphi(z)$ is given by $\phi(z) = \varphi(z)dz^{\otimes 2}$.

Proposition 6.8 in [BS15] is easily extended to our cases.

Proposition 7.3 ([BS15], Proposition 6.8). *Assume that for a sequence of framed differentials*

$$(\phi_k, \theta_k) \in Q(N, n)_*^\Gamma, \quad k \geq 1,$$

the periods $Z_{\phi_k} \circ \theta_k: \Gamma \rightarrow \mathbb{C}$ converge as $k \rightarrow \infty$. Further, there is some constant $L > 0$, which is independent of k , such that any saddle connection γ of ϕ_k satisfies

$l_{\phi_k}(\gamma) \geq L$. Then there is some subsequence of $\{(\phi_k, \theta_k)\}_{k \geq 1}$ which converges in $Q(N, n)_*^\Gamma$.

7.2. Stability conditions from saddle-free differentials. Following [BS15, Section 10.4], we construct a stability condition on \mathcal{D}_n^N from a saddle-free differential $\phi \in Q(N, n)$.

Lemma 7.4. $\phi \in B_0 \subset Q(N, n)$ be a saddle-free differential. Consider the corresponding heart $\mathcal{H}(\Delta_\phi) \in \text{EG}_N^2(\mathcal{H}_0)$ and define a linear map

$$Z: K(\mathcal{H}(\Delta_\phi)) \longrightarrow \mathbb{C}$$

by $Z := Z_\phi \circ \nu_\phi$ where $Z_\phi: H_\pm(\phi) \rightarrow \mathbb{C}$ is a period of ϕ and $\nu_\phi: K(\mathcal{H}(\Delta_\phi)) \rightarrow H_\pm(\phi)$ is an isomorphism given by Corollary 6.13. Then the pair $(Z, \mathcal{H}(\Delta_\phi))$ determines a stability condition $\sigma(\phi) = (Z, \mathcal{H}(\Delta_\phi)) \in \text{Stab}^\circ(\mathcal{D}_n^N)$.

Proof. By Proposition 5.5, we only need to check that $Z(E) \in H$ for any nonzero object $E \in \mathcal{H}(\Delta_\phi)$. Let h_1, \dots, h_n be horizontal strips determined by ϕ and S_1, \dots, S_n be corresponding simple objects in $\mathcal{H}(\Delta_\phi)$. By the definition of ν in Corollary 6.13, we have $\nu(S_i) = \gamma_i$ where γ_i is the standard saddle connection associated to the horizontal strip h_i . As explained in Section 3.4, the period of γ_i satisfies that $Z_\phi(\gamma_i) \in \mathbb{H} \subset H$. Hence $Z(S_i) \in H$. \square

7.3. Wall crossing. Let $\phi_1 \in F_1$ be a differential which has just one saddle trajectory. Write by Δ_{ϕ_1} the set of $(N-2)$ -diagonals obtained by taking one generic trajectory from each horizontal strip of ϕ_1 . By Lemma 3.2, $n-1$ horizontal strips appear in the decomposition of \mathbb{P}^1 by ϕ_1 , so the number of $(N-2)$ -diagonals in Δ_{ϕ_1} is $n-1$.

By Proposition 7.2, if we take $r > 0$ sufficiently small, then for any $0 < t \leq r$, the differentials

$$\phi(t) := e^{+i\pi t} \cdot \phi_1, \quad \phi^\sharp(t) := e^{-i\pi t} \cdot \phi_1$$

are saddle-free. Write $\phi := \phi(r)$ and $\phi^\sharp := \phi^\sharp(r)$.

Let Δ_ϕ and Δ_{ϕ^\sharp} be N -angulations determined by these saddle-free differentials. Write by δ_v a unique $(N-2)$ -diagonal of $\Delta_\phi = \{\delta_1, \dots, \delta_n\}$ which is not contained in Δ_{ϕ_1} :

$$\delta_v := \Delta_\phi \setminus \Delta_{\phi_1}.$$

We recall from Section 6.1 that a new N -angulation $\mu_{\delta_v}^\sharp(\Delta_\phi)$ is defined by rotating δ_v one step clockwise.

Lemma 7.5. Two N -angulations Δ_ϕ and Δ_{ϕ^\sharp} are related by

$$\mu_{\delta_v}^\sharp(\Delta_\phi) = \Delta_{\phi^\sharp}.$$

Proof. The center of Figure 8 illustrates a unique saddle trajectory of ϕ_1 and two boundaries of half planes or horizontal strips which contain the saddle trajectory.

By Lemma 3.1, the saddle trajectory maps to the saddle connection of phase $-\theta$ or $+\theta$ ($\theta := d_n^N r/2$) for $\phi = e^{-i\pi r} \cdot \phi_1$ or $\phi^\sharp = e^{+i\pi r} \cdot \phi_1$ as the left or right of Figure 8. Comparing the left and right of Figure 8, we have the result. \square

By Lemma 6.9 and Lemma 7.5, we have

$$\mu_v^\sharp \mathcal{Q}(\Delta_\phi) = \mathcal{Q}(\Delta_{\phi^\sharp}).$$




FIGURE 8. Wall crossing.

As a result, we have a natural isomorphism of colored quiver lattices $L_{\mu_v^\# \mathcal{Q}(\Delta_\phi)} \cong L_{\mathcal{Q}(\Delta_{\phi^\#})}$.

For ϕ and $\phi^\#$, we fix the natural basis as follows:

$$\begin{aligned} L_{\mathcal{Q}(\Delta_\phi)} &\cong \bigoplus_{i=1}^n \mathbb{Z} \alpha_i, & L_{\mathcal{Q}(\Delta_{\phi^\#})} &\cong \bigoplus_{i=1}^n \mathbb{Z} \alpha'_i, \\ H_\pm(\phi) &\cong \bigoplus_{i=1}^n \mathbb{Z} \gamma_i, & H_\pm(\phi^\#) &\cong \bigoplus_{i=1}^n \mathbb{Z} \gamma'_i \end{aligned}$$

where $\gamma_1, \dots, \gamma_n$ and $\gamma'_1, \dots, \gamma'_n$ are standard saddle classes.

Let

$$F_v: L_{\mathcal{Q}(\Delta_{\phi^\#})} \longrightarrow L_{\mathcal{Q}(\Delta_\phi)}$$

be the linear isomorphism defined in Section 4.9.

The relationship between F_v and the Gauss-Manin connection

$$\text{GM}_c: H_\pm(\phi^\#) \longrightarrow H_\pm(\phi)$$

along the path $c(t) = e^{i\pi t} \cdot \phi_1$ ($t \in [-r, r]$) is given by the following diagram.

Lemma 7.6. *Let $\mu_\phi: L_{\mathcal{Q}(\Delta_\phi)} \rightarrow H_\pm(\phi)$ and $\mu_{\phi^\#}: L_{\mathcal{Q}(\Delta_{\phi^\#})} \rightarrow H_\pm(\phi^\#)$ be the isomorphisms given by Lemma 6.12. Then the diagram*

$$\begin{array}{ccc} L_{\mathcal{Q}(\Delta_{\phi^\#})} & \xrightarrow{F_v} & L_{\mathcal{Q}(\Delta_\phi)} \\ \mu_{\phi^\#} \downarrow & & \downarrow \mu_\phi \\ H_\pm(\phi^\#) & \xrightarrow{\text{GM}_c} & H_\pm(\phi) \end{array}$$

commutes.


Proof. As in Figure 9 if $q_{vj}^{(0)} = 1$, then we have

$$\text{GM}_c(\gamma'_j) = \gamma_j + \gamma_v, \quad \text{GM}_c(\gamma'_v) = -\gamma_v.$$

On the other hand if $q_{jv}^{(0)} = 0$, then we can check $\text{GM}_c(\gamma'_j) = \gamma_j$. Since $\mu_\phi(\alpha_j) = \gamma_j$ and $\mu_{\phi^\#}(\alpha'_j) = \gamma'_j$, the result follows. \square

For $0 < t \leq r$, we denote by

$$\sigma(t) := \sigma(\phi(t)), \quad \sigma^\#(t) := \sigma(\phi^\#(t))$$

FIGURE 9. Changes of cycles along the path c .

stability conditions in $\text{Stab}^\circ(\mathcal{D}_n^N)$ constructed by Lemma 7.4 for saddle-free differentials $\phi(t)$ and $\phi^\sharp(t)$.

The following result is the analogue of Proposition 10.7 in [BS15].

Proposition 7.7. *Let $\nu_\phi: K(\mathcal{D}_n^N) \rightarrow H_\pm(\phi)$ and $\nu_{\phi^\sharp}: K(\mathcal{D}_n^N) \rightarrow H_\pm(\phi^\sharp)$ be the isomorphisms given by Corollary 6.13. Then there is a unique spherical twist $\Phi \in \text{Br}(\mathcal{D}_n^N)$ such that the following (1) and (2) hold.*

(1) *The diagram*

$$\begin{array}{ccc} K(\mathcal{D}_n^N) & \xrightarrow{[\Phi]} & K(\mathcal{D}_n^N) \\ \nu_{\phi^\sharp} \downarrow & & \downarrow \nu_\phi \\ H_\pm(\phi^\sharp) & \xrightarrow{\text{GM}_c} & H_\pm(\phi) \end{array}$$

commutes.

(2)

$$\lim_{t \rightarrow +0} \sigma(t) = \lim_{t \rightarrow +0} \Phi \cdot \sigma^\sharp(t).$$

Proof. Part (1) immediately follows from Lemma 6.10 and Lemma 7.6. For the proof of part (2), we first note that by Lemma 6.6 and Lemma 7.5, there is a unique spherical twist $\Phi \in \text{Br}(\mathcal{D}_n^N)$ such that

$$\Phi \cdot \mathcal{H}(\Delta_{\phi^\sharp}) = \mathcal{H}(\Delta_\phi)_{S_v}^\sharp.$$

By the definition of $\sigma(t)$ and $\sigma^\sharp(t)$ in Lemma 7.4, $\sigma(t)$ is contained in $\text{Stab}(\mathcal{H}(\Delta_\phi))$ and $\Phi \cdot \sigma^\sharp(t)$ is contained in $\text{Stab}(\mathcal{H}(\Delta_\phi)_{S_v}^\sharp)$. Further by Part (1), central charges of $\sigma(t)$ and $\Phi \cdot \sigma^\sharp(t)$ approach the same point in $\text{Hom}(K(\mathcal{D}_n^N), \mathbb{C})$ as $t \rightarrow +0$. Hence by Lemma 5.7, we can take some open subset

$$U \subset \text{Stab}(\mathcal{H}(\Delta_\phi)) \cup \text{Stab}(\mathcal{H}(\Delta_\phi)_{S_v}^\sharp)$$

such that U is mapped isomorphically on $\text{Hom}(K(\mathcal{D}_n^N), \mathbb{C})$, and U contains both $\sigma(t)$ and $\Phi \cdot \sigma^\sharp(t)$. This implies the result. \square

7.4. Construction of isomorphisms.

Definition 7.8. Let $(\phi, \theta) \in Q(N, n)^\Gamma$ be a framed saddle-free differential and $\nu_\phi: K(\mathcal{D}_n^N) \rightarrow H_\pm(\phi)$ be an isomorphism given by Corollary 6.13. We define the isomorphism κ_ϕ by the composition of θ and ν_ϕ^{-1} :

$$\kappa_\phi := \nu_\phi^{-1} \circ \theta: \Gamma \rightarrow K(\mathcal{H}(\Delta_\phi)) \cong K(\mathcal{D}_n^N).$$

Let $(\phi_0, \theta_0) \in Q(N, n)^\Gamma$ be the fixed framed differential in Section 2.7 and $Q(N, n)_*^\Gamma$ be the connected component which contains (ϕ_0, θ_0) . In the following, we also assume that (ϕ_0, θ_0) is saddle-free.

We identify $K(\mathcal{D}_n^N)$ with Γ by using the isomorphism $\kappa_0 := \kappa_{\phi_0}$ determined by Definition 7.8:

$$\kappa_0: \Gamma \xrightarrow{\sim} K(\mathcal{D}_n^N).$$

Lemma 7.9. *Let $(\phi, \theta) \in Q(N, n)_*^\Gamma$ be a framed saddle-free differential and $\kappa_\phi: \Gamma \rightarrow K(\mathcal{D}_n^N)$ be an isomorphism given by Definition 7.8. Then there is a spherical twist $\Phi \in \text{Br}(\mathcal{D}_n^N)$ such that the diagram*

$$\begin{array}{ccc} & \Gamma & \\ \kappa_\phi \swarrow & & \searrow \kappa_0 \\ K(\mathcal{D}_n^N) & \xrightarrow{[\Phi]} & K(\mathcal{D}_n^N). \end{array}$$

commutes. In addition, if there are two such spherical twists Φ_1 and Φ_2 , then the induced linear isomorphisms on $K(\mathcal{D}_n^N)$ are the same:

$$[\Phi_1] = [\Phi_2].$$

Proof. $[\Phi_1] = [\Phi_2]$ immediately follows from

$$[\Phi_1] \circ \kappa_\phi = \kappa_0 = [\Phi_2] \circ \kappa_\phi.$$

Therefore, we only need to show that such a spherical twist exists. Take a path $c: [0, 1] \rightarrow Q(N, n)_*^\Gamma$ from (ϕ, θ) to (ϕ_0, θ_0) . By Proposition 7.2, we can deform the path c to lie in D_1 and to intersect only finitely many points of F_1 , which consists of framed differentials with just one saddle trajectory.

By applying Proposition 7.7 to each of these points, we have a spherical twist $\Phi \in \text{Br}(\mathcal{D}_n^N)$ fitting into a diagram

$$\begin{array}{ccc} K(\mathcal{D}_n^N) & \xrightarrow{[\Phi]} & K(\mathcal{D}_n^N) \\ \nu_\phi \downarrow & & \downarrow \nu_0 \\ H_\pm(\phi) & \xrightarrow{\text{GM}_c} & H_\pm(\phi_0) \end{array}$$

where ν_ϕ and $\nu_0 := \nu_{\phi_0}$ are isomorphisms given by Corollary 6.13. Further, recall the diagram

$$\begin{array}{ccc} & \Gamma & \\ \theta \swarrow & & \searrow \theta_0 \\ H_\pm(\phi) & \xrightarrow{\text{GM}_c} & H_\pm(\phi_0) \end{array}$$

in Section 2.7. Combining the above two diagrams, we have the result. \square

Let $\mathrm{Br}^0(\mathcal{D}_n^N) \subset \mathrm{Br}(\mathcal{D}_n^N)$ be the subgroup of spherical twists which act as the identity on $K(\mathcal{D}_n^N)$. We recall that $\mathrm{Br}(\mathcal{D}_n^N) \cong B_{n+1}$ by Theorem 4.6, and $\mathrm{Br}(\mathcal{D}_n^N)$ acts as the group W_\pm on $K(\mathcal{D}_n^N)$ (see Section 4.3). Hence the group $\mathrm{Br}^0(\mathcal{D}_n^N)$ is isomorphic to the group P_\pm in Section 2.1.

By completely the same argument of Proposition 11.3 in [BS15], we can construct the following equivariant holomorphic map.

Proposition 7.10. *There is a W_\pm -equivariant and \mathbb{C} -equivariant holomorphic map of complex manifolds K_N such that the diagram*

$$\begin{array}{ccc} Q(N, n)_*^\Gamma & \xrightarrow{K_N} & \mathrm{Stab}^\circ(\mathcal{D}_n^N)/\mathrm{Br}^0(\mathcal{D}_n^N) \\ & \searrow \mathcal{W}_N & \swarrow \mathcal{Z} \\ & \mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}) & \end{array}$$

commutes where \mathcal{W}_N and \mathcal{Z} are local isomorphisms in Proposition 2.5 and Theorem 5.3 respectively.

Proof. First we see the \mathbb{C} -equivariant property of K_N . Define the action of $t \in \mathbb{C}$ on $\mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$ by $Z \mapsto e^{-i\pi t} \cdot Z$ for $Z \in \mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$. Then by Definition 2.7 and Lemma 3.1, the map \mathcal{W}_N is \mathbb{C} -equivariant. On the other hand, by the definition of \mathbb{C} -action on $\mathrm{Stab}^\circ(\mathcal{D}_n^N)$ given in Section 5.3, the map \mathcal{Z} is also \mathbb{C} -equivariant. Since both \mathcal{W}_N and \mathcal{Z} are local isomorphisms by Proposition 2.5 and Theorem 5.3, if K_N exists then K_N is \mathbb{C} -equivariant.

Next we construct the map K_N . Recall from Section 7.1 that there is a stratification

$$D_0 \subset D_1 \subset \cdots \subset D_n = Q(N, n).$$

We extend the stratification of $Q(N, n)$ to $Q(N, n)_*^\Gamma$ by the obvious way (and use the same notation).

Here we give the construction of K_N only on the stratum D_0 which consists of framed saddle-free differentials. By using Proposition 7.2 and Proposition 7.7, we can extend K_N on the stratum D_p ($p \geq 1$) by doing the same argument of Proposition 11.3 in [BS15].

Let $(\phi, \theta) \in D_0$ be a framed saddle-free differential. By Lemma 7.4, we can construct a stability condition $\sigma(\phi) \in \mathrm{Stab}^\circ(\mathcal{D}_n^N)$. Let $\Phi \in \mathrm{Br}(\mathcal{D}_n^N)$ be a spherical twist for (ϕ, θ) determined by Lemma 7.9. We note that Φ is determined up to the action of $\mathrm{Br}^0(\mathcal{D}_n^N) \subset \mathrm{Br}(\mathcal{D}_n^N)$. Therefore, if we define the map $K_N: D_0 \rightarrow \mathrm{Stab}^\circ(\mathcal{D}_n^N)/\mathrm{Br}^0(\mathcal{D}_n^N)$ by

$$K_N((\phi, \theta)) := \Phi(\sigma(\phi)) \in \mathrm{Stab}^\circ(\mathcal{D}_n^N)/\mathrm{Br}^0(\mathcal{D}_n^N),$$

then K_N is well-defined.

Finally we show that the map K_N is W_\pm -equivariant. Note that the action of W_\pm preserves the stratification of $Q(N, n)_*^\Gamma$ and commutes with the action of \mathbb{C} . Hence it is sufficient to prove that K_N is W_\pm -equivariant only on D_0 , and this is clear by the construction of K_N on D_0 . \square

Let $(\phi, \theta) \in Q(N, n)_*^\Gamma$ and we set $\sigma = K_N((\phi, \theta))$. The stability condition σ is determined up to the action of $\mathrm{Br}^0(\mathcal{D}_n^N) \subset \mathrm{Aut}(\mathcal{D}_n^N)$. For a class $0 \neq \alpha \in \Gamma$, let $\mathcal{M}_\sigma(\alpha)$ be the moduli space of σ -stable objects which have the class α and the phase in $(0, 1]$ (here, we treat $\mathcal{M}_\sigma(\alpha)$ as a set).

The same argument of Proposition 11.11 in [BS15] gives the following.

Proposition 7.11. *The map K_N in Proposition 7.10 is an isomorphism.*

Proof. Here we recall the argument of Proposition 11.11 in [BS15].

We first show that K_N is injective. Let $(\phi_1, \theta_1), (\phi_2, \theta_2) \in Q(N, n)_*^\Gamma$ such that $K_N((\phi_1, \theta_1)) = K_N((\phi_2, \theta_2))$. Since the maps \mathcal{W}_N and \mathcal{Z} are local isomorphisms by Proposition 2.5 and Theorem 5.3 and K_N commutes with these two maps, we can deform ϕ_1 and ϕ_2 to be saddle-free differentials.

By the construction of K_N in Proposition 7.10, we can write $K_N((\phi_i, \theta_i)) = \Phi_i(\sigma(\phi_i))$ for some $\Phi_i \in \text{Br}(\mathcal{D}_n^N)$ determined by Lemma 7.9. Since $\Phi_1(\sigma(\phi_1))$ and $\Phi_2(\sigma(\phi_2))$ have the same heart (up to the action of $\text{Br}^0(\mathcal{D}_n^N)$), we have

$$\Phi_1 \cdot \mathcal{H}(\Delta_{\phi_1}) \equiv \Phi_2 \cdot \mathcal{H}(\Delta_{\phi_2}) \pmod{\text{Br}^0(\mathcal{D}_n^N)}.$$

Recall Remark 6.5 that $\mathcal{H}(\Delta_{\phi_1})$ and $\mathcal{H}(\Delta_{\phi_2})$ are in the fundamental domain $\text{EG}_N^\circ(\mathcal{H}_0)$. Hence Theorem 4.18 implies that $\mathcal{H}(\Delta_{\phi_1}) = \mathcal{H}(\Delta_{\phi_2})$ and $[\Phi_1] = [\Phi_2]$ as linear maps on $K(\mathcal{D}_n^N)$. As a result, we have $\Delta_{\phi_1} = \Delta_{\phi_2}$ by Corollary 6.4. Thus ϕ_1 and ϕ_2 have the same N -angulations and the same periods. Then Proposition 4.9 in [BS15] implies that $\phi_1 = \phi_2$. On the other hand, $[\Phi_1] = [\Phi_2]$ together with the diagram of Lemma 7.9 implies that $\theta_1 = \theta_2$.

Next to prove the surjectivity of K_N , we show that the image of K_N is open and closed. The injectivity of K_N together with the local isomorphism property of \mathcal{W}_N and \mathcal{Z} implies that the image of K_N is open.

Set $\sigma_k := K_N((\phi_k, \theta_k)) \in \text{Stab}^\circ(\mathcal{D}_n^N) / \text{Br}^0(\mathcal{D}_n^N)$ for the sequence of framed differentials $\{(\phi_k, \theta_k)\}_{k \geq 1}$ in $Q(N, n)_*^\Gamma$, and assume that σ_k converge to the point σ_∞ as $k \rightarrow \infty$. We can assume that (ϕ_k, θ_k) is generic saddle-free since generic saddle-free differentials are dense in $Q(N, n)_*^\Gamma$. In this setting, we show that there is a subsequence of $\{(\phi_k, \theta_k)\}_{k \geq 1}$ which converges some point in $Q(N, n)_*^\Gamma$. First note that the periods $Z_{\phi_k} \circ \theta_k$ converge as $k \rightarrow \infty$ since the central charge Z_k of σ_k also converge as $k \rightarrow \infty$ by the assumption. Set

$$M_k := \inf\{|Z_k(E)| \in \mathbb{R}_{\geq 0} \mid [E] \in \mathcal{C}^{\text{ss}}(\sigma_k)\} \quad \text{for } k = 1, \dots, \infty$$

where Z_k is the central charge of σ_k . The support property of σ_k implies that $M_k > 0$. Since $M_k \rightarrow M_\infty$ as $k \rightarrow \infty$, there is some constant $L > 0$ such that $M_k \geq L$ for all k . By Theorem 11.6 and Proposition 11.7 in [BS15], the length $l_{\phi_k}(\gamma)$ of the saddle connection γ for ϕ_k coincides with the mass $|Z_k(S_\gamma)|$ of the corresponding stable object S_γ . Thus we can apply Proposition 7.3 for $\{(\phi_k, \theta_k)\}_{k \geq 1}$, and obtain the convergent subsequence. \square

Theorem 7.12. *There is a B_{n+1} -equivariant and \mathbb{C} -equivariant isomorphism of complex manifolds \widetilde{K}_N such that the diagram*

$$\begin{array}{ccc} \widetilde{M}_n & \xrightarrow{\widetilde{K}_N} & \text{Stab}^\circ(\mathcal{D}_n^N) \\ & \searrow \mathcal{W}_N & \swarrow \mathcal{Z} \\ & \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}) & \end{array}$$

commutes.

Proof. First recall from Corollary 2.9 that \widetilde{M}_n is the universal cover of the space $Q(N, n)_*^\Gamma$ and $\widetilde{M}_n / P_\pm \xrightarrow{\sim} Q(N, n)_*^\Gamma$. Further, by Proposition 7.10 and Proposition

7.11, we have the isomorphism

$$\widetilde{M}_n/P_{\pm} \xrightarrow{\sim} \text{Stab}^{\circ}(\mathcal{D}_n^N)/\text{Br}^0(\mathcal{D}_n^N).$$

Since $\text{Br}^0(\mathcal{D}_n^N) \cong P_{\pm}$ and the quotient map $\text{Stab}^{\circ}(\mathcal{D}_n^N) \rightarrow \text{Stab}^{\circ}(\mathcal{D}_n^N)/\text{Br}^0(\mathcal{D}_n^N)$ is a covering map by Lemma 5.10, we can lift the map K_N to the B_{n+1} -equivariant isomorphism

$$\widetilde{K}_N: \widetilde{M}_n \xrightarrow{\sim} \text{Stab}^{\circ}(\mathcal{D}_n^N).$$

□

Corollary 7.13. *The distinguished connected component $\text{Stab}^{\circ}(\mathcal{D}_n^N)$ is contractible.*

Proof. Since \widetilde{M}_n is contractible (for example, see Lemma 1.28 in [KT08]), the result follows. □

7.5. Autoequivalence groups. In this section, we determine the group of autoequivalences $\text{Aut}^{\circ}(\mathcal{D}_n^N)$ which preserves the distinguished connected component of the exchange graph $\text{EG}^{\circ}(\mathcal{D}_n^N)$.

Lemma 7.14. *Let $\Phi \in \text{Aut}^{\circ}(\mathcal{D}_n^N)$ an autoequivalence which acts trivially on $\text{EG}^{\circ}(\mathcal{D}_n^N)$. Then $\Phi \cong \text{id}$.*

Proof. Since Φ preserves all labeled edges in $\text{EG}^{\circ}(\mathcal{D}_n^N)$, it preserves simple objects R_1, \dots, R_n in the standard heart \mathcal{H}_0 , i.e. $R_i \cong \Phi(R_i)$. We note that the dimension of the homomorphism space between R_i and R_j is at most one because the number of arrows between two vertices in the A_n -quiver is at most one. Hence Φ preserves the homomorphism space $\text{Hom}_{\mathcal{D}_n^N}(R_i, R_j)$. Since \mathcal{D}_n^N is generated by R_1, \dots, R_n and their shifts, Φ is isomorphic to the identity functor. □

Let $\Pi_{d_n^N}$ be a d_n^N -gon as in Section 6.1. The mapping class group of $\Pi_{d_n^N}$ is given by a cyclic group $\mathbb{Z}/d_n^N\mathbb{Z}$. We assume that the generator $1 \in \mathbb{Z}/d_n^N\mathbb{Z}$ corresponds to one step clockwise rotation of $\Pi_{d_n^N}$.

Theorem 7.15 ([BS15], Theorem 9.10). *Assume that $n \geq 2$. Then there is a short exact sequence of groups*

$$1 \rightarrow \text{Br}(\mathcal{D}_n^N) \rightarrow \text{Aut}^{\circ}(\mathcal{D}_n^N) \rightarrow \mathbb{Z}/d_n^N\mathbb{Z} \rightarrow 1.$$

Proof. The same argument of Theorem 9.10 in [BS15] gives the following short exact sequence:

$$1 \rightarrow \text{Br}(\mathcal{D}_n^N) \rightarrow \text{Aut}^{\circ}(\mathcal{D}_n^N)/\text{Nil}^{\circ}(\mathcal{D}_n^N) \rightarrow \mathbb{Z}/d_n^N\mathbb{Z} \rightarrow 1$$

where $\text{Nil}^{\circ}(\mathcal{D}_n^N)$ is the subgroup of $\text{Aut}^{\circ}(\mathcal{D}_n^N)$ consisting of autoequivalences which act trivially on $\text{EG}^{\circ}(\mathcal{D}_n^N)$. Lemma 7.14 implies that $\text{Nil}^{\circ}(\mathcal{D}_n^N) = 1$. □

For an integer $m \neq 0$, let $\mathbb{Z}[m] \subset \text{Aut}^{\circ}(\mathcal{D}_n^N)$ be the infinite cyclic group generated by the degree m shift functor $[m] \in \text{Aut}^{\circ}(\mathcal{D}_n^N)$.

Recall from Theorem 4.6 that there is an isomorphism

$$B_{n+1} \xrightarrow{\sim} \text{Br}(\mathcal{D}_n^N), \quad \sigma_i \mapsto \Phi_{S_i}.$$

By Lemma 4.14 in [ST01], the generator of center $C \in B_{n+1}$ (see Section 2.1) is mapped to the shift functor $[-d_n^N] \in \text{Br}(\mathcal{D}_n^N)$ under the above isomorphism.

Define the diagonal action of the element $d_n^N \in d_n^N\mathbb{Z}$ on $B_{n+1} \times \mathbb{Z}$ by

$$(\sigma, l) \mapsto (\sigma \cdot C, l + d_n^N) \quad \text{for } (\sigma, l) \in B_{n+1} \times \mathbb{Z}.$$

Theorem 7.16. *Assume that $n \geq 2$. Then the group $\text{Aut}^\circ(\mathcal{D}_n^N)$ is given by*

$$\text{Aut}^\circ(\mathcal{D}_n^N) \cong (B_{n+1} \times \mathbb{Z}) / d_n^N \mathbb{Z}$$

where $1 \in \mathbb{Z}$ acts as the shift functor $[1] \in \text{Aut}^\circ(\mathcal{D}_n^N)$.

Proof. Consider the group homomorphism

$$\rho: B_{n+1} \times \mathbb{Z} \longrightarrow \text{Aut}^\circ(\mathcal{D}_n^N)$$

defined by $(\sigma_i, l) \mapsto \Phi_{S_i}[l]$.

By Definition 2.7, $1 \in \mathbb{C}$ acts as a rotation of the phase $2\pi/d_n^N$ in the clockwise direction on $Q(N, n)$. This implies that single shift functor $[1] \in \text{Aut}^\circ(\mathcal{D}_n^N)$ is mapped to the generator $1 \in \mathbb{Z}/d_n^N \mathbb{Z}$ under the map of the short exact sequence in Theorem 7.15. Hence ρ is surjective. On the other hand, the kernel of ρ is given by

$$\{(C^m, md_n^N) \in B_{n+1} \times \mathbb{Z} \mid m \in \mathbb{Z}\}$$

since $\rho(C) = [-d_n^N]$. □

Remark 7.17. *In the case $n = 1$, the direct calculation gives that $\text{Br}(\mathcal{D}_1^N) \cong \mathbb{Z}[-N + 1]$ and $\text{Aut}^\circ(\mathcal{D}_1^N) \cong \mathbb{Z}[1]$.*

REFERENCES

- [Ami09] C Amiot. Cluster categories for algebras of global dimension 2 and quivers with potential. *Ann. Inst. Fourier (Grenoble)*, 59(6):2525–2590, 2009.
- [BBD82] A. A. Beilinson, J. Bernstein, and P. Deligne. Faisceaux pervers. In *Analysis and topology on singular spaces, I (Luminy, 1981)*, volume 100 of *Astérisque*, pages 5–171. Soc. Math. France, Paris, 1982.
- [BM08] K. Baur and R. J. Marsh. A geometric description of m -cluster categories. *Trans. Amer. Math. Soc.*, 360(11):5789–5803, 2008.
- [Bri06] T. Bridgeland. Stability conditions on a non-compact Calabi-Yau threefold. *Comm. Math. Phys.*, 266(3):715–733, 2006.
- [Bri07] T. Bridgeland. Stability conditions on triangulated categories. *Ann. of Math.*, 166(2):317–345, 2007.
- [Bri09] T. Bridgeland. Spaces of stability conditions. In *Algebraic geometry—Seattle 2005. Part I*, volume 80 of *Proc. Sympos. Pure Math.*, pages 1–21. Amer. Math. Soc., Providence, RI, 2009.
- [BS15] T. Bridgeland and I. Smith. Quadratic differentials as stability conditions. *Publ. Math. de l’IHES.*, 121:155–278, 2015.
- [BQS] T. Bridgeland, Y. Qiu and T. Sutherland. Stability conditions and the A_2 quiver. arXiv:1406.2566.
- [BT09] A. B. Buan and H. Thomas. Coloured quiver mutation for higher cluster categories. *Adv. Math.*, 222(3):971–995, 2009.
- [Dou02] R. Douglas. Dirichlet branes, homological mirror symmetry, and stability. In *Proceedings of the International Congress of Mathematicians, Vol. III (Beijing, 2002)*, pages 395–408, Beijing, 2002. Higher Ed. Press.
- [Dub04] B. Dubrovin. On almost duality for Frobenius manifolds. In *Geometry, topology, and mathematical physics*, volume 212 of *Amer. Math. Soc. Transl. Ser. 2.*, pages 75–132. Amer. Math. Soc, Providence, RI, 2004.
- [Gin] V. G. Ginzburg. Calabi-Yau algebras. arXiv:math/0612139.
- [GMN] D. Gaiotto, G. Moore, and A. Neitzke. Wall-crossing, Hitchin Systems, and the WKB Approximation. arXiv:0907.3987.
- [HRS96] D. Happel, I. Reiten, and S. O. Smalø. Tilting in abelian categories and quasitilted algebras. *Mem. Amer. Math. Soc.*, 120(575):viii+ 88, 1996.
- [IUU10] A. Ishii, K. Ueda, and H. Uehara. Stability conditions on A_n -singularities. *J. Differential Geom.*, 84(1):87–126, 2010.
- [Kel11] B Keller. Deformed Calabi-Yau completions. *J. Reine Angew. Math.*, 654:125–180, 2011. With an appendix by Michel Van den Bergh.

- [KN13] B. Keller and P. Nicolás. Weight structures and simple dg modules for positive dg algebras *Int. Math. Res. Not.*, 1028–1078, 2013.
- [KQ] K. King and Y. Qiu. Exchange graphs and Ext quivers. to appear in *Adv. Math.*, arXiv:1109.2924v3.
- [KS] M. Kontsevich and Y. Soibelman. Stability structures, motivic Donaldson-Thomas invariants and cluster transformations. arXiv:0811.2435.
- [KS02] M. Khovanov and P. Seidel. Quivers, floer cohomology, and braid group actions. *J. Amer. Math. Soc.*, 15:203–271, 2002.
- [KT08] C. Kassel and V. Turaev. *Braid groups*, volume 247 of *Graduate Texts in Mathematics*. Springer, New York, 2008. With the graphical assistance of Olivier Dodane.
- [KY11] B. Keller and D. Yang. Derived equivalences from mutations of quivers with potential. *Adv. Math.*, 226(3):2118–2168, 2011.
- [Qiu15] Y. Qiu. Stability conditions and quantum dilogarithm identities for Dynkin quivers. *Adv. Math.*, 269:220–264, 2015.
- [Sai83] K. Saito. Period Mapping Associated to a Primitive Form. In *Publ. RIMS*. 19:1231–1264, 1983.
- [ST01] P. Seidel and R. P. Thomas. Braid group actions on derived categories of coherent sheaves. *Duke Math. J.*, 108(1):37–108, 2001.
- [Str84] K. Strebel. *Quadratic differentials*, volume 5 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1984.
- [Sut] T. Sutherland. The modular curve as the space of stability conditions of a CY3 algebra. arXiv:1111.4184.
- [Tho06] R. P. Thomas. Stability conditions and the braid group. *Comm. Anal. Geom.*, 14(1):135–161, 2006.
- [TY02] R.P. Thomas and S.T. Yau. Special Lagrangians, stable bundles and mean curvature flow. *Comm. Anal. Geom.*, 10:1075–1113, 2002.
- [Woo10] J. Woolf. Stability conditions, torsion theories and tilting. *J. Lon. Math. Soc.*, 82:663–682, 2010.

KAVLI INSTITUTE FOR THE PHYSICS AND MATHEMATICS OF THE UNIVERSE (WPI), UTIAS, THE UNIVERSITY OF TOKYO, KASHIWA, CHIBA 277-8583, JAPAN
E-mail address: akishi.ikeda@ipmu.jp