

# Three-Slit Interference: A Duality Relation

Mohd Asad Siddiqui and Tabish Qureshi

Centre for Theoretical Physics, Jamia Millia Islamia, New Delhi, India.

The issue of interference and which-way information is addressed in the context of 3-slit interference experiments. A new path distinguishability  $\mathcal{D}_Q$  is introduced, based on Unambiguous Quantum State Discrimination (UQSD). An inequality connecting the interference visibility and path distinguishability,  $\mathcal{V} + \frac{2\mathcal{D}_Q}{3-\mathcal{D}_Q} \leq 1$ , is derived which puts a bound on how much fringe visibility and which-way information can be simultaneously obtained. It is argued that this bound is tight. For 2-slit interference, we derive a new duality relation which reduces to Englert's duality relation and Greenberger-Yasin's duality relation, in different limits.

## 1. Introduction


The two-slit interference experiment with particles has become a cornerstone for studying wave-particle duality. So fundamental is the way in which the two-slit experiment captures the essence of quantum theory, that Feynman ventured to state that it is a phenomenon "which has in it the heart of quantum mechanics; in reality it contains the *only* mystery" of the theory [1]. That radiation and massive particles can exhibit both wave nature and particle nature in different experiment, had become quite clear in the early days of quantum mechanics. However, Niels Bohr emphasized that wave-nature, characterized by two-slit interference, and the particle-nature, characterized by the knowledge of which slit the particle passed through, are mutually exclusive [2]. In doing this he raised this concept to the level of a new fundamental principle.

Much later, this principle was made quantitatively precise by deriving a bound on the extent to which the two natures could be observed simultaneously, by Greenberger and Yasin [3] and later by Englert [4]. Greenberger and Yasin characterised the particle nature by the ability to correctly predict which slit the particle passed through [3]. This predictability was based only on the initial state of the particle, and not on any measurement on it. Englert characterised the particle nature by the ability to distinguish between the two paths of the particle, by an actual measurement [4]. He introduced a quantity  $\mathcal{D}$  for this purpose, which took values between 0 and 1. The wave nature was characterised by the visibility of the interference, given by  $\mathcal{V}$ . The relation putting a bound on the path distinguishability and fringe visibility is given by [4]

$$\mathcal{V}^2 + \mathcal{D}^2 \le 1. \tag{1}$$

Thus one can see that  $\mathcal{D}$  and  $\mathcal{V}$ , which can take values between 0 and 1, are dependent on each other. A full which-way information ( $\mathcal{D} = 1$ ) would surely wash out the interference ( $\mathcal{V} = 0$ ). Eqn. (1) can be thought to be a quantitative statement of Bohr's complementarity

<sup>\*</sup>E-mail: tabish@ctp-jamia.res.in



**Fig. 1** A schematic diagram of the 3-slit interference experiment, with a quantum which-path detector.

principle. It smoothly interpolates between the two extreme scenarios discussed by Bohr, namely, full which-way information and no which-way information.

A dramatic manifestation of Bohr's complementarity principle has been demonstrated in the so-called quantum eraser. Here, "erasing" the which-way information after the particle has passed through the slits, allows one to recover the lost interference fringes [5]. Wave-particle duality has also been connected to various other phenomena. Connection of wave-particle duality with uncertainty relations has been investigated [6–9]. An interesting complementarity between entanglement and the visibility of interference has been demonstrated for two-path interference [10]. That the entanglement between the particle and which-path detector is what affects the visibility of interference, is also the view we take in this investigation.

Various aspects of complementarity were also explored by Jaeger, Shimony and Vaidman [11] where they also explored the case where the particle can follow multiple path, and not just two, before interfering. Bohr's principle of complementarity should surely apply to multi-slit experiments too. However, one might wonder if one can find a quantitatively precise statement of it for multi-slit experiments. Various attempts have been made to formulate a quantitatively precise statement of complementarity, some kind of a duality relation, for the case of multibeam interferometers [12–16]. However, the issue is still not satisfactorily resolved. Beyond the well studied two-slit experiment, the simplest multi-beam case is the 3-slit interference experiment. Englert's duality relation was derived only in the context of 2-slit experiments, and one would like an analogous relation for the case of 3-slit experiments. That is the focus of this paper. Of late there has been a newly generated focus on the three-slit interference experiments [17–19], albeit for a different reason.

## 2. Three-slit interference

Three slit interference is somewhat more involved than its 2-slit counterpart simply due to the fact that while the two-slit interference is the result of interference between two parts coming from the two slits, in the 3-slit interference there are three parts which interfere in different ways. In the general case, if we assume that the separation between slits 1 and 2 is  $\ell_1$  and that between slits 2 and 3 is  $\ell_2$ , there are two interferences from slit 1 and 2 and from slit 2 and slit 3. In addition there is an interference between parts from slit 1 and 3, which involves a slit separation of  $\ell_1 + \ell_2$ . In the case where the two slit separations are the same,  $\ell_1 = \ell_2 = d$ , there are two interferences with slit separation d and one interference with slit separation d. Having more than two slits also allows, in principle, the possibility of having different geometrical arrangement of slits. However, we restrict ourselves to the case of three slits in a linear geometry, as shown in Fig. 1, as that is the geometry in which the experiment is usually done, and that is also the geometry which is used in previous investigations of multi-slit experiments.

We expect additional complicacy in interpreting Bohr's complementarity because if we know that the particle did not go through (say) slit 3, it may not imply complete loss of interference as there is still ambiguity regarding which of the other two slits, 1 or 2, the particle went through.

## 2.1. Which-way information

First we would like to have a way of knowing which of the three slits the particle passed through. Any which-path detector should have three states which should correlate with the particle passing through each slit. Let these states be  $|d_1\rangle, |d_2\rangle, |d_3\rangle$ , which correspond to particle passing through slits 1, 2 and 3, respectively. Without loss of generality we assume that the states  $|d_1\rangle, |d_2\rangle, |d_3\rangle$  are normalized, although they may not necessarily be mutually orthogonal. The combined state of the particle and the which-path detector can be written as

$$|\Psi\rangle = \sqrt{p_1}|\psi_1\rangle|d_1\rangle + \sqrt{p_2}|\psi_2\rangle|d_2\rangle + \sqrt{p_3}|\psi_3\rangle|d_3\rangle,\tag{2}$$

where  $\sqrt{p_1}|\psi_1\rangle$ ,  $\sqrt{p_2}|\psi_2\rangle$ ,  $\sqrt{p_3}|\psi_3\rangle$  are the amplitudes of the particle passing through the slit 1, 2 and 3, respectively. Particle passes through slits 1, 2, and 3 with probabilities  $p_1$ ,  $p_2$  and  $p_3$ , respectively.

If  $|d_1\rangle, |d_2\rangle, |d_3\rangle$  are mutually orthogonal, we can find a Hermitian operator (and thus, a measurable quantity) which will give us different eigenvalues corresponding to  $|d_1\rangle, |d_2\rangle, |d_3\rangle$ , and thus to the particle passing through each of the three slit. In this case, which slit the particle went through, can be known without ambiguity.

If  $|d_1\rangle, |d_2\rangle, |d_3\rangle$  are not mutually orthogonal, the three ways of particle going through the slits will not be fully distinguishable. One needs to define a *distinguishability* of the three different paths of the particle. Defining distinguishability in multi-slit experiments has been a thorny issue [11–15].

#### 2.2. Unambiguous Quantum State Discrimination

As one can see from (2), the problem of distinguishing the three paths of the particle boils down to distinguishing between the three states  $|d_1\rangle, |d_2\rangle, |d_3\rangle$ . In the following we describe a well established method of *unambiquously* discriminating between two non-orthogonal

quantum states, which goes by the name of Unambiguous Quantum State Discrimination (UQSD) [20–24]. If two states  $|p\rangle, |q\rangle$  are not orthogonal, it is impossible to distinguish between the two with certainty. What UQSD does is to separate the measurement results into two categories. First category is the one in which the discrimination fails. The second one distinguishes between the two states without any error. Let the first system, whose states are  $|p\rangle, |q\rangle$  interact with a second system which is initially in a state  $|s_0\rangle$ . The interaction and time evolution leads to the following entangled states

$$U|p\rangle|s_0\rangle = \alpha|p_1\rangle|s_1\rangle + \beta|p_2\rangle|s_2\rangle$$

$$U|q\rangle|s_0\rangle = \gamma|q_1\rangle|s_1\rangle + \delta|q_2\rangle|s_2\rangle, \tag{3}$$

where U is a unitary operator such that  $\langle s_1|s_2\rangle=0$  and  $\langle p_1|q_1\rangle=0$ . If one measures an observable of the second system which has two eigenstates  $|s_1\rangle, |s_2\rangle$  with different eigenvalues, the result  $|s_1\rangle$  lands us into a situation where the non-orthogonal states  $|p\rangle, |q\rangle$  have been replaced by orthogonal  $|p_1\rangle, |q_1\rangle$ . The orthogonal states  $|p_1\rangle, |q_1\rangle$  can be distinguished with hundred percent accuracy, thus distinguishing the original  $|p\rangle, |q\rangle$  without error. However, the other result for system 2,  $|s_2\rangle$ , leads us to states  $|p_2\rangle, |q_2\rangle$  which are not orthogonal, and the discrimination of  $|p\rangle, |q\rangle$  fails. So, either the process fails, or it distinguishes between  $|p\rangle, |q\rangle$  without error. The probability of successfully distinguishing between  $|p\rangle$  and  $|q\rangle$  depends on the constants  $\alpha, \beta, \gamma, \delta$ . It can be easily shown that the maximum probability of successfully distinguishing between  $|p\rangle$  and  $|q\rangle$  is given by [21]

$$P = 1 - |\langle p|q\rangle|. \tag{4}$$

This is called the IDP (Ivanovic-Dieks-Peres) limit, and  $|p\rangle$  and  $|q\rangle$  cannot be distinguished unambiguously with a probability larger than this, even in principle [20–22]. In other words, UQSD is the best bet for discriminating between two non-orthogonal states. This fact has also been experimentally demonstrated recently [25, 26]. Thus, the IDP limit sets a fundamental bound on the distinguishability of two non-orthogonal states.

#### 2.3. Distinguishability

The preceding analysis suggests a natural definition of path-distinguishability. In a two-slit experiment, if the combined state of the particle and the which way detector can be written as  $|\Psi\rangle = |\psi_1\rangle|d_1\rangle + |\psi_2\rangle|d_2\rangle$ , the two paths can be distinguished if the two states  $|d_1\rangle, |d_2\rangle$  can be distinguished. The probability of successfully telling which slit the particle went through is just the probability with which  $|d_1\rangle, |d_2\rangle$  can be unambiguously distinguished. We thus define distinguishability of the two paths in a double-slit experiment as the upper limit of the probability of unambiguously distinguishing between  $|d_1\rangle$  and  $|d_2\rangle$ . Thus we define a new path distinguishability for a two-slit experiment as

$$\mathcal{D}_Q \equiv 1 - |\langle d_1 | d_2 \rangle|. \tag{5}$$

We denote it by  $\mathcal{D}_Q$  to distinguish it from the  $\mathcal{D}$  used in (1).

In the 3-slit experiment, the entangled state (2) implies that, after the particle has passed through the triple slit, the which-path detector could be in state  $|d_1\rangle$  or  $|d_2\rangle$  or  $|d_3\rangle$ , with probabilities  $p_1$ ,  $p_2$  and  $p_3$ , respectively. The probability with which one can tell which of the three slits the particle went through, is just the probability with which one can

distinguish between  $|d_1\rangle, |d_2\rangle, |d_3\rangle$ . So, the problem of distinguishing between the three paths boils down to distinguishing between three non-orthogonal states  $|d_1\rangle, |d_2\rangle, |d_3\rangle$ , which occur with different probabilities.

UQSD has also been generalized to the case of N non-orthogonal states. The probability of unambiguously distinguishing between N non-orthogonal quantum states is bounded by [27, 28]

$$P_N \le 1 - \frac{1}{N-1} \sum_{i \ne j} \sqrt{p_i p_j} |\langle d_i | d_j \rangle|, \tag{6}$$

where  $\{|d_i\rangle\}$  are the N non-orthogonal states, and  $p_i$  are their respective a-priori probabilities. For three non-orthogonal states, the above reduces to

$$P_3 \le 1 - (\sqrt{p_1 p_2} |\langle d_1 | d_2 \rangle| + \sqrt{p_2 p_3} |\langle d_2 | d_3 \rangle| + \sqrt{p_1 p_3} |\langle d_1 | d_3 \rangle|). \tag{7}$$

We define the path-distinguishability for the 3-slit experiment as the upper limit of the probability with which one can distinguish between the three states  $|d_1\rangle, |d_2\rangle, |d_3\rangle$ , which is now given by

$$\mathcal{D}_Q \equiv 1 - (\sqrt{p_1 p_2} |\langle d_1 | d_2 \rangle| + \sqrt{p_2 p_3} |\langle d_2 | d_3 \rangle| + \sqrt{p_1 p_3} |\langle d_1 | d_3 \rangle|), \tag{8}$$

and whose value lies in the range  $0 \le \mathcal{D}_Q \le 1$ . Eqn. (8) should be seen as a generalization of the IDP limit to the case of three non-orthogonal states.

# 2.4. Interference and which-way information

In order to obtain a tight bound on the visibility of interference, given a particular amount of which-way information, we do a wave-packet analysis in the following way. We consider a particle traveling along the z-direction and passing through the triple-slit, with a slit separations  $\ell_1$  and  $\ell_2$ , and then interacting with a which-path detector through a unitary evolution. The evolution of the state is given by the time-dependent Schrödinger equation,

$$i\hbar\partial_t |\Psi(t)\rangle = \mathcal{H}|\Psi(t)\rangle,$$
 (9)

where  $\mathcal{H}$  is the Hamiltonian of the system (the energy operator).

Our strategy is the following. The motion of the particle along the z-axis is redundant, as it only translates the position of the particle from the triple-slit to the screen. What is relevant is the motion and dispersion of the particle along the x-direction. Without going into the details of the which-path detector, we assume that it is a device having 3 states  $|d_1\rangle, |d_2\rangle, |d_3\rangle$ , which get entangled with the three amplitudes of the particle passing through slits 1, 2 and 3, respectively. This entanglement is a necessary condition for the which-path detector to get any information about which slit the particle went through [7].

In the following we assume that the mass of the particle is m, and the three slits are located at  $x = +\ell_1$ , 0,  $-\ell_2$ . The wavefunction emerging from a particular slit is assumed to be a Gaussian wave-packet in the x-direction, of width  $\epsilon$ , localized at  $x = \ell_1$ , 0, or  $-\ell_2$ . A wavefunction of the form  $e^{-\frac{(x-\ell_1)^2}{4\epsilon^2}}$  should represent the particle emerging from a slit of width  $\epsilon$ , located at  $x = \ell_1$ , in a reasonable approximation. As we shall see later, the choice of the Gaussian form does not lead to any loss of generality because the width of the Gaussians does not appear in the final results.

Using this strategy, the combined state of the particle and the which-path detector, when the particle emerges from the triple-slit (time t = 0), will be of the form

$$\Psi(x,0) = A\left(\sqrt{p_1}|d_1\rangle e^{-\frac{(x-\ell_1)^2}{4\epsilon^2}} + \sqrt{p_2}|d_2\rangle e^{-\frac{x^2}{4\epsilon^2}} + \sqrt{p_3}|d_3\rangle e^{-\frac{(x+\ell_2)^2}{4\epsilon^2}}\right),\tag{10}$$

where  $A = (2\pi\epsilon^2)^{-1/4}$ , and  $p_1 + p_2 + p_3 = 1$ . It represents three Gaussian wave-packets localised at the three slits, namely  $x = \ell_1$ , x = 0 and  $x = -\ell_2$ , entangled with the three states of the which-path detector. The width of the three Gaussians is chosen to be the same because the widths of the three slits is assumed to be the same. It is not difficult to see, from the subsequent analysis, that even if we consider the three slit-widths to be different, e.g.,  $\epsilon_1$ ,  $\epsilon_2$ ,  $\epsilon_3$ , they would all drop out from our final result. In general one should consider a factor  $\sqrt{p_1}e^{i\theta}$  instead of  $\sqrt{p_1}$ , and likewise for  $\sqrt{p_2}$  and  $\sqrt{p_3}$ . We do not consider these phase factors simply because they will be absorbed in certain phases introduced later, in section 2.4. We consider the probabilities of all the three paths to be different for generality, but point out that unequal probabilities reduce the visibility of interference. Even in the two-slit interference, unequal beams reduce the visibility of interference.

After a time t, the state of the particle and the detector evolves via (9), with  $\mathcal{H} = \mathbf{p}_x^2/2m$ , to

$$\Psi(x,t) = A\left(\sqrt{p_1}|d_1\rangle e^{-\frac{(x-\ell_1)^2}{4\epsilon^2+2i\hbar t/m}} + \sqrt{p_2}|d_2\rangle e^{-\frac{x^2}{4\epsilon^2+2i\hbar t/m}} + \sqrt{p_3}|d_3\rangle e^{-\frac{(x+\ell_2)^2}{4\epsilon^2+2i\hbar t/m}}\right), (11)$$

where  $A = [\sqrt{2\pi}(\epsilon + i\hbar t/2m\epsilon)]^{-1/2}$ . It represents the three wave-packets spreading and overlapping with each other. The smaller the width  $\epsilon$  of the slits, the stronger is the overlap between the three wave-packets.

The probability (density) of finding the particle at a position x on the screen is given by

$$|\Psi(x,t)|^{2} = |A|^{2} \left( p_{1}e^{-\frac{(x-\ell_{1})^{2}}{2\sigma^{2}}} + p_{2}e^{-\frac{x^{2}}{2\sigma^{2}}} + p_{3}e^{-\frac{(x+\ell_{2})^{2}}{2\sigma^{2}}} \right)$$

$$+ \sqrt{p_{1}p_{2}}e^{-\frac{2x^{2}+\ell_{1}^{2}-2x\ell_{1}}{4\sigma^{2}}} \left[ \langle d_{1}|d_{2}\rangle e^{\frac{ix\ell_{1}\hbar t/m-i\hbar t\ell_{1}^{2}/2m}{4\Omega^{2}}} + \langle d_{2}|d_{1}\rangle e^{-\frac{ix\ell_{1}\hbar t/m-i\hbar t\ell_{1}^{2}/2m}{4\Omega^{2}}} \right]$$

$$+ \sqrt{p_{2}p_{3}}e^{-\frac{2x^{2}+\ell_{2}^{2}+2x\ell_{2}}{4\sigma^{2}}} \left[ \langle d_{2}|d_{3}\rangle e^{\frac{ix\ell_{2}\hbar t/m+i\hbar t\ell_{2}^{2}/2m}{4\Omega^{2}}} + \langle d_{3}|d_{2}\rangle e^{-\frac{ix\ell_{2}\hbar t/m+i\hbar t\ell_{2}^{2}/2m}{4\Omega^{2}}} \right]$$

$$+ \sqrt{p_{1}p_{3}}e^{-\frac{x^{2}+(\ell_{1}^{2}+\ell_{2}^{2})/2+x(\ell_{2}-\ell_{1})}{2\sigma^{2}}} \left[ \langle d_{1}|d_{3}\rangle e^{\frac{i[x(\ell_{1}+\ell_{2})+(\ell_{2}^{2}-\ell_{1}^{2})/2]\hbar t/m}{4\Omega^{2}}} + \langle d_{3}|d_{1}\rangle e^{-\frac{i[x(\ell_{1}+\ell_{2})+(\ell_{2}^{2}-\ell_{1}^{2})/2]\hbar t/m}{4\Omega^{2}}} \right]$$

$$+ \langle d_{3}|d_{1}\rangle e^{-\frac{i[x(\ell_{1}+\ell_{2})+(\ell_{2}^{2}-\ell_{1}^{2})/2]\hbar t/m}{4\Omega^{2}}} \right] , \qquad (12)$$

where  $\sigma^2 = \epsilon^2 + (\hbar t/2m\epsilon)^2$  and  $\Omega^2 = \epsilon^4 + (\hbar t/2m)^2$ . We write the overlaps of various detector states as:  $\langle d_1|d_2\rangle = |\langle d_1|d_2\rangle|e^{i\theta_1}$ ,  $\langle d_2|d_3\rangle = |\langle d_3|d_3\rangle|e^{i\theta_2}$ ,  $\langle d_1|d_3\rangle = |\langle d_1|d_3\rangle|e^{i\theta_3}$ . The phases  $\theta_1, \theta_2, \theta_3$  being arbitrary, any additional phases coming from the amplitudes of the initial

state (10) would be absorbed in them. The probability density then reduces to

$$|\Psi(x,t)|^{2} = |A|^{2} \left( e^{-\frac{x^{2}}{2\sigma^{2}}} \left( p_{1}e^{-\frac{\ell_{1}^{2}-2x\ell_{1}}{2\sigma^{2}}} + p_{2} + p_{3}e^{-\frac{\ell_{2}^{2}+2x\ell_{2}}{2\sigma^{2}}} \right) + 2\sqrt{p_{1}p_{2}} |\langle d_{1}|d_{2}\rangle| e^{-\frac{2x^{2}+\ell_{1}^{2}-2x\ell_{1}}{4\sigma^{2}}} \cos\left(\frac{x\ell_{1}\hbar t}{4m\Omega^{2}} - \beta_{1} + \theta_{1}\right) + 2\sqrt{p_{2}p_{3}} |\langle d_{2}|d_{3}\rangle| e^{-\frac{2x^{2}+\ell_{2}^{2}+2x\ell_{2}}{4\sigma^{2}}} \cos\left(\frac{x\ell_{2}\hbar t}{4m\Omega^{2}} + \beta_{2} + \theta_{2}\right) + 2\sqrt{p_{1}p_{3}} |\langle d_{1}|d_{3}\rangle| e^{-\frac{2x^{2}+\ell_{1}^{2}+\ell_{2}^{2}+2x(\ell_{2}-\ell_{1})}{4\sigma^{2}}} \cos\left(\frac{x(\ell_{1}+\ell_{2})\hbar t}{4m\Omega^{2}} + \beta_{3} + \theta_{3}\right)\right), (13)$$

where  $\beta_1 = \frac{\hbar t \ell_1^2}{8m\Omega^2}$ ,  $\beta_2 = \frac{\hbar t \ell_2^2}{8m\Omega^2}$ , and  $\beta_3 = \frac{\hbar t (\ell_2^2 - \ell_1^2)}{8m\Omega^2}$ .

Visibility of the interference fringes is conventionally defined as [29]

$$\mathcal{V} = \frac{I_{max} - I_{min}}{I_{max} + I_{min}},\tag{14}$$

where  $I_{max}$  and  $I_{min}$  represent the maximum and minimum intensity in neighbouring fringes, respectively. Since all the factors multiplying the cosine terms in (13) are non-negative, the maxima of the fringe pattern will occur where the value of all the cosines is +1. The minima will occur where value of all the cosines is -1/2 at the same time.

# 2.5. Equally spaced slits

In the usual three-slit interference experiments, the three slits are equally spaced. In that case  $\ell_1 = \ell_2 \equiv d$ , and  $\beta_1 = \beta_2 \equiv \beta$  and  $\beta_3 = 0$ . Provided we ignore  $\beta$ , the cosines can indeed have values all +1 or all -1/2 for certain values of x. Ignoring  $\beta$  amounts to ignoring d/2 in comparison to x, which is justified if one looks at any fringe except the central one (at x = 0), since fringe width on the screen is much larger than the slit separation d. The *ideal* visibility can then be written down as

$$\mathcal{V}_{I} = \frac{3\left(\sqrt{p_{1}p_{2}}|\langle d_{1}|d_{2}\rangle|e^{\frac{xd}{2\sigma^{2}}} + \sqrt{p_{1}p_{3}}|\langle d_{1}|d_{3}\rangle|e^{-\frac{d^{2}}{2\sigma^{2}}} + \sqrt{p_{2}p_{3}}|\langle d_{2}|d_{3}\rangle|e^{-\frac{xd}{2\sigma^{2}}}\right)}{\alpha + \sqrt{p_{1}p_{2}}|\langle d_{1}|d_{2}\rangle|e^{\frac{xd}{2\sigma^{2}}} + \sqrt{p_{1}p_{3}}|\langle d_{1}|d_{3}\rangle|e^{-\frac{d^{2}}{2\sigma^{2}}} + \sqrt{p_{2}p_{3}}|\langle d_{2}|d_{3}\rangle|e^{-\frac{xd}{2\sigma^{2}}}}.$$
(15)

where  $\alpha = 2\left(p_2 + \left[p_1e^{\frac{xd}{\sigma^2}} + p_3e^{\frac{-xd}{\sigma^2}}\right]e^{-\frac{d^2}{2\sigma^2}}\right)$ . In reality, fringe visibility will be reduced due many factors, including the width of the slits. For example, if the width of the slits is very large, the fringes may not be visible at all. It will also get reduced by the varying phases  $\theta_1, \theta_2, \theta_3$ . In an ideal situation, the maximum visibility one can theoretically get will be in the case when  $d \ll \sigma$ , which amounts to assuming that the spread of a particular wave-packet, when it reaches the screen, is so large that the separation between two neighboring slits d is negligible in comparison, and also when all the phases are either zero or their effect cancels out. Actual fringe visibility will be less than or equal to that, and can be written as

$$\mathcal{V} \le \frac{3\left(\sqrt{p_1 p_2} |\langle d_1 | d_2 \rangle| + \sqrt{p_1 p_3} |\langle d_1 | d_3 \rangle| + \sqrt{p_2 p_3} |\langle d_2 | d_3 \rangle|\right)}{2 + \sqrt{p_1 p_2} |\langle d_1 | d_2 \rangle| + \sqrt{p_1 p_3} |\langle d_1 | d_3 \rangle| + \sqrt{p_2 p_3} |\langle d_2 | d_3 \rangle|}.$$
(16)

Using (8) the above equation gives

$$\mathcal{V} + \frac{2\mathcal{D}_Q}{3 - \mathcal{D}_Q} \le 1. \tag{17}$$

Eqn. (17) is a new duality relation which puts a bound on how much which-way information we can obtain and how much fringe visibility we can get at the same time. It is straightforward to check that  $\mathcal{V}=1$  is possible only for  $\mathcal{D}_Q=0$ , and  $\mathcal{D}_Q=1$  implies  $\mathcal{V}=0$ . Note that (17) can also be expressed in another form

$$\mathcal{D}_Q + \frac{2\mathcal{V}}{3 - \mathcal{V}} \le 1. \tag{18}$$

## 2.6. Unequally spaced slits

When the three slits are equally spaced, maximum is achieved when the two  $\cos\left(\frac{xd\hbar t}{4m\Omega^2}\right)$  terms and the  $\cos\left(\frac{2xd\hbar t}{4m\Omega^2}\right)$  term have values +1 at the same time, which happens when  $\frac{xd\hbar t}{4m\Omega^2} = 2n\pi$ , provided that  $\theta_1, \theta_2, \theta_3$  are zero. When the three slits are unequal, the three terms  $\cos\left(\frac{x\ell_1\hbar t}{4m\Omega^2}\right)$ ,  $\cos\left(\frac{x\ell_2\hbar t}{4m\Omega^2}\right)$  and  $\cos\left(\frac{x(\ell_1+\ell_2)\hbar t}{4m\Omega^2}\right)$  cannot all have values +1 at the same time. Thus the maximum intensity will be smaller than that in the equally spaced case, or  $I_{max}^{unequal} < I_{max}^{equal}$ .

Minimum intensity in the equally spaced case is attained when all the cosine terms are equal to -1/2, which happens when  $\frac{xd\hbar t}{4m\Omega^2} = \frac{2n\pi}{3}$ , provided that  $\theta_1, \theta_2, \theta_3$  are zero. When the three slits are unequally spaced, the three terms  $\cos\left(\frac{x\ell_1\hbar t}{4m\Omega^2}\right)$ ,  $\cos\left(\frac{x\ell_2\hbar t}{4m\Omega^2}\right)$  and  $\cos\left(\frac{x(\ell_1+\ell_2)\hbar t}{4m\Omega^2}\right)$  cannot all have values -1/2 at the same time. Thus the minimum intensity will be larger than that in the equally spaced case, or  $I_{min}^{unequal} > I_{min}^{equal}$ . These two observations lead to the straightforward conclusion that fringe visibility in the case of unequal slits will be strictly smaller than that in the case of equal slits, other things being the same,

$$\mathcal{V}^{unequal} < \mathcal{V}^{equal}. \tag{19}$$

Using the above in conjunction with (16) we can write for the fringe visibility, for the case of unequal slits

$$\mathcal{V} < \frac{3\left(\sqrt{p_1 p_2} |\langle d_1 | d_2 \rangle| + \sqrt{p_1 p_3} |\langle d_1 | d_3 \rangle| + \sqrt{p_2 p_3} |\langle d_2 | d_3 \rangle|\right)}{2 + \sqrt{p_1 p_2} |\langle d_1 | d_2 \rangle| + \sqrt{p_1 p_3} |\langle d_1 | d_3 \rangle| + \sqrt{p_2 p_3} |\langle d_2 | d_3 \rangle|}.$$
 (20)

Using (8) the above equation gives

$$\mathcal{V} + \frac{2\mathcal{D}_Q}{3 - \mathcal{D}_Q} < 1. \tag{21}$$

Note the strictly less than sign in the above, in contrast with (17). The above relation implies that if the slits are unequally spaced, even if the distinguishability  $\mathcal{D}_Q$  is reduced to zero, the visibility of interference can never be 1. The reason of this behaviour lies in the fact that a certain amount of loss of interference visibility is rooted not in the path distinguishability, but in the unequal spacing of the three slits.

# 2.7. Specific cases

Let us look at some special cases arising from the fact that there are not two, but three slit. Suppose we have a which-path detector which can detect with certainty if the particle has passed through slit 1 or not. If the particle has not passed through slit 1, the detector is unable to say which of the other two slits, 2 or 3, has the particle taken. Such a scenario can occur, for example, if we have a tiny camera in front of slit 1 which, for each particle, can say if the particle has gone through slit 1 or not. In this case  $|d_1\rangle$  is orthogonal to both  $|d_2\rangle$  and

 $|d_3\rangle$ , and  $|d_2\rangle$ ,  $|d_3\rangle$  are parallel. Thus, in this case,  $|\langle d_2|d_3\rangle|=1$  and  $|\langle d_1|d_2\rangle|=|\langle d_1|d_3\rangle|=0$ . Assuming, for simplicity,  $p_1=p_2=p_3=\frac{1}{3}$ , the distinguishability  $\mathcal{D}_Q$ , in this case is

$$\mathcal{D}_Q \equiv 1 - \frac{1}{3} (|\langle d_1 | d_2 \rangle| + |\langle d_2 | d_3 \rangle| + |\langle d_1 | d_3 \rangle|) = \frac{2}{3}, \tag{22}$$

Consequently, the fringe visibility is limited by

$$V \le \frac{3}{7}.\tag{23}$$

Physically what is happening is the following. Particle going through slits 2 and 3 gives rise to a sharp interference pattern, however, particle going through slit 1 gives rise to a uniform background particle count, thus reducing the overall visibility of the fringes arising from slits 2 and 3.

Let us consider another case where  $|d_1\rangle$  and  $|d_2\rangle$  are orthogonal to each other, but both have equal overlap with  $|d_3\rangle$ . Such a case can be exemplified by  $|d_1\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle)$ ,  $|d_2\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle)$  and  $|d_3\rangle = |\uparrow\rangle$ , where  $|\uparrow\rangle$ ,  $|\downarrow\rangle$  form an orthonormal set. In this case (again, for simplicity, assuming  $p_1 = p_2 = p_3 = \frac{1}{3}$ ) the distinguishability is

$$\mathcal{D}_Q = 1 - \frac{1}{3}(|\langle d_1 | d_2 \rangle| + |\langle d_2 | d_3 \rangle| + |\langle d_1 | d_3 \rangle|) = 1 - \frac{\sqrt{2}}{3},\tag{24}$$

Consequently, the fringe visibility is limited by

$$\mathcal{V} \le \frac{3\sqrt{2}}{6+\sqrt{2}}.\tag{25}$$

## 2.8. The two-slit experiment

Just for completeness, here we wish to derive a duality relation for a two-slit experiment in the case where one defines path distinguishability based on UQSD. We define the distinguishability of two paths as the upper limit of (6) for N=2. The path distinguishability then reads

$$\mathcal{D}_O \equiv 1 - 2\sqrt{p_1 p_2} |\langle d_1 | d_2 \rangle|, \tag{26}$$

where  $p_1$  and  $p_2$  are the probabilities of the particle to go through the first and the second slit, respectively.

By carrying out an analysis similar to the one earlier in this section (essentially, putting  $p_3 = 0$  in (13) and redefining some constants), one can show that the distinguishability and fringe visibility, in the two-slit experiment, are bounded by

$$\mathcal{V} + \mathcal{D}_O \le 1. \tag{27}$$

The above relation is a simple wave-particle duality relation for a two-slit interference experiment where the beams may be unequal.

It can be connected to Englert's duality relation (1) for the case  $p_1 = p_2 = 1/2$ . In Englert's analysis, distinguishability is given by  $\mathcal{D} \equiv \sqrt{1 - |\langle d_1 | d_2 \rangle|^2}$  [4], which can be related to  $\mathcal{D}_Q$  in (26), for  $p_1 = p_2 = 1/2$ , by the relation

$$\mathcal{D}_Q = 1 - \sqrt{1 - \mathcal{D}^2}.\tag{28}$$

If one plugs in the above form of  $\mathcal{D}_Q$  in (27), the latter reduces to  $\mathcal{V}^2 + \mathcal{D}^2 \leq 1$ , which is just Englert's duality relation (1). The new relation (27) appears to be more versatile for two-slit

experiments, because it also applies to certain modified two-slit experiments in which the which-path detector is replaced by a "quantum device" [30].

Lastly we discuss a particular scenario in which the two states of the which-path detector are identical, namely  $|\langle d_1|d_2\rangle|=1$ . In such a situation, experimentally one cannot tell which slit the particle went through. However, if the probabilities of the particle for going through the two slits are known to be different, one can *predict* which slit the particle is most likely to have gone through. In this situation our  $\mathcal{D}_Q$ , given by (26), is reduced to

$$\overline{\mathcal{D}}_Q \equiv 1 - 2\sqrt{p_1 p_2}.\tag{29}$$

Interestingly the above reduced distinguishability is related to the *predictability*, defined by Greenberger and Yasin [3] as  $\mathcal{P} \equiv |p_1 - p_2|$ , by the following relation

$$\overline{\mathcal{D}}_Q = 1 - \sqrt{1 - \mathcal{P}^2}. (30)$$

So, for the case  $|\langle d_1|d_2\rangle|=1$ , our new duality relation (27) reduces to

$$\mathcal{P}^2 + \mathcal{V}^2 \le 1,\tag{31}$$

which is precisely the duality relation derived by Greenberger and Yasin [3].

So, the versatility of the new two-slit duality relation can be seen from the fact that for  $p_1 = p_2$  it reduces to Englert's duality relation dealing with *distinguishability*, and for  $|\langle d_1|d_2\rangle| = 1$ , it reduces to Greenberger and Yasin's duality relation dealing with *predictability*.

#### 3. Conclusion

In the analysis carried out in this paper, we have introduced a new path distinguishability  $\mathcal{D}_Q$ , based on UQSD, which is just the upper limit of the probability with which one can unambiquously distinguish between the quantum states of the which-path detector correlated with the paths of the particle. Consequently, it is the maximum probability with which one can unambiquously tell which slit the particle went through. We carried out a wavepacket evolution of a particle through a triple-slit. Calculating the fringe-visibility after a Schrödinger evolution, we relate it to the path distinguishability and derive a new duality relation  $V + \frac{2\mathcal{D}_Q}{3-\mathcal{D}_Q} \leq 1$ . The analysis is restricted to three slits of equal widths, in a linear geometry, as shown in Fig. 1. Starting from the triple-slit, the time evolution, leading to the probability density of the particle on the screen, is exact. Various approximations, in the subsequent analysis, are made only to obtain the maximum possible visibility of interference, given a particular  $\mathcal{D}_Q$ . Because of the way in which the analysis is carried out, this should be the tightest possible bound on distinguishability and fringe visibility for the 3-slit experiment. For two-slit interference, we derive a new duality relation which reduces to Englert's duality relation and Greenberger and Yasin's duality relation, in different limits. Lastly, we feel that (6) suggests a straightforward definition of distinguishability for N-slit interference experiments:

$$\mathcal{D}_Q = 1 - \frac{1}{N-1} \sum_{i \neq j} \sqrt{p_i p_j} |\langle d_i | d_j \rangle|, \tag{32}$$

where  $|d_i\rangle$  is the state of the path-detector correlated with the i'th of the N possible paths.

## Acknowledgement

Mohd Asad Siddiqui thanks the University Grants Commission, India for financial support.

#### References

- R.P. Feynman, R.B. Leighton, M. Sands, The Feynman Lectures on Physics (Addison-Wesley 1966) Vol. 3, pp. 1-1.
- [2] N. Bohr, "The quantum postulate and the recent development of atomic theory," Nature (London) 121, 580-591 (1928).
- [3] D.M. Greenberger, A. Yasin, "Simultaneous wave and particle knowledge in a neutron interferometer", Phys. Lett. A 128, 391 (1988).
- [4] B-G. Englert, "Fringe visibility and which-way information: an inequality", Phys. Rev. Lett. 77, 2154 (1996).
- [5] S.P. Walborn, M.O. Terra Cunha, S.Pádua, C.H. Monken, "Quantum erasure", American Scientist 91, 336-343 (2003).
- [6] S. Dürr, G. Rempe, "Can wave-particle duality be based on the uncertainty relation?," Am. J. Phys. 68, 1021-1024 (2000).
- [7] T. Qureshi, R. Vathsan, "Einstein's recoiling slit experiment, complementarity and uncertainty", Quanta 2, 58-65 (2013). DOI: 10.12743/quanta.v2i1.11
- [8] P.J. Coles, J. Kaniewski, S. Wehner, "Equivalence of wave-particle duality to entropic uncertainty," Nature Communications 5, 5814 (2014).
- [9] S. Tanimura, "Complementarity and the nature of uncertainty relations in Einstein-Bohr recoiling slit experiment," Quanta 4, 1-9 (2015). DOI: 10.12743/quanta.v4i1.35
- [10] A. Hosoya, A. Carlini, S. Okano, "Complementarity of entanglement and interference," Int. J. Mod. Phys. C 17, 493 (2006). DOI: 10.1142/S0129183106008716
- [11] G. Jaeger, A. Shimony, L. Vaidman, "Two interferometric complemenarities," Phys. Rev. A 51, 54 (1995).
- [12] S. Dürr, "Quantitative wave-particle duality in multibeam interferometers," Phys. Rev. A 64, 042113 (2001).
- [13] G. Bimonte, R. Musto, "Comment on 'Quantitative wave-particle duality in multibeam interferometers'," Phys. Rev. A 67, 066101 (2003).
- [14] G. Bimonte, R. Musto, "On interferometric duality in multibeam experiments" J. Phys. A: Math. Gen. 36, 11481 (2003). (2003).
- [15] B-G. Englert, "Wave-particle duality in multi-path interferometers: General concepts and three-path interferometers," Int. J. Quantum Inform. 6, 129 (2008).
- [16] M. Zawisky, M. Baron, R. Loidl, "Three-beam interference and which-way information in neutron interferometry," *Phys. Rev. A* **66**, 063608 (2002).
- [17] U. Sinha, C. Couteau, T. Jennewein, R. Laflamme, G. Weihs, "Ruling Out Multi-Order Interference in Quantum Mechanics", Science 329, 418-421 (2010).
- [18] H.D. Raedt, K. Michielsen, K. Hess, "Analysis of multipath interference in three-slit experiments", Phys. Rev. A 85, 012101 (2012).
- [19] R. Sawant, J. Samuel, A. Sinha, S. Sinha, U. Sinha, "Nonclassical paths in quantum interference experiments," Phys. Rev. Lett. 113, 120406 (2014).
- [20] I.D. Ivanovic, "How to differentiate between non-orthogonal states", Phys. Lett. A 123, 257 (1987).
- [21] D. Dieks, "Overlap and distinguishability of quantum states," Phys. Lett. A 126, 303 (1988).
- [22] A. Peres, "How to differentiate between non-orthogonal states," Phys. Lett. A 128, 19 (1988).
- [23] G. Jaeger, A. Shimony, "Optimal distinction between two non-orthogonal quantum states," *Phys. Lett.* A 197, 83 (1995).
- [24] J.A. Bergou, U. Herzog, M. Hillery, "Discrimination of quantum states," Lect. Notes Phys. 649, 417-465 (2004).
- [25] G. Waldherr, A.C. Dada, P. Neumann, F. Jelezko, E. Andersson, J. Wrachtrup, "Distinguishing between nonorthogonal quantum states of a single nuclear spin," Phys. Rev. Lett. 109, 180501 (2012).
- [26] M. Agnew, E. Bolduc, K.J. Resch, S. Franke-Arnold, J. Leach, "Discriminating single-photon states unambiguously in high dimensions," Phys. Rev. Lett. 113, 020501 (2014).
- [27] S. Zhang Y. Feng, X. Sun, M. Ying, "Upper bound for the success probability of unambiguous discrimination among quantum states," Phys. Rev. A 64, 062103 (2001).
- [28] D. Qiu, "Upper bound on the success probability for unambiguous discrimination," Phys. Lett. A 303, 140-146 (2002)
- [29] M. Born, E. Wolf, "Principles of Optics" (Cambridge University Press, UK, 2002), 7th edition.
- [30] T. Qureshi, "Quantum twist to complementarity: A duality relation", Prog. Theor. Exp. Phys. 2013,

041A01 (2013).