SEMISIMPLE QUANTUM COHOMOLOGY OF SOME FANO VARIETIES

NICOLAS PERRIN

ABSTRACT. We give sufficient conditions for the semisimplicity of quantum cohomology of Fano varieties of Picard rank 1. We apply these techniques to prove new semisimplicity results for some Fano varieties of Picard rank 1 and large index. We also give examples of Fano varieties having a non semisimple small quantum cohomology but a semisimple big quantum cohomology.

INTRODUCTION

Since Dubrovin's conjecture [20], the question whether the big quantum cohomology ring of a variety is semisimple is important and has been discussed in many articles [29, 1, 22, 23, 18, 14]. In particular necessary conditions for semisimplicity are given by Hertling, Manin and Teleman in [22]. For a few varieties X, for example some Fano threefolds [18], toric varieties [23] or some homogeneous spaces [14], it was proved that the small quantum cohomology ring QH(X) (see Subsection 1.4) is semisimple. However for some homogeneous spaces, it was proved in [14] and [16] that the small quantum cohomology does not need to be semisimple.

In this paper we give sufficient conditions for the small quantum cohomology ring QH(X) of a smooth complex Fano variety X to be semisimple. For X such a Fano variety and \star_0 the product in QH(X) (see Subsection 1.4), we define

 $Q_X(a,b) =$ sum of the coefficients of q^k for some k in the product $a \star_0 b$.

This is a quadratic form defined on QH(X) and if R(X) is the radical of QH(X) we prove (see Theorem 2.3.1)

Theorem 1. Let X be Fano with Picard number 1 and Q_X positive definite. Let h the class of a generator of the Picard group.

1. Then $R(X) \subset \{a \in QH(X) \mid h^k \star_0 a = 0 \text{ for some } k\}.$

2. If h is invertible in QH(X), then QH(X) is semisimple.

In the second part of this paper we give examples of varieties whose quadratic form Q_X is positive definite. In particular we obtain (See Theorem 3.4.1 and Proposition 3.5.1)

Theorem 2. Let X be a variety in the following list (See Subsection 3.3)

\mathbb{P}^n	Q_n	OG(5, 10)	E_{6}/P_{6}
$\operatorname{Gr}(2,n)$	LG(3,6)	OG(6, 12)	E_{7}/P_{7}

or an adjoint variety (See Subsection 3.5) and let Y be a general linear section of codimension k of X with $2c_1(Y) > \dim Y$. Then Q_Y is positive definite.

Date: 25.04.2014.

²⁰⁰⁰ Mathematics Subject Classification. 14N35, 14N15, 53D45.

In particular alsmost all Fano varieties of coindex 3 occur in the above list (see Example 3.4.3). In the third part we give a closer look at the product $h \star_0$ – with the generator of the Picard group and obtain the following semisimplicity result (See Theorem 4.2.1).

Theorem 3. Let Y be a general hyperplane section with $2c_1(Y) > \dim Y$ of a homogeneous space in the following list

\mathbb{P}^n	Q_n	LG(3,6)	F_{4}/P_{1}
Gr(2, 2n+1)	OG(5, 10)	OG(2, 2n+1)	$G_2/P_1.$

Then QH(Y) is semisimple.

In particular, this Theorem recovers in a uniform way semisimplicity results proved [16] and [34] and provides new semisimplicity results.

In the last section we consider two cases where the small quantum cohomology is not semisimple and prove using Theorem 1 that the big quantum cohomology ring (denoted BQH(X), see Section 1) is semisimple (see Theorem 5.2.2 and Theorem 5.3.2).

Theorem 4. Let X = IG(2, 2n) or $X = F_4/P_4$. Then QH(X) is not semisimple but BQH(X) is semisimple.

This result was the starting point of this work which came from discussions with A. Mellit and M. Smirnov. They obtain in [21] together with S. Galkin an independent proof of the semisimplicity of BQH(Y) for Y = IG(2, 6).

Let us say few words on Dubrovin's conjecture. Recall that the first part of this conjecture states that for X smooth projective, the semisimplicity of BQH(X) is equivalent to the existence of a full exceptional collection in $D^b(X)$, where $D^b(X)$ denotes the bounded derived category of coherent sheaves on X.

For all homogeneous spaces X appearing in Theorem 3 and Theorem 4 except F_4/P_1 and F_4/P_4 , it is known (see [25, Section 6] and [26]) that their derived category admits a full exceptional collection proving Dubrovin's conjecture in these cases.

The same is true for a hyperplane section Y = IG(2, 2n+1) of X = Gr(2, 2n+1) recovering results of [34].

Furthermore, for X = Gr(2, 5), X = OG(5, 10) or X = LG(3, 6) the hyperplane sections Y of X with $2c_1(Y) > \dim Y$ also have an exceptional collection (see [25, Section 6]) proving Dubrovin's conjecture in these cases.

Acknowledgement. I thank A. Mellit and M. Smirnov for enlightening discussions and email exchanges. I also thank the organisers of the conference *Quantum* cohomology and quantum K-theory held in Paris in January 2014 where this work started. Finally I thank P.-E. Chaput for the program [11] which was of great use in many computations.

CONTENTS

Int	roduction	1
1.	Big quantum cohomology	3
2.	localisation of the radical	5
3.	Varieties with Q_X positive definite	7
4.	Semisimplicity of the quantum cohomology	14

 $\mathbf{2}$

1. BIG QUANTUM COHOMOLOGY

In this section we recall few facts and fix notation for the quantum cohomology of a complex smooth projective variety X. We write H(X) for $H^*(X, \mathbb{R})$.

1.1. **Reminders.** Let X be a smooth projective variety, let Eff(X) be the cone of effective curves and let $\beta \in \text{Eff}(X)$. Denote by $\overline{M}_{0,n}(X,\beta)$ the Kontsevich moduli space of genus 0 stable maps to X of degree β with n marked points. This is a proper scheme and there are evaluation morphisms $\text{ev}_i : \overline{M}_{0,n}(X,\beta) \to X$ defined by evaluating the map at the n-th marked points. For $(\gamma_i)_{i \in [1,n]}$ cohomology classes on X, one defines the Gromov-Witten invariants as follows:

$$I_{0,n,\beta}(\gamma_1,\cdots,\gamma_n) = \int_{[\overline{\mathrm{M}}_{0,n}(X,\beta)]^{\mathrm{vir}}} \mathrm{ev}_1^* \, \gamma_1 \cup \cdots \cup \mathrm{ev}_n^* \, \gamma_n$$

where $[\overline{\mathrm{M}}_{0,n}(X,\beta)]^{\mathrm{vir}}$ is the virtual fundamental class as defined in [4]. When $\gamma_i = \gamma$ for all $i \in [1,n]$, we write $I_{0,n,\beta}(\gamma_1,\cdots,\gamma_n) = I_{0,n,\beta}(\gamma^n)$.

Let $N + 1 = \operatorname{rk}(\operatorname{H}(X))$ and $r = \operatorname{rk}(\operatorname{Pic}(X))$. Let $(e_i)_{i \in [0,N]}$ be a basis of $\operatorname{H}(X)$ such that $e_0 = 1$ is the fundamental class and (e_1, \dots, e_r) is a basis of $H^2(X, \mathbb{R})$. For $\gamma \in \operatorname{H}(X)$, write $\gamma = \sum_i x_i e_i$ and define the **Gromov-Witten potential** by

$$\Phi(\gamma) = \sum_{n \ge 0} \sum_{\beta \in \text{Eff}(X)} \frac{1}{n!} I_{0,n,\beta}(\gamma^n).$$

This is an element in $R = \mathbb{R}[[(x_i)_{i \in [0,N]}]]$. If (,) denotes the Poincaré pairing, then the **quantum product** \star is defined as follows

$$(e_i \star e_j, e_k) = \frac{\partial^3 \Phi}{\partial x_i \partial x_j \partial x_k}(\gamma)$$

and extended by bilinearity to any other classes. This actually defines a family of products parametrised by H(X). The main result of the theory states that these products are associative (see [3, 5, 24]). We write BQH(X) for the algebra $(H(X) \otimes_{\mathbb{R}} R, \star)$.

1.2. Virtual fundamental class. For our computations of Gromov-Witten invariants we shall use the following general result on the virtual fundamental class for smooth projective varieties (this was proved in [39] according to [2, Point (1.4)], we refer to [28, Proposition 2] for an algebraic proof).

Proposition 1.2.1. Let X be a smooth projective complex algebraic variety such that $\overline{M}_{g,n}(X,\beta)$ has the expected dimension, then the virtual class is the fundamental class.

1.3. Divisor axiom. One very useful property of Gromov-Witten invariants is that for degree 2 cohomology classes, they can be easily computed. Indeed we have the following divisor axiom. For $\gamma_1 \in H^2(X, \mathbb{R})$, we have

$$I_{0,n,\beta}(\gamma_1,\cdots,\gamma_n) = \left(\int_{\beta} \gamma_1\right) I_{0,n-1,\beta}(\gamma_2,\cdots,\gamma_n).$$

This gives a simplification of the potential (modulo terms of degree lower than 2):

$$\Phi(\gamma) = \gamma \cup \gamma \cup \gamma + \sum_{n \ge 0} \sum_{\beta \in \text{Eff}(X)} \frac{1}{n!} I_{0,n,\beta}(\bar{\gamma}^n) q^{\beta}$$

where $\bar{\gamma}$ is the projection of γ on the span of $(e_i)_{i \in [r+1,N]}$ and $q^{\beta} = q_1^{d_1} \cdots q_r^{d_r}$ with $d_i = \int_{\beta} e_i$ and $q_i = e^{x_i}$ for $i \in [1, r]$. Writing $\bar{\gamma} = \sum_{i=r+1}^N x_i e_i$ we get

$$\Phi(\gamma) = \gamma \cup \gamma \cup \gamma + \sum_{n \ge 0} \sum_{\beta \in \text{Eff}(X)} \sum_{n_{r+1} + \dots + n_N = n} \frac{x_{r+1}^{n_{r+1}} \cdots x_N^{n_N}}{n_{r+1}! \cdots n_N!} I_{0,n,\beta}(e_{r+1}^{n_{r+1}}, \cdots, e_N^{n_N}) q^{\beta}$$

1.4. Small quantum product. A very classical special product called the small quantum product and denoted by \star_0 in this paper is obtained as follows

$$(e_i \star_0 e_j, e_k) = \frac{\partial^3 \Phi}{\partial x_i \partial x_j \partial x_k}(\gamma)|_{\bar{\gamma}=0}$$

This is also a family of associative products parametrised by $H^2(X, \mathbb{R})$. This product is easier to compute and only involves 3-points Gromov-Witten invariants *i.e.* Gromov-Witten invariants with n = 3. Set $R_0 = \mathbb{R}[[(q_i)_{i \in [1,r]}]]$. We write QH(X)for the algebra $(H(X) \otimes_{\mathbb{R}} R_0, \star_0)$. Note that $R_0 = R/((x_i)_{i \in [r+1,N]})$.

1.5. Deformation in the τ -direction. Let $\tau \in H(X)$ and choose the basis $(e_i)_{i \in [0,N]}$ so that $e_{r+1} = \tau$. For $\gamma \in H(X)$, write $\hat{\gamma}$ for its projection in the span of $(e_i)_{i \in [r+2,N]}$. We define the product \star_{τ} as follows:

$$(e_i \star_\tau e_j, e_k) = \frac{\partial^3 \Phi}{\partial x_i \partial x_j \partial x_k}(\gamma)|_{\hat{\gamma}=0}$$

This is also a family of associative products parametrised by $H^2(X, \mathbb{R}) \oplus \mathbb{R}\tau$. Set $R_{\tau} = \mathbb{R}[[(q_i)_{i \in [1,r]}, x_{r+1}]]$. We write $\operatorname{BQH}_{\tau}(X)$ for the algebra $(\operatorname{H}(X) \otimes_{\mathbb{R}} R_{\tau}, \star_{\tau})$. Note that $R_{\tau} = R/((x_i)_{i \in [r+2,N]})$. Let us describe a general product in this algebra (we set $t = x_{r+1}$):

$$(e_i \star_{\tau} e_j, e_k) = (e_i \star_0 e_j, e_k) + t \sum_{\beta \in \text{Eff}(X)} I_{0,4,\beta}(e_i, e_j, e_k, e_{r+1}) q^{\beta} + O(t^2).$$

In particular when e_i is the class of a divisor this product takes a simple form. Denote by Ψ_i the endomorphism of $\mathbb{R}[[(q_i)_{i \in [1,r]}]]$ defined by

$$\Psi_i\left(\sum_{\beta\in \mathrm{Eff}(X)} z_\beta q^\beta\right) = \sum_{\beta\in \mathrm{Eff}(X)} d_i z_\beta q^\beta$$

with $d_i = \int_{\beta} e_i$ and extend Ψ_i by linearity on QH(X) via its actions on the scalars. We get the formula $e_i \star_{\tau} e_j = e_i \star_0 e_j + t \Psi_i(e_{r+1} \star_0 e_k) + O(t^2)$.

2. LOCALISATION OF THE RADICAL

In this section, we prove that the existence of positive definite hermitian or real forms imply semisimplicity or regularity results on finite dimensional commutative algebras. Let A be a finite dimensional commutative \mathbb{C} -algebra with 1. We write R(A) for the radical of A. For $a \in A$, we write $E_a \in \text{End}_A(A)$ for the endomorphism obtained by multiplication with a.

2.1. Semisimplicity and inner product. We first relate the semisimplicity of A with the existence of an inner product.

Proposition 2.1.1. The algebra A is semisimple if and only if there exists an algebra involution $a \mapsto \overline{a}$ and a linear form $\varphi : A \to \mathbb{C}$ with $\varphi(1) = 1$ such that the bilinear form defined by $Q(a, b) = \varphi(a\overline{b})$ is an inner product.

Proof. Let $n = \dim A$. Assume that A is semisimple, then for $a \in A$, the endomorphism $E_a \in \operatorname{End}_A(A)$ is semisimple. Since A is commutative, the endomorphisms $(E_a)_{a \in A}$ are simultaneously diagonalisable in a basis $(e_i)_{i \in [1,n]}$. These elements are orthogonal idempotents whose sum is 1. Define the linear form φ by $\varphi(a) = n^{-1}\operatorname{Tr}(E_a)$ and the involution $a \mapsto \overline{a}$ as the unique antilinear map with $\overline{e}_i = e_i$. This defines the desired inner product.

Conversely, assume that such an algebra involution and linear form exist. Then the endomorphisms E_a are normal for Q: we have $Q(E_a(b), c) = Q(ab, c) = \varphi(ab\bar{c}) = Q(b, \bar{a}c) = Q(b, E_{\bar{a}}(c))$. The adjoint of E_a is $E_{\bar{a}}$ and they commute. In particular the endomorphisms $(E_a)_{a \in A}$ are simultaneously diagonalisable in a basis $(e_i)_{i \in [1,n]}$ and these elements are orthogonal idempotents whose sum is 1. \Box

This result was first motivated by the following example.

Example 2.1.2. We refer to Subsection 3.3 for results on quantum cohomology of cominuscule homogeneous spaces. Let X be a cominuscule homogeneous space and let $A(X) = QH(X)_{q=1}$ be its small quantum cohomology with product \star_0 and with quantum parameter equal to 1. Let pt the the cohomology class of a point and let σ_u be a Schubert class. Consider PD(pt $\star_0 \sigma_u$) where PD stands for Poincaré duality. It was proved in [13] and [14], that this class is a Schubert class $\sigma_{\bar{u}}$ and that $\sigma_u \mapsto \sigma_{\bar{u}}$ defines an algebra involution. Define furthermore a linear form φ by $\varphi(\sigma_u) = \delta_{u,1}$ on the Schubert basis (recall that $1 = \sigma_1$). One easily checks that $Q(\sigma_u, \sigma_{\bar{v}}) = \delta_{u,v}$ proving that Q is an inner product. We recover this way a result of [14] relating the semisimplicity with the existence of an algebra involution.

2.2. Radical and positive definite forms. One of the major problems for applying the above result is that the algebra involution is not *a priori* given and is usually hard to produce (for an example see [16, Remark 6.6]). In this section we furthermore assume that A is a \mathbb{R} -algebra which is $\mathbb{Z}/c_1\mathbb{Z}$ -graded and denote by A_k the graded piece of degree k.

Definition 2.2.1. Let $E \in \text{End}(A)$. We set $A_{\lambda}(E) = \{a \in A \mid (E - \lambda \text{id}_A)^n (a) = 0 \text{ for } n \text{ large}\}$ and $m_E(\lambda) = \dim A_{\lambda}(E)$.

Lemma 2.2.2. Let $E \in \text{End}_A(A)$ with real eigenvalues. We have

$$R(A) \subset \bigoplus_{m_E(\lambda) > 1} A_\lambda(E).$$

In particular if E is semisimple regular, we have R(A) = 0.

Remark 2.2.3. Note that we assume here that *E* is *A*-linear. This is for example the case for $E = E_a$ with $a \in A$.

Proof. We have a decomposition $A = \bigoplus_{\lambda} A_{\lambda}(E)$. Furthermore by A-linearity, we have $A_{\lambda}(E) \cdot A_{\mu}(E) \subset A_{\lambda}(E) \cap A_{\mu}(E) = 0$ for $\lambda \neq \mu$. Let $a \in R(A)$ and write $a = \sum_{\lambda} a_{\lambda}$ with $a_{\lambda} \in A_{\lambda}(E)$. Then a_{λ} is nilpotent. Write $1 = \sum_{\lambda} 1_{\lambda}$ with $1_{\lambda} \in A_{\lambda}(E)$. Then 1_{λ} is an idempotent and for $m_{E}(\lambda) = 1$ we have $a_{\lambda} = x_{\lambda} 1_{\lambda}$ with $x_{\lambda} \in \mathbb{C}$. Since a_{λ} is nilpotent we have $x_{\lambda} = 0$ and the result follows. \Box

Proposition 2.2.4. Assume that there exists $\varphi_0 : A_0 \to \mathbb{C}$ a linear form with $\varphi_0(1) = 1$ such that the bilinear form defined by $Q_0(a, b) = \varphi(ab)$ for $a, b \in A_0$ is positive definite.

1. Then $R(A) \subset A_0(E_a)$ for all $a \in A_1$.

2. If there exists $a \in A_1$ with Ker $E_a = 0$, then A is semisimple.

3. If there exists $a \in A_1$ with $A_0 = A_0 \cap \operatorname{Ker} E_a \oplus A_0 \cap \operatorname{Im} E_a$, then $R(A) \subset \operatorname{Ker} E_a$, $A = \operatorname{Im} E_a \oplus \operatorname{Ker} E_a$ and the subalgebra generated by a is semisimple.

Proof. 1. Let $n = \dim A_0$. As in the proof of Proposition 2.1.1, the endomorphisms $E_a|_{A_0}$ are self-adjoint for Q_0 . In particular they have real eigenvalues and there exists a basis $(e_i)_{i \in [1,n]}$ of orthogonal idempotents whose sum is 1 in A_0 . Let $a \in A_1$. Then $a^{c_1} \in A_0$ and $E_{a^{c_1}|_{A_0}}$ is semisimple. Furthermore there exists $b \in A_0^{\times}$ such that $E_{(ab)^{c_1}|_{A_0}}$ is semisimple and has eigenvalues with multiplicity 1 except maybe for 0. The minimal polynomial μ of $E_{(ab)^{c_1}|_{A_0}}$ therefore has simple roots. Since $1 \in A_0$ this implies that $\mu((ab)^{c_1}) = 0$ and the minimal polynomial of $E_{(ab)^{c_1}}$ divides μ and thus has simple roots. In particular $E_{(ab)^{c_1}}$ is semisimple and has the same eigenvalues as $E_{(ab)^{c_1}}|_{A_0}$.

Let v be an eigenvector of E_{ab} with eigenvalue $\lambda \neq 0$. Write $v = \sum_k v_k$ with $v_k \in A_k$. Then v_k is an eigenvector of $E_{(ab)^{c_1}}$ for the eigenvalue λ^{c_1} . In particular v_0 is the unique (up to scalar) eigenvector of $E_{(ab)^{c_1}}$ with eigenvalue λ^{c_1} . From $E_{ab}(v) = \lambda v$ we deduce $E_{(ab)^{c_1-k}}(v_k) = \lambda^{c_1-k}v_0$. Applying $E_{(ab)^k}$ we get $\lambda^{c_1}v_k = E_{(ab)^{c_1}}(v_k) = \lambda^{c_1-k}E_{(ab)^k}(v_0)$. Finally we have

$$v = \sum_{k=0}^{c_1-1} \lambda^{-k} E_{(ab)^k}(v_0).$$

Note furthermore that for $\lambda \neq 0$ we have $A_{\lambda}(E_{ab}) = \text{Ker}(E_{ab} - \lambda \text{id})$ since $E_{(ab)^{c_1}}$ is semisimple. Therefore $m_{E_{ab}}(\lambda) = 1$ for $\lambda \neq 0$. By the previous Lemma, this implies $R(A) \subset A_0(E_{ab}) = A_0(E_a)$ (the last equality holds since b was chosen invertible).

2. If a is invertible, then $A_0(E_a) = 0$ and the result follows from 1.

3. Let $a \in A_1$. Let μ be the minimal polynomial of $E_{a^{c_1}}$. Since $E_{a^{c_1}}$ is semisimple, we have $\mu(X) = XP(X)$ with P such that P has only simple roots and $P(0) \neq 0$. The minimal polynomial of E_a divides $X^{c_1}P(X^{c_1})$. Let $b = P(a^{c_1}) \in A_0$ and write b = ac + d with $c \in A$ and $d \in A_0 \cap \operatorname{Ker} E_a$.

Lemma 2.2.5. We have $a^i b = 0$ for $i \ge 1$

Proof. Per descending induction on i. For $i \ge c_1$ the result follows from $0 = \mu(a^{c_1}) = a^{c_1}b$. Assume $a^ib = 0$ for i > 1. Then $a^iP(a^{c_1}) = 0$ and the minimal polynomial of E_a divides $X^iP(X)$. In particular Ker $E_{a^{i+1}} = \text{Ker } E_{a^i}$. We have $0 = a^ib = a^{i+1}c + a^id = a^{i+1}c$. Thus $c \in \text{Ker } E_{a^{i+1}} = \text{Ker } E_{a^i}$. This implies $a^{i-1}b = a^ic + a^{i-1}d = a^ic = 0$.

In particular the minimal polynomial of E_a divides $XP(X^{c_1})$ and therefore has only simple roots (over \mathbb{C}). This implies that the subalgebra generated by a is semisimple. It also implies that E_a is semisimple and $A = \text{Im}E_a \oplus \text{Ker} E_a$. This finally implies $A_0(E_a) = \text{Ker} E_a$ and the result follows from 1. \Box

Remark 2.2.6. The above proposition works actually for any $a \in A_k$ such that $gcd(k, c_1) = 1$

2.3. Application to quantum cohomology of Fano varieties. Let X be a smooth complex projective Fano variety of Picard rank 1 and let H be a divisor such that $\mathcal{O}_X(H)$ is an ample generator of the Picard group. The index $c_1(X)$ of X is defined via $-K_X = c_1(X)H$. We write $H(X) = H^*(X, \mathbb{R}), h \in H^2(X, \mathbb{Z})$ for the cohomology class of H and $[\text{pt}] \in H^{2\dim X}(X, \mathbb{Z})$ for the cohomology class of a point in X.

We denote by QH(X) the small quantum cohomology ring obtained using only 3-points Gromov-Witten invariants. We write \star_0 for the product in QH(X). Recall that as \mathbb{R} -vector space we have $QH(X) = H(X) \otimes_{\mathbb{R}} \mathbb{R}[q]$. The ring QH(X) is graded with $\deg(q) = 2c_1(X)$. We write $QH^k(X)$ for the degree k graded piece. We let $A = QH(X)_{q=1}$ be the algebra obtained from QH(X) by quotienting with the ideal (q-1). This algebra is $\mathbb{Z}/2c_1(X)\mathbb{Z}$ -graded and we write A_k for its degree k graded piece.

In particular we have a decomposition

$$A_0 = \bigoplus_{k \ge 0} H^{2kc_1(X)}(X, \mathbb{R}).$$

Denote by $\varphi_0 : A_0 \to \mathbb{R}$ the projection on the first factor and let $Q_X : A_0 \times A_0 \to \mathbb{R}$ be the quadratic form defined by $Q_X(\alpha, \beta) = \varphi_0(\alpha \star_0 \beta)$.

Theorem 2.3.1. Let X be Fano with Picard number 1 and Q_X positive definite. 1. Then $R(A) \subset A_0(E_h)$.

2. If Ker $E_h = 0$, then A is semisimple.

3. If $A_0 = A_0 \cap \operatorname{Ker} E_h \oplus A_0 \cap \operatorname{Im} E_h$, then $R(A) \subset \operatorname{Ker} E_h$, $A = \operatorname{Im} E_h \oplus \operatorname{Ker} E_h$ and the subalgebra generated by h is semisimple.

Proof. Follows from Proposition 2.2.4.

3. Varieties with Q_X positive definite

In this section we give examples of Fano varieties X with Q_X positive definite. These varieties are obtained as complete intersections of homogeneous spaces. The commun feature of our homogeneous spaces is the fact that the variety of conics passing through a point has a positive definite intersection form on its middle cohomology. Note that this positivity property implies strong topological properties (see for example [35] and [36]).

3.1. Fano of large index. We start with general remarks on Q_X for Fano varieties of large index. More precisely, we assume $2c_1(X) > \dim X$. Note that with this assumption we have

$$A_0 = H^0(X, \mathbb{R}) \oplus H^{2c_1(X)}(X, \mathbb{R}).$$

Furthermore the fact that $1 \in H^0(X, \mathbb{R})$ is a unit for the quantum cohomoloy implies $Q_X(1,1) = 1$ and $Q_X(1,\sigma) = 0$ for $\sigma \in H^{2c_1(X)}(X,\mathbb{R})$. To compute Q_X ,

we therefore only have to compute $\sigma \star_0 \sigma'$ for $\sigma, \sigma' \in H^{2c_1(X)}(X, \mathbb{R})$. By dimension arguments we have

$$\sigma \star_0 \sigma' = q \sigma'' + I_2(\sigma, \sigma', \mathrm{pt}) q^2$$

where $\sigma'' \in H^{2c_1(X)}(X, \mathbb{R})$ and $I_2(a, b, c)I_{0,3,2}(a, b, c) =$ is 3-points Gromov-Witten invariant of degree 2 in genus 0 for the classes a, b, c. We obtain the following result.

Lemma 3.1.1. Let X be a Fano variety with $2c_1(X) > \dim X$. Then Q_X is positive definite if and only if $I_2(-, -, \operatorname{pt})$ is positive definite on $H^{2c_1(X)}(X, \mathbb{R})$.

For computing $I_2(-, -, pt)$ we shall prove using Proposition 1.2.1 that the virtual fundamental class is the actual fundamental class.

3.2. Complete intersections in projective spaces. We first consider complete intersections in \mathbb{P}^{n+r} with large index. We reinterpret here results obtained by Beauville [2]. Let X be a smooth complete intersection of r hypersurfaces of degree (d_1, \dots, d_r) in \mathbb{P}^{n+r} with $n \ge 2$. Assume that $n \ge 2\sum (d_i - 1) - 1$.

Lemma 3.2.1. The form Q_X is positive definite.

Proof. This easily follows from the results in [2]. We have a basis $(1, h^{c_1(X)})$ of A_0 and $\varphi_0(h^{c_1(X)}) \neq 0$. Furthermore $h^{c_1(X)} \star_0 h^{c_1(X)} = d_1^{d_1} \cdots d_r^{d_r} h^{c_1(X)}$ proving the result.

Remark 3.2.2. Let X be a complete intersection as above.

1. By Theorem 2.3.1, we have $R(X) \subset A_0(E_h)$. Using the results of Beauville [2] the primitive classes $H_0^{\dim X}(X,\mathbb{R})$ in $H^{\dim X}(X,\mathbb{R})$ (defined as the kernel of E_h) are nilpotent as well as the class $h^{c_1(X)+1} - d_1^{d_1} \cdots d_r^{d_r} qh$. One can furthermore easily check that these classes generate the radical R(X) of QH(X). So QH(X) is not semisimple in general.

2. Tian and Xu [42] proved that the subalgebra generated by the hyperplane class in BQH(X) – the big quantum cohomology – is semisimple for any complete intersection as above.

3. We do not know in general whether BQH(X) is semisimple.

- By results of Hertling, Manin and Teleman [22], a variety has semisimple quantum cohomology only if its cohomology is even and pure of type (p, p). By results of Deligne [19] the only possible complete intersections are quadrics, the cubic surface and even-dimensional complete intersection of two quadrics.
- For the first three it is known that the (small) quantum cohomology is semisimple (see [14], [18]).
- For the last one, we do not know if BQH(X) is semisimple.

3.3. Cominuscule homogeneous spaces. Let G be a semisimple algebraic group. A parabolic subgroup P is called *cominuscule* if its unipotent radical U_P is abelian. This group theoretic condition has many nice implications on the geometry of X = G/P ([40], [37], [38], [41]). Table 1 gives a list of all cominuscule homogeneous spaces.

Recall that the vertices of the Dynkin diagram of G are the simple roots. The marked vertex is the simple root of G which is not a simple root of P. In the above table we denoted Gr(k, n) (resp. LG(n, 2n), OG(n, 2n)) the Graßmann variety of k-subspaces in \mathbb{C}^n (resp. isotropic *n*-subspaces for a symplectic or non degenerate

quadratic form in \mathbb{C}^{2n} , for $OG(n, 2n)$ we only consider a connected component of the
Graßmann variety). We wrote Q_n for a smooth <i>n</i> -dimensional quadric hypersurface
and $\mathbb{OP}^2 = E_6/P_6$ and E_7/P_7 are the Cayley plane and the Freudenthal variety.

Type	X	Diagram	Dimension	$c_1(X)$	$\dim(\Gamma_2)$
A_{n-1}	$\operatorname{Gr}(k,n)$	0000	k(n-k)	n	4
B_n	Q_{2n-1}	• <u> </u>	2n - 1	2n - 1	2n - 1
C_n	$\mathrm{LG}(n,2n)$	∘∘∘=•	$\frac{n(n+1)}{2}$	n+1	3
D_n	Q_{2n-2}		2n - 2	2n - 2	2n - 2
D_n	$\mathrm{OG}(n,2n)$	>	$\frac{n(n-1)}{2}$	2n - 2	6
E_6	\mathbb{OP}^2	•	16	12	8
E_7	E_{7}/P_{7}	°°•	27	18	10

Table 1. List of cominuscule homogeneous spaces.

3.3.1. Cominuscule varieties with Q_X positive definite. Let X be cominuscule. The following results on QH(X) were proved in [14] and [13]. Let pt the the cohomology class of a point and let σ_u be a Schubert class. Then

$$\operatorname{pt} \star_0 \sigma_u = q^{d(u)} \operatorname{PD}(\sigma_{\bar{u}})$$

where d(u) is a non negative integer and $u \mapsto \bar{u}$ is an involution on Schubert classes. It was also proved that $\sigma_u \mapsto q^{-d(u)}\sigma_{\bar{u}}$ defines an algebra involution. As explained in Example 2.1.2 this defines an inner product on $QH(X)_{q=1}$. Note that proving that Q_X is positive definite is equivalent to proving that the classes of degree a multiple of $c_1(X)$ are fixed by the above involution. Since the involution is explicit, an easy check gives the following result.

Proposition 3.3.1. Let X be cominuscule. The form Q_X is positive definite if and only if X is one of the following varieties:

\mathbb{P}^n	$LG(n, 2n)$ for $n \in [3, 4]$
$\operatorname{Gr}(2,n)$ for $n \geq 2$	$OG(n, 2n) \text{ for } n \in [1, 6]$
$Q_n \text{ for } n \geq 2$	\mathbb{OP}^2 or E_7/P_7 .

3.3.2. Geometric proof. We now give a geometric proof of the above result when $2c_1(X) > \dim X$ (this only excludes LG(4,8) of the list). According to Lemma 3.1.1, we only have to understand $I_2(-, -, \text{pt})$ on $H^{2c_1(X)}(X, \mathbb{R})$. We recall a geometric construction for cominuscule homogeneous spaces.

For $x, y \in X$, let d(x, y) be the minimal degree of a rational curve passing through x and y and $d_X(2) = \max\{d(x, y) \mid x, y \in X\}$. We denote by $\Gamma_d(x, y)$ the union of all degree d(x, y) rational curves passing through x and y. Note that $d_X(2) = 1$

if and only if X is a projective space so we may assume $d_X(2) \ge 2$. The following result was proved in [12].

Proposition 3.3.2. Let $d \in [0, d_X(2)]$ and let $x, y \in X$ with d(x, y) = d.

- 1. The variety $\Gamma_d(x, y)$ is a homogeneous Schubert variety in X.
- 2. Any degree d curve is contained in a G-translate of $\Gamma_d(x, y)$.
- 3. A generic degree d curve is contained in a unique G-translate of $\Gamma_d(x, y)$.
- 4. There passes a unique degree d curve through three general points in $\Gamma_d(x, y)$.

For d and $x, y \in X$ as in the former proposition, denote by $Y_d(X)$ the variety of all G-translates of $\Gamma_d(x, y)$. Since $\Gamma_d(x, y)$ is a Schubert variety, its stabiliser is a parabolic subgroup Q and $Y_d(X) = G/Q$. Without loss of generality, we may assume that $P \cap Q$ contains a Borel subgroup. Let $Z_d(X) = G/(P \cap Q)$ be the incidence variety. Write $M_d(X)$ for the moduli space of stable maps of genus 0 and degree d with 3 marked points to X. Let $B\ell_d(X) = \{(\Gamma_d, f) \in Y_d(X) \times M_d(X) \mid f \text{ factors through } \Gamma_d\}$ and $Z_d^{(3)}(X) = \{(\Gamma_d, x_1, x_2, x_3) \in Y_d(X) \times X^3 \mid x_i \in \Gamma_d \text{ for all } i \in [1,3]\}$. We have a diagram



where π is the natural projection, ϕ maps (Γ_d, f) to $(\Gamma_d, \text{ev}_1(f), \text{ev}_2(f), \text{ev}_3(f))$, e_i maps $(\Gamma_d, x_1, x_2, x_3)$ to (Γ_d, x_i) and p and q are the natural projections. The above proposition implies that π and ϕ are both birational. Note also that the third point of Proposition 3.3.2 implies that considering $\Gamma_d \in Y_d(X)$ as a smooth subvariety in X, we have $2 \dim(\Gamma_d) = dc_1(\Gamma_d)$ (see also [12, Formula (5) on Page 73]). For d = 2this implies dim $\Gamma_2 = c_1(\Gamma_2)$ so Γ_2 is a smooth quadric hypersurface (see also Table 1 for its dimension).

By Proposition 1.2.1 and since $M_d(X)$ has expected dimension, for $\sigma, \sigma', \sigma'' \in$ H(X), the Gromov-Witten invariant $I_d(\sigma, \sigma', \sigma'')$ is the push-forward to the point of the class $\operatorname{ev}_1^* \sigma \cup \operatorname{ev}_2^* \sigma' \cup \operatorname{ev}_3^* \sigma''$. Since π and ϕ are birational, an easy diagram chasing gives the following formula (usually called *quantum to classical principle*):

(1)
$$I_d(\sigma,\sigma',\sigma'') = q_*p^*\sigma \cup q_*p^*\sigma' \cup q_*p^*\sigma''.$$

The very first version of this result was proved in [8] for (maximal isotropic) Graßmann varietes and generalised in [12]. For the formal computation in the above setting (and even in equivariant K-theory), we refer to [10, Lemma 3.5].

Proposition 3.3.3. Let X be one of the following varieties

\mathbb{P}^n	$LG(n,2n)$ for $n \in [1,4]$
$\operatorname{Gr}(2,n)$ for $n \geq 2$	$OG(n, 2n)$ for $n \in [1, 6]$
$Q_n \text{ for } n \geq 2$	\mathbb{OP}^2 or E_7/P_7 .

Then $I_2(-, -, pt)$ is positive definite.

Proof. We apply formula (1). Let $\sigma'' = \text{pt}$, then $q_*p^*\text{pt} = j_*[F]$ where F is any fiber of p and $j: F \to Y_d(X)$ the inclusion. Projection formula gives

$$I_d(,\sigma,\sigma',\sigma'') = I_d(,\sigma,\sigma',\mathrm{pt}) = q_*p^*\sigma \cup q_*p^*\sigma' \cup q_*p^*\mathrm{pt} = j^*q_*p^*\sigma \cup j^*q_*p^*\sigma'.$$

The following table gives the list of the varieties F (see [12, Table on Page 71]).

$$\begin{array}{c|c|c} X \\ Y_2(X) \\ F \end{array} \begin{vmatrix} \operatorname{Gr}(2,n) & Q_n & \operatorname{LG}(n,2n) & \operatorname{OG}(n,2n) & E_6/P_6 & E_7/P_7 \\ \operatorname{Gr}(4,n) & \{\mathrm{pt}\} & \operatorname{IG}(n-2,2n) & \operatorname{OG}(n-4,2n) & E_6/P_1 & E_7/P_1 \\ \operatorname{Gr}(2,n-2) & \{\mathrm{pt}\} & \operatorname{Gr}(2,n) & \operatorname{Gr}(4,n) & Q_8 & E_6/P_1. \end{matrix}$$

Table 2. Varieties $Y_2(X)$ and F.

Note that since π and ϕ are birational, we have dim $X + 2c_1(X) = \dim Z_2^{(3)}(X) = \dim Z_2(X) + 2 \dim \Gamma_2 = \dim X + \dim F + 2 \dim \Gamma_2$. In particular we get

$$\dim F = 2(c_1(X) - \dim \Gamma_2).$$

Thus deg $j^*q_*p^*\sigma = \deg q_*p^*\sigma = 2c_1(X) - 2\dim \Gamma_2 = \dim F$ for $\sigma \in H^{2c_1(X)}(X,\mathbb{Z})$. We get an induced map

$$j^*q_*p^*: H^{2c_1(X)}(X,\mathbb{Z}) \to H^{\dim F}(F,\mathbb{Z}).$$

Now for the varieties in the list of the proposition one easily checks (using Schubert classes) that this map is an isomorphism. Furthermore the variety F is of even dimension and has positive definite Poincaré pairing on $H^{\dim F}(F,\mathbb{Z})$ (see [35, Table on Page 572] for the fact that the Poincaré pairing is positive definite on the middle cohomology of $\operatorname{Gr}(2, n)$, Q_8 and E_6/P_1). This finishes the proof.

3.4. Linear sections of cominuscule homogeneous spaces. In this subsection, we extend the result on cominuscule varieties to linear sections of cominuscule varieties. More precisely we prove

Theorem 3.4.1. Let X be a cominuscule variety with Q_X positive definite and let Y be a general linear section of codimension k of X with $2c_1(Y) > \dim Y$. Then Q_Y is positive definite.

Remark 3.4.2. The possible values of k in the above Theorem are as follows:

For \mathbb{P}^n we have $k \leq n$	For $OG(5, 10)$ we have $k \le 5$
For $Gr(2, n)$ we have $k \leq 3$	For OG(6, 12) we have $k \leq 4$
For Q_n we have $k \leq n$	For \mathbb{OP}^2 we have $k \leq 7$
For LG(3,6) we have $k \leq 1$	For E_7/P_7 we have $k \leq 8$.

We will prove this result in two steps. First note that since $2c_1(Y) > \dim Y$ it is enough to check (using Lemma 3.1.1) that $I_2(-, -, \operatorname{pt})$ is positive definite. Let L_k be the linear subspace of codimension $k = \dim X - \dim Y$ cuting Y out of X. Note that since $c_1(Y) = c_1(X) - k$ and $\dim Y = \dim X - k$, the condition $2c_1(Y) > \dim Y$ translates into $k < 2c_1(X) - \dim X$ and that this implies $k < \dim \Gamma_2$ (where Γ_2 is the fiber of $q: Z_2(X) \to Y_2(X)$). 3.4.1. Moduli space $M_2(Y)$. We first prove the following result asserting that $M_2(Y)$ has the expected dimension.

Proposition 3.4.3. We have dim $M_2(Y) = 2c_1(Y) + \dim Y$.

Proof. Note that we have a map of stacks $M_2(Y) \to \mathfrak{M}_{0,3}$ where $\mathfrak{M}_{0,3}$ is the stack of prestable curves of genus 0 with 3 marked points. The fibers of this map are the schemes of morphisms of degree 2 from a fixed prestable curve to X (see for example [3]). In particular the irreducible components of any of these fibers has dimension at least the corresponding expected dimension.

Set $B\ell_2(Y) = \pi^{-1}(M_2(Y))$. We first prove that map $B\ell_2(Y) \to Y_2$ is surjective. Indeed. let $\Gamma_2 \in Y_2$. Then Γ_2 is a quadric of dimension dim $\Gamma_2 > k$. Its intersection with L_k therefore contains a conic. In particular, there exists a stable map in $M_2(Y)$ factorising through Γ_2 proving the surjectivity. The fiber of the map $B\ell_2(Y) \to Y_2(X)$ over Γ_2 is therefore given by the genus zero stable maps of degree 2 and three marked points to $\Gamma_2 \cap L_k$. We thus need to understand this intersection more precisely. We shall consider L_k as the intersection of k hyperplanes H_1, \dots, H_k .

- 1. the intersection $\Gamma_2 \cap L_k$ is a smooth quadric of dimension dim $\Gamma_2 k$,
- 2. the intersection $\Gamma_2 \cap L_k$ is a quadric of dimension dim $\Gamma_2 k$ and rank $\ell < \dim \Gamma_2 k + 1$,
- 3. we have $\dim(\Gamma_2 \cap L_k) > \dim \Gamma_2 k$.

Since the dimension of the moduli space of genus zero stable maps of degree 2 to a quadric of given rank is well known, an easy check proves that the locus in $B\ell_2(Y)$ over points $\Gamma_2 \in Y_2(X)$ such that the intersection $\Gamma_2 \cap L_k$ is for each prestable curve in $\mathfrak{M}_{0,3}$ of dimension strictly less than the expected dimension. In particular irreducible components of $M_2(Y)$ come from irreducible components of $B\ell_2(Y)$ containing points mapping in Y_2 to an quadric Γ_2 such that $\Gamma_2 \cap L_k$ is a smooth quadric of dimension dim $\Gamma_2 - k$. Since any stable map to a such intersection $\Gamma_2 \cap L_k$ is a limit of a stable map from an irreducible curve we can consider only irreducible curves. This implies that $M_2(Y)$ is irreducible of expected dimension dim $Y + 2c_1(Y)$.

3.4.2. Proof of Theorem 3.4.1. Since $M_2(Y)$ has expected dimension, the virtual class is the fundamental class by Proposition 1.2.1. Let $Z_2(Y) = p^{-1}(Y)$ and denote by r and s the projections $r : Z_2(Y) \to Y$ and $s : Z_2(Y) \to Y_2(X)$. Since π and ϕ restricted to $B\ell_2(Y)$ are again birational, the same computation as in the cominuscule case gives the relation

$$I_2(\tau, \tau', \tau'')_Y = r_* s^* \tau \cup r_* s^* \tau' \cup r_* s^* \tau''$$

where $I_2(-, -, -)_Y$ denotes the Gromov-Witten invariants in degree 2 in Y. But the diagram

is Cartesian with p flat (it is a locally trivial fibration since X = G/P, see for example [7, Proposition 2.3]). In particular we have $i_*r^* = p^*j_*$. We thus get $s_*r^*\tau = q_*i_*r^*\tau = q_*p^*j_*\tau$. We deduce $I_2(\tau, \tau', \tau'')_Y = I_2(j_*\tau, j_*\tau', j_*\tau'')_X$. In particular the result follows since $I_2(-, -, \text{pt})_X$ is positive definite. **Remark 3.4.4.** Note that we proved more than Theorem 3.4.1. Indeed, for any cohomology classes $\tau, \tau', \tau'' \in H(Y)$ we have the equality

$$I_2(\tau, \tau', \tau'')_Y = I_2(j_*\tau, j_*\tau', j_*\tau'')_X.$$

3.4.3. *Examples.* Several linear sections satisfying Theorem 3.4.1 are classical.

1. Hyperplane sections of the Graßmann variety $\operatorname{Gr}(2, n)$. The Plücker embedding is given by the representation $\Lambda^2 \mathbb{C}^n$. A general linear section corresponds to a general symplectic form on \mathbb{C}^n and the hyperplane section is the subvariety of isotropic 2-dimensional subspaces.

For n = 2p even, the variety Y = IG(2, 2p) is homogeneous.

For n = 2p+1 odd, the variety Y = IG(2, 2p+1) is not homogeneous. This variety has two orbits under its automorphism group and is known as the odd symplectic Graßmann variety of lines. We refer to [30, 34, 32, 33] for several geometric results on this variety.

2. Hyperplane sections of $\mathbb{OP}^2 = E_6/P_6$ are homogeneous under the group F_4 . Actually we have $Y = F_4/P_4$ which is the coadjoint variety of type F_4 (see [27, 16]).

3. For X = Gr(2,5) and k = 3, then $Y = V_5$ is the *del Pezzo* threefold of index 2 and degree 5.

4. Note that we obtain almost all Fano varieties X of coindex 3 *i.e.* with $c_1(X) = \dim X - 2$: we obtain all Fano varieties X of coindex 3 with genus $g \in [7, 10]$ *i.e.* missing the extremal values g = 6 and g = 12 (see [31, Theorem 5.2.3]).

3.5. Adjoint varieties. The last family of varietes with Q_X positive definite are the adjoint varieties (see [16]). These are homogeneous spaces and can be defined, for G a semisimple group, as the closed G-orbit in $\mathbb{P}\mathfrak{g}$ where \mathfrak{g} is the Lie algebra of G and G acts on its Lie algebra by via the adjoint representation. The list of adjoint varieties is given in Table 3. Note that the following equality holds: $\dim X = 2c_1(X) - 1$ (except in type C_n).

The following result was proved in [16, Proof of Proposition 6.5]

Proposition 3.5.1. Let X be an adjoint variety. Then Q_X is positive definite.

Type	variety	diagram	dimension	index
A_n	$\operatorname{Fl}(1,n;n+1)$	• <u> </u> • <u> </u> • <u> </u> •	2n - 1	(n,n)
B_n	OG(2, 2n+1)	o —● —o—o — o — o	4n - 5	2n - 2
C_n	\mathbb{P}^{2n-1}	•ooooo	2n - 1	2n
D_n	OG(2,2n)	·	4n - 7	2n - 3
E_6	E_{6}/P_{2}	°>>	21	11
E_7	E_{7}/P_{1}	•	33	17
E_8	E_{8}/P_{8}	°	• 57	29
F_4	F_4/P_1	° ●—○ — ○—○	15	8
G_2	G_2/P_1	• = 0	5	3

Table 3. List of adjoint varietes.

4. Semisimplicity of the quantum cohomology

In this section we apply the results in Section 2 to the varieties of Section 3 and get results on the semisimplicity of their quantum cohomology.

4.1. Linear sections of cominuscules homogeneous spaces. We consider the varieties Y obtained as linear sections of a cominuscule homogeneous space X satisfying the assumptions of Theorem 3.4.1. These varieties are listed in Remark 3.4.2.

4.1.1. Multiplication with degree 2 classes. We want to understand the endomorphism E_h^Y of QH(Y) obtained by multiplication with h the hyperplane class. Let $j: Y \to X$ be the inclusion and let $\tau \in H(Y)$. We denote by h the hyperplane class in H(X) and H(Y) as well. Projection formula gives the following result.

Lemma 4.1.1. We have $j_*(h \cup \tau) = h \cup j_*\tau$.

As in the proof of Theorem 3.4.1 (see also Remark 3.4.4) we prove a result comparing Gromov-Witten invariants on X and Y. Write $I_d(-, -, -)_X$ and $I_d(-, -, -)_Y$ for Gromov-Witten invariants of degree d in X and Y.

Lemma 4.1.2. Let $\tau, \tau' \in H(Y)$ be cohomology classes such that the following conditions hold.

- $\deg \tau + \deg \tau' = 2 \dim Y + 2c_1(Y) 2.$
- There exists varieties S, S' in Y with $j_*\tau = [S], j_*\tau' = [S']$ which are in general position in X.

Then $I_1(\tau, \tau', h)_Y = I_1(j_*\tau, j_*\tau', h)_X$

Proof. Note that the equality on degrees is equivalent to $\operatorname{codim}_X S + \operatorname{codim}_X S' = \dim X + c_1(X) - 1$ and $\operatorname{codim}_Y S + \operatorname{codim}_Y S' = \dim Y + c_1(Y) - 1$. This together with the second condition imply the following property: the scheme of 2-points degree 1 stable maps to X passing through S and S' is finite and reduced. Remark that any degree 1 stable map to Y passing through S and S' is a degree 1 stable map to X passing through S and S' is a linear section of X. In particular the scheme of degree 1 stable maps to Y passing through S and S' is finite and reduced. This implies that the moduli space of degree 1 stable maps to Y has the expected dimension and that the above number of stable maps is equal to both $I_1(\tau, \tau', h)_Y$ and $I_1(j_*\tau, j_*\tau', h)_X$.

Proposition 4.1.3. Let a, b be integers in $[0, \dim Y]$ such that

- $a + b = \dim Y + c_1(Y) 1$.
- $k \ge c_1(Y)$.

Then there exists basis of classes τ, τ' in $j^*H^{2a}(X, \mathbb{R})$ and $j^*H^{2b}(X, \mathbb{R})$ such that the assumptions of Lemma 4.1.2 are satisfied.

Remark 4.1.4. Note that $a, b \ge c_1(Y) - 1$. By the Hard Lefschetz Theorem and since $2c_1(Y) > \dim Y$ we get $j^*H^{2a}(X, \mathbb{R}) = H^{2a}(Y, \mathbb{R})$ except if $a = c_1(Y) - 1$ and $\dim Y = 2c_1(Y) - 1$.

Proof. We may assume $a \ge b$. We shall construct general hyperplane sections $Y = X \cap L_k$ with L_k a general linear subspace of codimension k satisfying the proposition.

Case 1. We first deal with the case where $a = \dim Y$ or $\dim Y - 1$ *i.e.* there is a unique class τ in $H^{2a}(Y, \mathbb{R})$: the class pt or a point or the class ℓ of a line.

We prove that τ and the pull-back via j of the Schubert basis in $H^{2b}(X,\mathbb{Z})$ satisfy the proposition. Let S be a Schubert variety of codimension b in X and S'be a point or a line (depending on whether $a = \dim Y$ or $a = \dim Y - 1$). Let S be family of G-translates of S and \mathfrak{T} the family of G-translates of T. Let \mathcal{L}_k be the variety parametrising linear subspaces of codimension k in the Plücker embedding of X. We have a rational morphism $f: G \times \mathcal{L}_k \to \mathcal{H}$ where \mathcal{H} is a Hilbert scheme of subvarieties in X defined by $(g, L_k) \mapsto gS \cap L_k$. Let \mathcal{V} be the closure of its image.

We consider $I = \{(V, g'T, L_k) \in \mathcal{V} \times \mathcal{T} \times \mathcal{L}_k \mid V \subset L_k \supset g'T\}$. We have a diagram

Since $b = \dim Y + c_1(Y) - 1 - a$, the Schubert variety S is contained in a linear section of X of codimension $\dim Y + c_1(Y) - 1 - a$ and thus $V = gS \cap L_k$ is contained in a linear section of X of codimension $\dim Y + c_1(Y) - 1 - a + k$. In particular since the space of T has dimension $\dim Y - a$, the variety $V \cup g'T$ is contained in a linear section of codimension $c_1(Y) - 1 + k \ge k$. This proves that the map p is surjective. The map q is also surjective: for $L_k \in \mathcal{L}_k$ pick for g'T a point or a line in $X \cap L_k$, pick $g \in G$ general and set $V = gS \cap L_k$.

In particular, for $L_k \in \mathcal{L}_k$ general, there is a translate gS and a point or a line $T \subset X \cap L_k$ such that $V = gS \cap L_k$ and T are in general position in X. Setting $Y = X \cap L_k$ proves the result.

Case 2. We now consider the other cases. We prove that the pull-backs via j of the Schubert basis in $H^{2a}(X, \mathbb{R})$ and $H^{2b}(X, \mathbb{R})$ satisfy the proposition. Let S and S' be Schubert varieties in X of respective codimension a and b. Denote by S and S' the family of G-translates of S and S'. These are projective homogeneous spaces since the stabiliser of S and S' contain a Borel subgroup. Let \mathcal{L}_k be the projective variety of all linear subspaces of codimension k in the Plücker embedding of X.

We have a rational morphism $f : S \times S' \times \mathcal{L}_k \times \mathcal{L}_k \to \mathcal{H} \times \mathcal{H}'$, where \mathcal{H} and \mathcal{H}' are Hilbert schemes of subvarieties in X, defined by $(gS, g'S', L_k, L'_k) \mapsto (gS \cap L_k, g'S' \cap L'_k)$. Let $\mathcal{V} \times \mathcal{V}'$ be the closure of the image of f.

Lemma 4.1.5. Let $\Delta : \mathbb{S} \times \mathbb{S}' \times \mathcal{L}_k \to \mathbb{S} \times \mathbb{S}' \times \mathcal{L}_k \times \mathcal{L}_k$ be the map induced by the diagonal embedding of \mathcal{L}_k . Then $\mathcal{V} \times \mathcal{V}'$ is the image of $f \circ \Delta$.

Assume that the lemma holds. Let O be the open subset in $\mathcal{V} \times \mathcal{V}'$ of subvarieties in general position and let \mathcal{L}_k° be the open non empty subset in \mathcal{L}_k of linear subspaces having a smooth intersection with X. Then $(f \circ \Delta)^{-1}(O)$ is non empty and open in $\mathcal{S} \times \mathcal{S}' \times \mathcal{L}_k$ as well as $\operatorname{pr}_3^{-1}(\mathcal{L}_k^\circ)$ where pr_3 is the projection on the third factor in $\mathcal{S} \times \mathcal{S}' \times \mathcal{L}_k$. Since $\mathcal{S} \times \mathcal{S}' \times \mathcal{L}_k$ is irreducible, these subsets intersect. Let $(gS, g'S', L_k)$ be in the intersection. Then $Y = X \cap L_k$ and $V = Y \cap gS$, $V' = Y \cap g'S'$ satisfy the desired property since V and V' are in general position in X.

We are left to proving the lemma. Equivalently we need to prove that for $(gS, L_K) \in S \times \mathcal{L}_k$ and $(g'S', L'_k) \in S' \times \mathcal{L}_k$ such that $\operatorname{codim}_{gS} gS \cap L_k = k = \operatorname{codim}_{g'S'} g'S' \cap L'_k$, there exists $L''_k \in \mathcal{L}_k$ such that $gS \cap L_k \subset L''_k \supset g'S' \cap L'_k$. Let W be the vector space defining the Plücker embedding and let $\langle gS \rangle$ and $\langle g'S' \rangle$ be the spans in W of element whose classes are in gS and g'S'. It is enough to prove that $\dim \langle gS \rangle + \dim \langle g'S' \rangle \leq \dim W + k$. We prove this inequality by case by case analysis.

For X = LG(3, 6), we have k = 1 and $\dim Y + c_1(Y) - 1 = 7$ thus $a \ge \dim Y - 1$ and there is nothing to prove.

Note that in the other cases X is a minuscule homogeneous space. This means that the weights of W for a maximal torus form a unique orbit for the action of the Weyl group of G. This in particular implies that there is a correspondence between Schubert varieties and weights of W. As a consequence we get the equality

$$\dim \langle S \rangle = |\{S'' \subset S \mid S'' \text{ a Schubert variety}\}|.$$

In words: the dimension of the span $\langle S \rangle$ of a Schubert variety S is equal to the number of Schubert varieties contained in S. This translates the inequality $\dim \langle gS \rangle + \dim \langle g'S' \rangle \leq \dim W + k$ into a combinatorial computation and an easy case by case check gives the result. \Box

We shall now define maps between subspaces of QH(Y) and QH(X). Recall that we have morphisms $j^* : H^m(X, \mathbb{Z}) \to H^m(Y, \mathbb{Z})$ and $j_* : H_m(X, \mathbb{Z}) \to H_m(X, \mathbb{Z})$ which become isomorphisms for $m < \dim Y$ by Lefschetz Theorem. Let A(Y) and A(X) be the algebras obtained from QH(Y) and QH(X) by quotienting with the ideal (q-1). Recall that these algebras are respectively $\mathbb{Z}/2c_1(Y)\mathbb{Z}$ and $\mathbb{Z}/2c_1(X)\mathbb{Z}$ graded. We write $A_a(Y)$ and $A_a(X)$ for their degree a graded piece. We have

$$\begin{aligned} A_{a}(X) &= H^{a}(X, \mathbb{R}) \oplus H^{2c_{1}(X)+a}(X, \mathbb{R}) & \text{for } a \in [0, 2 \dim Y - 2c_{1}(Y)] \\ A_{a}(Y) &= H^{a}(Y, \mathbb{R}) \oplus H^{2c_{1}(Y)+a}(Y, \mathbb{R}) & \text{for } a \in [0, 2 \dim Y - 2c_{1}(Y)] \\ A_{2c_{1}(Y)-2}(X) &= H^{2c_{1}(Y)-2}(X, \mathbb{R}) & \text{for } 2c_{1}(Y) - 1 > \dim Y \\ A_{2c_{1}(Y)-2}(X) &= H^{2c_{1}(Y)-2}(Y, \mathbb{R}) \oplus H^{2\dim(X)}(X, \mathbb{R}) & \text{for } 2c_{1}(Y) - 1 > \dim Y \\ A_{2c_{1}(Y)-2}(X) &= H^{2c_{1}(Y)-2}(X, \mathbb{R}) \oplus H^{2\dim(X)}(X, \mathbb{R}) & \text{for } 2c_{1}(Y) - 1 = \dim Y \\ A_{2c_{1}(Y)-2}(Y) &= H^{2c_{1}(Y)-2}(Y, \mathbb{R}) \oplus H^{2\dim(Y)}(Y, \mathbb{R}) & \text{for } 2c_{1}(Y) - 1 = \dim Y \end{aligned}$$

We define a morphism J between these spaces as follows:

$$J = j^* \oplus j_*^{-1}: \quad A_a(X) \to A_a(Y) \qquad \text{for } a \in [2c_1(Y), 2 \dim Y]$$

$$J = j^*: \quad A_{2c_1(Y)-2}(X) \to A_{2c_1(Y)-2}(Y) \qquad \text{for } 2c_1(Y) - 1 > \dim Y$$

$$J = j^* \oplus j_*^{-1}: \quad A_{2c_1(Y)-2}(X) \to A_{2c_1(Y)-2}(Y) \qquad \text{for } 2c_1(Y) - 1 = \dim Y$$

Note that by Lefschetz Theorem and because $2c_1(Y) > \dim Y$, the first map is an isomorphism for all $a \in [2c_1(Y), 2 \dim Y]$.

Corollary 4.1.6. For all $a \in [2c_1(Y), 2 \dim Y - 2]$, we have a commutative diagrams

$$\begin{array}{c|c} A_{a}(X) \xrightarrow{J} A_{a}(Y) & and & A_{2c_{1}(Y)-2}(X) \xrightarrow{J} A_{2c_{1}(Y)-2}(Y) \\ \hline E_{h}^{X} & & \downarrow E_{h}^{Y} & E_{h^{k+1}}^{X} & \downarrow E_{h}^{Y} \\ A_{a+2}(X) \xrightarrow{J} A_{a+2}(Y) & & A_{0}(X) \xrightarrow{J} A_{0}(Y). \end{array}$$

Proof. For a quantum product $a \star_0 b$ we write $a \star_0 b = \sum_d q^d (a \star_0 b)_d$.

We start with the first square. Let $\sigma \in A_a(Y)$. We have deg $\sigma = a$ or deg $\sigma = 2c_1(X) + a$. In the first case $J\sigma = j^*\sigma$ and deg $J\sigma + 2 = \deg \sigma + 2 < 2c_1(Y) < 2c_1(X)$. In particular $E_X(\sigma) = h \cup \sigma$ and $E_h^Y(Js) = h \cup J\sigma$ and we get $E_h^YJ(\sigma) = h \cup j^*\sigma = j^*(h \cup \sigma) = JE_h^X(\sigma)$. In the second case we have $\sigma = j_*\tau$ with $\tau = J\sigma$. We get $JE_h^X(\sigma) = J(h\star_0 j_*\tau)$ and $E_h^YJ(\sigma) = E_h^Y(\tau) = h\star_0\tau$. By Lemma 4.1.1 we have $j_*(h\star_0\tau)_0 = (h\star_0 j_*\tau)_0$ so the result is true for the classical part of the quantum product. The quantum parts of $E_h^YJ(\sigma)$ and $JE_h^XJ(\sigma)$ are of the form

$$(h \star_0 \tau)_1 = I_1(h, \tau, \ell)_Y qh$$
 and $J((h \star_0 j_* \tau)_1) = I_1(h, j_* \tau, \ell)_X qh$.

Since $2c_1(Y) > \dim Y$, the Hard Lefschetz Theorem implies that there is a σ'' with $\tau = j^* \sigma''$. Applying the above proposition, we get $I_1(h, \tau, \ell)_Y = I_1(h, j_* \tau, \ell)_X$ proving the result.

We now consider the second square. The possible degrees for $\sigma \in A_{2c_1(Y)-2}(X)$ are $2c_1(Y) - 2$ or $2 \dim X$ if $\dim Y = 2c_1(Y) - 1$. First assume deg $\sigma = 2c_1(Y) - 2$ and let $\tau = J(\sigma) = j^* \sigma$. We have $JE_{h^{k+1}}^X(\sigma) = J(h^{k+1} \star_0 \sigma)$ and $E_h^Y J(\sigma) = h \star_0 \tau$. For degree reasons, we have $h^k = [Y]$ and $[Y] \star_0 \sigma = [Y] \cup \sigma$. We get $E_{h^{k+1}}^X(\sigma) = h^{k+1} \star_0 \sigma = h \star_0 ([Y] \cup \sigma) = h \star_0 j_* j^* \sigma$. By Lemma 4.1.1, we have $j_*(h \star_0 \tau)_0 = j_*(h \cup \tau) = h \cup j_* \tau = h \cup j_* j^* \sigma$ and the result is true for the classical part of the quantum product. The quantum part of $E_h^Y J(\sigma)$ and $JE_{h^{k+1}}^X(\sigma)$ are of the form

$$(h \star_0 \tau)_1 = I_1(h, \tau, \text{pt})_Y q \text{ and } J((h \star_0 j_* \tau)_1) = I_1(h, j_* \tau, \text{pt})_X q.$$

By the above proposition, we get $I_1(h, \tau, \text{pt})_Y = I_1(h, j_*\tau, \text{pt})_X$ proving the result for deg $\sigma = 2c_1(Y) - 2$.

Finally assume dim $Y = 2c_1(Y) - 1$ and deg $\sigma = 2 \dim X$. We have $\sigma = pt$ and $J(\sigma) = pt$. We get $JE_{h^{k+1}}^X(\sigma) = J(h^{k+1}\star_0 pt)$ and $E_h^Y J(\sigma) = h\star_0 pt$. We have $h\star_0 pt = q \sum_{\gamma} I_1(h, pt, \gamma)_Y PD(\gamma) + I_2(h, pt, pt)_Y q^2$ where the sum runs over a basis of $H^{2c_1(Y)-2}(Y, \mathbb{R})$. By Lefschetz Theorem the classes δ with $J\delta = j^*\delta = \gamma$ form a basis of $H^{2c_1(Y)-2}(X, \mathbb{R})$. By the above proposition, we have $I_1(h, pt, \gamma)_Y =$ $I_1(h, pt, j_*\gamma)_X$. On the other hand, applying Remark 3.4.4 we have $I_2(h, pt, pt)_Y =$ $I_2(j_*h, pt, pt)_X$. We get

$$\begin{split} E_h^Y J(\sigma) &= q \sum_{\sigma} I_1(h, \mathrm{pt}, j_* j^* \delta)_X \mathrm{PD}(j^* \delta) + I_2(j_* h, \mathrm{pt}, \mathrm{pt})_X q^2 \\ &= q \sum_{\sigma} I_1(h, \mathrm{pt}, [Y] \cup \delta)_X j^* \mathrm{PD}(\delta) + I_2([Y] \cup h, \mathrm{pt}, \mathrm{pt})_X q^2 \\ &= q \sum_{\sigma} I_1(h, \mathrm{pt}, [Y] \star_0 \delta)_X j^* \mathrm{PD}(\delta) + I_2([Y] \star_0 h, \mathrm{pt}, \mathrm{pt})_X q^2 \\ &= q \sum_{\sigma} I_1([Y] \star_0 h, \mathrm{pt}, \delta)_X j^* \mathrm{PD}(\delta) + I_2(h^{k+1}, \mathrm{pt}, \mathrm{pt})_X q^2 \\ &= q j^* \sum_{\sigma} I_1(h^{k+1}, \mathrm{pt}, \delta)_X \mathrm{PD}(\delta) + I_2(h^{k+1}, \mathrm{pt}, \mathrm{pt})_X q^2 \\ &= J E_{h^{k+1}}^X(\mathrm{pt}) \end{split}$$

This completes the proof.

Remark 4.1.7. For dim $Y < 2c_1(Y) - 2$ using the same arguments as above we get a slightly better factorisation. We have an isomorphism J defined by

$$J = j_*^{-1} : A_{-2}(X) = H^{2c_1(X)-2}(X,\mathbb{R}) \to A_{-2}(Y) = H^{2c_1(Y)-2}(Y,\mathbb{R}).$$

For dim $Y < 2c_1(Y) - 2$, we have a commutative diagram

$$\begin{array}{c|c} A_{-2}(X) \xrightarrow{J} & A_{-2}(Y) \\ \hline E_h^X & & \downarrow E_h^Y \\ \hline A_0(X) \xrightarrow{J} & A_0(Y). \end{array}$$

4.2. Semisimple small quantum cohomology. In this subsection, we prove the following semisimplicity result.

Theorem 4.2.1. Let Y be a general hyperplane section with $2c_1(Y) > \dim Y$ of the following homogeneous space X:

$$\begin{array}{lll} & {\rm Gr}(2,2n+1) & & F_4/P_1 \\ & {\rm OG}(5,10) & & {\rm OG}(2,2n+1) \\ & {\rm LG}(3,6) & & & G_2/P_1. \end{array}$$

Then QH(Y) is semisimple.

Proof. Note that for the varieties of the second column, we have $2c_1(X) - 1 = \dim X$ thus we must have Y = X. By Theorem 3.4.1 and Proposition 3.5.1, the quadratic form Q_Y is positive definite. By Theorem 2.3.1, it is enough to prove that Ker $E_h^Y = 0$ *i.e.* that E_h^Y is bijective. It is therefore enough to prove that h is invertible *i.e.* that E_h^Y is surjective onto $A_0(Y)$. By Corollary 4.1.6 it is enough to prove that E_h^X is surjective onto $A_0(X)$. This is now an easy check using [12] and [16].

Remark 4.2.2. Note that this result together with the result of [22] implies that the cohomology of the varieties Y above is even and of pure type (p, p).

18

4.3. Structure of the radical. In this subsection, we describe the radical of QH(Y) for some Fano varieties Y.

Proposition 4.3.1. Let Y be a general hyperplane section with $2c_1(Y) > \dim Y$ of the following homogeneous space X:

$\operatorname{Gr}(2,2n)$	OG(2,2n)
OG(6, 12)	E_{6}/P_{2}
E_{6}/P_{6}	E_{7}/P_{1}
E_{7}/P_{7}	$E_8/P_8.$

Then

$$R(Y) = \operatorname{Ker} E_h^Y \cap \bigoplus_{2c_1(Y) \not \mid k} \operatorname{QH}^k(Y).$$

Proof. Note that for the varieties of the second column, we have $2c_1(X) - 1 = \dim X$ thus we must have Y = X.

We first prove $R(Y) \subset \text{Ker} E_h^Y$. By Theorem 3.4.1 and Proposition 3.5.1, the quadratic form Q_Y is positive definite. By Theorem 2.3.1, it is enough to prove that $A_0(Y) = \text{Ker} E_h^Y \cap A_0(Y) \oplus \text{Im} E_h^Y \cap A_0(Y)$. By Corollary 4.1.6 this is equivalent to the same statement in $A_0(X)$. This is now an easy check using [12] and [16].

Note that in $\operatorname{QH}_0(Y)$, there is a unique vector $K \in \operatorname{Ker} E_h^Y$ and this vector is of the form $K = \lambda + v$ with $\lambda \in \mathbb{R} \setminus \{0\}$ and $v \ni \operatorname{Im} E_h^Y$. We thus have $K^n = \lambda^{n-1}K$ and K is not nilpotent. We now prove that any element $K \in$ $\operatorname{Ker} E_h^Y \cap \bigoplus_{2c_1(Y) \not \mid k} \operatorname{QH}^k(Y)$ is nilpotent. Remark that for degree reasons there will be a power K^n of K in $\operatorname{Im} E_h^Y$ (by Corollary 4.1.6 and case by case inspection on X. We refer to Subsection 5.2 for a detailled proof of this in the case $X = \operatorname{Gr}(2, 2n)$). But K^n is in $\operatorname{Ker} E_h^Y$ thus $K^n = 0$. \Box

Example 4.3.2. Let X = Gr(2, 6) and Y be a general hyperplane section. Then Y is isomorphic to an isotropic Grasmann variety IG(2, 6). We have dim Y = 7 and $c_1(Y) = 5$. The dimensions of the graded parts of A(Y) are

One easily checks that $\operatorname{Ker} E_h^Y$ has dimension 2. There is an element K_0 of degree 0 in $\operatorname{Ker} E_h^Y$ and an element K_4 of degree 4 in $\operatorname{Ker} E_h^Y$. The image $\operatorname{Im} E_h^Y$ is a complement of $\operatorname{Ker} E_h^Y$. As in the above proposition K_0 is not nilpotent and K_4^2 has degree 8 this is in $\operatorname{Im} E_h^Y$ so $K_4^2 = 0$ and we have $R(Y) = \langle K_4 \rangle$. We recover the example in [14, Section 7].

5. BIG QUANTUM COHOMOLOGY

In this section we consider the case of two Fano varieties Y obtained as hyperplane section of a cominuscule homogeneous space X whose small quantum cohomology QH(Y) is not semisimple. We however prove that the big quantum cohomology BQH(Y) is semisimple. These are the first examples of semisimplicity of the big quantum cohomology in the presence of non semisimple small quantum cohomology.

The varieties we consider are Y = IG(2, 2n) obtained as hyperplane section of X = Gr(2, 2n) and $Y = F_4/P_4$ obtained as hyperplane section of $X = E_6/P_6$. Note that both varieties are homogeneous and actually *coadjoint varieties* in the sense of

[16]. Their small quantum cohomology is not semisimple but well understood. We refer to [16] for more details. In the next subsection, we recall few fact on QH(Y) for Y coadjoint.

5.1. Quantum cohomology of coadjoint varieties. Let Y be one of the following two varieties Y = IG(2, 2n) or $Y = F_4/P_4$ which are homogeneous under the action of a reductive group G of type C_n of F_4 respectively.

5.1.1. Cohomology and short roots. The cohomology of a coadjoint variety Y homogeneous under the action of a reductive group G is easily described using Schubert classes. There are several indexing sets for Schubert varieties. We will choose the indexing set described in [16]. Let R_s be the set of short roots in R the root system of G. For each root $\alpha \in R_s$, there is a cohomology class σ_{α} and the family $(\sigma_{\alpha})_{\alpha \in R_s}$ form a basis of H(X).

Let *n* be the rank of the group (and of the roots system *R*). We choose $(\alpha_1, \dots, \alpha_n)$ a Basis of the root system with notation as in [6]. For a root α we have an expression

$$\alpha = \sum_{i=1}^{n} a_i \alpha_i$$

with $a_i \in \mathbb{Z}$ for all $i \in [1, n]$. We define the height $ht(\alpha)$ of $\alpha \in R$ by

$$\operatorname{ht}(\alpha) = \sum_{i=1}^{n} a_i$$

Let θ be the highest short root of R. The above indexing satisfies many nice properties. We have (see [16, Proposition 2.9])

$$\deg(\sigma_{\alpha}) = \begin{cases} 2(\operatorname{ht}(\theta) - \operatorname{ht}(\alpha)) & \text{for } \alpha \text{ positive,} \\ 2(\operatorname{ht}(\theta) - \operatorname{ht}(\alpha) - 1) & \text{for } \alpha \text{ negative.} \end{cases}$$

We will write 1 for the class σ_{θ} and h for the hyperplane class in the Plücker embedding of X. The Poincaré duality has a very simple form on roots: the Poincaré dual σ_{α}^{\vee} of σ_{α} is simply $\sigma_{\alpha}^{\vee} = \sigma_{-\alpha}$ (see [16, Proposition 2.9]).

5.1.2. Small quantum cohomology and affine short roots. The above parametrisation of Schubert classes by short roots can be extended to quantum monomials in $QH(X)_{loc} = QH(X)[q, q^{-1}]$. A quantum monomial is a element $q^d \sigma_{\alpha}$ where $d \in \mathbb{Z}$ and $\alpha \in R_s$. We write QM(X) for the set of all quantum monomials and we write $QH^k(X)_{loc}$ for the degree k part of $QH(X)_{loc}$.

Let \widehat{R} be the extended affine root system of R and let δ be the minimal positive imaginary root. The extended root system has basis $(\alpha_0, \dots, \alpha_n)$ and in this basis we have $\delta = \Theta + \alpha_0$ where Θ is the highest root of R. A short root of \widehat{R} is a root of the form $\alpha + d\delta$ for $\alpha \in R_s$ and $d \in \mathbb{Z}$. We write \widehat{R}_s for the set of short roots in \widehat{R} . There is a bijection

$$\eta: \hat{R}_s \to \mathrm{QM}(X)$$

defined by $\eta(\alpha - d\delta) = q^d \sigma_{\alpha}$. Note that we can extend the height function on \widehat{R} and that we have $\deg(q) = 2(\operatorname{ht}(\delta) - 1)$.

5.1.3. Multiplication with the hyperplane. We have the following very simple description of the small quantum product \star_0 with h (see [16, Theorem 3]):

$$h \star_0 \sigma_\alpha = \sum_{i \in [0,n], \ \langle \alpha_i^\vee, \alpha \rangle > 0} \langle \alpha_i^\vee, \alpha \rangle \eta(s_{\alpha_i}(\alpha))$$

5.2. Graßmann variety of lines. Let X = Gr(2, 2n) and let Y be a linear section of X of codimension 1. Note that Y = IG(2, 2n) is an isotropic Grasmann variety. The small quantum cohomology of this variety is described in [16]. We first prove the following result which improves Proposition 4.3.1.

Lemma 5.2.1. Let Y be a linear section of X = Gr(2, 2n) of codimension 1.

- 1. There exists a unique element K_{4n+2} modulo scalar in $\operatorname{QH}^{4n+2}(Y) \cap \operatorname{Ker} E_h^Y$. 2. The element K_{4n+2}^k is divible by q^k . 3. We have $R(Y) = \langle K_{4n+2}, \cdots, q^{-(n-2)} K_{4n+2}^{n-2} \rangle$ and $K_{4n+2}^{n-1} = 0$.

Proof. 1. Note that all odd dimension cohomolgy groups of X and Y vanish. Recall also that we have the inclusion $R(Y) \subset \operatorname{Ker} E_h^Y$. Using the description of Schubert classes with partitions having 2 parts, we have

$$\dim \operatorname{QH}^{2a}(X) = \begin{cases} n & \text{for } a \text{ even} \\ n-1 & \text{for } a \text{ odd.} \end{cases}$$

We deduce results on the dimension of the graded parts in QH(Y). Recall that multiplication with q induces an isomorphism $QH^{2a}(Y) \to QH^{2a+2c_1(Y)}(Y)$ so we only need to describe these dimensions for $0 \le a < c_1(Y) = 2n - 1$.

dim QH^{2a}(Y) =

$$\begin{cases}
n & \text{for } a \le 2n - 4 \text{ even} \\
n - 1 & \text{for } a = 2n - 2 \\
n - 1 & \text{for } a \le 2n - 3 \text{ odd.}
\end{cases}$$

We work modulo the ideal (q-1). This is enough since we can recover the powers of q by considering the degrees. An easy check gives that $E_h^X : A_{2a}(X) \to A_{2a+2}(X)$ is of (maximal) rank n-1. Corollary 4.1.6 implies that the same holds for E_h^Y . In particular Ker $E_{h_{-1}}^Y = \langle K_{4n-2}, K_{4n-2}, \cdots, K_{8n-10} \rangle$ for some $K_a \in A_a(Y) \cap \operatorname{Ker} E_h^Y$. Furthermore $\operatorname{Im} E_h^Y$ is a complement of this space. This implies that $K_{4n-2} = \lambda + v$ with $v \in \operatorname{Im} E_h^Y$ and since $\operatorname{Ker} E_h^Y \cap \operatorname{Im} E_h^Y = 0$, we have $K_{4n-2}^N = \lambda^{N-1} K_{4n-2} \neq 0$ and K_{4n-2} is not nilpotent. We claim that modulo scalar we have $K_{4n+2}^i = K_{4i}$ for all $i \in [1, n-2]$. Let $\sigma_{(1,1)} \in H^4(X, \mathbb{Z})$ be the Schubert class defined by the partition (1,1) (this Schubert class is also the top Chern class of the tautological subbundle of X). It is easy to check that $\sigma_{(1,1)}^{n-2} = \sigma_{(n-2,n-2)}$ this last class is the Schubert class associated to the partition (n-2, n-2). Indeed for degree reasons, this product is a classical cohomological product and the result follows from Littelwood-Richardson's rule. We get $h \cup \sigma_{(n-2,n-2)} \neq 0$. This implies that for $j: Y \to X$ the inclusion is rule. We get m = 0 or $(n-2, n-2) \neq 0$. This implies that for $j: Y \to X$ the inclusion, we have $j^* \sigma_{(1,1)}^{n-2} = j^* \sigma_{(n-2,n-2)}$ and that $j^* \sigma_{(1,1)}^{n-2} \cup h \neq 0$. In particular $j^* \sigma_{(1,1)}^{n-2} \notin \operatorname{Ker} E_h^Y$ thus $j^* \sigma_{(1,1)} \notin \operatorname{Ker} E_h^Y$ and we may write $j^* \sigma_{(1,1)}^{n-2} = \mu K_{8n-10} + w$ and $j^* \sigma_{(1,1)} = \lambda K_{4n+2} + v$ for some $\lambda, \mu \in \mathbb{R} \setminus \{0\}$ and $v, w \in \text{Im}E_h^Y$. This implies $\mu K_{8n-10} + w = j^* \sigma_{(1,1)}^{n-2} = \lambda^{n-2} K_{4n+2} + v^{n-2}$ and thus $\mu K_{8n-10} = \lambda^{n-2} K_{4n+2}^{n-2}$ proving the claim. Now $K_{4n+2}^{n-1} \in A_{4n-4}(Y) \subset \text{Im} E_h^Y$ (recall that we have a $\mathbb{Z}/2c_1(Y)\mathbb{Z} = \mathbb{Z}/(4n-2)\mathbb{Z}$ -grading) thus $K_{4n+2}^{n-1} = 0$ proving the result.

For technical reasons we have to distinguish the cases n = 3 and $n \ge 4$ in the proof of the next result.

Theorem 5.2.2. Let Y = IG(2, 2n). Then BQH(Y) is semisimple.

Proof. We will use the notation of [6] to index simple roots of the root system of type C_n . For classes $\sigma, \sigma' \in H(Y)$ we will write

$$\sigma \star_{\tau} \sigma' = \sigma \star_0 \sigma' + t(\sigma \star_1 \sigma') + O(t^2).$$

We start with the case n = 3. Let $\tau = \text{pt} = \sigma_{-\theta}$ where θ is the highest short root. We prove that $\text{BQH}_{\tau}(Y)$ (with notation as in Subsection 1.5) is semisimple.

Let $C \in BQH_{\tau}(Y)$ be a nilpotent element of order 2. If $C \neq 0$, up to dividing with the parameter t (see Subsection 1.5), we may assume that C is of the form

$$C = C_0 + tC_1 + O(t^2)$$
 with $C_0 \neq 0$

We have $0 = C \star_{\tau} C = C_0 \star_0 C_0 + O(t)$ thus $C_0 \in R(Y)$. Up to rescaling, we may assume that $C_0 = q\sigma_{\alpha_2+\alpha_3} - q\sigma_{\alpha_2+\alpha_3} + \sigma_{-\theta} \in \mathrm{QH}^{14}(Y)$. On the other hand, the product $h \star_{\tau} C$ has to be nilpotent. But we have

$$h \star_{\tau} C = h \star_0 C_0 + t(h \star_0 C_1 + h \star_1 C_0) + O(t^2) = t(h \star_0 C_1 + h \star_1 C_0) + O(t^2).$$

In particular $h \star_0 C_1 + h \star_1 C_0$ is nilpotent. Since $h \star_1 C_0$ is of degree 28 and since there is no nilpotent element of degree 28 in $\operatorname{QH}(Y)$, we must have $h \star_0 C_1 + h \star_1 C_0 =$ 0. This is possible since E_h^Y is surjective on degree 28. The element C_1 is thus uniquely determined by $h \star_0 C_1 = -h \star_1 C_0$. Note that $C_1 \in \operatorname{QH}^{26}(Y) \subset \operatorname{Im} E_h^Y$ and $C_0 \in \operatorname{Ker} E_h^Y$ thus $C_0 \star_0 C_1 = 0$.

We now remark the following equality $C_0 = 3\sigma_{-\theta} + (q\sigma_{\alpha_2+\alpha_3} - q\sigma_{\alpha_2+\alpha_3} - 2\sigma_{-\theta})$ where the second term is in the image of E_h^Y . Thus there exists $D_0 \in \mathrm{QH}^{12}(Y)$ with $h \star_0 D_0 = q\sigma_{\alpha_2+\alpha_3} - q\sigma_{\alpha_2+\alpha_3} - 2\sigma_{-\theta}$. An easy computation gives

$$D_0 = q\sigma_{\alpha_1 + \alpha_2 + \alpha_3} - 2\sigma_{-\alpha_1 - \alpha_2 - \alpha_3}.$$

We have $h \star_{\tau} D_0 = (q\sigma_{\alpha_2+\alpha_3} - q\sigma_{\alpha_2+\alpha_3} - 2\sigma_{-\theta}) + t(h \star_1 D_0) + O(t^2)$ with $h \star_1 D_0 \in$ QH²⁶(Y). Since E_h^Y is surjective on degree 26, there exists $D_1 \in$ QH²⁴(Y) with $h \star_0 D_1 = -h \star_1 D_0$. Setting $D = D_0 + tD_1$ we get $h \star_{\tau} D = q\sigma_{\alpha_2+\alpha_3} - q\sigma_{\alpha_2+\alpha_3} - 2\sigma_{-\theta} + O(t^2)$. This altogether gives $C_0 = 3\text{pt} + h \star_{\tau} D + O(t^2)$ and

$$3pt = C - h \star_{\tau} D - tC_1 + O(t^2).$$

Computing the square gives (recall that $C \star_{\tau} C = 0$, $h \star_{\tau} C = O(t^2)$ and $C_0 \star_0 C_1 = 0$)

9pt
$$\star_{\tau}$$
 pt = $h \star_{\tau} D \star_{\tau} h \star_{\tau} D - 2tC \star_{\tau} C_{1} + 2th \star_{\tau} D \star_{\tau} C_{1} + O(t^{2})$
= $h \star_{\tau} D \star_{\tau} h \star_{\tau} D - 2tC_{0} \star_{0} C_{1} + 2th \star_{0} D \star_{0} C_{1} + O(t^{2})$
= $h \star_{\tau} D \star_{\tau} h \star_{\tau} D + 2h \star_{0} D \star_{0} C_{1} + O(t^{2})$

Note that we have

$$h \star_{\tau} D \star_{\tau} h \star_{\tau} D = h \star_{0} h \star_{0} D_{0} \star_{0} D_{0} +2th \star_{0} h \star_{0} D_{0} \star_{0} D_{1} + th \star_{0} h \star_{0} (D_{0} \star_{1} D_{0}) +th \star_{0} (h \star_{1} (D_{0} \star_{0} D_{0})) + th \star_{1} (h \star_{0} D_{0} \star_{0} D_{0}) + O(t^{2}) = th \star_{1} (h \star_{0} D_{0} \star_{0} D_{0}) + t \cdot \operatorname{Im} E_{h}^{Y} + O(t^{2}).$$

Finally we obtain $9pt \star_{\tau} pt = 9pt \star_0 pt + th \star_1 (h \star_0 D_0 \star_0 D_0) + t \cdot Im E_h^Y + O(t^2)$. Since D_0 is explicitely given and since the endomorphism $h \star_1 -$ is understood using $pt \star_0 -$ (see Subsection 1.5) we get

9pt
$$\star_1$$
 pt = $12q^4 + 3q^3\sigma_{-\alpha_1-\alpha_2} - 3q^3\sigma_{-\alpha_2-\alpha_3} + \text{Im}E_h^Y = 6q^4 + \text{Im}E_h^Y$.

22

On the other hand we compute directly the product $pt \star_{\tau} pt$. We have

$$\operatorname{pt} \star_1 \operatorname{pt} = \sum_{d \ge 0} \sum_{\alpha \in R_s} q^d I_d(\operatorname{pt}, \operatorname{pt}, \operatorname{pt}, \sigma_{-\alpha}) \sigma_{\alpha}.$$

The invariant $I_d(\text{pt}, \text{pt}, \text{pt}, \sigma_{-\alpha})$ vanishes unless $3 \deg \text{pt} + \deg \sigma_{-\alpha} = 2 \dim Y + 2dc_1(Y) + 2$ thus $\deg \sigma_{-\alpha} = 10d - 26$. Since $\deg \sigma_{-\alpha}$ is even and contained in [0, 14], the invariant $I_d(\text{pt}, \text{pt}, \text{pt}, \sigma_{-\alpha})$ vanishes unless d = 3 and $\deg \sigma_{-\alpha} = 4$ or d = 4 and $\deg \sigma_{-\alpha} = 14$. But one easily check that there is no degree 3 curve in Y = IG(2, 6) passing through 3 points in general position. Thus $I_3(\text{pt}, \text{pt}, \text{pt}, -) = 0$. The only non vanishing invariant is thus $I_4(\text{pt}, \text{pt}, \text{pt}, \text{pt})$ and we have

$$9 \text{pt} \star_1 \text{pt} = 9I_4(\text{pt}, \text{pt}, \text{pt}, \text{pt})q^4 = 6q^4 + \text{Im}E_b^Y.$$

Since $q^4 \notin \text{Im}E_h^Y$ this implies $9I_4(\text{pt}, \text{pt}, \text{pt}, \text{pt}) = 6$ which is impossible since $I_4(\text{pt}, \text{pt}, \text{pt}, \text{pt}, \text{pt}) \in \mathbb{Z}_{\geq 0}$.

Assume $n \ge 4$. Let $\tau = \sigma_{\theta - \alpha_1 - \alpha_2} \in \text{QH}^4(Y)$ where θ is the highest short root. We prove that $\text{BQH}_{\tau}(Y)$ (with notation as in Subsection 1.5) is semisimple.

Let $C \in BQH_{\tau}(Y)$ be a nilpotent element of order 2. If $C \neq 0$, up to dividing with the parameter t (see Subsection 1.5), we may assume that C is of the form

$$C = C_0 + tC_1 + O(t^2)$$
 with $C_0 \neq 0$.

We have $0 = C \star_{\tau} C = C_0 \star_0 C_0 + O(t)$ thus $C_0 \in R(Y)$. Up to rescaling and multiplying with the generator of R(Y), we may assume that C_0 is the unique element in R(Y) of degree 8n - 10 such that the coefficient of $\sigma_{-\theta}$ in C_0 is 1. The product $h \star_{\tau} C$ has to be nilpotent. But we have

$$h \star_{\tau} C = h \star_0 C_0 + t(h \star_0 C_1 + h \star_1 C_0) + O(t^2) = t(h \star_0 C_1 + h \star_1 C_0) + O(t^2).$$

In particular $h \star_0 C_1 + h \star_1 C_0$ is nilpotent. Since $h \star_1 C_0$ is of degree 8n - 6and since there is no nilpotent element of degree 8n - 6 in QH(Y), we must have $h \star_0 C_1 + h \star_1 C_0 = 0$. This is possible since E_h^Y is surjective on degree 8n - 6. The element C_1 is thus uniquely determined by $h \star_0 C_1 = -h \star_1 C_0$. Note that $C_1 \in QH^{8n-4}(Y) \subset ImE_h^Y$ and $C_0 \in \operatorname{Ker} E_h^Y$ thus $C_0 \star_0 C_1 = 0$ (one can check that $h \star_1 C_0 \neq 0$ but we do not need this since if $h \star_1 C_0 = 0$, then $h \star_0 C_1$ has to be nilpotent which implies $h \star_0 C_1 = 0$ thus $C_1 \in R(X)$ and $C_0 \star_0 C_1 = 0$).

We now remark the following equality $C_0 = n\sigma_{-\theta} + v$ with $v \in \operatorname{Im} E_h^Y$. Thus there exists $D_0 \in \operatorname{QH}^{8n-12}(Y)$ with $h \star_0 D_0 = v$. We have $h \star_\tau D_0 = v + t(h \star_1 D_0) + O(t^2)$ with $h \star_1 D_0 \in \operatorname{QH}^{8n-8}(Y)$. Since E_h^Y is surjective on degree 8n - 8, there exists $D_1 \in \operatorname{QH}^{8n-10}(Y)$ with $h \star_0 D_1 = -h \star_1 D_0$. Setting $D = D_0 + tD_1$ we get $h \star_\tau D = v + O(t^2)$. This altogether gives $C_0 = n \text{ pt} + h \star_\tau D + O(t^2)$ and

$$n \text{ pt} = C - h \star_{\tau} D - tC_1 + O(t^2).$$

Computing the square gives

$$n^2$$
 pt \star_{τ} pt $= n^2$ pt \star_0 pt $+ th \star_1 (h \star_0 D_0 \star_0 D_0) + t \cdot \text{Im}E_h^Y + O(t^2).$

Now $h \star_0 D_0 \star_0 D_0 = v \star_0 D_0 = (C_0 - n\sigma_{-\theta}) \star_0 D_0$ and since $D_0 \in \operatorname{QH}^{8n-12}(Y) \subset \operatorname{Im} E_h^Y$ we have $C_0 \star_0 D_0 = 0$ thus $h \star_0 D_0 \star_0 D_0 = -n$ pt $\star_0 D_0$. Let $A = \operatorname{pt} \star_0 D_0$. To compute A we first compute $h \star_0 A = \operatorname{pt} \star_0 h \star_0 D_0 = \operatorname{pt} \star_0 (C_0 - n \text{ pt}) = -n$ pt \star_0 pt (since pt $\in \langle C_0, \operatorname{Im} E_h^Y \rangle$ and C_0 is orthogonal to both terms we have $C_0 \star_0 \operatorname{pt} = 0$). The Product pt $\star_0 \operatorname{pt} = q^2 \sigma$ with $\sigma = \sigma_{\alpha_1 + \alpha_2 + \alpha_3 - \theta}$ is easy to compute using for example the kernel and span technique presented in [9]. In particular

we get $h \star_0 A = -n q^2 \sigma$. Thus for $\sigma' = \sigma_{\alpha_1 + 2\alpha_2 + \alpha_3 - \theta}$ and K the unique element in Ker E_h^Y of degree $16n - 22 = \deg(A)$ with coefficient 1 on $q^2 \sigma'$, since $h \star_0 \sigma' = \sigma$, we have

$$A = \lambda K - n \ q^2 \sigma'$$

for some scalar λ . Let ℓ be the class of a line in Y. Note that the coefficient of ℓ in D_0 is 1-n. To compute λ , first note that $\operatorname{pt} \star_0 \operatorname{PD}(\sigma') = q^2 h$ (again obtained using the kernel and span technique). In particular, the only non vanishing invariant of the form $I_d(\operatorname{pt}, \operatorname{PD}(\sigma'), -)$ is the invariant $I_2(\operatorname{pt}, \operatorname{PD}(\sigma'), \ell) = 1$. This implies that the coefficient of $q^2\sigma'$ in $A = \operatorname{pt} \star_0 D_0$ is the coefficient of ℓ in D_0 and has value 1-n. We deduce that $\lambda = 1$ and

$$h \star_0 D_0 \star_0 D_0 = n^2 q^2 \sigma' - n K.$$

We now compute $h \star_1 (h \star_0 D_0 \star_0 D_0)$. We actually only need to compute modulo n^2 so that we only to consider $h \star_1 K$. An easy degree argument gives that there are only 2 Schubert classes appearing in K with non vanishing value under $h \star_1 -$: the classes $q^3\sigma_{a_1+\alpha_2+\alpha_3+\alpha_4-\theta}$ and $q^3\sigma_{\alpha_2+\alpha_3+\alpha_4+\alpha_5-\theta}$ (only the first one for n = 4). One then easily check using the kernel and span technique the following formula

$$h \star_1 K = q^3 \sigma_{\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5}$$
 (for $n = 4$ we have $h \star_1 K = \sigma_{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4}$).

Write $q^3\gamma = h \star_1 K$. The class γ is a Schubert class and is not contained in $\text{Im}E_h^Y$. Altogether working in the Schubert basis modulo n^2 and modulo $\text{Im}E_h^Y$ we get:

$$-nq^2\gamma \equiv n^2 \text{pt} \star_1 \text{pt} \equiv 0 \pmod{n^2} \text{ and } \text{Im}E_h^Y$$
.

A contradiction.

5.3. Cayley plane. In this subsection we consider the variety $Y = F_4/P_4$ obtained as hyperplane section of the Cayley plane $\mathbb{OP}^2 = E_6/P_6$. Since the arguments are very similar to the case Y = IG(2, 2n) and since the computer program [11] gives a complete description of the small quantum cohomology we shall only state the results and give a sketch of proof.

Lemma 5.3.1. Let Y be a linear section of $X = E_6/P_6$ of codimension 1. We have $\operatorname{Ker} E_h^Y = R(Y) = \langle K_8 \rangle$ for some element $K_8 \in \operatorname{QH}^8(Y)$.

Proof. Follows from the description of E_h^Y using short roots and the fact that the element $K_8 \in \operatorname{Ker} E_h^Y \cap \operatorname{QH}^8(Y)$ satisfies $K_8^2 \in A_5(Y) \subset E_h^Y$ and $K_8^2 \in \operatorname{Ker} E_h^Y$ thus $K_8^2 = 0$ since $\operatorname{Im} E_h^Y \cap \operatorname{Ker} E_h^Y = 0$.

Theorem 5.3.2. The big quantum cohomology BQH(Y) is semisimple.

Proof. For classes $\sigma, \sigma' \in H(Y)$ we will write

$$\sigma \star_{\tau} \sigma' = \sigma \star_0 \sigma' + t(\sigma \star_1 \sigma') + O(t^2).$$

Let $(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$ a system of simple roots of the root system of F_4 with α_3 and α_4 short. We have $\theta = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4$ (recall that θ is the highest short root). Let $\tau = \text{pt}$, we prove that $\text{BQH}_{\tau}(Y)$ is semisimple. Let $C \in \text{BQH}_{\tau}(Y)$ be nilpotent with $C \neq 0$ and $c \star_{\tau} C = 0$. We may assume that

$$C = C_0 + tC_1 + O(t^2)$$
 with $C_0 \neq 0$

We have $0 = C \star_{\tau} C = C_0 \star_0 C_0 + O(t)$ thus $C_0 \in R(Y)$ and modulo rescaling we may assume $C_0 = \sigma_{-\theta} - q\sigma_{\alpha} + q\sigma_{\beta} \in QH^{30}(Y)$ with $\alpha = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$ and $\beta = \alpha_2 + 2\alpha_3 + \alpha_4$. We have:

$$h \star_{\tau} C = h \star_0 C_0 + t(h \star_1 C_0 + h \star_0 C_1) + O(t^2) = t(h \star_1 C_0 + h \star_0 C_1) + O(t^2).$$

In particular since $h \star_{\tau} C$ is nilpotent, we get that $h \star_1 C_0 + h \star_0 C_1$ is nilpotent. Since $h \star_1 C_0 = \in \operatorname{QH}^{60}(Y) = q^2 \operatorname{QH}^{16}(Y)$ and since there is no nilpotent element in degree 60 we get $h \star_1 C_0 + h \star_0 C_1 = 0$. This is possible since E_h^Y is surjective on degree 60. The element C_1 is thus uniquely determined by $h \star_0 C_1 = -h \star_1 C_0$. Note that $C_1 \in \operatorname{QH}^{58}(Y) \subset \operatorname{Im} E_h^Y$ and $C_0 \in \operatorname{Ker} E_h^Y$ thus $C_0 \star_0 C_1 = 0$.

We now remark the following equality $C_0 = 3\sigma_{-\theta} + (q\sigma_\beta - q\sigma_\alpha - 2\sigma_{-\theta})$ where the second term is in the image of E_h^Y . Thus there exists $D_0 \in \text{QH}^{28}(Y)$ with $h \star_0 D_0 = q\sigma_\beta - q\sigma_\alpha - 2\sigma_{-\theta}$. An easy computation gives

$$D_0 = q\sigma_\gamma - 2\sigma_{-\delta}$$

with $\gamma = \alpha_1 + \alpha_2 + 2\alpha_3 + \alpha_4$ and $\delta = \alpha_1 + 2\alpha_2 + 3\alpha_3 + \alpha_4$. We have $h \star_{\tau} D_0 = (q\sigma_{\beta} - q\sigma_{\alpha} - 2\sigma_{-\theta}) + t(h \star_1 D_0) + O(t^2)$ with $h \star_1 D_0 \in \operatorname{QH}^{58}(Y)$. Since E_h^Y is surjective on degree 58, there exists $D_1 \in \operatorname{QH}^{56}(Y)$ with $h \star_0 D_1 = -h \star_1 D_0$. Setting $D = D_0 + tD_1$ we get $h \star_{\tau} D = q\sigma_{\beta} - q\sigma_{\alpha} - 2\sigma_{-\theta} + O(t^2)$. This altogether gives $C_0 = \operatorname{3pt} + h \star_{\tau} D + O(t^2)$ and

$$3pt = C - h \star_{\tau} D - tC_1 + O(t^2).$$

Computing the square gives as in the proof of Theorem 5.2.2

9pt
$$\star_{\tau}$$
 pt = pt \star_{0} pt + th \star_{1} (h $\star_{0} D_{0} \star_{0} D_{0}$) + t · Im E_{h}^{Y} + $O(t^{2})$.

Since D_0 is explicitly given and since the endomorphism $h \star_1 -$ is understood using pt $\star_0 -$ (see Subsection 1.5) we get

9pt
$$\star_1$$
 pt = $12q^4 + 12q^3\sigma_{-\alpha} + 6q^3\sigma_{-\beta} + \text{Im}E_h^Y = 6q^4 + \text{Im}E_h^Y$

On the other hand we compute directly the product $pt \star_{\tau} pt$. We have

$$\operatorname{pt} \star_1 \operatorname{pt} = \sum_{d \ge 0} \sum_{\zeta \in R_s} q^d I_d(\operatorname{pt}, \operatorname{pt}, \operatorname{pt}, \sigma_{-\zeta}) \sigma_{\zeta}.$$

The invariant $I_d(\text{pt}, \text{pt}, \text{pt}, \sigma_{-\zeta})$ vanishes unless $3 \deg \text{pt} + \deg \sigma_{-\zeta} = 2 \dim Y + 2dc_1(Y) + 2$ thus $\deg \sigma_{-\zeta} = 22d - 58$. Since $\deg \sigma_{-\zeta}$ is even and contained in [0, 30], the invariant $I_d(\text{pt}, \text{pt}, \text{pt}, \sigma_{-\zeta})$ vanishes unless d = 3 and $\deg \sigma_{-\zeta} = 8$ or d = 4 and $\deg \sigma_{-\zeta} = 30$. But one it was proved in [17, Proposition 2.13] that there is no degree 3 curve in Y passing through 3 points in general position. Thus $I_3(\text{pt}, \text{pt}, \text{pt}, -) = 0$. The only non vanishing invariant is thus $I_4(\text{pt}, \text{pt}, \text{pt}, \text{pt})$ and we have

$$9pt \star_1 pt = 9I_4(pt, pt, pt, pt)q^4 = 6q^4 + \operatorname{Im} E_h^Y.$$

Since $q^4 \notin \text{Im}E_h^Y$ this implies $9I_4(\text{pt}, \text{pt}, \text{pt}, \text{pt}) = 6$ which is impossible since $I_4(\text{pt}, \text{pt}, \text{pt}, \text{pt}, \text{pt}) \in \mathbb{Z}_{\geq 0}$.

References

- A. Bayer and Y.I. Manin, (Semi)simple exercises in quantum cohomology. The Fano Conference, 143–173, Univ. Torino, Turin, 2004.
- [2] A. Beauville, Quantum cohomology of complete intersections, Mat. Fiz. Anal. Geom. 2 (1995), no. 3-4, 384–398.
- [3] K. Behrend, Gromov-Witten invariants in algebraic geometry. Invent. Math. 127 (1997), no. 3, 601–617.
- [4] K. Behrend and B. Fantechi, The intrinsic normal cone. Invent. Math. 128 (1997), no. 1, 45–88.
- [5] K. Behrend and Y.I. Manin, Stacks of stable maps and Gromov-Witten invariants. Duke Math. J. 85 (1996), no. 1, 1–60.
- [6] N. Bourbaki, Groupes et algèbres de Lie. Hermann 1954.
- [7] A. S. Buch, P.-E. Chaput, L. C. Mihalcea, N. Perrin, Finiteness of cominuscule quantum K-theory. Annales de l'ÉNS, 46, (2013), no. 3, 477–494.
- [8] A. S. Buch, A. Kresch, and H. Tamvakis, Gromov-Witten invariants on Grassmannians, J. Amer. Math. Soc. 16 (2003), no. 4, 901–915 (electronic).
- [9] <u>Quantum Pieri rules for isotropic Grassmannians</u>. Invent. Math. 178 (2009), no. 2, 345–405.
- [10] A. S. Buch and L. C. Mihalcea, Quantum K-theory of Grassmannians, Duke Math. J. 156 (2011), no. 3, 501–538.
- [11] P.-E. Chaput, Computing in the quantum cohomology of some homogeneous spaces. Computer program available at http://www.iecn.u-nancy.fr/~chaput/quiver-demo.html
- [12] P.-E. Chaput, L. Manivel, and N. Perrin, Quantum cohomology of minuscule homogeneous spaces, Transform. Groups 13 (2008), no. 1, 47–89.
- [13] _____, Quantum cohomology of minuscule homogeneous spaces. II. Hidden symmetries. Int. Math. Res. Not. IMRN 2007, no. 22.
- [14] _____, Quantum cohomology of minuscule homogeneous spaces III : semi-simplicity and consequences, Canad. J. Math. 62 (2010), no. 6, 1246–1263.
- [15] _____, Affine symmetries of the equivariant quantum cohomology ring of rational homogeneous spaces. Math. Res. Lett. 16 (2009), no. 1, 7–21.
- [16] P.-E. Chaput and N. Perrin, On the quantum cohomology of adjoint varieties, Proc. Lond. Math. Soc. (3) 103 (2011), no. 2, 294–330.
- [17] _____, Rationality of some Gromov-Witten varieties and application to quantum K-theory. Commun. Contemp. Math. 13 (2011), no. 1, 67–90.
- [18] G. Ciolli, Computing the quantum cohomology of some Fano threefolds and its semisimplicity. Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) 7 (2004), no. 2, 511–517.
- [19] P. Deligne Cohomologie des intersections complètes. SGA 7, Exposé XI. LNM 340 Springer-Verlag (1973).
- [20] B. Dubrovin, Geometry and analytic theory of Frobenius manifolds, in: Proceedings of the International Congress of Mathematicians, Berlin, 1998, Doc. Math. Extra vol. II (1998) 315–326.
- [21] S. Galkin, M. Smirnov and A. Mellit, Dubrovin's conjecture for IG(2,6), preprint arXiv:1405.3857.
- [22] C. Hertling, Y.I. Manin and C. Teleman, An update on semisimple quantum cohomology and F-manifolds. Tr. Mat. Inst. Steklova 264 (2009), Mnogomernaya Algebraicheskaya Geometriya, 69–76; translation in Proc. Steklov Inst. Math. 264 (2009), no. 1, 62–69.
- [23] H. Iritani, Convergence of quantum cohomology by quantum Lefschetz. J. Reine Angew. Math. 610 (2007), 29–69.
- [24] M. Kontsevich and Y.I. Manin, Gromov-Witten classes, quantum cohomology, and enumerative geometry. Comm. Math. Phys. 164 (1994), no. 3, 525–562.
- [25] A.G. Kuznetsov, Hyperplane sections and derived categories. Izv. Math. 70 (2006), no. 3, 447–547.
- [26] <u>Exceptional collections for Grassmannians of isotropic lines.</u> Proc. Lond. Math. Soc.
 (3) 97 (2008), no. 1, 155–182.
- [27] J. M. Landsberg and L. Manivel, The projective geometry of Freudenthal's magic square. J. Algebra 239 (2001), no. 2, 477–512.
- [28] Y.-P. Lee, Quantum K-theory. I. Foundations. Duke Math. J. 121 (2004), no. 3, 389–424.

- [29] Y.I. Manin, Frobenius manifolds, quantum cohomology, and moduli spaces. American Mathematical Society Colloquium Publications, 47. American Mathematical Society, Providence, RI, 1999.
- [30] I. A. Mihai, Odd symplectic flag manifolds. Transform. Groups 12 (2007), no. 3, 573–599.
- [31] A. N. Parshin and I. R. Shafarevich, Algebraic geometry. V. Fano varieties. Encyclopaedia of Mathematical Sciences, 47. Springer-Verlag, Berlin, 1999.
- [32] B. Pasquier, On some smooth projective two-orbit varieties with Picard number 1. Math. Ann. 344 (2009), no. 4, 963–987.
- [33] B. Pasquier and N. Perrin, Local rigidity of quasi-regular varieties. Math. Z. 265 (2010), no. 3, 589–600.
- [34] C. Pech, Quantum cohomology of the odd symplectic Grassmannian of lines. J. Algebra 375 (2013), 188–215.
- [35] N. Perrin, Small codimension subvarieties in homogeneous spaces, Indag. Math. (N.S.) 20 (2009), no. 4, 557–581.
- [36] _____Small codimension smooth subvarieties in even-dimensional homogeneous spaces with Picard group Z, CRAS 345 (2007), no. 3, 155–160.
- [37] _____Small resolutions of minuscule Schubert varieties. Compos. Math. 143 (2007), no. 5, 1255–1312.
- [38] _____ The Gorenstein locus of minuscule Schubert varieties. Adv. Math. 220 (2009), no. 2, 505–522.
- [39] Y. Ruan and G. Tian, A mathematical theory of quantum cohomology. J. Differential Geom. 42 (1995), no. 2, 259–367.
- [40] C.S. Seshadri, Geometry of G/P. I. Theory of standard monomials for minuscule representations, C. P. Ramanujam – a tribute, pp. 207–239, Tata Inst. Fund. Res. Studies in Math., 8, Springer, Berlin-New York, 1978.
- [41] H. Thomas and A. Yong, A combinatorial rule for (co)minuscule Schubert calculus. Adv. Math. 222 (2009), no. 2, 596–620.
- [42] G. Tian and G. Xu, On the semi-simplicity of the quantum cohomology algebras of complete intersections, Math. Res. Lett. 4 (1997), no. 4, 481–488.

MATH. INSTITUT, HEINRICH-HEINE-UNIVERSITÄT, D-40204 DÜSSELDORF, GERMANY *E-mail address*: perrin@math.uni-duesseldorf.de