

Intersection numbers for subspace designs

Michael Kiermaier * Mario Osvin Pavčević †
University of Bayreuth *University of Zagreb*

September 3, 2018

Intersection numbers for subspace designs are introduced and q -analogs of the Mendelsohn and Köhler equations are given. As an application, we are able to determine the intersection structure of a putative q -analog of the Fano plane for any prime power q . It is shown that its existence implies the existence of a 2 - $(7, 3, q^4)_q$ subspace design. Furthermore, several simplified or alternative proofs concerning intersection numbers of ordinary block designs are discussed.

1 Introduction and preliminaries

1.1 History

The earliest reference for q -analogs of block designs (subspace designs) is [4]. However, the idea is older, since it is stated that “Several people have observed that the concept of a t -design can be generalised [...]”. They have also been mentioned in a more general context in [8]. An introduction can be found in [22, Day 4].

For $q = 2$, the first nontrivial subspace design with $t = 2$ has been constructed in [27] and generalized to arbitrary q in [23, 26]. The first nontrivial subspace design with $t = 3$ is found in [3].

*University of Bayreuth, Institute for Mathematics, 95440 Bayreuth, Germany

email: michael.kiermaier@uni-bayreuth.de

homepage: <http://www.mathe2.uni-bayreuth.de/michaelk/>

Research supported by ESF COST Action IC1104.

†University of Zagreb, Faculty of Electrical Engineering and Computing, Department of Applied Mathematics, Unska 3, 10000 Zagreb, Croatia

email: mario.pavcevic@fer.hr

Research supported by ESF COST Action IC1104.

In [21, Th. 1.2] it has been shown that for fixed parameters t, v and k and λ sufficiently large, each admissible parameter set t - (v, k, λ) is realizable as a subspace design with possibly repeated blocks. In [9] it has been proven that nontrivial simple subspace designs exist for any value of t .

Quite recently [2], a 2-analog of the Steiner triple system STS(13) has been found computationally, by applying the Kramer-Mesner method described in [3]. This discovery is a significant breakthrough, since it is the very first nontrivial q -Steiner system with $t > 1$ and refutes the earlier conjecture that no such q -Steiner system exists.

1.2 Gaussian binomial coefficients

We define the q -analog of a non-negative integer n as

$$[n] = [n]_q = \frac{q^n - 1}{q - 1} \in \mathbb{Z}[q]$$

and the q -factorial of n as

$$[n]! = [n]_q! = \prod_{i=1}^n [i] \in \mathbb{Z}[q].$$

The notion q -analog stems from the fact that the evaluation for $q = 1$ gives $[n]_1 = n$ and $[n]_1! = n!$. Using this notation, for $k \in \mathbb{Z}$ and $n \in \mathbb{N}$ the *Gaussian binomial coefficient* is the $\mathbb{Z}[q]$ -polynomial

$$\begin{bmatrix} n \\ k \end{bmatrix} = \begin{bmatrix} n \\ k \end{bmatrix}_q = \begin{cases} \frac{[n]!}{[k]! [n-k]!} & \text{if } k \in \{0, \dots, n\}, \\ 0 & \text{otherwise.} \end{cases}$$

Its evaluation for $q = 1$ gives the binomial coefficient $\binom{n}{k}$. For that reason, the Gaussian binomial coefficient is known as q -analog of the binomial coefficient. Many identities for binomial coefficients have q -analogs for the Gaussian binomial coefficients. As an example, we mention

$$\begin{bmatrix} n \\ k \end{bmatrix} = \begin{bmatrix} n \\ n-k \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} n \\ h \end{bmatrix} \begin{bmatrix} n-h \\ k \end{bmatrix} = \begin{bmatrix} n \\ k \end{bmatrix} \begin{bmatrix} n-k \\ h \end{bmatrix},$$

for $n \geq 1$ the q -Pascal triangle identities

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q + q^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_q = q^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q + \begin{bmatrix} n-1 \\ k \end{bmatrix}_q$$

and the identity (see for example [8, Eq. (3)])

$$\sum_{i=a}^b (-1)^{i-a} q^{\binom{i-a}{2}} \begin{bmatrix} b \\ i \end{bmatrix}_q \begin{bmatrix} i \\ a \end{bmatrix}_q = \begin{cases} 1 & \text{if } a = b, \\ 0 & \text{else.} \end{cases} \quad (1)$$

Another one is provided in the following

Lemma 1.1 *Let $n, k \in \mathbb{Z}$ with $n \geq 1$. Then*

$$\sum_{i=0}^k (-1)^i q^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix}_q = (-1)^k q^{\binom{k+1}{2}} \begin{bmatrix} n-1 \\ k \end{bmatrix}_q.$$

Proof. By the q -Pascal triangle identity,

$$\begin{aligned} \sum_{i=0}^k (-1)^i q^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix}_q &= \sum_{i=0}^k (-1)^i q^{\binom{i}{2}} \cdot q^i \begin{bmatrix} n-1 \\ i \end{bmatrix}_q + \sum_{i=0}^k (-1)^i q^{\binom{i}{2}} \cdot \begin{bmatrix} n-1 \\ i-1 \end{bmatrix}_q \\ &= \sum_{i=0}^k (-1)^i q^{\binom{i+1}{2}} \begin{bmatrix} n-1 \\ i \end{bmatrix}_q - \sum_{i=-1}^{k-1} (-1)^i q^{\binom{i+1}{2}} \begin{bmatrix} n-1 \\ i \end{bmatrix}_q \\ &= (-1)^k q^{\binom{k+1}{2}} \begin{bmatrix} n-1 \\ k \end{bmatrix}_q. \quad \square \end{aligned}$$

Remark 1.2 According to [10], q -analogs of non-negative integers were introduced in [12] and their binomial coefficients in [30]. For a deeper discussion of the Gaussian binomial coefficients, see [19, 10, 11, 5].

1.3 q -analogs of combinatorial structures

The set of k -element subsets (k -subsets) of a set V will be denoted by $\binom{V}{k}$ and the set of all k -dimensional subspaces (k -subspaces) of an \mathbb{F}_q -vector space will be denoted by $\begin{bmatrix} V \\ k \end{bmatrix}$ or $\begin{bmatrix} V \\ k \end{bmatrix}_q$. The latter is known as the *Graßmannian*. This notation is chosen for the fact that the size of $\binom{V}{k}$ equals the binomial coefficient $\binom{\#V}{k}$, and the size of $\begin{bmatrix} V \\ k \end{bmatrix}_q$ equals the Gaussian binomial coefficient $\begin{bmatrix} \dim(V) \\ k \end{bmatrix}_q$. There are good reasons to interpret the subspace lattice $\mathcal{L}(V)$ of a v -dimensional vector space V as the q -analog of the subset lattice of a v -element set V , which corresponds to $q = 1$ [5].

Many combinatorial areas, like design theory and coding theory, are based on the subset lattice of a v -element set V . Replacing the set-theoretic notions by their vector space counterparts gives rise to the study of their q -analogs, which are based on the subspace lattice of a v -dimensional vector space V . An important part of these theories is the investigation of results in the set-theoretic case for their applicability in the q -analog case. For example, in [14] q -analogs of derived and residual designs are studied. In this article, we will give a q -analog of the theory of intersection numbers of designs.

In the following, q -analogs of several well-known definitions and statements on ordinary block designs are given (Def. 1.3, Fact 1.4, Fact 1.5, Fact 1.7, Def. 2.1, Th. 2.4, Th. 2.6). This means that one gets back the original definition or statement if q is set to 1 and all vector space notions are replaced by their set-theoretic counterparts.

1.4 Subspace designs

Definition 1.3 Let q be a prime power, V an \mathbb{F}_q -vector space of finite dimension v and t, k, λ be non-negative integers. A set D of k -subspaces (blocks) of V is called a t - $(v, k, \lambda)_q$ (subspace) design if each t -subspace of V is contained in exactly λ blocks of D .

By the above discussion, an ordinary block design can be seen as the case $q = 1$ of a subspace design. For all $t \in \{0, \dots, k\}$, the full Grassmannian $\begin{bmatrix} V \\ k \end{bmatrix}$ forms the *trivial* t - $(v, k, \begin{bmatrix} v-t \\ k-t \end{bmatrix}_q)_q$ subspace design. It is clear that for any t - $(v, k, \lambda)_q$ subspace design D , the *complementary* design $\begin{bmatrix} V \\ k \end{bmatrix} \setminus D$ is a t - $(v, k, \begin{bmatrix} v-t \\ k-t \end{bmatrix}_q - \lambda)_q$ subspace design.

Of particular interest is the case $\lambda = 1$, where D is called a *Steiner system*. For $t = 1$, a 1 - $(v, k, 1)_q$ Steiner system is the same as a spread of $(k - 1)$ -flats in the projective geometry $\text{PG}(v - 1, q)$, which exists if and only if k divides v . The only known nontrivial q -analog of a Steiner system with $t \geq 2$ has the parameters 2 - $(13, 3, 1)_2$ [2].

By the fundamental theorem of projective geometry, the automorphism group of the lattice $\mathcal{L}(V)$ is given by the projective semilinear group $\text{P}\Gamma\text{L}(V)$ with its natural action on $\mathcal{L}(V)$. The automorphism group $\text{Aut}(D)$ of a subspace design D is defined as the stabilizer of D under the induced action of $\text{P}\Gamma\text{L}(V)$ on the power set of $\mathcal{L}(V)$.

The following fact is the q -analog of a well-known property of block designs:

Fact 1.4 ([24, Lemma 4.1(1)]) Let D be t - $(v, k, \lambda)_q$ subspace design. For each $i \in \{0, \dots, t\}$, D is an i - $(v, k, \lambda_i)_q$ subspace design with

$$\lambda_i = \frac{\begin{bmatrix} v-i \\ t-i \end{bmatrix}_q}{\begin{bmatrix} k-i \\ t-i \end{bmatrix}_q} \cdot \lambda = \frac{\begin{bmatrix} v-i \\ k-i \end{bmatrix}_q}{\begin{bmatrix} v-t \\ k-t \end{bmatrix}_q} \cdot \lambda.$$

In particular, the number of blocks is given by $\#D = \lambda_0$.

As a result, the existence of a t - $(v, k, \lambda)_q$ design implies the *integrality conditions* $\lambda_i \in \mathbb{Z}$ for all $i \in \{0, \dots, t\}$. Without requiring the actual existence of a corresponding design, any parameter set t - $(v, k, \lambda)_q$ fulfilling the integrality conditions will be called *admissible*.

The following fact describes a refinement of the numbers λ_i .

Fact 1.5 ([24, Lemma 2.1], [22, Lemma 4.2]) Let D be a t - $(v, k, \lambda)_q$ subspace design and i, j non-negative integers with $i + j \leq t$. Let $I \in \begin{bmatrix} V \\ i \end{bmatrix}$ and $J \in \begin{bmatrix} V \\ j \end{bmatrix}$ with $I \cap J = \{\mathbf{0}\}$. The number

$$\lambda_{i,j} = \{B \in D \mid I \leq B \text{ and } B \cap J = \{\mathbf{0}\}\}$$

is independent of the choice of I and J . They are determined by the recurrence relation

$$\lambda_{i,0} = \lambda_i \quad \text{and} \quad \lambda_{i,j+1} = \lambda_{i,j} - q^j \lambda_{i+1,j}.$$

In closed form,

$$\lambda_{i,j} = q^{j(k-i)} \frac{\begin{bmatrix} v-i-j \\ k-i \end{bmatrix}_q}{\begin{bmatrix} v-t \\ k-t \end{bmatrix}_q} \cdot \lambda.$$

Remark 1.6 (i) Fact 1.5 can also be found in [25, Lemma 2.2 and Lemma 2.6]. However, the exponents of q given there are not correct.

(ii) For ordinary block designs, the numbers $\lambda_{i,j}$ have been introduced in [20].

Fixing some non-singular bilinear form β on V , the *dual subspace* of a subspace $W \in \mathcal{L}(V)$ is defined as

$$W^\perp = \{x \in V \mid \beta(x, y) = 0 \text{ for all } y \in W\}.$$

Now for a t - $(v, k, \lambda)_q$ subspace design D , its *dual* subspace design is defined as

$$D^\perp = \{B^\perp \mid B \in \mathcal{B}\}.$$

Up to equivalence, this definition does not depend on the choice of β . The dual subspace design is the q -analog of the supplementary block design.

Fact 1.7 ([24, Lemma 4.2]) *Let D be a t - $(v, k, \lambda)_q$ subspace design. Then D^\perp is a subspace design with the parameters*

$$t - \left(v, v - k, \frac{\begin{bmatrix} v-t \\ k \end{bmatrix}_q}{\begin{bmatrix} v-t \\ k-t \end{bmatrix}_q} \cdot \lambda \right)_q.$$

2 Intersection numbers

2.1 Plain intersection numbers

Definition 2.1 *Let D be a t - $(v, k, \lambda)_q$ subspace design. For any subspace S of V and $i \in \{0, \dots, k\}$, we define the i -th intersection number of S in D as*

$$\alpha_i(S) = \#\{B \in D \mid \dim(B \cap S) = i\}.$$

If the set S is clear from the context, we use the abbreviation $\alpha_i = \alpha_i(S)$. Furthermore, the $(k+1)$ -tuple $\alpha(S) = (\alpha_0(S), \alpha_1(S), \dots, \alpha_k(S))$ will be called the intersection vector of S in D .

The intersection numbers are a q -analog of the intersection numbers defined in [17] for blocks S and independently as “ i -Treffers” for general sets S in [18].

First, we describe the relation to the intersection numbers α_i^\perp of the dual design D^\perp :

Lemma 2.2 *Let D be a t - $(v, k, \lambda)_q$ subspace design, $S \in \mathcal{L}(V)$, $s = \dim(S)$ and $i \in \{0, \dots, k\}$.*

(a) *For $i > s$ or $k - i > v - s$ we have*

$$\alpha_i(S) = 0.$$

(b) For $i \leq s$ and $k - i \leq v - s$ we have

$$\alpha_i(S) = \alpha_{(v-s)-(k-i)}^\perp(S^\perp).$$

Proof. Part (a) is a direct consequence of the dimension formula. For part (b), note that for all blocks $B \in D$

$$\begin{aligned} \dim(B \cap S) = i &\iff \dim(B + S) = k + s - i \\ &\iff \dim(B^\perp \cap S^\perp) = v - (k + s - i). \end{aligned} \quad \square$$

In the range where the dimension or the codimension of S in V is at most t , the intersection numbers are closely related to the numbers $\lambda_{i,j}$:

Lemma 2.3 *Let D be a t - $(v, k, \lambda)_q$ subspace design, S a subspace of V of dimension*

$$s = \dim(S) \in \{0, \dots, t\} \cup \{v - t, \dots, v\}$$

and $i \in \{0, \dots, k\}$. The intersection vector $\alpha(S)$ is uniquely determined by

$$\alpha_i(S) = q^{(s-i)(k-i)} \frac{\begin{bmatrix} s \\ i \end{bmatrix}_q \begin{bmatrix} v-s \\ k-i \end{bmatrix}_q}{\begin{bmatrix} v-t \\ k-t \end{bmatrix}_q} \cdot \lambda. \quad (2)$$

For $i \leq s \leq t$ we have

$$\alpha_i(S) = \begin{bmatrix} s \\ i \end{bmatrix}_q \cdot \lambda_{i,s-i} \quad (3)$$

and for $k - i \leq v - s \leq t$ we have

$$\alpha_i(S) = q^{sk-iv} \begin{bmatrix} v-s \\ k-i \end{bmatrix}_q \cdot \lambda_{k-i, (v-s)-(k-i)}. \quad (4)$$

Proof. From Lemma 2.2(a), $\alpha_i(S) = 0$ for $i > s$ or $k - i > v - s$, in agreement with equation (2). So we may assume $i \leq s$ and $k - i \leq v - s$.

Case 1 We first consider the case $s \leq t$. We count the set

$$X = \left\{ (B, I) \in D \times \begin{bmatrix} S \\ i \end{bmatrix}_q \mid B \cap S = I \right\}$$

in two ways.

There are $\alpha_i(S)$ blocks B with $\dim(B \cap S) = i$, each one uniquely determining $I = B \cap S$. This shows that $\#X$ equals the left hand side of equation (3).

On the other hand, there are $\begin{bmatrix} s \\ i \end{bmatrix}_q$ ways to select the subspace I of S . For fixed I , let J be a complement of I in S . Let $B \in D$ with $I \leq B$. If $B \cap S \neq I$, then $\dim(B \cap S) > i$, and because of $\dim(J) + \dim(B \cap S) > (s - i) + i = s$, we get that $\dim(J \cap B) > 1$. Hence $B \cap S = I$ is equivalent to $I \leq B$ and $J \cap B = \{\mathbf{0}\}$. So the number of blocks intersecting S in I is $\lambda_{\dim(I), \dim(J)} = \lambda_{i, s-i}$, showing that $\#X$ equals the right hand side of equation (3).

So equation (3) is shown, and replacing $\lambda_{i, s-i}$ with the formula given in Fact 1.5 yields formula (2).

Case 2 Now assume that $v - s \leq t$. By Fact 1.7, the dual design D^\perp has the parameters $t^\perp = t$, $v^\perp = v$, $k^\perp = v - k$ and $\lambda^\perp = \begin{bmatrix} v-t \\ k \end{bmatrix}_q / \begin{bmatrix} v-t \\ k-t \end{bmatrix}_q \cdot \lambda$. We further define $s^\perp = v - s$ and $i^\perp = (v - s) - (k - i)$. Now

$$\alpha_i(S) = \alpha_{i^\perp}^\perp(S^\perp) = q^{(s^\perp - i^\perp)(k^\perp - i^\perp)} \frac{\begin{bmatrix} s^\perp \\ i^\perp \end{bmatrix}_q \begin{bmatrix} v^\perp - s^\perp \\ k^\perp - i^\perp \end{bmatrix}_q}{\begin{bmatrix} v^\perp - t^\perp \\ k^\perp - t^\perp \end{bmatrix}_q} \cdot \lambda^\perp,$$

where the first equality is Lemma 2.2(b), and the second equality comes from applying equation (2) (because of $\dim(S^\perp) = s^\perp \leq t^\perp$ we are in case 1 that we have already shown). Plugging in the above defined expressions, this expression indeed simplifies to the right hand side of (2). Finally equation (4) can be verified using the formula from Fact 1.5. \square

Theorem 2.4 (q -analog of the Mendelsohn equations [17, Th. 1]) *Let D be a t - $(v, k, \lambda)_q$ subspace design, S a subspace of V and $s = \dim(S)$. For $i \in \{0, \dots, t\}$ we have the following equation on the intersection numbers of S in D :*

$$\sum_{j=i}^s \begin{bmatrix} j \\ i \end{bmatrix}_q \alpha_j = \begin{bmatrix} s \\ i \end{bmatrix}_q \lambda_i \quad (5)$$

Proof. We count the set

$$X = \left\{ (I, B) \in \begin{bmatrix} V \\ i \end{bmatrix}_q \times D \mid I \leq B \cap S \right\}$$

in two ways.

There are $\begin{bmatrix} s \\ i \end{bmatrix}_q$ possibilities for the choice of $I \in \begin{bmatrix} S \\ i \end{bmatrix}$. By Fact 1.4, there are λ_i blocks B such that $I \leq B$, which shows that $\#X$ equals the right hand side of equation (5).

Fixing a block B , the number of i -subspaces I of $B \cap S$ is $\begin{bmatrix} \dim(B \cap S) \\ i \end{bmatrix}_q$. Summing over the possibilities for $j = \dim(B \cap S)$, we see that $\#X$ also equals the left hand side of equation (5). \square

Remark 2.5 In [17, Th. 1], S was required to be a block. For general S , the equations were given independently in [18, Satz 2].

Theorem 2.6 (q -analog of the Köhler equations [15, Satz 1]) *Let D be a t - $(v, k, \lambda)_q$ subspace design, S a subspace of V and $s = \dim(S)$. For $i \in \{0, \dots, t\}$, a parametrization of the intersection number α_i by $\alpha_{t+1}, \dots, \alpha_k$ is given by*

$$\alpha_i = \begin{bmatrix} s \\ i \end{bmatrix}_q \sum_{j=i}^t (-1)^{j-i} q^{\binom{j-i}{2}} \begin{bmatrix} s-i \\ j-i \end{bmatrix}_q \lambda_j \\ + (-1)^{t+1-i} q^{\binom{t+1-i}{2}} \sum_{j=t+1}^k \begin{bmatrix} j \\ i \end{bmatrix}_q \begin{bmatrix} j-i-1 \\ t-i \end{bmatrix}_q \alpha_j.$$

Proof. The Mendelsohn equations (Th. 2.4) can be interpreted as a system of linear equations on the intersection vector of S in D :

$$\left(\begin{array}{cccc|ccc} \begin{bmatrix} 0 \\ 0 \end{bmatrix}_q & \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}_q & \begin{bmatrix} 2 \\ 0 \\ 1 \\ 2 \end{bmatrix}_q & \cdots & \begin{bmatrix} t \\ 0 \\ 1 \\ 2 \\ \vdots \\ t \end{bmatrix}_q & \begin{bmatrix} t+1 \\ 0 \\ 1 \\ 2 \\ \vdots \\ t \end{bmatrix}_q & \cdots & \begin{bmatrix} k \\ 0 \\ 1 \\ 2 \\ \vdots \\ t \end{bmatrix}_q \\ \alpha_0 \\ \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_k \end{array} \right) = \begin{pmatrix} \begin{bmatrix} s \\ 0 \\ 1 \\ 2 \\ \vdots \\ t \end{bmatrix}_q \lambda_0 \\ \begin{bmatrix} s \\ 1 \\ 2 \\ \vdots \\ t \end{bmatrix}_q \lambda_1 \\ \begin{bmatrix} s \\ 2 \\ \vdots \\ t \end{bmatrix}_q \lambda_2 \\ \vdots \\ \begin{bmatrix} s \\ t \end{bmatrix}_q \lambda_t \end{pmatrix}$$

This equation system has the form

$$(P_q | A) \cdot \mathbf{x} = \mathbf{b} \quad (6)$$

where

$$\begin{aligned} P_q &= \left(\begin{bmatrix} j \\ i \end{bmatrix}_q \right)_{i,j \in \{0, \dots, t\}} \in \mathbb{Z}^{(t+1) \times (t+1)}, \\ A &= \left(\begin{bmatrix} j \\ i \end{bmatrix}_q \right)_{i \in \{0, \dots, t\}, j \in \{t+1, \dots, k\}} \in \mathbb{Z}^{(t+1) \times (k-t)} \quad \text{and} \\ \mathbf{b} &= \left(\begin{bmatrix} s \\ i \end{bmatrix}_q \lambda_i \right)_{i \in \{0, \dots, t\}} \in \mathbb{Z}^{t+1}. \end{aligned}$$

The matrix P_q is known as the *upper triangular q -Pascal matrix*. By equation (1), P_q is invertible with the inverse

$$P_q^{-1} = \left((-1)^{j-i} q^{\binom{j-i}{2}} \begin{bmatrix} j \\ i \end{bmatrix}_q \right)_{i,j \in \{0, \dots, t\}}.$$

After left multiplication by P_q^{-1} , equation (6) is equivalent to

$$(I | P_q^{-1}A) \cdot \mathbf{x} = P_q^{-1}\mathbf{b}. \quad (7)$$

Numbering the columns of A with $t+1, \dots, k$, the entry in the i -th row and the j -th column of $P_q^{-1}A$ is

$$\begin{aligned} (P_q^{-1}A)_{i,j} &= \sum_{\nu=0}^t (-1)^{\nu-i} q^{\binom{\nu-i}{2}} \begin{bmatrix} \nu \\ i \end{bmatrix}_q \begin{bmatrix} j \\ \nu \end{bmatrix}_q \\ &= \sum_{\nu=i}^t (-1)^{\nu-i} q^{\binom{\nu-i}{2}} \begin{bmatrix} j \\ i \end{bmatrix}_q \begin{bmatrix} j-i \\ \nu-i \end{bmatrix}_q \\ &= \begin{bmatrix} j \\ i \end{bmatrix}_q \sum_{\nu=0}^{t-i} (-1)^{\nu} q^{\binom{\nu}{2}} \begin{bmatrix} j-i \\ \nu \end{bmatrix}_q \\ &= \begin{bmatrix} j \\ i \end{bmatrix}_q (-1)^{t-i} q^{\binom{t-i+1}{2}} \begin{bmatrix} j-i-1 \\ t-i \end{bmatrix}_q, \end{aligned}$$

where Lemma 1.1 was used in the last step. The i -th entry of $P_q^{-1}\mathbf{b}$ is

$$\begin{aligned} (P_q^{-1}\mathbf{b})_i &= \sum_{j=0}^t (-1)^{j-i} q^{\binom{j-i}{2}} \begin{bmatrix} j \\ i \end{bmatrix}_q \begin{bmatrix} s \\ j \end{bmatrix}_q \lambda_j \\ &= \begin{bmatrix} s \\ i \end{bmatrix}_q \sum_{j=i}^t (-1)^{j-i} q^{\binom{j-i}{2}} \begin{bmatrix} s-i \\ j-i \end{bmatrix}_q \lambda_j. \end{aligned}$$

Plugging these expressions into equation (7), its rows evaluate to the Köhler equations. \square

Remark 2.7 (a) For ordinary block designs, Theorem 2.6 was originally shown in [15] in a lengthy induction proof. The main result of the article [29] was a simplified proof based on the notion of “vectorproduct”. In a slightly more general context, another induction proof as well as a proof based on the principle of inclusion and exclusion was given in [28].

(b) Our proof can be interpreted as transforming the linear system of Mendelsohn equations to row reduced echelon form by Gauss reduction. Since this method is directly applicable also to block designs, it provides a short and systematic proof for the original Köhler equations.

2.2 High order intersection numbers

For block designs, “high order” versions of the numbers $\lambda_{i,j}$ [28] and the intersection numbers α_i [17] (see also [28, 1]) have been introduced. For that matter, some positive integer ℓ is fixed, and in the definitions the block B is replaced by the intersection of an ℓ -tuple of blocks.

The same is possible in our q -analog situation: For a t - $(v, k, \lambda)_q$ subspace design D , non-negative integers i and j with $i + j \leq t$ and subspaces $I \in \begin{bmatrix} V \\ i \end{bmatrix}$ and $J \in \begin{bmatrix} V \\ j \end{bmatrix}$ with $I \cap J = \{\mathbf{0}\}$, the number

$$\lambda_{i,j}^{(\ell)} = \# \left\{ \mathcal{B} \in \binom{D}{\ell} \mid I \leq \bigcap \mathcal{B} \text{ and } \bigcap \mathcal{B} \cap J = \{\mathbf{0}\} \right\}$$

does not depend on the choice of I and J . For $S \in \mathcal{L}(V)$ and $i \in \{0, \dots, k\}$ the i -th *high order intersection number* of S in D is defined as

$$\alpha_i^{(\ell)}(S) = \# \left\{ \mathcal{B} \in \binom{D}{\ell} \mid \dim \left(\bigcap \mathcal{B} \cap S \right) = i \right\}.$$

Clearly, $\lambda_{i,j} = \lambda_{i,j}^{(1)}$ and $\alpha_i(S) = \alpha_i^{(1)}(S)$, so the high order versions generalize the basic versions considered so far.

Replacing α_i by $\alpha_i^{(\ell)}$, $\lambda_{i,j}$ by $\lambda_{i,j}^{(\ell)}$ and λ_i by $\binom{\lambda_i}{\ell}$, we get high order versions of the statements of Fact 1.5 (except the closed formula), Th. 2.4 and Th. 2.6. A partial high order version of Lemma 2.3 is the following:

Lemma 2.8 *Let D be a t -(v, k, λ) $_q$ subspace design, S a subspace of V of dimension $s = \dim(S)$ and ℓ a positive integer. For $s \leq t$ or $s \geq v - t$, the high order intersection vector $\alpha^{(\ell)}(S)$ is uniquely determined. In the case $s \leq t$,*

$$\alpha_i^{(\ell)}(S) = \begin{cases} \begin{bmatrix} s \\ i \end{bmatrix}_q \cdot \lambda_{i, s-i}^{(\ell)} & \text{for all } i \in \{0, \dots, s\} \text{ and} \\ 0 & \text{for all } i \in \{s+1, \dots, k\}. \end{cases}$$

Using the high order Mendelsohn equations, it can be checked that in the range $s \geq v - t$, the intersection vector $\alpha_i^{(\ell)}(S)$ is still unique. However, the formula gets more complicated than in Lemma 2.3. This is indicated by the fact that for $k - i > v - s$, we don't necessarily get $\alpha_i^{(\ell)}(S) = 0$ any more.

Since the high order versions complicate the presentation, their benefit is not entirely clear and the proofs only need trivial adjustments, we decided to go with the basic versions in the main part.

3 Non-existence results for block designs

For ordinary block designs, the Mendelsohn equations have been used to show that certain admissible parameter sets are not realizable. Below, we give three such examples. These results are not new, but the proofs are new alternatives or simplify the previous ones.

Theorem 3.1 *The parameter set 3-(11, 5, 2) is admissible, but not realizable.*

Proof. The numbers

$$\lambda_0 = 33, \quad \lambda_1 = 15, \quad \lambda_2 = 5, \quad \lambda_3 = \lambda = 2$$

are all integers, so the parameter set is admissible.

To show that the parameter set is not realizable, let $V = \{1, \dots, 11\}$ and assume that there is a design on V of these parameters. The Köhler equations for the intersection vector of a block B are

$$\begin{aligned} \alpha_0 &= -2 + \alpha_4 + 4\alpha_5, & \alpha_1 &= 15 - 4\alpha_4 - 15\alpha_5, \\ \alpha_2 &= 6\alpha_4 + 20\alpha_5, & \alpha_3 &= 20 - 4\alpha_4 - 10\alpha_5. \end{aligned}$$

Since B is a block, $\alpha_5 = 1$, and because of $\alpha_1 \geq 0$, the second equation forces $\alpha_4 = 0$. So the unique intersection vector is

$$\alpha(B) = (2, 0, 20, 10, 0, 1).$$

In particular, there are $\alpha_0 = 2$ blocks contained in $V \setminus B$. Because of $\#(V \setminus B) = 11 - 5 = 6$, those two blocks intersect in exactly 4 points, which contradicts $\alpha_4 = 0$. \square

Remark 3.2 (i) For block designs, the considered parameter set 2-(11, 5, 2) is the smallest admissible parameter set (in terms of v) which is not realizable, compare [16, p. 36 ff.] and [13, Table 4.44].

(ii) Theorem 3.1 is the main result of [7], where it was shown using the same intersection vector and additionally the classification of 2-(10, 4, 2) designs. Our above proof simplifies this reasoning.

Theorem 3.3 *Let $n \geq 5$ be an integer such that $4 \nmid n$. Then the parameters*

$$t = 4, \quad v = \binom{n}{2} + 2, \quad k = n + 1, \quad \lambda = 2$$

are admissible, but not realizable.

Proof. We compute

$$\begin{aligned} \lambda_3 &= \lambda_4 \cdot \frac{v-3}{k-3} = 2 \cdot \frac{\frac{n(n-1)}{2} - 1}{n-2} = n+1, \\ \lambda_2 &= \lambda_3 \cdot \frac{v-2}{k-2} = \frac{(n+1)\frac{n(n-1)}{2}}{n-1} = \frac{n(n+1)}{2}, \\ \lambda_1 &= \lambda_2 \cdot \frac{v-1}{k-1} = \frac{n(n+1)}{2} \cdot \frac{\frac{n(n-1)}{2} + 1}{n} = \frac{(n+1)(n^2-n+2)}{4}, \\ \lambda_0 &= \lambda_1 \cdot \frac{v}{k} = \frac{(n+1)(n^2-n+2)}{4} \cdot \frac{\frac{n(n-1)}{2} + 2}{n+1} = \frac{(n^2-n+2)(n^2-n+4)}{8}. \end{aligned}$$

To see that the parameters are admissible, we have to check that the values λ_i are integral. This is clear for λ_4 , λ_3 and λ_2 . The integrality of λ_1 follows from checking the three possibilities $n \equiv 1$, $n \equiv 2$ and $n \equiv 3 \pmod{4}$. For λ_0 we note that $n^2 - n$ is always even, so one of the factors $n^2 - n + 2$ and $n^2 - n + 4$ is divisible by 4 and the other one is even.

We consider the Köhler equation with $i = 0$ for a block B (so $s = k = n + 1$ and $\alpha_k = 1$). Because of

$$\begin{aligned} \binom{s}{i} \sum_{j=i}^t (-1)^{j-i} \binom{s-i}{j-i} \lambda_j \\ &= \frac{(n^2-n+2)(n^2-n+4)}{8} - \frac{(n+1)^2(n^2-n+2)}{4} + \frac{n^2(n+1)^2}{4} \\ &\quad - \frac{(n-1)n(n+1)^2}{6} + \frac{(n-2)(n-1)n(n+1)}{12} \\ &= \frac{(n-1)(n-2)^2(n-3)}{24}, \end{aligned}$$

we get the contradiction

$$\begin{aligned} \alpha_0 &= \frac{(n-1)(n-2)^2(n-3)}{24} - \sum_{j=5}^n \binom{j-1}{4} \alpha_j - \binom{n}{4} \\ &= -\frac{(n-1)(n-2)(n-3)}{12} - \sum_{j=5}^n \binom{j-1}{4} \alpha_j < 0. \quad \square \end{aligned}$$

Remark 3.4 Designs with the parameters from Theorem 3.3 would be tight, since $t = 4 = 2s$ with $s = 2$ and $\lambda_0 = \binom{v}{s}$. The existence of this series was ruled out in [20, Cor. of Th. 5], the parameters are explicitly stated as $S_2(4, k, 2 + \frac{1}{2}(k-1)(k-2))$ in [20, p. 738].

Theorem 3.5 *Let $n \geq 2$ be an integer. Then the parameters*

$$t = 3, \quad v = (2n-1)(4n-1) + 1, \quad k = 4n-1, \quad \lambda = 1$$

are admissible, but not realizable.

Proof. We compute

$$\begin{aligned} \lambda_2 &= \lambda_3 \cdot \frac{v-2}{k-2} = \frac{(2n-1)(4n-1) - 1}{4n-3} = 2n, \\ \lambda_1 &= \lambda_2 \cdot \frac{v-1}{k-1} = 2n \cdot \frac{(2n-1)(4n-1)}{4n-2} = n(4n-1), \\ \lambda_0 &= \lambda_1 \cdot \frac{v}{k} = n(4n-1) \frac{(2n-1)(4n-1) + 1}{4n-1} = 2n(4n^2 - 3n + 1). \end{aligned}$$

So the parameters are admissible.

We consider the Köhler equation with $i = 1$ for a block S (so $s = k = 4n-1$ and $\alpha_k = 1$). Because of

$$\begin{aligned} \binom{s}{i} \sum_{j=i}^t (-1)^{j-i} \binom{s-i}{j-i} \lambda_j \\ = (4n-1)(n(4n-1) - (4n-2) \cdot 2n + (2n-1)(4n-3)) \\ = (n-1)(4n-1)(4n-3), \end{aligned}$$

we get the contradiction

$$\begin{aligned} \alpha_1 &= (n-1)(4n-1)(4n-3) - \sum_{j=4}^{4n-2} j \binom{j-2}{2} \alpha_j - (4n-1) \binom{4n-3}{2} \\ &= -(n-1)(4n-1)(4n-3) - \sum_{j=4}^{4n-2} j \binom{j-2}{2} \alpha_j < 0. \quad \square \end{aligned}$$

Remark 3.6 Alternatively, Theorem 3.5 can be shown as follows. According to [6, Th. 5.6], the existence of a $3-(v, k, 1)$ design implies $\binom{v}{3} \geq \frac{v}{k} \binom{v-1}{1} \binom{k}{3}$. In our case, this yields the contradiction

$$\frac{2}{3}n(2n-1)(4n-3)(4n-1)(4n^2 - 3n + 1) \geq \frac{2}{3}(2n-1)^2(4n-3)(4n-1)(4n^2 - 3n + 1).$$

4 Intersection structure of a q -analog of the Fano plane

It is a notorious open problem if for any prime power q , a q -analog of the Fano plane exists, which is a Steiner system with admissible parameters $2-(7, 3, 1)_q$. In this section, we compute the intersection vector distribution of such a Steiner system and discuss its implications. Thereby, $\Phi_n \in \mathbb{Z}[q]$ will denote the n -th cyclotomic polynomial in q . Since all Gaussian binomial coefficients are a product of cyclotomic polynomials, they often allow a compact representation of the arising polynomials in factorized form.

In the following, let D be a $2-(7, 3, 1)_q$ subspace design. We have

$$\begin{aligned}\lambda_0 &= q^8 + q^6 + q^5 + q^4 + q^3 + q^2 + 1 = \Phi_6\Phi_7, \\ \lambda_1 &= q^4 + q^2 + 1 = \Phi_2\Phi_4, \\ \lambda_2 &= 1.\end{aligned}$$

As a showcase, for $s = 4$ the Köhler equations yield

$$\begin{aligned}\alpha_0 &= (q^8 - q^7 + q^3) - q^3\alpha_3, \\ \alpha_1 &= (q^7 + q^6 + q^5 - q^3 - q^2 - q) + (q^3 + q^2 + q)\alpha_3, \\ \alpha_2 &= (q^4 + q^3 + 2q^2 + q + 1) - (q^2 + q + 1)\alpha_3.\end{aligned}$$

Since $\lambda = 1$, $S \in \begin{bmatrix} V \\ 4 \end{bmatrix}$ can contain at most 1 block, implying $\alpha_3 \in \{0, 1\}$. Thus, the two possible intersection vectors are

$$(q^8 - q^7 + q^3, q^7 + q^6 + q^5 - q^3 - q^2 - q, q^4 + q^3 + 2q^2 + q + 1, 0)$$

and

$$(q^8 - q^7, q^7 + q^6 + q^5, q^4 + q^3 + q^2, 1).$$

Let a_i ($i \in \{0, 1\}$) be the number of $S \in \begin{bmatrix} V \\ 4 \end{bmatrix}$ of the first and the second intersection vectors, respectively. Double counting the flags $(B, S) \in D \times \begin{bmatrix} V \\ 4 \end{bmatrix}$ with $B < D$ yields

$$a_1 = \#D \cdot \begin{bmatrix} 4 \\ 1 \end{bmatrix}_q = \Phi_2\Phi_4\Phi_6\Phi_7$$

and thus

$$a_0 = \begin{bmatrix} 7 \\ 4 \end{bmatrix}_q - a_1 = q^4\Phi_6\Phi_7.$$

For $s = 3$, the two possible intersection vectors and their frequencies are computed similarly. For each $s \in \{0, 1, 2, 5, 6, 7\}$, the intersection vector is uniquely determined by Lemma 2.3. The result is shown in Table 1. For the important special cases $q = 2$ and $q = 3$, the evaluated numbers are shown in Table 2 and Table 3, respectively. For $s = 3$, we denote the two possible types of subspaces S by 3_0 (those with $\alpha_3(S) = 0$) and 3_1 (the blocks with $\alpha_3(S) = 1$). Similarly, the two different types of blocks of dimension $s = 4$ will be denoted by 4_0 and 4_1 .

#subspaces S	s	$\alpha_0(S)$	$\alpha_1(S)$	$\alpha_2(S)$	$\alpha_3(S)$
1	7	0	0	0	$\Phi_6\Phi_7$
Φ_7	6	0	0	$q^4\Phi_3\Phi_6$	$\Phi_2\Phi_4\Phi_6$
$\Phi_3\Phi_6\Phi_7$	5	0	q^8	$q^3\Phi_2\Phi_4$	Φ_4
$\Phi_2\Phi_4\Phi_6\Phi_7$	4	$q^7\Phi_1$	$q^5\Phi_3$	$q^2\Phi_3$	1
$q^4\Phi_6\Phi_7$	4	$q^3(q^5 - q^4 + 1)$	$q\Phi_1\Phi_2\Phi_3\Phi_4$	$\Phi_3\Phi_4$	0
$\Phi_6\Phi_7$	3	$q^4\Phi_4\Phi_2\Phi_1$	$q^2\Phi_3\Phi_4$	0	1
$q\Phi_2\Phi_4\Phi_6\Phi_7$	3	$q^3(q^5 - q + 1)$	$q(q^3 + q - 1)\Phi_3$	Φ_3	0
$\Phi_3\Phi_6\Phi_7$	2	$q^6\Phi_4$	$q^2\Phi_2\Phi_4$	1	0
Φ_7	1	$q^3\Phi_2\Phi_4\Phi_6$	$\Phi_3\Phi_6$	0	0
1	0	$\Phi_6\Phi_7$	0	0	0

Table 1: Intersection vector distribution of a $2-(7, 3, 1)_q$ design

#subspaces S	s	$\alpha_0(S)$	$\alpha_1(S)$	$\alpha_2(S)$	$\alpha_3(S)$
1	7	0	0	0	381
127	6	0	0	336	45
2667	5	0	256	120	5
5715	4	128	224	28	1
6096	4	136	210	35	0
381	3	240	140	0	1
11430	3	248	126	7	0
2667	2	320	60	1	0
127	1	360	21	0	0
1	0	381	0	0	0

Table 2: Intersection vector distribution of a $2-(7, 3, 1)_2$ design

#subspaces S	s	$\alpha_0(S)$	$\alpha_1(S)$	$\alpha_2(S)$	$\alpha_3(S)$
1	7	0	0	0	7651
1093	6	0	0	7371	280
99463	5	0	6561	1080	10
306040	4	4374	3159	117	1
619731	4	4401	3120	130	0
7651	3	6480	1170	0	1
918120	3	6507	1131	13	0
99463	2	7290	360	1	0
1093	1	7560	91	0	0
1	0	7651	0	0	0

Table 3: Intersection vector distribution of a $2-(7, 3, 1)_3$ design

Theorem 4.1 *Let q be a prime power. The existence of a $2-(7, 3, 1)_q$ subspace design implies the existence of a $2-(7, 3, q^4)_q$ design.*

Proof. Let S be a 5-space in V . By the unique intersection vector for $s = 5$, there are $\Phi_4 = q^2 + 1$ blocks contained in S . A 4-space $W \leq S$ is of type 4_1 if and only if it contains one of those blocks. For each such block B , there are $\begin{bmatrix} 5-3 \\ 4-3 \end{bmatrix}_q = \begin{bmatrix} 2 \\ 1 \end{bmatrix}_q = q + 1$ intermediate 4-spaces W with $B \leq W \leq S$. This gives us the number of spaces of type 4_1 in S as $(q^2 + 1)(q + 1) = q^3 + q^2 + q + 1$. Therefore, the number of spaces of type 4_0 in S is $\begin{bmatrix} 5 \\ 4 \end{bmatrix}_q - (q^3 + q^2 + q + 1) = q^4$. This shows that $\{S^\perp \mid S \in \begin{bmatrix} V \\ 4 \end{bmatrix} \text{ of type } 4_0\}$ forms a $2-(7, 3, q^4)_q$ design. \square

Remark 4.2 The blocks of the original $2-(7, 3, 1)_q$ design are given by the spaces of type 3_1 , and the above proof shows that after dualization, the spaces of type 4_0 form the blocks of a $2-(7, 3, q^4)_q$ design. Similarly, the spaces of type 3_0 are the blocks of a $2-(7, 3, q^4 + q^3 + q^2 + q)_q$ design and after dualization, the spaces of type 4_1 are the blocks of a $2-(7, 3, q^3 + q^2 + q + 1)_q$ design. However, these are just the complementary designs of the ones arising from the spaces 3_1 and 4_0 , respectively.

The resulting “intersection structure” of a 2-analog of the Fano plane is shown in Figure 1. We explain by a few examples how to read this figure: The entry $(128, 224, 28, 1)^{5715}$ on the level $s = 4$ means that there are 5715 subspaces of dimension $s = 4$ having the intersection vector $(128, 224, 28, 1)$ (the subspaces of type 4_1). It is connected by a line to the intersection vector $(248, 126, 7, 0)$ (type 3_0) because a subspace of type 4_1 contains subspaces of type 3_0 . More precisely, the number 14 at the 4_1 -end of the line tells us that each subspace of type 4_1 contains exactly 14 subspaces of type 3_0 . Similarly, the number 7 at the 3_0 -end means that each subspace of type 3_0 is contained in exactly 7 subspaces of type 4_1 .

Acknowledgement

The authors would like to acknowledge the financial support provided by COST – *European Cooperation in Science and Technology*. Both authors are members of the Action IC1104 *Random Network Coding and Designs over $GF(q)$* . This research was carried out during a 6 week stay of the first author at the University of Zagreb, which was supported by an STSM grant of the COST project.

We would like to express our gratitude to R. Laue for pointing out this problem, giving us hints for references and sharing some very useful thoughts with us.

References

- [1] A. Betten. “Schnittzahlen von Designs”. PhD thesis. Universität Bayreuth, 1998.
- [2] M. Braun, T. Etzion, P. Östergård, A. Vardy, and A. Wassermann. *Existence of q -Analogues of Steiner Systems*. Apr. 4, 2013. arXiv:1304.1462.




Figure 1: Intersection structure of a $2-(7, 3, 1)_2$ design

- [3] M. Braun, A. Kerber, and R. Laue. “Systematic construction of q -analogs of t - (v, k, λ) -designs”. In: *Des. Codes Cryptogr.* 34.1 (2005), pp. 55–70.
- [4] P. J. Cameron. “Generalisation of Fisher’s inequality to fields with more than one element”. In: *Combinatorics. Proceedings of the British Combinatorial Conference 1973*. Ed. by T. P. McDonough and V. C. Mavron. London Mathematical Society Lecture Note Series 13. Cambridge: Cambridge University Press, 1974, pp. 9–13. ISBN: 0-521-20454-2.
- [5] H. Cohn. “Projective geometry over \mathbb{F}_1 and the Gaussian binomial coefficients”. In: *Amer. Math. Monthly* 111.6 (2004), pp. 487–495.
- [6] C. J. Colbourn and R. Mathon. “Steiner Systems”. In: *Handbook of Combinatorial Designs*. Ed. by C. J. Colbourn and J. H. Dinitz. 2nd ed. Chapman & Hall/CRC, 2007. Chap. II.5, pp. 102–119. ISBN: 1-58488-506-8.
- [7] M. Dehon. “Non-existence d’un 3-design de paramètres $\lambda = 2$, $k = 5$ et $v = 11$ ”. In: *Discrete Math.* 15.1 (1976), pp. 23–25.
- [8] P. Delsarte. “Association schemes and t -designs in regular semilattices”. In: *J. Combin. Theory Ser. A* 20.2 (1976), pp. 230–243.
- [9] A. Fazeli, S. Lovett, and A. Vardy. *Nontrivial t -designs over finite fields exist for all t* . June 10, 2013. arXiv:1306.2088.
- [10] R. D. Fray. “Congruence properties of ordinary and q -binomial coefficients”. In: *Duke Math. J.* 34.3 (1967), pp. 467–480.
- [11] J. Goldman and G.-C. Rota. “On the foundations of combinatorial theory. IV. Finite vector spaces and Eulerian generating functions”. In: *Stud. Appl. Math.* 49.3 (1970), pp. 239–258.
- [12] F. H. Jackson. “ q -difference equations”. In: *Amer. J. Math.* 32.3 (1910), pp. 305–314.
- [13] G. R. Khosrovshahi and R. Laue. “ t -designs with $t \geq 3$ ”. In: *Handbook of Combinatorial Designs*. Ed. by C. J. Colbourn and J. H. Dinitz. 2nd ed. Chapman & Hall/CRC, 2007. Chap. II.1, pp. 25–58. ISBN: 1-58488-506-8.
- [14] M. Kiermaier and R. Laue. *Derived and residual subspace designs*. May 21, 2014. arXiv:1405.5432.
- [15] E. Köhler. “Allgemeine Schnittzahlen in t -designs”. In: *Discrete Math.* 73.1–2 (1988–1989), pp. 133–142.
- [16] R. Mathon and A. Rosa. “2- (v, k, λ) designs of small order”. In: *Handbook of Combinatorial Designs*. Ed. by C. J. Colbourn and J. H. Dinitz. 2nd ed. Chapman & Hall/CRC, 2007. Chap. II.1, pp. 25–58. ISBN: 1-58488-506-8.
- [17] N. S. Mendelsohn. “Intersection numbers of t -designs”. In: *Studies in Pure Mathematics*. Ed. by L. Mirsky. London: Academic Press, 1971, pp. 145–150. ISBN: 0-12-498450-9.

- [18] W. Oberschelp. “Lotto-Garantiesysteme und Blockpläne”. In: *Math.-Phys. Semesterber., N. F.* 19 (1972), pp. 55–67.
- [19] G. Pólya and G. Alexanderson. “Gaussian binomial coefficients”. In: *Elem. Math.* 26 (1971), pp. 102–109. URL: <http://eudml.org/doc/141022>.
- [20] D. K. Ray-Chaudhuri and R. M. Wilson. “On t -designs”. In: *Osaka J. Math.* 12.3 (1975), pp. 737–744.
- [21] D. Ray-Chaudhuri and N. Singhi. “ q -analogues of t -designs and their existence”. In: *Linear Algebra Appl.* 114–115 (1989), pp. 57–68.
- [22] H. Suzuki. “Five Days Introduction to the Theory of Designs”. 1989. URL: <http://subsite.icu.ac.jp/people/hsuzuki/lecturenote/designtheory.pdf>.
- [23] H. Suzuki. “2-designs over $GF(2^m)$ ”. In: *Graphs Combin.* 6.3 (1990), pp. 293–296.
- [24] H. Suzuki. “On the inequalities of t -designs over a finite field”. In: *European J. Combin.* 11.6 (1990), pp. 601–607.
- [25] H. Suzuki. “‘ t -designs’ in $H(d, q)$ ”. In: *Hokkaido Math. J.* 19.3 (1990), pp. 403–415.
- [26] H. Suzuki. “2-designs over $GF(q)$ ”. In: *Graphs Combin.* 8.4 (1992), pp. 381–389.
- [27] S. Thomas. “Designs over finite fields”. In: *Geom. Dedicata* 24.2 (1987), pp. 237–242.
- [28] T. V. Trung, Q.-r. Wu, and D. M. Mesner. “High order intersection numbers of t -designs”. In: *J. Statist. Plann. Inference* 56.2 (1996), pp. 257–268.
- [29] C. de Vroedt. “Über einen Satz von Köhler”. In: *Discrete Math.* 97.1–3 (1991), pp. 161–165.
- [30] M. Ward. “A calculus of sequences”. In: *Amer. J. Math.* 58.2 (1936), pp. 255–266.