

HIGHER RESIDUE PAIRING ON CRYSTALLINE LOCAL SYSTEMS

MOHAMMAD REZA RAHMATI

ABSTRACT. We explain a generalization of the K. Saito higher residue pairing for local system of p -adic isocrystals.

1. INTRODUCTION

In complex Hodge theory the variation of Hodge structure (VHS) associated to an analytic projective fibration $f : X \rightarrow S$ between quasi-projective varieties with a finite number of critical points, locally admits a universal flat sesqui-linear nondegenerate pairing namely Higher residue pairing of K. Saito. The construction and properties of the Saito bilinear pairing was first studied in [S]. The flat non-degenerate pairing of an isolated singularity can also be stated using the duality between the twisted de Rham complex of f and the twisted complex of polyvector fields. This view of point has been considered in [LLS] with an insight toward mirror symmetry.

The crystalline cohomology of varieties defined over a quotient ring of Witt vectors also satisfies a similar Hodge-Tate decomposition. Therefore a family as above also defines a family of filtered isocrystals which are crystalline representations of the absolute Galois group of a p -adic field. The relative de Rham-Witt complex of the fibration f defines a variation of filtered crystalline cohomology over $\mathbf{C} := \widehat{\mathbf{Q}}$ where \mathbf{Q} is the quotient field of the Witt ring $W(k)$ of the perfect field k of characteristic p .

The family of crystalline cohomologies $H_{crys}^k(X_s/W) \otimes \mathbf{C}$, for each k , defines a variation of Hodge-Tate (HT)-structure $H^{(k)} = Rf_* \mathbf{C}_X$ together with a flat connection defined locally on S . The HT-structure defines the slope filtrations on the fibers $H_{\mathbf{C}}$. The p -adic Riemann-Hilbert correspondence guarantees that $H = \ker \nabla$ as a 1-1 functorial correspondence between p -adic local systems and flat connections over the base, cf.[RZ]. We screen this construction over the p -ring \mathbf{C} to state Kyoji Saito form S_f over this ring.

1991 *Mathematics Subject Classification.* 11R23, 11F80, 11S25, 11G40.

Key words and phrases. Ring of Witt vectors, Complete local rings, Higher residue pairing, Crystalline cohomology.

2. THE SET UP FOR RESIDUE PAIRING

Let k be a perfect field of char = p , and $W = W(k)$ be the ring of Witt vectors over k . As a set W can be considered as the infinite product k^∞ with specific sum and products given in each component by polynomials in char = p . Such a ring has characteristic 0. Let \mathbf{Q} be the quotient field of W and $\mathbf{C} = \widehat{\mathbf{Q}}$ the completion of its algebraic closure. We consider family of projective varieties X_s over \mathbf{C} defined by a projective map $f : X \rightarrow S$, where X and S are quasi-projective varieties. We assume $\dim X = n$, $\dim S = 1$ and $S = B(\mathbf{C})$ to be a small disc. Then the relative algebraic de Rham-Witt complex $W\Omega_{X/S}^\bullet$ of the fibration defines a Gauss-Manin system of p -adic isocrystals

$$(1) \quad (E = R^k f_* W\Omega_{X/S}^\bullet, \nabla), \quad \ker(\nabla) = R^k f_* \mathbf{C}_X$$

where $\nabla : E \rightarrow E \times W\Omega_S^1$ is a flat connection, with kernel the local system $\mathcal{V} = R^k f_* \mathbf{C}$. We may pass to the completion of the local rings over $B(\mathbf{C})$ and consider the twisted complex

$$(2) \quad \widehat{\Omega}^\bullet := (DR_X[[\pi]], \widehat{d} = d + \frac{df}{\pi} \wedge)$$

where π is the uniformizer of the local ring of B at 0. Set $\widehat{H}_f := H^*(\widehat{\Omega}^\bullet)$. The module \widehat{H}_f has a natural decreasing filtration F^k induced by multiplication with π^k . The trace map

$$(3) \quad Tr : W\widehat{H}_f \longrightarrow \mathcal{O}_{S,0}[[\pi]]$$

induces a $k[[\pi]]$ -homomorphism \widehat{Res}_f that can be written as

$$(4) \quad W\widehat{Res}_f = \sum_k W\widehat{Res}_f^{(k)}(\cdot)\pi^k$$

In fact putting $S_f(u, u') := Tr(\tilde{u}\overline{\tilde{u}'})$, where \tilde{u} is the lift of u in the complex $\widehat{\Omega}^\bullet$, defines a bilinear pairing. Therefore we can define the higher residue pairing;

Theorem 2.1. *(Higher residue pairing on crystalline site) Let $f : X \rightarrow S$ be a family of projective varieties defined over the p -ring \mathbf{C} . Let \widehat{H}_f be the cohomology of the twisted relative de Rham-Witt complex of the fibration. Then we have a \mathbf{C} -nondegenerate bilinear form*

$$(5) \quad WS_f(,) : W\widehat{H}_f \times W\widehat{H}_f \rightarrow \mathbf{C}[[\pi]]$$

Let u_1, u_2 be local sections of $W\widehat{H}_f$, then;

- $WS_f(u_1, u_2) = \overline{WS_f(u_2, u_1)}$.
- $WS_f(g(\pi)u_1, u_2) = WS_f(u_1, v(-\pi)u_2) = g(\pi)WS_f(u_1, u_2)$, $g(\pi) \in \mathbf{C}[[\pi]]$.
- $\partial_\pi WS_f(u_1, u_2) = WS_f(\partial_\pi u_1, u_2) + WS_f(u_1, \partial_\pi u_2)$.
- $(\pi\partial_\pi + n)WS_f(u_1, u_2) = WS_f(\pi\partial_\pi u_2, u_1) + WS_f(u_1, \pi\partial_\pi u_2)$
- *The induced pairing on $W\widehat{H}_f/\pi.W\widehat{H}_f \otimes W\widehat{H}_f/\pi.W\widehat{H}_f \rightarrow \mathbf{C}$ is the classical Grothendieck residue.*

The conjugation is formally done by $\overline{g(\pi) \otimes \eta} = g(-\pi).\eta$. The pairing is defined via the trace map on the cohomology of the complex $\widehat{\Omega}^\bullet$ by

$$(6) \quad S_f(g_1(\pi)u, g_2(\pi)u') := g_1(\pi)g_2(-\pi)Tr(\tilde{u}.\tilde{u}')$$

The flat pairing (5) is the universal pairing on the local system of isocrystals defined by the variation of crystalline cohomology of fibration. The pairing is called higher residue pairing due to the works by K. Saito [S].

Proof. The proof is analogous to the complex case in [S], see also [LLS]. The first four properties are straight forward by the definitions. \square

Remark 2.2. *The period isomorphism between crystalline and étale cohomology with coefficient in \mathbb{C}_p implies that a similar form can be defined in variation of p -adic étale cohomology.*

REFERENCES

- [LLS] Li C., Li S., Saito K., Primitive forms via polyvector fields, arXiv:1311.1659v3, 2014.
[LZ] Langer A., T. Zink, De Rham-Witt cohomology for a proper and smooth morphism, Journal of the Institute of Mathematics of Jussieu, 231 - 314, 2003
[RZ] L. Ruochuan, X. Zhu, Rigidity and a Riemann-Hilbert correspondence for p -adic local systems. Invent. Math. 207 (2017), no.1, 291-343
[S] Saito K., Period mapping associated to a primitive form, Publications of Research Inst. Math. Sci., Kyoto Univ., Vol 19, No 3, 1983
[ST] Stack Project, Crystalline cohomology, Cotangent complex

ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES, PAKISTAN

E-mail address: mrahmati@cimat.mx, rahmati@sms.edu.pk