

Exact values for the Grundy number of some graphs

Ali Mansouri^{*}, Mohamed salim bouhlef^{**}

mehermansouri@yahoo.fr
medsalim.bouhlef@enis.rnu.tn

Abstract: The Grundy number of a graph G is the maximum number k of colors used to color the vertices of G such that the coloring is proper and every vertex x colored with color i , $1 \leq i \leq k$, is adjacent to $(i - 1)$ vertices colored with each color j , $1 \leq j \leq i - 1$.

In this paper we give bounds for the Grundy number of some graphs and Cartesian products of graphs. In particular, we determine an exact value of this parameter for n -dimensional meshes and some n -dimensional toroidal meshes. Finally, we present an algorithm to generate all graphs for a given Grundy number.

Key words: algorithm, cartesian product of several graphs, graphs, bipartite graph, Grundy number,

INTRODUCTION

We consider graphs without loops or multiple edges. Let G be a graph on vertices $x_0; x_1; \dots; x_{n_G-1}$, with vertex set $V(G)$ and edge set $E(G)$. Let $d(x)$ be the degree of the vertex x of G and let $\Delta(G)$ be the maximum degree of G .

The Cartesian product of two graphs $G = (V_1; E_1)$ and $H = (V_2; E_2)$, denoted $G \square H$, has the vertex set $V_1 \times V_2$, and the neighborhood of each vertex $(x_1; x_2)$ is $NG \square H((x_1; x_2)) = (\{x_1\} \times NH(x_2)) \cup (NG(x_1) \times \{x_2\})$. Thus, in the graph $G \square H$ we find several copies of graphs G and H denoted by G_i and H_j , where i represents the rows of $G \square H$, with $0 \leq i \leq n_G - 1$, and j represents the columns of $G \square H$, with $0 \leq j \leq n_H - 1$. The set $\{x_0^j; x_1^j; \dots; x_{n_G-1}^j\}$ denotes the vertices of the j 'th copy of G in $G \square H$, with $0 \leq j \leq n_H - 1$.

Next, we define a k -coloring of G as a function c defined on $V(G)$ into a set of colors

$C = \{1, 2, \dots, k\}$ such that for each vertex x_i , with $0 \leq i \leq n_G - 1$, $c(x_i) \in C$. A proper k -coloring is a k -coloring verifying the condition $c(x) \neq c(y)$ for any pair of adjacent vertices $x; y \in V(G)$. A Grundy coloring is a proper k -coloring satisfying the following property P :

Every vertex v , colored with color i ($1 \leq i \leq k$), is adjacent to vertices colored by each color j such that $1 \leq j \leq i - 1$. The Grundy number $\Gamma(G)$ of a graph G is then defined as the maximum number of colors among all Grundy colorings of G . If we color only a set of vertices of a graph G , we will say that the coloring of G is partial.

This parameter was introduced by Christen and Selkow [CHS79] in 1979. They proved that determining the Grundy number is NP-complete for general graphs (also studied by McRae in 1994 [AMC94]). In [SST84], Hedetniemi et al. gave a linear

algorithm for the Grundy number of a tree and established a relation between the chromatic number, the Grundy number and the achromatic number: $\chi(G) \leq \Gamma(G) \leq \Psi(G)$, where the achromatic number $\Psi(G)$ is the maximum number of colors used for a proper coloring of G such that each pair of colors appears on at least one edge of G . In 1997, Telle and Proskurowski [GAP97] gave an algorithm for the Grundy number of partial k -trees in $O(n^3k^2)$ and bounded this parameter for these graphs by the value $1 + k \log_2 n$, where n is the graph order. In 2000, Dunbar et al. used the Grundy number to bound new parameters that they introduced in [GHK00], the chromatic and the achromatic numbers of a fall coloring. Recently, Germain and Kheddouci studied in [CGH03], [CGH06] the Grundy coloring of power graphs. They gave bounds for the Grundy number of the power graphs of a path, a cycle, a caterpillar and a complete binary tree. Such colorings are also explored for other graphs like chessboard graphs [CPR02].

The Cartesian product of graphs is widely studied in literature since it generates interesting classes of graphs like grids (obtained by the Cartesian product of paths and cycles) or hypercubes, which are used to model problems on interconnection networks or multiprocessor networks [CWE97],[SGR02],[GHW02], [FSA03]. In particular, several coloring parameters were evaluated for graphs resulting from the Cartesian product of graphs.

Cizek and Klavzar [NCK94] computed chromatic numbers of the Cartesian sum of two odd cycles. In [MK02], Kouider and Mahéo were interested in the b -chromatic number of the Cartesian product of two graphs. In [XZH92] and [XZH02], Zhu presented several bounds for respectively the star and the fractional chromatic number of graph products.

In our case, we are interested in the Grundy number of the Cartesian product of graphs. This

parameter has a lot of applications in fields like scheduling or multiprocessor architectures. For instance, suppose a set of processes such that a process P_i could be computed if processes $P_1; P_2; \dots; P_{i-1}$ are already computed. Such a law on the processes can be modeled by a Grundy coloring. Thus, if we consider a given architecture G , the study of the Grundy number brings a solution for two questions.

First, how many processes can we put into this architecture? This is directly given by the Grundy number $\Gamma(G)$ of the architecture G . Second, how many times must we load processes in the architecture to compute P_n ? The relation $n - \Gamma(G) / \Gamma(G - 1)$ gives a solution to this second question.

Thus in this paper we will decompose our study in several parts. First in Section 2, we will present some properties of the Grundy number, by comparing it to other graph parameters. Then we will discuss in Section 3 the Grundy number of several Cartesian products of two graphs (paths, cycles, complete and bipartite graphs. . .).

In Section 4, this parameter will be studied for the Cartesian product of several graphs. Thus, we will determine an exact value of the Grundy number for n -dimensional meshes and some n -dimensional toroidal meshes. In these sections, in addition to determining exact values and bounds for the Grundy number, we will also propose constructions of these colorings. Finally in Section 5, we will present an algorithm to generate graphs for a given Grundy number

1. Grundy number of a graph

The First, the following obvious fact enables us to find a proper coloring for a graph G from a proper coloring of G' where G' is a subgraph of G .

Fact 1 Let G' be an induced subgraph of G given by a set of vertices $V' \subseteq V$. Any proper coloring of G' can be extended to a proper coloring of G .

Proof We extend the coloring of G' to G as follows. Let x be a vertex of G such that $x \notin V'$. Let C be the set of colors of $NG'(x)$. Let c be the smallest color such that $c \notin C$. We put $c_x = c$ and $V' := V' \cup \{x\}$. Then, we repeat this process until $V' = V$.

Then, we present some results for the Grundy number of simple graphs.

Proposition 1 Let S_n, K_n, P_n, C_n and $K_{n;p}$ be respectively the stable graph, the complete graph, the path, the cycle on order n and the complete bipartite graph on $n + p$ vertices. Let G be a non connected graph with connected components $G_1; G_2; \dots; G_p$. Then, we have:

Proof. For cases 1 to 4, the proofs are obvious.

For the complete bipartite graph, we prove the

result by contradiction. Let $V(K_{n;p}) = A \cup B$, with $A = \{a_1; a_2; \dots; a_n\}$ and $B = \{b_1; b_2; \dots; b_p\}$. Let a_i be a vertex colored by a color c , with $1 \leq i \leq n$ and $c \geq 3$. There exist two vertices $b_j; b_k$ colored respectively by

1. $\Gamma(S_n) = 1$ and $\Gamma(K_n) = n$
2. $\Gamma(P_n) = 2$ if $2 \leq n \leq 3$ $\Gamma(P_n) = 3$ if $n \geq 4$
3. $\Gamma(C_n) = 2$ if $n = 4$ $\Gamma(C_n) = 3$ if $n \neq 4$
4. $\Gamma(G) \geq \max \{\Gamma(G_i) : 1 \leq i \leq p\}$
5. $\Gamma(K_{n;p}) = 2$

colors 1 and 2, with $1 \leq j \neq k \leq p$. Observe that b_k must be adjacent to color 1 on a vertex $a_{i'}$, with $1 \leq i' \neq i \leq n$. So $a_{i'}$ and b_j admit the same color 1, which is a contradiction.

Next, we show two results where the Grundy number of a graph G is bounded by other parameters of G . Firstly, we give a relation between the Grundy number and the stability number of G .

Theorem 1 Let G be a connected graph on order n and stability number α . Then $\Gamma(G) \leq n + 1 - \alpha$

Proof. Let A be an independent set in G of size α . Suppose that there is a Grundy coloring of G with $k \geq n - \alpha + 2$ colors. Then at least two color classes should be subsets of A . This contradicts the definition of a Grundy coloring.

Secondly, we present the inequality of "Nordhaus-Gaddum"-type for Grundy number of some graph classes.

Proposition 2 Let G be a graph on order n and G^- its complement. Let Γ and Γ^- be the Grundy number of G and G^- respectively. Let x and y be two vertices of G (noted x^- and y^- in G^-) such that $c_x = \Gamma$ and $c_y^- = \Gamma^-$. Then, $\Gamma + \Gamma^- \leq n + 1$ if one of these assertions is verified:

1. G is a k -regular graph, with $k \geq 1$
2. $d(x) \leq d(y)$
3. $c_x = \Gamma$ and $c_{x^-} = \Gamma^-$

Proof.

1. The graph G is k -regular. By definition we have $\Gamma(G) \leq \Delta(G) + 1$ and $\Gamma(G^-) \leq \Delta(G^-) + 1$. Since the graph is k -regular, we deduce that $\Delta(G) = k$ and $\Delta(G^-) = n - 1 - k$. Thus, $\Gamma(G) + \Gamma(G^-) \leq \Delta(G) + \Delta(G^-) + 2 \leq n + 1$.

2. By definition we have

$$d(x) \geq \Gamma - 1; (1)$$

And

$$d(y^-) \geq \Gamma^- - 1; (2)$$

Then, inequality (2) implies

$$d(y) = n - 1 - d(y^-) \leq n - \Gamma^- : (3)$$

As $d(x) \leq d(y)$, inequalities (1) and (3) give

$$\Gamma - 1 \leq d(x) \leq d(y) \leq n - \Gamma;$$

$$\Gamma + \Gamma^- \leq n + 1.$$

3. From inequalities (1) and (2) we can deduce

$$\Gamma + \Gamma \leq d(x) + d(x^-) + 2 = d(x) + (n - d(x) - 1) + 2 = n + 1.$$

Remark: Observe that this inequality is not verified for any graph. Indeed, consider a graph G on order n composed by a complete graph K_3 (vertices denoted x_1, x_2 and x_3), where every node x_i is the center of a star K_{1, n_i} ; n_i , with $1 \leq i \leq 3$ and $n_i \geq 1$. By coloring G such that $c(x_1) = 2, c(x_2) = 3, c(x_3) = 4$ and each endvertex of the stars is colored by 1 (see Figure 1.a), we find $\Gamma(G) \geq 4$. Then in the complement graph G^- , the endvertices of the stars form a clique (denoted $K_{n_1+n_2+n_3}$) and vertices x_1, x_2 and x_3 are the centers of the stars respectively $K_{1, n_2+n_3}, K_{1, n_1+n_3}$ and K_{1, n_1+n_2} .

Therefore by coloring $c(x_1) = c(x_2) = c(x_3) = 1$ and every vertex of the clique with a different color c , with $2 \leq c \leq n-2$ (see Figure 1.b); we have $\Gamma(G^-) \geq n-2$. Thus, $\Gamma(G) + \Gamma(G^-) \geq n + 2$. This remark presents a counterexample to the inequality of "Nordhaus-Gaddum"-type for any graph.




Figure 1: Grundy colorings of a) G and b) G^- where $2 \leq c_1 \neq c_2 \neq \dots \neq c_{n-3} \leq n-2$.

2. Grundy number of the Cartesian product of two graphs

In this section, we will discuss the Grundy number of the Cartesian product of two graphs. In particular we will study the Cartesian product of two paths, two cycles, a path by a cycle, a bipartite graph by other graphs and a complete graph by any graph G .

Proposition 3 Let G be a bipartite graph. Let P_n be a path on order $n, 3$ and C_m be a cycle on order $m \geq 4$. Then,

$\Gamma(G \square P_n) \geq \Gamma(G) + 2$
$\Gamma(G \square C_m) \geq \Gamma(G) + 2.$

Proof. On the copy G_1 , we put the coloring of G incremented by 2. Thus, on G_1 we find the colors 3 to

$\Gamma(G) + 2$. Let X_i and Y_i be the two independent sets of the copy G_i . We put the color 1 on every vertex of X_0 and Y_2 , and the color 2 on every vertex of X_2 and Y_0 . If some vertices are not colored, Fact 1 allows us to color them with a proper coloring. Thus we can deduce $\Gamma(G \square P_n) \geq \Gamma(G) + 2$ and $\Gamma(G \square C_m) \geq \Gamma(G) + 2$.

As we studied the Cartesian product of G by a path or a cycle, where G is a bipartite graph, we discuss the same products if G is not bipartite.

Proposition 4 Let G be a non bipartite graph on order n . Let P_n be a path on order $n, 4$ and C_m be a cycle on order $m, 4$. Then:

$\Gamma(G \square P_n) \geq \Gamma(G) + 1;$
$\Gamma(G \square C_m) \geq \Gamma(G) + 1.$

Proof. The proof is given by construction. We color G_0, G_1 and G_2 with the coloring of G . Let x_p^0 be a vertex of G_0 colored by $\Gamma(G)$. Then we put $c_{x_p^0} = \Gamma(G) + 1$.

$c_{x_p^1} = \Gamma(G)$ and $c_{x_p^2} = \Gamma(G) - 1$. Next we can remove the colors of vertices x_j^1 on G_1 for which $c_{x_j^1} = 1$ (with $0 \leq j \neq p \leq nG - 1$) and the colors of vertices x_i^2 on G_2 for which $c_{x_i^2} = \Gamma(G) - 1$ (with $0 \leq i \neq p \leq nG - 1$). Then for every remaining colored vertex on G_1 , we compute $c_{x_j^1} = c_{x_j^1} - 1$.

Finally, by Fact 1, we color the remaining non colored vertices with a proper coloring. Thus we can deduce:

$\Gamma(G \square P_n) \geq \Gamma(G) + 1$ and $\Gamma(G \square C_m) \geq \Gamma(G) + 1.$
--

Remark Let P_n and P_m be the paths on order n and m respectively. And let C_n, C_m and C_k be the cycles on respectively n, m and k vertices. Then,

1. $\Gamma(P_n \square P_m) = (\leq) 4$ if $n = 2$ or $n = m = 3$; 5 otherwise;
2. $\Gamma(P_n \square C_k) = 4$ if $n = 2$ or $n = k = 3$; 5 otherwise;
3. $\Gamma(C_n \square C_m) = (\leq) 4$ if $n = 3$ and $m = 4$ or $n = m = 3$; 5 otherwise.

Proof. In each case, the maximum degree is 4. So the Grundy number is less or equal to 5. The lower bounds are obvious; they are deduced from constructions (see Figure 2).

In the following theorems we study the Grundy number for the Cartesian product of a complete graph by another graph. Firstly, we present the Cartesian

product of a bipartite graph by a complete graph.

Theorem 2 Let G be a bipartite graph on order nG . Let K_p be a complete graph on order $p \geq 3$. Then,

$$\Gamma(G) + p - 1 \leq \Gamma(K_p \square G) \leq p + \Delta(G).$$




Figure 2: Grundy colorings of a) $P_4 \square P_5$, b) $P_5 \square C_4$, c) $C_4 \square C_4$ and d) $C_4 \square C_5$.

Proof. We prove by construction that $\Gamma(K_p \square G) \geq \Gamma(G) + p - 1$. Let X_i and Y_i be the two independent sets of the copy G_i . We color G_0 by $cx_0i = cxi + p - 1$ where cxi is the color of the vertex xi of G , with $0 \leq i \leq nG - 1$. Then, for each $1 \leq i \leq p - 1$, we color each vertex of X_i with the color i , and each vertex of Y_i with the color $i + 1 \pmod{p - 1}$. For example, Figure 3.b gives a Grundy coloring of $K_4 \square G$. As $\Delta(K_p \square G) = \Delta(G) + p - 1$, we can deduce that $\Gamma(K_p \square G) \leq \Delta(G) + p$.




Figure 3: Grundy colorings of a) G and b) $K_4 \square G$

Remark: For all the classes of bipartite graphs such that $\Gamma(G) = \Delta(G) + 1$, the equality $\Gamma(K_p \square G) = \Gamma(G) + p - 1$ holds.

Secondly we bound the Grundy number of the Cartesian product of a complete graph by any graph G .

Theorem 3 Let G be a graph on order nG . Let K_n be a complete graph on order n . Then,

$$\Gamma(K_n \square G) \geq \begin{cases} n + \Gamma(G) - 1 & \text{if } \Gamma(G) \leq n - 1. \\ 2n - 2 & \text{if } n \leq \Gamma(G) \leq 2n - 3. \\ \Gamma(G) & \text{if } \Gamma(G) \geq 2n - 2. \end{cases}$$

$$\Gamma(K_n \square G) \leq n + \Delta(G).$$

Proof. As $\Delta(K_n \square G) = \Delta(G) + n - 1$, we have $\Gamma(K_n \square G) \leq \Delta(K_n \square G) + 1 \leq n + \Delta(G)$. The proof of the lower bound is given by construction.

- $\Gamma(G) \leq n - 1$. Let $k = n + \Gamma(G) - 1$. For each vertex x_0^i with $0 \leq i \leq nG - 1$, we shift the color of the vertex xi on G by $(n - 1)$

Then, for each vertex x_1^i with $0 \leq i \leq nG - 1$, we put $cx_1^i = cx_i$. Next, for each remaining vertex, we put $cx_j^i = cx_{i-1}^i + 1 \pmod{n - 1}$, with $2 \leq i \leq n - 1$ and $0 \leq j \leq nG - 1$. Figure 4.b shows a Grundy coloring of $K_5 \square G$.

$n \leq \Gamma(G) \leq 2n - 3$. Let $k = 2n - 2$. Let x_i , with $0 \leq i \leq nG - 1$, be a vertex of G colored by $\Gamma(G)$. As $\Gamma(G) \geq n$, then x_i (resp. x_0^i) has at least $n - 1$ neighbors in G (resp. G_0). Let $\{v_1; v_2; \dots; v_{n-1}\}$ be a set of $n - 1$ vertices of $NG_0(x_0^i)$. To color $K_n \square G$ we put the colors $(k; k - 1; \dots; n)$ respectively on vertices $x_0^i; x_1^i; \dots; x_i$. Then, on each copy K_n^j containing a vertex vm , with $0 \leq j \leq nG - 1$ and $1 \leq m \leq n - 1$, we put the colors $(m; m + 1; \dots; m + n - 2)$, where each color is taken modulo $(n - 1)$, on vertices $x_0^j; x_1^j; \dots; x_{n-2}^j$ respectively.

- $\Gamma(G) \geq 2n - 2$. We put the coloring of G on G_0 and Fact 1 completes the coloring of $K_n \square G$ to have a proper coloring.

3. Grundy number of the Cartesian product of several graphs

The results presented in the previous sections enable us to deduce some results for the Cartesian product of several graphs. Firstly, we determine exact values for the Grundy number of the Cartesian product of several paths and the Cartesian product of several even cycles.

Proposition 5[BEH02] Let $P_{n_1}; P_{n_2}; \dots; P_{n_k}$ be the paths of sizes respectively $n_1; n_2; \dots; n_k$ such that $k \geq 2, n_i \geq 3$ for each $1 \leq i \leq k$ and $\max\{n_i : 1 \leq i \leq k\} > 3$. Let $C_{m_1}; C_{m_2}; \dots; C_{m_k}$ be the cycles of sizes respectively $m_1; m_2; \dots; m_k$, where $k \geq 2, m_i \geq 4$ and m_i is even, for every $1 \leq i \leq k$. Then,

$$\Gamma(P_{n_1} \square P_{n_2} \square \dots \square P_{n_k}) = 2k + 1$$

$$\Gamma(C_{m_1} \square C_{m_2} \square \dots \square C_{m_k}) = 2k + 1$$

Remark: The graph $P_{n_1} \square P_{n_2} \square \dots \square P_{n_k}$ is a n_k -dimensional mesh. Therefore, Proposition 10 gives the Grundy number of an n -dimensional mesh.

Secondly, we give a bound to the parameter for the Cartesian product of odd cycles.

Proposition 6 Let $C_3; C_5; \dots; C_{2k+1}$ be the cycles of size 3; 5... $2k + 1$ respectively. The Grundy number of $C_3 \square C_5 \square \dots \square C_{2k+1}$ is given by:

$$\Gamma(C_3 \square C_5 \square \dots \square C_{2k+1}) = 2k + 1$$




Figure 4: Partial colorings of a) $C_3 \square C_5$ and b) $C_3 \square C_5 \square C_7$.

Remark: The graph $C_{n1} \square C_{n2} \square \dots \square C_{nk}$ is a n -dimensional toroidal mesh. So Proposition 10 and Proposition 12 give the Grundy number of a n -dimensional toroidal mesh in two particular cases.

4. Generating algorithm

We present a simple recursive algorithm to generate all graphs G with the minimum number of edges such that $\Gamma(G) = k$. The main idea is the following: we start from a tree with $2k-1$ vertices and we join together some vertices having the same color. By computing all the possible groupings, we find a set of graphs with a Grundy number equals to k . Recursively we start again from each graph of this set while the computation can be done.

5. Conclusion

In this article, we first positioned the Grundy number of a graph G compared to other graph parameters (stability number, complement graph of G). Then we presented several bounds and values for the Grundy number of the Cartesian product of two graphs. In particular we studied the Cartesian product of a bipartite graph by a path or a cycle, a bipartite graph by a complete graph and a complete graph by any graph G . Then, we deduced exact values for the Grundy number of a n -dimensional mesh and particular cases of a n -dimensional toroidal mesh.

Thus, for every graph H containing an induced subgraph G presented in this paper, Fact 1 shows that $\Gamma(H) \geq \Gamma(G)$ and gives a proper coloring to H .

ACKNOWLEDGMENT

I think Mr. Brice Effantin for his collaboration

REFERENCES

[AMC94] A. McRae, NP-completeness Proof for Grundy Coloring, unpublished manuscript 1994. Personal communication.

[BEH02] B. Effantin, H.kheddouci Grundy number of graph, Mathematical Journal 44 (2002) 103-115.

[CHS79] C. A. Christen and S. M. Selkow, Some perfect coloring properties of graphs, Journal of Combinatorial Theory B27 (1979) 49-59.

[CGH03] C. Germain and H. Kheddouci, Grundy coloring for power caterpillars, Proceedings of International Optimization Conference INOC 2003, 243-247, October 2003 Evry/Paris France.

[CGH06] C. Germain and H. Kheddouci, Grundy coloring of powers of graphs, to appear in Discrete Mathematics 2006.

[CPR02] C. Parks and J. Rhyne, Grundy Coloring for Chessboard Graphs, Seventh North Carolina Mini-Conference on Graph Theory, Combinatorics, and Computing (2002).

[CWE97] C. Y. Chang, W. E. Clark and E.O. Hare, Domination Numbers of Complete Grid Graphs, I, Ars Combinatoria 36 (1994) 97-111.11

[FSA03] F. Ruskey and J. Sawada, Bent Hamilton Cycles in d -Dimensional Grid Graphs, the Electronic Journal of Combinatorics 10 (2003) #R1.

[JAP97] J.A. Telle, A. Proskurowski, Algorithms for vertex partitioning problems on partial k -trees, SIAM Journal on Discrete Mathematics 10(4) (1997) 529-550.

[JHW02] J.D. Horton and W.D. Wallis, Factoring the cartesian product of a cubic graph and a triangle, Discrete Mathematics, 259, Issues 1-3 (2002) 137-146.

[JHK00] J.E. Dunbar, S.M. Hedetniemi, S.T. Hedetniemi, D.P. Jacobs, J. Knisely, R.C. Laskar and D.F. Rall, Fall Colorings of Graphs, Journal of Combinatorial Mathematics and Combinatorial Computing 33 (2000) 257-273.

[MKA02] M. Kouider and M. Maheo, Some bound for the b -chromatic number of a graph, Discrete Mathematics, 256 (2002) 267-277.

[NCK94] N. Cizek and S. Klavzar, on the chromatic number of the lexicographic product and the cartesian sum of graphs, Discrete Mathematics, 134, Issues 1-3 (1994) 17-24.

[SGR02] S. Gravier, Total domination number of grid graphs, *Discrete Applied Mathematics*, 121, Issues 1-3 (2002) 119-128.

[SST84] S. M. Hedetniemi, S. T. Hedetniemi and T. Beyer, A linear algorithm for the Grundy (coloring) number of a tree, *Congr. Numer.* 36 (1982) 351-363.

[XZH92] X. Zhu, Star chromatic numbers and products of graphs, *Journal of Graph Theory* 16 (1992) 557-569.

[XZH02] X. Zhu, The fractional chromatic number of the direct product of graphs, *Glasgow Mathematical Journal* 44 (2002) 103-115.