

GENERALIZATIONS OF ANDREWS' CURIOUS IDENTITIES

CHUANAN WEI^A, XIAOXIA WANG^B

^A*Department of Information Technology
Hainan Medical College, Haikou 571199, China*

^B*Department of Mathematics
Shanghai University, Shanghai 200444, China*

ABSTRACT. According to the method of series rearrangement, we establish two generalizations of Andrews' curious q -series identity with an extra integer parameter. The limiting cases of them produce two extensions of Andrews' curious ${}_3F_2(\frac{3}{4})$ -series identity with an additional integer parameter. Meanwhile, several related results are also given.

1. INTRODUCTION

For a complex variable x , define the shifted-factorial by

$$(x)_n = \begin{cases} \prod_{k=0}^{n-1} (x+k), & n > 0; \\ 1, & n = 0; \\ \frac{(-1)^n}{\prod_{k=1}^n (k-x)}, & n < 0. \end{cases}$$

For simplifying the expressions, we shall use the symbol:

$$\left[\begin{matrix} a, & b, & \dots, & c \\ \alpha, & \beta, & \dots, & \gamma \end{matrix} \right]_n = \frac{(a)_n (b)_n \dots (c)_n}{(\alpha)_n (\beta)_n \dots (\gamma)_n}.$$

Following Bailey [3], define the hypergeometric series by

$${}_{1+r}F_s \left[\begin{matrix} a_0, & a_1, & \dots, & a_r \\ b_1, & \dots, & b_s \end{matrix} \middle| z \right] = \sum_{k=0}^{\infty} \frac{(a_0)_k (a_1)_k \dots (a_r)_k}{k! (b_1)_k \dots (b_s)_k} z^k,$$

where $\{a_i\}_{i \geq 0}$ and $\{b_j\}_{j \geq 1}$ are complex parameters such that no zero factors appear in the denominators of the summand on the right hand side. Then the curious ${}_3F_2(\frac{3}{4})$ -series identity due to Andrews [1, Equation (22b)] reads as

$${}_3F_2 \left[\begin{matrix} -n, & a, & 3a+n \\ \frac{3a}{2}, & \frac{1+3a}{2} \end{matrix} \middle| \frac{3}{4} \right] = \left[\begin{matrix} \frac{1}{3}, & \frac{2}{3} \\ \frac{1}{3}+a, & \frac{2}{3}+a \end{matrix} \right]_{\frac{n}{3}} \chi(n), \tag{1}$$

where the symbol $\chi(n)$ has been offered by

$$\chi(n) = \begin{cases} 1, & n = 3m; \\ 0, & \text{otherwise.} \end{cases}$$

The reversal of it is the ${}_3F_2(\frac{4}{3})$ -series identity:

$${}_3F_2 \left[\begin{matrix} -n, & \frac{3b}{2}, & \frac{1+3b}{2} \\ 3b, & b - \frac{n-2}{3} \end{matrix} \middle| \frac{4}{3} \right] = \left[\begin{matrix} \frac{1}{3}, & \frac{2}{3} \\ \frac{1}{3}-b, & \frac{2}{3}+b \end{matrix} \right]_{\frac{n}{3}} \chi(n). \tag{2}$$

2010 Mathematics Subject Classification: Primary 05A19 and Secondary 33C20.

Key words and phrases. Hypergeometric series; q -Series; Series rearrangement.

Email addresses: weichuanan78@163.com (C. Wei), xwang 913@126.com (X. Wang).

Some recent research work for hypergeometric series should be mentioned. In accordance with inversion techniques, Chen and Chu [4, 5] derived many ${}_3F_2(\frac{4}{3})$ -series identities related to (2). According to series rearrangement, Chu [6], Lavoie [7] and Lavoie et al. [8] explored ${}_3F_2(1)$ -series identities.

For two complex numbers x and q , define the q -shifted factorial by

$$(x; q)_n = \begin{cases} \prod_{i=0}^{n-1} (1 - xq^i), & n > 0; \\ 1, & n = 0; \\ \frac{1}{\prod_{j=n}^{-1} (1 - xq^j)}, & n < 0. \end{cases}$$

The fraction form of it reads as

$$\left[\begin{matrix} a, & b, & \cdots, & c \\ \alpha, & \beta, & \cdots, & \gamma \end{matrix} \middle| q \right]_n = \frac{(a; q)_n (b; q)_n \cdots (c; q)_n}{(\alpha; q)_n (\beta; q)_n \cdots (\gamma; q)_n}.$$

Following Gasper and Rahman [9], define the q -series by

$${}_{1+r}\phi_s \left[\begin{matrix} a_0, & a_1, & \cdots, & a_r \\ b_1, & \cdots, & b_s \end{matrix} \middle| q; z \right] = \sum_{k=0}^{\infty} \left[\begin{matrix} a_0, & a_1, & \cdots, & a_r \\ q, & b_1, & \cdots, & b_s \end{matrix} \middle| q \right]_k z^k,$$

where $\{a_i\}_{i \geq 0}$ and $\{b_j\}_{j \geq 1}$ are complex parameters such that no zero factors appear in the denominators of the summand on the right hand side. Then the terminating ${}_6\phi_5$ -series identity (cf. Gasper and Rahman [9, p. 42]) can be expressed as

$${}_6\phi_5 \left[\begin{matrix} a, & q\sqrt{a}, & -q\sqrt{a}, & b, & c, & q^{-\ell} \\ \sqrt{a}, & -\sqrt{a}, & qa/b, & qa/c, & q^{1+\ell}a \end{matrix} \middle| q; \frac{q^{1+\ell}a}{bc} \right] = \left[\begin{matrix} qa, & qa/bc \\ qa/b, & qa/c \end{matrix} \middle| q \right]_{\ell}. \quad (3)$$

Recall a curious q -series identity due to Andrews [2, Equation (4.7)]:

$$\sum_{k=0}^n (a; q^3)_k \left[\begin{matrix} q^{-n}, & q^n a \\ q, & \sqrt{a}, & -\sqrt{a}, & \sqrt{qa}, & -\sqrt{qa} \end{matrix} \middle| q \right]_k q^k = a^{n/3} \left[\begin{matrix} q, & q^2 \\ qa, & q^2 a \end{matrix} \middle| q^3 \right]_{\frac{n}{3}} \chi(n). \quad (4)$$

The reversal of it can be stated as

$$\sum_{k=0}^n \frac{1}{(q^{2-n}b; q^3)_k} \left[\begin{matrix} q^{-n}, & \sqrt{b}, & -\sqrt{b}, & \sqrt{qb}, & -\sqrt{qb} \\ q, & b \end{matrix} \middle| q \right]_k q^k = \left[\begin{matrix} q, & q^2 \\ q/b, & q^2 b \end{matrix} \middle| q^3 \right]_{\frac{n}{3}} \chi(n). \quad (5)$$

Inspired by the work of [4]-[8], we shall establish two generalizations of (4), which involve two extensions of (1), by means of series rearrangement in section 2. The reversal of them creates two generalizations of (5), which involve two extensions of (2), in section 3.

2. GENERALIZATIONS OF ANDREWS' CURIOUS IDENTITIES

Theorem 1. *For a nonnegative integer ℓ and a complex number a , there holds*

$$\begin{aligned} & \sum_{k=0}^n (a; q^3)_k \left[\begin{matrix} q^{-n}, & q^{n-\ell} a \\ q, & \sqrt{a}, & -\sqrt{a}, & \sqrt{qa}, & -\sqrt{qa} \end{matrix} \middle| q \right]_k q^k = \left[\begin{matrix} q^{n-\ell} a \\ q^{2n-\ell} a \end{matrix} \middle| q \right]_{\ell} \\ & \times \sum_{i=0}^{\ell} a^{\frac{n+2i}{3}} q^{(\ell+2n-i)i} \frac{a - q^{2i-2n}}{a - q^{-2n}} \left[\begin{matrix} q^{-\ell}, & q^{-n}, & q^{-2n}/a \\ q, & q^{1+\ell-2n}/a, & q^{n-i} a \end{matrix} \middle| q \right]_i \\ & \times \left[\begin{matrix} q, & q^2 \\ qa, & q^2 a \end{matrix} \middle| q^3 \right]_{\frac{n-i}{3}} \chi(n-i). \end{aligned}$$

Proof. Letting $a \rightarrow q^{-2n}/a$, $b \rightarrow q^{k-n}$ and $c \rightarrow 0$ for (3), we get the equation:

$$\left[\begin{matrix} q^{n-\ell+k} a \\ q^{2n-\ell} a \end{matrix} \middle| q \right]_{\ell} \sum_{i=0}^{\ell} a^i q^{(\ell+2n-i)i} \frac{a - q^{2i-2n}}{a - q^{-2n}} \left[\begin{matrix} q^{-\ell}, & q^{k-n}, & q^{-2n}/a \\ q, & q^{1+\ell-2n}/a, & q^{n+k-i} a \end{matrix} \middle| q \right]_i = 1.$$

Then there is the following relation:

$$\begin{aligned} & \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{qa}, -\sqrt{qa} \mid q \right]_k q^k \\ &= \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{qa}, -\sqrt{qa} \mid q \right]_k q^k \\ & \times \left[\begin{matrix} q^{n-\ell+k} a \\ q^{2n-\ell} a \end{matrix} \mid q \right]_{\ell} \sum_{i=0}^{\ell} a^i q^{(\ell+2n-i)i} \frac{a - q^{2i-2n}}{a - q^{-2n}} \left[\begin{matrix} q^{-\ell}, q^{k-n}, q^{-2n}/a \\ q, q^{1+\ell-2n}/a, q^{n+k-i} a \end{matrix} \mid q \right]_i. \end{aligned}$$

Interchange the summation order for the last double sum to achieve

$$\begin{aligned} & \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{qa}, -\sqrt{qa} \mid q \right]_k q^k \\ &= \left[\begin{matrix} q^{n-\ell} a \\ q^{2n-\ell} a \end{matrix} \mid q \right]_{\ell} \sum_{i=0}^{\ell} a^i q^{(\ell+2n-i)i} \frac{a - q^{2i-2n}}{a - q^{-2n}} \left[\begin{matrix} q^{-\ell}, q^{-n}, q^{-2n}/a \\ q, q^{1+\ell-2n}/a, q^{n-i} a \end{matrix} \mid q \right]_i \\ & \times \sum_{k=0}^{n-i} (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{qa}, -\sqrt{qa} \mid q \right]_k q^k. \end{aligned}$$

Calculating the series on the last line by (4), we attain Theorem 1 to complete the proof. \square

When $\ell = 0$, Theorem 1 reduces to (4) exactly. Other two examples are displayed as follows.

Example 1 ($\ell = 1$ in Theorem 1).

$$\begin{aligned} & \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{qa}, -\sqrt{qa} \mid q \right]_k q^k \\ &= \begin{cases} \frac{a^m(1-aq^{-1})}{1-aq^{6m-1}} \left[\begin{matrix} q, q^2 \\ qa, q^{-1}a \end{matrix} \mid q^3 \right]_m, & n = 3m; \\ \frac{a^{m+1}q^{3m}(1-q)}{1-aq^{6m+1}} \left[\begin{matrix} q^2, q^4 \\ qa, q^2a \end{matrix} \mid q^3 \right]_m, & n = 1 + 3m; \\ 0, & n = 2 + 3m. \end{cases} \end{aligned}$$

Example 2 ($\ell = 2$ in Theorem 1).

$$\begin{aligned} & \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{qa}, -\sqrt{qa} \mid q \right]_k q^k \\ &= \begin{cases} \frac{a^m(1-aq^{-2})(1-aq^{-1})}{(1-aq^{6m-2})(1-aq^{6m-1})} \left[\begin{matrix} q, q^2 \\ q^{-1}a, q^{-2}a \end{matrix} \mid q^3 \right]_m, & n = 3m; \\ \frac{a^{m+1}q^{3m-1}(1-q^2)(1-aq^{-1})}{(1-aq^{6m-1})(1-aq^{6m+1})} \left[\begin{matrix} q^2, q^4 \\ q^{-1}a, qa \end{matrix} \mid q^3 \right]_m, & n = 1 + 3m; \\ \frac{a^{m+2}q^{6m}(1-q)(1-q^2)}{(1-aq^{6m+1})(1-aq^{6m+2})} \left[\begin{matrix} q^4, q^5 \\ qa, q^2a \end{matrix} \mid q^3 \right]_m, & n = 2 + 3m. \end{cases} \end{aligned}$$

Performing the replacement $a \rightarrow q^{3a}$ in Theorem 1 and then letting $q \rightarrow 1$, we obtain the following equation.

Theorem 2. For a nonnegative integer ℓ and a complex number a , there holds

$$\begin{aligned} & {}_3F_2 \left[-n, \quad a, \quad 3a+n-\ell \mid \frac{3}{4} \right] = \left[\begin{matrix} 1-n-3a \\ 1-2n-3a \end{matrix} \right]_{\ell} \\ & \times \sum_{i=0}^{\ell} (-1)^i \frac{3a+2n-2i}{3a+2n} \left[\begin{matrix} -\ell, -n, -2n-3a \\ 1, 1-n-3a, 1+\ell-2n-3a \end{matrix} \right]_i \\ & \times \left[\begin{matrix} \frac{1}{3}, \quad \frac{2}{3} \\ \frac{1}{3}+a, \quad \frac{2}{3}+a \end{matrix} \right]_{\frac{n-i}{3}} \chi(n-i). \end{aligned}$$

When $\ell = 0$, Theorem 2 reduces to (1) exactly. Other two examples are laid out as follows.

Example 3 ($\ell = 1$ in Theorem 2).

$$\begin{aligned} & {}_3F_2 \left[-n, \quad a, \quad 3a+n-1 \mid \frac{3}{4} \right] \\ & = \begin{cases} \frac{3a-1}{3a+6m-1} \left[\begin{matrix} \frac{1}{3}, \quad \frac{2}{3} \\ -\frac{1}{3}+a, \quad \frac{1}{3}+a \end{matrix} \right]_m, & n=3m; \\ \frac{1}{3a+6m+1} \left[\begin{matrix} \frac{2}{3}, \quad \frac{4}{3} \\ \frac{1}{3}+a, \quad \frac{2}{3}+a \end{matrix} \right]_m, & n=1+3m; \\ 0, & n=2+3m. \end{cases} \end{aligned}$$

Example 4 ($\ell = 2$ in Theorem 2).

$$\begin{aligned} & {}_3F_2 \left[-n, \quad a, \quad 3a+n-2 \mid \frac{3}{4} \right] \\ & = \begin{cases} \frac{(3a-2)(3a-1)}{(3a+6m-2)(3a+6m-1)} \left[\begin{matrix} \frac{1}{3}, \quad \frac{2}{3} \\ -\frac{1}{3}+a, \quad -\frac{2}{3}+a \end{matrix} \right]_m, & n=3m; \\ \frac{2(3a-1)}{(3a+6m-1)(3a+6m+1)} \left[\begin{matrix} \frac{2}{3}, \quad \frac{4}{3} \\ -\frac{1}{3}+a, \quad \frac{1}{3}+a \end{matrix} \right]_m, & n=1+3m; \\ \frac{2}{(3a+6m+1)(3a+6m+2)} \left[\begin{matrix} \frac{4}{3}, \quad \frac{5}{3} \\ \frac{1}{3}+a, \quad \frac{2}{3}+a \end{matrix} \right]_m, & n=2+3m. \end{cases} \end{aligned}$$

Theorem 3. For a nonnegative integer ℓ and a complex number a , there holds

$$\begin{aligned} & \sum_{k=0}^n (a; q^3)_k \left[\begin{matrix} q^{-n}, q^{n-\ell} a \\ q, \sqrt{a}, -\sqrt{a}, \sqrt{q^{1-2\ell} a}, -\sqrt{qa} \end{matrix} \mid q \right]_k q^k = \left[\begin{matrix} q^{n-\ell} a, q^{\frac{1}{2}+n-\ell} \sqrt{a} \\ q^{2n-\ell} a, q^{\frac{1}{2}-\ell} \sqrt{a} \end{matrix} \mid q \right]_{\ell} \\ & \times \sum_{i=0}^{\ell} a^{\frac{2n-5i}{6}} q^{\frac{5}{2}+\ell} \frac{1-q^{2n-2i} a}{1-q^{2n} a} \left[\begin{matrix} q^{-\ell}, q^{-n}, q^{-2n}/a \\ q, q^{1+\ell-2n}/a, q^{1-n}/a \end{matrix} \mid q \right]_i \\ & \times \left[\begin{matrix} q, q^2 \\ qa, q^2 a \end{matrix} \mid q^3 \right]_{\frac{n-i}{3}} \chi(n-i). \end{aligned}$$

Proof. Letting $a \rightarrow q^{-2n}/a$, $b \rightarrow q^{k-n}$ and $c \rightarrow q^{\frac{1}{2}-n}/\sqrt{a}$ for (3), the resulting equation reads as

$$\begin{aligned} & \left[\begin{matrix} q^{n-\ell} a, q^{\frac{1}{2}+n-\ell} \sqrt{a} \\ q^{2n-\ell} a, q^{\frac{1}{2}+k-\ell} \sqrt{a} \end{matrix} \mid q \right]_{\ell} \sum_{i=0}^{\ell} \frac{q^{(5/2+\ell)i}}{a^{i/2}} \frac{1-q^{2n-2i} a}{1-q^{2n} a} \left[\begin{matrix} q^{-\ell}, q^{k-n}, q^{-2n}/a \\ q, q^{1+\ell-2n}/a, q^{1-n}/a \end{matrix} \mid q \right]_i \\ & \times \left[\begin{matrix} q^{n-\ell+k} a \\ q^{n-\ell} a \end{matrix} \mid q \right]_{\ell-i} = 1. \end{aligned}$$

Then we can proceed as follows:

$$\begin{aligned}
& \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{q^{1-2\ell}a}, -\sqrt{qa} \mid q \right]_k q^k \\
&= \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{q^{1-2\ell}a}, -\sqrt{qa} \mid q \right]_k q^k \\
&\times \left[q^{n-\ell}a, q^{\frac{1}{2}+n-\ell}\sqrt{a} \mid q \right]_{\ell} \sum_{i=0}^{\ell} \frac{q^{(5/2+\ell)i}}{a^{i/2}} \frac{1-q^{2n-2i}a}{1-q^{2n}a} \left[q^{-\ell}, q^{k-n}, q^{-2n}/a \mid q \right]_i \\
&\times \left[q^{n-\ell+k}a \mid q \right]_{\ell-i}.
\end{aligned}$$

Interchange the summation order for the last double sum to get

$$\begin{aligned}
& \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{q^{1-2\ell}a}, -\sqrt{qa} \mid q \right]_k q^k = \left[q^{n-\ell}a, q^{\frac{1}{2}+n-\ell}\sqrt{a} \mid q \right]_{\ell} \\
&\times \sum_{i=0}^{\ell} \frac{q^{(5/2+\ell)i}}{a^{i/2}} \frac{1-q^{2n-2i}a}{1-q^{2n}a} \left[q^{-\ell}, q^{-n}, q^{-2n}/a \mid q \right]_i \\
&\times \sum_{k=0}^{n-i} (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{qa}, -\sqrt{qa} \mid q \right]_k q^k.
\end{aligned}$$

Evaluating the series on the last line by (4), we achieve Theorem 3 to complete the proof. \square

When $\ell = 0$, Theorem 3 also reduces to (4) exactly. Other two examples are displayed as follows.

Example 5 ($\ell = 1$ in Theorem 3).

$$\begin{aligned}
& \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{q^{-1}a}, -\sqrt{qa} \mid q \right]_k q^k \\
&= \begin{cases} \frac{a^m(\sqrt{q}+\sqrt{a})}{\sqrt{q}+q^{3m}\sqrt{a}} \left[q, q^2 \mid q^3 \right]_m, & n = 3m; \\ \frac{a^{m+1/2}(q-1)}{(\sqrt{q}-\sqrt{a})(1+q^{3m+1/2}\sqrt{a})} \left[q^2, q^4 \mid q^3 \right]_m, & n = 1 + 3m; \\ 0, & n = 2 + 3m. \end{cases}
\end{aligned}$$

Example 6 ($\ell = 2$ in Theorem 3).

$$\begin{aligned}
& \sum_{k=0}^n (a; q^3)_k \left[q, \sqrt{a}, -\sqrt{a}, \sqrt{q^{-3}a}, -\sqrt{qa} \mid q \right]_k q^k \\
&= \begin{cases} \frac{a^m(q^2-a)(\sqrt{q}+\sqrt{a})(1-q^{3m-3/2}\sqrt{a})}{(q^2-q^{6m}a)(1-q^{-3/2}\sqrt{a})(\sqrt{q}+q^{3m}\sqrt{a})} \left[q, q^2 \mid q^3 \right]_m, & n = 3m; \\ \frac{a^{m+1/2}(q^2-1)(\sqrt{q}+\sqrt{a})}{(q^2-\sqrt{q}a)(1+q^{3m-1/2}\sqrt{a})(1+q^{3m+1/2}\sqrt{a})} \left[q^2, q^4 \mid q^3 \right]_m, & n = 1 + 3m; \\ \frac{a^{m+1}(1-q)(1-q^2)(1-q^{3m+3/2}\sqrt{a})}{(\sqrt{q}-\sqrt{a})(\sqrt{q^3}-\sqrt{a})(1+q^{3m+1/2}\sqrt{a})(1-q^{6m+2}a)} \left[q^4, q^5 \mid q^3 \right]_m, & n = 2 + 3m. \end{cases}
\end{aligned}$$

Employing the substitution $a \rightarrow q^{3a}$ in Theorem 3 and then letting $q \rightarrow 1$, we obtain the following equation.

Theorem 4. For a nonnegative integer ℓ and a complex number a , there holds

$$\begin{aligned} & {}_3F_2 \left[-n, \quad a, \quad 3a+n-\ell \mid \frac{3}{4} \right] = \left[\begin{matrix} 1-3a-n, \frac{1-3a}{2}-n \\ 1-3a-2n, \frac{1-3a}{2} \end{matrix} \right]_\ell \\ & \times \sum_{i=0}^{\ell} \frac{3a+2n-2i}{3a+2n} \left[\begin{matrix} -\ell, -n, -2n-3a \\ 1, 1-n-3a, 1+\ell-2n-3a \end{matrix} \right]_i \\ & \times \left[\begin{matrix} \frac{1}{3}, \quad \frac{2}{3} \\ \frac{1}{3}+a, \quad \frac{2}{3}+a \end{matrix} \right]_{\frac{n-i}{3}} \chi(n-i). \end{aligned}$$

When $\ell = 0$, Theorem 4 also reduces to (1) exactly. Other two examples are laid out as follows.

Example 7 ($\ell = 1$ in Theorem 4).

$$\begin{aligned} & {}_3F_2 \left[-n, \quad a, \quad 3a+n-1 \mid \frac{3}{4} \right] \\ & = \begin{cases} \left[\begin{matrix} \frac{1}{3}, \quad \frac{2}{3} \\ -\frac{1}{3}+a, \quad \frac{1}{3}+a \end{matrix} \right]_m, & n = 3m; \\ \frac{1}{1-3a} \left[\begin{matrix} \frac{2}{3}, \quad \frac{4}{3} \\ \frac{1}{3}+a, \quad \frac{2}{3}+a \end{matrix} \right]_m, & n = 1+3m; \\ 0, & n = 2+3m. \end{cases} \end{aligned}$$

Example 8 ($\ell = 2$ in Theorem 4).

$$\begin{aligned} & {}_3F_2 \left[-n, \quad a, \quad 3a+n-2 \mid \frac{3}{4} \right] \\ & = \begin{cases} \frac{(3a-2)(a+2m-1)}{(a-1)(3a+6m-2)} \left[\begin{matrix} \frac{1}{3}, \quad \frac{2}{3} \\ -\frac{1}{3}+a, \quad -\frac{2}{3}+a \end{matrix} \right]_m, & n = 3m; \\ \frac{2}{3(1-a)} \left[\begin{matrix} \frac{2}{3}, \quad \frac{4}{3} \\ -\frac{1}{3}+a, \quad \frac{1}{3}+a \end{matrix} \right]_m, & n = 1+3m; \\ \frac{2(a+2m+1)}{(a-1)(3a-1)(3a+6m+2)} \left[\begin{matrix} \frac{4}{3}, \quad \frac{5}{3} \\ \frac{1}{3}+a, \quad \frac{2}{3}+a \end{matrix} \right]_m, & n = 2+3m. \end{cases} \end{aligned}$$

3. SEVERAL IDENTITIES FROM REVERSAL

Performing the replacement $k \rightarrow n-k$ and $a \rightarrow q^{1-2n}/b$ in Theorem 1, we derive the following equation.

Theorem 5. For a nonnegative integer ℓ and a complex number b , there holds

$$\begin{aligned} & \sum_{k=0}^n \frac{1}{(q^{2-n}b; q^3)_k} \left[\begin{matrix} q^{-n}, \sqrt{b}, -\sqrt{b}, \sqrt{qb}, -\sqrt{qb} \\ q, q^\ell b \end{matrix} \mid q \right]_k q^{(1+\ell)k} = \frac{(q^{1-2n}/b; q)_n}{(q^{1-2n}/b; q^3)_n} \\ & \times \sum_{i=0}^{\ell} (-1)^n \frac{q^{\{(4+4n+6\ell-6i)i-n-n^2\}/6}}{b^{(n+2i)/3}} \frac{q-q^{2i}b}{q-b} \left[\begin{matrix} q^{-\ell}, q^{-n}, b/q \\ q, q^\ell b, q^{1-n-i}/b \end{matrix} \mid q \right]_i \\ & \times \left[\begin{matrix} q, q^2 \\ q^{2-2n}/b, q^{3-2n}/b \end{matrix} \mid q^3 \right]_{\frac{n-i}{3}} \chi(n-i). \end{aligned}$$

When $\ell = 0$, Theorem 5 reduces to (5) exactly. Other two examples are displayed as follows.

Example 9 ($\ell = 1$ in Theorem 5).

$$\begin{aligned} & \sum_{k=0}^n \frac{1}{(q^{2-n}b; q^3)_k} \left[\begin{matrix} q^{-n}, \sqrt{b}, -\sqrt{b}, \sqrt{qb}, -\sqrt{qb} \\ q, qb \end{matrix} \middle| q \right]_k q^{2k} \\ &= \begin{cases} \left[\begin{matrix} q, q^2 \\ q/b, q^2b \end{matrix} \middle| q^3 \right]_m, & n = 3m; \\ \frac{1-q}{1-qb} \left[\begin{matrix} q^2, q^4 \\ q^2/b, q^4b \end{matrix} \middle| q^3 \right]_m, & n = 1 + 3m; \\ 0, & n = 2 + 3m. \end{cases} \end{aligned}$$

Example 10 ($\ell = 2$ in Theorem 5).

$$\begin{aligned} & \sum_{k=0}^n \frac{1}{(q^{2-n}b; q^3)_k} \left[\begin{matrix} q^{-n}, \sqrt{b}, -\sqrt{b}, \sqrt{qb}, -\sqrt{qb} \\ q, q^2b \end{matrix} \middle| q \right]_k q^{3k} \\ &= \begin{cases} \left[\begin{matrix} q, q^2 \\ q/b, q^2b \end{matrix} \middle| q^3 \right]_m, & n = 3m; \\ \frac{1-q^2}{1-q^2b} \left[\begin{matrix} q^2, q^4 \\ q^2/b, q^4b \end{matrix} \middle| q^3 \right]_m, & n = 1 + 3m; \\ \frac{(1-q)(1-q^2)}{(1-q^2b)(1-q^3b)} \left[\begin{matrix} q^4, q^5 \\ q^3/b, q^6b \end{matrix} \middle| q^3 \right]_m, & n = 2 + 3m. \end{cases} \end{aligned}$$

Employing the substitution $a \rightarrow q^{3a}$ in Theorem 5 and then letting $q \rightarrow 1$, we get the following equation.

Theorem 6. For a nonnegative integer ℓ and a complex number a , there holds

$$\begin{aligned} & {}_3F_2 \left[\begin{matrix} -n, & \frac{3b}{2}, & \frac{1+3b}{2} \\ 3b + \ell, & b - \frac{n-2}{3} \end{matrix} \middle| \frac{4}{3} \right] = \left[\begin{matrix} 3b + n \\ \frac{1-2n}{3} - b \end{matrix} \right]_n \\ & \times \sum_{i=0}^{\ell} \frac{(-1)^i}{3^n} \frac{3b-1+2i}{3b-1} \left[\begin{matrix} -\ell, -n, 3b-1 \\ 1, 3b+n, 3b+\ell \end{matrix} \right]_i \\ & \times \left[\begin{matrix} \frac{1}{3}, & \frac{2}{3} \\ \frac{2-2n-3b}{3}, & \frac{3-2n-3b}{3} \end{matrix} \right]_{\frac{n-i}{3}} \chi(n-i). \end{aligned}$$

When $\ell = 0$, Theorem 5 reduces to (2) exactly. Other two examples are laid out as follows.

Example 11 ($\ell = 1$ in Theorem 5: Chen and Chu [5, Theorem 4]).

$$\begin{aligned} & {}_3F_2 \left[\begin{matrix} -n, & \frac{3b}{2}, & \frac{1+3b}{2} \\ 1 + 3b, & b - \frac{n-2}{3} \end{matrix} \middle| \frac{4}{3} \right] \\ &= \begin{cases} \left[\begin{matrix} \frac{1}{3}, & \frac{2}{3} \\ \frac{1}{3} - b, & \frac{2}{3} + b \end{matrix} \right]_m, & n = 3m; \\ \frac{1}{3b+1} \left[\begin{matrix} \frac{2}{3}, & \frac{4}{3} \\ \frac{2}{3} - b, & \frac{4}{3} + b \end{matrix} \right]_m, & n = 1 + 3m; \\ 0, & n = 2 + 3m. \end{cases} \end{aligned}$$

Example 12 ($\ell = 2$ in Theorem 5: Chen and Chu [5, Theorem 25]).

$$\begin{aligned} & {}_3F_2 \left[\begin{matrix} -n, & \frac{3b}{2}, & \frac{1+3b}{2} \\ 2+3b, & b - \frac{n-2}{3} \end{matrix} \middle| \frac{4}{3} \right] \\ &= \begin{cases} \left[\begin{matrix} \frac{1}{3}, & \frac{2}{3} \\ \frac{1}{3}-b, & \frac{2}{3}+b \end{matrix} \middle| q \right]_m, & n = 3m; \\ \frac{2}{3b+2} \left[\begin{matrix} \frac{2}{3}, & \frac{4}{3} \\ \frac{2}{3}-b, & \frac{4}{3}+b \end{matrix} \middle| q \right]_m, & n = 1+3m; \\ \frac{2}{3(b+1)(3b+2)} \left[\begin{matrix} \frac{4}{3}, & \frac{5}{3} \\ 1-b, & 2+b \end{matrix} \middle| q \right]_m, & n = 2+3m. \end{cases} \end{aligned}$$

Performing the replacement $k \rightarrow n - k$ and $a \rightarrow q^{1-2n}/b$ in Theorem 3, we deduce the following equation.

Theorem 7. For a nonnegative integer ℓ and a complex number b , there holds

$$\begin{aligned} & \sum_{k=0}^n \frac{1}{(q^{2-n}b; q^3)_k} \left[\begin{matrix} q^{-n}, q^\ell \sqrt{b}, -\sqrt{b}, \sqrt{qb}, -\sqrt{qb} \\ q, q^\ell b \end{matrix} \middle| q \right]_k q^k = \frac{(q^{1-2n}/b; q)_n}{(q^{1-2n}/b; q^3)_n} \\ & \times \sum_{i=0}^{\ell} (-1)^n \frac{q^{\{(10n+6\ell-2)i-n-n^2\}/6}}{b^{(2n-5i)/3}} \frac{q - q^{2i}b}{q - b} \left[\begin{matrix} q^{-\ell}, q^{-n}, b/q \\ q, q^\ell b, q^n b \end{matrix} \middle| q \right]_i \\ & \times \left[\begin{matrix} q, q^2 \\ q^{2-2n}/b, q^{3-2n}/b \end{matrix} \middle| q^3 \right]_{\frac{n-i}{3}} \chi(n-i). \end{aligned}$$

When $\ell = 0$, Theorem 7 reduces to (5) exactly. Other two examples are displayed as follows.

Example 13 ($\ell = 1$ in Theorem 7).

$$\begin{aligned} & \sum_{k=0}^n \frac{1}{(q^{2-n}b; q^3)_k} \left[\begin{matrix} q^{-n}, q\sqrt{b}, -\sqrt{b}, \sqrt{qb}, -\sqrt{qb} \\ q, qb \end{matrix} \middle| q \right]_k q^k \\ &= \begin{cases} \left[\begin{matrix} q, q^2 \\ q/b, q^2b \end{matrix} \middle| q^3 \right]_m, & n = 3m; \\ \frac{\sqrt{b}(q-1)}{1-qb} \left[\begin{matrix} q^2, q^4 \\ q^2/b, q^4b \end{matrix} \middle| q^3 \right]_m, & n = 1+3m; \\ 0, & n = 2+3m. \end{cases} \end{aligned}$$

Example 14 ($\ell = 2$ in Theorem 7).

$$\begin{aligned} & \sum_{k=0}^n \frac{1}{(q^{2-n}b; q^3)_k} \left[\begin{matrix} q^{-n}, q^2\sqrt{b}, -\sqrt{b}, \sqrt{qb}, -\sqrt{qb} \\ q, q^2b \end{matrix} \middle| q \right]_k q^k \\ &= \begin{cases} \left[\begin{matrix} q, q^2 \\ q/b, q^2b \end{matrix} \middle| q^3 \right]_m, & n = 3m; \\ \frac{\sqrt{b}(q^2-1)}{1-q^2b} \left[\begin{matrix} q^2, q^4 \\ q^2/b, q^4b \end{matrix} \middle| q^3 \right]_m, & n = 1+3m; \\ \frac{qb(1-q)(1-q^2)}{(1-q^2b)(1-q^3b)} \left[\begin{matrix} q^4, q^5 \\ q^3/b, q^6b \end{matrix} \middle| q^3 \right]_m, & n = 2+3m. \end{cases} \end{aligned}$$

Employing the substitution $a \rightarrow q^{3a}$ in Theorem 7 and then letting $q \rightarrow 1$, we obtain the following equation.

Theorem 8. For a nonnegative integer ℓ and a complex number a , there holds

$$\begin{aligned} & {}_3F_2 \left[-n, \frac{3b}{2} + \ell, \frac{1+3b}{2} \mid \frac{4}{3} \right] = \frac{1}{3^n} \left[\frac{3b+n}{\frac{1-2n}{3} - b} \right]_n \\ & \times \sum_{i=0}^{\ell} \frac{3b-1+2i}{3b-1} \left[\frac{-\ell, -n, 3b-1}{1, 3b+n, 3b+\ell} \right]_i \\ & \times \left[\frac{\frac{1}{3}, \frac{2}{3}}{\frac{2-2n-3b}{3}, \frac{3-2n-3b}{3}} \right]_{\frac{n-i}{3}} \chi(n-i). \end{aligned}$$

When $\ell = 0$, Theorem 8 reduces to (2) exactly. Other two examples are laid out as follows.

Example 15 ($\ell = 1$ in Theorem 8).

$$\begin{aligned} & {}_3F_2 \left[-n, \frac{1+3b}{2}, \frac{2+3b}{2} \mid \frac{4}{3} \right] \\ & = \begin{cases} \left[\frac{\frac{1}{3}, \frac{2}{3}}{\frac{1}{3}-b, \frac{2}{3}+b} \right]_m, & n = 3m; \\ \frac{-1}{3b+1} \left[\frac{\frac{2}{3}, \frac{4}{3}}{\frac{2}{3}-b, \frac{4}{3}+b} \right]_m, & n = 1+3m; \\ 0, & n = 2+3m. \end{cases} \end{aligned}$$

Example 16 ($\ell = 2$ in Theorem 8).

$$\begin{aligned} & {}_3F_2 \left[-n, \frac{1+3b}{2}, \frac{4+3b}{2} \mid \frac{4}{3} \right] \\ & = \begin{cases} \left[\frac{\frac{1}{3}, \frac{2}{3}}{\frac{1}{3}-b, \frac{2}{3}+b} \right]_m, & n = 3m; \\ \frac{-2}{3b+2} \left[\frac{\frac{2}{3}, \frac{4}{3}}{\frac{2}{3}-b, \frac{4}{3}+b} \right]_m, & n = 1+3m; \\ \frac{2}{3(b+1)(3b+2)} \left[\frac{\frac{4}{3}, \frac{5}{3}}{1-b, 2+b} \right]_m, & n = 2+3m. \end{cases} \end{aligned}$$

Although Theorems 6 and 8 can produce countless ${}_3F_2(\frac{4}{3})$ -series identities related to (2) with the change of ℓ , many known results due to Chu and Chen [4, 5] can't be covered by them.

Acknowledgments

The work is supported by the Natural Science Foundations of China (Nos. 11301120, 11201241 and 11201291).

REFERENCES

- [1] G.E. Andrews, Applications of basic hypergeometric functions, SIAM Rev. 16 (1974), 441-484.
- [2] G.E. Andrews, Connection coefficient problems and partitions, Proc. Symposia in Pure Math. 34 (1979), 1-24.
- [3] W.N. Bailey, Generalized Hypergeometric Series, Cambridge University Press, Cambridge, 1935.
- [4] X. Chen, W. Chu, The Pfaff-Saalschütz theorem and terminating ${}_3F_2(\frac{4}{3})$ -series identities, J. Math. Anal. Appl. 380 (2011), 440-454.
- [5] X. Chen, W. Chu, Further ${}_3F_2(\frac{4}{3})$ -series via Gould-Hsu inversions, Integral Transforms Spec. Funct. 24 (2013), 441-469.

- [6] W. Chu, Analytical formulae for extended ${}_3F_2$ -series of Watson-Whipple-Dixon with two extra integer parameters, *Math. Comp.* 81 (2012), 467-479.
- [7] J. L. Lavoie, Some summation formulas for the series ${}_3F_2(1)$, *Math. Comp.* 49 (1987), 269-274.
- [8] J. L. Lavoie, F. Grondin, A.K. Rathie, Generalizations of Watson's theorem on the sum of a ${}_3F_2$, *Indian J. Math.* 34 (1992), 23-32.
- [9] G. Gasper, M. Rahman, *Basic Hypergeometric Series* (2nd edition), Cambridge University Press, Cambridge, 2004.