ON THE ESSENTIAL SPECTRUM OF TWO-DIMENSIONAL PAULI OPERATORS WITH REPULSIVE POTENTIALS

JOSEF MEHRINGER

ABSTRACT. We investigate the spectrum of the two-dimensional Pauli operator, describing a spin- $\frac{1}{2}$ particle in a magnetic field B, with a negative scalar potential V such that |V| grows at infinity. In particular, we obtain criteria for discrete and dense pure-point spectrum.

1. Introduction

For modeling the kinetic energy of a non-relativistic spin- $\frac{1}{2}$ particle in the plane, moving under a magnetic field B in the perpendicular direction to the plane, one uses the two-dimensional Pauli operator

$$H_{\mathbf{A}} := \left[\boldsymbol{\sigma} \cdot \left(-i \nabla - \mathbf{A} \right) \right]^2 = \left(-i \nabla - \mathbf{A} \right)^2 - \sigma_3 B \quad \text{on } L^2(\mathbb{R}^2, \mathbb{C}^2),$$

where **A** is a vector potential associated to B, i.e. $B = \operatorname{curl} \mathbf{A} := \partial_1 A_2 - \partial_1 A_1$. Here, $\boldsymbol{\sigma} = (\sigma_1, \sigma_2)$ and σ_3 are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -\mathrm{i} \\ \mathrm{i} & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

To study the behaviour of such spin- $\frac{1}{2}$ particles (e.g. electrons) in presence of an additional electric potential V, we investigate the spectrum of the operator

$$H := H_{\mathbf{A}} + V = \left[\boldsymbol{\sigma} \cdot \left(-i \nabla - \mathbf{A} \right) \right]^2 + V \quad \text{on } L^2(\mathbb{R}^2, \mathbb{C}^2).$$

If V is non-negative or decays at infinity (e.g. potentials with Coulomb singularities), spectral properties of the magnetic Schrödinger operator $(-i\nabla - \mathbf{A})^2 + V$, as well as of the Pauli operator $H_{\mathbf{A}} + V$ (in dimension d = 2 or 3) have been widely studied over the last decades (see, e.g. [2], [4] or [5] for a latest overview). In this article, instead, we want to point out some interesting features of the spectrum of H for potentials V tending to $-\infty$ as $|\mathbf{x}| \to \infty$. Since such scalar potentials result in an operator H, unbounded from below, it is necessary to discuss questions related to the self-adjointness of H. We emphasize that, since we also consider unbounded magnetic fields B, the self-adjointness of H cannot simply be reduced to the one of the magnetic Schrödinger operator.

One motivation for the following considerations is an observation made in [10] for the two-dimensional massless magnetic Dirac operator coupled to an electric potential V; There, an accumulation process of spectral points has been observed, governed by the ratio $|V^2/B|$ at infinity. This phenomenon can be ascribed to the non-confining effect of V in the case of the Dirac operator. Regarding the Pauli

Date: March 7, 2022.

 $^{2010\ \}textit{Mathematics Subject Classification}.\ \text{Primary 81Q10; Secondary 47A25, 81Q80}.$

Key words and phrases. Pauli operator, dense pure point spectrum.

operator, the influence of an additional scalar potential V on the spectrum $\sigma(H)$ depends crucially on the sign of V. For simplicity, we outline this dependence in the case of a constant magnetic field $B(x) = B_0$: a positive potential V growing at infinity always leads to discrete spectrum of the operator H, independently of the field strength B_0 (see e.g. [9]). Such trapping potentials only enhance a localization effect caused by B that generates eigenvalues and spectral gaps (proportional to B_0). If we instead consider negative potentials V, the situation is quite different since the particle lowers its energy by moving in regions where V is small. A scalar potential V converging to $-\infty$ as $|\mathbf{x}| \to \infty$ has therefore a delocalizing effect, i.e. the particle tends to escape any compact region of the plane. Our results show that such negative potentials V (describing for example constant radial fields) counteract the localizing effect of "hard" magnetic fields B, as they close spectral gaps induced by B:

- If V converges to $-\infty$, but remains small compared to B, the spectrum $\sigma(H)$ is discrete, i.e. it consists only of eigenvalues of finite multiplicity.
- If V is comparable to B, more precisely $|V| \approx 2B$ at infinity, points in the essential spectrum occur.
- If V overtakes B, more explicitly $|V/B| \to \infty$ as $|\mathbf{x}| \to \infty$ (at least along a path), the spectrum $\sigma(H)$ covers the whole real line.

One may compare the third claim with the result in [11] on $H_{\mathbf{A}}$ for decaying magnetic fields. The precise statements of the claims above are contained in Theorems 1–4 of Sect. 3. We remark that the case $|V/B| \to \infty$ as $|\mathbf{x}| \to \infty$ is treated by Theorems 3 and 4. Unlike Theorem 4, which is only valid for constant magnetic fields, Theorem 3 covers also non-constant fields B, but requires stronger constraints on the growth of V. Thus, the important case $B = B_0$ is adressed by two theorems. The ideas of the proofs of Theorem 1–3 originate from those used to prove the results in [10]. However, since we work with a second-order operator, the proofs are technically more laborious. Theorem 4 is based on a further construction of a Weyl sequence, obtained by treating V locally as a potential of a constant electric field. This is a refined ansatz compared to the method used for the proof of Theorem 3.

The organization of this article is as follows: In the next section some known facts about the Pauli operator are recapitulated. We present our precise results in Section 3, provided with some remarks and important applications. In Section 4 we give the proof of Theorem 1. The proofs of Theorems 2 and 3 are contained in Section 5, while the proof of Theorem 4 can be found in the last section. In the appendix, attached to the main text, we give a proof of the essential self-adjointness of the Pauli operator.

2. Basic properties of the Pauli operator

In this section we point out some basic facts about the Pauli operator and the massless Dirac operator $D_{\mathbf{A}}$, whose square equals $H_{\mathbf{A}}$. For a vector potential $\mathbf{A} \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ generating the field $B = \text{curl } \mathbf{A} \in C(\mathbb{R}^2, \mathbb{R})$, the Hamiltonian $D_{\mathbf{A}}$ is defined as the closure of the operator

(1)
$$\boldsymbol{\sigma} \cdot \left(-i \nabla - \mathbf{A} \right) = \begin{pmatrix} 0 & d^* \\ d & 0 \end{pmatrix} \quad \text{on } C_0^{\infty}(\mathbb{R}^2, \mathbb{C}^2),$$

which is essentially self-adjoint on the given core (see [3]). In particular, d and d^* can be seen as closed operators, i.e. we use the notation

(2)
$$d = \overline{-i \partial_1 - A_1 + i \left(-i \partial_2 - A_2\right)} \upharpoonright_{C_0^{\infty}(\mathbb{R}^2, \mathbb{C})}$$

and analogously for d^* . One observes that d, d^* satisfy the commutation relation

(3)
$$[d, d^*]\varphi := (dd^* - d^*d)\varphi = 2B\varphi \quad \text{for } \varphi \in C_0^\infty(\mathbb{R}^2, \mathbb{C}).$$

We can write

$$H_{\mathbf{A}} = D_{\mathbf{A}}^2 = \begin{pmatrix} d^*d & 0 \\ 0 & dd^* \end{pmatrix}$$
 on $C_0^{\infty}(\mathbb{R}^2, \mathbb{C}^2)$

and consider $H_{\mathbf{A}}$ as a self-adjoint operator on $\{\psi \in \mathcal{D}(D_{\mathbf{A}}) \mid D_{\mathbf{A}}\psi \in \mathcal{D}(D_{\mathbf{A}})\}$ given by the Friedrichs extension. The two components dd^* and d^*d of $H_{\mathbf{A}}$ are unitarily equivalent on the orthogonal complement of $\ker(H_{\mathbf{A}}) = \ker(D_{\mathbf{A}})$. To verify this we note first that, due to the matrix structure of $D_{\mathbf{A}}$, we have

(4)
$$\operatorname{sgn}(D_{\mathbf{A}}) := \frac{D_{\mathbf{A}}}{|D_{\mathbf{A}}|} = \begin{pmatrix} 0 & s^* \\ s & 0 \end{pmatrix}$$

on $\ker(D_{\mathbf{A}})^{\perp} = \ker(d)^{\perp} \oplus \ker(d^*)^{\perp}$. Since $\operatorname{sgn}(D_{\mathbf{A}})^2 = \operatorname{Id}$ on $\ker(D_{\mathbf{A}})^{\perp}$, the maps

(5)
$$s: \ker(d)^{\perp} \to \ker(d^*)^{\perp}, \qquad s^*: \ker(d^*)^{\perp} \to \ker(d)^{\perp}$$

are unitary and conjugated to each other. By the operator identity $H_{\bf A}=D_{\bf A}^2={\rm sgn}(D_{\bf A})D_{\bf A}^2{\rm sgn}(D_{\bf A})$ one concludes that

(6)
$$\begin{pmatrix} d^*d & 0 \\ 0 & dd^* \end{pmatrix} \varphi = \begin{pmatrix} s^*dd^*s & 0 \\ 0 & sd^*ds^* \end{pmatrix} \varphi$$

for any $\varphi = (\varphi_1, \varphi_2)^{\mathrm{T}}$ with $\varphi_1 \in \mathcal{D}(d^*d) \cap \ker(d)^{\perp}$ and $\varphi_2 \in \mathcal{D}(dd^*) \cap \ker(d^*)^{\perp}$. Hence, on $\ker(d)^{\perp}$ the operator d^*d is unitarily equivalent to dd^* (considered as an operator on $\ker(d^*)^{\perp}$). Let us denote the orthogonal projection on $\ker(D_{\mathbf{A}})$ by P_0 and the orthogonal projections on $\ker(d)$, $\ker(d^*)$ by π , π_* . Further, we set

$$P_0^{\perp} := \mathbb{1} - P_0, \quad \pi^{\perp} := \mathbb{1} - \pi, \quad \pi_*^{\perp} := \mathbb{1} - \pi_*.$$

To define our full Hamiltonian let $\mathbf{A} \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ and $B, V \in C(\mathbb{R}^2, \mathbb{R})$ be such that $B = \text{curl } \mathbf{A}$, then H is given by

$$H\varphi = \left[D_{\mathbf{A}}^2 + V\right]\varphi = \left[\left(-i\nabla - \mathbf{A}\right)^2 - \sigma_3 B + V\right]\varphi \quad \text{for } \varphi \in C_0^{\infty}(\mathbb{R}^2, \mathbb{C}^2).$$

In general, the closure of this densely defined operator is not self-adjoint without any restriction on the growth rate of V at infinity. However, there are conditions, very similar to those for the classical Schrödinger operator, to ensure essential self-adjointness.

Proposition 1. Let $B, V \in C^1(\mathbb{R}^2, \mathbb{R})$ and $\mathbf{A} \in C^2(\mathbb{R}^2, \mathbb{R}^2)$ with $B = \text{curl } \mathbf{A}$. In addition, assume that V fulfills the lower bound

(7)
$$V(\mathbf{x}) \geq -c|\mathbf{x}|^2 + d, \quad \mathbf{x} \in \mathbb{R}^2,$$

for some constants c > 0, $d \in \mathbb{R}$. Then H is essentially self-adjoint on $C_0^{\infty}(\mathbb{R}^2, \mathbb{C}^2)$.

Remark 1. Following the lines of the proof given in Appendix A, we see that the regularity condition on B,V can be relaxed to B,V $\in C^{\alpha}_{loc}(\mathbb{R}^2,\mathbb{R})$, i.e. both only need to be locally α -Hölder continuous. By a perturbation argument one can also see that it suffices to assume that B,V are C^{α}_{loc} outside some compact set $K \subset \mathbb{R}^2$, while inside K they only need to be continuous.

Remark 2. The self-adjoint operator given by Proposition 1 is locally compact, i.e. for any characteristic function $\chi_{B_R(0)}$ on the ball $B_R(0)$ with radius R, the operator $\chi_{B_R(0)}(H-i)^{-1}$ is compact.

Remark 3. Considering the case V=0, we obtain that $H_{\mathbf{A}}$, dd^* and d^*d are essentially self-adjoint on $C_0^{\infty}(\mathbb{R}^2,\mathbb{C}^2)$, respectively on $C_0^{\infty}(\mathbb{R}^2,\mathbb{C})$.

Note that (7) is the same lower bound on V as one needs for the (magnetic) Schrödinger operator to ensure the essential self-adjointness, whereas no restriction on the growth of B is necessary. The regularity conditions on V and A are quite strong compared to those for the magnetic Schrödinger operator (see [12]). The reason is that due to the lack of a diamagnetic inequality for H_A , one uses a direct argument that requires more regularity on the potentials A and V. The interesting question remains: Could one relax these conditions for the Pauli operator as in the case of the magnetic Schrödinger operator?

3. Main results

In this section we assume that B, V and \mathbf{A} satisfy the conditions of Proposition 1. It is easy to see that in the following results $B, V \in C^1(\mathbb{R}^2, \mathbb{R})$ can be relaxed to hold only outside some compact set $K \subset \mathbb{R}^2$ as in Remark 1.

Theorem 1. Assume that

(8)
$$V(\mathbf{x}) \longrightarrow -\infty \quad as \quad |\mathbf{x}| \to \infty,$$

(9)
$$\left| \frac{\nabla V(\mathbf{x})}{V(\mathbf{x})} \right| \longrightarrow 0 \quad as \quad |\mathbf{x}| \to \infty,$$

(10)
$$\limsup_{|\mathbf{x}| \to \infty} \left| \frac{V(\mathbf{x})}{2B(\mathbf{x})} \right| < 1.$$

Then $\sigma_{\rm ess}(H) = \emptyset$, i.e. H has purely discrete spectrum.

Condition (9) is a restriction on the growth rate of V and rather of technical necessity. The interplay between B and V (as mentioned in the introduction) is described by condition (10). Thus, it is worthwhile to investigate further the dependence of $\sigma(H)$ on this quotient:

One can easily observe that if the quotient of (10) surpasses the constant 1, the spectrum of H changes its character. To see this pick $\Omega \in \ker(d^*d)$, then

$$(dd^* + V)\Omega = (d^*d + 2B + V)\Omega \approx 0$$

if $2B \approx -V$. Therefore, if $\ker(d^*d)$ contains enough functions (which is the case for fields B bounded form below by some positive constant), we obtain points in the essential spectrum of H. One can even show that the condition $2B \approx -V$ (at infinity) does not need to hold globally for obtaining $\sigma_{\rm ess}(H) \neq \emptyset$. We demonstrate this for a certain class of fields B and potentials V.

Definition 1. A function $f: \mathbb{R}^2 \to \mathbb{R}$ varies with rate $\nu \in [0,1]$ on a set $X \subset \mathbb{R}^2$ if there is a constant C > 0 such that for all $\mathbf{x} \in X$ it holds that

$$|f(\mathbf{x} + \mathbf{y})| \le C|f(\mathbf{x})|,$$

whenever $\mathbf{y} \in \mathbb{R}^2$ satisfies $|\mathbf{y}| \leq \frac{1}{2} |\mathbf{x}|^{\nu}$

Note that functions of the form $f_1(\mathbf{x}) = c|\mathbf{x}|^s$ and $f_2(\mathbf{x}) = c|x_1|^s$, with $c, s \in \mathbb{R}$, vary with any rate $\nu \in [0, 1]$ on $\mathbb{R}^2 \setminus B_1(0)$ and on $\mathbb{R}^2 \setminus [-1, 1] \times \mathbb{R}$ respectively.

Theorem 2. Assume that there is a sequence $(\mathbf{x}_n)_{n\in\mathbb{N}}$ with $|\mathbf{x}_n| \to \infty$ as $n \to \infty$ and constants $k \in \mathbb{N}$, $\varepsilon \in (0,1)$ such that $|\nabla V|, |\nabla B|$ vary with rate 0 on $(\mathbf{x}_n)_{n\in\mathbb{N}}$, as well as

(11)
$$V(\mathbf{x}_n) \longrightarrow -\infty$$
,

(12)
$$\frac{|\nabla B(\mathbf{x}_n)|^2}{|B(\mathbf{x}_n)|^{1-\varepsilon}}, \frac{|\nabla V(\mathbf{x}_n)|^2}{|V(\mathbf{x}_n)|^{1-\varepsilon}} \longrightarrow 0,$$

(13)
$$V(\mathbf{x}_n) + 2k|B(\mathbf{x}_n)| \longrightarrow 0$$

as $n \to \infty$. Then $0 \in \sigma_{ess}(H)$.

Let us now consider the case $V \gg B$ at infinity. The next two theorems state that the accumulation of eigenvalues intensifies, creating more points in the essential spectrum and closing spectral gaps.

Theorem 3. Assume that there is a continuous path $\gamma : \mathbb{R}^+ \to \mathbb{R}^2$, with $|\gamma(t)| \to \infty$ as $t \to \infty$, and constants $\epsilon > 0$, $\nu \in [0,1]$ such that $|\nabla V|, |\nabla B|$ vary with rate ν on $\mathrm{Im}(\gamma)$, as well as

(14)
$$\frac{V(\gamma(t))}{2|B(\gamma(t))|} \longrightarrow -\infty$$

(15)
$$\left(\frac{|\nabla B(\gamma(t))|}{|B(\gamma(t))|} + \frac{|\nabla V(\gamma(t))|}{|V(\gamma(t))|} \right) \left(\frac{|V(\gamma(t))|^3}{B^2(\gamma(t))} \right)^{\frac{1+\epsilon}{2}} \longrightarrow 0,$$

(16)
$$\frac{1}{|\gamma(t)|^{2\nu}} \left(\frac{|V(\gamma(t))|}{B^2(\gamma(t))}\right)^{1+\epsilon} \longrightarrow 0$$

as $t \to \infty$. In addition, suppose that for all $t \in (0, \infty)$ the inequality

(17)
$$B_0 \le |B(\gamma(t))| \le \alpha \exp\left(\kappa \left| \frac{V(\gamma(t))}{B(\gamma(t))} \right| \right)$$

holds with constants $\alpha, \kappa, B_0 > 0$. Then $\sigma_{ess}(H) = \mathbb{R}$.

For our main application, potentials of power-like growth (see discussion after the next theorem), condition (16) imposes unsatisfying restrictions on the growth rate of V/B. At least in the case of a constant magnetic field they can be weakened.

Theorem 4. Let $B = B_0 > 0$ and $V \in C^2(\mathbb{R}^2, \mathbb{R})$. Assume that there is a continuous path $\gamma : \mathbb{R}^+ \to \mathbb{R}^2$, with $|\gamma(t)| \to \infty$ as $t \to \infty$, and constants $\epsilon > 0$, $\nu \in [0,1]$ such that the matrix norm of the Hessian matrix $\|\text{Hess}(V)\|_2 : \mathbb{R}^2 \to \mathbb{R}$ varies with rate ν on $\text{Im}(\gamma)$, as well as

(18)
$$V(\gamma(t)) \longrightarrow -\infty,$$

(19)
$$\|\operatorname{Hess}(V)\|_{2}(\gamma(t))|V(\gamma(t))|^{1+\epsilon} \longrightarrow 0,$$

(20)
$$\frac{1}{|\gamma(t)|^{2\nu}} |V(\gamma(t))|^{1+\epsilon} \longrightarrow 0$$

as $t \to \infty$. In addition, let

(21)
$$\limsup_{t \to \infty} \frac{|\nabla V(\gamma(t))|^2}{|V(\gamma(t))|} < (2B_0)^2.$$

Then $\sigma_{\rm ess}(H) = \mathbb{R}$.

Remark 4. Note that a well-known, basic example for this last theorem is the case of a constant electric field \mathcal{E}_0 in x_1 -direction with the corresponding potential $V(\mathbf{x}) = \mathcal{E}_0 x_1$.

Remark 5. Results similar to that of Theorems 1-4 can be obtained for the magnetic Schrödinger operator with scalar potentials V by using the same techniques as in the proofs of Theorems 1-4.

Finally, we want to discuss some consequences of our results, in particular with respect to spherically symmetric fields B and potentials V, i.e. $B(\mathbf{x}) = b(|\mathbf{x}|)$, $V(\mathbf{x}) = v(|\mathbf{x}|)$ for $\mathbf{x} \in \mathbb{R}$. Using the rotational gauge

$$\mathbf{A}(\mathbf{x}) := \frac{A(r)}{r} \begin{pmatrix} -x_2 \\ x_1 \end{pmatrix}, \quad A(r) = \frac{1}{r} \int_0^r b(s) s \, \mathrm{d}s,$$

with $r = |\mathbf{x}|$, we decompose H in a direct sum of operators on the half-line. More explicitly, there is a unitary map

$$U: L^2(\mathbb{R}^2, \mathbb{C}^2) \to \bigoplus_{j \in \mathbb{Z}} L^2(\mathbb{R}^+, \mathbb{C}^2; dr)$$

such that $UHU^* = \bigoplus_{j \in \mathbb{Z}} h_j$, with

$$h_j := \begin{pmatrix} -\partial_r^2 + \frac{j^2 - 1/4}{r^2} & 0\\ 0 & -\partial_r^2 + \frac{(j+1)^2 - 1/4}{r^2} \end{pmatrix} + A^2(r) - \frac{m_j}{r} A(r) + \sigma_3 A'(r) + v(r)$$

on $L^2(\mathbb{R}^+, \mathbb{C}^2; dr)$, where $m_j = j + \frac{1}{2}$ (see e.g. [13]). It is easy to verify that if

$$\lim_{r \to \infty} \inf b(r) > 0,$$

(23)
$$A'(r)/A^2(r) \longrightarrow 0 \text{ as } r \to \infty,$$

$$\limsup_{r \to \infty} |v(r)|/A^2(r) < 1,$$

then h_j has purely discrete spectrum for every $j \in \mathbb{Z}$. As a consequence, one can use the relations

$$\sigma_{\#}(H) = \overline{\bigcup_{j \in \mathbb{Z}} \sigma_{\#}(h_j)}, \qquad \# \in \{\text{ac, sc, pp}\}$$

to conclude that $\sigma(H) = \sigma_{\rm pp}(H)$, $\sigma_{\rm ac}(H) = \sigma_{\rm sc}(H) = \emptyset$ if (22)-(24) are satisfied. To get more information on $\sigma(H) = \sigma_{\rm pp}(H)$, we employ Theorems 1-4 and obtain:

Corollary 1. Let $b(r) = b_0 r^s$, $v(r) = v_0 r^t$ with $v_0 < 0 < b_0$ and exponents $0 \le s$, $0 \le t \le 2$. Then

- a) $\sigma(H)$ is purely discrete if 0 < t < s or 0 < t = s and $|v_0| < 2b_0$,
- b) $0 \in \sigma_{\text{ess}}(H)$ if 0 < t = s and $|v_0| = 2kB_0$ for some $k \in \mathbb{N}$,
- c) $\sigma(H) = \mathbb{R}$ is dense pure point if 3s < 3t < 2(s+1),
- d) $\sigma(H) = \mathbb{R}$ is dense pure point if s = 0 and 0 < t < 1.

The origins of the strong restrictions on s, t in c), d) can easily be tracked back to conditions (15), (16) of Theorem 3 and (19) of Theorem 4. Unfortunately, even in the case of a constant magnetic field (s = 0) we cannot cover the full range of potentials $(0 < t \le 2)$ for which one might expect $\sigma(H) = \mathbb{R}$.

4. Proof of Theorem 1

Note that the assumptions imply that either $B(\mathbf{x}) \to \infty$ or $B(\mathbf{x}) \to -\infty$. It suffices to consider the case $B(\mathbf{x}) \to \infty$ as $|\mathbf{x}| \to \infty$ since otherwise we only have to interchange the roles of d and d^* in the proof. By modifying B and V on a compact set and comparing the corresponding resolvents, we may assume that B and V satisfy

$$(25) V(\mathbf{x}) \le -1/\delta,$$

(26)
$$|\nabla V(\mathbf{x})| \le |\delta V(\mathbf{x})|,$$

$$(27) |V(\mathbf{x})| \le 2(1 - \eta)B(\mathbf{x}),$$

where $\delta \in (0, \frac{1}{4})$ is fixed, but can be chosen arbitrarily small, and $\eta \in (0, 1)$ is a fixed (δ -independent) constant (c.f. [10] Appendix B).

Using the commutator relation (3), we see that

(28)
$$dd^* \ge 2B \ge (1 - \eta)^{-1} |V| \ge (1 - \eta)^{-1} \delta^{-1}$$

on $C_0^{\infty}(\mathbb{R}^2, \mathbb{C})$ and therefore on $\mathcal{D}(dd^*)$. Since dd^* and d^*d are isospectral away from 0, we obtain a spectral gap $(0,\beta) \subset \varrho(H_{\mathbf{A}})$, with $\beta = (1-\eta)^{-1}\delta^{-1}$. Thus, 0 can be regarded as an isolated point of the spectrum, which is used in the following commutator estimates.

Lemma 1. Let $V \in C^1(\mathbb{R}^2, \mathbb{R})$, $B \in C(\mathbb{R}^2, \mathbb{R})$ and $\mathbf{A} \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ with $B = \text{curl } \mathbf{A}$. Assume further that the conditions (25)-(27) are fulfilled for $\delta \in (0, \frac{1}{4})$ and $\eta \in (0, 1)$. Then:

a) The operators $\left[P_0^{\perp}, V^{-1}\right] V$, $V\left[P_0^{\perp}, V^{-1}\right]$ are well-defined on $C_0^{\infty}(\mathbb{R}^2, \mathbb{C}^2)$ and extend to bounded operators on $L^2(\mathbb{R}^2, \mathbb{C}^2)$ with

$$||V[P_0^{\perp}, V^{-1}]||, ||[P_0^{\perp}, V^{-1}]V|| \le 4\delta^{\frac{3}{2}}.$$

The same holds true if we replace P_0^{\perp} above by P_0 .

b) $P_0\mathcal{D}(V), P_0^{\perp}\mathcal{D}(V) \subset \mathcal{D}(V)$.

Lemma 2. Let $V \in C^1(\mathbb{R}^2, \mathbb{R})$, $B \in C(\mathbb{R}^2, \mathbb{R})$ and $\mathbf{A} \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ with $B = \operatorname{curl} \mathbf{A}$. Assume further that the conditions (25)–(27) are fulfilled for $\delta \in (0, \frac{1}{4})$ and $\eta \in (0, 1)$. Then $\left[\operatorname{sgn}(D_{\mathbf{A}})P_0^{\perp}, V^{-1}\right]$ maps $L^2(\mathbb{R}^2, \mathbb{C}^2)$ into $\mathcal{D}(V)$ and

(29)
$$||V\left[\operatorname{sgn}(D_{\mathbf{A}})P_0^{\perp}, V^{-1}\right]|| \le 4\delta^{\frac{3}{2}}.$$

The proofs of these commutator estimates can be found in [10]. Since $D_{\mathbf{A}}$ is a first-order operator, it is much more convenient to commute V with functions of $D_{\mathbf{A}}$ instead of with functions of $H_{\mathbf{A}}$. For proving Theorem 1, it suffices to find a constant c > 0 such that

(30)
$$||H\varphi|| \ge c||V\varphi||, \quad \varphi \in C_0^{\infty}(\mathbb{R}^2, \mathbb{C}^2)$$

holds (see Lemma 4 in the appendix).

Proof of Theorem 1. Let $\varphi \in C_0^{\infty}(\mathbb{R}^2, \mathbb{C}^2)$. By Lemma 1 we can split $||H\varphi||$ as

$$\begin{aligned} \left\| (H_{\mathbf{A}} + V)\varphi \right\|^2 &= \left\| (H_{\mathbf{A}} + V)(P_0 + P_0^{\perp})\varphi \right\|^2 \\ &= \left\| \left(VP_0 + (H_{\mathbf{A}} + V)P_0^{\perp})\varphi \right\|^2 \\ &= \left\| (H_{\mathbf{A}} + V)P_0^{\perp}\varphi \right\|^2 + 2\operatorname{Re}\langle (H_{\mathbf{A}} + V)P_0^{\perp}\varphi, VP_0\varphi \rangle + \left\| VP_0\varphi \right\|^2 \\ &= \left\| (H_{\mathbf{A}} + V)P_0^{\perp}\varphi \right\|^2 - \delta \left\| VP_0^{\perp}\varphi \right\|^2 \\ &+ 2\operatorname{Re}\langle VP_0\varphi, H_{\mathbf{A}}P_0^{\perp}\varphi \rangle + \left\| V\varphi \right\|^2 - (1 - \delta) \left\| VP_0^{\perp}\varphi \right\|^2. \end{aligned}$$

For the cross-term, condition (26) yields

$$\begin{aligned} |\langle VP_{0}\varphi, H_{\mathbf{A}}P_{0}^{\perp}\varphi\rangle| &= |\langle D_{\mathbf{A}}VP_{0}\varphi, D_{\mathbf{A}}P_{0}^{\perp}\varphi\rangle| \\ &= |\langle [D_{\mathbf{A}}, V] V^{-1}VP_{0}\varphi, D_{\mathbf{A}}P_{0}^{\perp}\varphi\rangle| \\ &\leq \frac{1}{2}\delta^{-\frac{1}{2}}\|(-\mathrm{i}\,\boldsymbol{\sigma}\nabla V)V^{-1}VP_{0}\varphi\|^{2} + \frac{1}{2}\delta^{\frac{1}{2}}\|D_{\mathbf{A}}P_{0}^{\perp}\varphi\|^{2} \\ &\leq \frac{1}{2}\delta^{\frac{3}{2}}\|VP_{0}\varphi\|^{2} + \frac{1}{2}\delta^{\frac{1}{2}}\|H_{\mathbf{A}}P_{0}^{\perp}\varphi\|\|\varphi\| \\ &\leq \frac{1}{4}\delta^{\frac{3}{2}}\|H_{\mathbf{A}}P_{0}^{\perp}\varphi\|^{2} + \frac{1}{4}\delta^{\frac{3}{2}}(\|V\varphi\|^{2} + 2\|VP_{0}\varphi\|^{2}). \end{aligned}$$

Applying Lemma 1 a) results in

$$||VP_0^{\perp}\varphi||, ||VP_0\varphi|| \le (1+4\delta^{\frac{3}{2}})||V\varphi||,$$

and therefore

(32)
$$\|V\varphi\|^{2} - (1-\delta) \|VP_{0}^{\perp}\varphi\|^{2} - \frac{1}{4}\delta^{\frac{3}{2}} \|V\varphi\|^{2} - \frac{1}{2}\delta^{\frac{3}{2}} \|VP_{0}\varphi\|^{2}$$
$$\geq (\delta - 14\delta^{\frac{3}{2}}) \|V\varphi\|^{2}.$$

Because

$$\begin{aligned} \left\| (H_{\mathbf{A}} + V) P_0^{\perp} \varphi \right\|^2 - \delta^{\frac{3}{2}} \| H_{\mathbf{A}} P_0^{\perp} \varphi \|^2 - \delta \| V P_0 \varphi \|^2 \\ & \geq (1 - \varepsilon - \delta^{\frac{3}{2}}) \| H_{\mathbf{A}} P_0^{\perp} \varphi \|^2 + (1 - \varepsilon^{-1} - \delta) \| V P_0^{\perp} \varphi \|^2 \end{aligned}$$

for any $\varepsilon \in (0,1)$, it suffices to show, in view of (31) and (32), that

(33)
$$\langle H_{\mathbf{A}} P_0^{\perp} \varphi, H_{\mathbf{A}} P_0^{\perp} \varphi \rangle + \frac{1 - \varepsilon^{-1} - \delta}{1 - \varepsilon - \delta^{\frac{3}{2}}} \langle V P_0^{\perp} \varphi, V P_0^{\perp} \varphi \rangle \ge 0$$

for $\delta > 0$ small enough and some $\varepsilon \in (0,1)$. We choose $\epsilon = 1 - \delta^{\frac{1}{2}}$, then

$$-\frac{1 - \epsilon^{-1} - \delta}{1 - \epsilon - \delta^{\frac{3}{2}}} = \frac{1}{1 - \delta} \left(\frac{1}{1 - \delta^{\frac{1}{2}}} + \delta^{\frac{1}{2}} \right) =: c_{\delta} > 0.$$

Since $dd^* \geq 2B$ and therefore $\ker(d^*) = \{0\}$, we have

$$(34) P_0^{\perp} = \begin{pmatrix} \pi^{\perp} & 0 \\ 0 & \pi_*^{\perp} \end{pmatrix} = \begin{pmatrix} \pi^{\perp} & 0 \\ 0 & 1 \end{pmatrix}.$$

Setting $\varphi = (\varphi_1, \varphi_2)^T$, one can rewrite (33) as

$$\|H_{\mathbf{A}}P_0^{\perp}\varphi\|^2 - c_{\delta} \|VP_0^{\perp}\varphi\|^2 = \|d^*d\pi^{\perp}\varphi_1\|^2 - c_{\delta} \|V\pi^{\perp}\varphi_1\|^2 + \|dd^*\varphi_2\|^2 - c_{\delta} \|V\varphi_2\|^2.$$

By using the isometries s, s^* given in (5), relation (6) and estimate (28), one obtains

$$\|dd^*\varphi_2\|^2 - c_{\delta} \|V\varphi_2\|^2 = \langle d^*\varphi_2, d^*dd^*\varphi_2 \rangle - c_{\delta} \langle \sqrt{-V}\varphi_2, |V|\sqrt{-V}\varphi_2 \rangle$$

$$= \langle sd^*\varphi_2, dd^*sd^*\varphi_2 \rangle - c_{\delta} \langle \sqrt{-V}\varphi_2, |V|\sqrt{-V}\varphi_2 \rangle$$

$$\geq \langle sd^*\varphi_2, 2Bsd^*\varphi_2 \rangle - c_{\delta} \langle d^*\sqrt{-V}\varphi_2, d^*\sqrt{-V}\varphi_2 \rangle$$

$$\geq \langle sd^*\varphi_2, 2Bsd^*\varphi_2 \rangle - c_{\delta} \langle \sqrt{-V}d^*\varphi_2, \sqrt{-V}d^*\varphi_2 \rangle$$

$$- c_{\delta} \langle [d^*, \sqrt{-V}]\varphi_2, \sqrt{-V}d^*\varphi_2 \rangle$$

$$- c_{\delta} \langle \sqrt{-V}d^*\varphi_2, [d^*, \sqrt{-V}]\varphi_2 \rangle$$

$$- c_{\delta} \langle [d^*, \sqrt{-V}]\varphi_2, [d^*, \sqrt{-V}]\varphi_2 \rangle$$

$$- c_{\delta} \langle [d^*, \sqrt{-V}]\varphi_2, [d^*, \sqrt{-V}]\varphi_2 \rangle$$

$$\geq \|\sqrt{2B}sd^*\varphi_2\|^2 - c_{\delta} \left(\|[d^*, \sqrt{-V}]\varphi_2\| + \|\sqrt{-V}d^*\varphi_2\| \right)^2$$

$$\geq \|\sqrt{2B}sd^*\varphi_2\|^2 - c_{\delta} \left(\delta \|sd^*\varphi_2\| + (1 + 4\delta^{\frac{3}{2}}) \|\sqrt{-V}sd^*\varphi_2\| \right)^2$$

$$\geq [1 - c_{\delta}(1 - \eta)(1 + 15\delta^{\frac{3}{2}})] \|\sqrt{2B}sd^*\varphi_2\|^2,$$

$$(35)$$

where we applied the bound $\left\|\sqrt{-V}\left[s\pi^{\perp},\sqrt{-V}^{-1}\right]\right\| \leq 4\delta^2$. For the latter we write

$$\sqrt{-V} \left[\operatorname{sgn}(D_{\mathbf{A}}) P_0^{\perp}, \sqrt{-V}^{-1} \right] = \begin{pmatrix} 0 & \sqrt{-V} \left[s^*, \sqrt{-V}^{-1} \right] \\ \sqrt{-V} \left[s \pi^{\perp}, \sqrt{-V}^{-1} \right] & 0 \end{pmatrix}$$

and therefore, by Lemma 2 with $\sqrt{-V}$ instead of V, we get

$$\left\|\sqrt{-V}\left[s\pi^{\perp}, \sqrt{-V}^{-1}\right]\right\| \leq \left\|\sqrt{-V}\left[\operatorname{sgn}(D_{\mathbf{A}})P_{0}^{\perp}, \sqrt{-V}^{-1}\right]\right\| \leq 4\delta^{\frac{3}{2}}.$$

Similarly, we obtain a lower bound for $\|d^*d\pi^{\perp}\varphi_1\|^2 - c_{\delta}\|V\pi^{\perp}\varphi_1\|^2$ by using again the upper relation of Equation (6). More precisely,

$$\begin{aligned} \|d^* d\pi^{\perp} \varphi_1\|^2 - c_{\delta} \|V \pi^{\perp} \varphi_1\|^2 &= \|d^* ds^* s \pi^{\perp} \varphi_1\|^2 - c_{\delta} \|V s^* s \pi^{\perp} \varphi_1\|^2 \\ &= \|d d^* s \pi^{\perp} \varphi_1\|^2 - c_{\delta} \|V s^* V^{-1} V s \pi^{\perp} \varphi_1\|^2 \\ &= \|d d^* s \pi^{\perp} \varphi_1\|^2 - c_{\delta} \|\left(s^* + V \left[s^*, V^{-1}\right]\right) V s \pi^{\perp} \varphi_1\|^2, \end{aligned}$$

where $V\left[s^*, V^{-1}\right]$ is one of the components of the operator

$$V\left[\mathrm{sgn}(D_{\mathbf{A}})P_{0}^{\perp},V^{-1}\right] = \left(\begin{array}{cc} 0 & V\left[s^{*},V^{-1}\right] \\ V\left[s\pi^{\perp},V^{-1}\right] & 0 \end{array}\right),$$

so Lemma 2 yields $||V[s^*, V^{-1}]|| \le 4\delta^{\frac{3}{2}}$. Thus,

$$\|d^*d\pi^{\perp}\varphi_1\|^2 - c_{\delta}\|V\pi^{\perp}\varphi_1\|^2 \ge \|dd^*s\pi^{\perp}\varphi_1\|^2 - c_{\delta}(1 + 10\delta^{\frac{3}{2}})\|Vs\pi^{\perp}\varphi_1\|^2.$$

We note that $s\pi^{\perp}\varphi_1 \subset \mathcal{D}(dd^*) \subset \mathcal{D}(V)$, hence we can use (35) (by approximating $s\pi^{\perp}\varphi_1$ through C_0^{∞} -functions in the graph norm of dd^*) to conclude that

$$||d^*d\pi^{\perp}\varphi_1||^2 - c_{\delta}||V\pi^{\perp}\varphi_1||^2$$

$$\geq \left[1 - c_{\delta}(1 - \eta)\left(1 + 10\delta^{\frac{3}{2}}\right)\left(1 + 15\delta^{\frac{3}{2}}\right)\right]||\sqrt{2B}sd^*s\pi^{\perp}\varphi_1||^2.$$

Combining this inequality with (35) leads to

$$\|H_{\mathbf{A}}P_{0}^{\perp}\varphi\|^{2} - c_{\delta} \|VP_{0}^{\perp}\varphi\|^{2}$$

$$\geq \left[1 - c_{\delta}(1 - \eta)(1 + 50\delta^{\frac{3}{2}})\right] \left(\|\sqrt{2B}sd^{*}\varphi_{2}\|^{2} + \|\sqrt{2B}d\pi^{\perp}\varphi_{1}\|^{2}\right),$$

where the r.h.s is non-negativ for δ small enough.

5. Proofs of Theorem 2 and 3

The basic strategy of the proofs is to represent B and V locally through constant values $V_n := V(\mathbf{x}_n)$ and $B_n := B(\mathbf{x}_n)$ along a sequence $(\mathbf{x}_n)_{n \in \mathbb{N}} \subset \mathbb{R}^2$. Since one also needs to compare vector potentials associated to B_n and B, we use the gauges

$$\mathbf{A}_n(\mathbf{x}) := \int_0^1 B_n \wedge (\mathbf{x} - \mathbf{x}_n) s ds = \frac{1}{2} B_n \wedge (\mathbf{x} - \mathbf{x}_n),$$
$$\widetilde{\mathbf{A}}_n(\mathbf{x}) := \int_0^1 B(\mathbf{x}_n + s(\mathbf{x} - \mathbf{x}_n)) \wedge (\mathbf{x} - \mathbf{x}_n) s ds,$$

where $a \wedge \mathbf{v} := a(-v_2, v_1)$ for $a \in \mathbb{R}$ and $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$. The two given vector potentials satisfy curl $\mathbf{A}_n = \text{curl } \widetilde{\mathbf{A}}_n = B$, hence for every $n \in \mathbb{N}$ there exists a function $g_n \in C^2(\mathbb{R}^2, \mathbb{R})$ such that $\nabla g_n = \mathbf{A} - \widetilde{\mathbf{A}}_n$. In addition, for every vector potential \mathbf{A}_n , representing the constant magnetic fields B_n , we obtain operators d_n and d_n^* , $n \in \mathbb{N}$, defined as in (2). For a sequence of natural numbers $(k_n)_{n \in \mathbb{N}}$ we set

$$\psi_n(\mathbf{x}) := \begin{pmatrix} (d_n^*)^{k_n} e^{-B_n |\mathbf{x} - \mathbf{x}_n|^2/4} \\ 0 \end{pmatrix}.$$

Iterating commutator relation (3) for d_n, d_n^* yields

(36)
$$d_n^* d_n \left[(d_n^*)^{k_n} e^{-B_n |\mathbf{x} - \mathbf{x}_n|^2/4} \right] = 2k_n B_n \left[(d_n^*)^{k_n} e^{-B_n |\mathbf{x} - \mathbf{x}_n|^2/4} \right], \quad n \in \mathbb{N},$$

i.e. ψ_n is an eigenfunction of $H_{\mathbf{A}_n}$ with the corresponding eigenvalue $2k_nB_n$. For the localization let $\chi \in C_0^{\infty}(\mathbb{R}^2, [0, 1])$ be such that $\chi(\mathbf{x}) = 1$ for $|\mathbf{x}| \leq 1$ and $\chi(\mathbf{x}) = 0$ for $|\mathbf{x}| \geq 2$. We set

$$\chi_n(\mathbf{x}) := \chi\left(\frac{\mathbf{x} - \mathbf{x}_n}{r_n}\right),$$

where the $r_n > 0$ will be chosen in the proofs. For the Weyl sequence we define the functions φ_n through

(37)
$$\varphi_n(\mathbf{x}) := e^{i g_n(\mathbf{x})} \chi_n(\mathbf{x}) \psi_n(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^2,$$

with $n \in \mathbb{N}$. Bounds on the norm of φ_n can be obtained as in [10]. They are given by:

Lemma 3. For all $n \in \mathbb{N}$ large enough we have

(38)
$$\|\varphi_n\|^2 \le \|\psi_n\|^2 = 2\pi \int_0^\infty (B_n r)^{2k_n} e^{-\frac{B_n}{2}r^2} dr = 2^{k_n+1} \pi B_n^{k_n-1} k_n!,$$

(39)
$$\|\varphi_n\|^2 \ge \|\psi_n\|^2 \left(1 - \frac{1}{k_n!} \int_{\frac{1}{2}B_n r_n^2}^{\infty} s^{k_n} e^{-s} ds\right).$$

Now $H\varphi_n$ can be written as

(40)
$$e^{-ig_{n}}(H_{\mathbf{A}}+V)\varphi_{n} = (H_{\widetilde{\mathbf{A}}_{n}}+V)\chi_{n}\psi_{n}$$

$$= (H_{\mathbf{A}_{n}}+V)\chi_{n}\psi_{n} + 2(\widetilde{\mathbf{A}}_{n}-\mathbf{A}_{n})(-i\nabla-\mathbf{A}_{n})\chi_{n}\psi_{n} + (\widetilde{\mathbf{A}}_{n}-\mathbf{A}_{n})^{2}\chi_{n}\psi_{n} - i\nabla\cdot(\widetilde{\mathbf{A}}_{n}-\mathbf{A}_{n})\chi_{n}\psi_{n} + (B-B_{n})\chi_{n}\psi_{n},$$

with the localization error

(41)
$$(H_{\mathbf{A}_n} + V)\chi_n\psi_n - \chi_n(H_{\mathbf{A}_n} + V)\psi_n$$

$$= -(\Delta \chi_n)\psi_n + 2(-\mathrm{i} \nabla \chi_n)(-\mathrm{i} \nabla - \mathbf{A}_n)\psi_n.$$

To prove Theorem 2 and Theorem 3, we estimate each term of (40) separately. For the proofs we use the notation $K_n := \{x \in \mathbb{R}^2 \mid r_n \leq |\mathbf{x} - \mathbf{x}_n| \leq 2r_n\}$ with $n \in \mathbb{N}$.

Proof of Theorem 2. We set $k_n = k$ and choose the radii to be $r_n^{-4} = B_n^{(2-\epsilon)}$. Then, for any $p \ge 0$,

$$\frac{(B_n)^p}{k_n!} \int_{\frac{1}{2}B_n r_n^2}^{\infty} s^{k_n} e^{-s} ds = \frac{(B_n)^p}{k!} \int_{\frac{1}{2}B_n^{\epsilon/2}}^{\infty} s^k e^{-s} ds \longrightarrow 0 \quad \text{as } n \to \infty.$$

Further, we have $\|\psi_n\|^2 \le 2\|\varphi_n\|^2$ for $n \in \mathbb{N}$ large enough. For treating the terms on the r.h.s. of (40), we estimate

$$\begin{split} & \| (\widetilde{\mathbf{A}}_{n} - \mathbf{A}_{n}) (-i \nabla - \mathbf{A}_{n}) \chi_{n} \psi_{n} \|^{2} \\ & \leq C_{1} r_{n}^{4} |\nabla B(\mathbf{x}_{n})|^{2} \| (-i \nabla - \mathbf{A}_{n}) \chi_{n} \psi_{n} \|^{2} \\ & \leq 2 C_{1} r_{n}^{4} |\nabla B(\mathbf{x}_{n})|^{2} \left[(2k+1) B_{n} \|\psi_{n}\|^{2} + r_{n}^{-2} \|\nabla \chi\|_{\infty}^{2} \int_{K_{n}} |\psi_{n}(\mathbf{x})|^{2} d^{2}x \right] \\ & \leq 16k C_{1} B_{n} \frac{|\nabla B(\mathbf{x}_{n})|^{2}}{B_{n}^{2-\epsilon}} \|\psi_{n}\|^{2} + 4C_{1} \|\nabla \chi\|_{\infty}^{2} \|\psi_{n}\|^{2} B_{n}^{(2-\epsilon)/2} \frac{1}{k!} \int_{\frac{1}{4}B_{n}^{\epsilon/2}}^{\infty} s^{k} e^{-s} ds. \end{split}$$

In addition,

$$\begin{split} \|(\widetilde{\mathbf{A}}_{n} - \mathbf{A}_{n})^{2} \chi_{n} \psi_{n}\|^{2} &\leq C_{2} r_{n}^{4} |\nabla B(\mathbf{x}_{n})|^{4} \|\psi_{n}\|^{2} \leq C_{2} \frac{|\nabla B(\mathbf{x}_{n})|^{4}}{B_{n}^{2(1-\epsilon)}} B_{n}^{-\epsilon} \|\psi_{n}\|^{2}, \\ \|\operatorname{div}(\widetilde{\mathbf{A}}_{n} - \mathbf{A}_{n}) \chi_{n} \psi_{n}\|^{2} &= \|\operatorname{div}\widetilde{\mathbf{A}}_{n} \chi_{n} \psi_{n}\|^{2} \leq C_{3} r_{n}^{2} |\nabla B(\mathbf{x}_{n})|^{2} \|\psi_{n}\|^{2} \\ &\leq C_{3} \frac{|\nabla B(\mathbf{x}_{n})|^{2}}{B_{n}^{1-\epsilon}} B_{n}^{-\epsilon/2} \|\psi_{n}\|^{2}, \\ \|(B - B_{n}) \chi_{n} \psi_{n}\|^{2} \leq C_{4} r_{n}^{2} |\nabla B(\mathbf{x}_{n})|^{2} \|\psi_{n}\|^{2} \leq C_{4} \frac{|\nabla B(\mathbf{x}_{n})|^{2}}{B_{n}^{1-\epsilon}} B_{n}^{-\epsilon/2} \|\psi_{n}\|^{2}. \end{split}$$

For the first term of the r.h.s of (40) we get, due to (41),

$$||(H_{\mathbf{A}_{n}} + V)\chi_{n}\psi_{n}|| \le ||\chi_{n}(H_{\mathbf{A}_{n}} + V)\psi_{n}|| + ||(\Delta\chi_{n})\psi_{n}|| + 2||(-i\nabla\chi_{n})(-i\nabla - \mathbf{A}_{n})\psi_{n}||,$$

with

$$\|(\Delta \chi_n)\psi_n\|^2 \le r_n^{-4} \|\Delta \chi\|_{\infty}^2 \int_{K_n} |\psi_n(\mathbf{x})|^2 d^2 x$$

$$\le 2\|\Delta \chi\|_{\infty}^2 \|\psi_n\|^2 \frac{1}{k!} B_n^{2-\epsilon} \int_{\frac{1}{2}B_n^{\epsilon/2}}^{\infty} s^k e^{-s} ds,$$

and

$$\begin{split} \|(-\mathrm{i}\,\nabla\chi_n)(-\mathrm{i}\,\nabla-\mathbf{A}_n)\psi_n\|^2 &\leq r_n^{-2}\|\nabla\chi\|_{\infty}^2 \int_{K_n} \left|(-\mathrm{i}\,\nabla-\mathbf{A}_n)\psi_n(\mathbf{x})\right|^2 \mathrm{d}^2x \\ &\leq \|\nabla\chi\|_{\infty}^2 B_n^{(2-\epsilon)/2} (2k+1) B_n \int_{K_n} \left|\psi_n(\mathbf{x})\right|^2 \mathrm{d}^2x \\ &\leq \|\nabla\chi\|_{\infty}^2 \|\psi_n\|^2 (2k+1) B_n^{2-\epsilon/2} \int_{\frac{1}{4}B_n^{\epsilon/2}}^{\infty} s^k e^{-s} \mathrm{d}s. \end{split}$$

Because of (36) and since $|\nabla V|$ vary with rate 0, we conclude by the mean value theorem that

$$\begin{split} \|\chi_n(H_{\mathbf{A}_n} + V)\psi_n\|^2 &\leq \|\chi_n(V + 2kB_n)\psi_n\|^2 \\ &\leq C_5 |\nabla V(\mathbf{x}_n)|^2 r_n^2 \|\chi_n\psi_n\|^2 + (2kB_n + V_n)^2 \|\chi_n\psi_n\|^2 \\ &\leq C_5 (4k)^{1-\epsilon} \frac{|\nabla V(\mathbf{x}_n)|^2}{|V_n|^{1-\epsilon}} \|\varphi_n\|^2 + (2kB_n + V_n)^2 \|\varphi_n\|^2 \,. \end{split}$$

Hence, by (40) and conditions (11)-(13), we see that $||(H_{\mathbf{A}} + V)\varphi_n||/||\varphi_n|| \to 0$ as $n \to \infty$. In addition, note that $r_n \to 0$ as $n \to \infty$, so we can assume that the φ_n have mutually disjoint support, i.e. $(\varphi_n)_{n \in \mathbb{N}}$ is a Weyl sequence for 0.

Proof of Theorem 3. We first note that it suffices to proof $0 \in \sigma_{\mathrm{ess}}(H)$ since for $E \in \mathbb{R}$ we consider $V_E := V - E$ instead of V, which also fulfills (14)-(17) along γ . Because $\mathbb{R}^+ \ni t \mapsto V(\gamma(t))/B((\gamma(t)))$ is continuous and (14) holds, we find points $(\mathbf{x}_n)_{n \in \mathbb{N}} \subset \mathrm{Im}(\gamma)$, with $|\mathbf{x}_n| \to \infty$ as $n \to \infty$, such that $2nB(\mathbf{x}_n) = -V(\mathbf{x}_n)$. We choose $k_n = n$ and set

$$(42) r_n := \sqrt{2n^{1+\epsilon}/B_n}.$$

Note that $r_n/|\mathbf{x}_n|^{\nu} \to 0$ as $n \to \infty$ by (16). In particular, we might assume the φ_n 's to have mutually disjoint support. Further, for any $\lambda \geq 0$,

$$\frac{e^{\lambda n}}{n!} \int_{n^{1+\epsilon}}^{\infty} s^n e^{-s} ds \le e^{\lambda n} \exp(n \ln(2n) - n^{1+\epsilon}/2 + n) \longrightarrow 0 \quad \text{as } n \to \infty.$$

Hence, we can choose $N \in \mathbb{N}$ so large that $\|\varphi_n\|^2 \leq \|\psi_n\|^2 \leq 2\|\varphi_n\|^2$ for $n \geq N$. Proceeding as in the proof of Theorem 2, we obtain

$$\begin{split} &\|(\widetilde{\mathbf{A}}_{n} - \mathbf{A}_{n})(-\mathrm{i}\,\nabla - \mathbf{A}_{n})\chi_{n}\psi_{n}\|^{2} \\ &\leq C_{6}r_{n}^{4}|\nabla B(\mathbf{x}_{n})|^{2}(2n+1)B_{n}\bigg[\|\psi_{n}\|^{2} + r_{n}^{-2}\|\nabla\chi\|_{\infty}^{2}\int_{K_{n}}|\psi_{n}(\mathbf{x})|^{2}\mathrm{d}^{2}x\bigg] \\ &\leq 16C_{6}n^{3+2\epsilon}B_{n}\frac{|\nabla B(\mathbf{x}_{n})|^{2}}{B_{n}^{2}}\|\psi_{n}\|^{2} + C_{6}\|\nabla\chi\|_{\infty}^{2}\|\psi_{n}\|^{2}B_{n}\frac{1}{n!}\int_{n^{1+\epsilon}}^{\infty}s^{n}e^{-s}\mathrm{d}s \\ &\leq \widetilde{C}_{6}\left(\frac{|V_{n}|^{3}}{B_{n}^{2}}\right)^{1+\epsilon}\frac{|\nabla B(\mathbf{x}_{n})|^{2}}{B_{n}^{2}}\|\psi_{n}\|^{2} + \alpha C_{6}\|\nabla\chi\|_{\infty}^{2}\|\psi_{n}\|^{2}\frac{e^{2\epsilon n}}{n!}\int_{n^{1+\epsilon}}^{\infty}s^{n}e^{-s}\mathrm{d}s. \end{split}$$

Using (as in the first inequality above) that $|\nabla B|$ vary with rate ν , we conclude

$$\|(\widetilde{\mathbf{A}}_{n} - \mathbf{A}_{n})^{2} \chi_{n} \psi_{n}\|^{2} \leq C_{7} r_{n}^{4} |\nabla B(\mathbf{x}_{n})|^{4} \|\psi_{n}\|^{2} \leq \frac{C_{7}}{4} \frac{|V_{n}|^{6}}{B_{n}^{4}} \frac{|\nabla B(\mathbf{x}_{n})|^{4}}{B_{n}^{4}} \|\psi_{n}\|^{2},$$

$$\|\operatorname{div}(\widetilde{\mathbf{A}}_{n} - \mathbf{A}_{n}) \chi_{n} \psi_{n}\|^{2} \leq C_{8} r_{n}^{2} |\nabla B(\mathbf{x}_{n})|^{2} \|\psi_{n}\|^{2} \leq \frac{C_{8}}{2} \frac{|V_{n}|^{3}}{B_{n}^{2}} \frac{|\nabla B(\mathbf{x}_{n})|^{2}}{B_{n}^{2}} \|\psi_{n}\|^{2},$$

$$\|(B - B_n)\chi_n\psi_n\|^2 \le C_9 r_n^2 |\nabla B(\mathbf{x}_n)|^2 \|\psi_n\|^2 \le \frac{C_9}{2} \frac{|V_n|^3}{B_n^2} \frac{|\nabla B(\mathbf{x}_n)|^2}{B_n^2} \|\psi_n\|^2.$$

We estimate, using equality (41), the first term on the r.h.s. of (40) by

$$||(H_{\mathbf{A}_n} + V)\chi_n \psi_n|| \le ||(V - V_n)\chi_n \psi_n|| + ||(\Delta \chi_n)\psi_n|| + 2||(-i\nabla \chi_n)(-i\nabla - \mathbf{A}_n)\psi_n||,$$

with

$$\|(\Delta \chi_n)\psi_n\|^2 \le \|\Delta \chi\|_{\infty}^2 \|\psi_n\|^2 \frac{B_n^2}{n^{2+2\epsilon}} \frac{1}{n!} \int_{n^{1+\epsilon}}^{\infty} s^n e^{-s} ds$$
$$\le \alpha^2 \|\Delta \chi\|_{\infty}^2 \|\psi_n\|^2 \frac{e^{4\kappa n}}{n^{2+2\epsilon}} \frac{1}{n!} \int_{n^{1+\epsilon}}^{\infty} s^n e^{-s} ds,$$

and

$$\|(-\mathrm{i}\,\nabla\chi_n)(-\mathrm{i}\,\nabla-\mathbf{A}_n)\psi_n\|^2 \le \|\nabla\chi\|_{\infty}^2 \frac{B_n}{n^{1+\epsilon}} \int_{K_n} (2n+1)B_n |\psi_n(\mathbf{x})|^2 \mathrm{d}^2 x$$
$$\le \alpha \|\nabla\chi\|_{\infty}^2 \|\psi_n\|^2 \frac{e^{2\kappa n}}{n^{1+\epsilon}} \frac{1}{n!} \int_{n^{1+\epsilon}}^{\infty} s^n e^{-s} \mathrm{d}s.$$

Since $|\nabla V|$ vary with rate ν , we have

$$\|\chi_n(H_{\mathbf{A}_n} + V)\psi_n\|^2 \le \|\chi_n(V - V_n)\psi_n\|^2$$

$$\leq C_{10}|\nabla V(\mathbf{x}_n)|^2 r_n^2 \|\psi_n\|^2 \leq \widetilde{C}_{10} \left(\frac{|V_n|^3}{B_n^2}\right)^{1+\epsilon} \frac{|\nabla V(\mathbf{x}_n)|^2}{|V_n|^2} \|\psi_n\|^2.$$

We see that $\|(H_{\mathbf{A}_n}+V)\varphi_n\|/\|\varphi_n\|\to 0$ as $n\to\infty$ and therefore, by (40) and the estimates above, that $\|(H_{\mathbf{A}}+V)\varphi_n\|/\|\varphi_n\|\to 0$ as $n\to\infty$.

6. Proof of Theorem 4

Throughout this section we consider the case of a constant magnetic field $B(\mathbf{x}) = B_0$. In addition, we assume that **A** is in the rotational gauge, i.e.

$$\mathbf{A}(\mathbf{x}) = \frac{B_0}{2} \begin{pmatrix} -x_2 \\ x_1 \end{pmatrix}.$$

Note that $H_{\mathbf{A}}$ is invariant under rotations. More precisely, for a special orthogonal matrix $\mathcal{R} \in SO(2,\mathbb{R})$ define the unitary map

$$U_{\mathcal{R}}: L^2(\mathbb{R}^2, \mathbb{C}^2) \to L^2(\mathbb{R}^2, \mathbb{C}^2), \qquad \psi(\cdot) \mapsto \psi(\mathcal{R}^{-1} \cdot),$$

then $U_{\mathcal{R}}^{-1}H_{\mathbf{A}}U_{\mathcal{R}}=H_{\mathbf{A}}$ and therefore

$$U_{\mathcal{R}}^{-1}(H_{\mathbf{A}} + V)U_{\mathcal{R}} = H_{\mathbf{A}} + V_{\mathcal{R}} \text{ with } V_{\mathcal{R}}(\cdot) = V(\mathcal{R} \cdot).$$

To construct a Weyl sequence, consider a second gauge $\tilde{\mathbf{A}}(\mathbf{x}) = B_0 x_1 \hat{e}_2$, called the Landau gauge. Then our Hamiltonian reads

(43)
$$H_{\tilde{\mathbf{A}}} + V = -\partial_1^2 + (-\mathrm{i}\,\partial_2 - B_0 x_1)^2 - \sigma_3 B_0 + V$$
$$= \tilde{d}^* \tilde{d} + B_0 - \sigma_3 B_0 + V,$$

with

$$\tilde{d} = -i \partial_1 + i (-i \partial_2 - B_0 x_1), \quad \tilde{d}^* = -i \partial_1 - i (-i \partial_2 - B_0 x_1).$$

For electric fields of the form $V(\mathbf{x}) = V_0 + \mathcal{E}_0(x_1 - \zeta)$, with constants $V_0, \mathcal{E}_0, \zeta \in \mathbb{R}$, we can write

$$H_{\widetilde{\mathbf{A}}} + V = -\partial_1^2 + (-\mathrm{i}\,\partial_2 - B_0 x_1)^2 - \sigma_3 B_0 + V_0 + \mathcal{E}_0(x_1 - \zeta)$$

$$= -\partial_1^2 + B_0^2 \left(x_1 - \frac{1}{B_0} \left(-\mathrm{i}\,\partial_2 - \frac{\mathcal{E}_0}{2B_0}\right)\right)^2 + \frac{\mathcal{E}_0}{B_0} \left(-\mathrm{i}\,\partial_2 - \frac{\mathcal{E}_0}{2B_0}\right) - \mathcal{E}_0 \zeta - \sigma_3 B_0 + V_0 + \left(\frac{\mathcal{E}_0}{2B_0}\right)^2.$$

Performing a Fourier transform in x_2 , we obtain the direct integral representation

$$H_{\widetilde{\mathbf{A}}} + V \cong \int_{\mathbb{R}}^{\oplus} h(\xi) d\xi$$

on $L^2(\mathbb{R}_{\xi}, L^2(\mathbb{R}, \mathbb{C}^2))$, with

$$h(\xi) = -\partial_1^2 + B_0^2 \left(x_1 - \frac{1}{B_0} \left(\xi - \frac{\mathcal{E}_0}{2B_0} \right) \right)^2 + \frac{\mathcal{E}_0}{B_0} \left(\xi - \frac{\mathcal{E}_0}{2B_0} \right) - \mathcal{E}_0 \zeta - \sigma_3 B_0 + V_0 + \left(\frac{\mathcal{E}_0}{2B_0} \right)^2$$

= $-\partial_1^2 + B_0^2 \left(x_1 - \hat{\zeta} \right)^2 + \mathcal{E}_0 \hat{\zeta} - \mathcal{E}_0 \zeta - \sigma_3 B_0 + V_0 + \left(\frac{\mathcal{E}_0}{2B_0} \right)^2.$

Here we set $\hat{\zeta} = \frac{1}{B_0} \left(\xi - \frac{\mathcal{E}_0}{2B_0} \right)$. Note that $h(\xi)$ is the Hamiltonian of a shifted harmonic oscillator. Thus, we define for $n \in \mathbb{N}_0$

$$\phi_n(x) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} \vartheta_n(x) e^{-x^2/2}, \quad x \in \mathbb{R},$$

where ϑ_n denotes the n-th Hermite polynomial. The normalized functions

$$\widehat{\psi}_{\mathcal{E}_0,n,\xi}(x_1) := \sqrt[4]{B_0} \begin{pmatrix} \phi_n \left(\sqrt{B_0} \left(x_1 - \frac{1}{B_0} \left(\xi - \frac{\mathcal{E}_0}{2B_0} \right) \right) \right) \\ 0 \end{pmatrix}$$

fulfill the equation

$$h(\xi)\widehat{\psi}_{\mathcal{E}_0,n,\xi} = \left(2nB_0 + \mathcal{E}_0\left(\frac{1}{B_0}\left(\xi - \frac{\mathcal{E}_0}{2B_0}\right) - \zeta\right) + V_0 + \left(\frac{\mathcal{E}_0}{2B_0}\right)^2\right)\widehat{\psi}_{\mathcal{E}_0,n,\xi}.$$

Hence.

(45)
$$\psi_{\mathcal{E}_0,n,\xi}(x_1,x_2) := e^{i\,\xi x_2} \widehat{\psi}_{\mathcal{E}_0,n,\xi}(x_1)$$

satisfies

$$(46) \quad \left[H_{\widetilde{\mathbf{A}}} + V\right] \psi_{\mathcal{E}_0, n, \xi} = \left(2nB_0 + \mathcal{E}_0\left(\frac{1}{B_0}\left(\xi - \frac{\mathcal{E}_0}{2B_0}\right) - \zeta\right) + V_0 + \left(\frac{\mathcal{E}_0}{2B_0}\right)^2\right) \psi_{\mathcal{E}_0, n, \xi}$$

for $\xi \in \mathbb{R}$ and $n \in \mathbb{N}_0$, seen as a differential equation. In addition, we have

(47)
$$\tilde{d}\psi_{\mathcal{E}_0,n,\xi} = -i\sqrt{2nB_0}\psi_{\mathcal{E}_0,n-1,\xi} + i\frac{\mathcal{E}_0}{2B_0}\psi_{\mathcal{E}_0,n,\xi},$$

(48)
$$\tilde{d}^* \psi_{\mathcal{E}_0, n, \xi} = i \sqrt{2(n+1)B_0} \psi_{\mathcal{E}_0, n+1, \xi} - i \frac{\mathcal{E}_0}{2B_0} \psi_{\mathcal{E}_0, n, \xi}.$$

Proof of Theorem 4. As argumented in the proof of Theorem 3, it suffices to find a Weyl sequence for E=0. Because of (18) and (21), there exists a sequence $\{\mathbf{y}_n\}_{n\in\mathbb{N}}\subset\mathrm{Im}(\gamma)$ such that

(49)
$$V(\mathbf{y}_n) = -2nB_0 - \left(\frac{|\nabla V(\mathbf{y}_n)|}{2B_0}\right)^2.$$

Further, there are rotations $\mathcal{R}_n \in SO(2,\mathbb{R})$ such that $\nabla V_{\mathcal{R}_n}(\mathbf{x}_n) = |\nabla V_{\mathcal{R}_n}(\mathbf{x}_n)|\hat{e}_1$, with $\mathbf{x}_n = \mathcal{R}_n^{-1}\mathbf{y}_n = (x_{n,1}, x_{n,2})^T$ for $n \in \mathbb{N}$. We set

$$(50) V_n := V(\mathbf{y}_n) = V_{\mathcal{R}_n}(\mathbf{x}_n),$$

(51)
$$\mathcal{E}_n := |\nabla V(\mathbf{y}_n)| = |\nabla V_{\mathcal{R}_n}(\mathbf{x}_n)|,$$

$$\xi_n := B_0 x_{n,1} + \frac{\mathcal{E}_n}{2B_0}.$$

For the Weyl functions let $\chi \in C_0^{\infty}(\mathbb{R}, [0, 1])$ with $\chi(x) = 1$ for $|x| \leq 1$ and $\chi(x) = 0$ for $|x| \geq 2$. Define

$$\chi_{n,j}(x) := \chi\left(\frac{x - x_{n,j}}{r_n}\right),$$

for j = 1, 2, and

$$\varphi_n(\mathbf{x}) := \chi_{n,1}(x_1)\chi_{n,2}(x_2)\psi_{\mathcal{E}_n,n,\xi_n}(x_1,x_2) = \chi\left(\frac{x_2 - x_{n,2}}{r_n}\right)e^{-i\,\xi_n x_2}\chi\left(\frac{x_1 - x_{n,1}}{r_n}\right)\left(\sqrt[4]{B_0}\phi_n\left(\sqrt{B_0}(x_1 - x_{n,1})\right)\right),$$

where the localization radii r_n are chosen to be $r_n := \sqrt{n^{1+\epsilon}/B_0}$. Note that

$$(53) r_n \le 2r_n \int_{-\sqrt{n^{1+\epsilon}}}^{\sqrt{n^{1+\epsilon}}} |\phi_n(x)|^2 dx \le \|\varphi_n\|^2 \le 4r_n \int_{-2\sqrt{n^{1+\epsilon}}}^{2\sqrt{n^{1+\epsilon}}} |\phi_n(x)|^2 dx \le 4r_n$$

for $n \in \mathbb{N}$ large enough (see Lemma 5 in the appendix). By denoting $g(\mathbf{x}) = \frac{B_0}{2}x_1x_2$ for $\mathbf{x} \in \mathbb{R}^2$, we get, due to (43), (46), (49) and (52), that

(54)
$$HU_{\mathcal{R}_{n}}e^{-ig}\varphi_{n} = U_{\mathcal{R}_{n}}e^{-ig}\left[H_{\widetilde{\mathbf{A}}} + V_{\mathcal{R}_{n}}\right]\varphi_{n}$$

$$= U_{\mathcal{R}_{n}}e^{-ig}\left(\tilde{d}^{*}\tilde{d}\varphi_{n} - \chi_{n,1}\chi_{n,2}\tilde{d}^{*}\tilde{d}\psi_{\mathcal{E}_{n},n,\xi_{n}}\right) + U_{\mathcal{R}_{n}}e^{-ig}\left[V_{\mathcal{R}_{n}} - V_{n} - \mathcal{E}_{n}(x_{1} - x_{1,n})\right]\varphi_{n}.$$

The localization error results in

$$\begin{split} \tilde{d}^* \tilde{d} \varphi_n - \chi_{n,1} \chi_{n,2} \tilde{d}^* \tilde{d} \psi_{\mathcal{E}_n,n,\xi_n} \; &= \; \big[-\mathrm{i} \, \chi_{n,2} \partial_1 \chi_{n,1} + \chi_{n,1} \partial_2 \chi_{n,2} \big] \tilde{d}^* \psi_{\mathcal{E}_n,n,\xi_n} \; + \\ & \quad \big[-\mathrm{i} \, \chi_{n,2} \partial_1 \chi_{n,1} - \chi_{n,1} \partial_2 \chi_{n,2} \big] \tilde{d} \psi_{\mathcal{E}_n,n,\xi_n} \; + \\ & \quad \big[- \chi_{n,2} \partial_1^2 \chi_{n,1} - \chi_{n,1} \partial_2^2 \chi_{n,2} \big] \psi_{\mathcal{E}_n,n,\xi_n}, \end{split}$$

with, using (47) and (48),

$$\begin{split} \left\| \left[- \mathrm{i} \, \chi_{n,2} \partial_1 \chi_{n,1} + \chi_{n,1} \partial_2 \chi_{n,2} \right] \tilde{d}^* \psi_{\mathcal{E}_n,n,\xi_n} \right\| \\ & \leq \sqrt{2(n+1)B_0} \left\| \left[- \mathrm{i} \, \chi_{n,2} \partial_1 \chi_{n,1} + \chi_{n,1} \partial_2 \chi_{n,2} \right] \psi_{\mathcal{E}_n,n+1,\xi_n} \right\| \\ & + \frac{\mathcal{E}_n}{2B_0} \left\| \left[- \mathrm{i} \, \chi_{n,2} \partial_1 \chi_{n,1} + \chi_{n,1} \partial_2 \chi_{n,2} \right] \psi_{\mathcal{E}_n,n,\xi_n} \right\| \\ & \leq 2\sqrt{2(n+1)B_0} r_n^{-1} \|\chi'\|_{\infty} \sqrt{2r_n} \|\phi_{n+1}\| + 2\frac{\mathcal{E}_n}{2B_0} r_n^{-1} \|\chi'\|_{\infty} \sqrt{2r_n} \|\phi_n\| \\ & \leq 2\sqrt{2} \|\chi'\|_{\infty} \left(B_0 \sqrt{\frac{2n+2}{n^{1+\epsilon}}} + \frac{\mathcal{E}_n}{2} \sqrt{\frac{B_0}{n^{1+\epsilon}}} \right) \sqrt{r_n} \,, \end{split}$$

$$\begin{split} \left\| \left[-\mathrm{i}\,\chi_{n,2}\partial_1\chi_{n,1} - \chi_{n,1}\partial_2\chi_{n,2} \right] \tilde{d}\psi_{\mathcal{E}_n,n,\xi_n} \right\| \\ & \leq \sqrt{2nB_0} \left\| \left[-\mathrm{i}\,\chi_{n,2}\partial_1\chi_{n,1} - \chi_{n,1}\partial_2\chi_{n,2} \right] \psi_{\mathcal{E}_n,n-1,\xi_n} \right\| \\ & + \frac{\mathcal{E}_n}{2B_0} \left\| \left[-\mathrm{i}\,\chi_{n,2}\partial_1\chi_{n,1} - \chi_{n,1}\partial_2\chi_{n,2} \right] \psi_{\mathcal{E}_n,n,\xi_n} \right\| \\ & \leq 2\sqrt{2} \|\chi'\|_{\infty} \left(B_0 \sqrt{\frac{2n}{n^{1+\epsilon}}} + \frac{\mathcal{E}_n}{2} \sqrt{\frac{B_0}{n^{1+\epsilon}}} \right) \sqrt{r_n} \end{split}$$

and

$$\| \left[-\chi_{n,2} \partial_1^2 \chi_{n,1} - \chi_{n,1} \partial_2^2 \chi_{n,2} \right] \psi_{\mathcal{E}_n,n,\xi_n} \| \le 2\sqrt{2} \| \chi'' \|_{\infty} r_n^{-2} \sqrt{r_n} \,.$$

Thus, in view of condition (21) and estimate (53), we get

(55)
$$\|\tilde{d}^*\tilde{d}\varphi_n - \chi_{n,1}\chi_{n,2}\tilde{d}^*\tilde{d}\psi_{\mathcal{E}_n,n,\xi_n}\|/\|\varphi_n\| \longrightarrow 0 \text{ as } n \to \infty.$$

For estimating the remaining term on the r.h.s of (54), we expand $V_{\mathcal{R}_n}$ up to second order and obtain, by (50) and (51), that

$$|[V_{\mathcal{R}_n}(\mathbf{x}) - V_n - \mathcal{E}_n(x_1 - x_{1,n})]\varphi_n(\mathbf{x})| \le ||\operatorname{Hess}(V_{\mathcal{R}_n})||_2(\eta_{x,x_n})|\mathbf{x} - \mathbf{x}_n|^2|\varphi_n(\mathbf{x})|,$$

with $\eta_{x,x_n} \in [\mathbf{x}, \mathbf{x}_n]$. Because \mathcal{R}_n are rotations, we have that $\|\operatorname{Hess}(V_{\mathcal{R}_n})\|_2(\cdot) = \|\operatorname{Hess}(V)\|_2(\mathcal{R}_n \cdot)$ for $n \in \mathbb{N}$. Since $\|\operatorname{Hess}(V)\|_2$ varies with rate ν along $\operatorname{Im}(\gamma)$ and since, by (20) and (49), $r_n/|\mathbf{x}_n|^{\nu} \to 0$ as $n \to \infty$, we find a constant $C_{11} > 0$ such that for $n \in \mathbb{N}$ large enough

$$\|\operatorname{Hess}(V_{\mathcal{R}_n})\|_2(\eta) \le C_{11} \|\operatorname{Hess}(V_{\mathcal{R}_n})\|_2(\mathbf{x}_n), \quad \eta \in B_{2r_n}(\mathbf{x}_n)$$

holds. As a consequence,

$$\begin{aligned} \left\| U_{\mathcal{R}_n} e^{-\mathrm{i}\,g} \left[V_{\mathcal{R}_n} - V_n - \mathcal{E}_n(x_1 - x_{1,n}) \right] \varphi_n \right\| &\leq 4C_{11} r_n^2 \| \mathrm{Hess}(V) \|_2(\mathcal{R}_n \mathbf{x}_n) \| \varphi_n \| \\ &\leq 4C_{11} \| \mathrm{Hess}(V) \|_2(\mathbf{y}_n) \left(\frac{|V_n|}{B_0} \right)^{1+\epsilon} \| \varphi_n \| \end{aligned}$$

for $n \in \mathbb{N}$ large enough. In view of (19), we conclude that $(U_{\mathcal{R}_n}e^{-ig}\varphi_n)_{n\in\mathbb{N}}$ is a Weyl sequence for 0.

Acknowledgments. This work has been supported by SFB-TR12 "Symmetries and Universality in Mesoscopic Systems" of the DFG. The author also wants to thank Edgardo Stockmeyer for useful discussions and remarks as well as the *Faculdad de Física de la Pontificia Universidad Católica de Chile* for the great hospitality.

APPENDIX A. ESSENTIAL SELF-ADJOINTNESS OF THE PAULI OPERATOR

In this section we recapitulate an argument, originally given in [8], for proving the essential self-adjointness of the Pauli operator. As we will see, this argumentation works also for the relaxed regularity conditions on B and V of Proposition 1.

For the proof we first note that for $\varphi \in C_0^{\infty}(\mathbb{R}^2, \mathbb{C})$ we can write

$$[(-i\nabla - \mathbf{A})^2 + B]\varphi = \sum_{k,l=1}^{2} (-i\partial_k - A_k)\overline{C_{k,l}}(-i\partial_l - A_l)\varphi,$$
$$[(-i\nabla - \mathbf{A})^2 - B]\varphi = \sum_{k,l=1}^{2} (-i\partial_k - A_k)C_{k,l}(-i\partial_l - A_l)\varphi,$$

where $C_{k,l}$ denote the coefficients of the symmetric non-negative definite matrix

$$C = \mathbb{1} - \sigma_2 = \begin{pmatrix} 1 & \mathbf{i} \\ -\mathbf{i} & 1 \end{pmatrix} = C^*.$$

Furthermore, along the proof we use the notation $B_R := \{ \mathbf{x} \in \mathbb{R}^2 | |\mathbf{x}| \leq R \}$ and $S_R := \{ \mathbf{x} \in \mathbb{R}^2 | |\mathbf{x}| = R \}.$

Proof of Proposition 1. Since H is a diagonal matrix operator, it suffices to show that both operators on the diagonal,

$$Q_{\pm} := [(-\mathrm{i}\,\nabla - \mathbf{A})^2 \pm B + V],$$

are essentially self-adjoint on $C_0^{\infty}(\mathbb{R}^2, \mathbb{C})$. Because Q_{\pm} are symmetric on $C_0^{\infty}(\mathbb{R}^2, \mathbb{C})$, we have to show $Q_+^*\varphi = \pm i \varphi$ implies $\varphi \equiv 0$ for $\varphi \in \mathcal{D}(Q_+^*)$, respectively $Q_-^*\varphi = \pm i \varphi$ implies $\varphi \equiv 0$ for $\varphi \in \mathcal{D}(Q_-^*)$. We only treat the case $Q_-^*\varphi = i \varphi$ since the others are completely analogous. Let $\varphi \in \mathcal{D}(Q_-^*)$ be such that $Q_-^*\varphi = i \varphi$, then

(56)
$$[(-i\nabla - \mathbf{A})^2 - B + V]\varphi = i\varphi$$

holds in distributional sense. Due to elliptic regularity theory (see e.g. [6], [7]), we obtain that $\varphi \in C^2(\mathbb{R}^2, \mathbb{C})$ and that (56) holds strongly. Applying integration by parts results in

$$\int_{B_R} \left[\sum_{k,l=1}^{2} (-\mathrm{i}\,\partial_k - A_k) C_{k,l} (-\mathrm{i}\,\partial_l - A_l) \varphi \right] \bar{\varphi} \,\mathrm{d}^2 x$$

$$= \int_{B_R} \left[\sum_{k,l=1}^{2} (-\mathrm{i}\,\partial_k - A_k) \varphi C_{k,l} \overline{(-\mathrm{i}\,\partial_l - A_l) \varphi} \right] \mathrm{d}^2 x$$

$$+ \mathrm{i} \int_{S_R} \left[\sum_{k,l=1}^{2} \nu_k C_{k,l} (-\mathrm{i}\,\partial_l - A_l) \varphi \right] \bar{\varphi} \,\mathrm{d}S,$$

with R > 0, where $\nu_k(\mathbf{x}) = x_k/|\mathbf{x}|$ for k = 1, 2. By taking the imaginary part of (57), we conclude with (56) that

$$\int_{B_R} |\varphi|^2 d^2 x = \int_{S_R} \left[\sum_{k,l=1}^2 \nu_k C_{k,l} (-i \partial_l - A_l) \varphi \right] \bar{\varphi} dS$$

for any R > 0. The Cauchy-Schwarz inequality yields

$$\int_{B_R} |\varphi|^2 d^2x \le \left(\int_{S_R} \sum_{k,l=1}^2 (-i \partial_k - A_k) \varphi C_{k,l} \overline{(-i \partial_l - A_l) \varphi} dS \right)^{1/2} \left(\int_{S_R} |\varphi|^2 dS \right)^{1/2}.$$

Hence, it suffices to show that

(58)
$$\int_{\mathbb{R}^2} \sum_{k,l=1}^2 \frac{(-\mathrm{i}\,\partial_k - A_k)\varphi C_{k,l}\overline{(-\mathrm{i}\,\partial_l - A_l)\varphi}}{|\mathbf{x}|^2 + 1} \,\mathrm{d}^2 x < \infty$$

since this implies that $(1, \infty) \ni r \mapsto r^{-1} \int_{B_r} |\varphi|^2 d\mathbf{x}$ is an L^1 -function, i.e. $\varphi \equiv 0$. For (58) we consider the function

$$f(R) := \int_{B_R} \sum_{k,l=1}^{2} \frac{(-\mathrm{i}\,\partial_k - A_k)\varphi C_{k,l} \overline{(-\mathrm{i}\,\partial_l - A_l)\varphi}}{|\mathbf{x}|^2 + 1} \,\mathrm{d}^2 x\,,$$

with R > 0. Using Equation (56) and integration by parts, we obtain, with $\zeta(\mathbf{x}) = (|\mathbf{x}|^2 + 1)^{-1}$ and $M \ge c + |d|$, that

$$\begin{split} f(R) - M \|\varphi\|^2 &\leq f(R) + \int_{B_R} \zeta V |\varphi|^2 \,\mathrm{d}^2 x \\ &= \int_{B_R} \zeta (Q_-^* \varphi) \bar{\varphi} \,\mathrm{d}^2 x - \mathrm{i} \int_{B_R} \left[\sum_{k,l=1}^2 (\partial_l \zeta) C_{k,l} (-\mathrm{i} \, \partial_k - A_k) \varphi \right] \bar{\varphi} \,\mathrm{d}^2 x \\ &+ \mathrm{i} \int_{S_R} \zeta \left[\sum_{k,l=1}^2 \nu_l C_{k,l} (-\mathrm{i} \, \partial_k - A_k) \varphi \right] \bar{\varphi} \,\mathrm{d} S. \end{split}$$

By the estimates

$$\left| \int_{B_R} \left[\sum_{k,l=1}^2 (\partial_l \zeta) C_{k,l} (-\mathrm{i} \, \partial_k - A_k) \varphi \right] \bar{\varphi} \, \mathrm{d}^2 x \right|$$

$$\leq \int_{B_R} 2\zeta^{1/2} \left| \sum_{k,l=1}^2 \nu_l C_{k,l} (-\mathrm{i} \, \partial_k - A_k) \varphi \right| |\varphi| \, \mathrm{d}^2 x$$

$$\leq 2 \int_{B_R} \left[\sum_{k,l=1}^2 \zeta (-\mathrm{i} \, \partial_k - A_k) \varphi C_{k,l} \overline{(-\mathrm{i} \, \partial_l - A_l) \varphi} \right]^{1/2} |\varphi| \, \mathrm{d}^2 x$$

$$\leq 2 [f(R)]^{1/2} \|\varphi\| \leq \frac{1}{2} f(R) + 2 \|\varphi\|^2$$

and

$$\left| \int_{S_R} \zeta \left[\sum_{k,l=1}^2 \nu_k C_{k,l} (-i \,\partial_l - A_l) \varphi \right] \bar{\varphi} \, \mathrm{d}S \right|$$

$$\leq \int_{S_R} \zeta \left[\sum_{k,l=1}^2 (-i \,\partial_k - A_k) \varphi C_{k,l} \overline{(-i \,\partial_l - A_l) \varphi} \right]^{1/2} |\varphi| \, \mathrm{d}S$$

$$\leq \left(\int_{S_R} \sum_{k,l=1}^2 \zeta (-i \,\partial_k - A_k) \varphi C_{k,l} \overline{(-i \,\partial_l - A_l) \varphi} \, \mathrm{d}S \right)^{1/2} \left(\int_{S_R} |\varphi|^2 \, \mathrm{d}S \right)^{1/2}$$

$$= \left(f'(R) \int_{S_R} |\varphi|^2 \, \mathrm{d}S \right)^{1/2},$$

we conclude that

$$f(R) \le 2(3+c)\|\varphi\|^2 + 2\left(f'(R)\int_{S_R} |\varphi|^2 dS\right)^{1/2}.$$

If f(R)=0 for all R>0, then clearly (58) holds. If $f(R_0)>0$ for some $R_0>0$, then f(R)>0 for all $R>R_0$ and $f'(R)/f^2(R)\in L^1((R_0,\infty))$, implying that

$$\left(\frac{f'(R)}{f^2(R)}\int_{S_R} |\varphi|^2 dS\right)^{1/2} \in L^1((R_0, \infty)).$$

Hence, there exists a sequence $(R_n)_{n\in\mathbb{N}}\subset(0,\infty)$ such that $R_n\to\infty$ as $n\to\infty$ and

$$\left(f'(R_n)\int_{S_{R_n}} |\varphi|^2 dS\right)^{1/2} \le \frac{1}{4} f(R_n).$$

Therefore, we have $f(R_n) \leq 4(3+c)\|\varphi\|^2$ for all $n \in \mathbb{N}$, which implies (58) since f(R) is a monotonically increasing function.

APPENDIX B. REMARKS ON LOCALLY COMPACT OPERATORS

Lemma 4. Let A be a locally compact, self-adjoint operator on $L^2(\mathbb{R}^n, \mathbb{C}^m)$ with $n, m \geq 1$. Assume there is a function $W \in L^{\infty}_{loc}(\mathbb{R}^n, [0, \infty))$, with $W(\mathbf{x}) \to \infty$ as $|\mathbf{x}| \to \infty$, such that

$$||A\varphi|| \ge ||W\varphi|| \quad for \ \varphi \in \mathcal{D}(A).$$

Then $\sigma_{\text{ess}}(A) = \emptyset$, i.e. A has only discrete spectrum.

Proof. Assume $\lambda \in \sigma_{\text{ess}}(A) \subset \mathbb{R}$. Then there is a normalized sequence $(\varphi_n)_{n \in \mathbb{N}} \subset \mathcal{D}(A)$ such that $\varphi_n \to 0$ and $\|(A - \lambda)\varphi_n\| \to 0$ as $n \to \infty$. Let R > 0 be a fixed constant. We have

$$\|\chi_R \varphi_n\| = \|\chi_R (A - i - \lambda)^{-1} (A - i - \lambda) \varphi_n\|$$

$$\leq \|\chi_R (A - i - \lambda)^{-1}\| \|(A - \lambda) \varphi_n\| + \|\chi_R (A - i - \lambda)^{-1} \varphi_n\|,$$

using the notation $\chi_R := \chi_{B_R(0)}$. Since $\chi_R(A - \mathbf{i} - \lambda)^{-1}$ is compact, this inequality implies that $\|\chi_R \varphi_n\| \to 0$ as $n \to \infty$. Let R > 0 be so large that $W(\mathbf{x}) \ge 5|\lambda| + 1$ if $|\mathbf{x}| \ge R$. Choosing $N \in \mathbb{N}$ large enough, we can estimate, for $n \ge N$,

$$||(A - \lambda)\varphi_n|| \ge ||W\varphi_n|| - |\lambda|$$

$$\ge ||W(\mathbb{1} - \chi_R)\varphi_n|| - ||W\chi_R||_{\infty} ||\chi_R\varphi_n|| - |\lambda|$$

$$\ge (5|\lambda| + 1)||(\mathbb{1} - \chi_R)\varphi_n|| - ||W\chi_R||_{\infty} ||\chi_R\varphi_n|| - |\lambda|$$

$$\ge (|\lambda| + 1/2) - ||W\chi_R||_{\infty} ||\chi_R\varphi_n||.$$

Hence, $\|(A-\lambda)\varphi_n\| \to 0$ as $n \to \infty$, which is a contradiction.

APPENDIX C. INTEGRAL ESTIMATES

Lemma 5. Let ϑ_n be the n-th Hermite polynomial. Then, for any $\epsilon > 0$, it holds that

$$\frac{1}{2^n n!} \int_{\sqrt{n^{1+\epsilon}}}^{\infty} |\vartheta_n(x)|^2 e^{-x^2} dx \longrightarrow 0 \quad as \ n \to \infty,$$

$$\frac{1}{2^n n!} \int_{-\sqrt{n^{1+\epsilon}}}^{-\sqrt{n^{1+\epsilon}}} |\vartheta_n(x)|^2 e^{-x^2} dx \longrightarrow 0 \quad as \ n \to \infty.$$

 $2^n n! \int_{-\infty}^{\infty} e^{-n(x)} e^{-n(x)} dx$ where $e^{-n(x)}$ is the first case since the second claim can be deduced from the

$$\vartheta_n(x) = (-1)^n \sum_{k_1 + 2k_2 - n} \frac{n!}{k_1! k_2!} (-1)^{k_1 + k_2} (2x)^{k_1}$$

for the n-th Hermite polynomial (see e.g. [1]), we obtain for $|x| \geq 1$ the estimate

$$|\vartheta_n(x)| \le \sum_{k_1+2k_2=n} \frac{n!}{k_1!k_2!} (2|x|)^{k_1} \le \frac{n+1}{2} 2^n n! |x|^n.$$

Thus, for $n \in \mathbb{N}$ large enough we have

$$\frac{1}{2^{n} n!} \int_{\sqrt{n^{1+\epsilon}}}^{\infty} |\vartheta_{n}(x)|^{2} e^{-x^{2}} dx \leq \frac{(n+1)^{2}}{4} 2^{n} n! \int_{\sqrt{n^{1+\epsilon}}}^{\infty} x^{2n} e^{-x^{2}} dx
\leq \frac{(n+1)^{2}}{4} 2^{n} n^{1+\epsilon} n! \exp(-n^{1+\epsilon}/2) \int_{\sqrt{n^{1+\epsilon}}}^{\infty} e^{-x^{2}/2} dx,$$

and the r.h.s. tends to 0 as $n \to \infty$ by Stirling's Formula.

first one by a symmetry argument. Due to the identity

References

- M. Abramowitz and I. A. Stegun. Handbook of mathematical functions with formulas, graphs, and mathematical tables. National Bureau of Standards Applied Mathematics Series, vol. 55. Washington, D.C., 1964.
- [2] J. Avron, I. Herbst and B. Simon. Schrödinger operators with magnetic fields. I. General interactions. *Duke Math. J.*, 45(4):847–883, 1978.
- [3] P. R. Chernoff. Schrödinger and Dirac operators with singular potentials and hyperbolic equations. *Pacific J. Math.*, 72(2):361–382, 1977.
- [4] H. L. Cycon, R. G. Froese, W. Kirsch, and B. Simon. Schrödinger operators with application to quantum mechanics and global geometry. Texts and Monographs in Physics. Springer-Verlag, Berlin, study edition, 1987.
- [5] L. Erdős. Recent developments in quantum mechanics with magnetic fields. In Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon's 60th birthday, volume 76 of Proc. Sympos. Pure Math., pages 401–428. Amer. Math. Soc., Providence, RI, 2007.
- [6] D. Gilbarg and N. S. Trudinger. Elliptic partial differential equations of second order, volume 224 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2nd edition, 1983.
- [7] G. Hellwig. Partial differential equations: an introduction. B. G. Teubner, Stuttgart, 2nd edition, 1977. Translated from the German by Eberhard Gerlach, Mathematische Leitfäden.
- [8] A. Iwatsuka. Essential selfadjointness of the Schrödinger operators with magnetic fields diverging at infinity. Publ. Res. Inst. Math. Sci., 26(5):841–860, 1990.
- [9] V. Kondratiev, V. Maz'ya, and M. Shubin. Discreteness of spectrum and strict positivity criteria for magnetic Schrödinger operators. *Comm. Part. Diff. Eq.*, 29(3-4):489–521, 2005.
- [10] J. Mehringer and E. Stockmeyer. Confinement-deconfinement transitions for two-dimensional Dirac particles. J. Funct. Anal., 266(4):2225–2250, 2014.
- [11] K. Miller and B. Simon. Quantum magnetic Hamiltonians with remarkable spectral properties. Phys. Rev. Lett., 44:1706–1707, 1980.
- [12] M. Reed and B. Simon. Methods of modern mathematical physics. II. Fourier analysis, self-adjointness. Academic Press, New York, London, 1975.
- [13] B. Thaller. *The Dirac equation*. Texts and Monographs in Physics. Springer-Verlag, Berlin,

Josef Mehringer, Fakultät für Mathematik, Ludwig-Maximilians-Universität München, Theresienstrasse 39, 80333 München, Germany.

 $E ext{-}mail\ address: josef.mehringer@gmail.com}$