

Strong duality in Lasserre's hierarchy for polynomial optimization

Cédric Josz^{1,2}, Didier Henrion^{3,4,5}

Draft of January 24, 2018

Abstract

A polynomial optimization problem (POP) consists of minimizing a multivariate real polynomial on a semi-algebraic set K described by polynomial inequalities and equations. In its full generality it is a non-convex, multi-extremal, difficult global optimization problem. More than an decade ago, J. B. Lasserre proposed to solve POPs by a hierarchy of convex semidefinite programming (SDP) relaxations of increasing size. Each problem in the hierarchy has a primal SDP formulation (a relaxation of a moment problem) and a dual SDP formulation (a sum-of-squares representation of a polynomial Lagrangian of the POP). In this note, when the POP feasibility set K is compact, we show that there is no duality gap between each primal and dual SDP problem in Lasserre's hierarchy, provided a redundant ball constraint is added to the description of set K . Our proof uses elementary results on SDP duality, and it does not assume that K has an interior point.

1 Introduction

Consider the following polynomial optimization problem (POP)

$$\begin{aligned} \inf_x \quad & f(x) := \sum_{\alpha} f_{\alpha} x^{\alpha} \\ \text{s.t.} \quad & g_i(x) := \sum_{\alpha} g_{i,\alpha} x^{\alpha} \geq 0, \quad i = 1, \dots, m \end{aligned} \quad (1)$$

where we use the multi-index notation $x^{\alpha} := x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ for $x \in \mathbb{R}^n$, $\alpha \in \mathbb{N}^n$ and where the data are polynomials $f, g_1, \dots, g_m \in \mathbb{R}[x]$ so that in the above sums only a finite number of coefficients f_{α} and $g_{i,\alpha}$ are nonzero. Assume that the feasibility set

$$K := \{x \in \mathbb{R}^n : g_i(x) \geq 0, i = 1, \dots, m\}$$

¹French transmission system operator Réseau de Transport d'Electricité (RTE), 9 rue de la Porte de Buc, BP 561, F-78000 Versailles, France.

²INRIA Paris-Rocquencourt, BP 105, F-78153 Le Chesnay, France. cedric.josz@inria.fr

³CNRS, LAAS, 7 avenue du colonel Roche, F-31400 Toulouse, France. henrion@laas.fr

⁴Université de Toulouse, LAAS, F-31400 Toulouse, France.

⁵Faculty of Electrical Engineering, Czech Technical University in Prague, Technická 2, CZ-16626 Prague, Czech Republic

is nonempty and bounded, so that the above infimum is attained. To solve POP (1), Lasserre [3, 4] proposed a semidefinite programming (SDP) relaxation hierarchy with guarantees of global and generically finite convergence [5] provided an algebraic assumption holds:

Assumption 1 *There exists a polynomial $u \in \mathbb{R}[x]$ such that $\{x \in \mathbb{R}^n : u(x) \geq 0\}$ is bounded and $u = u_0 + \sum_{i=1}^m u_i g_i$ where polynomials $u_i \in \mathbb{R}[x]$, $i = 0, 1, \dots, m$ are sums of squares (SOS) of other polynomials.*

Assumption 1 can be difficult to check computationally (as the degrees of the SOS multipliers can be arbitrarily large), and it is often replaced by the following slightly stronger assumption:

Assumption 2 *Let $R > 0$ be the radius of an Euclidean ball including set K , and add the redundant ball constraint $g_{m+1}(x) = R^2 - \sum_{i=1}^n x_i^2 \geq 0$ to the description of K .*

Indeed, under Assumption 2, simply choose $u = g_{m+1}$, $u_1 = \dots = u_m = 0$, and $u_{m+1} = 1$ to conclude that Assumption 1 holds as well. In practice, it is often easy to identify a bound R on the radius of the feasibility set K .

Each problem in Lasserre's hierarchy consists of a primal-dual SDP pair, called SDP relaxation, where the primal corresponds to a convex moment relaxation of the original (typically nonconvex) POP, and the dual corresponds to a SOS representation of a polynomial Lagrangian of the POP. The question arises of the absence of duality gap in each SDP relaxation. This is of practical importance because numerical algorithms to solve SDP problems are guaranteed to converge only where there is no duality gap, and sometimes under the stronger assumption that there is a primal or/and dual SDP interior point.

In [6, Example 4.9], Schweighofer provides a two-dimensional POP with bounded K with no interior point for which Assumption 1 holds, yet a duality gap exists at the first SDP relaxation: $\inf_{x_1 x_2} \text{ s.t. } x \in K = \{x \in \mathbb{R}^2 : -1 \leq x_1 \leq 1, x_2^2 \leq 0\}$, with primal SDP value equal to zero and dual SDP value equal to minus infinity. This shows that a stronger assumption is required to ensure no SDP duality gap. A sufficient condition for strong duality has been given in [4]: the interior of the POP feasibility set K should be nonempty. However, this may be too restrictive: in the proof of Lemma 1 in [2] the authors use notationally awkward arguments involving truncated moment matrices to prove the absence of SDP duality gap for a set K with no interior point. This shows that absence of an interior point for K is not necessary for no SDP duality gap, and a weaker assumption is welcome.

Motivated by these observations, in this note we prove that under the basic Assumption 2 on the description of set K , there is no duality gap in the SDP hierarchy. In particular, this covers the cases when K has an empty interior. Our interpretation of this result, and the main message of this contribution, is that in the context of Lasserre's hierarchy for POP, a practically relevant description of a bounded semialgebraic feasibility set must include a redundant ball constraint.

2 Proof of strong duality

For notational convenience, let $g_0(x) = 1 \in \mathbb{R}[x]$ denote the unit polynomial. Define the localizing moment matrix

$$M_{d-d_i}(g_i y) := \left(\sum_{\gamma} g_{i,\gamma} y_{\alpha+\beta+\gamma} \right)_{|\alpha|,|\beta| \leq d-d_i} = \sum_{|\alpha| \leq 2d} A_{i,\alpha} y_{\alpha}$$

where d_i is the smallest integer greater than or equal to half the degree of g_i , for $i = 0, 1, \dots, m$. For $d \geq d_{\min} := \max_{i=0,1,\dots,m} d_i$, the Lasserre hierarchy for POP (1) consists of a primal moment SDP problem

$$(P_d) : \quad \begin{aligned} & \inf_y \quad \sum_{\alpha} f_{\alpha} y_{\alpha} \\ & \text{s.t.} \quad y_0 = 1 \\ & \quad \quad M_{d-d_i}(g_i y) \succeq 0, \quad i = 0, 1, \dots, m \end{aligned}$$

and a dual SOS SDP problem

$$(D_d) : \quad \begin{aligned} & \sup_{z, Z} \quad z \\ & \text{s.t.} \quad f_0 - z = \sum_{i=0}^m \langle A_{i,0}, Z_i \rangle \\ & \quad \quad f_{\alpha} = \sum_{i=0}^m \langle A_{i,\alpha}, Z_i \rangle, \quad 0 < |\alpha| \leq 2d \\ & \quad \quad Z_i \succeq 0, \quad i = 0, 1, \dots, m \end{aligned}$$

where $A \succeq 0$ stands for matrix A positive semidefinite, $\langle A, B \rangle = \text{trace } AB$ is the inner product between two matrices. The primal-dual pair (P_d, D_d) is called the SDP relaxation of order d for POP (1).

Let us define the following sets:

- \mathcal{P}_d : feasible points for P_d ;
- \mathcal{D}_d : feasible points for D_d ;
- $\text{int } \mathcal{P}_d$: strictly feasible points for P_d ;
- $\text{int } \mathcal{D}_d$: strictly feasible points for D_d ;
- \mathcal{P}_d^* : optimal solutions for P_d ;
- \mathcal{D}_d^* : optimal solutions for D_d .

Finally, let us denote by $\text{val } P_d$ the infimum in problem P_d and by $\text{val } D_d$ the supremum in problem D_d . Strong duality holds whenever $\text{val } P_d = \text{val } D_d < \infty$.

Lemma 1 *int \mathcal{P}_d nonempty or int \mathcal{D}_d nonempty implies $\text{val } P_d = \text{val } D_d$.*

Lemma 1 is classical in convex optimization, and it is generally called Slater's condition, see e.g. [7, Theorem 4.1.3].

Lemma 2 \mathcal{P}_d is nonempty and $\text{int } \mathcal{D}_d$ is nonempty if and only if \mathcal{P}_d^* is nonempty and bounded.

A proof of Lemma 2 can be found in [8]. According to Lemmas 1 and 2, \mathcal{P}_d^* nonempty and bounded implies strong duality. This result is also mentioned without proof at the end of [7, Section 4.1.2].

Lemma 3 Under Assumption 2, set \mathcal{P}_d is included in the Euclidean ball of radius $\sum_{k=0}^d R^{2k}$.

Proof: Consider a feasible point $(y_\alpha)_{|\alpha| \leq 2d} \in \mathcal{P}_d$. Let $k \in \mathbb{N}$ be such that $1 \leq k \leq d$. In the SDP problem P_k , the localizing matrix associated to the redundant ball constraint $g_{m+1}(x) = R^2 - \sum_{i=1}^n x_i^2 \geq 0$ reads

$$M_{k-1}(g_{m+1}y) = \left(\sum_{\gamma} g_{m+1,\gamma} y_{\alpha+\beta+\gamma} \right)_{|\alpha|, |\beta| \leq k-1}$$

with trace equal to

$$\begin{aligned} \text{trace } M_{k-1}(g_{m+1}y) &= \sum_{|\alpha| \leq k-1} \sum_{\gamma} g_{m+1,\gamma} y_{2\alpha+\gamma} \\ &= \sum_{|\alpha| \leq k-1} \left(g_{m+1,0} y_{2\alpha} + \sum_{|\gamma|=1} g_{m+1,2\gamma} y_{2\alpha+2\gamma} \right) \\ &= \sum_{|\alpha| \leq k-1} \left(R^2 y_{2\alpha} - \sum_{|\gamma|=1} y_{2(\alpha+\gamma)} \right) \\ &= \sum_{|\alpha| \leq k-1} R^2 y_{2\alpha} - \sum_{|\alpha| \leq k-1, |\gamma|=1} y_{2(\alpha+\gamma)} \\ &= R^2 \left(\sum_{|\alpha| \leq k-1} y_{2\alpha} \right) + y_0 - \sum_{|\alpha| \leq k} y_{2\alpha} \\ &= R^2 \text{trace } M_{k-1}(y) + 1 - \text{trace } M_k(y). \end{aligned}$$

From the structure of the localizing matrix, it holds $M_{k-1}(g_{m+1}y) \succeq 0$ hence $\text{trace } M_{k-1}(g_{m+1}y) \geq 0$ and

$$\text{trace } M_k(y) \leq 1 + R^2 \text{trace } M_{k-1}(y)$$

from which we derive

$$\text{trace } M_d(y) \leq \sum_{k=1}^d R^{2(k-1)} + R^{2d} \text{trace } M_0(y) = \sum_{k=0}^d R^{2k}$$

since $\text{trace } M_0(y) = y_0 = 1$. The norm $\|M_d(y)\|_2$, equal to the maximum eigenvalue of $M_d(y)$, is upper bounded by $\text{trace } M_d(y)$, the sum of the eigenvalues of $M_d(y)$, which are all nonnegative. Moreover

$$\begin{aligned} \|M_d(y)\|_2^2 &= \left\langle \sum_{|\alpha| \leq 2d} A_\alpha y_\alpha, \sum_{|\alpha| \leq 2d} A_\alpha y_\alpha \right\rangle \\ &= \sum_{|\alpha| \leq 2d} \langle A_\alpha, A_\alpha \rangle y_\alpha^2 \quad \text{by orthogonality of the matrices } (A_\alpha)_{|\alpha| \leq 2d} \\ &\geq \sum_{|\alpha| \leq 2d} y_\alpha^2 \quad \text{because } \langle A_\alpha, A_\alpha \rangle \geq 1. \end{aligned}$$

The proof follows then from

$$\sqrt{\sum_{|\alpha| \leq 2d} y_\alpha^2} \leq \|M_d(y)\|_2 \leq \sum_{k=0}^d R^{2k}.$$

□

Theorem 1 *Under Assumption 2, strong duality holds for SDP relaxations of all orders.*

Proof: Let us first show that \mathcal{P}_d is nonempty. Let $v_d(x) := (x^\alpha)_{|\alpha| \leq d}$ and consider a feasible point $x^* \in K$ for POP (1). Then $y^* = v_{2d}(x^*) \in \mathcal{P}_d$ since by construction $M_{d-d_i}(g_i y^*) = g_i(x^*) v_{d-d_i}(x^*) v_{d-d_i}(x^*)^T \succeq 0$ for all i . From Lemma 3, \mathcal{P}_d is bounded and closed, and the objective function in P_d is linear, so we conclude that \mathcal{P}_d^* is nonempty and bounded. According to Lemma 2, \mathcal{D}_d is nonempty, and from Lemma 1 strong duality holds. □

3 Conclusion

We prove that strong duality always holds in Lasserre’s SDP hierarchy for POP on bounded semi-algebraic sets after adding a redundant ball constraint. To preclude numerical troubles with SDP solvers, we advise to systematically add such a ball constraint, combined with an appropriate scaling so that all scaled variables belong to the unit sphere. Without scaling, numerical troubles can occur as well, but they are not due to the presence of a duality gap, see [1] and also the example of [9, Section 6].

References

- [1] D. Henrion, J. B. Lasserre. Detecting global optimality and extracting solutions in GloptiPoly. In: D. Henrion, A. Garulli (eds.). *Positive Polynomials in Control*. Lecture Notes on Control and Information Sciences 312, Springer, Berlin, 2005.
- [2] D. Henrion, J. B. Lasserre. Inner approximations for polynomial matrix inequalities and robust stability regions. *IEEE Transactions on Automatic Control*, 57(6):1456–1467, 2012.
- [3] J. B. Lasserre. Optimisation globale et théorie des moments. *C. R. Acad. Sci. Paris, Série I*, 331:929–934, 2000.
- [4] J. B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM Journal on Optimization*, 11:796–817, 2001.
- [5] J. Nie. Optimality conditions and finite convergence of Lasserre’s hierarchy. *Mathematical Programming*, Ser. A, to appear, 2014.

- [6] M. Schweighofer. Optimization of polynomials on compact semialgebraic sets. *SIAM Journal on Optimization*, 15:805–825, 2005.
- [7] A. Shapiro. *Duality, optimality conditions and perturbation analysis*. In: H. Wolkowicz, R. Saigal, L. Vandenberghe (eds.). *Handbook of Semidefinite Programming. Theory, Algorithms, and Applications*. Kluwer Academic Publishers, Boston, 2000.
- [8] M. Trnovská. Strong duality conditions in semidefinite programming. *Journal of Electrical Engineering*, 56:1–5, 2005.
- [9] H. Waki, M. Nakata, M. Muramatsu. Strange behaviors of interior-point methods for solving semidefinite programming problems in polynomial optimization. *Computational Optimization and Applications*, 53(3):823–844, 2012.