# Fundamental Tensor Operations for Large-Scale Data Analysis in Tensor Train Formats

Namgil Lee<sup>a</sup> and Andrzej Cichocki<sup>a</sup>

<sup>a</sup>Laboratory for Advanced Brain Signal Processing, RIKEN Brain Science Institute, Wako-shi, Saitama, 3510198, Japan

#### Abstract

We discuss extended definitions of linear and multilinear operations such as Kronecker, Hadamard, and contracted products, and establish links between them for tensor calculus. Then we introduce effective low-rank tensor approximation techniques including Candecomp/Parafac (CP), Tucker, and tensor train (TT) decompositions with a number of mathematical and graphical representations. We also provide a brief review of mathematical properties of the TT decomposition as a low-rank approximation technique. With the aim of breaking the curseof-dimensionality in large-scale numerical analysis, we describe basic operations on large-scale vectors, matrices, and high-order tensors represented by TT decomposition. The proposed representations can be used for describing numerical methods based on TT decomposition for solving large-scale optimization problems such as systems of linear equations and symmetric eigenvalue problems.

KEY WORDS: tensor networks; tensor train; matrix product state; matrix product operator; generalized Tucker model; strong Kronecker product; contracted product; multilinear operator; numerical analysis; tensor calculus

## 1 Introduction

Multi-dimensional or multi-way data is prevalent nowadays, which can be represented by tensors. An Nth-order tensor is a multi-way array of size  $I_1 \times I_2 \times \cdots \times I_N$ , where the *n*th dimension or mode is of size  $I_n$ . For example, a tensor can be induced by the discretization of a multivariate function [26]. Given a multivariate function  $f(x_1, \ldots, x_N)$  defined on a domain  $[0, 1]^N$ , we can get a tensor with entries containing the function values at grid points. For another example, we can obtain tensors based on observed data [4, 24]. We can collect and integrate measurements from different modalities by neuroimaging technologies such as functional magnetic resonance imaging (fMRI) and electroencephalography (EEG): subjects, time, frequency, electrodes, task conditions, trials, and so on. Furthermore, high-order tensors can be created by a process called tensorization or quantization [25], by which a large-scale vectors and matrices are reshaped into higher-order tensors.

However, it is impossible to store a high-order tensor because the number of entries,  $I^N$  when  $I = I_1 = I_2 = \cdots = I_N$ , grows exponentially as the order N increases. This is called the "curse-of-dimensionality". Even for I = 2, with N = 50 we obtain  $2^{50} \approx 10^{15}$  entries. Such a huge storage and computational costs required for high-dimensional problems prohibit the use of standard numerical algorithms. To make high-dimensional problems tractable, there have been developed approximation methods including sparse grids [1, 37] and low-rank tensor approximations [15, 16, 26]. In this paper,

we focus on the latter approach, where computational operations are performed on tensor formats, i.e., low-parametric representations of tensors.

In this paper, we consider several tensor decompositions (formats), especially the tensor train (TT) decomposition, which is one of the simplest tensor networks developed with the aim of overcoming the curse-of-dimensionality. Extensive overviews of the modern low-rank tensor approximation techniques are presented in [15, 26]. The TT decomposition is equivalent to the matrix product states (MPS) for open boundary conditions proposed in computational physics, and it has taken a key role in density matrix renormalization group (DMRG) methods for simulating quantum manybody systems [36, 40]. It was later re-discovered in numerical analysis community [20, 33, 35]. The TT-based numerical algorithms can accomplish algorithmic stability and adaptive determination of ranks by employing the singular value decomposition (SVD) [33]. Its scope of application is quickly expanding for addressing high-dimensional problems such as multi-dimensional integrals, stochastic and parametric PDEs, computational finance, and machine learning [15]. On the other hand, a comprehensive survey on traditional low-rank tensor approximation techniques such as Candecomp/Parafac (CP) and Tucker decompositions is presented in [6, 24].

Despite the large interest in high-order tensors in TT format, mathematical representations of the TT tensors are usually limited to the representations based on scalar operations on matrices and vectors, which leads to complex and tedious index notation in the tensor calculus. For example, a TT tensor is defined by each entry represented as products of matrices [20, 33]. On the other hand, representations of traditional low-rank tensor decompositions have been developed based on multilinear operations such as the Kronecker product, Khatri-Rao product, Hadamard product, and mode-*n* multilinear product [4, 24], which enables coordinate-free notation. Through the utilization of the multilinear operations, the traditional tensor decompositions expanded the area of application to chemometrics, signal processing, numerical linear algebra, computer vision, data mining, graph analysis, and neuroscience [24].

In this work, we develop extended definitions of multilinear operations on tensors. Based on the tensor operations, we provide a number of new and useful representations of the TT decomposition. We also provide graphical representations of TT decompositions, motivated by [20], which are help-ful in understanding the underlying principles and TT-based numerical algorithms. Based on the TT representations of large-scale vectors and matrices, we show that the basic numerical operations such as the addition, contraction, matrix-by-vector product, and quadratic form are conveniently described by the proposed representations. We demonstrate the usefulness of the proposed tensor operations in tensor calculus by giving a proof of orthonormality of the so-called frame matrices. Moreover, we derive explicit representations of localized linear maps in TT format that have been implicitly presented in matrix forms in the literature in the context of alternating linear scheme (ALS) for solving various optimization problems. The suggested mathematical operations and TT representations can be further applied to describing TT-based numerical methods such as the solutions to large-scale systems of linear equations and eigenvalue problems [20].

This paper is organized as follows. In Section 2, we introduce notations for tensors and definitions for tensor operations. In Section 3, we provide the mathematical and graphical representations of the TT decomposition. We also review mathematical properties of the TT decomposition as a low-rank approximation. In Section 4, we describe basic numerical operations on tensors represented by TT decomposition such as the addition, Hadamard product, matrix-by-vector multiplication, and quadratic form in terms of the multilinear operations and TT representations. Discussion and conclusions are given in Section 5.

Table 1: Notations for tensors

Notation	Description
$\underline{\mathbf{X}} \in \mathbb{R}^{I_1  imes I_2  imes \cdots  imes I_N}$	Nth-order tensor of size $I_1 \times I_2 \times \cdots \times I_N$
$x, \mathbf{x}, \mathbf{X}$	scalar, vector, and matrix
$x_{i_1,i_2,\ldots,i_N}, \underline{\mathbf{X}}(i_1,i_2,\ldots,i_N)$	$(i_1, i_2, \ldots, i_N)$ th entry of <b>X</b>
$\mathbf{x}_{:,i_2,i_3,\ldots,i_N},  \underline{\mathbf{X}}(:,i_2,i_3,\ldots,i_N)$	mode-1 fiber of $\underline{\mathbf{X}}$
$\mathbf{X}_{:,:,i_3,i_4,\ldots,i_N},  \underline{\mathbf{X}}(:,:,i_3,i_4,\ldots,i_N)$	frontal slice of $\underline{\mathbf{X}}$
$\mathbf{X}_{(n)} \in \mathbb{R}^{I_n  imes I_1 I_2 \cdots I_{n-1} I_{n+1} \cdots I_N}$	mode- $n$ unfolding of $\underline{\mathbf{X}}$
$\mathbf{X}_{< n>} \in \mathbb{R}^{I_{1}I_{2}\cdots I_{n}  imes I_{n+1}\cdots I_{N}}$	mode- $(1, 2, \ldots, n)$ unfolding of $\underline{\mathbf{X}}$
$\underline{\mathbf{G}}, \underline{\mathbf{G}}^{(n)}, \underline{\mathbf{X}}^{(n)}, \underline{\mathbf{A}}^{(n)}$	core tensors and factor matrices/tensors
$R, R_n$	ranks
$\overline{i_1 i_2 \cdots i_N}$	multi-index, $i_N + (i_{N-1} - 1)I_N + \dots + (i_1 - 1)I_2I_3 \cdots I_N$

## 2 Notations for tensors and tensor operations

The notations in this paper follow the convention provided by [4, 24]. Table 1 summarizes the notations for tensors. Scalars, vectors, and matrices are denoted by lowercase, lowercase bold, and uppercase bold letters  $\mathbf{x}$ ,  $\mathbf{x}$ , and  $\mathbf{X}$ , respectively. Tensors are denoted by underlined uppercase bold letters  $\mathbf{X}$ . The  $(i_1, i_2, \ldots, i_N)$ th entry of a tensor  $\mathbf{X}$  of size  $I_1 \times I_2 \times \cdots \times I_N$  is denoted by  $x_{i_1,i_2,\ldots,i_N}$  or  $\mathbf{X}(i_1, i_2, \ldots, i_N)$ . A subtensor of  $\mathbf{X}$  obtained by fixing the indices  $i_3, i_4, \ldots, i_N$  is denoted by  $\mathbf{X}_{:,:,i_3,i_4,\ldots,i_N}$  or  $\mathbf{X}(:,:,i_3,i_4,\ldots,i_N)$ . We may omit ':' as  $\mathbf{X}_{i_3,i_4,\ldots,i_N}$  if the omitted indices are clear to readers. We define the multi-index by  $i_1i_2\cdots i_N = i_N + (i_{N-1} - 1)I_N + \cdots + (i_1 - 1)I_2I_3\cdots I_N$  for the indices  $i_n = 1, 2, \ldots, I_n$ ,  $n = 1, 2, \ldots, N$ . By using this notation, we can write an entry of a Kronecker product as  $(\mathbf{a} \otimes \mathbf{b})_{ii} = a_i b_j$ .

Moreover, it is important to note that in this paper the vectorization and matricization are defined in accordance with the multi-index notation. That is, for a tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$ , let  $\operatorname{vec}(\underline{\mathbf{X}}) \in \mathbb{R}^{I_1 I_2 \cdots I_N}$ ,  $\mathbf{X}_{(n)} \in \mathbb{R}^{I_n \times I_1 I_2 \cdots I_{n-1} I_{n+1} \cdots I_N}$ , and  $\mathbf{X}_{\langle n \rangle} \in \mathbb{R}^{I_1 I_2 \cdots I_n \times I_{n+1} \cdots I_N}$  denote the vectorization, mode-*n* matricization, and mode- $(1, 2, \ldots, n)$  matricization of  $\underline{\mathbf{X}}$ , whose entries are defined by

$$\left(\operatorname{vec}(\underline{\mathbf{X}})\right)_{\overline{i_1i_2\cdots i_N}} = \left(\mathbf{X}_{(n)}\right)_{i_n,\overline{i_1i_2\cdots i_{n-1}i_{n+1}\cdots i_N}} = \left(\mathbf{X}_{}\right)_{\overline{i_1i_2\cdots i_n},\overline{i_{n+1}\cdots i_N}} = \underline{\mathbf{X}}(i_1,i_2,\ldots,i_N)$$
(1)

for n = 1, ..., N.

Table 2 summarizes the notations and definitions for basic tensor operations used in this paper.

### 2.1 Graphical representations of tensors

It is quite useful to visualize tensors and related operations by tensor network diagrams, e.g., see [20], and references in [15]. Figure 1(a), (b), and (c) illustrate tensor network diagrams representing a vector, a matrix, and a 3rd-order tensor. In each graph, the number of edges connected to a node indicates the order of the tensor, and the mode size can be shown by the label on each edge. Figure 1(d) represents the singular value decomposition of a matrix. The orthonormalized matrices are represented by half-filled circles and the diagonal matrix by a circle with slash inside. Figure 1(e) represents the mode-3 product,  $\underline{\mathbf{A}} \times_3 \mathbf{B}$ , of a tensor  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times I_3}$  with a matrix  $\mathbf{B} \in \mathbb{R}^{J_1 \times J_2}$  ( $I_3 = J_2$ ). Figure 1(f) represents the contracted product,  $\underline{\mathbf{A}} \times^1 \underline{\mathbf{B}}$ , of a tensor  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times I_3}$  with a tensor  $\underline{\mathbf{B}} \in \mathbb{R}^{J_1 \times J_2 \times J_3}$  ( $I_3 = J_1$ ).

Table 2: Notations and definitions for basic tensor operations

Notation	Description
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \otimes \underline{\mathbf{B}}$	Kronecker product of $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$ and $\underline{\mathbf{B}} \in \mathbb{R}^{J_1 \times \cdots \times J_N}$ yields a
	tensor $\underline{\mathbf{C}}$ of size $I_1 J_1 \times \cdots \times I_N J_N$ with entries $\underline{\mathbf{C}}(\overline{i_1 j_1}, \ldots, \overline{i_N j_N}) =$
	$\underline{\mathbf{A}}(i_1,\ldots,i_N)\underline{\mathbf{B}}(j_1,\ldots,j_N)$
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \circledast \underline{\mathbf{B}}$	Hadamard (elementwise) product of $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$ and $\underline{\mathbf{B}} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$
	yields a tensor $\underline{\mathbf{C}}$ of size $I_1 \times \cdots \times I_N$ with entries $\underline{\mathbf{C}}(i_1, \ldots, i_N) =$
	$\underline{\mathbf{A}}(i_1,\ldots,i_N)\underline{\mathbf{B}}(i_1,\ldots,i_N)$
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \circ \underline{\mathbf{B}}$	Outer product of $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times \cdots \times I_M}$ and $\underline{\mathbf{B}} \in \mathbb{R}^{J_1 \times \cdots \times J_N}$ yields a tensor $\underline{\mathbf{C}}$ of
	size $I_1 \times \cdots \times I_M \times J_1 \times \cdots \times J_N$ with entries $\underline{\mathbf{C}}(i_1, \ldots, i_M, j_1, \ldots, j_N) =$
	$\underline{\mathbf{A}}(i_1,\ldots,i_M)\underline{\mathbf{B}}(j_1,\ldots,j_N)$
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \oplus \underline{\mathbf{B}}$	Direct sum of $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$ and $\underline{\mathbf{B}} \in \mathbb{R}^{J_1 \times \cdots \times J_N}$ yields a tensor $\underline{\mathbf{C}}$ of size
	$(I_1 + J_1) \times \cdots \times (I_N + J_N)$ with entries $\underline{\mathbf{C}}(k_1, \ldots, k_N) = \underline{\mathbf{A}}(k_1, \ldots, k_N)$ if
	$1 \leq k_n \leq I_n \ \forall n, \ \underline{\mathbf{C}}(k_1, \dots, k_N) = \underline{\mathbf{B}}(k_1 - I_1, \dots, k_N - I_N) \ \text{if} \ I_n < k_n \leq I_n$
	$I_n + J_n \ \forall n, \text{ and } \underline{\mathbf{C}}(k_1, \dots, k_N) = 0 \text{ otherwise}$
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \boxtimes \underline{\mathbf{B}}$	Partial Kronecker product of factor tensors $\underline{\mathbf{A}} \in \mathbb{R}^{R_1 \times \cdots \times R_M \times I_1 \times \cdots \times I_N}$ and
	$\underline{\mathbf{B}} \in \mathbb{R}^{S_1 \times \cdots \times S_M \times I_1 \times \cdots \times I_N} \text{ yields a tensor } \underline{\mathbf{C}} \text{ of size } R_1 S_1 \times \cdots \times R_M S_M \times I_1 \times \cdots \times I_N S_M \times I_1 \times \cdots \times S_M \times S_M \times I_1 \times \cdots \times S_M \times S_M$
	$\cdots \times I_N$ with subtensors $\underline{\mathbf{C}}(:,\ldots,:,i_1,\ldots,i_N) = \underline{\mathbf{A}}(:,\ldots,:,i_1,\ldots,i_N) \otimes \underline{\mathbf{B}}(:$
	$,\ldots,:,i_1,\ldots,i_N)$
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \boxplus \underline{\mathbf{B}}$	Partial direct sum of factor tensors $\underline{\mathbf{A}} \in \mathbb{R}^{K_1 \times \cdots \times K_M \times I_1 \times \cdots \times I_N}$ and $\underline{\mathbf{B}} \in \mathbb{R}^{K_1 \times \cdots \times K_M \times I_1 \times \cdots \times I_N}$
	$\mathbb{R}^{S_1 \times \cdots \times S_M \times I_1 \times \cdots \times I_N}$ yields a tensor <u>C</u> of size $(R_1 + S_1) \times \cdots \times (R_M + S_M) \times$
	$I_1 \times \cdots \times I_N$ with subtensors $\underline{\mathbf{C}}(:, \dots, :, i_1, \dots, i_N) = \underline{\mathbf{A}}(:, \dots, :, i_1, \dots, i_N) \oplus$
	$\underline{\mathbf{B}}(:,\ldots,:,i_1,\ldots,i_N)$
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \times_n \mathbf{B}$	Mode- <i>n</i> product of tensor $\underline{\mathbf{A}} \in \mathbb{R}^{n \times n \times n}$ and matrix $\mathbf{B} \in \mathbb{R}^{n \times n \times n}$ yields a
	tensor $\underline{C}$ of size $I_1 \times \cdots \times I_{n-1} \times J \times I_{n+1} \times \cdots \times I_N$ with mode- <i>n</i> fibers
$C  A \overline{X} $ h	$\underline{\mathbf{C}}(i_1,\ldots,i_{n-1},:,i_{n+1},\ldots,i_N) = \mathbf{B}\underline{\mathbf{A}}(i_1,\ldots,i_{n-1},:,i_{n+1},\ldots,i_N)$ Mode m (noter) module of tensor $\mathbf{A} \in \mathbb{D}^{I_1 \times \cdots \times I_N}$ and note the $\in \mathbb{D}^{I_n}$
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \times_n \mathbf{D}$	wields a topsor $\mathbf{C}$ of size $L \times \dots \times L \times \times L$ with entries
	yields a tensor $\underline{\underline{C}}$ of size $I_1 \times \cdots \times I_{n-1} \times I_{n+1} \times \cdots \times I_N$ with entries $C(i_1, \dots, i_{n-1}) = \mathbf{b}^\top \mathbf{A}(i_1, \dots, i_{n-1})$
$\begin{bmatrix} \mathbf{a} & \mathbf{i} \\ \mathbf{k} \end{bmatrix}$	$\underline{\mathbf{U}}(\iota_1,\ldots,\iota_{n-1},\iota_{n+1},\ldots,\iota_N) = \mathbf{U}  \underline{\mathbf{A}}(\iota_1,\ldots,\iota_{n-1},\ldots,\iota_N)$
$\left[\underline{\mathbf{G}};\underline{\mathbf{A}}^{(1)},\ldots,\underline{\mathbf{A}}^{(1)}\right]$	Multilinear operator for tensors <u><b>G</b></u> and <u><b>A</b></u> <sup>(<math>n</math>)</sup> , $n = 1,, N$ , defined by (4)
$\underline{\mathbf{C}} = \underline{\mathbf{A}} \times^{1} \underline{\mathbf{B}}$	Mode- $(M, 1)$ contracted product of tensors $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times \cdots \times I_M}$ and $\underline{\mathbf{B}} \in \mathbb{R}^{I_1 \times \cdots \times I_M}$
	$\mathbb{R}^{J_1 \times J_2 \times J_3 \times \cdots \times J_N}$ with $I_M = J_1$ yields a tensor $\underline{\mathbf{C}}$ of size $I_1 \times \mathbf{C}$
	$\cdots \times I_{M-1} \times J_2 \times \cdots \times J_N$ with entries $\underline{\mathbf{C}}(i_1, \ldots, i_{M-1}, j_2, \ldots, j_N) =$
	$\sum_{i_M=1}^{I_M} \underline{\mathbf{A}}(i_1, \dots, i_M) \underline{\mathbf{B}}(i_M, j_2, \dots, j_N)$
$\mathbf{C} = \mathbf{A}   \otimes   \mathbf{B}$	Strong Kronecker product of two block matrices $\mathbf{A} = [\mathbf{A}_{r_1, r_2}] \in \mathbb{R}^{R_1 I_1 \times R_2 I_2}$
	and $\mathbf{B} = [\mathbf{B}_{r_2,r_3}] \in \mathbb{R}^{R_2 J_1 \times R_3 J_2}$ yields a block matrix $\mathbf{C} = [\mathbf{C}_{r_1,r_3}] \in$
	$\mathbb{R}^{R_1I_1J_1 \times R_3I_2J_2}$ with blocks $\mathbf{C}_{r_1,r_3} = \sum_{r_2=1}^{R_2} \mathbf{A}_{r_1,r_2} \otimes \mathbf{B}_{r_2,r_3}$
$\underline{\mathbf{C}} = \underline{\mathbf{A}}   \otimes   \underline{\mathbf{B}}$	Strong Kronecker product of two block tensors $\underline{\mathbf{A}} = [\underline{\mathbf{A}}_{r_1,r_2}] \in$
	$\mathbb{R}^{R_1I_1 \times R_2I_2 \times I_3}$ and $\underline{\mathbf{B}} = [\underline{\mathbf{B}}_{r_2,r_3}] \in \mathbb{R}^{R_2J_1 \times R_3J_2 \times J_3}$ yields a block tensor $\underline{\mathbf{C}} =$
	$[\underline{\mathbf{C}}_{r_1,r_3}] \in \mathbb{R}^{R_1 I_1 J_1 \times R_3 I_2 J_2 \times I_3 J_3}$ with blocks $\underline{\mathbf{C}}_{r_1,r_2} = \sum_{r_2-1}^{R_2} \underline{\mathbf{A}}_{r_1,r_2} \otimes \underline{\mathbf{B}}_{r_2,r_3}$
$Tr\left(\mathbf{X}\right)$	Partial trace operator $Tr: \mathbb{R}^{R \times I_1 \times \cdots \times I_N \times R} \to \mathbb{R}^{I_1 \times \overline{I_2} \times \cdots \times \overline{I_N}}$ is defined by
. ,	$Tr(\underline{\mathbf{X}}) = \sum_{r=1}^{R} \underline{\mathbf{X}}(r, i_1, \dots, i_N, r)$



Figure 1: Graphical representations for (a) a vector (b) a matrix, (c) a 3rd-order tensor, (d) singular value decomposition of an  $I \times J$  matrix, (e) mode-3 product of a 3rd-order tensor with a matrix, and (f) contracted product of two 3rd-order tensors.

## 2.2 Kronecker, Hadamard, and outer products

Definitions for traditional matrix-matrix product operations such as the Kronecker, Hadamard, and outer products can immediately be generalized to tensor-tensor products.

**Definition 2.1** (Kronecker product). The Kronecker product of tensors  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  and  $\underline{\mathbf{B}} \in \mathbb{R}^{J_1 \times J_2 \times \cdots \times J_N}$  is defined by

$$\mathbf{C} = \mathbf{A} \otimes \mathbf{B} \in \mathbb{R}^{I_1 J_1 \times I_2 J_2 \times \cdots \times I_N J_N}$$

with entries

$$\underline{\mathbf{C}}(\overline{i_1j_1},\overline{i_2j_2},\ldots,\overline{i_Nj_N}) = \underline{\mathbf{A}}(i_1,i_2,\ldots,i_N)\underline{\mathbf{B}}(j_1,j_2,\ldots,j_N)$$

for  $i_n = 1, 2, \dots, I_n, j_n = 1, 2, \dots, J_n, n = 1, \dots, N$ .

**Definition 2.2** (Hadamard product). The Hadamard (elementwise) product of  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$ and  $\underline{\mathbf{B}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  is defined by

$$\underline{\mathbf{C}} = \underline{\mathbf{A}} \circledast \underline{\mathbf{B}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$$

with entries

$$\underline{\mathbf{C}}(i_1, i_2, \dots, i_N) = \underline{\mathbf{A}}(i_1, i_2, \dots, i_N) \underline{\mathbf{B}}(i_1, i_2, \dots, i_N)$$

for  $i_n = 1, 2, \dots, I_n, n = 1, \dots, N$ .

**Definition 2.3** (Outer product). The outer product of  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_M}$  and  $\underline{\mathbf{B}} \in \mathbb{R}^{J_1 \times J_2 \times \cdots \times J_N}$  is defined by

$$\underline{\mathbf{C}} = \underline{\mathbf{A}} \circ \underline{\mathbf{B}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_M \times J_1 \times J_2 \times \cdots \times J_N}$$

with entries

$$\underline{\mathbf{C}}(i_1, i_2, \dots, i_M, j_1, j_2, \dots, j_N) = \underline{\mathbf{A}}(i_1, i_2, \dots, i_M)\underline{\mathbf{B}}(j_1, j_2, \dots, j_N)$$

for  $i_m = 1, 2, \dots, I_m, m = 1, \dots, M, j_n = 1, 2, \dots, J_n, n = 1, \dots, N.$ 

Note that an *N*th-order tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  is rank-one if it is written as the outer product of *N* vectors

$$\underline{\mathbf{X}} = \mathbf{x}^{(1)} \circ \mathbf{x}^{(2)} \circ \cdots \circ \mathbf{x}^{(N)}.$$

In general, Nth-order tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  can be represented as a sum of rank-one tensors as (so called CP or PARAFAC [24])

$$\underline{\mathbf{X}} = \sum_{r=1}^{R} \mathbf{x}_{r}^{(1)} \circ \mathbf{x}_{r}^{(2)} \circ \cdots \circ \mathbf{x}_{r}^{(N)}$$

The smallest number R of the rank-one tensors that produce  $\underline{\mathbf{X}}$  is called the tensor rank of  $\underline{\mathbf{X}}$  [24]. We can define a tensor operation between rank-one tensors and generalize it to sums of rank-one tensors. For example, let  $\underline{\mathbf{A}} = \mathbf{a}^{(1)} \circ \mathbf{a}^{(2)} \circ \cdots \circ \mathbf{a}^{(N)}$  and  $\underline{\mathbf{B}} = \mathbf{b}^{(1)} \circ \mathbf{b}^{(2)} \circ \cdots \circ \mathbf{b}^{(N)}$  denote two rank-one tensors, and let  $\mathbf{a} \circledast \mathbf{b} = (a_i b_i)$  denote Hadamard (elementwise) product of vectors, then,

• the Kronecker product  $\underline{\mathbf{A}} \otimes \underline{\mathbf{B}}$  can be expressed by

$$\underline{\mathbf{A}} \otimes \underline{\mathbf{B}} = \left(\mathbf{a}^{(1)} \otimes \mathbf{b}^{(1)}\right) \circ \left(\mathbf{a}^{(2)} \otimes \mathbf{b}^{(2)}\right) \circ \cdots \circ \left(\mathbf{a}^{(N)} \otimes \mathbf{b}^{(N)}\right),$$

• the Hadamard product by

$$\underline{\mathbf{A}} \circledast \underline{\mathbf{B}} = \left(\mathbf{a}^{(1)} \circledast \mathbf{b}^{(1)}\right) \circ \left(\mathbf{a}^{(2)} \circledast \mathbf{b}^{(2)}\right) \circ \cdots \circ \left(\mathbf{a}^{(N)} \circledast \mathbf{b}^{(N)}\right),$$

• and the outer product by

$$\underline{\mathbf{A}} \circ \underline{\mathbf{B}} = \mathbf{a}^{(1)} \circ \cdots \circ \mathbf{a}^{(N)} \circ \mathbf{b}^{(1)} \circ \cdots \circ \mathbf{b}^{(N)}$$

However, the problem of determining the tensor rank of a specific tensor is NP-hard in general if the order is larger than 2 [18]. So, for practical applications, we will define tensor operations by using index notation and provide examples with rank-one tensors.

### 2.3 Direct sum

The direct sum of matrices **A** and **B** is defined by

$$\mathbf{A} \oplus \mathbf{B} = \operatorname{diag}(\mathbf{A}, \mathbf{B}) = \begin{bmatrix} \mathbf{A} & \mathbf{0} \\ \mathbf{0} & \mathbf{B} \end{bmatrix}.$$

A generalization of the direct sum to tensors is defined as follows.

**Definition 2.4** (Direct sum). The direct sum of tensors  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  and  $\underline{\mathbf{B}} \in \mathbb{R}^{J_1 \times J_2 \times \cdots \times J_N}$  is defined by

$$\underline{\mathbf{C}} = \underline{\mathbf{A}} \oplus \underline{\mathbf{B}} \in \mathbb{R}^{(I_1 + J_1) \times (I_2 + J_2) \times \dots \times (I_N + J_N)}$$

with entries

$$\underline{\mathbf{C}}(k_1, k_2, \dots, k_N) = \begin{cases} \underline{\mathbf{A}}(k_1, k_2, \dots, k_N) & \text{if } 1 \le k_n \le I_n \ \forall n \\ \underline{\mathbf{B}}(k_1 - I_1, k_2 - I_2, \dots, k_N - I_N) & \text{if } I_n < k_n \le I_n + J_n \ \forall n \\ 0 & \text{otherwise.} \end{cases}$$

As special cases, the direct sum of vectors  $\mathbf{a} \in \mathbb{R}^{I}$  and  $\mathbf{b} \in \mathbb{R}^{J}$  is the concatenated vector  $\mathbf{a} \oplus \mathbf{b} \in \mathbb{R}^{I+J}$ , and the direct sum of matrices  $\mathbf{A} \in \mathbb{R}^{I_1 \times I_2}$  and  $\mathbf{B} \in \mathbb{R}^{J_1 \times J_2}$  is the block diagonal matrix  $\mathbf{A} \oplus \mathbf{B} = \text{diag}(\mathbf{A}, \mathbf{B}) \in \mathbb{R}^{(I_1+J_1) \times (J_1+J_2)}$ . We suppose that the direct sum of scalars  $a, b \in \mathbb{R}$  is the addition  $a \oplus b = a + b \in \mathbb{R}$ . Similarly, direct sum of two 3rd-order tensors is a block diagonal 3rd-order tensor, as illustrated in Figure 2.



Figure 2: Illustration of direct sum  $\underline{\mathbf{A}} \oplus \underline{\mathbf{B}}$  of 3rd-order tenosors  $\underline{\mathbf{A}}$  and  $\underline{\mathbf{B}}$ .

## 2.4 Partial Kronecker product and partial direct sum: General operations for factor tensors

A low-rank tensor decomposition method approximately represents a tensor as contraction of a collection of factor (core) tensors, which helps to reduce the number of representation parameters. A *factor tensor* refers to a tensor which forms such a collection in a low-rank tensor representation [11]. In general, each mode of a factor tensor can be classified as either a physical (spatial) mode or an auxiliary mode [5, 12]. In this paper, sizes of auxiliary modes are denoted by R or S, and the corresponding indices are denoted by r or s. For example, if a factor tensor  $\underline{X}$  is mentioned to have size  $R_1 \times I_1 \times R_2 \times I_2$ , then it implies that the modes 1 and 3 are auxiliary modes and the rest are physical modes. A factor tensor may have both types of or only one type of the physical and auxiliary modes.

The partial Kronecker product and the partial direct sum, which will be defined below, are generalizations of the Kronecker product and the direct sum to factor tensors.

**Definition 2.5** (Partial Kronecker product and partial direct sum). Let  $\underline{\mathbf{A}} \in \mathbb{R}^{R_1 \times R_2 \times \cdots \times R_M \times I_1 \times \cdots \times I_N}$ and  $\underline{\mathbf{B}} \in \mathbb{R}^{S_1 \times S_2 \times \cdots \times S_M \times I_1 \times \cdots \times I_N}$  be two factor tensors with M auxiliary modes and N physical modes. The partial Kronecker product of  $\underline{\mathbf{A}}$  and  $\underline{\mathbf{B}}$  is defined by

$$\mathbf{C} = \mathbf{A} \boxtimes \mathbf{B} \in \mathbb{R}^{R_1 S_1 \times R_2 S_2 \times \dots \times R_M S_M \times I_1 \times \dots \times I_N}$$

with subtensors

$$\underline{\mathbf{C}}(:,\ldots,:,i_1,\ldots,i_N) = \underline{\mathbf{A}}(:,\ldots,:,i_1,\ldots,i_N) \otimes \underline{\mathbf{B}}(:,:,\ldots,:,i_1,\ldots,i_N)$$

for  $i_n = 1, 2, \ldots, I_n$ ,  $n = 1, \ldots, N$ . Similarly, the partial direct sum of <u>A</u> and <u>B</u> is defined by

$$\mathbf{C} = \mathbf{A} \boxplus \mathbf{B} \in \mathbb{R}^{(R_1 + S_1) \times (R_2 + S_2) \times \dots \times (R_M + S_M) \times I_1 \times \dots \times I_N}$$

with subtensors

$$\underline{\mathbf{C}}(:,\ldots,:,i_1,\ldots,i_N) = \underline{\mathbf{A}}(:,\ldots,:,i_1,\ldots,i_N) \oplus \underline{\mathbf{B}}(:,:,\ldots,:,i_1,\ldots,i_N)$$

for  $i_n = 1, 2, \dots, I_n, n = 1, \dots, N$ .

In the above definition, if M = 0, i.e., the tensors  $\underline{\mathbf{A}}$  and  $\underline{\mathbf{B}}$  have only physical modes, then the partial Kronecker product and the partial direct sum are equivalent to the Hadamard (elementwise) product and the elementwise addition, respectively. On the other hand, if N = 0, i.e., there are only auxiliary modes, then the partial Kronecker product and the partial direct sum are equivalent to the Kronecker product and the direct sum, respectively.

## 2.5 Multilinear operator

The mode-*n* product of a tensor  $\underline{\mathbf{G}} \in \mathbb{R}^{R_1 \times R_2 \times \cdots \times R_N}$  and a matrix  $\mathbf{A} \in \mathbb{R}^{I_n \times R_n}$  is a multilinear operator defined by [24]

$$\underline{\mathbf{X}} = \underline{\mathbf{G}} \times_n \mathbf{A} \in \mathbb{R}^{R_1 \times R_2 \times \dots \times R_{n-1} \times I_n \times R_{n+1} \times \dots \times R_N}$$
(2)

with entries

$$\underline{\mathbf{X}}(r_1, r_2, \dots, r_{n-1}, i_n, r_{n+1}, \dots, r_N) = \sum_{r_n=1}^{R_n} \underline{\mathbf{G}}(r_1, r_2, \dots, r_N) \mathbf{A}(i_n, r_n).$$

A few selected properties of the mode-n product are listed as follows<sup>1</sup>.

**Proposition 2.1** ([7, 23, 24]). Let  $\mathbf{G} \in \mathbb{R}^{R_1 \times R_2 \times \cdots \times R_N}$  be an Nth-order tensor. Then

- (a)  $\underline{\mathbf{G}} \times_m \mathbf{A} \times_n \mathbf{B} = \underline{\mathbf{G}} \times_n \mathbf{B} \times_m \mathbf{A}$  for  $m \neq n$ .
- (b)  $\underline{\mathbf{G}} \times_n \mathbf{A} \times_n \mathbf{B} = \underline{\mathbf{G}} \times_n \mathbf{B}\mathbf{A}$ .
- (c) If  $\mathbf{A}$  has full column rank, then

$$\underline{\mathbf{X}} = \underline{\mathbf{G}} \times_n \mathbf{A} \Rightarrow \underline{\mathbf{G}} = \underline{\mathbf{X}} \times_n \mathbf{A}^{\dagger},$$

where  $\mathbf{A}^{\dagger}$  is the Moore-Penrose pseudoinverse of  $\mathbf{A}$ .

(d) If  $\mathbf{A} \in \mathbb{R}^{I \times R_n}$ , then

$$\underline{\mathbf{X}} = \underline{\mathbf{G}} \times_n \mathbf{A} \Leftrightarrow \mathbf{X}_{(n)} = \mathbf{A}\mathbf{G}_{(n)}.$$

(e) If  $\mathbf{A}^{(n)} \in \mathbb{R}^{I_n \times R_n}$  for all  $n = 1, \dots, N$ , we have, for  $n \in \{1, \dots, N\}$ ,

$$\underline{\mathbf{X}} = \underline{\mathbf{G}} \times_1 \mathbf{A}^{(1)} \times_2 \mathbf{A}^{(2)} \cdots \times_N \mathbf{A}^{(N)} \Leftrightarrow$$
$$\mathbf{X}_{(n)} = \mathbf{A}^{(n)} \mathbf{G}_{(n)} \left( \mathbf{A}^{(1)} \otimes \cdots \otimes \mathbf{A}^{(n-1)} \otimes \mathbf{A}^{(n+1)} \otimes \cdots \otimes \mathbf{A}^{(N)} \right)^\top.$$

(f) If  $\mathbf{A}^{(n)} \in \mathbb{R}^{I_n \times R_n}$  for all  $n = 1, \dots, N$ , we have, for  $n \in \{1, \dots, N\}$ ,

$$\underline{\mathbf{X}} = \underline{\mathbf{G}} \times_1 \mathbf{A}^{(1)} \times_2 \mathbf{A}^{(2)} \cdots \times_N \mathbf{A}^{(N)} \Leftrightarrow$$
$$\mathbf{X}_{\langle n \rangle} = \left(\mathbf{A}^{(1)} \otimes \cdots \otimes \mathbf{A}^{(n)}\right) \mathbf{G}_{\langle n \rangle} \left(\mathbf{A}^{(n+1)} \otimes \cdots \otimes \mathbf{A}^{(N)}\right)^\top.$$

Kolda and Bader [24] further introduced a multilinear operator called the Tucker operator [23] to simplify the expression for the mode-*n* product. The Tucker operator of a tensor  $\underline{\mathbf{G}} \in \mathbb{R}^{R_1 \times R_2 \times \cdots \times R_N}$  and matrices  $\mathbf{A}^{(n)} \in \mathbb{R}^{I_n \times R_n}$ ,  $n = 1, \ldots, N$ , is defined by

$$\left[\!\left[\underline{\mathbf{G}};\mathbf{A}^{(1)},\mathbf{A}^{(2)},\ldots,\mathbf{A}^{(N)}\right]\!\right] = \underline{\mathbf{G}} \times_1 \mathbf{A}^{(1)} \times_2 \mathbf{A}^{(2)} \times_3 \cdots \times_N \mathbf{A}^{(N)} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}.$$
(3)

Here, we generalize this to a multilinear operator between tensors.

<sup>&</sup>lt;sup>1</sup> In Proposition 2.1(e) and (f), factors of Kronecker products are in a different (reversed) order, e.g., increasing from 1 to N, compared to the order in the literature [7, 23, 24], e.g., decreasing from N to 1, due to the definition of matricizations in (1).



Figure 3: Tensor network diagrams for (a) multilinear operator  $[\![\underline{\mathbf{G}};\underline{\mathbf{A}}^{(1)},\ldots,\underline{\mathbf{A}}^{(N)}]\!]$  with Nth-order tensors  $\underline{\mathbf{G}}$  and 4th-order tensors  $\underline{\mathbf{A}}^{(1)},\ldots,\underline{\mathbf{A}}^{(N)}$ , and (b) multilinear operator  $[\![\underline{\mathbf{G}};\mathbf{I}_{R_1},\underline{\mathbf{A}},\mathbf{I}_{R_3}]\!]$  with rank-one tensors  $\underline{\mathbf{G}} = \mathbf{g}^{(1)} \circ \mathbf{g}^{(2)} \circ \mathbf{g}^{(3)}$  and  $\underline{\mathbf{A}} = \mathbf{a}^{(1)} \circ \mathbf{a}^{(2)} \circ \cdots \circ \mathbf{a}^{(5)}$  as described in Example 2.1.

**Definition 2.6** (Multilinear operator). Let  $N \ge 1$  and  $M_n \ge 0$  for n = 1, ..., N. For an Nthorder tensor  $\underline{\mathbf{G}} \in \mathbb{R}^{R_1 \times R_2 \times \cdots \times R_N}$  and  $(M_n + 1)$ th-order tensors  $\underline{\mathbf{A}}^{(n)} \in \mathbb{R}^{I_{n,1} \times I_{n,2} \times \cdots \times I_{n,M_n} \times R_n}$ , n = 1, ..., N, the multilinear operator is defined by the  $(M_1 + M_2 + \cdots + M_N)$ th-order tensor

$$\underline{\mathbf{X}} = \left[\!\left[\underline{\mathbf{G}}; \underline{\mathbf{A}}^{(1)}, \dots, \underline{\mathbf{A}}^{(N)}\right]\!\right] \in \mathbb{R}^{I_{1,1} \times \dots \times I_{1,M_1} \times \dots \times I_{N,1} \times \dots \times I_{N,M_N}}$$
(4)

with entries

$$\underline{\mathbf{X}}(\mathbf{i}_1,\mathbf{i}_2,\ldots,\mathbf{i}_N) = \sum_{r_1=1}^{R_1} \sum_{r_2=1}^{R_2} \cdots \sum_{r_N=1}^{R_N} \underline{\mathbf{G}}(r_1,r_2,\ldots,r_N) \underline{\mathbf{A}}^{(1)}(\mathbf{i}_1,r_1) \underline{\mathbf{A}}^{(2)}(\mathbf{i}_2,r_2) \cdots \underline{\mathbf{A}}^{(N)}(\mathbf{i}_N,r_N),$$

where  $\mathbf{i}_n = (i_{n,1}, i_{n,2}, \dots, i_{n,M_n})$  is the ordered indices.

Figure 3(a) illustrates the tensor network diagram for multilinear operator  $[\underline{\mathbf{G}}; \underline{\mathbf{A}}^{(1)}, \dots, \underline{\mathbf{A}}^{(N)}]$  with an Nth-order tensor  $\underline{\mathbf{G}}$  and 4th-order tensors  $\underline{\mathbf{A}}^{(1)}, \dots, \underline{\mathbf{A}}^{(N)}$ .

As a special case, if  $\underline{\mathbf{A}}^{(n)}$  are matrices, i.e.,  $M_n = 1$  for all n, then the multilinear operator (4) is equivalent to the standard Tucker operator (3). Moreover, in the case of vectors  $\mathbf{a}^{(n)} \in \mathbb{R}^{R_n}$ , i.e.,  $M_n = 0$ , we have the scalar

$$\left[\!\left[\underline{\mathbf{G}};\mathbf{a}^{(1)},\mathbf{a}^{(2)},\ldots,\mathbf{a}^{(N)}\right]\!\right] = \underline{\mathbf{G}}\,\overline{\times}_1\,\mathbf{a}^{(1)}\,\overline{\times}_2\,\mathbf{a}^{(2)}\,\overline{\times}_3\,\cdots\,\overline{\times}_N\,\mathbf{a}^{(N)} \in \mathbb{R},$$

where  $\overline{\times}_n$  is the mode-*n* (vector) product [24].

**Example 2.1.** Let  $\underline{\mathbf{G}} = \mathbf{g}^{(1)} \circ \mathbf{g}^{(2)} \circ \mathbf{g}^{(3)} \in \mathbb{R}^{R_1 \times R_2 \times R_3}$  and  $\underline{\mathbf{A}} = \mathbf{a}^{(1)} \circ \mathbf{a}^{(2)} \circ \mathbf{a}^{(3)} \circ \mathbf{a}^{(4)} \circ \mathbf{a}^{(5)} \in \mathbb{R}^{I_1 \times I_2 \times I_3 \times I_4 \times R_2}$  be rank-one tensors. Then,

$$\llbracket \underline{\mathbf{G}}; \mathbf{I}_{R_1}, \underline{\mathbf{A}}, \mathbf{I}_{R_3} \rrbracket = \langle \mathbf{g}^{(2)}, \mathbf{a}^{(5)} \rangle \cdot \mathbf{g}^{(1)} \circ \mathbf{a}^{(1)} \circ \mathbf{a}^{(2)} \circ \mathbf{a}^{(3)} \circ \mathbf{a}^{(4)} \circ \mathbf{g}^{(3)} \in \mathbb{R}^{R_1 \times I_1 \times I_2 \times I_3 \times I_4 \times R_3},$$

where  $\langle \mathbf{v}, \mathbf{w} \rangle = \mathbf{v}^{\top} \mathbf{w}$  is the innerproduct of vectors, see Figure 3(b).

We can derive the following properties.

**Proposition 2.2.** Let  $N \ge 1$  and  $M_n \ge 0$  for n = 1, ..., N. Let  $\underline{\mathbf{G}}_A$  and  $\underline{\mathbf{G}}_B$  be Nth-order tensors and  $\underline{\mathbf{A}}^{(n)}$  and  $\underline{\mathbf{B}}^{(n)}$  be  $(M_n + 1)$ th-order factor tensors whose  $(M_n + 1)$ th modes are auxiliary modes for n = 1, ..., N. Let

$$\underline{\mathbf{A}} = \begin{bmatrix} \underline{\mathbf{G}}_A; \underline{\mathbf{A}}^{(1)}, \dots, \underline{\mathbf{A}}^{(N)} \end{bmatrix} \quad and \quad \underline{\mathbf{B}} = \begin{bmatrix} \underline{\mathbf{G}}_B; \underline{\mathbf{B}}^{(1)}, \dots, \underline{\mathbf{B}}^{(N)} \end{bmatrix}.$$

Then  
(a) 
$$\underline{\mathbf{A}} + \underline{\mathbf{B}} = \begin{bmatrix} \underline{\mathbf{G}}_A \oplus \underline{\mathbf{G}}_B; \underline{\mathbf{A}}^{(1)} \boxplus \underline{\mathbf{B}}^{(1)}, \dots, \underline{\mathbf{A}}^{(N)} \boxplus \underline{\mathbf{B}}^{(N)} \end{bmatrix}$$
 if  $\underline{\mathbf{A}}$  and  $\underline{\mathbf{B}}$  have the same size.  
(b)  $\underline{\mathbf{A}} \oplus \underline{\mathbf{B}} = \begin{bmatrix} \underline{\mathbf{G}}_A \oplus \underline{\mathbf{G}}_B; \underline{\mathbf{A}}^{(1)} \oplus \underline{\mathbf{B}}^{(1)}, \dots, \underline{\mathbf{A}}^{(N)} \oplus \underline{\mathbf{B}}^{(N)} \end{bmatrix}$ .  
(c)  $\underline{\mathbf{A}} \circledast \underline{\mathbf{B}} = \begin{bmatrix} \underline{\mathbf{G}}_A \otimes \underline{\mathbf{G}}_B; \underline{\mathbf{A}}^{(1)} \boxtimes \underline{\mathbf{B}}^{(1)}, \dots, \underline{\mathbf{A}}^{(N)} \boxtimes \underline{\mathbf{B}}^{(N)} \end{bmatrix}$  if  $\underline{\mathbf{A}}$  and  $\underline{\mathbf{B}}$  have the same size.  
(d)  $\underline{\mathbf{A}} \otimes \underline{\mathbf{B}} = \begin{bmatrix} \underline{\mathbf{G}}_A \otimes \underline{\mathbf{G}}_B; \underline{\mathbf{A}}^{(1)} \otimes \underline{\mathbf{B}}^{(1)}, \dots, \underline{\mathbf{A}}^{(N)} \otimes \underline{\mathbf{B}}^{(N)} \end{bmatrix}$ .

*Proof.* (a) to (d) can be derived from the definitions of the corresponding operations and algebraic manipulation.  $\Box$ 

**Example 2.2.** We consider the examples where  $\underline{\mathbf{A}}^{(n)}$  and  $\underline{\mathbf{B}}^{(n)}$  are either factor matrices or vectors.

1. Let  $M_n = 1$  for n = 1, ..., N, i.e., the tensors <u>A</u> and <u>B</u> have the form (which is the Tucker decomposition, to be introduced in Section 3.1)

$$\underline{\mathbf{A}} = \left[\!\left[\underline{\mathbf{G}}_{A}; \mathbf{A}^{(1)}, \dots, \mathbf{A}^{(N)}\right]\!\right] \in \mathbb{R}^{I_{1} \times I_{2} \times \dots \times I_{N}},$$
$$\underline{\mathbf{B}} = \left[\!\left[\underline{\mathbf{G}}_{B}; \mathbf{B}^{(1)}, \dots, \mathbf{B}^{(N)}\right]\!\right] \in \mathbb{R}^{I_{1} \times I_{2} \times \dots \times I_{N}}.$$

It follows that the Kronecker product, Hadamard product, direct sum, and addition lead to tensors in the same form:

- (a)  $\underline{\mathbf{A}} + \underline{\mathbf{B}} = \llbracket \underline{\mathbf{G}}_A \oplus \underline{\mathbf{G}}_B; \mathbf{A}^{(1)} \boxplus \mathbf{B}^{(1)}, \dots, \mathbf{A}^{(N)} \boxplus \mathbf{B}^{(N)} \rrbracket.$
- (b)  $\underline{\mathbf{A}} \oplus \underline{\mathbf{B}} = \llbracket \underline{\mathbf{G}}_A \oplus \underline{\mathbf{G}}_B; \mathbf{A}^{(1)} \oplus \mathbf{B}^{(1)}, \dots, \mathbf{A}^{(N)} \oplus \mathbf{B}^{(N)} \rrbracket.$
- (c)  $\underline{\mathbf{A}} \otimes \underline{\mathbf{B}} = \llbracket \underline{\mathbf{G}}_A \otimes \underline{\mathbf{G}}_B; \mathbf{A}^{(1)} \boxtimes \mathbf{B}^{(1)}, \dots, \mathbf{A}^{(N)} \boxtimes \mathbf{B}^{(N)} \rrbracket.$
- (d)  $\underline{\mathbf{A}} \otimes \underline{\mathbf{B}} = \llbracket \underline{\mathbf{G}}_A \otimes \underline{\mathbf{G}}_B; \mathbf{A}^{(1)} \otimes \mathbf{B}^{(1)}, \dots, \mathbf{A}^{(N)} \otimes \mathbf{B}^{(N)} \rrbracket.$

Moreover, if the core tensors  $\underline{\mathbf{G}}_A$  and  $\underline{\mathbf{G}}_B$  are superdiagonal tensors, which is the case of the CP decomposition (see Section 3.1,) then the results are also given as the CP decomposition because the Kronecker product and the direct sum of superdiagonal core tensors are superdiagonal tensors as well.

2. Let  $M_n = 0$  for n = 1, ..., N, then we have the scalars

$$a = \left[ \left[ \underline{\mathbf{G}}_{A}; \mathbf{a}^{(1)}, \dots, \mathbf{a}^{(N)} \right] \right] \in \mathbb{R},$$
$$b = \left[ \left[ \underline{\mathbf{G}}_{B}; \mathbf{b}^{(1)}, \dots, \mathbf{b}^{(N)} \right] \right] \in \mathbb{R}.$$

The addition and multiplication are given in the form

(a)  $a + b = a \oplus b = \llbracket \underline{\mathbf{G}}_A \oplus \underline{\mathbf{G}}_B; \mathbf{a}^{(1)} \oplus \mathbf{b}^{(1)}, \dots, \mathbf{a}^{(N)} \oplus \mathbf{b}^{(N)} \rrbracket,$ (b)  $ab = a \otimes b = a \circledast b = \llbracket \underline{\mathbf{G}}_A \otimes \underline{\mathbf{G}}_B; \mathbf{a}^{(1)} \otimes \mathbf{b}^{(1)}, \dots, \mathbf{a}^{(N)} \otimes \mathbf{b}^{(N)} \rrbracket.$ 

## 2.6 Contracted product

The mode-n product of a tensor with a matrix in (2) can be extended to a product between tensors of any orders. We define one of the simplest cases of the tensor-by-tensor contracted product as follows.

**Definition 2.7** (Mode-(M, 1) contracted product). Let  $M, N \ge 1$ . The mode-(M, 1) contracted product of tensors  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_M}$  and  $\underline{\mathbf{B}} \in \mathbb{R}^{J_1 \times J_2 \times \cdots \times J_N}$  with  $I_M = J_1$  is defined by

$$\underline{\mathbf{C}} = \underline{\mathbf{A}} \times^1 \underline{\mathbf{B}} \in \mathbb{R}^{I_1 \times \cdots \times I_{M-1} \times J_2 \times \cdots \times J_N}$$

with entries

$$\underline{\mathbf{C}}(i_1,\ldots,i_{M-1},j_2,\ldots,j_N) = \sum_{i_M=1}^{I_M} \underline{\mathbf{A}}(i_1,\ldots,i_M) \underline{\mathbf{B}}(i_M,j_2,\ldots,j_N)$$

for all  $i_m, j_n, m = 1, \dots, M - 1, n = 2, \dots, N$ .

We note that the tensor-by-tensor contracted product defined above is a natural generalization of the matrix multiplication as  $\mathbf{AB} = \mathbf{A} \times^1 \mathbf{B}$ , and the vector innerproduct as  $\langle \mathbf{a}, \mathbf{b} \rangle = \mathbf{a} \times^1 \mathbf{b}$ . Especially, the contracted product between a tensor  $\mathbf{A} \in \mathbb{R}^{I_1 \times \cdots \times I_M}$  and a vector  $\mathbf{p} \in \mathbb{R}^{I_1}$  or  $\mathbf{q} \in \mathbb{R}^{I_M}$  produces a tensor of smaller order as

$$\mathbf{p} \times^{1} \underline{\mathbf{A}} \in \mathbb{R}^{I_{2} \times \cdots \times I_{M}},$$
$$\underline{\mathbf{A}} \times^{1} \mathbf{q} \in \mathbb{R}^{I_{1} \times \cdots \times I_{M-1}}.$$

**Example 2.3.** The contracted product of rank-one tensors yields

$$\left(\mathbf{a}^{(1)}\circ\cdots\circ\mathbf{a}^{(M)}\right)\times^{1}\left(\mathbf{b}^{(1)}\circ\cdots\circ\mathbf{b}^{(N)}\right)=\left\langle\mathbf{a}^{(M)},\mathbf{b}^{(1)}\right\rangle\cdot\mathbf{a}^{(1)}\circ\cdots\circ\mathbf{a}^{(M-1)}\circ\mathbf{b}^{(2)}\circ\cdots\circ\mathbf{b}^{(N)}.$$

In general, we have the following properties:

**Proposition 2.3.** Let  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_M}$ ,  $\underline{\mathbf{B}} \in \mathbb{R}^{I_M \times J_2 \times \cdots \times J_N}$ ,  $\underline{\mathbf{C}} \in \mathbb{R}^{J_N \times K_2 \times \cdots \times K_L}$ ,  $\underline{\mathbf{G}} \in \mathbb{R}^{R_1 \times \cdots \times R_N}$ ,  $\mathbf{P} \in \mathbb{R}^{I \times R_1}$ , and  $\mathbf{Q} \in \mathbb{R}^{R_N \times J}$ . Then,

- (a)  $\underline{\mathbf{A}} \times^1 \underline{\mathbf{B}} = [\![\underline{\mathbf{B}}; \underline{\mathbf{A}}, \mathbf{I}_{J_2}, \dots, \mathbf{I}_{J_N}]\!].$
- (b)  $(\underline{\mathbf{A}} \times^1 \underline{\mathbf{B}}) \times^1 \underline{\mathbf{C}} = \underline{\mathbf{A}} \times^1 (\underline{\mathbf{B}} \times^1 \underline{\mathbf{C}}).$
- (c)  $\mathbf{P} \times^1 \underline{\mathbf{G}} = \underline{\mathbf{G}} \times_1 \mathbf{P}$ .
- (d)  $\underline{\mathbf{G}} \times^1 \mathbf{Q} = \underline{\mathbf{G}} \times_N \mathbf{Q}^\top$ .
- (e)  $\left(\underline{\mathbf{A}} \times^{1} \underline{\mathbf{B}}\right)_{<m>} = \mathbf{A}_{<m>} \left(\mathbf{I}_{I_{m+1}I_{m+2}\cdots I_{M-1}} \otimes \mathbf{B}_{(1)}\right)$  for  $m = 1, 2, \dots, M-1$ .

(f) 
$$(\underline{\mathbf{A}} \times^1 \underline{\mathbf{B}})_{< M+n-2>} = (\mathbf{A}_{< M-1>} \otimes \mathbf{I}_{J_2 J_3 \cdots J_n}) \mathbf{B}_{< n>} \text{ for } n = 2, \dots, N.$$

$$(g) \ vec\left(\underline{\mathbf{A}} \times^{1} \underline{\mathbf{B}}\right) = \left(\mathbf{I}_{I_{1}I_{2}\cdots I_{M-1}} \otimes \mathbf{B}_{(1)}^{\top}\right) vec\left(\underline{\mathbf{A}}\right) = \left(\mathbf{A}_{(M)}^{\top} \otimes \mathbf{I}_{J_{2}J_{3}\cdots J_{N}}\right) vec\left(\underline{\mathbf{B}}\right)$$

*Proof.* (a) to (d) follow immediately from the definitions of the corresponding operations. We can prove (e) and the first equality of (g) as follows. Let  $\mathbf{Y}_{<0>} \equiv \operatorname{vec}(\underline{\mathbf{Y}})^{\top}$  denote the row vector for a

tensor  $\underline{\mathbf{Y}}$ . Note that  $\underline{\mathbf{A}} \times^1 \underline{\mathbf{B}} \in \mathbb{R}^{I_1 \times \cdots \times I_{M-1} \times J_2 \times \cdots \times J_N}$ . For  $0 \le m \le M-1$ , we have

$$\begin{aligned} \left(\underline{\mathbf{A}} \times^{1} \underline{\mathbf{B}}\right)_{} \left(\overline{i_{1} \cdots i_{m}}, \overline{i_{m+1} \cdots i_{M-1} j_{2} \cdots j_{N}}\right) &= \left(\underline{\mathbf{A}} \times^{1} \underline{\mathbf{B}}\right) \left(i_{1}, \dots, i_{M-1}, j_{2}, \dots, j_{N}\right) \\ &= \sum_{i_{M}=1}^{I_{M}} \underline{\mathbf{A}} (i_{1}, \dots, i_{M}) \underline{\mathbf{B}} (i_{M}, j_{2}, \dots, j_{N}) \\ &= \sum_{i_{M}=1}^{I_{M}} \underline{\mathbf{A}} (i_{1}, \dots, i_{M}) \mathbf{B}_{(1)} (i_{M}, \overline{j_{2} \cdots j_{N}}) \\ &= \left(\underline{\mathbf{A}} \times_{M} \mathbf{B}_{(1)}^{\top}\right) (i_{1}, \dots, i_{M-1}, \overline{j_{2} \cdots j_{N}}). \end{aligned}$$

From Proposition 2.1(f), we have

$$\underline{\mathbf{X}} = \underline{\mathbf{A}} \times_M \mathbf{B}_{(1)}^\top \Leftrightarrow \mathbf{X}_{} = \mathbf{A}_{} \left( \mathbf{I}_{I_{m+1}} \otimes \cdots \otimes \mathbf{I}_{I_{M-1}} \otimes \mathbf{B}_{(1)} \right),$$

and since

$$\underline{\mathbf{X}}(i_1,\ldots,i_{M-1},\overline{j_2\cdots j_N}) = \mathbf{X}_{}(\overline{i_1\cdots i_m},\overline{i_{m+1}\cdots i_{M-1}j_2\cdots j_N}),$$

the results in (e) and (g) follow.

We can prove (f) and the second equality of (g) similarly. We have

$$\left(\underline{\mathbf{A}}\times^{1}\underline{\mathbf{B}}\right)_{< M+n-2>} (\overline{i_{1}\cdots i_{M-1}j_{2}\cdots j_{n}}, \overline{j_{n+1}\cdots j_{N}}) = (\underline{\mathbf{B}}\times_{1}\mathbf{A}_{< M-1>}) (\overline{i_{1}\cdots i_{M-1}}, j_{2}, \dots, j_{N}).$$

From Proposition 2.1(f), we have

$$\underline{\mathbf{X}} = \underline{\mathbf{B}} \times_1 \mathbf{A}_{\langle M-1 \rangle} \Leftrightarrow \mathbf{X}_{\langle n \rangle} = (\mathbf{A}_{\langle M-1 \rangle} \otimes \mathbf{I}_{J_2} \otimes \mathbf{I}_{J_3} \otimes \cdots \otimes \mathbf{I}_{J_n}) \mathbf{B}_{\langle n \rangle},$$

and the results in (f) and (g) follow from

$$\underline{\mathbf{X}}(\overline{i_1\cdots i_{M-1}}, j_2, \dots, j_N) = \mathbf{X}_{\langle n \rangle}(\overline{i_1\cdots i_{M-1}j_2\cdots j_n}, \overline{j_{n+1}\cdots j_N}).$$

Moreover, the following property states that several binary operations (addition, direct sum, Hadamard product, and Kronecker product) preserve the form of sequential contracted products of factor (core) tensors. The form will be introduced as the TT decomposition in Section 3.2.

**Proposition 2.4.** Let  $N \ge 2$  and

$$\underline{\mathbf{A}} = \mathbf{A}^{(1)} \times^{1} \underline{\mathbf{A}}^{(2)} \times^{1} \cdots \times^{1} \underline{\mathbf{A}}^{(N-1)} \times^{1} \mathbf{A}^{(N)} \in \mathbb{R}^{I_{1} \times I_{2} \times \cdots \times I_{N}},$$
$$\underline{\mathbf{B}} = \mathbf{B}^{(1)} \times^{1} \underline{\mathbf{B}}^{(2)} \times^{1} \cdots \times^{1} \underline{\mathbf{B}}^{(N-1)} \times^{1} \mathbf{B}^{(N)} \in \mathbb{R}^{J_{1} \times J_{2} \times \cdots \times J_{N}},$$

where  $\underline{\mathbf{A}}^{(n)}$  and  $\underline{\mathbf{B}}^{(n)}$  are factor (core) tensors with sizes

$$\mathbf{A}^{(1)} \in \mathbb{R}^{I_1 \times R_1}, \quad \underline{\mathbf{A}}^{(n)} \in \mathbb{R}^{R_{n-1} \times I_n \times R_n}, \ n = 2, \dots, N-1, \quad \mathbf{A}^{(N)} \in \mathbb{R}^{R_{N-1} \times I_N}, \\ \mathbf{B}^{(1)} \in \mathbb{R}^{J_1 \times S_1}, \quad \underline{\mathbf{B}}^{(n)} \in \mathbb{R}^{S_{n-1} \times J_n \times S_n}, \ n = 2, \dots, N-1, \quad \mathbf{B}^{(N)} \in \mathbb{R}^{S_{N-1} \times J_N}.$$

Then,

(a) 
$$\underline{\mathbf{A}} + \underline{\mathbf{B}} = (\mathbf{A}^{(1)} \boxplus \mathbf{B}^{(1)}) \times^1 (\underline{\mathbf{A}}^{(2)} \boxplus \underline{\mathbf{B}}^{(2)}) \times^1 \cdots \times^1 (\mathbf{A}^{(N)} \boxplus \mathbf{B}^{(N)})$$
 if  $\underline{\mathbf{A}}$  and  $\underline{\mathbf{B}}$  have the same size.

$$(b) \ \underline{\mathbf{A}} \oplus \underline{\mathbf{B}} = \left(\mathbf{A}^{(1)} \oplus \mathbf{B}^{(1)}\right) \times^{1} \left(\underline{\mathbf{A}}^{(2)} \oplus \underline{\mathbf{B}}^{(2)}\right) \times^{1} \cdots \times^{1} \left(\mathbf{A}^{(N)} \oplus \mathbf{B}^{(N)}\right).$$

$$(c) \ \underline{\mathbf{A}} \otimes \underline{\mathbf{B}} = \left(\mathbf{A}^{(1)} \boxtimes \mathbf{B}^{(1)}\right) \times^{1} \left(\underline{\mathbf{A}}^{(2)} \boxtimes \underline{\mathbf{B}}^{(2)}\right) \times^{1} \cdots \times^{1} \left(\mathbf{A}^{(N)} \boxtimes \mathbf{B}^{(N)}\right) \text{ if } \underline{\mathbf{A}} \text{ and } \underline{\mathbf{B}} \text{ have the same}$$

$$(d) \ \underline{\mathbf{A}} \otimes \underline{\mathbf{B}} = \left(\mathbf{A}^{(1)} \otimes \mathbf{B}^{(1)}\right) \times^{1} \left(\underline{\mathbf{A}}^{(2)} \otimes \underline{\mathbf{B}}^{(2)}\right) \times^{1} \cdots \times^{1} \left(\mathbf{A}^{(N)} \otimes \mathbf{B}^{(N)}\right).$$

*Proof.* (a) to (d) can be derived by algebraic manipulation and the definitions of the corresponding operations.  $\Box$ 

The contracted product of tensors defined above can be further generalized to a contracted product of block tensors as follows. In the following definition, the tensors are given in partitioned form and the contracted product is performed between each pair of blocks.

**Definition 2.8** (Mode-(M, 1) contracted product for block tensors). Let tensors  $\underline{\widetilde{\mathbf{A}}} = [\underline{\mathbf{A}}_{r_1, r_2}]$ and  $\underline{\widetilde{\mathbf{B}}} = [\underline{\mathbf{B}}_{s_1, s_2}]$  be block tensors partitioned with *M*th-order tensors  $\underline{\mathbf{A}}_{r_1, r_2} \in \mathbb{R}^{I_1 \times \cdots \times I_M}$ ,  $r_1 = 1, \ldots, R_1, r_2 = 1, \ldots, R_2$ , and *N*th-order tensors  $\underline{\mathbf{B}}_{s_1, s_2} \in \mathbb{R}^{J_1 \times \cdots \times J_N}$  with  $I_M = J_1, s_1 = 1, \ldots, S_1$ ,  $s_2 = 1, \ldots, S_2$ , respectively, i.e.,

$$\widetilde{\underline{\mathbf{A}}} = \begin{bmatrix} \underline{\mathbf{A}}_{1,1} & \cdots & \underline{\mathbf{A}}_{1,R_2} \\ \vdots & \ddots & \vdots \\ \underline{\mathbf{A}}_{R_1,1} & \cdots & \underline{\mathbf{A}}_{R_1,R_2} \end{bmatrix}, \qquad \widetilde{\underline{\mathbf{B}}} = \begin{bmatrix} \underline{\mathbf{B}}_{1,1} & \cdots & \underline{\mathbf{B}}_{1,S_2} \\ \vdots & \ddots & \vdots \\ \underline{\mathbf{B}}_{S_1,1} & \cdots & \underline{\mathbf{B}}_{S_1,S_2} \end{bmatrix}$$

The mode-(M, 1) contracted product of  $\underline{\widetilde{A}}$  and  $\underline{\widetilde{B}}$  is defined by the block tensor

$$\underline{\widetilde{\mathbf{C}}} = \left[\underline{\mathbf{C}}_{t_1, t_2}\right] = \underline{\widetilde{\mathbf{A}}} \times^1 \underline{\widetilde{\mathbf{B}}}$$

partitioned with the (M + N - 1)th-order tensors

$$\underline{\mathbf{C}}_{t_1,t_2} = \underline{\mathbf{A}}_{r_1,r_2} \times^1 \underline{\mathbf{B}}_{s_1,s_2} \in \mathbb{R}^{I_1 \times \dots \times I_{M-1} \times J_2 \times \dots \times J_N}, \quad t_1 = \overline{r_1 s_1}, \ t_2 = \overline{r_2 s_2}$$

for all  $t_1 = 1, \ldots, R_1 S_1, t_2 = 1, \ldots, R_2 S_2$ .

**Example 2.4.** For example, the contracted product of block matrices  $\widetilde{\mathbf{A}}$  and  $\widetilde{\mathbf{B}}$  with 2 row partitions and 2 column partitions yields a block matrix with 4 row partitions and 4 column partitions as

$$\begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{bmatrix} \times^{1} \begin{bmatrix} \mathbf{B}_{11} & \mathbf{B}_{12} \\ \mathbf{B}_{21} & \mathbf{B}_{22} \end{bmatrix} = \begin{bmatrix} \mathbf{A}_{11}\mathbf{B}_{11} & \mathbf{A}_{11}\mathbf{B}_{12} & \mathbf{A}_{12}\mathbf{B}_{11} & \mathbf{A}_{12}\mathbf{B}_{12} \\ \mathbf{A}_{11}\mathbf{B}_{21} & \mathbf{A}_{11}\mathbf{B}_{22} & \mathbf{A}_{12}\mathbf{B}_{21} & \mathbf{A}_{12}\mathbf{B}_{22} \\ \mathbf{A}_{21}\mathbf{B}_{11} & \mathbf{A}_{21}\mathbf{B}_{12} & \mathbf{A}_{22}\mathbf{B}_{11} & \mathbf{A}_{22}\mathbf{B}_{12} \\ \mathbf{A}_{21}\mathbf{B}_{21} & \mathbf{A}_{21}\mathbf{B}_{22} & \mathbf{A}_{22}\mathbf{B}_{21} & \mathbf{A}_{22}\mathbf{B}_{22} \\ \end{bmatrix}$$

Note that the above definition of contracted product for block matrices/block tensors is different from the standard matrix-by-matrix product. That is, in the same example, the standard matrix-by-matrix product produces a block matrix with 2 row partitions and 2 column partitions with blocks  $\mathbf{C}_{t_1,t_2} = \sum_{r=1}^{2} \mathbf{A}_{t_1,r} \mathbf{B}_{r,t_2}$ .

## 2.7 Strong Kronecker product

The strong Kronecker product is an important tool for representation of low-rank TT decompositions of large-scale vectors, matrices, and low-order tensors. The original definition of the strong Kronecker product for block matrices [29] is presented below, together with its generalization to block tensors.

**Definition 2.9** (Strong Kronecker product, [29]). Let matrices  $\mathbf{A} = [\mathbf{A}_{r_1,r_2}] \in \mathbb{R}^{R_1 I_1 \times R_2 J_1}$  and  $\mathbf{B} = [\mathbf{B}_{r_2,r_3}] \in \mathbb{R}^{R_2 I_2 \times R_3 J_2}$  be block matrices partitioned with  $\mathbf{A}_{r_1,r_2} \in \mathbb{R}^{I_1 \times J_1}$  and  $\mathbf{B}_{r_2,r_3} \in \mathbb{R}^{I_2 \times J_2}$ , respectively. The strong Kronecker product of  $\mathbf{A}$  and  $\mathbf{B}$  is defined by the block matrix

$$\mathbf{C} = \begin{bmatrix} \mathbf{C}_{r_1, r_3} \end{bmatrix} = \mathbf{A} \mid \otimes \mid \mathbf{B} \in \mathbb{R}^{R_1 I_1 I_2 \times R_3 J_1 J_2}$$

partitioned with the  $I_1I_2 \times J_1J_2$  matrices

$$\mathbf{C}_{r_1,r_3} = \sum_{r_2=1}^{R_2} \mathbf{A}_{r_1,r_2} \otimes \mathbf{B}_{r_2,r_3} \in \mathbb{R}^{I_1 I_2 \times J_1 J_2},$$

for  $r_1 = 1, \ldots, R_1, r_3 = 1, \ldots, R_3$ .

More generally, let tensors  $\underline{\mathbf{A}} = [\underline{\mathbf{A}}_{r_1, r_2}] \in \mathbb{R}^{R_1 I_1 \times R_2 J_1 \times K_1}$  and  $\underline{\mathbf{B}} = [\underline{\mathbf{B}}_{r_2, r_3}] \in \mathbb{R}^{R_2 I_2 \times R_3 J_2 \times K_2}$ be block tensors partitioned with 3rd-order tensors  $\underline{\mathbf{A}}_{r_1, r_2} \in \mathbb{R}^{I_1 \times J_1 \times K_1}$  and  $\underline{\mathbf{B}}_{r_2, r_3} \in \mathbb{R}^{I_2 \times J_2 \times K_2}$ . The strong Kronecker product of  $\underline{\mathbf{A}}$  and  $\underline{\mathbf{B}}$  is defined by the block tensor

$$\underline{\mathbf{C}} = \left[\underline{\mathbf{C}}_{r_1, r_3}\right] = \underline{\mathbf{A}} |\otimes| \underline{\mathbf{B}} \in \mathbb{R}^{R_1 I_1 I_2 \times R_3 J_1 J_2 \times K_1 K_2},$$

partitioned with the  $I_1I_2 \times J_1J_2 \times K_1K_2$  tensors

$$\underline{\mathbf{C}}_{r_1,r_3} = \sum_{r_2=1}^{R_2} \underline{\mathbf{A}}_{r_1,r_2} \otimes \underline{\mathbf{B}}_{r_2,r_3} \in \mathbb{R}^{I_1 I_2 \times J_1 J_2 \times K_1 K_2},$$

for  $r_1 = 1, \ldots, R_1, r_3 = 1, \ldots, R_3$ .

**Example 2.5.** The strong Kronecker product has a similarity with the matrix-by-matrix multiplication. For example,

$$\begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{bmatrix} | \otimes | \begin{bmatrix} \mathbf{B}_{11} & \mathbf{B}_{12} \\ \mathbf{B}_{21} & \mathbf{B}_{22} \end{bmatrix} = \begin{bmatrix} \mathbf{A}_{11} \otimes \mathbf{B}_{11} + \mathbf{A}_{12} \otimes \mathbf{B}_{21} & \mathbf{A}_{11} \otimes \mathbf{B}_{12} + \mathbf{A}_{12} \otimes \mathbf{B}_{22} \\ \mathbf{A}_{21} \otimes \mathbf{B}_{11} + \mathbf{A}_{22} \otimes \mathbf{B}_{21} & \mathbf{A}_{21} \otimes \mathbf{B}_{12} + \mathbf{A}_{22} \otimes \mathbf{B}_{22} \end{bmatrix}.$$

## 2.8 Partial trace operator

We will define a linear operator, Tr, called as the partial trace, which generalizes the trace on matrices to tensors.

**Definition 2.10** (Partial trace operator). The partial trace,  $Tr : \mathbb{R}^{R \times I_1 \times I_2 \times \cdots \times I_N \times R} \to \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$ ,  $N \ge 2$ , is a linear operator defined by the tensor  $\underline{\mathbf{Y}} = Tr(\underline{\mathbf{X}})$  with entries

$$\underline{\mathbf{Y}}(i_1, i_2, \dots, i_N) = \sum_{r=1}^R \underline{\mathbf{X}}(r, i_1, i_2, \dots, i_N, r).$$
(5)

A more formal definition can be given by using the contracted product as

$$Tr: \underline{\mathbf{X}} \mapsto Tr(\underline{\mathbf{X}}) = \sum_{r=1}^{R} \mathbf{e}_r \times^1 \underline{\mathbf{X}} \times^1 \mathbf{e}_r, \tag{6}$$

where  $\mathbf{e}_r = [0, \dots, 0, 1, 0, \dots, 0]^{\top} \in \mathbb{R}^R$  is the *r*th standard basis vector. The partial trace is a generalization of the matrix trace: for a matrix  $\mathbf{A} \in \mathbb{R}^{R \times R}$ ,  $Tr(\mathbf{A}) = \operatorname{trace}(\mathbf{A}) = \sum_{r=1}^{R} a_{rr} \in \mathbb{R}$ . For a tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{R \times I_1 \times I_2 \times \dots \times I_N \times R}$ , the  $(i_1, i_2, \dots, i_N)$ th entry of  $Tr(\underline{\mathbf{X}})$  equals to the matrix trace of the  $(i_1, i_2, \dots, i_N)$ th slice as

$$(Tr(\underline{\mathbf{X}}))_{i_1,i_2,\ldots,i_N} = \operatorname{trace}\left(\mathbf{X}_{:,i_1,i_2,\ldots,i_N,:}\right).$$
(7)

Figure 4(a) illustrates a tensor network diagram representing the partial trace of a 7th-order tensor.



Figure 4: Tensor network diagrams for (a) partial trace,  $Tr(\underline{\mathbf{X}})$ , of a tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{R \times I_1 \times \cdots \times I_5 \times R}$ , and (b) partial trace of a rank-one tensor of order N = 7 as described in Example 2.6.

**Example 2.6.** The partial trace of a rank-one tensor can be expressed by, for  $N \ge 1$ ,

$$Tr\left(\mathbf{a}^{(1)} \circ \mathbf{a}^{(2)} \circ \cdots \circ \mathbf{a}^{(N)}\right) = \langle \mathbf{a}^{(1)}, \mathbf{a}^{(N)} \rangle \cdot \mathbf{a}^{(2)} \circ \cdots \circ \mathbf{a}^{(N-1)},$$

where  $\langle \mathbf{v}, \mathbf{w} \rangle = \mathbf{v}^{\top} \mathbf{w}$ , see, e.g., Figure 4(b).

**Example 2.7.** If  $\underline{\mathbf{A}} \in \mathbb{R}^{R_1 \times I \times R_2}$ ,  $\underline{\mathbf{B}} \in \mathbb{R}^{R_2 \times J \times R_3}$ , and  $\underline{\mathbf{C}} \in \mathbb{R}^{R_3 \times K \times R_1}$ , then,

$$\left(Tr(\underline{\mathbf{A}}\times^{1}\underline{\mathbf{B}}\times^{1}\underline{\mathbf{C}})\right)_{ijk} = \operatorname{trace}(\mathbf{A}_{:,i,:}\mathbf{B}_{:,j,:}\mathbf{C}_{:,k,:}) = \left(Tr(\underline{\mathbf{C}}\times^{1}\underline{\mathbf{B}}\times^{1}\underline{\mathbf{A}})\right)_{kji}$$

## 3 Tensor decompositions

Tensor decomposition is an approximate representation of a tensor as contraction of a set of factor (core) tensors. See, e.g., [11, 16] for more general definitions of tensor format and tensor representation in tensor product space. We will introduce several tensor decompositions (CP, Tucker, TT) in a consistent way, based on the notations and tensor operations defined in the previous section.

## 3.1 CP and Tucker decompositions

The CANDECOMP/PARAFAC (CP) decomposition expresses a tensor as a sum of rank-one tensors: a tensor  $\underline{\mathbf{X}}$  of size  $I_1 \times I_2 \times \cdots \times I_N$  is written as

$$\underline{\mathbf{X}} = \sum_{r=1}^{R} \lambda_r \mathbf{a}_r^{(1)} \circ \mathbf{a}_r^{(2)} \circ \dots \circ \mathbf{a}_r^{(N)}, \tag{8}$$

where  $\mathbf{a}_r^{(n)} \in \mathbb{R}^{I_n}, r = 1, \dots, R, n = 1, \dots, N$ , are normalized vectors,  $\lambda_r, r = 1, \dots, R$ , are weights, and  $R \in \mathbb{N}$  is called the rank of the CP decomposition (8). The above expression can be equivalently re-written in form of multilinear products as

$$\underline{\mathbf{X}} = \underline{\mathbf{\Lambda}} \times_1 \mathbf{A}^{(1)} \times_2 \mathbf{A}^{(2)} \dots \times_N \mathbf{A}^{(N)} = \left[\!\!\left[\underline{\mathbf{\Lambda}}; \mathbf{A}^{(1)}, \mathbf{A}^{(2)}, \dots, \mathbf{A}^{(N)}\right]\!\!\right],\tag{9}$$

where  $\underline{\mathbf{\Lambda}} \in \mathbb{R}^{R \times \cdots \times R}$  is the superdiagonal tensor with diagonals  $\lambda_1, \ldots, \lambda_R$ , and  $\mathbf{A}^{(n)} = [\mathbf{a}_1^{(n)}, \ldots, \mathbf{a}_R^{(n)}]$ are  $I_n \times R_n$  factor matrices. Other alternative representations are summarized in Table 3. The Khatri-Rao (columnwise Kronecker) product of two matrices  $\mathbf{A} \in \mathbb{R}^{I \times R}$  and  $\mathbf{B} \in \mathbb{R}^{J \times R}$  is denoted by  $\mathbf{C} = \mathbf{A} \odot \mathbf{B} \in \mathbb{R}^{IJ \times R}$  with columns  $\mathbf{c}_r = \mathbf{a}_r \otimes \mathbf{b}_r$ ,  $r = 1, \ldots, R$ .

CP	Tucker
Multil	inear product
$\underline{\mathbf{X}} = \underline{\mathbf{\Lambda}} \times_1 \mathbf{A}^{(1)} \cdots \times_N \mathbf{A}^{(N)}$	$\underline{\mathbf{X}} = \underline{\mathbf{G}} \times_1 \mathbf{A}^{(1)} \cdots \times_N \mathbf{A}^{(N)}$
$\underline{\mathbf{X}} = \left[\!\!\left[\underline{\mathbf{\Lambda}}; \mathbf{A}^{(1)}, \dots, \mathbf{A}^{(N)}\right]\!\!\right]$	$\underline{\mathbf{X}} = \left[\!\!\left[\underline{\mathbf{G}}; \mathbf{A}^{(1)}, \dots, \mathbf{A}^{(N)} ight]\!\!\right]$
Out	ter product
$\underline{\mathbf{X}} = \sum_{r=1}^{R} \lambda_r \mathbf{a}_r^{(1)} \circ \dots \circ \mathbf{a}_r^{(N)}$	$\underline{\mathbf{X}} = \sum_{r_1=1}^{R_1} \cdots \sum_{r_N=1}^{R_N} g_{r_1,\dots,r_N} \mathbf{a}_{r_1}^{(1)} \circ \dots \circ \mathbf{a}_{r_N}^{(N)}$
Sca	lar product
$x_{i_1,\dots,i_N} = \sum_{r=1}^R \lambda_r a_{i_1,r}^{(1)} \cdots a_{i_N,r}^{(N)}$	$x_{i_1,\dots,i_N} = \sum_{r_1=1}^{R_1} \cdots \sum_{r_N=1}^{R_N} g_{r_1,\dots,r_N} a_{i_1,r_1}^{(1)} \cdots a_{i_N,r_N}^{(N)}$
Slice re	epresentation*
$\mathbf{X}_{:,:,i_3,,i_N} = \mathbf{A}^{(1)} \widetilde{\mathbf{D}}_{i_3,,i_N} \mathbf{A}^{(2)\mathrm{T}}$	$\mathbf{X}_{:,:,i_3,,i_N} = \mathbf{A}^{(1)} \widetilde{\mathbf{G}}_{i_3,,i_N} \mathbf{A}^{(2)\mathrm{T}}$
Vect	torization**
$\operatorname{vec}(\underline{\mathbf{X}}) = \left( \bigotimes_{n=1}^{N} \mathbf{A}^{(n)} \right) \boldsymbol{\lambda}$	$\operatorname{vec}(\underline{\mathbf{X}}) = \left(\bigotimes_{n=1}^{N} \mathbf{A}^{(n)}\right) \operatorname{vec}(\underline{\mathbf{G}})$
Mat	ricization**
$\mathbf{X}_{(n)} = \mathbf{A}^{(n)} \mathbf{\Lambda} \left( \bigotimes_{\substack{m=1 \\ m \neq n}}^{N} \mathbf{A}^{(m)} \right)^{\mathrm{T}}$	$\mathbf{X}_{(n)} = \mathbf{A}^{(n)} \mathbf{G}_{(n)} \left( \bigotimes_{\substack{m=1\\m \neq n}}^{N} \mathbf{A}^{(m)} \right)^{\mathrm{T}}$
$\mathbf{X}_{\langle n \rangle} = \left( \bigodot_{m=1}^{n} \mathbf{A}^{(m)} \right) \mathbf{\Lambda} \left( \bigodot_{m=n+1}^{N} \mathbf{A}^{(m)} \right)^{\mathrm{T}}$	$\mathbf{X}_{\langle n \rangle} = \left(\bigotimes_{m=1}^{n} \mathbf{A}^{(m)}\right) \mathbf{G}_{\langle n \rangle} \left(\bigotimes_{m=n+1}^{N} \mathbf{A}^{(m)}\right)^{\mathrm{T}}$
* $\widetilde{\mathbf{D}}_{i_2,\ldots,i_N} = \operatorname{diag}(\widetilde{d}_{11},\ldots,\widetilde{d}_{RR}) \in \mathbb{R}^{R \times R}$ w	with diagonals $\tilde{d}_{rr} = \lambda_r a^{(3)}(i_3, r) \cdots a^{(N)}(i_N, r).$
* $\widetilde{\mathbf{G}}_{i}$ $= \sum \cdots \sum a_{i}^{(3)} \cdots a_{i}^{(N)}$	$\mathbf{F}_{\text{transmitter}}$ is the sum of frontal slices.
** $\mathbf{A} \odot \mathbf{B}$ stands for the Khatri-Rao (colu	$m_{\text{number}}$ model of the state of the s

Table 3: Various representations for the CP and Tucker decompositions of a tensor  $\underline{\mathbf{X}}$  of size  $I_1 \times I_2 \times \cdots \times I_N$ .



Figure 5: Tensor network diagrams for the (a) CP and (b) Tucker decompositions of an Nth-order tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$ . It is shown that the CP has a superdiagonal core tensor (denoted by a white circle with slash).

The Tucker decomposition [38] decomposes a tensor into a core tensor multiplied by a factor matrix on each mode as

$$\underline{\mathbf{X}} = \underline{\mathbf{G}} \times_1 \mathbf{A}^{(1)} \times_2 \mathbf{A}^{(2)} \cdots \times_N \mathbf{A}^{(N)} = \left[\!\!\left[\underline{\mathbf{G}}; \mathbf{A}^{(1)}, \mathbf{A}^{(2)}, \dots, \mathbf{A}^{(N)}\right]\!\!\right],$$
(10)

where  $\underline{\mathbf{G}} \in \mathbb{R}^{R_1 \times R_2 \times \cdots \times R_N}$  is a core tensor,  $\mathbf{A}^{(n)} \in \mathbb{R}^{I_n \times R_n}$  are factor matrices, and  $(R_1, \ldots, R_N)$  is called the (multilinear) rank of the Tucker decomposition (10). The Tucker decomposition can also be represented as a sum of rank-one tensors as

$$\underline{\mathbf{X}} = \sum_{r_1=1}^{R_1} \sum_{r_2=1}^{R_2} \cdots \sum_{r_N=1}^{R_N} g_{r_1, r_2, \dots, r_N} \mathbf{a}_{r_1}^{(1)} \circ \mathbf{a}_{r_2}^{(2)} \circ \cdots \circ \mathbf{a}_{r_N}^{(N)},$$
(11)

where  $\mathbf{a}_{r_n}^{(n)} \in \mathbb{R}^{I_n}$  is the  $r_n$ th column of  $\mathbf{A}^{(n)}$ . Alternative representations for the Tucker decomposition are summarized in Table 3.

In general, the CP can be regarded as a special case of the Tucker in the sense that the CP is a Tucker decomposition with a superdiagonal core tensor, e.g., see (9) and (10), where the multilinear rank is  $(R, R, \ldots, R)$ .

The CP and Tucker decompositions can be illustrated by tensor network diagrams as in Figure 5. Although the CP decomposition was illustrated by a network diagram in the figure, it is often not classified as a *tensor network format* when  $N \geq 3$  and  $R \geq 2$  in a strict sense [11], due to the superdiagonality of the core tensor  $\underline{\Lambda}$ .

On the other hand, the higher-order singular value decomposition (HOSVD) [7] of a tensor produces a Tucker decomposition with orthogonalized factor matrices and a core tensor. All-orthogonality of the core tensor of Tucker decomposition is defined as follows:

**Definition 3.1** (All-orthogonality, [7]). An *N*th-order tensor  $\underline{\mathbf{G}} \in \mathbb{R}^{R_1 \times \cdots \times R_N}$  is called all-orthogonal if

$$\mathbf{G}_{(n)}\mathbf{G}_{(n)}^{ op} = \mathbf{\Lambda}^{(n)}$$

for some diagonal matrices  $\mathbf{\Lambda}^{(n)} = \operatorname{diag}(\lambda_1^{(n)}, \dots, \lambda_{R_n}^{(n)}) \in \mathbb{R}^{R_n \times R_n}$  with diagonals  $\lambda_1^{(n)} \ge \dots \ge \lambda_{R_n}^{(n)} \ge 0$  for all  $n = 1, \dots, N$ .



Figure 6: Tensor network diagram for TT decomposition of a 7th-order tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_7}$ .

## 3.2 Tensor train (TT) decomposition

By the tensor train (TT) decomposition, a tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  is represented as contracted products

$$\underline{\mathbf{X}} = \underline{\mathbf{G}}^{(1)} \times^1 \underline{\mathbf{G}}^{(2)} \times^1 \cdots \times^1 \underline{\mathbf{G}}^{(N)}, \tag{12}$$

where  $\underline{\mathbf{G}}^{(n)}$  are 3rd-order core (factor) tensors called the TT-cores with sizes  $R_{n-1} \times I_n \times R_n$ ,  $n = 1, \ldots, N$ , the integers  $R_1, \ldots, R_{N-1}$  are called the TT-ranks, and we assume that  $R_0 = R_N = 1$ . For notational convenience, we consider the 1st and the Nth TT-cores as 3rd-order tensors unless stated otherwise. Figure 6 illustrates the tensor network diagram for the TT decomposition of a 7th-order tensor.

The TT decomposition can alternatively be written entrywise as products of slice matrices

$$x_{i_1,i_2,\dots,i_N} = \mathbf{G}_{i_1}^{(1)} \mathbf{G}_{i_2}^{(2)} \cdots \mathbf{G}_{i_N}^{(N)},$$
(13)

where  $\mathbf{G}_{i_n}^{(n)} = \underline{\mathbf{G}}^{(n)}(:, i_n, :) \in \mathbb{R}^{R_{n-1} \times R_n}$  is the lateral slice of the *n*th TT-core,  $n = 1, \ldots, N$ . In this case, we suppose that  $\mathbf{G}_{i_1}^{(1)} \in \mathbb{R}^{1 \times R_1}$  and  $\mathbf{G}_{i_N}^{(N)} \in \mathbb{R}^{R_{N-1} \times 1}$  are row and column vectors. The TT decomposition can also be represented by a sum of outer products

$$\underline{\mathbf{X}} = \sum_{r_1=1}^{R_1} \sum_{r_2=1}^{R_2} \cdots \sum_{r_{N-1}=1}^{R_{N-1}} \mathbf{g}_{1,r_1}^{(1)} \circ \mathbf{g}_{r_1,r_2}^{(2)} \circ \cdots \circ \mathbf{g}_{r_{N-2},r_{N-1}}^{(N-1)} \circ \mathbf{g}_{r_{N-1},1}^{(N)},$$
(14)

where  $\mathbf{g}_{r_{n-1},r_n}^{(n)} = \underline{\mathbf{G}}^{(n)}(r_{n-1},:,r_n) \in \mathbb{R}^{I_n}$  are the mode-2 fibers.

Table 4 summarizes the various representations for the TT decomposition.

#### **3.2.1** Recursive representations and vectorizations for TT decomposition

The TT decomposition can be expressed in a recursive manner, which is summarized in Table 4. Given the TT-cores  $\underline{\mathbf{G}}^{(n)}, n = 1, \dots, N$ , from a TT decomposition (12), we define the partial contracted products  $\underline{\mathbf{G}}^{\leq n}$  and  $\underline{\mathbf{G}}^{\geq n}$  as

$$\underline{\mathbf{G}}^{\leq n} = \underline{\mathbf{G}}^{(1)} \times^{1} \underline{\mathbf{G}}^{(2)} \times^{1} \cdots \times^{1} \underline{\mathbf{G}}^{(n)} \in \mathbb{R}^{I_{1} \times \cdots \times I_{n} \times R_{n}},$$

$$\underline{\mathbf{G}}^{\geq n} = \underline{\mathbf{G}}^{(n)} \times^{1} \underline{\mathbf{G}}^{(n+1)} \times^{1} \cdots \times^{1} \underline{\mathbf{G}}^{(N)} \in \mathbb{R}^{R_{n} \times I_{n+1} \times \cdots \times I_{N}},$$
(15)

for n = 1, ..., N.  $\underline{\mathbf{G}}^{< n}$  and  $\underline{\mathbf{G}}^{> n}$  are defined in the same way. For completeness, we define  $\underline{\mathbf{G}}^{< 1} = \underline{\mathbf{G}}^{> N} = 1$ . Vectorization of the partial contracted products yield the following recursive equations

Table 4: Various representations for the TT decomposition of a tensor  $\underline{\mathbf{X}}$  of size  $I_1 \times \cdots \times I_N$ , either in global representation (left column) or in recursive representation (right column).  $\underline{\mathbf{X}} = \underline{\mathbf{G}}^{\leq N} = \underline{\mathbf{G}}^{\geq 1}$ .

(Iedola) TT	TT (recursive)
Contracted pro	oduct
$\mathbf{X} - \mathbf{G}^{(1)} \times^1 \mathbf{G}^{(2)} \times^1 \cdots \times^1 \mathbf{G}^{(N)}$	$\mathbf{G}^{\leq n} - \mathbf{G}^{\leq n-1} \times^1 \mathbf{G}^{(n)}$
$\underline{\mathbf{x}} = \underline{\mathbf{u}} \times \underline{\mathbf{u}} \times \underline{\mathbf{u}} \times \underline{\mathbf{u}}$	$\underline{\mathbf{G}}^{\geq n} = \underline{\mathbf{G}}^{(n)} \wedge \underline{\mathbf{G}}^{\geq n+1}$
	$\underline{\mathbf{G}}^{\pm n} = \underline{\mathbf{G}}^{(n)} \times^{\mathbf{T}} \underline{\mathbf{G}}^{\pm n+\mathbf{T}}$
Outer produ	lct
$\mathbf{v} = \sum_{n_1,\dots,n_{N-1}} \mathbf{r}^{(1)} \circ \mathbf{r}^{(2)} \circ \cdots \circ \mathbf{r}^{(N)}$	$\mathbf{C}^{\leq n} - \sum_{n=1}^{n-1} \mathbf{C}^{\leq n-1} \circ \mathbf{r}^{(n)}$
$\underline{\mathbf{A}} = \sum_{\mathbf{m}_{1}, \mathbf{r}_{1}} \mathbf{g}_{1, \mathbf{r}_{1}} \circ \mathbf{g}_{\mathbf{r}_{1}, \mathbf{r}_{2}} \circ \cdots \circ \mathbf{g}_{\mathbf{r}_{N-1}, 1}$	$\underline{\mathbf{G}}_{\overline{r}_{n}} = \sum_{r_{n-1}=1} \underline{\mathbf{G}}_{\overline{r}_{n-1}} \circ \mathbf{g}_{\overline{r}_{n-1},r_{n}}$
/1,,/N-1-1	$R_n$
	$\underline{\mathbf{G}}_{r_n-1}^{\geq n} = \sum \mathbf{g}_{r_n-1,r_n}^{(n)} \circ \underline{\mathbf{G}}_{r_n}^{\geq n+1}$
	$r_n=1$
Scalar produ	ıct
$x_{i_1,i_2,\ldots,i_N} =$	$g_{1,i_1,\ldots,i_n,r_n}^{\leq n} =$
$R_1, \dots, R_{N-1}$ (1) (2) (N)	$\sum_{n=1}^{n-1} (n)$
$\sum \qquad g_{1,i_1,r_1}^{(1)} g_{r_1,i_2,r_2}^{(2)} \cdots g_{r_{N-1},i_N,1}^{(N)}$	$\sum g_{1,i_1,\dots,i_{n-1},r_{n-1}}^{\leq n-1} g_{r_{n-1},i_n,r_n}^{(n)}$
$r_1,, r_{N-1} = 1$	$r_{n-1}=1$
	$g_{r_{n-1},i_n,,i_N,1}^- =$
	$\sum_{n=1}^{n} a^{(n)}$ $a^{\geq n+1}$
	$\sum_{r=1}^{J} g_{r_{n-1},i_n,r_n} g_{r_n,i_{n+1},\dots,i_N,1}$
Matrix produ	uct
$x_{i_1,i_2,\dots,i_N} = \mathbf{G}_i^{(1)} \mathbf{G}_i^{(2)} \cdots \mathbf{G}_i^{(N)}$	$\mathbf{G}_{i}^{\leq n}$ , $=\mathbf{G}_{i}^{\leq n-1}$ $\mathbf{G}_{i}^{(n)}$
$i_1, i_2, \dots, i_N = i_1 = i_2 = i_N$	$\mathbf{C}^{\geq n} = \mathbf{C}^{(n)} \mathbf{C}^{\geq n+1}$
<b>T</b> T , <b>1</b> , <b>1</b>	$\mathbf{G}_{i_n,\dots,i_N} - \mathbf{G}_{i_n} \ \mathbf{G}_{i_{n+1},\dots,i_N}$
Vectorizatio	Dn
$\operatorname{vec}(\mathbf{X}) = \prod_{i=1}^{n} \left( \mathbf{I}_{i,i} \otimes \mathbf{G}_{i,i}^{(n)\top} \right)$	$\operatorname{vec}(\mathbf{G}^{\leq n}) =$
$\prod_{n=N} \left( \prod_{i=1}^{n-1} \bigcirc \bigcirc (1) \right)$	
	$\left(\mathbf{I}_{I_{1},I_{2},,I_{n-1}} \otimes \mathbf{G}_{(1)}^{(n)\top}\right) \operatorname{vec}\left(\mathbf{G}^{\leq n-1}\right)$
N	$\left(\begin{array}{c} I_{1}I_{2}\cdots I_{n-1} \\ \end{array}\right)$
$\operatorname{vec}(\mathbf{X}) = \prod_{i=1}^{N} \left( \mathbf{G}_{i}^{(n)\top} \otimes \mathbf{I}_{i} - \mathbf{I}_{i} \right)$	$\operatorname{vec}\left(\mathbf{G}^{\geq n}\right) =$
$\prod_{n=1} \left( \mathbf{C}_{(3)} \otimes \mathbf{I}_{n+1} \cdots \mathbf{I}_{N} \right)$	
	$\left(\mathbf{G}_{(n)}^{(n)\top}\otimes\mathbf{I}_{I_{l-1}I_{l-1}}\right)\operatorname{vec}\left(\mathbf{G}^{\geq n+1}\right)$
	$\begin{pmatrix} (3) & -i_{n+1}i_{n+2} & i_N \end{pmatrix} \begin{pmatrix} - & - \\ - & - \end{pmatrix}$
$\operatorname{vec}(\underline{\mathbf{X}}) = \left( (\mathbf{G}_{(n)}^{< n})^{\top} \otimes \mathbf{I}_{I_n} \otimes (\mathbf{G}_{(1)}^{> n})^{\top} \right) \operatorname{vec}\left(\underline{\mathbf{G}}^{(n)}\right)$	
Matricizatio	on
$\mathbf{X}_{(n)} = \mathbf{G}_{(n)}^{(n)} \left( \mathbf{G}_{(n)}^{< n} \otimes \mathbf{G}_{(n)}^{> n} \right)$	$\mathbf{G}^{< n}_{(n)} = \mathbf{G}^{(n-1)}_{(n)} \left( \mathbf{G}^{< n-1}_{(n-1)} \otimes \mathbf{I}_{I_{n-1}} \right)$
$(n) = (2) \left( -(n) = -(1) \right)$	$(n) \qquad (3) \qquad (n-1) \stackrel{\frown}{\smile} \stackrel{-r_{n-1}}{\frown} $
$\mathbf{X}_{} = \mathbf{G}_{}^{\le n} \mathbf{G}_{(1)}^{>n} = \left(\mathbf{G}_{(n+1)}^{n}$	$\mathbf{G}_{(1)}^{>n} = \mathbf{G}_{(1)}^{(n+1)} \left( \mathbf{I}_{I_{n+1}} \otimes \mathbf{G}_{(1)}^{>n+1}  ight)$

(see also Proposition 2.3(g)):

$$\operatorname{vec}\left(\underline{\mathbf{G}}^{\leq n}\right) = \operatorname{vec}\left(\underline{\mathbf{G}}^{\leq n-1} \times^{1} \underline{\mathbf{G}}^{(n)}\right)$$
$$= \left(\mathbf{I}_{I_{1}I_{2}\cdots I_{n-1}} \otimes \mathbf{G}_{(1)}^{(n)\top}\right) \operatorname{vec}\left(\underline{\mathbf{G}}^{\leq n-1}\right), \quad n = 2, 3, \dots, N,$$
$$\operatorname{vec}\left(\underline{\mathbf{G}}^{\geq n}\right) = \operatorname{vec}\left(\underline{\mathbf{G}}^{(n)} \times^{1} \underline{\mathbf{G}}^{\geq n+1}\right)$$
$$= \left(\mathbf{G}_{(3)}^{(n)\top} \otimes \mathbf{I}_{I_{n+1}I_{n+2}\cdots I_{N}}\right) \operatorname{vec}\left(\underline{\mathbf{G}}^{\geq n+1}\right), \quad n = 1, 2, \dots, N-1.$$
(16)

Since  $\operatorname{vec}(\underline{\mathbf{X}}) = \operatorname{vec}(\underline{\mathbf{G}}^{\leq N}) = \operatorname{vec}(\underline{\mathbf{G}}^{\geq 1})$ , we can obtain the following formulas. **Proposition 3.1.** Let  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  admit the TT decomposition in (12). Then

(a) 
$$\operatorname{vec}(\underline{\mathbf{X}}) = \prod_{n=N}^{1} \left( \mathbf{I}_{I_{1}I_{2}\cdots I_{n-1}} \otimes \mathbf{G}_{(1)}^{(n)\top} \right), \text{ with } \mathbf{I}_{I_{1}I_{0}} = 1.$$
  
(b)  $\operatorname{vec}(\underline{\mathbf{X}}) = \prod_{n=1}^{N} \left( \mathbf{G}_{(3)}^{(n)\top} \otimes \mathbf{I}_{I_{n+1}I_{n+2}\cdots I_{N}} \right), \text{ with } \mathbf{I}_{I_{N+1}I_{N}} = 1.$   
(c)  $\operatorname{vec}(\underline{\mathbf{X}}) = \left( \left( \mathbf{G}_{(n)}^{< n} \right)^{\top} \otimes \mathbf{I}_{I_{n}} \otimes \left( \mathbf{G}_{(1)}^{> n} \right)^{\top} \right) \operatorname{vec}\left( \underline{\mathbf{G}}^{(n)} \right).$ 

*Proof.* (a) and (b) follow immediately from (16) and that  $\mathbf{G}_{(1)}^{(1)\top} = \operatorname{vec}(\underline{\mathbf{G}}^{(1)}), \ \mathbf{G}_{(3)}^{(N)\top} = \operatorname{vec}(\underline{\mathbf{G}}^{(N)}).$  (c) can be derived by using Proposition 2.3(g) as

$$\operatorname{vec}\left(\underline{\mathbf{X}}\right) = \operatorname{vec}\left(\underline{\mathbf{G}}^{n}\right)\right)$$
$$= \left(\left(\mathbf{G}_{(n)}^{n}\right)$$
$$= \left(\left(\mathbf{G}_{(n)}^{n}\right)^{\top}\right) \operatorname{vec}\left(\underline{\mathbf{G}}^{(n)}\right)$$
$$= \left(\left(\mathbf{G}_{(n)}^{n}\right)^{\top}\right) \operatorname{vec}\left(\underline{\mathbf{G}}^{(n)}\right).$$

The expression in Proposition 3.1(c) shows that the *n*th TT-core can be separated from the others, see, e.g., [9, 28]. By defining the so-called frame matrix

$$\mathbf{X}^{\neq n} = \left(\mathbf{G}_{(n)}^{< n}\right)^{\top} \otimes \mathbf{I}_{I_n} \otimes \left(\mathbf{G}_{(1)}^{> n}\right)^{\top} \in \mathbb{R}^{I_1 I_2 \cdots I_N \times R_{n-1} I_n R_n}$$
(17)

for  $n = 1, \ldots, N$ , it is simplified as

$$\operatorname{vec}(\underline{\mathbf{X}}) = \mathbf{X}^{\neq n} \operatorname{vec}(\underline{\mathbf{G}}^{(n)}).$$
 (18)

,

We can obtain a similar expression for the vectorization as in (18), where two neighboring TT-cores are separated from the other TT-cores as (see, e.g., [28])

$$\operatorname{vec}(\underline{\mathbf{X}}) = \mathbf{X}^{\neq n, n+1} \operatorname{vec}\left(\underline{\mathbf{G}}^{(n)} \times^{1} \underline{\mathbf{G}}^{(n+1)}\right)$$

where

$$\mathbf{X}^{\neq n,n+1} = \left(\mathbf{G}_{(n)}^{< n}\right)^{\top} \otimes \mathbf{I}_{I_n} \otimes \mathbf{I}_{I_{n+1}} \otimes \left(\mathbf{G}_{(1)}^{> n+1}\right)^{\top} \in \mathbb{R}^{I_1 I_2 \cdots I_N \times R_{n-1} I_n I_{n+1} R_{n+1}}$$
(19)

for  $n = 1, 2, \dots, N - 1$ .

#### **3.2.2** Orthogonalization of core tensors

Left- or right-orthogonalization of 3rd-order TT-cores  $\underline{\mathbf{G}}^{(n)}$  of TT decomposition is defined in the following way, which helps improve convergence of an iterative method using TT decomposition and reduces computational costs [20].

**Definition 3.2** (Left- or right-orthogonality, [19]). For a fixed n = 1, ..., N, the *n*th TT-core  $\underline{\mathbf{G}}^{(n)} \in \mathbb{R}^{R_{n-1} \times I_n \times R_n}$  is called left-orthogonal if

$$\mathbf{G}_{(3)}^{(n)}\mathbf{G}_{(3)}^{(n)\top} = \mathbf{I}_{R_n},\tag{20}$$

and right-orthogonal if

$$\mathbf{G}_{(1)}^{(n)}\mathbf{G}_{(1)}^{(n)\top} = \mathbf{I}_{R_{n-1}}.$$
(21)

During iterations in numerical algorithms, the TT-cores are kept either left- or right-orthogonal by SVD or QR decomposition [20]. For a fixed n, if all the TT-cores on the left, i.e.,  $\underline{\mathbf{G}}^{(m)}, m =$  $1, \ldots, n-1$ , are left-orthogonal, then  $\mathbf{G}_{(n)}^{< n} \in \mathbb{R}^{R_n \times I_1 I_2 \cdots I_{n-1}}$  has orthonormal rows. In the same way, if all the TT-cores on the right, i.e.,  $\underline{\mathbf{G}}^{(m)}, m = n + 1, \ldots, N$ , are right-orthogonal, then  $\mathbf{G}_{(1)}^{> n} \in \mathbb{R}^{R_n \times I_{n+1} I_{n+2} \cdots I_N}$  has orthonormal rows. The proof of the orthonormality of  $\mathbf{G}_{(n)}^{< n}$  and  $\mathbf{G}_{(1)}^{> n}$ can be given by using the tensor operations described in the previous section as follows.

**Proposition 3.2.** Let  $n = 1, 2, \ldots, N$  be fixed.

- (a) If  $\underline{\mathbf{G}}^{(1)}, \underline{\mathbf{G}}^{(2)}, \dots, \underline{\mathbf{G}}^{(n-1)}$  are left-orthogonal, then  $\mathbf{G}_{(n)}^{< n}$  has orthonormal rows.
- (b) If  $\underline{\mathbf{G}}^{(n+1)}, \underline{\mathbf{G}}^{(n+2)}, \dots, \underline{\mathbf{G}}^{(N)}$  are right-orthogonal, then  $\mathbf{G}_{(1)}^{>n}$  has orthonormal rows.

*Proof.* We will prove (a) by induction. (b) can be proved similarly. If n = 1, then  $\mathbf{G}_{(n)}^{< n} = \underline{\mathbf{G}}^{< 1} = 1$ , so we have  $\mathbf{G}_{(n)}^{< n} (\mathbf{G}_{(n)}^{< n})^{\top} = 1 = \mathbf{I}_{R_0}$  for n = 1. Next, we suppose that  $\mathbf{G}_{(k)}^{< k} (\mathbf{G}_{(k)}^{< k})^{\top} = \mathbf{I}_{R_{k-1}}$  under the left-orthogonality of  $\underline{\mathbf{G}}^{(1)}, \ldots, \underline{\mathbf{G}}^{(k-1)}$ . Since  $\underline{\mathbf{G}}^{< k+1} = \underline{\mathbf{G}}^{< k} \times^1 \underline{\mathbf{G}}^{(k)}$  for  $k \leq N$ , we can get the recursive expression (see also Proposition 2.3(f) and Table 4)

$$\mathbf{G}_{(k+1)}^{< k+1} = \mathbf{G}_{(3)}^{(k)} \left( \mathbf{G}_{(k)}^{< k} \otimes \mathbf{I}_{I_k} \right).$$

$$(22)$$

Since  $\mathbf{G}_{(k)}^{< k} (\mathbf{G}_{(k)}^{< k})^{\top} = \mathbf{I}_{R_{k-1}}$ , we have

$$\mathbf{G}_{(k+1)}^{< k+1} \left( \mathbf{G}_{(k+1)}^{< k+1} \right)^{\top} = \mathbf{G}_{(3)}^{(k)} \left( \mathbf{G}_{(k)}^{< k} (\mathbf{G}_{(k)}^{< k})^{\top} \otimes \mathbf{I}_{I_k} \right) \mathbf{G}_{(3)}^{(k)\top} = \mathbf{G}_{(3)}^{(k)} \mathbf{G}_{(3)}^{(k)\top}.$$

From the definition of the left-orthogonality (20) of  $\underline{\mathbf{G}}^{(k)}$ , we can conclude that  $\mathbf{G}_{(k+1)}^{< k+1} (\mathbf{G}_{(k+1)}^{< k+1})^{\top} = \mathbf{I}_{R_k}$  under the left-orthogonality of  $\underline{\mathbf{G}}^{(1)}, \ldots, \underline{\mathbf{G}}^{(k)}$ .

Note that the frame matrix  $\mathbf{X}^{\neq n}$  (17) will have orthonormal columns if both  $\mathbf{G}_{(n)}^{< n}$  and  $\mathbf{G}_{(1)}^{> n}$  have orthonormal rows. The same argument holds for the frame matrix  $\mathbf{X}^{\neq n,n+1}$  (19) with  $\mathbf{G}_{(n)}^{< n}$  and  $\mathbf{G}_{(1)}^{> n+1}$ .

Figure 7(a),(b), and (c) show tensor network diagrams for the TT decomposition of a 4th-order tensor. The left-orthogonalized or right-orthogonalized core tensors are represented by half-filled circles. For example, Figure 7(a) shows that the three among the four core tensors have been left-orthogonalized.

$$\underbrace{\mathbf{G}^{(1)}}_{I_1} \underbrace{\mathbf{G}^{(2)}}_{I_2} \underbrace{\mathbf{G}^{(3)}}_{I_3} \underbrace{\mathbf{G}^{(4)}}_{I_4} \underbrace{\mathbf{G}^{(1)}}_{I_1} \underbrace{\mathbf{G}^{(2)}}_{I_2} \underbrace{\mathbf{G}^{(3)}}_{I_3} \underbrace{\mathbf{G}^{(4)}}_{I_1} \underbrace{\mathbf{G}^{(1)}}_{I_2} \underbrace{\mathbf{G}^{(2)}}_{I_3} \underbrace{\mathbf{G}^{(2)}}_{I_4} \underbrace{\mathbf{G}^{(1)}}_{I_1} \underbrace{\mathbf{G}^{(2)}}_{I_2} \underbrace{\mathbf{G}^{(3)}}_{I_4} \underbrace{\mathbf{G}^{(4)}}_{I_1} \underbrace{\mathbf{G}^{(1)}}_{I_2} \underbrace{\mathbf{G}^{(2)}}_{I_3} \underbrace{\mathbf{G}^{(3)}}_{I_4} \underbrace{\mathbf{G}^{(4)}}_{I_1} \underbrace{\mathbf{G}^{(2)}}_{I_2} \underbrace{\mathbf{G}^{(3)}}_{I_3} \underbrace{\mathbf{G}^{(4)}}_{I_4} \underbrace{\mathbf{G}^{(4)}}_{I_1} \underbrace{\mathbf{G}^{(4)}}_{I_2} \underbrace{\mathbf{G}^{(4)}}_{I_3} \underbrace{\mathbf{G}^{(4)}}_{I_4} \underbrace{\mathbf{G}^{(4)}}_{I_1} \underbrace{\mathbf{G}^{(2)}}_{I_2} \underbrace{\mathbf{G}^{(3)}}_{I_3} \underbrace{\mathbf{G}^{(4)}}_{I_4} \underbrace{\mathbf{G}^{(4)}}_{I_1} \underbrace{\mathbf{G}^{(4)}}_{I_2} \underbrace{\mathbf{G}^{(4)}}_{I_3} \underbrace{\mathbf{G}^{(4)}}_{I_4} \underbrace{\mathbf$$

Figure 7: Tensor network diagrams for TT decomposition of a 4th-order tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times I_2 \times I_3 \times I_4}$ . In each figure, three among the four TT-cores have been (a) left-orthogonalized, (b) right-orthogonalized, or (c) left- or right-orthogonalized.

#### 3.2.3 Uniqueness of TT-ranks and minimal TT decomposition

Note that the TT-ranks  $R_1, \ldots, R_{N-1}$  in the TT decomposition (12) may not be unique, and depend on the decomposition rather than the tensor  $\underline{\mathbf{X}}$  itself.

Mathematical (geometric) properties of TT decomposition are closely related to the so-called separation ranks [19]. The *n*th separation rank,  $S_n$ , of a tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$  is defined as the rank of the *n*th canonical unfolding (i.e., mode- $(1, \ldots, n)$  matricization)  $\mathbf{X}_{< n>}$ , i.e.,

$$S_n = \operatorname{rank}(\mathbf{X}_{< n >})$$

By looking at the canonical unfolding of the TT decomposition (12), we can show the following relationship between the separation ranks and the TT-ranks:

$$S_n \leq R_n$$

for n = 1, ..., N - 1.

The TT decomposition (12) is called minimal or fulfilling the full-rank condition [19] if all TTcores have full left and right ranks, i.e.,

$$R_n = \operatorname{rank}\left(\mathbf{G}_{(3)}^{(n)}\right) \quad \text{and} \quad R_{n-1} = \operatorname{rank}\left(\mathbf{G}_{(1)}^{(n)}\right)$$

for all n = 1, ..., N. Holtz, Rohwedder, and Schneider [19] proved that the TT-ranks of minimal TT decompositions for a tensor  $\underline{\mathbf{X}}$  are unique, that is, if  $\underline{\mathbf{X}}$  admits for a minimal TT decomposition with TT-ranks  $R_n$ , then it holds that

$$R_n = S_n$$

for n = 1, ..., N - 1. Therefore, the TT-ranks of a tensor  $\underline{\mathbf{X}}$  can be defined as the TT-ranks of a minimal TT decomposition of  $\underline{\mathbf{X}}$ , i.e.,  $R_n = \operatorname{rank}(\mathbf{X}_{< n >})$ .

Let  $\mathbb{T}_{\mathrm{TT}}(R_1,\ldots,R_{N-1})$  denote the set of tensors of TT-ranks bounded by  $(R_1,\ldots,R_{N-1})$ . It has been shown that  $\mathbb{T}_{\mathrm{TT}}(R_1,\ldots,R_{N-1})$  is closed [13, 16], which implies that there exists a best TT-ranks- $(R_1,\ldots,R_{N-1})$  approximation of any tensor. Oseledets [33] proved that the TT-SVD algorithm proposed in [33] can return a quasi-optimal TT approximation of a given tensor  $\underline{\mathbf{Y}} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$ in principle. That is, for a tensor  $\underline{\mathbf{Y}}$ , the TT-SVD algorithm returns  $\underline{\hat{\mathbf{X}}} \in \mathbb{T}_{\mathrm{TT}}(R_1,\ldots,R_{N-1})$  such that

$$\left\|\underline{\mathbf{Y}} - \underline{\widehat{\mathbf{X}}}\right\|_{\mathrm{F}} \leq \sqrt{N-1} \cdot \min_{\underline{\mathbf{X}} \in \mathbb{T}_{\mathrm{TT}}(R_1, \dots, R_{N-1})} \left\|\underline{\mathbf{Y}} - \underline{\mathbf{X}}\right\|_{\mathrm{F}},$$

where  $\|\cdot\|_{\rm F}$  is the Frobenius norm. Moreover, the TT-SVD algorithm yields a minimal TT decomposition of a tensor <u>**Y**</u> [19, 33].

Unlike the Tucker decomposition, the TT decomposition requires a storage cost of  $\mathcal{O}(NIR^2)$ , for  $I = \max(I_n)$  and  $R = \max(R_n)$ , that is linear with the order N. Many algorithms based on TT

decomposition such as the approximation, truncation, contraction, and solution to linear systems also have computational complexity linear with N. Further mathematical properties of TT-based algorithms will be developed in the next section.

## 3.3 TT decomposition for large-scale vectors and matrices

We will introduce variations of TT decomposition for representing large-scale vectors, matrices, and low-order tensors. Various representations for TT decomposition of vectors and matrices are summarized in Table 5.

A large-scale vector  $\mathbf{x}$  of length  $I_1 I_2 \cdots I_N$  can be considered to be reshaped into a higher-order tensor  $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  and approximately represented by TT decomposition as

$$\underline{\mathbf{X}} = \underline{\mathbf{X}}^{(1)} \times^{1} \underline{\mathbf{X}}^{(2)} \times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(N)},$$
(23)

where  $\underline{\mathbf{X}}^{(n)} \in \mathbb{R}^{R_{n-1} \times I_n \times R_n}$  are 3rd-order TT-cores. Vectorization of the above TT decomposition yields the TT decomposition for  $\mathbf{x}$ , expressed as a sum of Kronecker products (e.g., see (14))

$$\mathbf{x} = \sum_{r_1=1}^{R_1} \sum_{r_2=1}^{R_2} \cdots \sum_{r_{N-1}=1}^{R_{N-1}} \mathbf{x}_{1,r_1}^{(1)} \otimes \mathbf{x}_{r_1,r_2}^{(2)} \otimes \cdots \otimes \mathbf{x}_{r_{N-2},r_{N-1}}^{(N-1)} \otimes \mathbf{x}_{r_{N-1},1}^{(N)},$$
(24)

where  $\mathbf{x}_{r_{n-1},r_n}^{(n)} = \underline{\mathbf{X}}^{(n)}(r_{n-1},:,r_n) \in \mathbb{R}^{I_n}$  for all  $r_{n-1},r_n$ , and n. The above form can be compactly represented as strong Kronecker products of block matrices

$$\mathbf{x} = \widetilde{\mathbf{X}}^{(1)} |\otimes| \widetilde{\mathbf{X}}^{(2)} |\otimes| \cdots |\otimes| \widetilde{\mathbf{X}}^{(N)},$$
(25)

where  $\widetilde{\mathbf{X}}^{(n)} = [\mathbf{x}_{r_{n-1},r_n}^{(n)}] \in \mathbb{R}^{R_{n-1}I_n \times R_n}$  are block matrices partitioned with  $\mathbf{x}_{r_{n-1},r_n}^{(n)} \in \mathbb{R}^{I_n}$  as

$$\widetilde{\mathbf{X}}^{(n)} = \begin{bmatrix} \mathbf{x}_{1,1}^{(n)} & \cdots & \mathbf{x}_{1,R_n}^{(n)} \\ \vdots & & \vdots \\ \mathbf{x}_{R_{n-1},1}^{(n)} & \cdots & \mathbf{x}_{R_{n-1},R_n}^{(n)} \end{bmatrix} \in \mathbb{R}^{R_{n-1}I_n \times R_n}$$

TT decomposition can also be extended to a representation for linear operators and matrices [32]. Suppose that a large-scale matrix  $\mathbf{A}$  of size  $I_1 I_2 \cdots I_N \times J_1 J_2 \cdots J_N$  is reshaped and permuted into a higher-order tensor  $\underline{\mathbf{A}}$  of size  $I_1 \times J_1 \times I_2 \times J_2 \times \cdots \times I_N \times J_N$ . Similarly to (12), a TT decomposition for  $\underline{\mathbf{A}}$  can be represented as contracted products

$$\underline{\mathbf{A}} = \underline{\mathbf{A}}^{(1)} \times^{1} \underline{\mathbf{A}}^{(2)} \times^{1} \cdots \times^{1} \underline{\mathbf{A}}^{(N)}, \tag{26}$$

where  $\underline{\mathbf{A}}^{(n)} \in \mathbb{R}^{R_{n-1}^A \times I_n \times J_n \times R_n^A}$  are 4th-order TT-cores and  $R_1^A, \ldots, R_{N-1}^A$  are TT-ranks. We assume that  $R_0^A = R_N^A = 1$ . The TT decomposition for the tensor  $\underline{\mathbf{A}}$  in (26) can be equivalently expressed as outer products of slice matrices

$$\underline{\mathbf{A}} = \sum_{r_1^A=1}^{R_1^A} \sum_{r_2^A=1}^{R_2^A} \cdots \sum_{r_{N-1}^A=1}^{R_{N-1}^A} \mathbf{A}_{1,r_1^A}^{(1)} \circ \mathbf{A}_{r_1^A,r_2^A}^{(2)} \circ \cdots \circ \mathbf{A}_{r_{N-2}^A,r_{N-1}^A}^{(N-1)} \circ \mathbf{A}_{r_{N-1}^A,1}^{(N)},$$

where  $\mathbf{A}_{r_{n-1}^A, r_n^A}^{(n)} = \underline{\mathbf{A}}^{(n)}(r_{n-1}^A; ;; ; r_n^A) \in \mathbb{R}^{I_n \times J_n}$  is a slice of the 4th-order core tensor  $\underline{\mathbf{A}}^{(n)} \in \mathbb{R}^{R_{n-1} \times I_n \times J_n \times R_n}$ . We can derive that TT decomposition for the matrix  $\mathbf{A} \in \mathbb{R}^{I_1 \cdots I_N \times J_1 \cdots J_N}$  can be represented as a sum of Kronecker products

$$\mathbf{A} = \sum_{r_1^A=1}^{R_1^A} \sum_{r_2^A=1}^{R_2^A} \cdots \sum_{r_{N-1}^A=1}^{R_{N-1}^A} \mathbf{A}_{1,r_1^A}^{(1)} \otimes \mathbf{A}_{r_1^A,r_2^A}^{(2)} \otimes \cdots \otimes \mathbf{A}_{r_{N-2}^A,r_{N-1}^A}^{(N-1)} \otimes \mathbf{A}_{r_{N-1}^A,1}^{(N)},$$
(27)

Vector TT	Matrix TT	
Strong Kronecker product		
$\mathbf{x} = \widetilde{\mathbf{X}}^{(1)}  \otimes  \widetilde{\mathbf{X}}^{(2)}  \otimes  \cdots  \otimes  \widetilde{\mathbf{X}}^{(N)},$	$\mathbf{A} = \widetilde{\mathbf{A}}^{(1)}  \otimes  \widetilde{\mathbf{A}}^{(2)}  \otimes  \cdots  \otimes  \widetilde{\mathbf{A}}^{(N)},$	
$\widetilde{\mathbf{X}}^{(n)} = \begin{bmatrix} \mathbf{x}_{r_{n-1},r_n}^{(n)} \end{bmatrix} \in \mathbb{R}^{R_{n-1}I_n \times R_n}$	$\widetilde{\mathbf{A}}^{(n)} = \left[\mathbf{A}_{r_{n-1}^A, r_n^A}^{(n)} ight] \in \mathbb{R}^{R_{n-1}^A I_n  imes R_n^A J_n}$	
with each block $\mathbf{x}_{r_{n-1},r_n}^{(n)} \in \mathbb{R}^{I_n}$	with each block $\mathbf{A}_{r_{n-1}^{A},r_{n}^{A}}^{(n)} \in \mathbb{R}^{I_{n} \times J_{n}}$	
Krone	cker product	
$R_1,, R_{N-1}$	$R_{1}^{A},,R_{N-1}^{A}$	
$\mathbf{x} = \sum_{r_1, r_2, r_3} \mathbf{x}_{1, r_1}^{(1)} \otimes \mathbf{x}_{r_1, r_2}^{(2)} \otimes \cdots \otimes \mathbf{x}_{r_{N-1}, 1}^{(N)}$	$\mathbf{A} = \sum_{A = \mathbf{A}} \mathbf{A}_{1,r_1^A}^{(1)} \otimes \mathbf{A}_{r_1^A,r_2^A}^{(2)} \otimes \cdots \otimes \mathbf{A}_{r_{N-1}^A,1}^{(N)}$	
71,,7N-1-1	$r_1^{i_1},,r_{N-1}^{i_N}=1$	
$\operatorname{Contra}_{\mathcal{C}}$	cted product $(1)$ $(2)$ $(1)$ $(N)$	
$\underline{\mathbf{X}} = \underline{\mathbf{X}}^{(1)} \times^{1} \underline{\mathbf{X}}^{(2)} \times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(1)}$	$\underline{\mathbf{A}} = \underline{\mathbf{A}}^{(1)} \times^{1} \underline{\mathbf{A}}^{(2)} \times^{1} \cdots \times^{1} \underline{\mathbf{A}}^{(N)}$	
Oute	er product	
$\underline{\mathbf{X}} = \sum_{r_1, \dots, r_{N-1}=1}^{R_1, \dots, R_{N-1}} \mathbf{x}_{1, r_1}^{(1)} \circ \mathbf{x}_{r_1, r_2}^{(2)} \circ \dots \circ \mathbf{x}_{r_{N-1}, 1}^{(N)}$	$\underline{\mathbf{A}} = \sum_{r_1^A, \dots, r_{N-1}^A = 1}^{R_1^A, \dots, R_{N-1}^A} \mathbf{A}_{1, r_1^A}^{(1)} \circ \mathbf{A}_{r_1^A, r_2^A}^{(2)} \circ \dots \circ \mathbf{A}_{r_{N-1}^A, 1}^{(N)}$	
Scala	ar product	
$x_{i_1,\dots,i_N} = \mathbf{x}(\overline{i_1\cdots i_N}) = \\ R_1,\dots,R_{N-1}$	$a_{i_1,j_1,i_2,j_2,\ldots,i_N,j_N}^{I} = \mathbf{A}(\overline{i_1\cdots i_N},\overline{j_1\cdots j_N}) = a_{\substack{R_1^A,\ldots,R_{N-1}^A}}^{I}$	
$\sum_{r_1,\dots,r_{N-1}=1}^{N} x_{1,i_1,r_1}^{(1)} x_{r_1,i_2,r_2}^{(2)} \cdots x_{r_{N-1},i_N,1}^{(N)}$	$\sum_{\substack{r_1^A,\dots,r_{N-1}^A \\ r_1^A,i_1,j_1,r_1^A}} a_{r_1^A,i_2,j_2,r_2^A}^{(2)} \cdots a_{r_{N-1}^A,i_N,j_N,1}^{(N)}$	
Matr	ix product	
$x_{i_1,\dots,i_N} = \mathbf{X}_{i_1}^{(1)} \mathbf{X}_{i_2}^{(2)} \cdots \mathbf{X}_{i_N}^{(N)}$	$a_{i_1,j_1,\dots,i_N,j_N} = \mathbf{A}_{i_1,j_1}^{(1)} \mathbf{A}_{i_2,j_2}^{(2)} \cdots \mathbf{A}_{i_N,j_N}^{(N)}$	
Vector/mat	rix representation	
$\mathbf{x} = \prod_{\substack{n=N \ i \in N}}^{1} \left( \mathbf{I}_{I_1 I_2 \cdots I_{n-1}} \otimes \mathbf{X}_{(1)}^{(n)  op}  ight)$	$\mathbf{A} = \prod_{n=1}^{N} \left( \mathbf{I}_{J_1 J_2 \cdots J_{n-1}} \otimes \mathbf{A}_{<2>}^{(n)} \otimes \mathbf{I}_{I_{n+1} I_{n+2} \cdots I_N} \right)$	
$\mathbf{x} = \prod_{n=1}^{N} \left( \mathbf{X}_{(3)}^{(n) op} \otimes \mathbf{I}_{I_{n+1} \cdots I_N}  ight)$		
$\mathbf{x} = \left( (\mathbf{X}_{(n)}^{< n})^\top \otimes \mathbf{I}_{I_n} \otimes (\mathbf{X}_{(1)}^{> n})^\top \right) \operatorname{vec} \left( \underline{\mathbf{X}}^{(n)} \right)$		

Table 5: Various representations for TT decomposition of a large-scale vector  $\mathbf{x} \in \mathbb{R}^{I_1 \cdots I_N}$  and a matrix  $\mathbf{A} \in \mathbb{R}^{I_1 \cdots I_N \times J_1 \cdots J_N}$ .

where  $\mathbf{A}_{r_{n-1}^A,r_n^A}^{(n)} = \underline{\mathbf{A}}^{(n)}(r_{n-1}^A; :, :, r_n^A) \in \mathbb{R}^{I_n \times J_n}$ . The representation in (27) can be equivalently expressed as strong Kronecker product of block matrices

$$\mathbf{A} = \widetilde{\mathbf{A}}^{(1)} \otimes \widetilde{\mathbf{A}}^{(2)} \otimes \cdots \otimes \widetilde{\mathbf{A}}^{(N)}, \tag{28}$$

where  $\widetilde{\mathbf{A}}^{(n)} = [\mathbf{A}_{r_{n-1}^A, r_n^A}^{(n)}] \in \mathbb{R}^{R_{n-1}^A I_n \times R_n^A J_n}$  are block matrices partitioned with the slice matrices  $\mathbf{A}_{r_{n-1}^A, r_n^A}^{(n)} \in \mathbb{R}^{I_n \times J_n}$  as

$$\widetilde{\mathbf{A}}^{(n)} = \begin{bmatrix} \mathbf{A}_{1,1}^{(n)} & \cdots & \mathbf{A}_{1,R_n^A}^{(n)} \\ \vdots & & \vdots \\ \mathbf{A}_{R_{n-1}^A,1}^{(n)} & \cdots & \mathbf{A}_{R_{n-1}^A,R_n^A}^{(n)} \end{bmatrix} \in \mathbb{R}^{R_{n-1}^A I_n \times R_n^A J_n}$$

The strong Kronecker product representation (28) for TT decomposition is useful, especially for representing large-scale high-dimensional operators such as discrete Laplace operator and multilevel (hierarchical) Toeplitz, Hankel, circulant, banded diagonal matrices [21, 22].

We call the TT decomposition in (24) and (25) for vectors as the vector TT decomposition, and the TT decomposition in (27) and (28) for matrices as the matrix TT decomposition. Note that the blocks  $\mathbf{x}_{r_{n-1},r_n}^{(n)} \in \mathbb{R}^{I_n}$  and  $\mathbf{A}_{r_{n-1},r_n}^{(n)} \in \mathbb{R}^{I_n \times J_n}$  in the expressions can be further generalized to higher-order tensors, which leads to TT decomposition for multilinear operators and higher-order tensors.

## 4 Basic operations using TT decomposition

Based on the introduced TT decompositions, basic operations on large-scale vectors and matrices such as matrix-by-vector multiplication can be performed fast and conveniently. For basic algebraic operations on tensors represented by TT decomposition, it is important to perform all the operations based on the factor (core) tensors in TT decomposition and avoid the explicit calculation of the full tensors. In this section we present basic operations on vectors, matrices, and higher-order tensors represented by TT decomposition, in simple and very efficient forms by using the notations introduced in the previous sections. The basic operations include addition, scalar multiplication, direct sum, Hadamard product, Kronecker product, contraction, matrix-by-vector product, matrixby-matrix product, and quadratic form. Figure 8 illustrates tensor network diagrams for the matrix TT decomposition, the matrix-by-vector multiplication, and the quadratic form, represented by TT decomposition.

Note that such operations usually increase the TT-ranks, which requires truncation (rounding) in the following step [33]. In addition, the matrix-by-vector product and quadratic form are very important for computational algorithms in optimization problems such as linear equations [10, 20, 34] and eigenvalue problems [9, 27, 28]. Tables 6 and 7 summarize the representations for the basic operations on the tensors represented by TT decomposition.

## 4.1 Addition and scalar multiplication

Let  $\underline{\mathbf{X}} = \underline{\mathbf{X}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{X}}^{(N)} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$  and  $\underline{\mathbf{Y}} = \underline{\mathbf{Y}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{Y}}^{(N)} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$  be TT tensors with TT-ranks  $\{R_n^X\}$  and  $\{R_n^Y\}$ . Under the assumption that the 1st and the Nth TT-cores are matrices (i.e., 2nd-order tensors,) the sum  $\underline{\mathbf{Z}} = \underline{\mathbf{X}} + \underline{\mathbf{Y}} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$  can be expressed by the TT decomposition (see, Proposition 2.4)

$$\underline{\mathbf{Z}} = \left(\mathbf{X}^{(1)} \boxplus \mathbf{Y}^{(1)}\right) \times^{1} \left(\underline{\mathbf{X}}^{(2)} \boxplus \underline{\mathbf{Y}}^{(2)}\right) \times^{1} \cdots \times^{1} \left(\mathbf{X}^{(N)} \boxplus \mathbf{Y}^{(N)}\right).$$
(29)

Table 6: TT representations for basic operations on tensors represented by TT decomposition.

Operation	TT-cores
TT (global)*	
$\underline{\mathbf{Z}} = \underline{\mathbf{X}} + \underline{\mathbf{Y}} = (\mathbf{X}^{(2)})$	$\mathbf{Y}^{(1)} \boxplus \mathbf{Y}^{(1)} $ $ imes^1 \left( \underline{\mathbf{X}}^{(2)} \boxplus \underline{\mathbf{Y}}^{(2)} \right)  imes^1 \cdots  imes^1 \left( \mathbf{X}^{(n)} \boxplus \mathbf{Y}^{(n)} \right)$
Ten	$\underline{\mathbf{Z}}^{(n)} = \underline{\mathbf{X}}^{(n)} \stackrel{\frown}{\boxplus} \underline{\mathbf{Y}}^{(n)}$
Mat	$\mathbf{Z}_{i_n}^{(n)} = \mathbf{X}_{i_n}^{(n)} \oplus \mathbf{Y}_{i_n}^{(n)}$
Vec	$\mathbf{z}_{s_{n-1},s_n}^{(n)} = \mathbf{x}_{s_{n-1},s_n}^{(n)};  \mathbf{y}_{s_{n-1}-R_{n-1}^X,s_n-R_n^X}^{(n)};  0_{I_n}$
$\underline{\mathbf{Z}} = \underline{\mathbf{X}} \oplus \underline{\mathbf{Y}} = (\mathbf{X}^{(2)})$	$(\mathbf{X}^{(1)} \oplus \mathbf{Y}^{(1)})  imes^1 \left( \underline{\mathbf{X}}^{(2)} \oplus \underline{\mathbf{Y}}^{(2)}  ight)  imes^1 \cdots  imes^1 \left( \mathbf{X}^{(n)} \oplus \mathbf{Y}^{(n)}  ight)$
Ten	$\underline{\mathbf{Z}}^{(n)} = \underline{\mathbf{X}}^{(n)} \oplus \underline{\mathbf{Y}}^{(n)}$
Mat	$\mathbf{Z}_{k_n}^{(n)} = \begin{cases} \mathbf{X}_{k_n}^{(n)} \oplus 0_{R_{n-1}^Y \times R_n^Y} & \text{if } 1 \le k_n \le I_n \\ 0 & \mathbf{y}_{n-1}^{(n)} & \text{if } I \le k_n \le I_n \end{cases}$
3.7	$(0_{R_{n-1}^{X} \times R_{n}^{X}} \oplus 1_{k_{n}-I_{n}} \text{ if } I_{n} < k_{n} \leq I_{n} + J_{n}$
vec	$\mathbf{z}_{s_{n-1},s_n} = \mathbf{x}_{s_{n-1},s_n} \oplus \mathbf{U}_{J_n}; 0_{I_n} \oplus \mathbf{y}_{s_{n-1}-R_{n-1},s_n-R_n^X}; 0_{I_n+J_n}$
$\underline{\mathbf{Z}} = \underline{\mathbf{X}} \circledast \underline{\mathbf{Y}} = \left(\underline{\mathbf{X}}^{U}\right)$	$\overset{(1)}{\longrightarrow} \underbrace{\mathbf{Y}}^{(1)} \times \overset{(1)}{\longrightarrow} \underbrace{(\underline{\mathbf{X}}^{(2)} \boxtimes \underline{\mathbf{Y}}^{(2)})}_{(1)} \times \overset{(1)}{\longrightarrow} \times \overset{(1)}{\longrightarrow} \underbrace{(\underline{\mathbf{X}}^{(n)} \boxtimes \underline{\mathbf{Y}}^{(n)})}_{(1)}$
Ten	$\underline{\mathbf{Z}}^{(n)} = \underline{\mathbf{X}}^{(n)} \boxtimes \underline{\mathbf{Y}}^{(n)}$
Mat	$\mathbf{Z}_{i_n}^{(n)} = \mathbf{X}_{i_n}^{(n)} \otimes \mathbf{Y}_{i_n}^{(n)} $
Vec	$\mathbf{z}_{s_{n-1},s_n}^{x} = \mathbf{x}_{r_{n-1},r_n}^{x} \circledast \mathbf{y}_{r_{n-1},r_n}^{y}  (s_n = r_n^{A} r_n^{n})$
$\underline{\mathbf{Z}} = \underline{\mathbf{X}} \otimes \underline{\mathbf{Y}} = \left(\underline{\mathbf{X}}^{(1)}\right)$	$\mathbf{Y}^{(1)} \otimes \mathbf{Y}^{(1)} $ $ imes^1 \left( \mathbf{X}^{(2)} \otimes \mathbf{Y}^{(2)} \right)  imes^1 \cdots  imes^1 \left( \mathbf{X}^{(n)} \otimes \mathbf{Y}^{(n)} \right)$
Ten	$\underline{\mathbf{Z}}^{(n)} = \underline{\mathbf{X}}^{(n)} \otimes \underline{\mathbf{Y}}^{(n)}$
Mat	$\mathbf{Z}_{k_n}^{(n)} = \mathbf{X}_{i_n}^{(n)} \otimes \mathbf{Y}_{j_n}^{(n)}  (k_n = \overline{i_n j_n})$
Vec	$\mathbf{z}_{s_{n-1},s_n}^{(n)} = \mathbf{x}_{r_{n-1}^X,r_n^X}^{(n)} \otimes \mathbf{y}_{r_{n-1}^Y,r_n^Y}^{(n)}  (s_n = \overline{r_n^X r_n^Y})$
$\underline{\mathbf{Z}} = \underline{\mathbf{X}} \circ \underline{\mathbf{Y}} = \underline{\mathbf{X}}^{(1)}$	$\times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(N)} \times^{1} \underline{\mathbf{Y}}^{(1)} \times^{1} \cdots \times^{1} \underline{\mathbf{Y}}^{(N)}$
Ten	$\underline{\mathbf{Z}}^{(n)} = \underline{\mathbf{X}}^{(n)} \ (n \le N); \ \underline{\mathbf{Y}}^{(n-N)} \ (n > N)$
Mat	$\mathbf{Z}_{i_n}^{(n)} = \mathbf{X}_{i_n}^{(n)}  (n \le N); \ \mathbf{Y}_{i_n}^{(n-N)}  (n > N)$
Vec	$\mathbf{z}_{s_{n-1},s_n}^{(n)} = \mathbf{x}_{s_{n-1},s_n}^{(n)} \ (n \le N); \ \mathbf{y}_{s_{n-1},s_n}^{(n-N)} \ (n > N)$
$\underline{\mathbf{Z}} = \underline{\mathbf{X}} \times_n \mathbf{A} = \underline{\mathbf{X}}^{(1)}$	$\mathbf{X}^{(1)} \times^{1} \cdots \times^{1} \mathbf{X}^{(n-1)} \times^{1} \left( \mathbf{X}^{(n)} \times_{2} \mathbf{A} \right) \times^{1} \mathbf{X}^{(n+1)} \times^{1} \cdots \times^{1} \mathbf{X}^{(N)}$
Ten	$\underline{\mathbf{Z}}^{(m)} = \underline{\mathbf{X}}^{(m)}  (m \neq n); \ \underline{\mathbf{X}}^{(m)} \times_2 \mathbf{A} \ (m = n)$
Mat	$\mathbf{Z}_{i_m}^{(m)} = \mathbf{X}_{i_m}^{(m)}  (m \neq n); \ \underline{\mathbf{X}}^{(m)} \overline{\mathbf{X}}_2 \mathbf{a}_{i_m,:} \ (m = n)$
Vec	$\mathbf{z}_{s_{m-1},s_m}^{(m)} = \mathbf{x}_{s_{m-1},s_m}^{(m)} \ (m \neq n); \ \mathbf{A}\mathbf{x}_{s_{m-1},s_m}^{(m)} \ (m = n)$
$\underline{\mathbf{Z}} = \underline{\mathbf{X}} \times^1 \underline{\mathbf{Y}} = \underline{\mathbf{X}}^{(1)}$	$\mathbf{X}^{(1)} \times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(N-1)} \times^{1} \left( \mathbf{X}^{(N)} \mathbf{Y}^{(1)} \times^{1} \underline{\mathbf{Y}}^{(2)} \right) \times^{1} \underline{\mathbf{Y}}^{(3)} \times^{1} \cdots \times^{1} \underline{\mathbf{Y}}^{(N)}$
Ten	$\underline{\mathbf{Z}}^{(n)} = \underline{\mathbf{X}}^{(n)} \ (n < N); \ \mathbf{X}^{(N)} \mathbf{Y}^{(1)} \times^{1} \underline{\mathbf{Y}}^{(2)}; \ \underline{\mathbf{Y}}^{(n-N+2)} \ (N < n)$
Mat	$\mathbf{Z}_{i_n}^{(n)} = \mathbf{X}_{i_n}^{(n)} \ (n < N); \ \mathbf{X}^{(N)} \mathbf{Y}^{(1)} \mathbf{Y}_{i_n}^{(2)}; \ \mathbf{Y}_{i_n}^{(n-N+2)} \ (N < n)$
Vec	$\mathbf{z}_{s_{n-1},s_n}^{(n)} = \mathbf{x}_{s_{n-1},s_n}^{(n)} \ (n < N); \ \mathbf{x}_{s_{n-1}}^{(N) \top} \mathbf{Y}^{(1)} \mathbf{Y}^{(2)}_{:,:,s_n}; \ \mathbf{y}_{s_{n-1},s_n}^{(n-N+2)} \ (N < n)$
$z = \langle \underline{\mathbf{X}}, \underline{\mathbf{Y}} \rangle$	$\mathbf{r}(n) = \left(\mathbf{r}(n) - \mathbf{r}(n)\right) - \mathbf{r}$
Ten	$\mathbf{Z}^{(n)} = \left(\underline{\mathbf{X}}^{(n)} \boxtimes \underline{\mathbf{Y}}^{(n)}\right) \times_2 1_{I_n}$
Mat	$\mathbf{Z}^{(n)} = \sum_{i_n} \mathbf{X}^{(n)}_{i_n} \otimes \mathbf{Y}^{(n)}_{i_n}$
Vec	$z_{s_{n-1},s_n}^{(n)} = \left\langle \mathbf{x}_{r_{n-1},r_n^X}^{(n)}, \mathbf{y}_{r_{n-1},r_n^Y}^{(n)} \right\rangle  \left( s_n = r_n^X r_n^Y \right)$

<sup>\*</sup> Ten = Tensors (cores), Mat = Matrices (slices), Vec = Vectors (fibers)

Table 7: TT representations for basic operations on tensors represented by vector TT and matrix TT decompositions. \_\_\_\_

Operation	TT-cores
Vector TT & Ma	trix TT*
$\mathbf{Z} = \mathbf{A} + \mathbf{B} = \begin{bmatrix} \hat{A} \end{bmatrix}$	$\begin{split} \widetilde{\mathbf{A}}^{(1)}  \widetilde{\mathbf{B}}^{(1)} \end{bmatrix} \ket{\otimes} \begin{bmatrix} \widetilde{\mathbf{A}}^{(2)} & 0 \\ 0 & \widetilde{\mathbf{B}}^{(2)} \end{bmatrix} \ket{\otimes} \cdots \ket{\otimes} \begin{bmatrix} \widetilde{\mathbf{A}}^{(N-1)} & 0 \\ 0 & \widetilde{\mathbf{B}}^{(N-1)} \end{bmatrix} \ket{\otimes} \begin{bmatrix} \widetilde{\mathbf{A}}^{(N)} \\ \widetilde{\mathbf{B}}^{(N)} \end{bmatrix} = \mathbf{A}^{(n)}$
DIK	$\mathbf{Z}^{(n)} = \mathbf{A}^{(n)} \boxplus \mathbf{B}^{(n)}$ $= \mathbf{Z}^{(n)} - \mathbf{A}^{(n)} \boxplus \mathbf{B}^{(n)}$
Vec	$\mathbf{Z}_{s_{n},j_{n}}^{(n)} = \mathbf{A}_{s_{n},j_{n}}^{(n)} \oplus \mathbf{D}_{s_{n},j_{n}}^{(n)} \\ \mathbf{Z}_{s_{n-1},s_{n}}^{(n)} = \mathbf{A}_{s_{n-1},s_{n}}^{(n)}; \mathbf{B}_{s_{n-1}-R_{n-1},s_{n}-R_{n}^{A}}^{(n)}; 0$
$\mathbf{Z} = \mathbf{A} \otimes \mathbf{B} = \widetilde{\mathbf{A}}$	$\overset{(1)}{\cong} \otimes \cdots \otimes \widetilde{\mathbf{A}}^{(N)} \otimes \widetilde{\mathbf{B}}^{(1)} \otimes \cdots \otimes \widetilde{\mathbf{B}}^{(N)} \overset{\widetilde{\mathbf{B}}}{\cong} \overset{\widetilde{\mathbf{B}}^{(N)}}{\boxtimes}$
Blk	$\widetilde{\mathbf{Z}}^{(n)} = \widetilde{\mathbf{A}}^{(n)}  (n \le N); \ \widetilde{\mathbf{B}}^{(n-N)}  (n > N)$
Mat	$\mathbf{Z}_{i_{n},j_{n}}^{(n)} = \mathbf{A}_{i_{n},j_{n}}^{(n)}  (n \le N); \ \mathbf{B}_{i_{n},j_{n}}^{(n-N)}  (n > N)$
Vec	$\mathbf{Z}_{s_{n-1},s_n}^{(n)} = \mathbf{A}_{s_{n-1},s_n}^{(n)} \ (n \le N); \ \mathbf{B}_{s_{n-1},s_n}^{(n-N)} \ (n > N)$
$\underline{\mathbf{Z}} = \underline{\mathbf{X}} \times_n \mathbf{A} = \Big($	$\left( \underline{\widetilde{\mathbf{X}}}^{(1)} \times_n \widetilde{\mathbf{A}}^{(1)} \right)  \otimes  \cdots  \otimes  \left( \underline{\widetilde{\mathbf{X}}}^{(N)} \times_n \widetilde{\mathbf{A}}^{(N)} \right)$
Bll	$\widetilde{\boldsymbol{z}}^{(n)} - \widetilde{\boldsymbol{x}}^{(n)} \vee \widetilde{\boldsymbol{\lambda}}^{(n)}$
Dik Mət	$\underline{\underline{\boldsymbol{\mathcal{L}}}}_{n}^{(n)} - \sum_{\boldsymbol{\mathcal{X}}} \mathbf{X}^{(n)} \otimes \mathbf{A}^{(n)}$
Vec	$\mathbf{Z}_{ijk}^{(n)} = \mathbf{X}_{ijt}^{(n)} \qquad \mathbf{X}_{kt}^{(n)} \qquad (s_r = \overline{r^X r^A})$
T ( )	$ \frac{\underline{\neg}}{\mathbf{x}_{n-1}, s_n} \xrightarrow{\mathbf{x}_n} r_{n-1}^X, r_n^X \land n \stackrel{\mathbf{x}_n}{\mathbf{x}_{n-1}}, r_n^A \xrightarrow{(o_n, \dots, o_n, \dots, o_n)} $
$z = \mathbf{x} \cdot \mathbf{y} = \langle \mathbf{x}, \mathbf{y} \rangle$	$\widetilde{\mathbf{A}}^{(1)} \times \widetilde{\mathbf{Y}}^{(1)} )  \otimes  \cdots  \otimes  \left( \mathbf{A}^{(1)} \times \widetilde{\mathbf{Y}}^{(1)} \right)$ $\widetilde{\mathbf{C}}^{(n)} = \widetilde{\mathbf{T}}^{(n)}  \otimes  1 \widetilde{\mathbf{T}}^{(n)} $
Blk	$\mathbf{Z}^{(n)} = \mathbf{X}^{(n)} \times^{1} \mathbf{Y}^{(n)}$ $\mathbf{Z}^{(n)} = \sum_{k=1}^{n} \mathbf{X}^{(n)} \odot \mathbf{X}^{(n)}$
Mat	$\mathbf{\Sigma}^{(i)} = \sum_{i_n} \mathbf{X}_{i_n}^{(i)} \otimes \mathbf{Y}_{i_n}^{(i)} $
Vec	$z_{s_{n-1},s_n}^{\star} = \left\langle \mathbf{x}_{r_{n-1},r_n^{\star}}^{\star}, \mathbf{y}_{r_{n-1},r_n^{\star}}^{\star} \right\rangle (s_n = r_n^{\star} r_n^{\star})$
$\mathbf{z} = \mathbf{A}\mathbf{x} = (\mathbf{A}^{(1)})$	$ \overset{\times^{1} \mathbf{X}^{(1)}}{\underset{\sim}{\sim}}  \otimes  \cdots  \otimes  \left( \mathbf{A}^{(N)} \times^{1} \mathbf{X}^{(N)} \right) $
Blk	$\widetilde{\mathbf{Z}}^{(n)} = \widetilde{\mathbf{A}}^{(n)} \times {}^{1} \widetilde{\mathbf{X}}^{(n)} $
Mat	$\mathbf{Z}_{i_n}^{(n)} = \sum_{j_n} \mathbf{A}_{i_n, j_n}^{(n)} \otimes \mathbf{X}_{j_n}^{(n)}$
Vec	$\mathbf{z}_{s_{n-1},s_n}^{(n)} = \mathbf{A}_{r_{n-1}^A,r_n^A}^{(n)} \mathbf{x}_{r_{n-1},r_n}^{(n)}  (s_n = r_n^A r_n)$
$\mathbf{Z} = \mathbf{A}\mathbf{B} = \left(\widetilde{\mathbf{A}}^{(1)}\right)$	$ \mathbf{X}^{(1)} \times^1 \widetilde{\mathbf{B}}^{(1)} \rangle  \otimes  \cdots \otimes  \langle \widetilde{\mathbf{A}}^{(N)} \times^1 \widetilde{\mathbf{B}}^{(N)} \rangle$
Blk	$\widetilde{\mathbf{Z}}^{(n)} = \widetilde{\mathbf{A}}^{(n)} \times^1 \widetilde{\mathbf{B}}^{(n)}$
Mat	$\mathbf{Z}_{i_n,j_n}^{(n)} = \sum_{k_n} \mathbf{A}_{i_n,k_n}^{(n)} \otimes \mathbf{B}_{k_n,j_n}^{(n)}$
Vec	$\mathbf{Z}_{s_{n-1},s_n}^{(n)} = \mathbf{A}_{r_n^A \ 1,r_n^A}^{(n)} \mathbf{B}_{r_n^B \ 1,r_n^B}^{(n)} \left(s_n = \overline{r_n^A r_n^B}\right)$
$z = \mathbf{x}^\top \mathbf{A} \mathbf{x} = \langle \mathbf{x},$	$\mathbf{A}\mathbf{x}\rangle = \left(\widetilde{\mathbf{X}}^{(1)} \times^{1} \widetilde{\mathbf{A}}^{(1)} \times^{1} \widetilde{\mathbf{X}}^{(1)}\right)  \otimes  \cdots  \otimes  \left(\widetilde{\mathbf{X}}^{(N)} \times^{1} \widetilde{\mathbf{A}}^{(N)} \times^{1} \widetilde{\mathbf{X}}^{(N)}\right)$
Blk	$\widetilde{\mathbf{Z}}^{(n)} = \widetilde{\mathbf{X}}^{(n)} \times^{1} \widetilde{\mathbf{A}}^{(n)} \times^{1} \widetilde{\mathbf{X}}^{(n)}$
Mat	$\mathbf{Z}^{(n)} = \sum_{i_n} \sum_{j_n} \mathbf{X}^{(n)}_{i_n} \otimes \mathbf{A}^{(n)}_{i_n, j_n} \otimes \mathbf{X}^{(n)}_{j_n}$
Vec	$z_{s_{n-1},s_n}^{(n)} = \left\langle \mathbf{x}_{r_{n-1}',r_n'}^{(n)}, \mathbf{A}_{r_{n-1}',r_n^A}^{(n)} \mathbf{x}_{r_{n-1},r_n}^{(n)} \right\rangle  (s_n = \overline{r_n' r_n^A r_n})$

\* Blk = Block matrices, Mat = Matrices (slices), Vec = Vectors (fibers)



Figure 8: (a) A matrix  $\mathbf{A} \in \mathbb{R}^{I_1 I_2 I_3 I_4 \times J_1 J_2 J_3 J_4}$  in matrix TT decomposition, (b) matrix-by-vector product  $\mathbf{y} = \mathbf{A}\mathbf{x}$  in TT decomposition, and (c) quadratic form  $\mathbf{x}^{\top}\mathbf{A}\mathbf{x}$  when  $I_n = J_n, n = 1, \dots, 4$ .

Note that each TT-core of the sum  $\underline{\mathbf{Z}} = \underline{\mathbf{X}} + \underline{\mathbf{Y}}$  is written as the partial direct sum of the corresponding TT-cores.

Alternatively, each entry of the sum can be represented as products of slice matrices of TT-cores

$$egin{aligned} & z_{i_1,i_2,...,i_N} = igg( \mathbf{X}_{i_1}^{(1)} \oplus \mathbf{Y}_{i_1}^{(1)} igg) igg( \mathbf{X}_{i_2}^{(2)} \oplus \mathbf{Y}_{i_2}^{(2)} igg) \cdots igg( \mathbf{X}_{i_N}^{(N)} \oplus \mathbf{Y}_{i_N}^{(N)} igg) \ & = igg[ \mathbf{X}_{i_1}^{(1)} \quad \mathbf{Y}_{i_1}^{(1)} igg] igg[ \mathbf{X}_{i_2}^{(2)} & \ & \mathbf{Y}_{i_2}^{(2)} igg] \cdots igg[ \mathbf{X}_{i_N}^{(N)} \ & \mathbf{Y}_{i_N}^{(2)} igg], \end{aligned}$$

where the first and the last factor matrices are assumed row and column vectors, respectively. The TT-ranks of the above TT decomposition for  $\underline{\mathbf{X}} + \underline{\mathbf{Y}}$  are the sums,  $\{R_n^X + R_n^Y\}$ .

On the other hand, multiplication of a TT tensor  $\underline{\mathbf{X}}$  with a scalar  $c \in \mathbb{R}$  can be obtained by simply multiplying one core, e.g.,  $\underline{\mathbf{X}}^{(1)}$ , with c as  $c\underline{\mathbf{X}}^{(1)}$ . This does not increase the TT-ranks.

We note that that the set of tensors with TT-ranks bounded by  $\{R_n\}$  is not convex, since a linear combination  $c\mathbf{X} + (1-c)\mathbf{Y}$  generally increases the TT-ranks, which may exceed  $\{R_n\}$ .

### 4.2 Direct sum

Let  $\underline{\mathbf{X}} = \underline{\mathbf{X}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{X}}^{(N)} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$  and  $\underline{\mathbf{Y}} = \underline{\mathbf{Y}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{Y}}^{(N)} \in \mathbb{R}^{J_1 \times \cdots \times J_N}$  be TT tensors with TT-ranks  $\{R_n^X\}$  and  $\{R_n^Y\}$ . Under the assumption that the 1st and the Nth TT-cores are matrices (i.e., 2nd-order tensors,) the direct sum  $\underline{\mathbf{Z}} = \underline{\mathbf{X}} \oplus \underline{\mathbf{Y}}$  can be expressed by the TT decomposition (see, Proposition 2.4)

$$\underline{\mathbf{Z}} = \left(\mathbf{X}^{(1)} \oplus \mathbf{Y}^{(1)}\right) \times^{1} \left(\underline{\mathbf{X}}^{(2)} \oplus \underline{\mathbf{Y}}^{(2)}\right) \times^{1} \cdots \times^{1} \left(\mathbf{X}^{(N)} \oplus \mathbf{Y}^{(N)}\right).$$
(30)

The TT-ranks of the above TT decomposition for  $\underline{\mathbf{X}} \oplus \underline{\mathbf{Y}}$  are the sums,  $\{R_n^X + R_n^Y\}$ .

### 4.3 Hadamard product

The Hadamard (elementwise) product  $\underline{\mathbf{Z}} = \underline{\mathbf{X}} \otimes \underline{\mathbf{Y}}$  of two TT tensors  $\underline{\mathbf{X}} = \underline{\mathbf{X}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{X}}^{(N)} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$  and  $\underline{\mathbf{Y}} = \underline{\mathbf{Y}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{Y}}^{(N)} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$  can be expressed by TT decomposition as (see, Proposition 2.4)

$$\underline{\mathbf{Z}} = \left(\underline{\mathbf{X}}^{(1)} \boxtimes \underline{\mathbf{Y}}^{(1)}\right) \times^{1} \left(\underline{\mathbf{X}}^{(2)} \boxtimes \underline{\mathbf{Y}}^{(2)}\right) \times^{1} \cdots \times^{1} \left(\underline{\mathbf{X}}^{(N)} \boxtimes \underline{\mathbf{Y}}^{(N)}\right).$$
(31)

As an alternative representation, each entry can be written as products of slice matrices of TT-cores

$$z_{i_1,i_2,\ldots,i_N} = \left(\mathbf{X}_{i_1}^{(1)} \otimes \mathbf{Y}_{i_1}^{(1)}\right) \left(\mathbf{X}_{i_2}^{(2)} \otimes \mathbf{Y}_{i_2}^{(2)}\right) \cdots \left(\mathbf{X}_{i_N}^{(N)} \otimes \mathbf{Y}_{i_N}^{(N)}\right)$$

The TT-ranks for the above Hadamard product representation are the multiplications of individual ranks,  $\{R_n^X R_n^Y\}$ .

### 4.4 Kronecker product

The Kronecker product  $\underline{\mathbf{Z}} = \underline{\mathbf{X}} \otimes \underline{\mathbf{Y}}$  of two TT tensors  $\underline{\mathbf{X}} = \underline{\mathbf{X}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{X}}^{(N)} \in \mathbb{R}^{I_1 \times \cdots \times I_N}$  and  $\underline{\mathbf{Y}} = \underline{\mathbf{Y}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{Y}}^{(N)} \in \mathbb{R}^{J_1 \times \cdots \times J_N}$  with TT-ranks  $\{R_n^X\}$  and  $\{R_n^Y\}$  can be expressed by TT decomposition as (see, Proposition 2.4)

$$\underline{\mathbf{Z}} = \left(\underline{\mathbf{X}}^{(1)} \otimes \underline{\mathbf{Y}}^{(1)}\right) \times^{1} \left(\underline{\mathbf{X}}^{(2)} \otimes \underline{\mathbf{Y}}^{(2)}\right) \times^{1} \cdots \times^{1} \left(\underline{\mathbf{X}}^{(N)} \otimes \underline{\mathbf{Y}}^{(N)}\right).$$
(32)

The TT-ranks of the above TT decomposition for  $\underline{\mathbf{X}} \otimes \underline{\mathbf{Y}}$  are the multiplications of individual ranks,  $\{R_n^X R_n^Y\}$ .

## 4.5 Full contraction: Inner product

The contraction of two tensors  $\underline{\mathbf{A}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  and  $\underline{\mathbf{B}} \in \mathbb{R}^{I_1 \times I_2 \times \cdots \times I_N}$  is defined by

$$\langle \underline{\mathbf{A}}, \underline{\mathbf{B}} \rangle = \sum_{i_1=1}^{I_1} \sum_{i_2=1}^{I_2} \cdots \sum_{i_N=1}^{I_N} \underline{\mathbf{A}}(i_1, i_2, \dots, i_N) \underline{\mathbf{B}}(i_1, i_2, \dots, i_N).$$

**Example 4.1.** The contraction of a TT tensor  $\underline{\mathbf{X}} = \underline{\mathbf{X}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{X}}^{(N)}$  with a rank-one tensor  $\mathbf{u}^{(1)} \circ \cdots \circ \mathbf{u}^{(N)}$  can be simplified as

$$\left\langle \underline{\mathbf{X}}, \mathbf{u}^{(1)} \circ \cdots \circ \mathbf{u}^{(N)} \right\rangle = \left( \underline{\mathbf{X}}^{(1)} \times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(N)} \right) \overline{\times}_{1} \mathbf{u}^{(1)} \cdots \overline{\times}_{N} \mathbf{u}^{(N)}$$

$$= \left( \underline{\mathbf{X}}^{(1)} \overline{\times}_{2} \mathbf{u}^{(1)} \right) \left( \underline{\mathbf{X}}^{(2)} \overline{\times}_{2} \mathbf{u}^{(2)} \right) \cdots \left( \underline{\mathbf{X}}^{(N)} \overline{\times}_{2} \mathbf{u}^{(N)} \right).$$

$$(33)$$

The full contraction of two TT tensors  $\underline{\mathbf{X}} = \underline{\mathbf{X}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{X}}^{(N)}$  and  $\underline{\mathbf{Y}} = \underline{\mathbf{Y}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{Y}}^{(N)}$  can be calculated by combining the Hadamard product and the contraction with the rank-one tensor  $\mathbf{1}_{I_1} \circ \cdots \circ \mathbf{1}_{I_N}$  as

$$\langle \underline{\mathbf{X}}, \underline{\mathbf{Y}} \rangle = \langle \underline{\mathbf{X}} \circledast \underline{\mathbf{Y}}, \mathbf{1}_{I_1} \circ \cdots \circ \mathbf{1}_{I_N} \rangle = \mathbf{Z}_1 \cdots \mathbf{Z}_N,$$

where

$$\mathbf{Z}_{n} = (\mathbf{\underline{X}}^{(n)} \boxtimes \mathbf{\underline{Y}}^{(n)}) \overline{\times}_{2} \mathbf{1}_{I_{n}}$$

$$= \sum_{i_{n}=1}^{I_{n}} \mathbf{X}_{i_{n}}^{(n)} \otimes \mathbf{Y}_{i_{n}}^{(n)} \in \mathbb{R}^{R_{n-1}^{X}R_{n-1}^{Y} \times R_{n}^{X}R_{n}^{Y}}, \quad n = 1, \dots, N.$$
(34)

The computational cost for calculating the contraction  $\langle \underline{\mathbf{X}}, \underline{\mathbf{Y}} \rangle$  of two TT tensors can be reduced to  $\mathcal{O}(NIR^3)$  where  $I = \max(I_n)$  and  $R = \max(\{R_n^X\}, \{R_n^Y\})$ , which is linear with N [33].

## 4.6 Matrix-by-vector product

2

The matrix-by-vector product can also be efficiently represented by vector TT and matrix TT decompositions. Let  $\mathbf{x} \in \mathbb{R}^{J_1 J_2 \cdots J_N}$  and  $\mathbf{A} \in \mathbb{R}^{I_1 I_2 \cdots I_N \times J_1 J_2 \cdots J_N}$  be a vector and a matrix represented by vector TT (25) and matrix TT (28) decompositions, i.e.,

$$\mathbf{x} = \widetilde{\mathbf{X}}^{(1)} |\otimes| \cdots |\otimes| \widetilde{\mathbf{X}}^{(N)},$$
$$\mathbf{A} = \widetilde{\mathbf{A}}^{(1)} |\otimes| \cdots |\otimes| \widetilde{\mathbf{A}}^{(N)},$$

with TT-ranks  $\{R_n^X\}$  and  $\{R_n^A\}$ , respectively. The matrix-by-vector product can be represented by vector TT decomposition as

$$\mathbf{A}\mathbf{x} = \mathbf{A} \times^{1} \mathbf{x} = \left(\widetilde{\mathbf{A}}^{(1)} \times^{1} \widetilde{\mathbf{X}}^{(1)}\right) |\otimes| \cdots |\otimes| \left(\widetilde{\mathbf{A}}^{(N)} \times^{1} \widetilde{\mathbf{X}}^{(N)}\right).$$
(35)

That is, the *n*th block matrix of the vector TT decomposition (35) of the product  $\mathbf{A}\mathbf{x}$  is expressed by contracted product of the corresponding block matrices. It can be re-written as

$$\widetilde{\mathbf{A}}^{(n)} \times^{1} \widetilde{\mathbf{X}}^{(n)} = \begin{bmatrix} \mathbf{z}_{s_{n-1}, s_{n}}^{(n)} \end{bmatrix} \in \mathbb{R}^{R_{n-1}^{A}R_{n-1}I_{n} \times R_{n}^{A}R_{n}},$$

where each block is a vector

$$\mathbf{z}_{s_{n-1},s_n}^{(n)} = \mathbf{A}_{r_{n-1}^A,r_n^A}^{(n)} \mathbf{x}_{r_{n-1},r_n}^{(n)} \in \mathbb{R}^{I_n}, \quad s_{n-1} = \overline{r_{n-1}^A r_{n-1}}, \ s_n = \overline{r_n^A r_n}, \tag{36}$$

for all  $s_{n-1} = 1, \dots, R_{n-1}^A R_{n-1}, s_n = 1, \dots, R_n^A R_n, n = 1, \dots, N.$ 

In TT decomposition, the vector (36) is considered as a fiber of a 3rd-order TT-core  $\underline{\mathbf{Z}}^{(n)} \in \mathbb{R}^{R_{n-1}^{A}R_{n-1} \times I_n \times R_n^{A}R_n}$ , i.e.,  $\mathbf{z}_{s_{n-1},s_n}^{(n)} = \underline{\mathbf{Z}}^{(n)}(s_{n-1}, :, s_n)$ , e.g., see Section 3.3. Slice matrices of the TT-core  $\underline{\mathbf{Z}}^{(n)}$  can be written as

$$\mathbf{Z}_{i_{n}}^{(n)} = \underline{\mathbf{Z}}^{(n)}(:, i_{n}, :) = \sum_{j_{n}=1}^{J_{n}} \mathbf{A}_{i_{n}, j_{n}}^{(n)} \otimes \mathbf{X}_{j_{n}}^{(n)} \in \mathbb{R}^{R_{n-1}^{A}R_{n-1} \times R_{n}^{A}R_{n}},$$
(37)

where  $\mathbf{A}_{i_n,j_n}^{(n)} = \underline{\mathbf{A}}^{(n)}(:, i_n, j_n, :)$  and  $\mathbf{X}_{j_n}^{(n)} = \underline{\mathbf{X}}^{(n)}(:, j_n :)$  are the slice matrices of the *n*th TT-cores of **A** and **x**, respectively. As a result, the entries of the product  $\mathbf{z} = \mathbf{A}\mathbf{x}$  can be written as products of slice matrices of TT-cores

$$\mathbf{z}(\overline{i_1,\ldots,i_N}) = \mathbf{Z}_{i_1}^{(1)} \mathbf{Z}_{i_2}^{(2)} \cdots \mathbf{Z}_{i_N}^{(N)}.$$

Note that  $\mathbf{Z}_{i_1}^{(1)} \in \mathbb{R}^{1 \times R_1^A R_1}$  and  $\mathbf{Z}_{i_N}^{(N)} \in \mathbb{R}^{R_{N-1}^A R_{N-1} \times 1}$  are row and column vectors.

The computational cost for computing a matrix-by-vector product of a matrix  $\mathbf{A} \in \mathbb{R}^{I_1 \cdots I_N \times J_1 \cdots J_N}$ with a vector  $\mathbf{x} \in \mathbb{R}^{J_1 \cdots J_N}$  can be  $\mathcal{O}(NI^2R^4)$  where  $I = \max(\{I_n\}, \{J_n\}), R = \max(\{R_n\}, \{R_n^A\}),$ by using TT decomposition [33].

On the other hand, recall that, in (17) and (18),

$$\mathbf{x} = \operatorname{vec}\left(\underline{\mathbf{X}}\right) = \mathbf{X}^{\neq n} \mathbf{x}^{(n)},$$

where  $\mathbf{X}^{\neq n}$  is the frame matrix and  $\mathbf{x}^{(n)} = \operatorname{vec}(\mathbf{X}^{(n)})$ . A large-scale matrix-by-vector multiplication can be reduced to a smaller matrix-by-vector multiplication as  $\mathbf{A}\mathbf{x} = \mathbf{A}\mathbf{X}^{\neq n}\mathbf{x}^{(n)} \equiv \widetilde{\mathbf{A}}_{n}\mathbf{x}^{(n)}$ , where

$$\widetilde{\mathbf{A}}_{n} = \mathbf{A}\mathbf{X}^{\neq n} \in \mathbb{R}^{I_{1}I_{2}\cdots I_{N} \times R_{n-1}J_{n}R_{n}}.$$
(38)

We often cannot calculate the matrix  $\mathbf{A}_n$  by matrix-by-matrix multiplication (38) for a large matrix  $\mathbf{A}$  due to high storage and computational costs. Instead, by using the distributed representation (35), a matrix-by-vector product,  $\widetilde{\mathbf{A}}_n \mathbf{w}^{(n)}$  for some vector  $\mathbf{w}^{(n)}$ , can be calculated by recursive core contractions as follows.

**Proposition 4.1.** Let a vector  $\mathbf{x} \in \mathbb{R}^{J_1 \cdots J_N}$  and a matrix  $\mathbf{A} \in \mathbb{R}^{I_1 \cdots I_N \times J_1 \cdots J_N}$  be represented by vector TT and matrix TT decompositions with block matrices  $\widetilde{\mathbf{X}}^{(n)}$  and  $\widetilde{\mathbf{A}}^{(n)}$ , respectively. For a fixed  $n = 1, \ldots, N$ , let  $\widetilde{\mathbf{A}}_n$  be the matrix defined by (38). Then, for any vector  $\mathbf{w}^{(n)} \in \mathbb{R}^{R_{n-1}J_nR_n}$ ,

$$\widetilde{\mathbf{A}}_{n}\mathbf{w}^{(n)} = \widetilde{\mathbf{Z}}^{(1)} \ket{\otimes} \cdots \ket{\otimes} \widetilde{\mathbf{Z}}^{(N)} \in \mathbb{R}^{I_{1}I_{2}\cdots I_{N}}$$

with block matrices

$$\widetilde{\mathbf{Z}}^{(m)} = \widetilde{\mathbf{A}}^{(m)} \times^{1} \widetilde{\mathbf{X}}^{(m)} \in \mathbb{R}^{R_{m-1}^{A}R_{m-1}I_{m} \times R_{m}^{A}R_{m}}, \quad m = 1, \dots, n-1, n+1, \dots, N,$$
$$\widetilde{\mathbf{Z}}^{(n)} = \widetilde{\mathbf{A}}^{(n)} \times^{1} \widetilde{\mathbf{W}}^{(n)} \in \mathbb{R}^{R_{n-1}^{A}R_{n-1}I_{n} \times R_{n}^{A}R_{n}},$$

where  $\underline{\mathbf{W}}^{(n)} \in \mathbb{R}^{R_{n-1} \times J_n \times R_n}$  is the 3rd-order tensor such that  $\mathbf{w}^{(n)} = vec(\underline{\mathbf{W}}^{(n)})$ , and  $\widetilde{\mathbf{W}}^{(n)} = [\underline{\mathbf{W}}^{(n)}(r_{n-1},:,r_n)] \in \mathbb{R}^{R_{n-1}J_n \times R_n}$  is the block matrix partitioned with the fiber vectors  $\underline{\mathbf{W}}^{(n)}(r_{n-1},:,r_n) \in \mathbb{R}^{J_n}$ .



Figure 9: Tensor network diagram for the matrix-by-vector product  $\widetilde{\mathbf{A}}_n \mathbf{x}^{(n)}$  represented by the TT decomposition (35), where  $\mathbf{x}^{(n)} = \operatorname{vec}(\underline{\mathbf{X}}^{(n)})$ .

*Proof.* Let  $\underline{\mathbf{X}}^{(n)} \in \mathbb{R}^{R_{n-1} \times J_n \times R_n}$  denote the TT-cores corresponding to the block matrices  $\widetilde{\mathbf{X}}^{(n)} \in \mathbb{R}^{R_{n-1}J_n \times R_n}$ . Let  $\underline{\mathbf{W}}$  be an *N*th-order tensor defined by  $\underline{\mathbf{W}} = \underline{\mathbf{X}}^{(1)} \times^1 \cdots \times^1 \underline{\mathbf{X}}^{(n-1)} \times^1 \underline{\mathbf{W}}^{(n)} \times^1 \underbrace{\mathbf{X}}^{(n+1)} \times^1 \cdots \times^1 \underline{\mathbf{X}}^{(n)}$ , then

$$\mathbf{w} \equiv \operatorname{vec}\left(\underline{\mathbf{W}}\right) = \mathbf{X}^{\neq n} \mathbf{w}^{(n)}.$$

As a result, we have

$$\widetilde{\mathbf{A}}_{n}\mathbf{w}^{(n)} = \mathbf{A}\mathbf{X}^{\neq n}\mathbf{w}^{(n)} = \mathbf{A}\mathbf{w}.$$

Note that  $\mathbf{w} = \operatorname{vec}(\underline{\mathbf{W}})$  can be represented by vector TT decomposition with block matrices  $\widetilde{\mathbf{X}}^{(1)}, \ldots, \widetilde{\mathbf{W}}^{(n)}, \ldots, \widetilde{\mathbf{X}}^{(N)}$ . The result follows from the expression (35).

Figure 9 illustrates the tensor network diagram for the product  $\widetilde{\mathbf{A}}_{n}\mathbf{x}^{(n)}$ . For each  $k = 1, \ldots, n - 1, n+1, \ldots, N$ , the node for the TT-core  $\underline{\mathbf{A}}^{(k)}$  is connected to the node for  $\underline{\mathbf{X}}^{(k)}$ , which is represented as  $\widetilde{\mathbf{A}}^{(k)} \times^{1} \widetilde{\mathbf{X}}^{(k)}$  in (35).

## 4.7 Quadratic form

The quadratic form  $\mathbf{x}^{\top} \mathbf{A} \mathbf{x}$  for a symmetric and very large-scale matrix  $\mathbf{A} \in \mathbb{R}^{I_1 I_2 \cdots I_N \times I_1 I_2 \cdots I_N}$  can be represented by TT decomposition as follows. Let  $\mathbf{x} \in \mathbb{R}^{I_1 I_2 \cdots I_N}$  and  $\mathbf{A} \in \mathbb{R}^{I_1 I_2 \cdots I_N \times I_1 I_2 \cdots I_N}$  be a vector and a matrix represented by vector TT (25) and matrix TT (28) decompositions, i.e.,

$$\mathbf{x} = \widetilde{\mathbf{X}}^{(1)} |\otimes| \cdots |\otimes| \widetilde{\mathbf{X}}^{(N)},$$
$$\mathbf{A} = \widetilde{\mathbf{A}}^{(1)} |\otimes| \cdots |\otimes| \widetilde{\mathbf{A}}^{(N)},$$

with TT-ranks  $\{R_n^X\}$  and  $\{R_n^A\}$ , respectively. The quadratic form can be represented as strong Kronecker products of block matrices

$$\mathbf{x}^{\top} \mathbf{A} \mathbf{x} = \mathbf{x} \times^{1} \mathbf{A} \times^{1} \mathbf{x} = \left( \widetilde{\mathbf{X}}^{(1)} \times^{1} \widetilde{\mathbf{A}}^{(1)} \times^{1} \widetilde{\mathbf{X}}^{(1)} \right) |\otimes| \cdots |\otimes| \left( \widetilde{\mathbf{X}}^{(N)} \times^{1} \widetilde{\mathbf{A}}^{(N)} \times^{1} \widetilde{\mathbf{X}}^{(N)} \right).$$
(39)

That is, the *n*th block matrix in (39) is expressed by contracted product of the corresponding block matrices. It can be re-written as

$$\widetilde{\mathbf{X}}^{(n)} \times^{1} \widetilde{\mathbf{A}}^{(n)} \times^{1} \widetilde{\mathbf{X}}^{(n)} = \left[ z_{s_{n-1},s_{n}}^{(n)} \right] \in \mathbb{R}^{R_{n-1}^{A}R_{n-1}^{2} \times R_{n}^{A}R_{n}^{2}},$$

where each block is a scalar

$$z_{s_{n-1},s_n}^{(n)} = \mathbf{x}_{r_{n-1}',r_n'}^{(n)} \mathbf{A}_{r_{n-1},r_n^A}^{(n)} \mathbf{x}_{r_{n-1},r_n}^{(n)} \in \mathbb{R}, \quad s_{n-1} = \overline{r_{n-1}'r_{n-1}^A r_{n-1}}, \ s_n = \overline{r_n'r_n^A r_n}, \tag{40}$$

for all  $s_{n-1} = 1, \dots, R_{n-1}^A R_{n-1}^2$ ,  $s_n = 1, \dots, R_n^A R_n^2$ ,  $n = 1, \dots, N$ .

Since each block of the block matrices in (39) is a scalar, the quadratic form can be re-written as product of matrices as

$$\mathbf{x}^{\top} \mathbf{A} \mathbf{x} = \mathbf{Z}^{(1)} \cdots \mathbf{Z}^{(N)},$$

where

$$\mathbf{Z}^{(n)} = \left[ z_{s_{n-1},s_n}^{(n)} \right] = \sum_{i_n=1}^{I_n} \sum_{j_n=1}^{I_n} \mathbf{X}_{i_n}^{(n)} \otimes \mathbf{A}_{i_n,j_n}^{(n)} \otimes \mathbf{X}_{j_n}^{(n)}, \quad n = 1, \dots, N,$$

where  $\mathbf{X}_{i_n}^{(n)} = \underline{\mathbf{X}}^{(n)}(:, i_n, :)$  and  $\mathbf{A}_{i_n, j_n}^{(n)} = \underline{\mathbf{A}}^{(n)}(:, i_n, j_n, :)$  are slice matrices of TT-cores, and  $\mathbf{Z}^{(1)} \in \mathbb{R}^{1 \times R_1^A R_1^2}$  and  $\mathbf{Z}^{(N)} \in \mathbb{R}^{R_{N-1}^A R_{N-1}^2 \times 1}$  are row and column vectors, respectively.

On the other hand, recall that

$$\mathbf{x} = \operatorname{vec}\left(\underline{\mathbf{X}}\right) = \mathbf{X}^{\neq n} \mathbf{x}^{(n)},$$

with  $\mathbf{x}^{(n)} = \operatorname{vec}(\underline{\mathbf{X}}^{(n)})$ , so the quadratic form  $\mathbf{x}^{\top} \mathbf{A} \mathbf{x}$  reduces to

$$\mathbf{x}^{\top} \mathbf{A} \mathbf{x} = \mathbf{x}^{(n)\top} (\mathbf{X}^{\neq n})^{\top} \mathbf{A} \mathbf{X}^{\neq n} \mathbf{x}^{(n)} \equiv \mathbf{x}^{(n)\top} \overline{\mathbf{A}}_n \mathbf{x}^{(n)},$$

where the matrix

$$\overline{\mathbf{A}}_{n} = (\mathbf{X}^{\neq n})^{\top} \mathbf{A} \mathbf{X}^{\neq n} \in \mathbb{R}^{R_{n-1}I_{n}R_{n} \times R_{n-1}I_{n}R_{n}}$$
(41)

is a much smaller matrix than  $\mathbf{A}$  when TT-ranks  $R_{n-1}$  and  $R_n$  are sufficiently small. Since  $\overline{\mathbf{A}}_n$  often cannot be calculated by matrix-by-matrix multiplication for a large matrix  $\mathbf{A}$ , we calculate it iteratively by recursive core contractions based on the distributed representation (39) as follows.

**Proposition 4.2.** Let a vector  $\mathbf{x} \in \mathbb{R}^{I_1 \cdots I_N}$  and a matrix  $\mathbf{A} \in \mathbb{R}^{I_1 \cdots I_N \times I_1 \cdots I_N}$  be represented by vector TT and matrix TT decompositions with block matrices  $\widetilde{\mathbf{X}}^{(n)}$  and  $\widetilde{\mathbf{A}}^{(n)}$ , respectively. For a fixed  $n = 1, \ldots, N$ , let  $\overline{\mathbf{A}}_n$  be the matrix defined by (41). Then, for any vector  $\mathbf{y}^{(n)}, \mathbf{w}^{(n)} \in \mathbb{R}^{R_{n-1}I_nR_n}$ ,

$$\mathbf{y}^{(n)\top}\overline{\mathbf{A}}_{n}\mathbf{w}^{(n)} = \widetilde{\mathbf{Z}}^{(1)} |\otimes| \cdots |\otimes| \widetilde{\mathbf{Z}}^{(N)} \in \mathbb{R}^{R_{n-1}I_{n}R_{n}}$$

with block matrices

$$\begin{split} \widetilde{\mathbf{Z}}^{(m)} &= \widetilde{\mathbf{X}}^{(m)} \times^{1} \widetilde{\mathbf{A}}^{(m)} \times^{1} \widetilde{\mathbf{X}}^{(m)} \in \mathbb{R}^{R_{m-1}^{A} R_{m-1}^{2} \times R_{m}^{A} R_{m}^{2}}, \quad m = 1, \dots, n-1, n+1, \dots, N, \\ \widetilde{\mathbf{Z}}^{(n)} &= \widetilde{\mathbf{Y}}^{(n)} \times^{1} \widetilde{\mathbf{A}}^{(n)} \times^{1} \widetilde{\mathbf{W}}^{(n)} \in \mathbb{R}^{R_{n-1}^{A} R_{n-1}^{2} I_{n} \times R_{n}^{A} R_{n}^{2}}, \end{split}$$

where  $\underline{\mathbf{Y}}^{(n)}, \underline{\mathbf{W}}^{(n)} \in \mathbb{R}^{R_{n-1} \times I_n \times R_n}$  are the 3rd-order tensors such that  $\mathbf{y}^{(n)} = vec(\underline{\mathbf{Y}}^{(n)}), \mathbf{w}^{(n)} = vec(\underline{\mathbf{W}}^{(n)}),$  and  $\widetilde{\mathbf{Y}}^{(n)} = [\underline{\mathbf{Y}}^{(n)}(r_{n-1}, :, r_n)], \quad \widetilde{\mathbf{W}}^{(n)} = [\underline{\mathbf{W}}^{(n)}(r_{n-1}, :, r_n)] \in \mathbb{R}^{R_{n-1}I_n \times R_n}$  are the block matrices partitioned with the fiber vectors  $\underline{\mathbf{Y}}^{(n)}(r_{n-1}, :, r_n), \quad \underline{\mathbf{W}}^{(n)}(r_{n-1}, :, r_n) \in \mathbb{R}^{I_n}$ .

*Proof.* Let  $\underline{\mathbf{X}}^{(n)} \in \mathbb{R}^{R_{n-1} \times I_n \times R_n}$  denote the TT-cores corresponding to the block matrices  $\widetilde{\mathbf{X}}^{(n)} \in \mathbb{R}^{R_{n-1}I_n \times R_n}$ . Let  $\underline{\mathbf{Y}}, \underline{\mathbf{W}}$  be Nth-order tensors defined by

$$\underline{\mathbf{Y}} = \underline{\mathbf{X}}^{(1)} \times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(n-1)} \times^{1} \underline{\mathbf{Y}}^{(n)} \times^{1} \underline{\mathbf{X}}^{(n+1)} \times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(N)},$$
  
$$\underline{\mathbf{W}} = \underline{\mathbf{X}}^{(1)} \times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(n-1)} \times^{1} \underline{\mathbf{W}}^{(n)} \times^{1} \underline{\mathbf{X}}^{(n+1)} \times^{1} \cdots \times^{1} \underline{\mathbf{X}}^{(N)},$$

then

$$\mathbf{y} \equiv \operatorname{vec}(\underline{\mathbf{Y}}) = \mathbf{X}^{\neq n} \mathbf{y}^{(n)}, \quad \mathbf{w} \equiv \operatorname{vec}(\underline{\mathbf{W}}) = \mathbf{X}^{\neq n} \mathbf{w}^{(n)}.$$

As a result, we have

$$\mathbf{y}^{(n)\top}\overline{\mathbf{A}}_{n}\mathbf{w}^{(n)} = \mathbf{y}^{(n)\top}(\mathbf{X}^{\neq n})^{\top}\mathbf{A}\mathbf{X}^{\neq n}\mathbf{w}^{(n)} = \mathbf{y}^{\top}\mathbf{A}\mathbf{w}$$

Since  $\mathbf{y} = \operatorname{vec}(\underline{\mathbf{Y}})$  (resp.  $\mathbf{w} = \operatorname{vec}(\underline{\mathbf{W}})$ ) can be represented by vector TT decomposition with block matrices  $\widetilde{\mathbf{X}}^{(1)}, \ldots, \widetilde{\mathbf{Y}}^{(n)}, \ldots, \widetilde{\mathbf{X}}^{(N)}$  (resp.  $\widetilde{\mathbf{X}}^{(1)}, \ldots, \widetilde{\mathbf{W}}^{(n)}, \ldots, \widetilde{\mathbf{X}}^{(N)}$ ), the result follows from the expression (39).



Figure 10: Tensor network diagram for the quadratic form  $\mathbf{x}^{(n)\top}\overline{\mathbf{A}}_n\mathbf{x}^{(n)}$  represented by the TT decomposition (39), where  $\mathbf{x}^{(n)} = \operatorname{vec}(\underline{\mathbf{X}}^{(n)})$ .

Figure 10 illustrates the tensor network diagram for the quadratic form  $\mathbf{x}^{(n)\top}\overline{\mathbf{A}}_n\mathbf{x}^{(n)}$ . It is clear that each node for core tensor  $\underline{\mathbf{A}}^{(k)}, k = 1, \ldots, n-1, n+1, \ldots, N$ , is connected to the node for core tensor  $\underline{\mathbf{X}}^{(k)}$ , which is represented as  $\widetilde{\mathbf{X}}^{(k)} \times^1 \widetilde{\mathbf{A}}^{(k)} \times^1 \widetilde{\mathbf{X}}^{(k)}$  in (39).

## 5 Discussion and conclusions

In this paper, we proposed several extended mathematical operations on tensors and developed their multilinear algebraic properties and their links with tensor network formats, especially TT decompositions. We generalized the standard matrix-based operations such as the Kronecker product, Hadamard product, and direct sum, and proposed tensor-based operations such as the partial trace and contracted product for block tensors. We have shown that the tensor-based operations are able to not only simplify traditional index notation for TT representations but also describe important basic operations which are very useful for computational algorithms using large-scale vectors, matrices, and higher-order tensors.

The partial trace operator can be used for describing the tensor chain (TC) decomposition [12, 26] simply by slightly modifying the suggested TT representations. Properties of TC decomposition should be more investigated in the future work. Moreover, the definitions and properties of partial Kronecker product, partial direct sum, and contracted product can also be generalized to any tensor network decompositions such as hierarchical Tucker (HT) [14, 16, 17] and hybrid formats [15, 26].

The partial contracted products of either the left or right core tensors of TT decomposition are matricized and used as a building block of the frame matrices. We have shown that the suggested tensor operations can be used to prove the orthonormality of the frame matrices, which have been proved only by using index notation in the literature. The developed relationships also play a key role in the alternating linear scheme (ALS) and modified alternating linear scheme (MALS) algorithms [20] for reducing the large-scale optimizations to iterative smaller scale problems. Recent studies adjust the frame matrices in order to incorporate rank adaptivity and improve convergence for the ALS [10, 28]. In this work, we have derived explicit representations of the localized linear maps  $\underline{\tilde{A}}_n$ and  $\underline{\tilde{A}}_n$  by the proposed tensor operations, which are important for TT-based iterative methods for breaking the curse-of-dimensionality [2], while the global convergence of the methods remains as a future work. In addition, it is important to keep the TT-ranks moderate for a feasible computational cost, which is a crucial issue for real world applications of TT decompositions, see, e.g., [39].

## References

- [1] Bungatrz, H.-J., & Griebel, M. (2004). Sparse grids. Acta Numer., 13, 147–269.
- [2] Cichocki, A. (2014a). Era of big data processing: A new approach via tensor networks and tensor decompositions. arXiv:1403.2048.
- [3] Cichocki, A. (2014b). Tensor networks for big data analytics and large-scale optimization problems. arXiv:1407.3124.
- [4] Cichocki, A., Zdunek, R., Phan, A. H., & Amari, S. (2009). Nonnegative Matrix and Tensor Factorizations: Applications to Exploratory Multi-way Data Analysis and Blind Source Separation. Chichester: Wiley.
- [5] Cirac, J.I., Michalakis, S., Pérez-García, D., & Schuch, N. (2013). Robustness in projected entangled pair states. *Phys. Rev. B*, 88, 115108.
- [6] De Lathauwer, L. (2009). A survey of tensor methods. In 2009 IEEE International Symposium on Circuits and Systems (ISCAS 2009) (pp. 2773–2776).
- [7] De Lathauwer, L., De Moor, B., & Vandewalle, J. (2000). A multilinear singular value decomposition. SIAM J. Matrix Anal. Appl., 21, 1253–1278.
- [8] De Silva, V., & Lim, L.-H. (2008). Tensor rank and the ill-posedness of the best low-rank approximation problem. *SIAM J. Matrix Anal. Appl.*, 30(3), 1084–1127.
- [9] Dolgov, S.V., Khoromskij, B.N., Oseledets, I.V., & Savostyanov, D.V. (2014). Computation of extreme eigenvalues in higher dimensions using block tensor train format. *Comput. Phys. Commun.*, 185(4), 1207–1216.
- [10] Dolgov, S.V., & Savostyanov, D.V. (2014). Alternating minimal energy methods for linear systems in higher dimensions. SIAM J. Sci. Comput., 36, A2248–A2271.
- [11] Espig, M., Hackbusch, W., Handschuh, S., & Schneider, R. (2011). Optimization problems in contracted tensor networks. *Comput. Vis. Sci.*, 14, 271–285.
- [12] Espig, M., Naraparaju, K.K., & Schneider, J. (2012). A note on tensor chain approximation. Comput. Vis. Sci., 15, 331–344.
- [13] Falcó, A., & Hackbusch, W. (2012). On minimal subspaces in tensor representations. Found. Comput. Math., 12, 765–803.
- [14] Grasedyck, L. (2010). Hierarchical singular value decomposition of tensors. SIAM J. Matrix Anal. Appl., 31(4), 2029–2054.
- [15] Grasedyck, L., Kressner, D., & Tobler, C. (2013). A literature survey of low-rank tensor approximation techniques. GAMM-Mitt., 36(1), 53–78.
- [16] Hackbusch, W. (2012). Tensor Spaces and Numerical Tensor Calculus. Berlin: Springer.
- [17] Hackbusch, W., & Kühn, S. (2009). A new scheme for the tensor representation. J. Fourier Anal. Appl., 15(5), 706–722.
- [18] Håstad, J. (1990). Tensor rank is NP-complete. J. Algorithms, 11(4), 644–654.
- [19] Holtz, S., Rohwedder, T., & Schneider, R. (2012a). On manifolds of tensors of fixed TT-rank. Numer. Math., 120(4), 701–731.

- [20] Holtz, S., Rohwedder, T., & Schneider, R. (2012b). The alternating linear scheme for tensor optimization in the tensor train format. SIAM J. Sci. Comput. 34(2), A683–A713.
- [21] Kazeev, V.A., & Khoromskij, B.N. (2012). Low-rank explicit QTT representation of the Laplace operator and its inverse. SIAM J. Matrix Anal. Appl., 33(3), 742–758.
- [22] Kazeev, V.A., Khoromskij, B.N., & Tyrtyshnikov, E.E. (2013). Multilevel Toeplitz matrices generated by tensor-structured vectors and convolution with logarithmic complexity. SIAM J. Sci. Comput., 35(3), A1511–A1536.
- [23] Kolda, T.G. (2006). Multilinear operators for higher-order decompositions. Technical Report SAND2006-2081, Sandia National Laboratories.
- [24] Kolda, T.G., & Bader, B.W. (2009). Tensor decompositions and applications. SIAM Rev., 51(3), 455–500.
- [25] Khoromskij, B.N. (2011). O(d log N)-quantics approximation of N-d tensors in high-dimensional numerical modeling. Constr. Approx., 34, 257–280.
- [26] Khoromskij, B.N. (2012). Tensors-structured numerical methods in scientific computing: Survey on recent advances. *Chemometr. Intell. Lab. Syst.*, 110(1), 1–19.
- [27] Khoromskij, B.N., & Oseledets, I.V. (2010). DMRG+QTT approach to computation of the ground state for the molecular Schrödinger operator. Preprint 69, MPI MiS, Leipzig.
- [28] Kressner, D., Steinlechner, M., & Uschmajew, A. (2014). Low-rank tensor methods with subspace correction for symmetric eigenvalue problems. SIAM J. Sci. Comput., 36, A2346–A2368.
- [29] Launey, W.D., & Seberry, J. (1994). The strong Kronecker product. J. Combin. Theory Ser. A, 66(2), 192–213.
- [30] Lee, N., & Cichocki, A. (2014). Big data matrix singular value decomposition based on low-rank tensor train decomposition. In Z. Zeng, Y. Li, I. King (Eds.), Advances in Neural Networks -ISNN 2014, LNCS 8866 (pp. 121–130). Springer.
- [31] Lee, N., & Cichocki, A. (2015). Estimating a few extreme singular values and vectors for largescale matrices in tensor train format. SIAM J. Matrix Anal. Appl., 36(3), 994–1014.
- [32] Oseledets, I.V. (2010). Approximation of  $2^d \times 2^d$  matrices using tensor decomposition. SIAM J. Matrix Anal. Appl., 31(4), 2130–2145.
- [33] Oseledets, I.V. (2011). Tensor-train decomposition. SIAM J. Sci. Comput., 33(5), 2295–2317.
- [34] Oseledets, I.V., & Dolgov, S.V. (2012). Solution of linear systems and matrix inversion in the TT-format. SIAM J. Sci. Comput., 34, A2718–A2739.
- [35] Oseledets, I.V., & Tyrtyshnikov, E.E. (2009). Breaking the curse of dimensionality, or how to use SVD in many dimensions. SIAM J. Sci. Comput., 31(5), 3744–3759.
- [36] Schollwöck, U. (2011). The density-matrix renormalization group in the age of matrix product states. Ann. Physics, 326(1), 96–192.
- [37] Smolyak, S.A. (1963). Quadrature and interpolation formulas for tensor products of certain classes of functions. Soviet Math. Dokl., 4, 240–243.
- [38] Tucker, L.R. (1966). Some mathematical notes on three-mode factor analysis. Psychometrika, 31(3), 279–311.

- [39] Vervliet, N., Debals, O., Sorber, L., & De Lathauwer, L. (2014). Breaking the curse of dimensionality using decompositions of incomplete tensors: Tensor-based scientific computing in big data analysis. *IEEE Signal Process. Mag.*, 31(5), 71–79.
- [40] White, S.R. (1993). Density-matrix algorithms for quantum renormalization groups. Phys. Rev. B., 48(14), 10345–10356.