

Notes on branched coverings of Seifert manifolds *

Hong Huang

Abstract

In a paper published in 2002, the author gave a criterion to determine whether there is a fiber-preserving branched covering with given degree between two given connected, closed, orientable Seifert manifolds with orientable bases. Here we supply some details of the proof of two claims in that paper. We give an explicit construction of fiber-preserving branched covering between two Seifert fibered solid tori when their Seifert invariants satisfy certain relation, and we show the factorability of fiber-preserving branched coverings between two closed Seifert manifolds.

Key words: Seifert manifolds, fiber-preserving branched coverings, factorization of branched coverings

Mathematics Subject Classification 2010: 57N10

In a paper [2] published in 2002, the author gave a criterion to determine whether there is a fiber-preserving branched covering with given degree between two given connected, closed, orientable Seifert manifolds with orientable bases. There are two statements used in [2] without detailed proof: One is on the existence of fiber-preserving branched covering between two Seifert fibered solid tori when their Seifert invariants satisfy certain relation, another is on the factorability of fiber-preserving branched coverings between two closed Seifert manifolds. Here we'll supply some details of the proof of these two statements.

Let V be a Seifert fibered solid torus. Let (q, h) be a (cross section, fiber) basis of $H_1(\partial V)$. We will abuse notation to use $q, h, etc.$ to denote both a closed curve and the homology class that it represents. If a meridian m of V is homologous to $\alpha q + \beta h$ in $H_1(\partial V)$, where $\alpha > 0, \beta$ are coprime integers, then we say that the fibered solid torus V has Seifert invariant (α, β) (w.r.t. (q, h)); see [5].

The following result is contained in [2], but the details of the proof of sufficiency are omitted there.

Proposition 0.1. *Let V_1, V_2 be two Seifert fibered solid tori. Suppose the Seifert invariant of V_2 w.r.t. some (cross section, fiber) basis of $H_1(\partial V_2)$ is (α_2, β_2) , and k is a positive integer. Then there is a fiber-preserving k -fold branched covering*

*Research supported by NSFC (No. 11171025) and by Laboratory of Mathematics and Complex Systems, Ministry of Education.

$f : V_1 \rightarrow V_2$ with fiber degree one and with upstairs and downstairs branched set the central fiber of V_1 and V_2 respectively if and only if there is some (cross section, fiber) basis of $H_1(\partial V_1)$ such that the Seifert invariant (α_1, β_1) of V_1 w.r.t. this basis satisfies $\beta_1/\alpha_1 = k\beta_2/\alpha_2$.

Proof First we show the necessity as in [2]. Let $f : V_1 \rightarrow V_2$ be a fiber-preserving k -fold branched covering with fiber degree one and with upstairs and downstairs branched set the central fiber of V_1 and V_2 respectively. (By upstairs branched set we mean the preimage of the downstairs branched set under f .) Let (q_2, h_2) be the (cross section, fiber) basis of $H_1(\partial V_2)$ w.r.t. which the Seifert invariant of V_2 is (α_2, β_2) . Let $q_1 = f^{-1}(q_2)$. Since ∂V_1 is the union of fibers through all points of q_1 and the fiber degree of f is one, q_1 must be connected as ∂V_1 is. Then we see that q_1 is a cross section of the Seifert fibration of ∂V_1 , and $f|_*(q_1) = kq_2$, where $f|_*$ is the restriction of f to ∂V_1 .

Let the Seifert invariant of V_1 w.r.t. the (cross section, fiber) basis (q_1, h_1) be (α_1, β_1) . By assumption we have $f|_*(h_1) = h_2$. By definition of branched covering, there is a nonzero integer s such that $f|_*(m_1) = sm_2$, where m_1 (resp. m_2) is a meridian of V_1 (resp. V_2). So $f|_*(\alpha_1 q_1 + \beta_1 h_1) = s(\alpha_2 q_2 + \beta_2 h_2)$, and $\alpha_1 k q_2 + \beta_1 h_2 = s\alpha_2 q_2 + s\beta_2 h_2$, which implies that $\alpha_1 k = s\alpha_2$, and $\beta_1 = s\beta_2$. It follows that $\beta_1/\alpha_1 = k\beta_2/\alpha_2$.

Then we show the sufficiency. Suppose there is some (cross section, fiber) basis (q_1, h_1) of $H_1(\partial V_1)$ such that the Seifert invariant (α_1, β_1) of V_1 w.r.t. this basis satisfies $\beta_1/\alpha_1 = k\beta_2/\alpha_2$. If $\beta_2 = 0$, then $\alpha_2 = 1$, $\beta_1 = 0$, and $\alpha_1 = 1$. In this case, the result is clear. Below we assume that $\beta_2 \neq 0$. From the relation $\beta_1/\alpha_1 = k\beta_2/\alpha_2$, using that α_j and β_j are coprime, we get that $\alpha_1|\alpha_2$, and $\beta_2|\beta_1$.

Now it is clear that there exists a fiber-preserving k -fold branched covering $f : V_1 \rightarrow V_2$ with fiber degree one and with upstairs and downstairs branched set the central fiber of V_1 and V_2 respectively. The reason is as follows. First we can construct a fiber preserving covering $f| : \partial V_1 \rightarrow \partial V_2$ such that $f|_*(q_1) = kq_2$, and $f|_*(h_1) = h_2$. We have, using the relation $\beta_1/\alpha_1 = k\beta_2/\alpha_2$,

$$f|_*(m_1) = f|_*(\alpha_1 q_1 + \beta_1 h_1) = \alpha_1 k q_2 + \beta_1 h_2 = \frac{\beta_1}{\beta_2}(\alpha_2 q_2 + \beta_2 h_2) = \frac{\beta_1}{\beta_2} m_2.$$

Then by shrinking $\partial V_1, \partial V_2$ and $f|$ respectively we see that $f|$ extends to a fiber-preserving branched covering $f : V_1 \rightarrow V_2$ as desired. (Compare the hint to Exercise 3 on p. 312 of [4].)

Below we give a more explicit construction. Let α'_j, β'_j be integers such that $\alpha_j \beta'_j - \alpha'_j \beta_j = 1$, $j = 1, 2$; note that α'_j (resp. β'_j) is determined by α_j and β_j up to multiples of α_j (resp. β_j). Now we have $\alpha_1 \beta'_1 - (\frac{\beta_1}{\beta_2} \alpha'_1) \beta_2 = 1$, and $\alpha_1 (\frac{\alpha_2}{\alpha_1} \beta'_2) - \alpha'_2 \beta_2 = 1$. (As noted above, both $\frac{\alpha_2}{\alpha_1}$ and $\frac{\beta_1}{\beta_2}$ are integers.) So $\alpha_1 (\beta'_1 - \frac{\alpha_2}{\alpha_1} \beta'_2) = (\frac{\beta_1}{\beta_2} \alpha'_1 - \alpha'_2) \beta_2$. We see that $\alpha'_2 - \frac{\beta_1}{\beta_2} \alpha'_1$ is a multiple of α_1 , since α_1 and β_2 are coprime.

For $j = 1, 2$, consider the universal covering $p_j : D^2 \times \mathbb{R} \rightarrow V_j$ with deck transformation group generated by the element

$$\begin{aligned}\tau_j &: D^2 \times \mathbb{R} \rightarrow D^2 \times \mathbb{R}, \\ (re^{2\pi i\theta}, t) &\mapsto (re^{2\pi i(\theta - \frac{\alpha'_j}{\alpha_j})}, t + 1).\end{aligned}$$

Here we think of V_j as the quotient of $D^2 \times \mathbb{R}$ by the action of the corresponding deck transformation group; cf. [5].

First we construct a map $\tilde{f} : D^2 \times \mathbb{R} \rightarrow D^2 \times \mathbb{R}$ as follows,

$$\tilde{f}(re^{2\pi i\theta}, t) = (re^{2\pi i\frac{\beta_1}{\beta_2}\theta}, \frac{\alpha_2}{\alpha_1}t).$$

Note that

$$\begin{aligned}\tilde{f}(\tau_1(re^{2\pi i\theta}, t)) &= \tilde{f}(re^{2\pi i(\theta - \frac{\alpha'_1}{\alpha_1})}, t + 1) \\ &= (re^{2\pi i\frac{\beta_1}{\beta_2}(\theta - \frac{\alpha'_1}{\alpha_1})}, \frac{\alpha_2}{\alpha_1}(t + 1)) \\ &= (re^{2\pi i(\frac{\beta_1}{\beta_2}\theta - \frac{\alpha_2}{\alpha_1}\frac{\alpha'_1}{\alpha_2})}, \frac{\alpha_2}{\alpha_1}t + \frac{\alpha_2}{\alpha_1}) \\ &= \tau_2^{\alpha_2/\alpha_1}(re^{2\pi i\frac{\beta_1}{\beta_2}\theta}, \frac{\alpha_2}{\alpha_1}t) \\ &= \tau_2^{\alpha_2/\alpha_1}(\tilde{f}(re^{2\pi i\theta}, t)).\end{aligned}$$

Here in the third equality we have used the fact $\alpha_1|(\alpha'_2 - \frac{\beta_1}{\beta_2}\alpha'_1)$.

It follows that \tilde{f} descends to a map $f : V_1 \rightarrow V_2$. One can easily check that $f : V_1 \rightarrow V_2$ is the desired branched covering. \square

The following result was stated in the proof of Theorem 2.2 in [2] without proof.

Proposition 0.2. *Let $f : M_1 \rightarrow M_2$ be a fiber-preserving d -fold branched covering between two (connected) closed Seifert manifolds (both orientable and with orientable base). Then there exists a positive integer d_1 with $d_1|d$ such that $f = f_2 \circ f_1$, where $f_1 : M_1 \rightarrow M_1/\mathbb{Z}_{d_1}$ (here the \mathbb{Z}_{d_1} -action on M_1 is inside the S^1 -action on M_1) is the natural projection, and $f_2 : M_1/\mathbb{Z}_{d_1} \rightarrow M_2$ is a fiber-preserving branched covering with fiber degree one.*

Proof Let $\{O_1, \dots, O_n\}$ be a nonempty collection of finitely many fibers in M_2 , including all singular fibers of the Seifert fibration of M_2 and downstairs branch set of f , and such that all the singular fibers of the Seifert fibration of M_1 are contained in $f^{-1}(O_1 \cup \dots \cup O_n)$. Let V_1, \dots, V_n be pairwise disjoint Seifert fibered solid tori in M_2 with central fibers O_1, \dots, O_n respectively, and $N_2 = M_2 \setminus \text{int}(V_1 \cup \dots \cup V_n)$. Let $N_1 = f^{-1}(N_2)$. Then both N_1 and N_2 are trivial S^1 -bundles (over compact, connected, orientable surfaces with boundary), and $f|_{N_1} : N_1 \rightarrow N_2$ is a fiber-preserving d -fold unbranched covering. When restricted to a fiber of N_1 , f is a d_1 -fold covering of a fiber in N_2 for some positive integer d_1 with $d_1|d$. Now we see that $f|_{N_1} = f'_2 \circ f'_1$, where $f'_1 : N_1 \rightarrow N_1/\mathbb{Z}_{d_1}$ (here the \mathbb{Z}_{d_1} -action on N_1 is inside the S^1 -action on N_1 coming from that on M_1) is the natural projection, and $f'_2 : N_1/\mathbb{Z}_{d_1} \rightarrow N_2$ is a fiber-preserving unbranched covering with fiber degree one. (By the way, one can easily check that f'_1 (resp. f'_2) sends a meridian in any

boundary torus of N_1 (resp. N_1/\mathbb{Z}_{d_1}) to a multiple of a meridian in the image torus.) Now we let V_1, \dots, V_n be shrunk to the central fibers O_1, \dots, O_n (at the same time), and let f_1 (resp. f_2) be the limit of the mappings f'_1 (resp. f'_2) thus gotten in the process. Then we are done. (Compare [1].) \square

Remark. See Theorem 3.3.1 in [3] for a result similar to Proposition 0.2 above. Our proof here is slightly different, and is along the lines of proof of Lemma 2.1 in [2].

References

- [1] R. H. Fox, Covering spaces with singularities, In Algebraic Geometry and Topology, A Symposium in Honor of S. Lefschetz, Edited by R. H. Fox, D. C. Spencer, and A. W. Tucker, pp. 243-257, Princeton University Press, 1957.
- [2] H. Huang, Branched coverings and nonzero degree maps between Seifert manifolds, Proc. Amer. Math. Soc. 130 (2002), no. 8, 2443-2449.
- [3] J. Remigio-Juárez, Coverings of Seifert manifolds branched along fibers, Ph.D. Thesis, 2008.
- [4] D. Rolfsen, Knots and links, AMS Chelsea Publishing, 2003.
- [5] H. Seifert, Topologie dreidimensionaler gefaserner Räume, Acta Math. 60 (1933), 147-238. (There is an English translation by W. Heil appended to H. Seifert and W. Threlfall, A textbook of topology, Academic Press, 1980.)

School of Mathematical Sciences, Beijing Normal University,
Laboratory of Mathematics and Complex Systems, Ministry of Education,
Beijing 100875, P.R. China
E-mail address: hhuang@bnu.edu.cn