

A Dutch Book theorem for partial subjective probability

Maurizio Negri,
Università di Torino

September 2, 2018

Abstract

The aim of this paper is to show that partial probability can be justified from the standpoint of subjective probability in much the same way as classical probability does. The seminal works of Ramsey and De Finetti have furnished a method for assessing subjective probabilities: ask about the bets the decision-maker would be willing to place. So we introduce the concept of partial bet and partial Dutch Book and prove for partial probability a result similar to the Ramsey-De Finetti theorem. Finally, we make a comparison between two concepts of bet: we can bet our money on a sentence describing an event, or we can bet our money on the event itself, generally conceived as a set. These two ways of understanding a bet are equivalent in classical probability, but not in partial probability.

Keywords: Dutch Book theorem; non-classical probability; Kleene's Logic.

1 Partial probability

Classical probability theory is grounded on the concept of probability space, a triple (A, \mathcal{C}_A, p) where A is a sample space, \mathcal{C}_A is a field of sets on A and $p : \mathcal{C}_A \rightarrow [0, 1]$ is a probability measure, i.e. a function satisfying Kolmogoroff's axioms: 1) $p(A) = 1$, 2) $p(X \cup Y) = p(X) + p(Y)$, when $X \cap Y = \emptyset$. In this way probability is seen as the measure of an event represented by a set of outcomes. Partial probability theory arises when we substitute classical events with partial events (see [6]). If S is the sample space of an experiment, we define the set of all *partial sets on S* as the set $D(S) = \{(A, B) : A, B \subseteq S, \text{ and } A \cap B = \emptyset\}$. Every $x \in A$ (resp. B) is said to be a *positive* (resp. *negative*) element of the

partial set (A, B) . Partial sets are structured by the following operations:

$$\begin{aligned}
(A, B) \sqcap (C, D) &= (A \cap C, B \cup D), \\
(A, B) \sqcup (C, D) &= (A \cup C, B \cap D), \\
-(A, B) &= (B, A), \\
0 &= (\emptyset, S), \\
1 &= (S, \emptyset), \\
n &= (\emptyset, \emptyset).
\end{aligned}$$

The algebra $\mathcal{D}(S) = (D(S), \sqcap, \sqcup, -, 0, n, 1)$ is the *algebra of all partial sets* on S . A *field of partial set* is any subalgebra $\mathcal{G}_S \subseteq \mathcal{D}(S)$. A partial set (A, B) is a *Boolean partial set* if $A = S - B$. The set of Boolean partial sets is a Boolean algebra isomorphic to $\mathcal{P}(S)$, the classical power-set algebra. We define a binary relation between partial sets setting $(A, B) \sqsubseteq (C, D)$ iff $A \subseteq C$ and $D \subseteq B$. It can be easily proved that $(D(S), \sqsubseteq)$ is a partially ordered set with (S, \emptyset) as top and (\emptyset, S) as bottom element.

When S is a sample space, we say that $\mathcal{D}(S)$ is the *algebra of partial events* on S . In relation with the experimental result $s \in S$, we say that (A, B) occurs positively if $s \in A$, occurs negatively if $s \in B$, is uncertain otherwise. Events of classical probability theory are to be identified with Boolean partial sets and will be called *Boolean* or *classical events*.

We define on R^2 the relation \preceq setting $(x, y) \preceq (w, z)$ iff $x \leq w$ e $z \leq y$. (Note that the natural order of R is reversed on the second elements of the ordered pairs.) It can be easily shown that (R^2, \preceq) is a partially ordered set. The operations $x + y$, xy and $-x$ in R are extended pointwise to the product R^2 , i.e. $(x, y) + (z, w) = (x + z, y + w)$, $(x, y)(z, w) = (xz, yw)$ and $-(x, y) = (-x, -y)$. We define the set of *partial probability values* T as a subset of R^2 , setting $T = \{(x, y) \in [0, 1]^2 : x + y \leq 1\}$. The probability value of a partial event (A, B) is a pair $(x, y) \in T$. The set of partial probability values T is partially ordered by \preceq with a maximum $(1, 0)$ and a minimum $(0, 1)$.

Given a partial field of set \mathcal{G}_S on S and a function $\mu : \mathcal{G}_S \rightarrow T$, we say that μ is a *measure of partial probability* when the following axioms are satisfied:

1. $\mu(S, \emptyset) = (1, 0)$.
2. $\mu(A, B) + \mu(C, D) = \mu((A, B) \sqcup (C, D)) - \mu((A, B) \sqcap (C, D))$.
3. $\mu(-(A, B)) = \sigma(\mu(A, B))$,
4. $(\emptyset, \emptyset) \sqsubseteq (A, B)$ implies $(0, 0) \preceq \mu(A, B)$,

where $\sigma : [0, 1]^2 \rightarrow [0, 1]^2$ is defined by $\sigma(x, y) = (y, x)$. As a consequence of axiom 4, we have $(0, 0) \preceq \mu(A, \emptyset)$, for all $(A, \emptyset) \in \mathcal{G}_S$. A *partial probability space* is a triple (S, \mathcal{G}_S, μ) where \mathcal{G}_S is a field of partial sets on the sample space S and μ is a measure of partial probability.

Given a classical probability space $(S, P(S), p)$, we define the *partial probability space associated to* $(S, P(S), p)$ as follows: we define $\mu : D(S) \rightarrow T$ setting

$$\mu(A, B) = (p(A), p(B)).$$

As (A, B) is a partial event, $A \cap B = \emptyset$ so $p(A) + p(B) = p(A \cup B)$: this proves that $\mu(A, B) \in T$. It can be easily proved that μ satisfies the four axioms above, so $(S, \mathcal{D}(S), \mu)$ is a partial probability space. For instance, we can start from the classical probability space $(S, P(S), p)$, where $S = \{n : 1 \leq n \leq 6\}$ and $p(n) = 1/6$ for all $n \in S$. If (A, B) is the Boolean event with $A = \{2, 4, 6\}$ and $B = \{1, 3, 5\}$, then $\mu(A, B) = (1/2, 1/2)$, and if (C, D) is the partial event with $C = \{2, 4\}$ and $D = \{5\}$, then $\mu(C, D) = (1/3, 1/6)$. If the outcome of the experiment is 3, then we say that (A, B) does not happen and that (C, D) is uncertain (neither happens nor does not happen). Some events like (A, \emptyset) have only positive occurrences (may only happen), others like (\emptyset, A) have only negative occurrences, (\emptyset, \emptyset) is the absolutely undefined event.

We list some differences between classical and partial probability.

1. (A, B) or not- (A, B) is no more the certain event, unless (A, B) is Boolean, i.e. $B = S - A$. In general we can only say that $\mu((A, B) \sqcup -(A, B)) = \mu(A \cup B, \emptyset) \succeq (0, 0)$. For the same reason, (A, B) and not- (A, B) is not, in general, the impossible event.
2. Additivity holds in the form of axiom 2, but now we have different kinds of disjointness. In a field of partial sets $\mathcal{D}(S)$, the partial sets like (\emptyset, X) , belonging to the interval $[(\emptyset, S), (\emptyset, \emptyset)]$, are generalizations of the empty set (they have no positive elements); so we can say that (A, B) is disjoint from (C, D) if $(A, B) \cap (C, D) = (\emptyset, X)$, for some X . Now the partial probability of a sum of disjoint sets is the sum of their probabilities, $\mu(A, B) + \mu(C, D) = \mu((A, B) \sqcup (C, D))$, only when (A, B) and (C, D) have the maximum degree of disjointness (\emptyset, \emptyset) .
3. Partial probability values are only partially ordered by \preceq , so we cannot say, in general, whether a partial event is more probable than another partial event or not. However, it can be easily seen that the set of partial probability values of Boolean events is totally ordered by \preceq .
4. Conditioning on partial events is possible, but with a slightly different meaning with respect to the classical case. See [6] par. 4 and [7] par. 9, 13

The aim of this paper is to show that partial probability can be justified from the standpoint of subjective probability in much the same way as classical probability does. The seminal works of Ramsey and De Finetti have furnished a method for assessing subjective probabilities: ask about the bets the decision-maker would be willing to place. In par. 2 we introduce the concept of classical bet and shortly review the proof of Ramsey-De Finetti theorem for classical probability. The remaining part of this paper is devoted to prove an analogous

result in the case of partial subjective probability. To this scope we introduce in par. 3 the fundamentals concepts of Kleene logic. In par. 4 we introduce the concept of partial bet and partial Dutch Book and prove for partial probability a result similar to Ramsey-De Finetti theorem. In par. 5 we make a comparison between two concepts of bet: we can bet our money on a sentence describing an event, or we can bet our money on the event itself, generally conceived as a set. These two ways of understanding a bet are equivalent in the classical case, but not in the partial case.

2 Classical bets

In the subjective approach to probability, the bearers of probability are no more sets but sentences and the probability value is conceived as the degree of belief in some event represented by a sentence. So we introduce a sentential n -ary language L_n based on the sentential variables $P_n = \{p_1, \dots, p_n\}$, the connectives $\{\neg, \wedge, \vee\}$ and the constants $\{0, 1\}$. (Through this text we deal only with finite languages.) We denote with F_n the set of formulas of L_n and with \mathcal{F}_n the algebra of formulas. (We simply write L , P and F when no confusion is possible.) We say that a function $\pi : F \rightarrow [0, 1]$ is a *probability function*, if the following axioms are satisfied:

1. if $\models \alpha$ then $\pi(\alpha) = 1$,
2. if $\models \neg(\alpha \wedge \beta)$ then $\pi(\alpha \vee \beta) = \pi(\alpha) + \pi(\beta)$,

where \models is the consequence relation of bivalent logic. (See, for instance, [5] or [8].) Both the measure-theoretic and the subjectivistic approach to classical probability are deeply grounded on Boolean algebras. On one side the events of a probability space constitute a Boolean algebra, on the other side the definition of a probability function requires the semantics of bivalent logic and Boolean truth-functions. Owing to this common ground, these two ways of presenting probability can be translated one into the other (see, for instance, [7], par.4).

When does a function $b : F \rightarrow [0, 1]$, representing the decision-maker's belief, satisfy the axioms of probability function? The theorem that we are going to prove, commonly attributed to Ramsey and De Finetti (see [3], p. 53), gives a sufficient condition through the concept of bet. We understand a *bet* as a triple (α, x, r) where $\alpha \in F$, $x \in [0, 1]$ and $r \in R$. The formula α represents the event on which the bet is placed, x is the betting quotient and r the stake. By accepting this bet, we agree to pay out rx to receive r from the bookmaker if the event described by α takes place, and nothing otherwise. The buyer's payoff can be described by the following table:

$$\begin{array}{ll} r - rx = r(1 - x) & \text{if } \alpha \text{ is true in the actual world,} \\ -rx & \text{if } \alpha \text{ is false in the actual world.} \end{array}$$

We can obtain the bookmaker's payoff by changing the sign of the stake. In general, when we talk of 'payoff', we understand the buyer's payoff.

The payoff is a function of α , x , r and of the state of the world that makes α true or false. The definition of such a function requires some basic concepts in the semantics of classical logic. We identify with 2^n the set of all possible worlds, where $2 = \{0, 1\}$ is the set of truth-values of classical logic. Then, for every sentential variable p_i , we can set $V_w(p_i) = w_i$: the truth-value of p_i in the world w . As \mathcal{F} is the absolutely free algebra, we know that V_w can be extended to a homomorphism $V_w : \mathcal{F} \rightarrow \mathbf{2}$, where $\mathbf{2}$ denotes the two-element Boolean algebra: $V_w(\alpha)$ is the truth-value of α in the world w . We say that α is a consequence of γ , in symbols $\gamma \models \alpha$, when $V_w(\gamma) \leq V_w(\alpha)$ for all $w \in 2^n$. We say that α is a consequence of the set of formulas Γ , and write $\Gamma \models \alpha$, when $\inf\{V_w(\gamma) : \gamma \in \Gamma\} \leq V_w(\alpha)$ for all $w \in 2^n$. When $\Gamma = \emptyset$ we write simply $\models \alpha$ and this amounts to $V_w(\alpha) = 1$, for all $w \in 2^n$, because $\inf\{V_w(\gamma) : \gamma \in \emptyset\} = \inf(\emptyset) = 1$. We say that two formulas α and β are *equivalent* and write $\alpha \equiv \beta$ iff $\alpha \models \beta$ and $\beta \models \alpha$.

The semantics of classical logic can also be defined through a function that associates to every formula α its meaning $M(\alpha)$, conceived as the set of worlds in which α holds true. To this end we define $M(p_i) = \{w \in 2^n : w_i = 1\}$, for all sentential variable p_i : intuitively, $M(p_i)$ is the set of all worlds making p_i true. As \mathcal{F} is the absolutely free algebra, we can extend M to a homomorphism from \mathcal{F} to the Boolean algebra $\mathcal{P}(2^n)$. The functions V_w and M are related in this way: given V_w , we can define $M(\alpha)$ as $\{w \in 2^n : V_w(\alpha) = 1\}$ and given M we can define $V_w(\alpha) = 1$ iff $w \in M(\alpha)$ and $V_w(\alpha) = 0$ iff $w \notin M(\alpha)$. So $\alpha \models \beta$ iff $M(\alpha) \subseteq M(\beta)$ and $\alpha \equiv \beta$ iff $M(\alpha) = M(\beta)$.

Now we return to the definition of the payoff. We associate to every bet (α, x, r) a function $[\alpha, x, r] : 2^n \rightarrow R$ that gives the *payoff of (α, x, r) in the world w* by

$$[\alpha, x, r](w) = r(V_w(\alpha) - x) = \begin{cases} r(1 - x) & \text{if } V_w(\alpha) = 1 \\ -rx & \text{if } V_w(\alpha) = 0. \end{cases}$$

This definition can be naturally extended to finite sets of bets as follows. If $B = \{(\alpha_i, x_i, r_i) : 1 \leq i \leq n\}$, then we define the payoff of the set B in the world w as

$$[B](w) = \sum_{i=1}^n \{[\alpha_i, x_i, r_i](w)\}.$$

We say that a set of bets B is a *Dutch Book* if, for all $w \in 2^n$, $[B](w) < 0$. Intuitively, a Dutch Book is a set of bets that guarantees a sure loss for the buyer. There is, however, a weaker notion of Dutch Book that will be useful in the following, where the set of bets can at best break even and, in at least one possible world, has a net loss. We say that a set of bets B is a *Weak Dutch Book* if, for all $w \in 2^n$, $[B](w) \leq 0$ and in at least one world w , $[B](w) < 0$. (This kind of Dutch Book has been introduced in [9].) If we take the values of $b : F \rightarrow [0, 1]$ as betting quotients, i.e. the bets in B are of kind $(\alpha_i, b(\alpha_i), r)$, and if a Dutch Book is available with such betting rates, then we say that there is a *Dutch Book for b* .

Lemma 1 For all $w \in 2^n$, $V_w(\alpha \vee \beta) = V_w(\alpha) + V_w(\beta) - V_w(\alpha \wedge \beta)$.

Proof. The proof is an easy calculation with truth-tables and is left to the reader. ■

Theorem 2 *If $b : F \rightarrow [0, 1]$ and there is no Dutch Book for b , then b is a probability function.*

Proof. Axiom 1, if $\models \alpha$ then $b(\alpha) = 1$. If $b(\alpha) < 1$, then $B = \{(a, b(\alpha), -1)\}$ is a Dutch Book because, for all $w \in 2^n$, $V_w(\alpha) = 1$ and then

$$[B](w) = [\alpha, b(\alpha), -1](w) = -1(V_w(\alpha) - b(\alpha)) = b(\alpha) - 1 < 0.$$

Axiom 2, if $\models \neg(\alpha \wedge \beta)$ then $b(\alpha \vee \beta) = b(\alpha) + b(\beta)$. Firstly we prove that

$$\text{if } \models \neg\alpha \text{ then } b(\alpha) = 0. \quad (1)$$

If $0 < b(\alpha)$, then $B = \{(a, b(\alpha), 1)\}$ is a Dutch Book because, for all $w \in 2^n$, $V_w(\alpha) = 0$ and then $[B](w) = [\alpha, x, 1](w) = 1(V_w(\alpha) - x) = 0 - x < 0$.

Secondly we show that

$$b(\alpha \vee \beta) + b(\alpha \wedge \beta) = b(\alpha) + b(\beta). \quad (2)$$

Let $b(\alpha) = x$, $b(\beta) = y$, $b(\alpha \vee \beta) = z$, and $b(\alpha \wedge \beta) = w$. If $z + w > x + y$, we define $B = \{(\alpha \vee \beta, z, 1), (\alpha \wedge \beta, w, 1), (\alpha, x, -1), (\beta, y, -1)\}$. We show that B is a Dutch Book. For all $w \in 2^n$, we have

$$\begin{aligned} [B](w) &= (V_w(\alpha \vee \beta) - z) + (V_w(\alpha \wedge \beta) - w) + (x - V_w(\alpha)) + (y - V_w(\beta)) \\ &= (V_w(\alpha \vee \beta) + V_w(\alpha \wedge \beta) - V_w(\alpha) - V_w(\beta)) + (x + y - z - w) \\ &= x + y - z - w \\ &< 0, \end{aligned}$$

because $V_w(\alpha \vee \beta) + V_w(\alpha \wedge \beta) - V_w(\alpha) - V_w(\beta) = 0$ by the above lemma. If $z + w < x + y$, just change the sign of the stakes.

Now we can easily prove axiom 2: if $\models \neg(\alpha \wedge \beta)$ then $b(\alpha \wedge \beta) = 0$, by 1, so $b(\alpha \vee \beta) = b(\alpha) + b(\beta)$ follows from 2. ■

3 Partial subjective probability and Kleene logic

As we have seen in par. 2, classic probability can be understood as a degree of belief in a sentence. We can do the same with partial probability: we have only to shift from bivalent logic to Kleene logic and from probability values in $[0, 1]$ to probability values in T . This should justify a brief digression in the semantics of Kleene logic.

The language of Kleene n -ary logic is $L_n^* = L_n \cup \{n\}$, where L_n is the n -ary language of classical logic introduced in par. 2. We denote with F_n^* the set of formulas of L_n^* and with \mathcal{F}_n^* the algebra of n -ary formula. We write simply L^* , F^* and \mathcal{F}^* when no confusion is possible. We denote with K the set $\{0, n, 1\}$ of truth-values of Kleene logic, where n stands for ‘neutral’ or ‘uncertain’. Every

$w \in K^n$ can be seen as an instantaneous description of the world, at the level of the atomic facts represented by sentential variables, so we can define a function V_w from $\{p_i : i \leq n\}$ to K setting $V_w(p_i) = w_i$. When $w_i = n$, the atomic fact represented by p_i neither happens nor does not happen. This uncertainty may be of an epistemic kind, related to a lack of knowledge, or may be deeply rooted in the reality. The next step is extending V_w to all formulas and to this end we give an algebraic structure to the set of truth-values as follows: we define on K the total order $0 < n < 1$ and define $x \wedge y = \min(x, y)$ and $x \vee y = \max(x, y)$, what amounts to giving the following truth-tables:

\wedge	1	0	n	\vee	1	0	n
1	1	0	n	1	1	1	1
0	0	0	0	0	1	0	n
n	n	0	n	n	1	n	n

As for negation, we set $\neg(n) = n$, $\neg(0) = 1$, $\neg(1) = 0$. Finally, we denote with \mathcal{K} the algebra $(K, \wedge, \vee, \neg, 0, 1, n)$. As \mathcal{F}^* is the absolutely free algebra, we can extend V_w to an homomorphism $V_w : \mathcal{F} \rightarrow \mathcal{K}$. So we say that α is *true in w* iff $V_w(\alpha) = 1$, *false* if $V_w(\alpha) = 0$ and *neutral* if $V_w(\alpha) = n$. We say that two formulas α and β are *equivalent* and write $\alpha \equiv \beta$ iff $V_w(\alpha)_w = V_w(\beta)_w$, for all $w \in K^n$.

The semantics of Kleene logic can also be defined through a function that associates to every formula α its meaning $M(\alpha)$ conceived as a partial set in $D(K^n)$, where $M(\alpha)_0$ and $M(\alpha)_1$ are the set of worlds in which α is respectively true and false, the positive and the negative models of α . Firstly, we define a function $M : \{p_i : i \leq n\} \rightarrow D(K^n)$ setting

$$M(p_i) = (\{w \in K^n : w_i = 1\}, \{w \in K^n : w_i = 0\}).$$

As \mathcal{F}^* is free, we extend M to a homomorphism $M : \mathcal{F} \rightarrow \mathcal{D}(K^n)$. Then the meanings of formulas can be recursively defined by the following equations:

$$\begin{aligned} M(\alpha \wedge \beta) &= M(\alpha) \sqcap M(\beta), \\ M(\alpha \vee \beta) &= M(\alpha) \sqcup M(\beta), \\ M(\neg\alpha) &= -M(\alpha), \\ M(0) &= (\emptyset, K^n), \\ M(1) &= (K^n, \emptyset), \\ M(n) &= (\emptyset, \emptyset). \end{aligned}$$

These two ways of giving a semantics are equivalent: if we take the notion of meaning given by M as primitive, then we can define

$$V_w(\alpha) = \begin{cases} 0 & \text{if } w \in M(\alpha)_1, \\ 1 & \text{if } w \in M(\alpha)_0, \\ n & \text{if } w \in K^n - (M(\alpha)_0 \cup M(\alpha)_1); \end{cases}$$

if we take the notion of truth in the possible world w given by V_w as primitive, then

$$M(\alpha) = (\{w \in K^n : V_w(\alpha) = 1\}, \{w \in K^n : V_w(\alpha) = 0\}).$$

We define the notion of *logical consequence* as follows: $\alpha \models \beta$ iff $M(\alpha) \sqsubseteq M(\beta)$ iff $M(\alpha)_0 \subseteq M(\beta)_0$ and $M(\beta)_1 \subseteq M(\alpha)_1$ iff every positive model of α is a positive model of β and every negative model of β is a negative model of α . The notion of logical consequence can be generalized to

$$\Gamma \models \alpha \text{ iff } \bigcap \{M(\gamma) : \gamma \in \Gamma\} \sqsubseteq M(\alpha).$$

In terms of V_s , the definition runs as follows:

$$\Gamma \models \alpha \text{ iff, for all } s \in K^n, \inf\{V_s(\gamma) : \gamma \in \Gamma\} \leq V_s(\alpha).$$

As in the case of classical logic, an easy calculation based on truth-tables gives the following theorem.

Theorem 3 *For all $w \in K^n$, $V_w(\alpha \vee \beta) = V_w(\alpha) + V_w(\beta) - V_w(\alpha \wedge \beta)$.*

Now we can define the notion of partial probability function. As the axioms of probability function (see par. 2) are modeled on Kolmogoroff's axioms, so the axioms of partial probability function are modeled on the axioms of measure of partial probability introduced in par. 1. We say that $\pi : F^* \rightarrow T$ is a *partial probability function* if the following axioms are satisfied, where \models denotes logical consequence in Kleene logic:

1. $1 \models \alpha$ implies $\pi(\alpha) = (1, 0)$,
2. $\pi(\alpha \vee \beta) = \pi(\alpha) + \pi(\beta) - \pi(\alpha \wedge \beta)$,
3. $\pi(\neg\alpha) = \sigma(\pi(\alpha))$,
4. $n \models \alpha$ implies $(0, 0) \preceq \pi(\alpha)$.

We call $\pi(\alpha)$ the *partial degree of belief* in α of our decision-maker. The following theorem shows some fundamental properties of π .

Theorem 4 *If π is a partial probability function on L_n^* , then*

1. $\pi(n) = (0, 0)$.
2. $\alpha \models 0$ implies $\pi(\alpha) = (0, 1)$,
3. $\alpha \models n$ implies $\pi(\alpha) \preceq (0, 0)$,
4. $\pi(\alpha) = \pi(\alpha \vee n) + \pi(\alpha \wedge n)$.

Proof. 1. From $n \models n$ we have $(0,0) \preceq \pi(n)$, by axiom 4. In Kleene logic we have $n \models \neg n$ and so $(0,0) \preceq \pi(\neg n)$. In general we have $(x,y) \preceq (x',y')$ iff $\sigma(x',y') \preceq \sigma(x,y)$, thus $\sigma(\pi(\neg(n))) \preceq (0,0)$. By axiom 3, $\sigma(\pi(\neg(n))) = \sigma(\sigma(\pi(n))) = \pi(n)$ holds. Thus $\pi(n) \preceq (0,0)$.

2. In Kleene logic $\alpha \models 0$ implies $1 \models \neg\alpha$ and so, by axiom 1, $\pi(\neg\alpha) = (1,0)$. Thus, by axiom 3, $\sigma(\pi(\alpha)) = (1,0) \in \pi(\alpha) = (0,1)$.

3. In Kleene logic $\alpha \models n$ implies $n \models \neg\alpha$, so $(0,0) \preceq \pi(\neg\alpha) = \sigma(\pi(\alpha))$ and $\pi(\alpha) \preceq (0,0)$.

4. $\pi(\alpha \vee n) = \pi(\alpha) + \pi(n) - \pi(\alpha \wedge n) = \pi(\alpha) - \pi(\alpha \wedge n)$ by axiom 2 and point 1). ■

4 Partial bets

A classical bet is a mechanism that receives a triple constituted by a sentence α in the language L of classical logic, a betting quotient $x \in [0,1]$ and a stake $r \in R$ as input, and gives a payoff, represented by a real number as output. We observe that both payoffs and stakes belong to R , besides, the set of truth-values of sentences $\{0,1\}$, betting rates $[0,1]$ and stakes R are related by $\{0,1\} \subseteq [0,1] \subseteq R$. As a result we have been able to define the payoff $[\alpha, x, r](w)$ of a classical bet in the world w as $r(V_w(\alpha) - x)$.

When we consider a partial bet, we begin with a sentence α in the language L^* of Kleene and a betting quotient $(x,y) \in T$. The first step is finding a common ground for truth-values of Kleene logic and partial probability values, so we make $K \subseteq T$ by giving a new definition of K . So, from now on, we set $K = \{(0,1), (0,0), (1,0)\}$. After all, the essential nature of truth-values is immaterial, as long as their formal properties remain unchanged. If we restrict to K the partial order \preceq defined on T , we have $(0,1) \preceq (0,0) \preceq (1,0)$ where $(0,1)$ stands for ‘false’, $(0,0)$ for ‘neutral’ and $(1,0)$ for ‘true’. We define the algebra \mathcal{K} of truth values as above. For reader’s convenience we write the new truth-tables:

\wedge	(1,0)	(0,1)	(0,0)	\vee	(1,0)	(0,1)	(0,0)
(1,0)	(1,0)	(0,1)	(0,0)	(1,0)	(1,0)	(1,0)	(1,0)
(0,1)	(0,1)	(0,1)	(0,1)	(0,1)	(1,0)	(0,1)	(0,0)
(0,0)	(0,0)	(0,1)	(0,0)	(0,0)	(1,0)	(0,0)	(0,0)

As for \neg , we have $\neg(0,1) = (1,0)$ and $\neg(0,0) = (0,0)$. Finally we get a homomorphism $V_w : \mathcal{F}^* \rightarrow \mathcal{K}$ and $V_w(\alpha)$ will be the truth-value of α in the world $w \in K^n$. The proof of lemma 3 remains obviously unchanged.

The second step is finding a common ground for partial probability values and stakes, so we define a stake as a pair (h,k) in R^2 . The payoff of a partial bet will be an element of R^2 too. Stakes and payoffs are ordered by the same partial order \preceq defined on partial probability values. So R^2 is the common ground for truth-values, partial probability values, stakes and payoffs

In view of the particular partial ordering of payoffs, some considerations are in order. (And the same holds for stakes.) A payoff (h, k) has a positive part h and a negative part k . If $h > 0$ then h is a reward and if $h < 0$ then h is a loss, but on the second elements the order is reversed, so $k > 0$ is a true loss and $k < 0$ is a true reward. The positive and negative parts may come from completely different domains, so we may think the first value h representing the gain or loss of money and the second value k representing a degree of physical pain or gratification. What is essential is that we cannot in principle strike a balance between the first and the second component of (h, k) and reduce the pair to a single number.

As \preceq is a partial order, we cannot say in general whether (h, k) is better than (h', k') , but we can make the following distinctions. We can partition R^2 in three exhaustive and disjoint subsets: the diagonal $\delta = \{(x, x) : x \in R\}$, the pairs under the diagonal $\delta^+ = \{(x, y) : x > y\}$ and the pairs over the diagonal $\delta^- = \{(x, y) : x < y\}$. The elements of δ are ‘neutral payoffs’: if the first component h is positive and so can be seen as a gain, then the second element h represents a loss of the same intensity. The situation is reversed when h is negative. The elements of δ^+ give more reward than punishment and the elements of δ^- behave in the opposite way, so a payoff in δ^+ can be seen as ‘good’ and a payoff in δ^- as ‘bad’.

Now we can introduce the concept of *partial bet* as a triple $(\alpha, (x, y), (h, k))$ where α is a formula of L_n^* , $(x, y) \in T$, $(h, k) \in R^2$. We can describe the payoff of the buyer for $(\alpha, (x, y), (h, k))$ by the following table,

$$\begin{aligned} (h, k)((1, 0) - (x, y)) &= (h(1 - x), -ky) && \text{if } \alpha \text{ is true in the actual world} \\ (h, k)((0, 0) - (x, y)) &= (-hx, -ky) && \text{if } \alpha \text{ is neutral in the actual world} \\ (h, k)((0, 1) - (x, y)) &= (-hx, k(1 - y)) && \text{if } \alpha \text{ is false in the actual world.} \end{aligned}$$

The net gain varies as the actual world w varies in the set K^n of the possible worlds, so we define a function $[\alpha, (xy), (h, k)]:K^n \rightarrow R^2$ that gives the *payoff* of $(\alpha, (xy), (h, k))$ in the world w by

$$[\alpha, (xy), (h, k)](w) = (h.k)(V_w(\alpha) - (x, y)).$$

We have $[\alpha, (xy), (h, k)](w) =$

$$\begin{cases} (h, k)((1, 0) - (x, y)) = (h(1 - x), -ky) & \text{if } V_w(\alpha) = (1, 0) \\ (h, k)((0, 0) - (x, y)) = (-hx, -ky) & \text{if } V_w(\alpha) = (0, 0) \\ (h, k)((0, 1) - (x, y)) = (-hx, k(1 - y)) & \text{if } V_w(\alpha) = (0, 1). \end{cases}$$

As in the case of classical bets, this definition can be naturally extended to finite sets of bets as follows. If $B = \{(\alpha_i, (x_i, y_i), (h_i, k_i)) : 1 \leq i \leq n\}$, then we define the net gain of the set B in the world w as

$$[B](w) = \sum_{i=1}^n \{[\alpha_i, (x_i, y_i), (h_i, k_i)]\}.$$

The definition of Dutch Book is slightly different from the classical case. The payoffs of classical bets are in R with its natural order \leq , so 0 is an equilibrium point between loss, negative real numbers, and rewards, positive natural numbers. The payoffs of partial bets are in R^2 partially ordered by \preceq , where δ is a natural separation between loss, the pairs in δ^- , and reward, the pairs in δ^+ . So we say that a set of partial bets B is a *Dutch Book* if, for all $w \in 2^n$, $[B](w) \in \delta^-$. Intuitively, a Dutch Book is a set of bets that guarantees more loss than reward for the buyer. We say that a set of partial bets B is a *Weak Dutch Book* if, for all $w \in 2^n$, $[B](w) \in \delta^- \cup \delta$ and there is at least one possible world w such that $[B](w) \in \delta^-$: so in all cases we have no gain and in some cases a sure loss. If B is a Dutch Book then B is obviously a Weak Dutch Book too.

Given a function b from the set F^* of formulas of Kleene logic to the set T of partial probability values, we say that B is a *(Weak) Dutch Book for π* if B is a (Weak) Dutch Book and the bets in B are of kind $(\alpha_i, b(\alpha_i), (h_i, k_i))$, where the betting rates are given by b . Now we can prove a result similar to theorem 2.

Lemma 5 *For all $(x, y), (z, w) \in T$, if $(y, x) \neq (z, w)$ and $x + z = y + w$, then there are $h, h', k, k' \in R$ such that*

1. $hx + h'z = ky + k'w$,
2. $h < k'$ and $h' < k$.

Proof. Firstly, we observe that x, y, z and w all belong to $[0, 1]$ because (x, y) and (z, w) belong to T . Secondly, if we set $k' = q + h$ and $k = t + h'$, then we reduce ourselves to prove that there are q, t, h, h' , with $q, t > 0$ such that

$$hx + h'z = (t + h')y + (q + h)w. \quad (1)$$

Now we distinguish two cases.

Case1, $0 < y, w$. The above equation can be rewritten as $qw + ty = h(x - w) - h'(y - z)$. From our hypothesis $x + z = y + w$ we have $x - w = y - z$, so we obtain $qw + ty = h(x - w) - h'(x - w)$ that can also be written as

$$q = -\frac{y}{w}t + \frac{(h - h')(x - w)}{w}, \quad (2)$$

a linear equation in two unknowns q, t and parameters h, h', w, x, y , with $w, y \neq 0$. This equation can be plotted as a line with negative slope, as t varies on the horizontal axis and q on the vertical axis. We can choose h and h' such that the set of pairs (t, q) satisfying (1), with $t, q > 0$, is not empty. To this scope we make the t -intercept positive and the q -intercept positive, i.e.

$$t = \frac{(h - h')(x - w)}{y} > 0 \text{ and } q = \frac{(h - h')(x - w)}{w} > 0,$$

by choosing : i) $h > h'$ if $x > w$, ii) $h' > h$ if $w > x$. (Remember that $w, y \neq 0$.) With such a choice of h and h' , any pair (t, q) such that (2) holds and

$$0 \leq t \leq \frac{(h - h')(x - w)}{y} \text{ and } 0 \leq q \leq \frac{(h - h')(x - w)}{w}$$

gives a solution. (The case $x = w$ is impossible, because from $x = w$ and the hypothesis $x + z = y + w$ we have $y = z$, that is contrary to our hypothesis $(y, x) \neq (z, w)$.)

Case 2, $y = 0$, or $w = 0$.

Subcase a), $y = 0$. Firstly we prove that $w \neq 0$ and $z \neq 0$. If $w = 0$ then, from our hypothesis $x + z = y + w$ we have $x + z = 0$ and so $x = z = 0$. Then x, y, z, w are all 0 and an absurd follows our hypothesis $(y, x) \neq (z, w)$. If $z = 0$ then $(x, y) = (x, 0)$ and $(z, w) = (0, w)$, so from our hypothesis $(y, x) \neq (z, w)$ we have $x \neq w$. But $x + z = y + w$ holds by hypothesis, so $x + 0 = 0 + w$ and $x = w$, that is absurd. Now equation (1) reduces $hx + h'z = (q + h)w$, because $y = 0$ by hypothesis, that can be rewritten as

$$h(x - w) + h'z = qw.$$

By hypothesis, $x + z = y + w = w$, so $x - w = -z$, so the above equation becomes $-hz + h'z = qw$, that can be rewritten as

$$q = \frac{(h' - h)z}{w},$$

where $w > 0$ and $z > 0$, so we get $q > 0$ by choosing $h' > h$.

Subcase b), $w = 0$. The proof is similar to subcase a). ■

Theorem 6 *If $b : F \rightarrow T$ and there is no Weak Dutch Book for, then b is a partial probability function.*

Proof. We prove that b satisfies the four axioms of partial probability function.

Axiom 1. We must prove that $1 \models \alpha$ implies $b(\alpha) = (1, 0)$, so we suppose $b(\alpha) = (x, y) \neq (1, 0)$ and set $B = \{(\alpha, (x, y)), (-1, -1)\}$. We have, for all $w \in K^n$,

$$[B](w) = (-1, -1)(V_w(\alpha) - (x, y)) = (-1, -1)((1, 0) - (x, y)) = (x - 1, y).$$

From $(x, y) \neq (1, 0)$ we have $x < 1$. (As $(x, y) \in T$, $(x, y) \neq (1, 0)$ means $(x, y) \prec (1, 0)$, so $x < 1$.) So we have $x - 1 < 0$ and $0 \leq y$ and then $(x - 1, y) \prec (0, 0)$. So $[B](w) \in \delta^-$ for all $w \in K^n$ and B is a Dutch Book for b .

Axiom 2. We must prove that $b(\alpha \vee \beta) = b(\alpha) + b(\beta) - b(\alpha \wedge \beta)$.

Case 1: $b(\alpha) + b(\beta) \prec b(\alpha \vee \beta) + b(\alpha \wedge \beta)$. We set

$$B = \{(\alpha \vee \beta, b(\alpha \vee \beta), (1, 1)), ((\alpha \wedge \beta, b(\alpha \wedge \beta), (1, 1)), (\alpha, b(\alpha), (-1, -1)), (\beta, b(\beta), (-1, -1))\}.$$

Then we have, for all $w \in K^n$,

$$\begin{aligned}
[B](w) &= (1, 1)(V_w(\alpha \vee \beta) - b(\alpha \vee \beta)) + (1, 1)(V_w(\alpha \wedge \beta) - b(\alpha \wedge \beta)) + \\
&\quad (-1, -1)(V_w(\alpha) - b(\alpha)) + (-1, -1)(V_w(\beta) - b(\beta)) \\
&= V_w(\alpha \vee \beta) - b(\alpha \vee \beta) + V_w(\alpha \wedge \beta) - b(\alpha \wedge \beta) + b(\alpha) - \\
&\quad V_w(\alpha) + b(\beta) - V_w(\beta) \\
&= (V_w(\alpha \vee \beta) + V_w(\alpha \wedge \beta) - V_w(\alpha) - V_w(\beta)) + \\
&\quad (b(\alpha) + b(\beta) - b(\alpha \vee \beta) - b(\alpha \wedge \beta)) \\
&= b(\alpha) + b(\beta) - b(\alpha \vee \beta) - b(\alpha \wedge \beta) \\
&\prec (0, 0)
\end{aligned}$$

where the next to last line follows from lemma 3 and the last line from our hypothesis. As $[B](w) \in \delta^-$ for all $w \in K^n$, B is a Dutch Book for b .

Case 2: $b(\alpha \vee \beta) + b(\alpha \wedge \beta) \prec b(\alpha) + b(\beta)$. We set

$$\begin{aligned}
B &= \{(\alpha \vee \beta, b(\alpha \vee \beta), (-1, -1)), ((\alpha \wedge \beta, b(\alpha \wedge \beta), (-1, -1)), \\
&\quad (\alpha, b(\alpha), (1, 1)), (\beta, b(\beta), (1, 1))\}
\end{aligned}$$

and we get $[B](w) = b(\alpha \vee \beta) + b(\alpha \wedge \beta) - b(\alpha) - b(\beta) \prec (0, 0)$, so $[B](w) \in \delta^-$.

Axiom 3. We must prove that $b(\neg\alpha) = \sigma(b(\alpha))$, so we suppose that $b(\neg\alpha) \neq \sigma(b(\alpha))$. Let $b(\alpha) = (x, y)$ and $b(\neg\alpha) = (z, w)$, where $(z, w) \neq (y, x) = \sigma(b(\alpha))$. We distinguish three cases.

Case 1, $x + z < y + w$. If we set

$$B = \{(\alpha, (x, y), (-1, -1)), (\neg\alpha, (z, w), (-1, -1))\},$$

then we have, for all $w \in K^n$,

$$\begin{aligned}
[B](w) &= (-1, -1)(V_w(\alpha) - (x, y)) + (-1, -1)(V_w(\neg\alpha) - (z, w)) \\
&= (-1, -1)(V_w(\alpha) + \sigma(V_w(\alpha)) - (x + z, y + w)).
\end{aligned}$$

We note that

$$V_w(\alpha) + \sigma(V_w(\alpha)) = \begin{cases} (1, 1) & \text{if } V_w(\alpha) = (1, 0) \text{ or } V_w(\alpha) = (0, 1), \\ (0, 0) & \text{if } V_w(\alpha) = (0, 0), \end{cases}$$

so we have

$$[B](w) = \begin{cases} (-1, -1)((1, 1) - (x + z, y + w)) = ((x + z) - 1, (y + w) - 1), \\ (-1, -1)((0, 0) - (x + z, y + w)) = (x + z, y + w). \end{cases}$$

By hypothesis $x + z < y + w$, so in both cases $[B](w) \in \delta^-$ and B is a Dutch Book.

Case 2, $y + w < x + z$. We take B as in case 1, but change the stake $(-1, -1)$ in $(1, 1)$, then we have

$$[B](w) = \begin{cases} (1 - (x + z), 1 - (y + w)), \\ (-(x + z), -(y + w)). \end{cases}$$

In both cases we have a point in δ^- because, by hypothesis, $-(x+z) < -(y+w)$.

Case 3, $x+z = y+w$. We show that there are stakes (h, k) and (h', k') such that, setting

$$B = \{(\alpha, (x, y), (h, k)), (-\alpha, (z, w), (h', k'))\},$$

we have $[B](w) \in \delta^-$ for all $w \in K^n$. Firstly we observe that

$$\begin{aligned} [B](w) &= (h, k)(V_w(\alpha) - (x, y)) + (h', k')(V_w(-\alpha) - (z, w)) \\ &= \begin{cases} (h, k)((1, 0) - (x, y)) + (h', k')((0, 1) - (z, w)) \\ (h, k)((0, 0) - (x, y)) + (h', k')((0, 0) - (z, w)) \\ (h, k)((0, 1) - (x, y)) + (h', k')((1, 0) - (z, w)) \end{cases} \\ &= \begin{cases} (h(1-x), -ky) + (-h'z, k'(1-w)) \\ (-hx, -ky) + (-h'z, -k'w) \\ (-hx, k(1-y)) + (h'(1-z), -k'w) \end{cases} \\ &= \begin{cases} (h - (hx + h'z), k' - (ky + k'w)) \\ (hx + h'z, ky + k'w) \\ (h' - (hx + h'z), k - (ky + k'w)). \end{cases} \end{aligned}$$

Then, by the above lemma, we can choose h, h', k' such that: i) $hx + h'z = ky + k'w$, and ii) $h < k'$ and $h' > k$. So in the first and third case we have a point in δ^- and in the second case we have a point in δ . This proves that $[B](w) \in \delta^- \cup \delta$ and B is a Weak Dutch Book.

Axiom 4. We must prove that $n \models \alpha$ implies $(0, 0) \preceq b(\alpha)$. We suppose that $b(\alpha) = (x, y)$ and $(0, 0) \not\preceq (x, y)$. We set $B = \{(a, (x, y), (0, -1))\}$. From the hypothesis $n \models \alpha$ we know that, for all $w \in K^n$, $V_w(n) \leq V_w(\alpha)$. As $V_w(n) = (0, 0)$, we have $V_w(\alpha) = (0, 0)$ or $V_w(\alpha) = (1, 0)$. Then, for all $w \in K^n$,

$$[B](w) = (0, 1)(V_w(\alpha) - (x, y)) = \begin{cases} (0, -1)((1, 0) - (x, y)) & \text{if } V_w(\alpha) = (1, 0), \\ (0, -1)((0, 0) - (x, y)) & \text{if } V_w(\alpha) = (0, 0). \end{cases}$$

In both cases we have $[B](w) = (0, y)$. As (x, y) is a partial probability value, $0 \leq y \leq 1$; if $y = 0$ then $(0, 0) \preceq (x, y)$, but by hypothesis $(0, 0) \not\preceq (x, y)$, so $0 < y$. Then we have $(0, y) \prec (0, 0)$, so $[B](w) \in \delta^-$ and B is a Dutch Book. ■

5 Equivalence

We conclude our work with some remarks about the concept of bet. We start with classical bets. If $b : F \rightarrow [0, 1]$ is to represent the degree of belief of a decision-maker, logically equivalent formulas of classical logic should receive the same probability value. We can show that this is the case, when there is no Dutch Book for b . We suppose that $\alpha \equiv \beta$ in classical logic and $b(\alpha) \neq b(\beta)$. If $b(\beta) < b(\alpha)$ we set $B = \{(\alpha, b(\alpha), 1), (\beta, b(\beta), -1)\}$, so we have $[B](w) = V_w(\alpha) - V_w(\beta) + b(\beta) - b(\alpha) = b(\beta) - b(\alpha) < 0$ and B is a Dutch Book for b . If $b(\alpha) < b(\beta)$ we choose a stake -1 for the bet on α and a stake 1 for the bet on β .

From this observation we can see that the concept of bet introduced above is substantially equivalent to another that is common in the literature (see, for instance, [4] p. 20-23). In fact, we can define a bet as a triple (U, x, r) , where $U \subseteq 2^n$, $x \in [0, 1]$ and $r \in R$, so in this way we are betting on an event, a set of possible world U , instead of betting on a formula α . If we denote with \mathcal{B} the set of all bets of kind (α, x, r) and define a relation $(\alpha, x, r) \sim (\beta, x, r)$ iff $\alpha \equiv \beta$, then \sim is an equivalence relation and we can take the quotient \mathcal{B}/\sim . Now there is a bijection f between \mathcal{B}/\sim and the set of all bets of kind (U, x, r) : just set $f(|(\alpha, x, r)|) = (M(\alpha), x, r)$, where $M(\alpha)$ denotes the set of possible worlds that is the meaning of α , as defined in 2. The function f is well defined because if $(\alpha, x, r) \sim (\beta, x, r)$ then $\alpha \equiv \beta$ and $M(\alpha) = M(\beta)$. The function is obviously injective and is surjective because every $U \subseteq 2^n$ is $M(\alpha)$ for some α . (This follows from the theorem on disjunctive normal form of classical logic.) As a consequence of the observation above, all bets in the class $|(\alpha, x, r)|$ behave in the same way with respect to b , when no Dutch Book is possible for b , so we can attach the probability value $b(\alpha)$ to the class $|(\alpha, x, r)|$ and this value can be transferred to $M(\alpha)$.

The same problem can be posed for partial bets: if $b : F^* \rightarrow T$ is to represent the partial degree of belief of a decision-maker, logically equivalent formulas of Kleene logic should receive the same partial probability value. The following theorem proves that this is the case when there is no Dutch Book for b .

Theorem 7 *Let $b : F^* \rightarrow T$ and suppose that there is no Dutch Book for b . If $\alpha \equiv \beta$ in Kleene logic then $b(\alpha) = b(\beta)$.*

Proof. We assume that $\alpha \equiv \beta$ and $b(\alpha) = (x, y) \neq (z, w) = b(\beta)$, then we show that there is a Dutch Book for b .

Case 1, $z - x < w - y$. We set $B = \{(\alpha, (x, y), (1, 1)), (\beta, (z, w), (-1, -1))\}$
Then

$$\begin{aligned} [B](w) &= (1, 1)(V_w(\alpha) - (x, y)) + (-1, -1)(V_w(\beta) - (z, w)) \\ &= (1, 1)V_w(\alpha) - (1, 1)(x, y) + (-1, -1)V_w(\beta) - (-1, -1)(z, w) \\ &= -(x, y) - (-z, -w) \\ &= (z - x, w - y). \end{aligned}$$

As $(z - x, w - y) \prec (0, 0)$, we have $[B](w) \in \delta^-$ and B is a Dutch Book.

Case 2, $w - y < z - x$. We set $B = \{(\alpha, (x, y), (-1, -1)), (\beta, (z, w), (1, 1))\}$
so that $[B](w) = (x - z, y - w) \in \delta^-$.

Case 3, $z - x = w - y$. Firstly we observe that $z - x \neq 0$. (If $z - x = 0$ then $w - y = 0$ and x, y, z and w are all 0, that is contrary to our hypothesis $(x, y) \neq (z, w)$.) Then we distinguish two cases.

Subcase a), $z - x < 0$. Then $w - y < 0$ and $y - w > 0$, so $z - x < y - w$ and $(z - x < y - w) \in \delta^-$. We obtain a Dutch Book choosing $B = \{(\alpha, (x, y), (1, -1)), (\beta, (z, w), (-1, 1))\}$, because in this case $[B](w) = (z - x, y - w)$.

Subcase b), $0 < z - x$. We set $B = \{(\alpha, (x, y), (-1, 1)), (\beta, (z, w), (1, -1))\}$.

■

Now we can take equivalence classes of partial bets, as we did in the case of classical bets. As all bets in the class $|(\alpha, x, r)|$ behave in the same way with respect to b , when no Dutch Book is possible for b , we can attach the partial probability value $b(\alpha)$ to the class $|(\alpha, (x, x'), (r, r'))|$, but here the analogy breaks off. If we define a concept of bet where the formula α is replaced by a partial set of possible worlds, i.e. we define a bet as a triple $((U, U'), (x, x'), (r, r'))$, where $(U, U') \subseteq \mathcal{D}(K^n)$, $(x, x') \in T$ and $(r, r') \in R^2$, we obtain a more comprehensive concept of partial bet: there are bets on partial events that cannot be simulated by a bet on a sentence. In fact, we may define a function $f(|(\alpha, (x, x'), (r, r'))|) = (M(\alpha), (x, x'), (r, r'))$ as above, where $M(\alpha)$ is the partial set of positive and negative models of α defined in 3, but now f is strictly injective, because there are partial sets in $\mathcal{D}(K^n)$ that cannot be denoted by a formula α . We end this work with a proof of this theorem.

The elements of K , that have been up to now understood as truth values, may also be conceived as values assigned to the amount of information carried by a proposition, to the degree of exactness of an assertion. (In this section we take the original definition of K as $\{0, n, 1\}$.) To this end, we define on K a partial order \preceq setting $x \preceq y$ iff $x = y$ or $(x = n \text{ and } y = 0)$ or $(x = n \text{ and } y = 1)$. This partial order can be pointwise extended to K^n , so given $s, t \in K^n$, $s \preceq t$ iff for all $i < n$, $s_i \preceq t_i$. The partially ordered set (K^n, \preceq) has a minimum \bar{n} , the sequence that takes always n as value, and 2^n maximal elements, the sequences that take always 0 or 1 as value.

If s and t are seen as possible worlds, then $s \preceq t$ means that the situation represented by t is at least so defined as the one described by s . What in s has been definitely settled (marked with 0 or 1) remains unchanged in the passage to t ; what has not been definitely settled in s (marked with n) may be (positively or negatively) settled in t . Whereas $V_s(\alpha) \leq V_t(\alpha)$ means that there is no loss in truth-value in the passage from α to β , $V_s(\alpha) \preceq V_t(\alpha)$ means that there is no loss of information.

Theorem 8 For all $s, t \in K^n$ and all $\alpha \in F^*$,

1. if $s \preceq t$ then $V_s(\alpha) \preceq V_t(\alpha)$,
2. if $s \preceq t$ and $s \in M(\alpha)_i$ then $t \in M(\alpha)_i$, for $i = 0, 1$.

Proof. 1. Firstly, we observe that the functions \neg, \wedge, \vee defined on K (see 3) are isotone. (This can be easily verified by the reader.) Then we suppose $s \preceq t$ and prove $V_s(\alpha) \preceq V_t(\alpha)$ by induction on α . If $\alpha = p_i$ then $V_s(p_i) \preceq V_t(p_i)$ holds, because $s_i \preceq t_i$. If $\alpha = \beta \wedge \gamma$ then by induction hypothesis we have $V_s(\beta) \preceq V_t(\beta)$ and $V_s(\gamma) \preceq V_t(\gamma)$, so the result follows by isotonicity of \wedge with respect to \preceq . The same proof can be given for \vee and \neg .

2. As we have seen in 3, $s \in M(\alpha)_0$ iff $Val_s(\alpha) = 1$. So from our hypothesis $s \preceq t$ and point 1) above, $s \in M(\alpha)_0$ implies $Val_t(\alpha) = 1$ and $t \in M(\alpha)_0$. The same proof holds for $M(\alpha)_1$. ■

Corollary 9 *The function $M : F^* \rightarrow D(K^n)$ is not surjective.*

Proof. For all $s \in K^n$ we have $s \leq \bar{n}$, so if $\bar{n} \in M(\alpha)_i$ then, for all $s \in K^n$, $s \in M(\alpha)_i$, by the above theorem, and $M(\alpha)_i = K^n$. Let (X, Y) be a Boolean partial set, different from (\emptyset, K^n) and (K^n, \emptyset) . Then we have $\bar{n} \in X$ or $\bar{n} \in Y$ and $(X, Y) \neq M(\alpha)$, for all α . ■

References

- [1] G. Birkhoff, *Lattice Theory*, A.M.S. v. 25, 1967.
- [2] B. A. Davey, H. A. Priestley, *Introduction to Lattices and Order*, Cambridge U.P., 1990.
- [3] D. Gillies, *Philosophical Theories of Probability*, Routledge, 2000.
- [4] J. Y. Halpern, *Reasoning about Uncertainty*, The MIT Press, 2003.
- [5] C. Howson, P. Urbach, *Scientific Reasoning: The Bayesian Approach*, Open Court Publishing Company, 1989.
- [6] M. Negri, A probability measure for partial events, *Studia Logica*, vol. 94, 2, pp. 271-290, 2013.
- [7] M. Negri, Partial probability and Kleene Logic, arXiv:1310.6172, 2013.
- [8] J. B. Paris, A note on the Dutch Book method, in *Proceedings of the Second International Symposium on Imprecise Probabilities and their Applications (ISIPTA 2001)*, pp. 301-309. Available at: www.maths.man.ac.uk/DeptWeb/Homepage/jbp/.
- [9] A. Shimony, *Coherence and the Axioms of Confirmation*, *The Journal of Symbolic Logic*, vol. 20, 1 pp. 1-28, 1955.