

NEW WEIGHTED MULTILINEAR OPERATORS AND COMMUTATORS OF HARDY-CESÀRO TYPE

HA DUY HUNG AND LUONG DANG KY

ABSTRACT. A general class of weighted multilinear Hardy-Cesàro operators that acts on the product of Lebesgue spaces and central Morrey spaces. Their sharp bounds are also obtained. In addition, we obtain sufficient and necessary conditions on weight functions so that the commutators of these weighted multilinear Hardy-Cesàro operators (with symbols in central BMO space) are bounded on the product of central Morrey spaces. These results extends known results on multilinear Hardy operators.

1. INTRODUCTION

The Hardy integral inequality and its variants play an important role in various branches of analysis such as approximation theory, differential equations, theory of function spaces etc (see [4, 10, 18, 20, 29, 30] and references therein). Therefore, there are various papers studying Hardy integral inequalities for operator S and its generalizations. The classical Hardy operator, its variants and extensions were appeared in various papers (we refer to [4, 5, 13, 14, 15, 20, 21, 29, 30, 31] for surveys and historical details about these different aspects of the subject). On the other hand, the study of multilinear operators is not motivated by a mere quest to generalize the theory of linear operators but rather by their natural appearance in analysis. Coifman and Meyer in their pioneer work in the 1970s were one of the first to adopt a multilinear point of view in their study of certain singular integral operators, such as the Calderón commutators, paraproducts, and pseudodifferential operators.

Let $\psi : [0, 1] \rightarrow [0, \infty)$ be a measurable function. The weighted Hardy operator U_ψ is defined on all complex-valued measurable functions f on \mathbb{R}^d as

$$(1.1) \quad U_\psi f(x) = \int_0^1 f(tx)\psi(t)dt.$$

When $\psi = 1$, this operator is reduced to the usual Hardy operator S defined by $Sf(x) = \frac{1}{x} \int_0^x f(t)dt$. Results on the boundedness of U_ψ on $L^p(\mathbb{R}^d)$ were

2010 *Mathematics Subject Classification.* 42B35 (46E30, 42B15, 42B30).

Key words and phrases. Hardy-Cesàro operators, Hardy's inequality, weighted Hardy-Littlewood averages.

first proved by Carton-Lebrun and Fosset [5]. Under certain conditions on ψ , the authors [5] found that U_ψ is bounded from $BMO(\mathbb{R}^d)$ into itself. Furthermore, U_ψ commutes with the Hilbert transform in the case $n = 1$ and with a certain Calderón-Zygmund singular integral operator (and thus with the Riesz transform) in the case $n \geq 2$. A deeper extension of the results obtained in [5] was due to Jie Xiao [31].

Theorem 1.1. ([31]) *Let $1 < p < \infty$ and $\psi : [0, 1] \rightarrow [0, \infty)$ be a measurable function. Then, U_ψ is bounded on $L^p(\mathbb{R}^d)$ if and only if*

$$(1.2) \quad \int_0^1 t^{-n/p} \psi(t) dt < \infty.$$

Furthermore,

$$(1.3) \quad \|U_\psi\|_{L^p(\mathbb{R}^d) \rightarrow L^p(\mathbb{R}^d)} = \int_0^1 t^{-n/p} \psi(t) dt < \infty.$$

Theorem 1.1 implies immediately the following celebrated integral inequality, due to Hardy [20]

$$(1.4) \quad \|Sf\|_{L^p(\mathbb{R})} \leq \frac{p}{p-1} \|f\|_{L^p(\mathbb{R})}.$$

For further applications of Theorem 1.1, for examples the sharp bounds of classical Riemann-Liouville integral operator, see [13, 14]. Recently, Chuong and Hung [7] introduced a more general form of U_ψ operator as follows.

$$(1.5) \quad U_{\psi,s}f(x) = \int_0^1 f(s(t)x) \psi(t) dt.$$

Here $\psi : [0, 1] \rightarrow [0, \infty)$, $s : [0, 1] \rightarrow \mathbb{R}$ are measurable functions and f is a measurable complex valued function on \mathbb{R}^d . The authors in [7] obtained sharp bounds of $U_{\psi,s}$ on weighted Lebesgue and BMO spaces, where weights are of homogeneous type. A characterization on weight functions so that the commutator of $U_{\psi,s}$ is bounded on Lebesgue spaces with symbols in BMO . We notice that also in [7], the authors proved the boundedness on Lebesgue spaces of the following operator (see Theorem 3.2 [7])

$$\mathcal{H}_{\psi,s}f(x) = \int_0^\infty f(s(t)x) \psi(t) dt.$$

When $d = 1$ and $s(t) = \frac{1}{t}$, then $\mathcal{H}_{\psi,s}$ reduces to the classical weighted Hausdorff operator (see [25, 26] and references therein).

Very recently, the multilinear version of U_ψ operators was introduced by Z.W. Fu, S.L Gong, S.Z. Lu and W. Yuan [14]. They defined the weighted multilinear Hardy operator as

$$(1.6) \quad U_\psi^m(f_1, \dots, f_m) = \int_{0 < t_1, \dots, t_m < 1} \left(\prod_{i=1}^m f_i(t_i x) \right) \psi(t_1, \dots, t_m) dt_1 \dots dt_m.$$

As showed in [14], when $d = 1$ and $\psi(t_1, \dots, t_m) = \frac{1}{\Gamma(\alpha)|(1-t_1, \dots, 1-t_m)|^{m-\alpha}}$, where $\alpha \in (0; m)$, then $U_\psi^m(f_1, \dots, f_m)(x) = |x|^\alpha I_\alpha^m(f_1, \dots, f_m)(x)$. The operator I_α^m turns out to be the one-sided analogous to the one-dimensional multilinear Riesz operator \mathcal{J}_α^m studied by Kenig and Stein [23], where

$$\mathcal{J}_\alpha^m(f_1, \dots, f_m)(x) = \int_{t_1, \dots, t_m \in \mathbb{R}} \frac{\prod_{k=1}^m f_k(t_k)}{|(x - t_1, \dots, x - t_m)|^{m-\alpha}} d_1 \dots dt_m.$$

In [14], the authors obtain the sharp bounds of U_ψ^m on the product of Lebesgue spaces and central Morrey spaces. They also proved sufficient and necessary conditions of the weight functions so that the commutators of U_ψ^m , with symbols in central BMO spaces, are bounded on the product of central Morrey spaces. In [17], S. Gong, Z. Fu, and B. Ma studied the bounds of U_ψ^m on the product Herz spaces and the product Morrey- Herz spaces. Notice that there is another multilinear version of Hardy operators which was also studied in [15, 28].

Motivated from [7, 14, 13, 17, 21], this paper aims to study the boundedness of a more general multilinear operator of Hardy type as follows.

Definition 1.1. Let $m, n \in \mathbb{N}$, $\psi : [0, 1]^n \rightarrow [0, \infty)$, $s_1, \dots, s_m : [0, 1]^n \rightarrow \mathbb{R}$. Given $f_1, \dots, f_m : \mathbb{R}^d \rightarrow \mathbb{C}$ be measurable functions, we define the weighted multilinear Hardy-Cesàro operator $U_{\psi, \vec{s}}^{m,n}$ by

$$(1.7) \quad U_{\psi, \vec{s}}^{m,n}(f_1, \dots, f_m)(x) := \int_{[0,1]^n} \left(\prod_{k=1}^m f_k(s_k(t)x) \right) \psi(t) dt,$$

where $\vec{s} = (s_1, \dots, s_m)$.

A multilinear version of $\mathcal{H}_{\psi, s}$ can be defined as

$$(1.8) \quad \mathcal{H}_{\psi, \vec{s}}^{m,n}(f_1, \dots, f_m)(x) := \int_{\mathbb{R}_+^n} \left(\prod_{k=1}^m f_k(s_k(t)x) \right) \psi(t) dt,$$

where \mathbb{R}_+ is the set of all positive real numbers.

It is obviously that, when $n = m$ and $s_k(t) = t_k$, $U_{\psi, \vec{s}}^{m,n}$ is reduced to U_ψ^m as defined in (1.5). The main aim of the paper is to establish the sharp bounds

of $U_{\psi, \vec{s}}^{m,n}$ and $\mathcal{H}_{\psi, \vec{s}}^{m,n}$ on the product of weighted Lebesgue spaces and weighted central Morrey spaces, with weights of homogeneous types. In addition, we prove sufficient and necessary conditions of the weight functions so that commutators of such weighted multilinear Hardy-Cesàro operators (with symbols in λ -central BMO space) are bounded on the product of central Morrey spaces. Since $U_{\psi, \vec{s}}^{m,n}$ is trivially more general than known operators U_ψ , U_ψ^m , our results can be used to recover main results of [7, 13, 14, 31].

Throughout the whole paper, the letter C will indicate an absolute constant, probably different at different occurrences. With χ_E we will denote the characteristic function of a set E and $B(x, r)$ will be a ball centered at x with radius r . With $|A|$ we will denote the Lebesgue measure of a measurable set E , and E^c will be the set $\mathbb{R}^d \setminus E$. With $\omega(E)$ we will denote by $\int_E \omega(x)dx$ and S_d will be the unit ball $\{x \in \mathbb{R}^d : |x| = 1\}$.

2. NOTATIONS AND DEFINITIONS

Throughout this paper, $\omega(x)$ will be denote a nonnegative measurable function on \mathbb{R}^d . Let us recall that a measurable function f belongs to $L_\omega^p(\mathbb{R}^d)$ if

$$(2.1) \quad \|f\|_{L_\omega^p} = \left(\int_{\mathbb{R}^d} |f(x)|^p \omega(x) dx \right)^{1/p} < \infty.$$

The weighted *BMO* space $BMO(\omega)$ is defined as the set of all functions f , which are of bounded mean oscillation with weight ω , that is,

$$(2.2) \quad \|f\|_{BMO(\omega)} = \sup_B \frac{1}{\omega(B)} \int_B |f(x) - f_{B,\omega}| \omega(x) dx < \infty,$$

where supremum is taken over all the balls $B \subset \mathbb{R}^d$. Here $f_{B,\omega}$ is the mean value of f on B with weight ω :

$$f_{B,\omega} = \frac{1}{\omega(B)} \int_B f(x) \omega(x) dx.$$

The case $\omega \equiv 1$ of (2.2) corresponds to the class of functions of bounded mean oscillation of F. John and L. Nirenberg [22]. We observe that $L^\infty(\mathbb{R}^d) \subset BMO(\omega)$.

Next we recall the definition of Morrey spaces. It is well-known that Morrey spaces are useful to study the local behavior of solutions to second-order elliptic partial differential equations and the boundedness of Hardy-Littlewood maximal operator, the fractional integral operators, singular integral operators (see [1, 6, 24]). We notice that the weighted Morrey spaces were first

introduced by Komori and Shirai [24], where they used them to study the boundedness of some important classical operators in harmonic analysis like as Hardy-Littlewood maximal operator, Calderón-Zygmund operators.

Definition 2.1. Let $\lambda \in \mathbb{R}$, $1 \leq p < \infty$ and ω be an weight function. The weighted Morrey space $L_{\omega}^{p,\lambda}(\mathbb{R}^d)$ is defined by the set of all locally p -integrable functions f satisfying

$$(2.3) \quad \|f\|_{L_{\omega}^{p,\lambda}(\mathbb{R}^d)} = \sup_{a \in \mathbb{R}^d, R > 0} \left(\frac{1}{\omega(B(a, R))^{1+\lambda p}} \int_{B(0, R)} |f(x)|^p \omega(x) dx \right)^{1/p} < \infty.$$

The spaces of bounded central mean oscillation $CMO^q(\mathbb{R}^d)$ appears naturally when considering the dual spaces of the homogeneous Herz type Hardy spaces and were introduced by Lu and Yang (see [27]). The relationships between central BMO spaces and Morrey spaces were studied by Alvarez, Guzmán-Partida and Lakey [2]. Furthermore, they introduced λ -central BMO spaces and central Morrey spaces as follows.

Definition 2.2. Let $\lambda \in \mathbb{R}$ and $1 < p < \infty$. The weighted central Morrey space $\dot{B}_{\omega}^{p,\lambda}(\mathbb{R}^d)$ is defined by the set of all locally p -integrable functions f satisfying

$$(2.4) \quad \|f\|_{\dot{B}_{\omega}^{p,\lambda}(\mathbb{R}^d)} = \sup_{R > 0} \left(\frac{1}{\omega(B(0, R))^{1+\lambda p}} \int_{B(0, R)} |f(x)|^p \omega(x) dx \right)^{1/p} < \infty.$$

Obviously, $\dot{B}_{\omega}^{p,\lambda}(\mathbb{R}^d)$ is a Banach space and one can easily check that $\dot{B}_{\omega}^{p,\lambda}(\mathbb{R}^d) = \{0\}$ if $\lambda < -\frac{1}{q}$. Similar to the classical Morrey space, we only consider the case $-1/p \leq \lambda < 0$.

Definition 2.3. Let $\lambda < \frac{1}{n}$ and $1 < q < \infty$ be two real numbers. The weighted space $CMO_{\omega}^{q,\lambda}(\mathbb{R}^d)$ is defined as the set of all function $f \in L_{\omega,loc}^q(\mathbb{R}^d)$ such that

$$(2.5) \quad \|f\|_{CMO_{\omega}^{q,\lambda}(\mathbb{R}^d)} = \sup_{R > 0} \left(\frac{1}{\omega(B(0, R))^{1+\lambda q}} \int_{B(0, R)} |f(x) - f_{B(0, R), \omega}|^q \omega(x) dx \right)^{1/q} < \infty.$$

Let ρ be the measure on $(0, \infty)$ so that $\rho(E) = \int_E r^{d-1} dr$ and the map $\Phi(x) = \left(|x|, \frac{x}{|x|}\right)$. Then there exists an unique Borel measure σ on S_n such that $\rho \times \sigma$ is the Borel measure induced by Φ from Lebesgue measure on \mathbb{R}^d ($d > 1$). (see [12, page 78] for more details). In one dimension case, it is conventional that $\int_{S_d} \omega(x) d\sigma(x)$ refers to $2\omega(1)$.

Definition 2.4. Let α be a real number. Let \mathcal{W}_α be the set of all functions ω on \mathbb{R}^d , which are measurable, $\omega(x) > 0$ for almost everywhere $x \in \mathbb{R}^d$, $0 < \int_{S_d} \omega(y) d\sigma(y) < \infty$, and are absolutely homogeneous of degree α , that is $\omega(tx) = |t|^\alpha \omega(x)$, for all $t \in \mathbb{R} \setminus \{0\}$, $x \in \mathbb{R}^d$.

Let us describe some typical examples and properties of \mathcal{W}_α . Note that, a weight $\omega \in \mathcal{W}_\alpha$ may not need to belong to $L^1_{\text{loc}}(\mathbb{R}^d)$. In fact, we observe that if $\omega \in \mathcal{W}_\alpha$, then $\omega \in L^1_{\text{loc}}(\mathbb{R}^d)$ if and only if $\alpha > -d$. If $d = 1$, then $\omega(x) = c|x|^\alpha$, for some positive constant c . For $d \geq 1$ and $\alpha \neq 0$, $\omega(x) = |x|^\alpha$ is in \mathcal{W}_α . If ω_1, ω_2 is in \mathcal{W}_α , so is $\theta\omega_1 + \lambda\omega_2$ for all $\theta, \lambda > 0$. There are many other examples in case $d > 1$ and $\alpha \neq 0$, namely $\omega(x_1, \dots, x_d) = |x_1|^\alpha$. In case $d > 1$ and $\alpha = 0$, we can construct a non-trivial example as the following: let ϕ be any positive, even and locally integrable function on $S_d = \{x \in \mathbb{R}^d : |x| = 1\}$, then

$$\omega(x) = \begin{cases} \phi\left(\frac{x}{|x|}\right) & \text{if } x \neq 0, \\ 0 & \text{if } x = 0, \end{cases}$$

is in \mathcal{W}_0 .

3. SHARP BOUNDEDNESS OF $U_{\psi, \vec{s}}^{m,n}$ ON THE PRODUCT OF WEIGHTED LEBESGUE SPACES

Before stating our results we give some notations and definitions.

Definition 3.1. For m exponents $1 \leq p_j < \infty$, $j = 1, \dots, m$ and $\alpha_1, \dots, \alpha_m > -d$ we will often write p for the number given by $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$ and

$$(3.1) \quad \alpha := \frac{p\alpha_1}{p_1} + \dots + \frac{p\alpha_m}{p_m}.$$

Definition 3.2. Given $\omega_k \in \mathcal{W}_{\alpha_k}$, $k = 1, \dots, m$, set

$$(3.2) \quad \omega(x) := \prod_{k=1}^m \omega_k^{p/p_k}(x).$$

It is obvious that $\omega \in \mathcal{W}_\alpha$. We say that $(\omega_1, \dots, \omega_m)$ satisfies the $\mathcal{W}_{\vec{\alpha}}$ condition if

$$(3.3) \quad \omega(S_d) \geq \prod_{k=1}^m \omega_k(S_d)^{\frac{p}{p_k}}.$$

Notice that the weights ω as defined in (3.2) were used in [19] to obtain multilinear extrapolation results. On the other hand, Condition (3.3) holds for power weights. In fact, (3.3) becomes to equality for $\omega_k(x) = |x|^{\alpha_k}$, $k = 1, \dots, m$.

First we will show the following result which can be viewed as an extension of Theorem 1.1 to the multilinear case.

Theorem 3.1. *Let $s_1, \dots, s_m : [0, 1]^n \rightarrow \mathbb{R}$ be measurable functions such that for every $k = 1, \dots, m$ then $|s_k(t_1, \dots, t_n)| \geq \min\{t_1^\beta, \dots, t_n^\beta\}$ almost everywhere $(t_1, \dots, t_n) \in [0, 1]^n$, $k = 1, \dots, m$, for some $\beta > 0$. Let $1 \leq p_k < \infty$, $k = 1, \dots, m$ and $\vec{s} = (s_1, \dots, s_m)$. Assume that $(\omega_1, \dots, \omega_m)$ satisfies $\mathcal{W}_{\vec{\alpha}}$ condition. Then $U_{\psi, \vec{s}}^{m, n}$ is bounded from $L_{\omega_1}^{p_1}(\mathbb{R}^d) \times \dots \times L_{\omega_m}^{p_m}(\mathbb{R}^d)$ to $L_{\omega}^p(\mathbb{R}^d)$ if and only if*

$$(3.4) \quad \mathcal{A} = \int_{[0,1]^n} \left(\prod_{k=1}^m |s_k(t)|^{-\frac{d+\alpha_k}{p_k}} \right) \psi(t) dt < \infty.$$

Furthermore,

$$(3.5) \quad \left\| U_{\psi, \vec{s}}^{m, n} \right\|_{L_{\omega_1}^{p_1}(\mathbb{R}^d) \times \dots \times L_{\omega_m}^{p_m}(\mathbb{R}^d) \rightarrow L_{\omega}^p(\mathbb{R}^d)} = \mathcal{A}.$$

Proof. First we take $\alpha = \sum_{k=1}^m \frac{p\alpha_k}{p_k}$ as in (3.1), then it is trivial that $\omega \in \mathcal{W}_{\alpha}$.

Suppose that \mathcal{A} is finite. Let $f_k \in L_{\omega_k}^{p_k}(\mathbb{R}^d)$. Applying Minkowski's inequality, Hölder's inequality and change of variable, we have

$$\begin{aligned} \left\| U_{\psi, \vec{s}}^{m, n}(f_1, \dots, f_m) \right\|_{L_{\omega}^p(\mathbb{R}^d)} &\leq \left(\int_{\mathbb{R}^d} \left(\int_{[0,1]^n} \prod_{k=1}^m |f_k(s_k(t)x)| \psi(t) dt \right)^p \omega(x) dx \right)^{1/p} \\ &\leq \int_{[0,1]^n} \left(\int_{\mathbb{R}^d} \prod_{k=1}^m |f_k(s_k(t)x)|^p \omega(x) dx \right)^{1/p} \psi(t) dt \\ &\leq \int_{[0,1]^n} \prod_{k=1}^m \left(\int_{\mathbb{R}^d} |f_k(s_k(t)x)|^{p_k} \omega_k(x) dx \right)^{1/p_k} \psi(t) dt \\ &= \mathcal{A} \cdot \prod_{k=1}^m \|f_k\|_{p_k, \omega_k}. \end{aligned}$$

The last inequality implies that $U_{\psi, \vec{s}}^{m, n}$ is bounded from $L_{\omega_1}^{p_1}(\mathbb{R}^d) \times \dots \times L_{\omega_m}^{p_m}(\mathbb{R}^d)$ to $L_{\omega}^p(\mathbb{R}^d)$ and

$$(3.6) \quad \left\| U_{\psi, \vec{s}}^{m, n} \right\|_{L_{\omega_1}^{p_1}(\mathbb{R}^d) \times \dots \times L_{\omega_m}^{p_m}(\mathbb{R}^d) \rightarrow L_{\omega}^p(\mathbb{R}^d)} \leq \mathcal{A}.$$

In order to prove the converse of the theorem, we first need the following lemma.

Lemma 3.1. *Let $w \in \mathcal{W}_\alpha$, $\alpha > -d$ and $\varepsilon > 0$. Then the function*

$$f_{p,\varepsilon}(x) = \begin{cases} 0 & \text{if } |x| \leq 1 \\ |x|^{-\frac{d+\alpha}{p}-\varepsilon} & \text{if } |x| \geq 1, \end{cases}$$

belongs to $L_w^p(\mathbb{R}^d)$ and $\|f_{p,\varepsilon}\|_{p,w} = \left(\frac{w(S_d)}{p\varepsilon}\right)^{1/p}$.

Since the proof of the lemma is straightforward, we omit it. Now we shall prove the theorem. Let ε be an arbitrary positive number and for each $k = 1, \dots, m$ we set $\varepsilon_k = \frac{p\varepsilon}{p_k}$ and

$$f_{p_k,\varepsilon_k}(x) = \begin{cases} 0 & \text{if } |x| \leq 1 \\ |x|^{-\frac{d+\alpha_k}{p_k}-\varepsilon_k} & \text{if } |x| \geq 1. \end{cases}$$

Lemma 3.1 implies that $f_{p_k,\varepsilon_k} \in L_{\omega_k}^{p_k}(\mathbb{R}^d)$ and

$$\|f_{p_k,\varepsilon_k}\|_{p_k,\omega_k} = \left(\frac{\omega_k(S_d)}{p_k\varepsilon_k}\right)^{1/p_k} = \left(\frac{\omega_k(S_d)}{p\varepsilon}\right)^{1/p_k} > 0,$$

for each $k = 1, \dots, m$. For each $x \in \mathbb{R}^d$ which $|x| \geq 1$, let

$$S_x = \bigcap_{k=1}^m \{t \in [0, 1]^n : |s_k(t)x| > 1\}.$$

From the assumption $|s_k(t_1, \dots, t_n)| \geq \min\{t_1^\beta, \dots, t_n^\beta\}$ a.e $t = (t_1, \dots, t_n) \in [0, 1]^n$, there exists a null subset E of $[0, 1]^n$ so that S_x contains $[1/|x|^{1/\beta}, 1]^n \setminus E$.

From (2.1), we have

$$\begin{aligned}
& \|U_{\psi, \vec{s}}^{m,n}(f_{p_1, \varepsilon_1}, \dots, f_{p_m, \varepsilon_m})\|_{L_\omega^p(\mathbb{R}^d)}^p = \int_{\mathbb{R}^d} \left| \int_{[0,1]^n} \left(\prod_{k=1}^m f_k(s_k(t)x) \right) \psi(t) dt \right|^p \omega(x) dx \\
&= \int_{\mathbb{R}^d} \prod_{k=1}^m |x|^{-\frac{d+\alpha_k}{p_k} - \varepsilon_k} \omega(x) \left| \int_{S_x} \prod_{k=1}^m |s_k(t)|^{-\frac{d+\alpha_k}{p_k} - \varepsilon_k} \psi(t) dt \right|^p dx \\
&\geq \int_{|x| \geq \varepsilon^{-\beta}} |x|^{-(d+\alpha+\varepsilon p)} \omega(x) \left| \int_{[\varepsilon,1]^n} \prod_{k=1}^m |s_k(t)|^{-\frac{d+\alpha_k}{p_k} - \varepsilon_k} \psi(t) dt \right|^p dx \\
&= \varepsilon^{p\beta\varepsilon} \left(\int_{|x| \geq 1} |x|^{-(d+\alpha+\varepsilon p)} \omega(x) dx \right) \left| \int_{[\varepsilon,1]^n} \prod_{k=1}^m |s_k(t)|^{-\frac{d+\alpha_k}{p_k} - \varepsilon_k} \psi(t) dt \right|^p \\
&= \varepsilon^{p\beta\varepsilon} \left(\frac{\omega(S_d)}{\varepsilon^p} \right) \left| \int_{[\varepsilon,1]^n} \prod_{k=1}^m |s_k(t)|^{-\frac{d+\alpha_k}{p_k} - \varepsilon_k} \psi(t) dt \right|^p \\
&\geq \varepsilon^{p\beta\varepsilon} \prod_{k=1}^m \|f_{p_k, \varepsilon_k}\|_{p_k, \varepsilon_k}^{p/p_k} \left| \int_{[\varepsilon,1]^n} \prod_{k=1}^m |s_k(t)|^{-\frac{d+\alpha_k}{p_k} - \varepsilon_k} \psi(t) dt \right|^p.
\end{aligned}$$

This implies that

$$\left\| U_{\psi, \vec{s}}^{m,n} \right\|_{L_{\omega_1}^{p_1}(\mathbb{R}^d) \times \dots \times L_{\omega_m}^{p_m}(\mathbb{R}^d) \rightarrow L_\omega^p(\mathbb{R}^d)} \geq \varepsilon^{\beta\varepsilon} \cdot \left| \int_{[\varepsilon,1]^n} \prod_{k=1}^m |s_k(t)|^{-\frac{d+\alpha_k}{p_k} - \varepsilon_k} \psi(t) dt \right|.$$

Notice that

$$|s_k(t)|^{-\varepsilon} \leq \min\{t_1, \dots, t_n\}^{-\varepsilon\beta} \leq \varepsilon^{-\varepsilon\beta} \rightarrow 1 \quad \text{when } \varepsilon \rightarrow 0^+,$$

Thus, letting $\varepsilon \rightarrow 0^+$ and by Lebesgue's dominated convergence theorem, we obtain

$$(3.7) \quad \left\| U_{\psi, \vec{s}}^{m,n} \right\|_{L_{\omega_1}^{p_1}(\mathbb{R}^d) \times \dots \times L_{\omega_m}^{p_m}(\mathbb{R}^d) \rightarrow L_\omega^p(\mathbb{R}^d)} \geq \mathcal{A}.$$

Combine (3.6) and (3.7), we obtain the result. \square

Analogous to the proof of Theorem 3.1, one can prove the following result.

Theorem 3.2. *Let $s_1, \dots, s_m : [0, 1]^n \rightarrow \mathbb{R}$ be measurable functions such that for every $k = 1, \dots, m$ then $|s_k(t_1, \dots, t_n)| \geq \min\{t_1^\beta, \dots, t_n^\beta\}$ almost everywhere $(t_1, \dots, t_n) \in [0, 1]^n$, $k = 1, \dots, m$, for some $\beta > 0$. Let $1 \leq p_k < \infty$, $k = 1, \dots, m$ and $\vec{s} = (s_1, \dots, s_m)$. Assume that $(\omega_1, \dots, \omega_m)$ satisfies $\mathcal{W}_{\vec{\alpha}}$ condition. Then $\mathcal{H}_{\psi, \vec{s}}^{m, n}$ is bounded from $L_{\omega_1}^{p_1}(\mathbb{R}^d) \times \dots \times L_{\omega_m}^{p_m}(\mathbb{R}^d)$ to $L_{\omega}^p(\mathbb{R}^d)$ if and only if*

$$(3.8) \quad \mathcal{A}_* = \int_{\mathbb{R}_+^n} \left(\prod_{k=1}^m |s_k(t)|^{-\frac{d+\alpha_k}{p_k}} \right) \psi(t) dt < \infty.$$

Furthermore,

$$(3.9) \quad \left\| \mathcal{H}_{\psi, \vec{s}}^{m, n} \right\|_{L_{\omega_1}^{p_1}(\mathbb{R}^d) \times \dots \times L_{\omega_m}^{p_m}(\mathbb{R}^d) \rightarrow L_{\omega}^p(\mathbb{R}^d)} = \mathcal{A}_*.$$

Theorem 3.3. *Let $1 \leq p, p_k < \infty$, $\lambda, \alpha_k, \lambda_k$ be real numbers such that $-\frac{1}{p_k} \leq \lambda_k < 0$, $k = 1, \dots, m$, $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$, and $\lambda = \frac{d+\alpha_1}{d+\alpha} \lambda_1 + \dots + \frac{d+\alpha_m}{d+\alpha} \lambda_m$. Let $\omega_k \in \mathcal{W}_{\alpha_k}$ for $k = 1, \dots, m$.*

(i) *If in addition*

$$(3.10) \quad \left(\frac{\omega(S_d)}{d+\alpha} \right)^{\frac{1+\lambda p}{p}} \geq \prod_{k=1}^m \left(\frac{\omega_k(S_d)}{d+\alpha_k} \right)^{\frac{1+\lambda_k p_k}{p_k}}$$

and

$$(3.11) \quad \mathcal{B} = \int_{[0, 1]^n} \left(\prod_{k=1}^m |s_k(t)|^{-\frac{(d+\alpha_k)\lambda_k}{p_k}} \right) \psi(t) dt < \infty,$$

then $U_{\psi, \vec{s}}^{m, n}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d)$ to $\dot{B}_{\omega}^{p, \lambda}(\mathbb{R}^d)$.

Furthermore, the operator norm of $U_{\psi, \vec{s}}^{m, n}$ not more than \mathcal{B} .

(ii) *Conversely, if*

$$(3.12) \quad \left(\frac{\omega(S_d)}{d+\alpha} \right)^{\lambda} (1+\lambda p)^{1/p} \leq \prod_{k=1}^m \left(\frac{\omega_k(S_d)}{d+\alpha_k} \right)^{\lambda_k} (1+\lambda_k p_k)^{1/p_k}$$

and $U_{\psi, \vec{s}}^{m, n}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d)$ to $\dot{B}_{\omega}^{p, \lambda}(\mathbb{R}^d)$, then \mathcal{B} is finite. Furthermore, we have

$$(3.13) \quad \left\| U_{\psi, \vec{s}}^{m, n} \right\|_{\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d) \rightarrow \dot{B}_{\omega}^{p, \lambda}(\mathbb{R}^d)} \geq \mathcal{B}.$$

Let $\omega_k \equiv 1$, we have that $\alpha_k = 0$ thus (3.15) becomes equality and (3.17) holds only when $\lambda_1 p_1 = \dots = \lambda_m p_m$. Hence, we obtain Theorem 2.1 in [14].

Proof. Since $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$, by Minkowski's inequality and Hölder's inequality, we see that, for all balls $B = B(0, R)$,

$$\begin{aligned}
& \left(\frac{1}{\omega(B)^{1+\lambda p}} \int_B \left| U_{\psi, \vec{s}}^{m,n}(f_1, \dots, f_m)(x) \right|^p \omega(x) dx \right)^{1/p} \\
& \leq \int_{[0,1]^n} \left(\frac{1}{\omega(B)^{1+\lambda p}} \int_B \left| \prod_{k=1}^m f_k(s_k(t)x) \right|^p \omega(x) dx \right)^{1/p} \psi(t) dt \\
& \leq \int_{[0,1]^n} \prod_{k=1}^m \left(\frac{1}{\omega_k(B)^{1+\lambda_k p_k}} \int_B |f_k(s_k(t)x)|^{p_k} \omega_k(x) dx \right)^{1/p_k} \psi(t) dt \\
& \leq \int_{[0,1]^n} \prod_{k=1}^m \left(\frac{1}{|s_k(t)B|^{1+\lambda_k p_k}} \int_B |f_k(y)|^{p_k} \omega_k(y) dy \right)^{1/p_k} \left(\prod_{k=1}^m |s_k(t)|^{(d+\alpha_k)\lambda_k} \right) \psi(t) dt \\
& \leq \prod_{k=1}^m \|f_k\|_{\dot{B}_{\omega_k}^{p_k, \lambda_k}(\mathbb{R}^d)} \cdot \mathcal{B}.
\end{aligned}$$

This means that

$$(3.14) \quad \left\| U_{\psi, \vec{s}}^{m,n} \right\|_{\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d) \rightarrow \dot{B}_{\omega}^{p, \lambda}(\mathbb{R}^d)} \leq \mathcal{B}.$$

Now we assume that (3.17) holds and $U_{\psi, s}^{m,n}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}$ to $\dot{B}_{\omega}^{p, \lambda}(\mathbb{R}^d)$. Let $f_k(x) = |x|^{(d+\alpha_k)\lambda_k}$, then an elementary computation shows that $f_k \in \dot{B}_{\omega_k}^{p_k, \lambda_k}(\mathbb{R}^d)$ and

$$\|f_k\|_{\dot{B}_{\omega_k}^{p_k, \lambda_k}(\mathbb{R}^d)} = \left(\frac{d + \alpha_k}{\omega(S_d)} \right)^{\lambda_k} (1 + \lambda_k p_k)^{-1/p_k}.$$

Thus,

$$\begin{aligned}
\prod_{k=1}^m \|f_k\|_{\dot{B}_{\omega_k}^{p_k, \lambda_k}(\mathbb{R}^d)} &= \prod_{k=1}^m \left(\frac{d + \alpha_k}{\omega(S_d)} \right)^{\lambda_k} (1 + \lambda_k p_k)^{-1/p_k} \\
&\leq \left(\frac{d + \alpha}{\omega(S_d)} \right)^{\lambda} (1 + \lambda p)^{-1/p}.
\end{aligned}$$

This leads us to

$$\|U_{\psi, \vec{s}}^{m,n}(f_1, \dots, f_m)\|_{\dot{B}_\omega^{p,\lambda}(\mathbb{R}^d)} = \mathcal{B} \cdot \left(\frac{d+\alpha}{\omega(S_d)} \right)^\lambda (1+\lambda p)^{-1/p} \geq \mathcal{B} \cdot \prod_{k=1}^m \|f_k\|_{\dot{B}_{\omega_k}^{p_k, \lambda_k}(\mathbb{R}^d)}.$$

Therefore,

$$\left\| U_{\psi, \vec{s}}^{m,n} \right\|_{\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d) \rightarrow L_\omega^p(\mathbb{R}^d)} \geq \mathcal{B}.$$

□

From the proof of Theorem 3.4, we also obtain the similar result to $\mathcal{H}_{\psi, \vec{s}}^{m,n}$ operator.

Theorem 3.4. *Let $1 \leq p, p_k < \infty$, $\lambda, \alpha_k, \lambda_k$ be real numbers such that $-\frac{1}{p_k} \leq \lambda_k < 0$, $k = 1, \dots, m$, $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$, and $\lambda = \frac{d+\alpha_1}{d+\alpha} \lambda_1 + \dots + \frac{d+\alpha_m}{d+\alpha} \lambda_m$. Let $\omega_k \in \mathcal{W}_{\alpha_k}$ for $k = 1, \dots, m$.*

(i) *If in addition*

$$(3.15) \quad \left(\frac{\omega(S_d)}{d+\alpha} \right)^{\frac{1+\lambda p}{p}} \geq \prod_{k=1}^m \left(\frac{\omega_k(S_d)}{d+\alpha_k} \right)^{\frac{1+\lambda_k p_k}{p_k}}$$

and

$$(3.16) \quad \mathcal{B}_\star = \int_{\mathbb{R}_+^n} \left(\prod_{k=1}^m |s_k(t)|^{-\frac{(d+\alpha_k)\lambda_k}{p_k}} \right) \psi(t) dt < \infty,$$

then $\mathcal{H}_{\psi, \vec{s}}^{m,n}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d)$ to $\dot{B}_\omega^{p, \lambda}(\mathbb{R}^d)$.

Furthermore, the operator norm of $\mathcal{H}_{\psi, \vec{s}}^{m,n}$ not more than \mathcal{B}_\star .

(ii) *Conversely, if*

$$(3.17) \quad \left(\frac{\omega(S_d)}{d+\alpha} \right)^\lambda (1+\lambda p)^{1/p} \leq \prod_{k=1}^m \left(\frac{\omega(S_d)}{d+\alpha_k} \right)^{\lambda_k} (1+\lambda_k p_k)^{1/p_k}$$

and $\mathcal{H}_{\psi, \vec{s}}^{m,n}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d)$ to $\dot{B}_\omega^{p, \lambda}(\mathbb{R}^d)$, then \mathcal{B}_\star is finite. Furthermore, we have

$$(3.18) \quad \left\| \mathcal{H}_{\psi, \vec{s}}^{m,n} \right\|_{\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d) \rightarrow L_\omega^p(\mathbb{R}^d)} \geq \mathcal{B}_\star.$$

4. COMMUTATORS OF WEIGHTED MULTILINEAR HARDY-CESÀRO OPERATOR

We use some analogous tools to study a second set of problems related now to multilinear versions of the commutators of Coifman, Rochberg and Weiss [9] In this section, we consider the sharp estimates of the multilinear commutator generated by $U_{\psi, \vec{s}}^{m,n}$ with symbols in $CMO^q(\mathbb{R}^d)$.

Definition 4.1. Let $m, n \in \mathbb{N}$, $\psi : [0, 1]^n \rightarrow [0, \infty)$, $s_1, \dots, s_m : [0, 1]^n \rightarrow \mathbb{R}$, b_1, \dots, b_m be locally integrable functions on \mathbb{R}^d and $f_1, \dots, f_m : \mathbb{R}^d \rightarrow \mathbb{C}$ be measurable functions. The commutator of weighted multilinear Hardy-Cesàro operator $U_{\psi, \vec{s}}^{m, n}$ is defined as

$$(4.1) \quad U_{\psi, \vec{s}}^{m, n, \vec{b}}(f_1, \dots, f_m)(x) := \int_{[0, 1]^n} \left(\prod_{k=1}^m f_k(s_k(t)x) \right) \left(\prod_{k=1}^m (b_k(x) - b_k(s_k(t)x)) \right) \psi(t) dt.$$

In what follows, we set

$$(4.2) \quad \mathcal{C} = \int_{[0, 1]^n} \left(\prod_{k=1}^m |s_k(t)|^{(d+\alpha_k)\lambda_k} \right) \psi(t) dt.$$

$$(4.3) \quad \mathcal{D} = \int_{[0, 1]^n} \left(\prod_{k=1}^m |s_k(t)|^{(d+\alpha_k)\lambda_k} \right) \left(\prod_{k=1}^m |\log |s_k(t)|| \right) \psi(t) dt.$$

Theorem 4.1. Let $1 < p < p_k < \infty$, $1 < q_k < \infty$, $-\frac{1}{p_k} < \lambda_k < 0$, $k = 1, \dots, m$ such that

$$\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m} + \frac{1}{q_1} + \dots + \frac{1}{q_m}$$

and $\lambda = \lambda_1 + \dots + \lambda_m$.

(i) If both \mathcal{C} and \mathcal{D} are finite then for any $b = (b_1, \dots, b_m) \in CMO_{\omega_1}^{q_1} \times CMO_{\omega_m}^{q_m}$ then $U_{\psi, \vec{s}}^{m, n, \vec{b}}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d)$ to $\dot{B}_{\omega}^{p, \lambda}(\mathbb{R}^d)$.

(ii) If for any $b = (b_1, \dots, b_m) \in CMO_{\omega_1}^{q_1} \times \dots \times CMO_{\omega_m}^{q_m}$, $U_{\psi, \vec{s}}^{m, n, \vec{b}}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d)$ to $\dot{B}_{\omega}^{p, \lambda}(\mathbb{R}^d)$, then \mathcal{D} is finite.

We note here that \mathcal{D} is finite is not enough to imply \mathcal{C} is finite (see [13, 14]). But it is easy to see that, if we assume in addition that for each $k = 1, \dots, m$ such that $|s_k(t)| \geq c > 1$ for all $t \in [0, 1]^n$ or $|s_k(t)| \leq c < 1$ for all $t \in [0, 1]^n$, then \mathcal{C} is finite if and only if \mathcal{D} is finite. Thus, Theorem 4.1 implies immediately that

Corollary 4.1. Let $1 < p < p_k < \infty$, $1 < q_k < \infty$, $-\frac{1}{p_k} < \lambda_k < 0$, $k = 1, \dots, m$ such that

$$\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m} + \frac{1}{q_1} + \dots + \frac{1}{q_m}$$

and $\lambda = \lambda_1 + \dots + \lambda_m$. Furthermore, suppose that for each $k = 1, \dots, m$ then $|s_k(t)| \geq c > 1$ for all $t \in [0, 1]^n$ or $|s_k(t)| \leq c < 1$ for all $t \in [0, 1]^n$, for each $k = 1, \dots, m$. Then, $U_{\psi, \vec{s}}^{m, n, \vec{b}}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dots \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d)$ to $\dot{B}_{\omega}^{p, \lambda}(\mathbb{R}^d)$ if and only if \mathcal{C} is finite.

Now we will give the proof for Theorem 4.1.

Proof. First we assume that \mathcal{C}_m is finite. We shall give a details analysis for the case $m = 2$ and by similarity, the general case is proved in the same way. We denote $B = B(0, R)$ for short. Let $\vec{b} = (b_1, b_2) \in CMO_{\omega_1}^{q_1}(\mathbb{R}^d) \times CMO_{\omega_2}^{q_2}(\mathbb{R}^d)$. By Minkowski's inequality, we have

$$\begin{aligned} & \left(\frac{1}{\omega(B)} \int_B \left| U_{\psi, s}^{2, n, b}(f_1, f_2)(x) \right|^p \omega(x) dx \right)^{1/p} \\ & \leq \left(\frac{1}{\omega(B)} \int_B \left(\int_{[0,1]^n} \left(\prod_{k=1}^2 |f_k(s_k(t)x)| \right) \left(\prod_{k=1}^2 |b_k(x) - b_k(s_k(t)x)| \right) \psi(t) dt \right)^p \omega(x) dx \right)^{1/p} \\ & \leq \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\prod_{k=1}^2 |f_k(s_k(t)x)| \cdot \prod_{k=1}^2 |b_k(x) - b_k(s_k(t)x)| \omega(x) \right)^p dx \right)^{1/p} \psi(t) dt \\ & =: I. \end{aligned}$$

For any $x_i, y_i, z_i, t_i \in \mathbb{C}$ with $i = 1, 2$, we have the following elementary inequality

$$\begin{aligned} & \prod_{i=1}^2 |x_i - y_i| \leq \prod_{i=1}^2 |x_i - z_i| + \prod_{i=1}^2 |y_i - t_i| + \prod_{i=1}^2 |z_i - t_i| \\ & + (|(x_1 - z_1)(z_2 - t_2)| + |(x_2 - z_2)(z_1 - t_1)|) + (|(x_1 - z_1)(y_2 - t_2)| + |(x_2 - z_2)(y_1 - t_1)|) \\ & + (|(z_1 - t_1)(y_2 - t_2)| + |(z_2 - t_2)(y_1 - t_1)|). \end{aligned}$$

It is convenient to denote by $b_{i, \omega, B}$ the integrals $\int_B \frac{1}{\omega(B)} b_i(x) \omega(x) dx$ for $i = 1, 2$. Now applying the inequality with $x_i = b_i(x)$, $y_i = b_i(s_i(t)x)$, $z_i = b_{i, B}$, $t_i = b_{i, s_i(t)B}$ and using Minkowski's inequality, we get that

$$I \leq I_1 + I_2 + I_3 + I_4 + I_5 + I_6,$$

where, if we set $\bar{f}(x) = \prod_{k=1}^2 |f_k(s_k(t)x)|$, then

$$\begin{aligned} I_1 &= \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\bar{f}(x) \cdot \prod_{k=1}^2 |b_k(x) - b_{k, \omega_k, B}| \right)^p \omega(x) dx \right)^{1/p} \psi(t) dt, \\ I_2 &= \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\bar{f}(x) \cdot \prod_{k=1}^2 |b_k(s_k(t)x) - b_{k, \omega_k, s_k(t)B}| \right)^p \omega(x) dx \right)^{1/p} \psi(t) dt, \end{aligned}$$

$$I_3 = \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\bar{f}(x) \cdot \prod_{k=1}^2 |b_{k,\omega_k,B} - b_{k,\omega_k,s_k(t)B}| \right)^p \omega(x) dx \right)^{1/p} \psi(t) dt,$$

$$I_4 = \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\bar{f}(x) \cdot \sum_{\substack{i \neq j \\ i,j=1,2}} |(b_i(x) - b_{i,B})(b_j,B - b_{j,\omega_j,s_j(t)B})| \right)^p \omega(x) dx \right)^{1/p} \psi(t) dt,$$

$$I_5 = \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\bar{f}(x) \cdot \sum_{\substack{i \neq j \\ i,j=1,2}} |(b_i(x) - b_{i,B})(b_j(s_j(t)x) - b_{j,\omega_j,s_j(t)B})| \right)^p \omega(x) dx \right)^{1/p} \psi(t) dt,$$

$$I_6 = \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\bar{f}(x) \cdot \sum_{\substack{i \neq j \\ i,j=1,2}} |(b_{i,B} - b_{i,s_i(t)B})(b_j(s_j(t)x) - b_{j,\omega_j,s_j(t)B})| \right)^p \omega(x) dx \right)^{1/p} \psi(t) dt.$$

Choose now $p < s_1 < \infty$ and $p < s_2 < \infty$ such that $\frac{1}{s_1} = \frac{1}{p_1} + \frac{1}{q_1}$ and $\frac{1}{s_2} = \frac{1}{p_2} + \frac{1}{q_2}$. Notice that $\frac{1}{s_1} + \frac{1}{s_2} = \frac{1}{p}$. Then by Hölder's inequality, we have that

$$\begin{aligned} I_1 &\leq \int_{[0,1]^n} \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)} \int_B |f_k(s_k(t)x|^{p_k} \omega_k(x) dx \right)^{1/p_k} \times \\ &\quad \times \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)} \int_B |b_k(x) - b_{k,\omega_k,B}|^{q_k} \omega_k(x) \right)^{1/q_k} \psi(t) dt \\ &\leq \int_{[0,1]^n} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \left(\frac{1}{\omega_k(s_k(t)B)^{1+\lambda_k p_k}} \int_{s_k(t)B} |f_k(y)|^{p_k} \omega_k(y) dy \right)^{1/p_k} \times \\ &\quad \times \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)} \int_B |b_k(x) - b_{k,\omega_k,B}|^{q_k} \omega_k(x) \right)^{1/q_k} \psi(t) dt \\ &\leq \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|b_k\|_{CMO_{\omega_k}^{q_k}} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \times \int_{[0,1]^n} \left(\prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \right) \psi(t) dt. \end{aligned}$$

Similarly to the estimate of I_1 , we have that

$$\begin{aligned}
I_2 &\leq \int_{[0,1]^n} \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)} \int_B |f_k(s_k(t)x)|^{p_k} \omega_k(x) dx \right)^{1/p_k} \times \\
&\quad \times \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)} \int_B |b_k(s_k(t)x) - b_{k,\omega_k,s_k(t)B}|^{q_k} \omega(x) dx \right)^{1/q_k} \psi(t) dt \\
&\leq \int_{[0,1]^n} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)^{1+\lambda_k p_k}} \int_{s_k(t)B} |f_k(y)|^{p_k} \omega_k(y) dy \right)^{1/p_k} \times \\
&\quad \times \prod_{k=1}^2 \left(\frac{1}{\omega_k(s_k(t)B)} \int_{s_k(t)B} |b_k(y) - b_{k,\omega_k,s_k(t)B}|^{q_k} \omega(y) dy \right)^{1/q_k} \psi(t) dt \\
&\leq \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|b_k\|_{CMO_{\omega_k}^{q_k}} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k,\lambda_k}} \times \int_{[0,1]^n} \left(\prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \right) \psi(t) dt.
\end{aligned}$$

Now we give the estimate for I_3 , applying Hölder's inequality, we get that

$$\begin{aligned}
I_3 &= \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\prod_{k=1}^2 |f_k(s_k(t)x)| \omega(x) \right)^p dx \right)^{1/p} \prod_{k=1}^2 |b_{k,\omega_k,B} - b_{k,\omega_k,s_k(t)B}| \psi(t) dt \\
&\leq \int_{[0,1]^n} \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)} \int_B |f_k(s_k(t)x)|^{s_k} \omega_k(x) dx \right)^{1/s_k} \prod_{k=1}^2 |b_{k,\omega_k,B} - b_{k,\omega_k,s_k(t)B}| \psi(t) dt \\
&\leq \int_{[0,1]^n} \prod_{k=1}^2 \left(\frac{1}{\omega_k(s_k(t)B)^{1+p_k\lambda_k}} \int_{s_k(t)B} |f_k(y)|^{p_k} \omega_k(y) dy \right)^{1/p_k} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \omega_k(B)^{\lambda_k} \times \\
&\quad \times \prod_{k=1}^2 |b_{k,\omega_k,B} - b_{k,\omega_k,s_k(t)B}| \psi(t) dt \\
&\leq \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k,\lambda_k}} \int_{[0,1]^n} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \prod_{k=1}^2 |b_{k,\omega_k,B} - b_{k,\omega_k,s_k(t)B}| \psi(t) dt
\end{aligned}$$

Notice that $[0, 1]^n$ is the union of pairwise disjoint subsets $S_{\ell_1, \dots, \ell_m}$, where

$$S_{\ell_1, \dots, \ell_m} = \bigcap_{i=1}^m \{t \in [0, 1]^n : 2^{-\ell_i-1} < |s_i(t)| \leq 2^{-\ell_i}\}.$$

Thus we obtain that

$$\begin{aligned} I_3 &\leq \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \sum_{\ell_1, \ell_2=0}^{\infty} \int_{S_{\ell_1, \ell_2}} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \psi(t) \times \\ &\quad \times \prod_{k=1}^2 \left(\sum_{j=0}^{\ell_k} |b_{k, \omega_k, 2^{-j-1}B} - b_{k, \omega_k, 2^{-j}B}| + |b_{k, \omega_k, 2^{-\ell_k-1}B} - b_{k, \omega_k, s_k(t)B}| \right) dt \\ &\leq C \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \sum_{\ell_1, \ell_2=0}^{\infty} \int_{S_{\ell_1, \ell_2}} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \times \\ &\quad \times \prod_{k=1}^2 (\ell_k + 2) \|b_k\|_{CMO_{\omega_k}^{q_k}} \times \psi(t) dt \\ &\leq C \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \prod_{k=1}^2 \|b_k\|_{CMO_{\omega_k}^{q_k}} \int_{[0, 1]^n} \prod_{k=1}^2 \left(|s_k(t)|^{(d+\alpha_k)\lambda_k} \log \frac{4}{|s_k(t)|} \right) \psi(t) dt. \end{aligned}$$

Now we give the estimate for I_4 . Similarly, we choose $1 < s < \infty$ such that $\frac{1}{s} = \frac{1}{q_1} + \frac{1}{q_2}$. Let

$$\overline{b_{i,j}}(x) := |(b_i(x) - b_{i,B})(b_{j,B} - b_{j, \omega_j, s_j(t)B})|,$$

then Minkowski's inequality and Hölders inequality imply that

$$\begin{aligned}
I_4 &= \int_{[0,1]^n} \left(\frac{1}{\omega(B)} \int_B \left(\left(\prod_{k=1}^2 |f_k(s_k(t)x)| \right) \left(\sum_{\substack{i \neq j \\ i,j=1,2}} \overline{b_{i,j}}(x) \right) \right)^p \omega(x) dx \right)^{1/p} \psi(t) dt \\
&\leq \int_{[0,1]^n} \sum_{\substack{i \neq j \\ i,j=1,2}} \left(\frac{1}{\omega(B)} \int_B \left(\left(\prod_{k=1}^2 |f_k(s_k(t)x)| \right) \cdot \overline{b_{i,j}}(x) \right)^p \omega(x) dx \right)^{1/p} \psi(t) dt \\
&\leq \int_{[0,1]^n} \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)} \int_B |f_k(s_k(t)x)|^{p_k} \omega_k(x) dx \right)^{1/p_k} \\
&\quad \times \left(\sum_{\substack{i \neq j \\ i,j=1,2}} \left(\frac{1}{\omega_i(B)} \int_B |b_i(x) - b_{i,B}|^s \omega_i(x) dx \right)^{1/s} |b_{j,B} - b_{j,\omega_j,s_j(t)B}| \right) \\
&\leq C \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \int_{[0,1]^n} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \prod_{k=1}^2 \left(\frac{1}{\omega_k(B)^{1+\lambda_k p_k}} \int_{s_k(t)B} |f_k(x)|^{p_k} \omega_k(x) dx \right)^{1/p_k} \\
&\quad \times \left(\sum_{\substack{i \neq j \\ i,j=1,2}} \left(\frac{1}{\omega_i(B)} \int_B |b_i(x) - b_{i,B}|^s \omega_i(x) dx \right)^{1/s} |b_{j,B} - b_{j,\omega_j,s_j(t)B}| \right) \psi(t) dt \\
&\leq C \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \int_{[0,1]^n} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \psi(t) \\
&\quad \times \left(\sum_{\substack{i \neq j \\ i,j=1,2}} \left(\frac{1}{\omega_i(B)} \int_B |b_i(x) - b_{i,B}|^s \omega_i(x) dx \right)^{1/s} |b_{j,B} - b_{j,\omega_j,s_j(t)B}| \right) dt
\end{aligned}$$

From the estimates of I_1 and I_3 , we deduce that

$$\begin{aligned}
I_4 &\leq C \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \prod_{k=1}^2 \|b_k\|_{C\dot{M}O_k^{q_k}} \\
&\quad \times \int_{[0,1]^n} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \left(1 + \sum_{k=1}^2 \log \frac{2}{|s_k(t)|} \right) \psi(t) dt.
\end{aligned}$$

It can be deduced from the estimates of I_1, I_2, I_3, I_4 that

$$I_5 \leq C \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \prod_{k=1}^2 \|b_k\|_{CMO_{\omega_k}^{q_k}} \int_{[0,1]^n} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \psi(t) dt.$$

$$I_6 \leq C \prod_{k=1}^2 \omega_k(B)^{\lambda_k} \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \prod_{k=1}^2 \|b_k\|_{CMO_{\omega_k}^{q_k}} \\ \times \int_{[0,1]^n} \prod_{k=1}^2 |s_k(t)|^{(d+\alpha_k)\lambda_k} \left(1 + \sum_{k=1}^2 \log \frac{2}{|s_k(t)|} \right) \psi(t) dt.$$

Combining the estimates of I_1, I_2, I_3, I_4, I_5 and I_6 gives

$$\left(\frac{1}{\omega(B)} \int_B |U_{\psi, s}^{2, n, b}(f_1, f_2)(x)|^p \omega(x) dx \right)^{1/p} \leq C \omega(B)^\lambda \prod_{k=1}^2 \|f_k\|_{\dot{B}^{p_k, \lambda_k}} \prod_{k=1}^2 \|b_k\|_{CMO_{\omega_k}^{q_k}} \\ \times \int_{[0,1]^n} \prod_{k=1}^2 \left(|s_k(t)|^{(d+\alpha_k)\lambda_k} \log \frac{4}{|s_k(t)|} \right) \psi(t) dt.$$

This proves (i).

Now we prove the necessity in (ii). Assume that $U_{\psi, s}^{2, n, b}$ is bounded from $\dot{B}_{\omega_1}^{p_1, \lambda_1}(\mathbb{R}^d) \times \dot{B}_{\omega_m}^{p_m, \lambda_m}(\mathbb{R}^d)$ to $\dot{B}_\omega^{p, \lambda}(\mathbb{R}^d)$. Firstly, we need the following lemmas.

Lemma 4.1. *If $\omega \in \mathcal{W}_\alpha$ and has doubling property, then $\log|x| \in BMO(\omega)$.*

This lemma was proved in [7], for convenience of the reader, the proof to the lemma is presented here.

Proof. To prove $\log|x| \in BMO(\omega)$, for any $x_0 \in \mathbb{R}^d$ and $r > 0$, we must find a constant $c_{x_0, r}$, such that $\frac{1}{\omega(B(x_0, r))} \int_{|x-x_0| \leq r} |\log|x| - c_{x_0, r}| \omega(x) dx$ is uniformly bounded. Since

$$\begin{aligned} & \frac{1}{\omega(B(x_0, r))} \int_{|x-x_0| \leq r} |\log|x| - c_{x_0, r}| \omega(x) dx \\ &= \frac{r^{\alpha+n}}{\omega(B(x_0, r))} \int_{|z-r^{-1}x_0| \leq 1} |\log|z| + \log r - c_{x_0, r}| \omega(z) dz \\ &= \frac{1}{\omega(B(r^{-1}x_0, 1))} \int_{|z-r^{-1}x_0| \leq 1} |\log|z| + \log r - c_{x_0, r}| \omega(z) dz, \end{aligned}$$

we may take $c_{x_0,r} = c_{r^{-1}x_0,1} - \log r$, and so things reduce to the case that $r = 1$ and x_0 is arbitrary. Let

$$A_{x_0} = \frac{1}{\omega(B(x_0,1))} \int_{|z-x_0|\leq 1} |\log |z| - c_{x_0,1}| \omega(z) dz.$$

If $|x_0| \leq 2$, we take $c_{x_0,1} = 0$, and observe that

$$\begin{aligned} A_{x_0} &\leq \frac{1}{\omega(B(x_0,1))} \int_{|z|\leq 3} \log 3 \cdot \omega(z) dz = \log 3 \cdot \frac{\omega(B(0,3))}{\omega(B(x_0,1))} \\ &\leq \log 3 \cdot \frac{\omega(B(x_0,6))}{\omega(B(x_0,1))} \leq C < \infty, \end{aligned}$$

where the last inequality comes from the assumption that ω has doubling property.

If $|x_0| \geq 2$, take $c_{x_0,1} = \log |x_0|$. In this case, notice that

$$\begin{aligned} A_{x_0} &= \frac{1}{\omega(B(x_0,1))} \int_{B(x_0,1)} \left| \log \frac{|z|}{|x_0|} \right| \omega(z) dz \\ &\leq \frac{1}{\omega(B(x_0,1))} \int_{B(x_0,1)} \max \left\{ \log \frac{|x_0|+1}{|x_0|}, \log \frac{|x_0|}{|x_0|-1} \right\} \cdot \omega(z) dz \\ &\leq \max \left\{ \log \frac{|x_0|+1}{|x_0|}, \log \frac{|x_0|}{|x_0|-1} \right\} \leq \log 2. \end{aligned}$$

Thus $\log |x|$ belongs to $BMO_\omega(\mathbb{R}^d)$. \square

Lemma 4.2. *Let $\omega \in \mathcal{W}_\alpha$, where $\alpha > -d$, $1 < p < \infty$ and $-\frac{1}{p} < \lambda$. The the function $f_0(x) = |x|^{(d+\alpha)\lambda}$ belongs to $\dot{B}_\omega^{p,\lambda}$ and*

$$\|f\|_{\dot{B}_\omega^{p,\lambda}} = (\omega_k(S_d))^{-\lambda_k} \left(\frac{1}{(d+\alpha)(1+\lambda p)} \right)^{1/p}.$$

Since the proof of this lemma is straightforward we omit it. Now we set $b_1(x) = b_2(x) = \log |x|$. Lemma 4.1 gives b_1, b_2 belong to $BMO_\omega(\mathbb{R}^d)$. Since $BMO_\omega(\mathbb{R}^d) \subset CMO_\omega^q(\mathbb{R}^d)$ for any $1 < q < \infty$, we obtain that $b_k \in CMO_{\omega_k}^{q_k}(\mathbb{R}^d)$. Define $f_k(x) = |x|^{(d+\alpha_k)\lambda_k}$ if $x \in \mathbb{R}^d \setminus \{0\}$ and $f_k(0) = 0$. From lemma 4.2, we get that

$$(4.4) \quad \|f_k\|_{\dot{B}_{\omega_k}^{p_k,\lambda_k}} = (\omega_k(S_d))^{-\lambda_k} \left(\frac{1}{(d+\alpha_k)(1+\lambda_k p_k)} \right)^{1/p_k}.$$

Also we have

$$U_{\psi,s}^{2,n,b}(f_1, f_2)(x) = \prod_{k=1}^2 |x|^{(d+\alpha_k)\lambda_k} \int_{[0,1]^n} \prod_{k=1}^2 \left(|s_k(t)|^{(d+\alpha_k)\lambda_k} \log \frac{1}{|s_k(t)|} \right) dt.$$

Let $B = B(0, R)$ be any ball of \mathbb{R}^d , then

$$\begin{aligned} & \left(\frac{1}{\omega(B)^{1+\lambda p}} \int_B \left| U_{\psi,s}^{2,n,b}(f_1, f_2)(x) \right|^p \omega(x) dx \right)^{1/p} \\ & \leq \left(\frac{1}{\omega(B)^{1+\lambda p}} \int_B |x|^{(d+\alpha)\lambda p} \omega(x) dx \right)^{1/p} \left(\int_{[0,1]^n} \prod_{k=1}^2 \left(|s_k(t)|^{(d+\alpha_k)\lambda_k} \log \frac{1}{|s_k(t)|} \right) \psi(t) dt \right)^{1/p} \\ & = (\omega_k(S_d))^{-\lambda_k} \left(\frac{1}{(d+\alpha)(1+\lambda p)} \right)^{1/p} \left(\int_{[0,1]^n} \prod_{k=1}^2 \left(|s_k(t)|^{(d+\alpha_k)\lambda_k} \log \frac{1}{|s_k(t)|} \right) \psi(t) dt \right)^{1/p} \\ & = \prod_{k=1}^2 \|f_k\|_{\dot{B}_{\omega_k}^{p_k, \lambda_k}} \left(\int_{[0,1]^n} \prod_{k=1}^2 \left(|s_k(t)|^{(d+\alpha_k)\lambda_k} \log \frac{1}{|s_k(t)|} \right) \psi(t) dt \right)^{1/p}. \end{aligned}$$

Taking the supremum over $R > 0$, we get that

$$\int_{[0,1]^n} \prod_{k=1}^2 \left(|s_k(t)|^{(d+\alpha_k)\lambda_k} \log \frac{1}{|s_k(t)|} \right) \psi(t) dt < \infty.$$

□

Acknowledgements. The authors would like to thank Prof. Nguyen Minh Chuong for many helpful suggestions and discussions. This paper is granted by Vietnam NAFOSTED (National Foundation for Science and Technology Development).

REFERENCES

- [1] D. R. Adams, A note on Riesz potentials. *Duke Math. J.* **42** (1975), no. 4, 765–778.
- [2] J. Alvarez, J. Lakey, and M. Guzmán-Partida, Spaces of bounded λ -central mean oscillation, Morrey spaces, and λ -central Carleson measures. *Collect. Math.* **51** (2000), no. 1, 147.
- [3] K. F. Andersen, Boundedness of Hausdorff operator on $L^p(\mathbb{R}^d)$, $H^1(\mathbb{R}^d)$ and $BMO(\mathbb{R}^d)$. *Acta Sci. Math. (Szeged)* **69** (2003), no. 1-2, 409–418.
- [4] J. S. Bradley, Hardy inequalities with mixed norms. *Canad. Math. Bull.* **21** (1978), no. 4, 405–408.

- [5] C. Carton-Lebrun and M. Fosset, Moyennes et quotients de Taylor dans BMO . Bull. Soc. Roy. Sci. Liège **53** (1984), no. 2, 85–87.
- [6] F. Chiarenza and M. Frasca, Morrey spaces and Hardy-Littlewood maximal function. Rend. Math. Appl. **7** (1987), no. 3-4, 273–279.
- [7] N. M. Chuong and H. D. Hung, Bounds of weighted Hardy-Cesàro operators on weighted Lebesgue and BMO spaces. Integral Transforms Spec. Funct. **25** (2014), no. 9, 697–710.
- [8] M. Christ and L. Grafakos, Best constants for two nonconvolution inequalities. Proc. Amer. Math. Soc. **123** (1995), no. 6, 1687–1693.
- [9] R. R. Coifman, R. Rochberg and G. Weiss, Factorization theorems for Hardy spaces in several variables. Ann. of Math. (2) **103** (1976), no. 3, 611–635.
- [10] D. E. Edmunds and W. D. Evans, Hardy operators, function spaces and embeddings. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2004. xii+326 pp.
- [11] W. G. Faris, Weak Lebesgue spaces and quantum mechanical binding. Duke Math. J. **43** (1976), no. 2, 365–373.
- [12] G. B. Folland, Real Analysis: Modern techniques and their applications: Pure and Applied Mathematics. Second edition. New York: John Wiley & Sons, Inc.1999.
- [13] Z. W. Fu, Z. G. Liu and S. Z. Lu, Commutators of weighted Hardy operators on \mathbb{R}^d . Proc. Amer. Math. Soc. **137** (2009), no. 10, 3319–3328.
- [14] Z. W. Fu, S. L. Gong, S. Z. Lu and W. Yuan, Weighted multilinear Hardy operators and commutators, Forum Math. 2014, DOI: 10.1515/forum-2013-0064.
- [15] Z. W. Fu, L. Grafakos, S. Z. Lu, F. Zhao, Sharp bounds for m -linear Hardy and Hilbert operators. Houston J. Math. **38** (2012), no. 1, 225–244.
- [16] R. R. Goldberg, Convolutions and general transforms on L^p . Duke Math. J. **27** (1960), 251–259.
- [17] S.L. Gong , Z.W. Fu and B. Ma, Weighted Multilinear Hardy Operators on Herz Type Spaces, The Scientific World Journal Volume 2014, 10 pages.
- [18] L. Grafakos, Classical Fourier analysis. Second edition. Graduate Texts in Mathematics, 249. Springer, New York, 2008. xvi+489 pp.
- [19] L. Grafakos and J. M. Martell, Extrapolation of weighted norm inequalities for multivariable operators and applications. J. Geom. Anal. **14** (2004), no. 1, 19–46.
- [20] G. H. Hardy, J. E. Littlewood and G. Pólya, Inequalities: Second Edition. Cambridge: the University Press; 1952.
- [21] H. D. Hung, The P -adic weighted Hardy-Cesàro operator and an application to discrete Hardy inequalities. J. Math. Anal. Appl. **409** (2014), 868–879.
- [22] F. John and L. Nirenberg, On functions of bounded mean oscillation. Comm. Pure. Appl. Math. **14** (1961), 415–426.
- [23] C. Kenig and E. M. Stein, Multilinear estimates and fractional integration. Math. Res. Lett. **6** (1999), 1–15.
- [24] Y. Komori and S. Shirai, Weighted Morrey spaces and a singular integral operator. Math. Nachr. **282** (2009), no. 2, 219–231.
- [25] E. Liflyand and F. Móricz, The Hausdorff operator is bounded on the real Hardy space $H^1(\mathbb{R})$. Proc. Amer. Math. Soc. **128** (2000), no. 5, 1391–1396.
- [26] A. K. Lerner and E. Liflyand, Multidimensional Hausdorff operators on the real Hardy space. J. Aust. Math. Soc. **83** (2007), no. 1, 79–86.
- [27] S. Z. Lu and D. C. Yang, The central BMO spaces and Littlewood-Paley operators. Approx. Theory Appl. (N.S.) **11** (1995), no. 3, 72–94.

- [28] S. Z. Lu, D. Yan and F. Zhao, Sharp bounds for Hardy type operators on higher-dimensional product spaces. *J. Inequal. Appl.* 2013, 2013:148, 11 pp.
- [29] B. Muckenhoupt, Hardy's inequalities with weights. *Studia Math.* **44** (1972), 31–38.
- [30] E. Sawyer, Weighted Lebesgue and Lorentz norm inequalities for the Hardy operator. *Trans. Amer. Math. Soc.* **281** (1984), no. 1, 329–337.
- [31] J. Xiao, L^p and BMO bounds of weighted Hardy-Littlewood averages. *J. Math. Anal. Appl.* **262** (2001), no. 2, 660–666.

HIGH SCHOOL FOR GIFTED STUDENTS, HANOI NATIONAL UNIVERSITY OF EDUCATION,
136 XUAN THUY, HANOI, VIETNAM

E-mail address: `hunghaduy@gmail.com`

DEPARTMENT OF MATHEMATICS, QUY NHON UNIVERSITY, 170 AN DUONG VUONG,
QUY NHON, BINH DINH, VIET NAM

E-mail address: `dangky@math.cnrs.fr`