

Defining Incomplete Gamma Type Function with Negative Arguments and Polygamma functions $\psi^{(n)}(-m)$

Emin Özçağ and İnci Ege

Abstract. In this paper the incomplete gamma function $\gamma(\alpha, x)$ and its derivative is considered for negative values of α and the incomplete gamma type function $\gamma_*(\alpha, x_-)$ is introduced. Further the polygamma functions $\psi^{(n)}(x)$ are defined for negative integers via the neutrix setting.

AMS Mathematics Subject Classification (2010): 33B15, 33B20, 33D05

Key words and phrases: Gamma function, incomplete gamma function, Digamma Function, Polygamma Function, neutrix, neutrix limit.

1. Introduction

The incomplete gamma function of real variable and its complement are defined via integrals

$$\gamma(\alpha, x) = \int_0^x u^{\alpha-1} e^{-u} du \quad (1)$$

and

$$\Gamma(\alpha, x) = \int_x^\infty u^{\alpha-1} e^{-u} du \quad (2)$$

respectively, for $\alpha > 0$ and $x \geq 0$ see [1, 8].

Note that the definition of $\gamma(\alpha, x)$ does not valid for negative α , but it can be extended by the identity

$$\gamma(\alpha, x) = \Gamma(\alpha) - \Gamma(\alpha, x).$$

The incomplete gamma function does not exists for negative integers α or zero [13]. However, Fisher et. al. [4] defined $\gamma(0, x)$ by

$$\gamma(0, x) = \int_0^x u^{-1}(e^{-u} - 1) du + \ln x.$$

The problem of evaluating incomplete gamma function is subject of some earlier articles. The general problem in which α and x are complex was considered by Winitzki in [15], but no method is found to be satisfactory in cases where $Re[x] < 0$. The incomplete gamma function with negative arguments are difficult to compute, see [6]. In [14] Thompson gave the algorithm for accurately computing the incomplete gamma function $\gamma(\alpha, x)$ in the cases where $\alpha = n + 1/2, n \in \mathbb{Z}$ and $x < 0$.

2. Incomplete Gamma Function

A series expansion of incomplete gamma function can be obtained by replacing the exponential in (1) with its Maclaurin series, the result is that

$$\gamma(\alpha, x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!(\alpha+k)} x^{\alpha+k}. \quad (3)$$

A single integration by parts in (1) yields the recurrence relation

$$\gamma(\alpha+1, x) = \alpha\gamma(\alpha, x) - x^\alpha e^{-x} \quad (4)$$

and thus the incomplete gamma function can be defined for $\alpha < 0$ and $\alpha \neq -1, -2, \dots$

By repeatedly applying recurrence relation, we obtain

$$\gamma(\alpha, x) = x^{\alpha-1} e^{-x} \sum_{k=0}^{\infty} \frac{x^k}{(\alpha)_k} + \frac{1}{(\alpha)_m} \gamma(\alpha+m, x)$$

where $(\cdot)_k$ is Pochhammer's symbol [1]. The last term on the right-hand side disappears in the limit $m \rightarrow \infty$.

By regularization we have

$$\gamma(\alpha, x) = \int_0^x u^{\alpha-1} \left[e^{-u} - \sum_{i=0}^{m-1} \frac{(-u)^i}{i!} \right] du + \sum_{i=0}^{m-1} \frac{(-1)^i x^{\alpha+i}}{(\alpha+i)i!}. \quad (5)$$

for $-m < \alpha < -m+1$ and $x > 0$. It follows from the definition of gamma function that

$$\lim_{x \rightarrow \infty} \gamma(\alpha, x) = \Gamma(\alpha)$$

for $\alpha \neq 0, -1, -2, \dots$, see [7, 11].

In the following we let N be the neutrix [2, 3, 4] having domain $N' = \{\varepsilon : 0 < \varepsilon < \infty\}$ and range N'' the real numbers, with negligible functions finite linear sums of the functions

$$\varepsilon^\lambda \ln^{r-1} \varepsilon, \quad \ln^r \varepsilon \quad (\lambda < 0, \quad r \in \mathbb{Z}^+) \quad (6)$$

and all functions of ε which converge to zero in the normal sense as ε tends to zero.

If $f(\varepsilon)$ is a real (or complex) valued function defined on N' and if it is possible to find a constant c such that $f(\varepsilon) - c$ is in N , then c is called the neutrix limit of $f(\varepsilon)$ as $\varepsilon \rightarrow 0$ and we write $N\text{-}\lim_{\varepsilon \rightarrow 0} f(\varepsilon) = c$.

Note that if a function $f(\varepsilon)$ tends to c in the normal sense as ε tends to zero, it converges to c in the neutrix sense.

On using equation (5), the incomplete gamma function $\gamma(\alpha, x)$ was also defined by

$$\gamma(\alpha, x) = N\text{-}\lim_{\varepsilon \rightarrow 0} \int_\varepsilon^x u^{\alpha-1} e^{-u} du \quad (7)$$

for all $\alpha \in \mathbb{R}$ and $x > 0$, and it was shown that $\lim_{x \rightarrow \infty} \gamma(-m, x) = \Gamma(-m)$ for $m \in \mathbb{N}$, see [4].

Further, the r -th derivative of $\gamma(\alpha, x)$ was similarly defined by

$$\gamma^{(r)}(\alpha, x) = \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^x u^{\alpha-1} \ln^r u e^{-u} du \quad (8)$$

for all α and $r = 0, 1, 2, \dots$, provided that the neutrix limit exists, see [11].

By using the neutrix N given in equation(6), the interesting formula

$$\gamma(-m, x) + \frac{1}{m} \gamma(-m+1, x) = \frac{(-1)^m}{mm!} - \frac{1}{m} e^{-x} x^{-m}$$

was obtained in [11] for $m \in \mathbb{N}$.

It can be easily seen that equation (3) is also valid for $\alpha < 0$ and $\alpha \neq -1, -2, \dots$. In fact it follows from the definition that

$$\begin{aligned} \gamma(\alpha, x) &= \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^x u^{\alpha-1} e^{-u} du = \text{N-}\lim_{\varepsilon \rightarrow 0} \left[\sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \int_{\varepsilon}^x u^{\alpha+k-1} du \right] \\ &= \text{N-}\lim_{\varepsilon \rightarrow 0} \sum_{k=0}^{\infty} \frac{(-1)^k}{k!(\alpha+k)} [x^{\alpha+k} - \varepsilon^{\alpha+k}] \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k}{k!(\alpha+k)} x^{\alpha+k}. \end{aligned}$$

Now assume that $\alpha \in \mathbb{Z}^-$, then writing

$$\begin{aligned} \int_{\varepsilon}^x u^{-m-1} e^{-u} du &= \sum_{\substack{k=0, \\ k \neq m}}^{\infty} \frac{(-1)^k}{k!} \int_{\varepsilon}^x u^{-m+k-1} du + \frac{(-1)^m}{m!} \int_{\varepsilon}^x u^{-1} du \\ &= \sum_{\substack{k=0, \\ k \neq m}}^{\infty} \frac{(-1)^k}{k!(k-m)} x^{k-m} + \frac{(-1)^m}{m!} \ln x + O(\varepsilon) \end{aligned}$$

where $O(\varepsilon)$ is negligible function, and taking the neutrix limit we obtain

$$\gamma(-m, x) = \sum_{\substack{k=0, \\ k \neq m}}^{\infty} \frac{(-1)^k}{k!(k-m)} x^{k-m} + \frac{(-1)^m}{m!} \ln x$$

for $x > 0$ and $m \in \mathbb{N}$. If $m = 0$ we particularly have

$$\gamma(0, x) = \sum_{k=1}^{\infty} \frac{(-x)^k}{kk!} + \ln x$$

and thus it follows by the calculation of Mathematica that

$$\gamma(0, 1/2) \approx -1.13699, \quad \gamma(0, 3/4) \approx -0.917556, \quad \gamma(0, 1) \approx -0.7966.$$

Similar equation for $\gamma^{(r)}(\alpha, x)$ can be obtained using the following lemma.

Lemma 2.1

$$\begin{aligned} \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^x u^{-1} \ln^r u \, du &= \frac{\ln^{r+1} x}{r+1} \\ \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^x u^{\alpha} \ln^r u \, du &= \sum_{k=0}^{r-1} \frac{(-1)^k r! x^{\alpha+1} \ln^{r-k} x}{(\alpha+1)^{k+1} (r-k)!} + \frac{(-1)^r r! x^{\alpha+1}}{(\alpha+1)^{r+1}} \end{aligned}$$

for all $\alpha \neq -1, x > 0$ and $r \in \mathbb{N}$.

Proof. Straight forward.

Now let $\alpha < 0$ and $\alpha \neq 0, -1, -2, \dots$. Then it follows from lemma 2.1 that

$$\begin{aligned} \gamma^{(r)}(\alpha, x) &= \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^x u^{\alpha-1} \ln^r u e^{-u} \, du \\ &= \text{N-}\lim_{\varepsilon \rightarrow 0} \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \int_{\varepsilon}^x u^{\alpha+k-1} \ln^r u \, du \\ &= \sum_{k=0}^{\infty} \sum_{i=0}^{r-1} \frac{(-1)^{i+k} r!}{k! (r-i)! (\alpha+k)^{i+1}} x^{\alpha+k} \ln^{r-i} x + \\ &\quad + \sum_{k=0}^{\infty} \frac{(-1)^{k+r} r!}{k! (\alpha+k)^{r+1}} x^{\alpha+k} \end{aligned}$$

and similarly

$$\begin{aligned} \gamma^{(r)}(-m, x) &= \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^x u^{-m-1} \ln^r u e^{-u} \, du \\ &= \sum_{\substack{k=0, \\ k \neq m}}^{\infty} \sum_{i=0}^{r-1} \frac{(-1)^{i+k} r!}{k! (r-i)! (m-k)^{i+1}} x^{k-m} \ln^{r-i} x + \\ &\quad + \sum_{\substack{k=0, \\ k \neq m}}^{\infty} \frac{(-1)^{k+r} r!}{k! (m-k)^{r+1}} x^{k-m} + \frac{(-1)^m}{(r+1)m!} \ln^{r+1} x \end{aligned}$$

for $r = 1, 2, \dots$ and $m \in \mathbb{N}$.

It was indicated in [3] that equation (1) could be replaced by the equation

$$\gamma(\alpha, x) = \int_0^x |u|^{\alpha-1} e^{-u} \, du \tag{9}$$

and this equation was used to define $\gamma(\alpha, x)$ for all x and $\alpha > 0$, the integral again diverging for $\alpha \leq 0$.

The locally summable function $\gamma_*(\alpha, x_-)$ for $\alpha > 0$ was then defined by

$$\begin{aligned}\gamma_*(\alpha, x_-) &= \begin{cases} \int_0^x |u|^{\alpha-1} e^{-u} du, & x \leq 0, \\ 0, & x > 0 \end{cases} \\ &= \int_0^{-x_-} |u|^{\alpha-1} e^{-u} du.\end{aligned}\tag{10}$$

If $\alpha > 0$, then the recurrence relation

$$\gamma_*(\alpha + 1, x_-) = -\alpha \gamma_*(\alpha, x_-) - x_-^\alpha e^{-x_-}.\tag{11}$$

holds. So we can use equation (11) to extend the definition of $\gamma_*(\alpha, x_-)$ to negative non-integer values of α .

More generally it can be easily proved that

$$\gamma_*(\alpha, x_-) = \int_0^{-x_-} |u|^{\alpha-1} \left[e^{-u} - \sum_{i=0}^{m-1} \frac{(-u)^i}{i!} \right] du - \sum_{i=0}^{m-1} \frac{x_-^{\alpha+i}}{(\alpha+i)!}.\tag{12}$$

if $-m < \alpha < -m + 1$ for $m = 1, 2, \dots$

Now if $-m < \alpha < -m + 1$, $m \in \mathbb{N}$ and $x < 0$, we have

$$\int_{-\varepsilon}^{-x_-} |u|^{\alpha-1} e^{-u} du = \int_{-\varepsilon}^{-x_-} |u|^{\alpha-1} \left[e^{-u} - \sum_{i=0}^{m-1} \frac{(-u)^i}{i!} \right] du - \sum_{i=0}^{m-1} \frac{[|x|^{-\alpha+i} - \varepsilon^{-\alpha+i}]}{(\alpha+i)!}$$

and thus

$$\gamma_*(\alpha, x_-) = \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{\alpha-1} e^{-u} du$$

on using equation (12). This suggests that we define $\gamma_*(-m, x_-)$ by

$$\gamma_*(-m, x_-) = \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{-m-1} e^{-u} du\tag{13}$$

for $x < 0$ and $m \in \mathbb{N}$.

If $x < 0$, then we simply write

$$\int_{-\varepsilon}^{-x_-} |u|^{-1} e^{-u} du = \int_{-\varepsilon}^{-x_-} |u|^{-1} (e^{-u} - 1) du - \ln |x| + \ln \varepsilon$$

thus we have

$$\begin{aligned}\text{N-}\lim_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{-1} e^{-u} du &= \int_0^{-x_-} |u|^{-1} (e^{-u} - 1) du - \ln x_- \\ &= \gamma_*(0, x_-).\end{aligned}$$

Next, writing similarly

$$\begin{aligned} \int_{-\varepsilon}^{-x_-} |u|^{-m-1} e^{-u} du &= \int_{-\varepsilon}^{-x_-} |u|^{-m-1} \left[e^{-u} - \sum_{i=0}^m \frac{(-u)^i}{i!} \right] du \\ &\quad - \sum_{i=0}^{m-1} \frac{[|x|^{i-m} - \varepsilon^{i-m}]}{(m-i)!} - \frac{1}{m!} [\ln |x| - \ln \varepsilon] \end{aligned}$$

for $x < 0$ and $m \in \mathbb{N}$, then it follows that

$$\begin{aligned} \gamma_*(-m, x_-) &= \text{N-lim}_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{-m-1} e^{-u} du \\ &= \int_0^{-x_-} |u|^{-m-1} \left[e^{-u} - \sum_{i=0}^m \frac{(-u)^i}{i!} \right] du - \sum_{i=0}^{m-1} \frac{x_-^{i-m}}{(m-i)!} - \frac{1}{m!} \ln x_- \end{aligned} \quad (14)$$

Equation (14) can be regarded as the definition of incomplete gamma function $\gamma_*(-m, x_-)$ for negative integers and also written in the form

$$\begin{aligned} \gamma_*(-m, x_-) &= \int_0^{-1} |u|^{-m-1} \left[e^{-u} - \sum_{i=0}^m \frac{(-u)^i}{i!} \right] du \\ &\quad + \int_{-1}^{-x_-} |u|^{-m-1} e^{-u} du + \sum_{i=0}^{m-1} \frac{1}{(m-i)!} \end{aligned} \quad (15)$$

Replacing e^{-x} by its Maclaurin series yields in equation (13)

$$\gamma_*(-m, x_-) = \sum_{\substack{k=0, \\ k \neq m}}^{\infty} \frac{x_-^{k-m}}{(m-k)!} - \frac{1}{m!} \ln x_- \quad (m \in \mathbb{N}). \quad (16)$$

If $m = 0$ we particular have

$$\gamma_*(0, x_-) = - \sum_{k=1}^{\infty} \frac{x_-^k}{k k!} - \ln x_- \quad (17)$$

and by Mathematica

$$\begin{aligned} \gamma_*(0, -1/3) &\approx 0.735308 & \gamma_*(0, -1/2) &\approx 0.122996 \\ \gamma_*(0, -3/4) &\approx -0.630117 & \gamma_*(0, -1) &\approx -1.3179. \end{aligned}$$

Differentiating equation (12) we get

$$\begin{aligned} \gamma'_*(\alpha, x_-) &= \int_0^{-x_-} |u|^{\alpha-1} \ln |u| \left[e^{-u} - \sum_{i=0}^{m-1} \frac{(-u)^i}{i!} \right] du \\ &\quad - \sum_{i=0}^{m-1} \frac{(\alpha+i)x_-^{\alpha+i} \ln x_- - x_-^{\alpha+i}}{(\alpha+i)^2 i!} \end{aligned}$$

for $-m < \alpha < -m + 1$ and $x < 0$.

On the other hand, arranging the following integral and taking the neutrix limit

$$\begin{aligned}
\int_{-\varepsilon}^{-x_-} |u|^{\alpha-1} \ln |u| e^{-u} du &= \int_{-\varepsilon}^{-x_-} |u|^{\alpha-1} \ln |u| \left[e^{-u} - \sum_{i=0}^{m-1} \frac{(-u)^i}{i!} \right] du \\
&\quad + \sum_{i=0}^{m-1} \frac{1}{i!} \int_{-\varepsilon}^{-x_-} |u|^{\alpha+i-1} \ln |u| du \\
&= \int_{-\varepsilon}^{-x_-} |u|^{\alpha-1} \ln |u| \left[e^{-u} - \sum_{i=0}^{m-1} \frac{(-u)^i}{i!} \right] du \\
&\quad - \sum_{i=0}^{m-1} \frac{1}{i!(\alpha+i)} \left[x_-^{\alpha+i} \ln x_- - \varepsilon^{\alpha+i} \ln \varepsilon \right] \\
&\quad + \sum_{i=0}^{m-1} \frac{1}{i!(\alpha+i)^2} \left[x_-^{\alpha+i} - \varepsilon^{\alpha+i} \right].
\end{aligned}$$

Thus

$$\gamma'_*(\alpha, x_-) = \text{N-lim}_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{\alpha-1} \ln |u| e^{-u} du$$

for $\alpha \neq 0, -1, -2, \dots$ and $x < 0$.

More generally it can be shown that

$$\gamma_*^{(r)}(\alpha, x_-) = \text{N-lim}_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{\alpha-1} \ln^r |u| e^{-u} du \quad (18)$$

for $\alpha \neq 0, -1, -2, \dots, r = 0, 1, 2, \dots$ and $x < 0$. This suggests the following definition.

Definition 2.2 The r -th derivative of incomplete gamma function $\gamma_*^{(r)}(-m, x_-)$ is defined by

$$\gamma_*^{(r)}(-m, x_-) = \text{N-lim}_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{-m-1} \ln^r |u| e^{-u} du \quad (19)$$

for $r, m = 0, 1, 2, \dots$ and $x < 0$ provided that the neutrix limit exists.

Equation(18) will then define $\gamma_*^{(r)}(\alpha, x_-)$ for all α and $r = 0, 1, 2, \dots$. To prove the neutrix limit above exists we need the following lemma.

Lemma 2.3

$$\int \ln^r |u| du = \sum_{i=0}^{r-1} \frac{(-1)^i r!}{(r-i)!} u \ln^{r-i} |u| + (-1)^r r! u \quad (20)$$

and

$$\int u^{-s-1} \ln^r |u| du = - \sum_{i=0}^{r-1} \frac{r!}{(r-i)!} s^{-i-1} u^{-s} \ln^{r-i} |u| - r! s^{-r-1} u^{-s} \quad (21)$$

for $r, s = 1, 2, \dots$

Proof. Equation (20) follows by induction and equation (21) follows on using equation (20) and making the substitution $w = u^{-s}$.

Theorem 2.4 The functions $\gamma_*^{(r)}(0, x_-)$ and $\gamma_*^{(r)}(-m, x_-)$ exist for $x < 0$ and

$$\gamma_*^{(r)}(0, x_-) = \int_{-1}^{-x_-} |u|^{-1} \ln^r |u| e^{-u} du + \int_0^{-1} |u|^{-1} \ln^r |u| [e^{-u} - 1] du \quad (22)$$

for $r = 0, 1, 2, \dots$ and

$$\begin{aligned} \gamma_*^{(r)}(-m, x_-) &= \int_{-1}^{-x_-} |u|^{-m-1} \ln^r |u| e^{-u} du + \\ &+ \int_0^{-1} |u|^{-m-1} \ln^r |u| \left[e^{-u} - \sum_{i=0}^m \frac{(-u)^i}{i!} \right] du + \sum_{i=0}^{m-1} \frac{r!(m-i)^{-r-1}}{i!} \end{aligned} \quad (23)$$

for $r, m = 1, 2, \dots$

Proof. We have

$$\begin{aligned} &\int_{-\varepsilon}^{-x_-} |u|^{-1} \ln^r |u| e^{-u} du = \\ &= \int_{-1}^{-x_-} |u|^{-1} \ln^r |u| e^{-u} du + \int_{-\varepsilon}^{-1} |u|^{-1} \ln^r |u| [e^{-u} - 1] du + \int_{-\varepsilon}^{-1} |u|^{-1} \ln^r |u| du \\ &= \int_{-1}^{-x_-} |u|^{-1} \ln^r |u| e^{-u} du + \int_{-\varepsilon}^{-1} |u|^{-1} \ln^r |u| [e^{-u} - 1] du + \frac{\ln^{r+1} \varepsilon}{r+1}. \end{aligned}$$

for $r = 0, 1, 2, \dots$. It follows that

$$\begin{aligned} \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{-1} \ln^r |u| e^{-u} du &= \\ &= \int_{-1}^{-x_-} |u|^{-1} \ln^r |u| e^{-u} du + \int_0^{-1} |u|^{-1} \ln^r |u| [e^{-u} - 1] du. \end{aligned}$$

and so equation (22) follows.

Now let us consider

$$\begin{aligned} \int_{-\varepsilon}^{-x_-} |u|^{-m-1} \ln^r |u| e^{-u} du &= \int_{-1}^{-x_-} |u|^{-m-1} \ln^r |u| e^{-u} du + \\ &+ \int_{-\varepsilon}^{-1} |u|^{-m-1} \ln^r |u| \left[e^{-u} - \sum_{i=0}^m \frac{(-u)^i}{i!} \right] du \\ &+ \sum_{i=0}^m \frac{(-1)^i}{i!} \int_{-\varepsilon}^{-1} |u|^{-m+i-1} \ln^r |u| du \end{aligned}$$

$$\begin{aligned}
&= \int_{-1}^{-x_-} |u|^{-m-1} \ln^r |u| e^{-u} du + \int_{-\varepsilon}^{-1} |u|^{-m-1} \ln^r |u| \left[e^{-u} - \sum_{i=0}^m \frac{(-u)^i}{i!} \right] du + \\
&\quad + \sum_{i=0}^{m-1} \sum_{j=0}^{r-1} \frac{(-1)^{i-m-1} r!}{i! (r-j)!} (m-i)^{-j-1} \varepsilon^{m-i} \ln^{r-j} \varepsilon + \\
&\quad - \sum_{i=0}^{m-1} \frac{(-1)^{i-m-1} (m-i)^{-r-1} r!}{i!} \left[(-1)^{m-i} - \varepsilon^{m-i} \right] + \frac{\ln^{r+1} \varepsilon}{m!(r+1)}.
\end{aligned}$$

Thus

$$\begin{aligned}
\gamma_*^{(r)}(-m, x_-) &= \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{-\varepsilon}^{-x_-} |u|^{-m-1} \ln^r |u| e^{-u} du \\
&= \int_{-1}^{-x_-} |u|^{-m-1} \ln^r |u| e^{-u} du + \int_0^{-1} |u|^{-m-1} \ln^r |u| \left[e^{-u} - \sum_{i=0}^m \frac{(-u)^i}{i!} \right] du \\
&\quad + \sum_{i=0}^{m-1} \frac{r! (m-i)^{-r-1}}{i!}.
\end{aligned}$$

for $r, m = 1, 2, \dots$. Equation (23) follows.

3. Polygamma Functions $\psi^{(n)}(-m)$

The polygamma function is defined by

$$\psi^{(n)}(x) = \frac{d^n}{dx^n} \psi(x) = \frac{d^{n+1}}{dx^{n+1}} \ln \Gamma(x) \quad (x > 0). \quad (24)$$

It may be represented as

$$\psi^{(n)}(x) = (-1)^{n+1} \int_0^\infty \frac{t^n e^{-xt}}{1 - e^{-t}} dt$$

which holds for $x > 0$, and

$$\psi^{(n)}(x) = - \int_0^1 \frac{t^{x-1} \ln^n t}{1-t} dt. \quad (25)$$

It satisfies the recurrence relation

$$\psi^{(n)}(x+1) = \psi^{(n)}(x) + \frac{(-1)^n n!}{x^{n+1}} \quad (26)$$

see [8]. This is used to define the polygamma function for negative non-integer values of x . Thus if $-m < x < -m+1$, $m = 1, 2, \dots$, then

$$\psi^{(n)}(x) = - \int_0^1 \frac{t^{x+m-1}}{1-t} \ln^n t dt - \sum_{k=0}^{m-1} \frac{(-1)^n n!}{(x+k)^{n+1}}. \quad (27)$$

Kölbig gave the formulae for the integral $\int_0^1 t^{\lambda-1}(1-t)^{-\nu} \ln^m t dt$ for integer and half-integer values of λ and ν in [10]. As the integral representation of the polygamma function is similar to the integral mentioned above, we prove the existence of the integral in equation (25) for all values of x by using the neutrix limit.

Now we let N be a neutrix having domain the open interval $\{\epsilon : 0 < \epsilon < \frac{1}{2}\}$ with the same negligible functions as in equation (6). We first of all need the following lemma.

Lemma 3.1. The neutrix limits as ϵ tends to zero of the functions

$$\int_{\epsilon}^{1/2} t^x \ln^n t \ln^r(1-t) dt, \quad \int_{1/2}^{1-\epsilon} (1-t)^x \ln^n t \ln^r(1-t) dt$$

exists for $n, r = 0, 1, 2, \dots$ and all x .

Proof. Suppose first of all that $n = r = 0$. Then

$$\int_{\epsilon}^{1/2} t^x dt = \begin{cases} \frac{2^{-x-1} - \epsilon^{x+1}}{x+1}, & x \neq -1, \\ -\ln 2 - \ln \epsilon, & x = -1 \end{cases}$$

and so $N\text{-}\lim_{\epsilon \rightarrow 0} \int_{\epsilon}^{1/2} t^x dt$ exists for all x .

Now suppose that $r = 0$ and that $N\text{-}\lim_{\epsilon \rightarrow 0} \int_{\epsilon}^{1/2} t^x \ln^n t dt$ exists for some nonnegative integer n and all x . Then

$$\int_{\epsilon}^{1/2} t^x \ln^{n+1} t dt = \begin{cases} \frac{-2^{-x-1} \ln^{n+1} 2 - \epsilon^{x+1} \ln^{n+1} \epsilon}{x+1} - \frac{n+1}{x+1} \int_{\epsilon}^{1/2} t^x \ln^n t dt, & x \neq -1, \\ \frac{(-1)^n \ln^{n+2} 2 - \ln^{n+2} \epsilon}{n+2}, & x = -1 \end{cases}$$

and it follows by induction that $N\text{-}\lim_{\epsilon \rightarrow 0} \int_{\epsilon}^{1/2} t^x \ln^n t dt$ exists for $n = 0, 1, 2, \dots$ and all x .

Finally we note that we can write

$$\ln^r(1-t) = \sum_{i=1}^{\infty} \alpha_{ir} t^i$$

for $r = 1, 2, \dots$, the expansion being valid for $|t| < 1$. Choosing a positive integer k such that $x + k > -1$, we have

$$\begin{aligned} & \int_{\epsilon}^{1/2} t^x \ln^n t \ln^r(1-t) dt = \\ & = \sum_{i=1}^{k-1} \alpha_{in} \int_{\epsilon}^{1/2} t^{x+i} \ln^n t dt + \sum_{i=k}^{\infty} \alpha_{in} \int_{\epsilon}^{1/2} t^{x+i} \ln^n t dt. \end{aligned}$$

It follows from what we have just proved that

$$N\text{-}\lim_{\epsilon \rightarrow 0} \sum_{i=1}^{k-1} \alpha_{in} \int_{\epsilon}^{1/2} t^{x+i} \ln^n t dt$$

exists and further

$$\begin{aligned} \text{N-}\lim_{\varepsilon \rightarrow 0} \sum_{i=k}^{\infty} \alpha_{in} \int_{\varepsilon}^{1/2} t^{x+i} \ln^n t \, dt &= \lim_{\varepsilon \rightarrow 0} \sum_{i=k}^{\infty} \alpha_{in} \int_{\varepsilon}^{1/2} t^{x+i} \ln^n t \, dt \\ &= \sum_{i=k}^{\infty} \alpha_{in} \int_0^{1/2} t^{x+i} \ln^n t \, dt, \end{aligned}$$

proving that the neutrix limit of $\int_{\varepsilon}^{1/2} t^x \ln^n t \ln^r(1-t) \, dt$ exists for $n, r = 0, 1, 2, \dots$ and all x . Making the substitution $1-t = u$ in

$$\int_{1/2}^{1-\varepsilon} (1-t)^x \ln^n t \ln^r t \, dt,$$

it follows that $\int_{1/2}^{1-\varepsilon} (1-t)^x \ln^n t \ln^r t \, dt$ also exists for $n, r = 0, 1, 2, \dots$ and all x .

Lemma 3.2. The neutrix limit as $\varepsilon \rightarrow 0$ of the integral $\int_{\varepsilon}^1 t^{-m-1} \ln^n t \, dt$ exists for $m, n = 1, 2, \dots$ and

$$\text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 t^{-m-1} \ln^n t \, dt = -\frac{n!}{m^{n+1}}. \quad (28)$$

Proof. Integrating by parts, we have

$$\int_{\varepsilon}^1 t^{-m-1} \ln t \, dt = m^{-1} \varepsilon^{-m} \ln \varepsilon + m^{-1} \int_{\varepsilon}^1 t^{-m-1} \, dt$$

and so

$$\text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 t^{-m-1} \ln t \, dt = -\frac{1}{n^2}$$

proving equation (28) for $n = 1$ and $m = 1, 2, \dots$

Now assume that equation (28) holds for some m and $n = 1, 2, \dots$. Then

$$\begin{aligned} \int_{\varepsilon}^1 t^{-m-2} \ln^n t \, dt &= (m+1)^{-1} \varepsilon^{-m-1} \ln^n \varepsilon + \frac{n}{m+1} \int_{\varepsilon}^1 t^{-m-2} \ln^{n-1} t \, dt \\ &= (m+1)^{-1} \varepsilon^{-m-1} \ln^n \varepsilon + \frac{n}{m+1} \frac{-(n-1)!}{(m+1)^n} \end{aligned}$$

and it follows that

$$\text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 t^{-m-2} \ln^n t \, dt = -\frac{n!}{(m+1)^{n+1}}$$

proving equation (28) for $m+1$ and $n = 1, 2, \dots$

Using the regularization and the neutrix limit, we prove the following theorem.

Theorem 3.3. The function $\psi^{(n)}(x)$ exists for $n = 0, 1, 2, \dots$, and all x .

Proof. Choose positive integer r such that $x > -r$. Then we can write

$$\begin{aligned} \int_{\varepsilon}^{1-\varepsilon} \frac{t^{x-1}}{1-t} \ln^n t \, dt &= \int_{\varepsilon}^{1/2} t^{x-1} \ln^n t \left[\frac{1}{1-t} - \sum_{i=0}^{r-1} (-1)^i t^i \right] dt \\ &\quad + \sum_{i=0}^{r-1} (-1)^i \int_{\varepsilon}^{1/2} t^{x+i-1} \ln^n t \, dt + \int_{1/2}^{1-\varepsilon} \frac{t^{x-1}}{1-t} \ln^n t \, dt. \end{aligned}$$

We have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^{1/2} \frac{t^{x-1}}{1-t} \ln^n t \left[\frac{1}{1-t} - \sum_{i=0}^{r-1} (-1)^i t^i \right] dt &= \\ &= \int_0^{1/2} \frac{t^{x-1}}{1-t} \ln^n t \left[\frac{1}{1-t} - \sum_{i=0}^{r-1} (-1)^i t^i \right] dt \end{aligned}$$

and

$$\lim_{\varepsilon \rightarrow 0} \int_{1/2}^{1-\varepsilon} \frac{t^{x-1}}{1-t} \ln^n t \, dt = \int_{1/2}^1 \frac{t^{x-1}}{1-t} \ln^n t \, dt$$

the integrals being convergent. Further, from the Lemma 3.1 we see that the neutrix limit of the function

$$\sum_{i=0}^{r-1} (-1)^i \int_{\varepsilon}^{1/2} t^{x+i-1} \ln^n t \, dt$$

exists and implying that

$$\text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^{1-\varepsilon} \frac{t^{x-1}}{1-t} \ln^n t \, dt$$

exists. This proves the existence of the function $\psi^{(n)}(x)$ for $n = 0, 1, 2, \dots$, and all x .

Before giving our main theorem, we note that

$$\psi^{(n)}(x) = -\text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 \frac{t^{x-1}}{1-t} \ln^n t \, dt$$

since the integral is convergent in the neighborhood of the point $t = 1$.

Theorem 3.4. The function $\psi^{(n)}(-m)$ exists and

$$\psi^{(n)}(-m) = \sum_{i=1}^m \frac{n!}{i^{n+1}} + (-1)^{n+1} n! \zeta(n+1) \quad (29)$$

for $n = 1, 2, \dots$ and $m = 0, 1, 2, \dots$, where $\zeta(n)$ denotes zeta function.

Proof. From Theorem 3.3, we have

$$\begin{aligned}\psi^{(n)}(-m) &= -\text{N-lim}_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 \frac{t^{-m-1} \ln^n t}{1-t} dt \\ &= -\text{N-lim}_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 \left[\sum_{i=1}^{m+1} t^{-i} + (1-t)^{-1} \right] \ln^n t dt.\end{aligned}\quad (30)$$

We first of all evaluate the neutrix limit of integral $\int_{\varepsilon}^1 t^{-i} \ln^n t dt$ for $i = 1, 2, \dots$ and $n = 1, 2, \dots$

It follows from Lemma 3.2 that

$$\text{N-lim}_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 t^{-i} \ln^n t dt = -\frac{n!}{(i-1)^{n+1}} \quad (i > 1). \quad (31)$$

For $i = 1$, we have

$$\int_{\varepsilon}^1 t^{-1} \ln^n t dt = O(\varepsilon).$$

Next

$$\begin{aligned}\int_{\varepsilon}^1 \frac{\ln^n t}{1-t} dt &= \int_0^1 \frac{\ln^n t}{1-t} dt = \sum_{k=0}^{\infty} \int_0^1 t^k \ln^n t dt \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k n!}{(k+1)^{n+1}} \\ &= (-1)^n n! \zeta(n+1),\end{aligned}\quad (32)$$

where

$$\int_0^1 t^k \ln^n t dt = \frac{(-1)^k n!}{(k+1)^{n+1}}.$$

It now follows from equations (30), (31) and (32) that

$$\begin{aligned}\psi^{(n)}(-m) &= -\text{N-lim}_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 \frac{t^{-m-1} \ln^n t}{1-t} dt \\ &= -\sum_{i=1}^{m+1} \text{N-lim}_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 t^{-i} \ln^n t dt - \text{N-lim}_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 \frac{\ln^n t}{(1-t)} dt. \\ &= \sum_{i=1}^m \frac{n!}{i^{n+1}} + (-1)^{n+1} n! \zeta(n+1)\end{aligned}$$

implying equation (29).

Note that the digamma function $\psi(x)$ can be defined by

$$\psi(x) = -\gamma + \text{N-lim}_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 \frac{1-t^{x-1}}{1-t} dt \quad (33)$$

for all x .

Also note that using Lemma 3.1 we have

$$\begin{aligned} \int_{\varepsilon}^1 \frac{1-t^{-m-1}}{1-t} dt &= -\sum_{i=1}^{m+1} \int_{\varepsilon}^1 t^{-i} dt \\ &= -[\ln 1 - \ln \varepsilon] - \sum_{i=2}^{m+1} \frac{[1 - \varepsilon^{-i+1}]}{-i+1} \end{aligned}$$

and it follows from equation (33) that

$$\begin{aligned} \psi(-m) &= -\gamma + \text{N-}\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 \frac{1-t^{-m-1}}{1-t} dt = -\gamma + \sum_{i=1}^m i^{-1} \\ &= -\gamma + \phi(m) \end{aligned} \tag{34}$$

which was also obtained in [5] and [9]. Equation (34) can be regarded as the definition of the digamma function for negative integers.

References

- [1] M. Abramowitz and I. Stegun, Hand of Mathematical Functions, Dover Publications, Newyork, 1965.
- [2] J.G. van der Corput, Introduction to the neutrix calculus, J. Analyse Math., **7**(1959), 291-398.
- [3] B. Fisher, On defining the incomplete gamma function $\gamma(-m, x_-)$, Integral Trans. Spec. Funct., 15(6)(2004), 467-476.
- [4] B. Fisher, B. Jolevska-Tuneska and A. Kılıçman, On defining the incomplete gamma function, Integral Trans. Spec. Funct., 14(4)(2003), 293-299.
- [5] Fisher, B. and Kuribayashi, Y., Some results on the Gamma function, J. Fac. Ed. Tottori Univ. Mat. Sci., 3(2)(1988) 111-117.
- [6] W. Gautschi, the incoplete gamma function since Tricomi. Tricomi's Ideas Contemporary Applied Mathematics. Number 147 in Atti dei Convegni Lincei. Roma : Accademia nazionale dei Lincel, Roma, Italy, 1998, 203-237.
- [7] I. M. Gel'fand and G. E. Shilov, Generalized Functions, Vol. I., Academic Press, Newyork/London, (1964).
- [8] I. S. Gradshteyn, and I. M. Ryzhik, Tables of integrals, Series, and Products, Academic Press, San Diego, 2000.

- [9] Jolevska-Tuneska, B. and Jolevski, I., Some results on the digamma function, *Appl. Math. Inf. Sci.*, 7(2013) 167-170.
- [10] Kölbig, K. S., On the integral $\int_0^1 x^{\nu-1}(1-x)^{-\lambda} \ln^m x dx$, *J. Comput. Appl. Math.*, 18(1987) 369-394.
- [11] E. Özçağ, İ. Ege, H. Gürçay and B. Jolevska-Tuneska, Some remarks on the incomplete gamma function, in: Kenan Taş et al. (Eds.), *Mathematical Methods in Engineering*, Springer, Dordrecht, 2007, pp. 97-108.
- [12] Özçağ, E., Ege, İ., Gürçay, H. and Jolevska-Tuneska, B., On partial derivatives of the incomplete beta function, *Appl. Math. Lett.*, 21(2008) 675-681.
- [13] I. Thompson, A note on real zeros of the incomplete gamma function, *Integral Trans. Spec. Funct.*, 23(6)(2012), 445-453.
- [14] I. Thompson, Algorithm 926: Incomplete Gamma Function with Negative Arguments, *ACM Trans. Math. Softw.* 39(2)(2013), 9 pp.
- [15] S. Winitzki, computing the incomplete gamma function to arbitrary precision, In *Proceedings of International Conference on Computational Science and Its Applications (ICCSA'03)*. *Lecturer Notes in Computer Science*, Vol. 2667 (2003), 790-798.

Emin Özçağ

Department of Mathematics
 Hacettepe University, Beytepe, Ankara, Turkey
 e-mail: ozcag1@hacettepe.edu.tr

İnci Ege

Department of Mathemacis
 Adnan Menderes University, Aydın, Turkey
 e-mail : iege@adu.edu.tr