

On the Roman Bondage Number of Graphs on surfaces

Vladimir Samodivkin

Department of Mathematics, University of Architecture Civil Engineering and Geodesy
Hristo Smirnenski 1 Blv., 1046 Sofia, Bulgaria
vlsam_fte@uacg.bg

July 16, 2018

Abstract

A Roman dominating function on a graph G is a labeling $f : V(G) \rightarrow \{0, 1, 2\}$ such that every vertex with label 0 has a neighbor with label 2. The Roman domination number, $\gamma_R(G)$, of G is the minimum of $\sum_{v \in V(G)} f(v)$ over such functions. The Roman bondage number $b_R(G)$ is the cardinality of a smallest set of edges whose removal from G results in a graph with Roman domination number not equal to $\gamma_R(G)$. In this paper we obtain upper bounds on $b_R(G)$ in terms of (a) the average degree and maximum degree, and (b) Euler characteristic, girth and maximum degree. We also show that the Roman bondage number of every graph which admits a 2-cell embedding on a surface with non negative Euler characteristic does not exceed 15.

Keywords: Roman domination, Roman bondage number, girth, average degree, Euler characteristic.
MSC 2010 05C69.

1 Introduction

All graphs considered in this article are finite, undirected, without loops or multiple edges. We denote the vertex set and the edge set of a graph G by $V(G)$ and $E(G)$, respectively. Let P_n denote the path with n vertices. For any vertex x of a graph G , $N_G(x)$ denotes the set of all neighbors of x in G , $N_G[x] = N_G(x) \cup \{x\}$ and the degree of x is $d_G(x) = |N_G(x)|$. The minimum and maximum degrees of the graph G are denoted by $\delta(G)$ and $\Delta(G)$, respectively. For a graph G , let $x \in X \subseteq V(G)$. A vertex $y \in V(G)$ is a X -private neighbor of x if $N_G[y] \cap X = \{x\}$. The X -private neighborhood of x , denoted $pn_G[x, X]$, is the set of all X -private neighbors of x . An orientable compact 2-manifold \mathbb{S}_h or orientable surface \mathbb{S}_h (see [13]) of genus h is obtained from the sphere by adding h handles. Correspondingly, a non-orientable compact 2-manifold \mathbb{N}_q or non-orientable surface \mathbb{N}_q of genus q is obtained from the sphere by adding q crosscaps. Compact 2-manifolds are called simply surfaces throughout the paper. The Euler characteristic is defined by $\chi(\mathbb{S}_h) = 2 - 2h$, $h \geq 0$, and $\chi(\mathbb{N}_q) = 2 - q$, $q \geq 1$. The Euclidean plane \mathbb{S}_0 , the projective plane \mathbb{N}_1 , the torus \mathbb{S}_1 , and the Klein bottle \mathbb{N}_2 are all the surfaces of nonnegative Euler characteristic.

A dominating set for a graph G is a subset $D \subseteq V(G)$ of vertices such that every vertex not in D is adjacent to at least one vertex in D . The minimum cardinality of a dominating set is called the domination number of G and is denoted by $\gamma(G)$. A variation of domination called Roman domination was introduced by ReVelle [10, 11]. Also see ReVelle and Rosing [12] for an integer programming formulation of the problem. The concept of Roman domination can be formulated in terms of graphs. A Roman dominating function

(RDF) on a graph G is a vertex labeling $f : V(G) \rightarrow \{0, 1, 2\}$ such that every vertex with label 0 has a neighbor with label 2. For a RDF f , let $V_i^f = \{v \in V(G) : f(v) = i\}$ for $i = 0, 1, 2$. Since this partition determines f , we can equivalently write $f = (V_0^f; V_1^f; V_2^f)$. The weight $f(V(G))$ of a RDF f on G is the value $\sum_{v \in V(G)} f(v)$, which equals $|V_1^f| + 2|V_2^f|$. The Roman domination number of a graph G , denoted by $\gamma_R(G)$, is the minimum weight of a Roman dominating function on G . A function $f = (V_0^f; V_1^f; V_2^f)$ is called a γ_R -function on G , if it is a Roman dominating function and $f(V(G)) = \gamma_R(G)$. One measure of the stability of the Roman domination number of a graph G under edge removal is the Roman bondage number $b_R(G)$, defined by Jafari Rad and Volkmann in [7], as the cardinality of a smallest set of edges whose removal from G results in a graph with Roman domination number not equal to $\gamma_R(G)$. For more information we refer the reader to [1, 2, 4, 7, 9, 15].

In this paper we obtain upper bounds for $b_R(G)$ in terms of (a) average degree and maximum degree, and (b) Euler characteristic, girth and maximum degree. We also prove that the Roman bondage number of every graph which admits a 2-cell embedding on a surface with non negative Euler characteristic does not exceed 15.

2 Some known results

The following results are important for our investigations.

Theorem A. *Let G be a connected graph embeddable on a surface \mathbb{M} whose Euler characteristic $\chi(\mathbb{M})$ is nonnegative and let $\delta(G) \geq 5$. Then G contains an edge $e = xy$ with $d_G(x) + d_G(y) \leq 11$ if one of the following holds:*

- (i) (Wernicke [18] and Sanders [17], respectively) $\mathbb{M} \in \{\mathbb{S}_0, \mathbb{N}_1\}$.
- (ii) (Jendrol' and Voss [6]) $\mathbb{M} \in \{\mathbb{S}_1, \mathbb{N}_2\}$ and $\Delta(G) \geq 7$.

Lemma B (Rad and Volkmann [8]). *If G is a graph, then $\gamma_R(G) \leq \gamma_R(G - e) \leq \gamma_R(G) + 1$ for any edge $e \in E(G)$.*

According to the effects of vertex removal on the Roman domination number of a graph G , let

- $V_R^+(G) = \{v \in V(G) \mid \gamma_R(G - v) > \gamma_R(G)\}$,
- $V_R^-(G) = \{v \in V(G) \mid \gamma_R(G - v) < \gamma_R(G)\}$,
- $V_R^0(G) = \{v \in V(G) \mid \gamma_R(G - v) = \gamma_R(G)\}$.

Clearly $\{V_R^-(G), V_R^0(G), V_R^+(G)\}$ is a partition of $V(G)$.

Theorem C (Rad and Volkmann [8]). *Let G be a graph of order at least 2.*

- (i) *If $v \in V_R^+(G)$ then for every γ_R -function $f = (V_0^f; V_1^f; V_2^f)$ on G , $|pn_G[v, V_2^f] \cap V_0^f| \geq 3$ and $f(v) = 2$.*
- (ii) *For any vertex $u \in V(G)$, $\gamma_R(G) - 1 \leq \gamma_R(G - u)$.*

Theorem D (Hansberg, Rad and Volkmann [5]). *Let v be a vertex of a graph G . Then $\gamma_R(G - v) < \gamma_R(G)$ if and only if there is a γ_R -function $f = (V_0, V_1, V_2)$ on G such that $v \in V_1$.*

Theorem E (Rad and Volkmann [7]). *If G is a graph, and x, y, z is a path of length 2 in G , then*

$$b_R(G) \leq d_G(x) + d_G(y) + d_G(z) - 3 - |N_G(x) \cap N_G(y)|.$$

The average degree $ad(G)$ of a graph G is defined as $ad(G) = 2|E(G)|/|V(G)|$.

Theorem F (Hartnell and Rall [3]). *For any connected nontrivial graph G , there exists a pair of vertices, say u and v , that are either adjacent or at distance 2 from each other, with the property that $d_G(u) + d_G(v) \leq 2ad(G)$.*

The girth of a graph G is the length of a shortest cycle in G ; the girth of a forest is ∞ .

Lemma G (Samodivkin [16]). *Let G be a connected graph embeddable on a surface \mathbb{M} whose Euler characteristic χ is as large as possible and let the girth of G is $k < \infty$. Then:*

$$ad(G) \leq \frac{2k}{k-2} \left(1 - \frac{\chi}{|V(G)|}\right).$$

Given a graph G of order n , let \widehat{G} be the graph of order $5n$ obtained from G by attaching the central vertex of a copy of P_5 , to each vertex of G .

Lemma H (S.Akbari, M. Khatirinejad and S. Qajar [1]). *Let G be a graph of order n , $n \geq 2$. Then $\gamma(\widehat{G}) = 2n$, $\gamma_R(\widehat{G}) = 4n$ and $b_R(\widehat{G}) = \delta(G) + 2$.*

3 Upper bounds

A graph G of order at least two is Roman domination vertex critical if removing any vertex of G decreases the Roman domination number. By \mathcal{RCV} we denote the class of all Roman domination vertex critical graphs. Results on this class can be found in Rad and Volkmann [8] and Hansberg et al. [5].

Theorem 1. *Let G be a connected graph.*

- (i) *If $V_R^-(G) \neq V(G)$ then $b_R(G) \leq \min\{d_G(u) - \gamma_R(G - u) + \gamma_R(G) \mid u \in V_R^0(G) \cup V_R^+(G)\} \leq \Delta(G)$.*
- (ii) *If $b_R(G) > \Delta(G)$ then G is in \mathcal{RCV} .*

Remark 2. Let G be any connected graph of order $n \geq 2$. Denote by S the set of all vertices of \widehat{G} each of which is adjacent to a vertex of degree 1. Then $f = (V(\widehat{G}) - S; \emptyset; S)$ is a RDF on \widehat{G} . Since the weight of f is $4n$, by Lemma H it follows that f is a γ_R -function on \widehat{G} . Theorem C(i) now implies $V(\widehat{G}) = V_R^-(\widehat{G}) \cup V_R^0(\widehat{G})$. Since $\gamma_R(P_5) = 4$ and since the central vertex of P_5 is in $V_R^0(P_5)$, $V(G) \subset V_R^0(\widehat{G})$. Labelling the vertices of each P_5 of \widehat{G} with $(1, 1, 0, 2, 0)$ yields a γ_R -function on \widehat{G} . It follows by Theorem D that $V(G) = V_R^0(\widehat{G})$. All this together with Lemma H shows that the bound in Theorem 1(i) is attainable for all graphs \widehat{G} . Furthermore, for any graph \widehat{G} the bound in Theorem E is attainable too.

To prove Theorem 1, we need the following lemma:

Lemma 3. *Let G be a connected graph. For any subset $U \subsetneq V(G)$, let E_U denote the set of all edges each of which joins U and $V(G) - U$.*

- (i) *If $v \in V_R^0(G) \cup V_R^+(G)$ then $\gamma_R(G - E_{\{v\}}) > \gamma_R(G)$.*
- (ii) *If $x \in V_R^+(G)$ then $1 \leq \gamma_R(G - x) - \gamma_R(G) \leq d_G(x) - 2$ and for any subset $S \subseteq E_{\{x\}}$ with $|S| \geq d_G(x) - \gamma_R(G - v) + \gamma_R(G)$, $\gamma_R(G - S) > \gamma_R(G)$.*

Proof. (i) We have $\gamma_R(G - E_{\{v\}}) = \gamma_R(G - v) + 1 > \gamma_R(G)$.

(ii) Denote $p = \gamma_R(G - x) - \gamma_R(G)$. Let f be any γ_R -function on G . Since $p > 0$, by Theorem C(i) it follows that $f(x) = 2$. Hence $h = (V_0^f - N_G(x); V_1^f \cup (N_G(x) - V_2^f); V_2^f - \{x\})$ is a RDF on $G - x$. But then $\gamma_R(G) + p = \gamma_R(G - x) \leq h(V(G - x)) \leq \gamma_R(G) + d_G(x) - 2$. Hence $1 \leq p \leq d_G(x) - 2$. For any set $S \subseteq E_{\{x\}}$ with $|S| \geq d_G(x) - p$ we have $\gamma_R(G - S) \geq \gamma_R(G - E_{\{x\}}) - |E_{\{x\}}| + |S| \geq (\gamma_R(G - x) + 1) - d_G(x) + (d_G(x) - p) = \gamma_R(G) + 1$, where the first inequality follows from Lemma B. \square

Proof of Theorem 1. (i) The result follows immediately by Lemma 3.

(ii) Immediately by (i). \square

Rad and Volkman [9] as well as Akbari et al. [1] gave upper bounds on the Roman bondage number of planar graphs. Upper bounds on the Roman bondage number of graphs 2-cell embeddable on topological surfaces in terms of orientable/non orientable genus and maximum degree, are obtained by the present author in [15].

Theorem 4. *Let G be a connected graph with $\Delta(G) \geq 2$.*

(i) *Then $b_R(G) \leq 2ad(G) + \Delta(G) - 3$.*

(ii) *Let G be embeddable on a surface \mathbb{M} whose Euler characteristic χ is as large as possible. If G has order n and girth $k < \infty$ then:*

$$b_R(G) \leq \frac{4k}{k-2} \left(1 - \frac{\chi}{n}\right) + \Delta(G) - 3 \leq -\frac{12\chi}{n} + \Delta(G) + 9.$$

Proof. (i) If G is a complete graph then the result is obvious. Hence we may assume G has nonadjacent vertices. Theorem F implies that there are 2 vertices, say x and y , that are either adjacent or at distance 2 from each other, with the property that $d_G(x) + d_G(y) \leq 2ad(G)$. Since G is connected and $\Delta(G) \geq 2$, there is a vertex z such that xyz or xzy is a path. In either case by Theorem E we have $b_R(G) \leq d_G(x) + d_G(y) + d_G(z) - 3 \leq 2ad(G) + \Delta(G) - 3$.

(ii) Lemma G and (i) together imply the result. \square

Remark 5. Let \mathbb{M} be a surface. Denote $\delta_{max}^{\mathbb{M}} = \max\{\delta(H) \mid \text{a graph } H \text{ is 2-cell embedded in } \mathbb{M}\}$. Let G be a connected graph 2-cell embeddable on \mathbb{M} and $\delta(G) = \delta_{max}^{\mathbb{M}}$. By Lemma H it immediately follows $b_R(\widehat{G}) = \delta_{max}^{\mathbb{M}} + 2$. Note that (a) if $\chi(\mathbb{M}) \leq 1$ then $\delta_{max}^{\mathbb{M}} \leq \left\lfloor (5 + \sqrt{49 - 24\chi(\mathbb{M})})/2 \right\rfloor$ (see Sachs [14], pp. 226-227), and (b) it is well known that $\delta_{max}^{\mathbb{S}_0} = \delta_{max}^{\mathbb{N}_1} = 5$ and $\delta_{max}^{\mathbb{N}_2} = \delta_{max}^{\mathbb{N}_3} = \delta_{max}^{\mathbb{S}_1} = 6$.

In [1], Akbari, Khatirinejad and Qajar recently prove that $b_R(G) \leq 15$ provided G is a planar graph. As the next result shows, more is true.

Theorem 6. *Let G be a connected graph 2-cell embedded on a surface \mathbb{M} with non negative Euler characteristic and let $\Delta(G) \geq 2$. Then $b_R(G) \leq 15$.*

Proof. If $2 \leq \Delta(G) \leq 6$ then $b_R(G) \leq 3\Delta(G) - 3 \leq 15$, because of Theorem E. So, assume $\Delta(G) \geq 7$. Denote $V_{\leq 5} = \{v \in V(G) \mid d_G(v) \leq 5\}$ and $G_{\geq 6} = G - V_{\leq 5}$. Since $\chi(\mathbb{M}) \geq 0$, $\delta(G) \leq 6$ (see Remark 5). If $\delta(G) = 6$ then G is a 6-regular triangulation on the torus or in the Klein Bottle, a contradiction with $\Delta(G) \geq 7$. So, $\delta(G) \leq 5$ and then $V_{\leq 5}$ is not empty. Since $G_{\geq 6}$ is embedded without crossings on \mathbb{M} and $\chi(\mathbb{M}) \geq 0$, there is a vertex $u \in V(G_{\geq 6})$ with $d_{G_{\geq 6}}(u) \leq 6$. If u has exactly 2 neighbors belonging to $V_{\leq 5}$ then again by Theorem E, $b_R(G) \leq 15$. Now let all $v_1, v_2, v_3 \in V_{\leq 5}$ be adjacent to u . Denote by E_1 the set of all edges of G which are incident to at least one of v_1, v_2 and v_3 . Since v_1, v_2 and v_3 are isolated vertices in $G - E_1$, for any γ_{PR} -function g on $G - E_1$, $g(v_1) = g(v_2) = g(v_3) = 1$. Define now $f : V(G) \rightarrow \{0, 1, 2\}$ by $f(v_1) = f(v_2) = f(v_3) = 0$, $f(u) = 2$ and $f(v) = g(v)$ for every $v \in V(G) - \{u, v_1, v_2, v_3\}$. Clearly f is a RDF on G with $\gamma_R(G) \leq f(V(G)) < g(V(G - E_1)) = \gamma_R(G - E_1)$. Thus, $b_R(G) \leq |E_1| \leq d_G(v_1) + d_G(v_2) + d_G(v_3) \leq 15$.

So, it remains to consider the case where each vertex of degree at most 6 in $G_{\geq 6}$ has no more than one neighbor in $V_{\leq 5}$. It immediately follows that $\delta(G_{\geq 6}) \geq 5$. First assume $\delta(G_{\geq 6}) = 5$. By Theorem A it follows that there is an edge $xy \in E(G_{\geq 6})$ such that $d_{G_{\geq 6}}(x) + d_{G_{\geq 6}}(y) \leq 11$. Hence $d_G(x) + d_G(y) \leq 13$. Let without loss of generality $d_{G_{\geq 6}}(x) \leq d_{G_{\geq 6}}(y)$. Then x has exactly one neighbor in $V_{\leq 5}$, say v . By Theorem E applied to the path v, x, y we have $b_R(G) \leq 5 + 13 - 3 = 15$. Now let $\delta(G_{\geq 6}) \geq 6$. But then $G_{\geq 6}$ is a 6-regular triangulation on the torus or in the Klein bottle. Since $\Delta(G) \geq 7$, $G \neq G_{\geq 6}$ and there is a path x, y, z in G , where $d_G(z) \leq 5$, and both x and y are in $V(G_{\geq 6})$. Since clearly $|N(x) \cap N(y)| \geq 2$, again using Theorem E we obtain $b_{PR}(G) \leq 7 + 7 + 5 - 3 - 2 = 14$. \square

References

- [1] S.Akbari, M. Khatirinejad and S. Qajar, *A Note on the Roman Bondage Number of Planar Graphs*, Graphs and Combinatorics, 29(2013), 327–331
- [2] A. Bahremandpour, Fu-Tao Hu, S.M. Sheikholeslami and Jun-Ming Xu. On the Roman Bondage number of a graph, Discrete Mathematics, Algorithms and Applications Vol. 5, No. 1 (2013) 1350001 (15 pages)
- [3] B.L. Hartnell and D.F. Rall, *A bound on the size of a graph with given order and bondage number*, Discrete Math. 197/198 (1999), 409–413.
- [4] F.-T. Hu and J.-M. Xu, Roman bondage numbers of some graphs, Australas. J. Comb. 58(1) (2014), 106–118
- [5] A. Hansberg, N.J. Rad and L. Volkmann, Vertex and edge critical Roman domination in graphs, To appear in Utilitas Mathematica.
- [6] S. Jendrol', H.-J. Voss, Light subgraphs of order at most 3 in large maps of minimum degree 5 on compact 2-manifolds, European Journal of Combinatorics 26 (2005) 457–471.
- [7] N. J. Rad and L. Volkmann, Roman bondage in graphs, Discussiones Mathematicae: Graph Theory; 2011, Vol. 31 Issue 4, 763–773
- [8] N. J. Rad and L. Volkmann, Changing and unchanging the Roman domination number of a graph, Util. Math. 89 (2012) 79-95
- [9] N. J. Rad and L. Volkmann, *On the roman bondage number of planar graphs*. Graphs Comb. 27 (2011), 531–538.
- [10] C.S. ReVelle, Can you protect the Roman Empire? Johns Hopkins Mag 49(2):40
- [11] ReVelle CS (1997b) Test your solution to "Can you protect the Roman Empire" Johns Hopkins Mag 49(3):70
- [12] ReVelle CS, Rosing KE (2000) Defendens Imperium Romanum: a classical problem in military. Am Math Mon 107(7):585–594
- [13] G. Ringel, Map Color Theorem, Springer-Verlag, Berlin, 1974.
- [14] H. Sachs, Einführung in die Theorie der endlichen Graphen, Teil II, Teubner, Leipzig, 1972 (in German).
- [15] V. Samodivkin, Upper bounds for the domination subdivision and bondage numbers of graphs on topological surfaces, Czechoslovak Math. Journal, 63(1), 2013, 191–204.
- [16] V. Samodivkin, Note on the bondage number of graphs on topological surfaces, Australasian J. Combin., 56 (2013), 183–186.
- [17] D. P. Sanders, On Light Edges and Triangles in Projective Planar Graphs, J. of Graph Theory, 21(1996), 335-342.
- [18] P. Wernicke, Über den kartographischen Vierfarbensatz, Math. Ann., 58 (1904), 413 -426