

THE GINZBURG-LANDAU FUNCTIONAL WITH VANISHING MAGNETIC FIELD

BERNARD HELFFER AND AYMAN KACHMAR

ABSTRACT. We study the infimum of the Ginzburg-Landau functional in the case of a vanishing external magnetic field in a two dimensional simply connected domain. We obtain an energy asymptotics which is valid when the Ginzburg-Landau parameter is large and the strength of the external field is comparable with the third critical field. Compared with the known results when the external magnetic field does not vanish, we show in this regime a concentration of the energy near the zero set of the external magnetic field. Our results complete former results obtained by K. Attar and X-B. Pan–K-H. Kwek.

1. INTRODUCTION

In a two dimensional bounded and simply connected domain Ω with smooth boundary, the Ginzburg-Landau functional is defined over configurations $(\psi, \mathbf{A}) \in H^1(\Omega; \mathbb{C}) \times H^1(\Omega; \mathbb{R}^2)$ by,

$$\mathcal{E}(\psi, \mathbf{A}) = \int_{\Omega} e_{\kappa, H}(\psi, \mathbf{A}) dx \quad (1.1)$$

where

$$e_{\kappa, H}(\psi, \mathbf{A}) := |(\nabla - i\kappa H \mathbf{A})\psi|^2 - \kappa^2 |\psi|^2 + \frac{\kappa^2}{2} |\psi|^4 + (\kappa H)^2 |\operatorname{curl} \mathbf{A} - B_0|^2.$$

The modulus of the wave function ψ measures the density of the superconducting electrons; the curl of the vector field \mathbf{A} measures the induced magnetic field; the parameter H measures the intensity of the external magnetic field and the parameter κ ($\kappa > 0$) is a characteristic of the superconducting material; dx is the Lebesgue measure $dx_1 dx_2$. The function B_0 represents the profile of the external magnetic field in Ω and is allowed to vanish non-degenerately on a smooth curve. We suppose that B_0 is defined and C^∞ in a neighborhood of $\bar{\Omega}$ and satisfies,

$$|B_0| + |\nabla B_0| \geq c > 0 \quad \text{in } \bar{\Omega}, \quad (1.2)$$

and that the set

$$\Gamma = \{x \in \bar{\Omega} : B_0(x) = 0\} \quad (1.3)$$

consists of a finite number of simple smooth curves.

We also assume that:

$$\Gamma \cap \partial\Omega \text{ is a finite set.} \quad (1.4)$$

The ground state energy of the functional is,

$$E_{\text{gs}}(\kappa, H) = \inf\{\mathcal{E}(\psi, \mathbf{A}) : (\psi, \mathbf{A}) \in H^1(\Omega; \mathbb{C}) \times H^1(\Omega; \mathbb{R}^2)\}. \quad (1.5)$$

We focus on the regime where H satisfies

$$H = \sigma \kappa^2, \quad \sigma \in (0, \infty). \quad (1.6)$$

Our results allow for σ to be a function of κ satisfying $\sigma \gg \kappa^{-1}$. Earlier results corresponding to vanishing magnetic fields have been obtained recently in [1, 2]. The assumption on the strength of the magnetic field was $H \leq C\kappa$, where C is a constant. In the regime of large κ , K. Attar has obtained in [1, 2] parallel results to those known for the constant magnetic field in [25]. However, it is proved in [1] that if

$$H = b\kappa, \quad (1.7)$$

Date: July 10, 2021.

and b is a constant, then when b is large enough, the energy and the superconducting density are concentrated near the set Γ with a length scale $\frac{1}{b}$. Essentially, that is a consequence of the following asymptotics of the energy ($\kappa \rightarrow \infty$),

$$E_{\text{gs}}(\kappa, H) = \kappa^2 \int_{\Omega} g\left(\frac{H}{\kappa}|B_0(x)|\right) dx + o\left(\kappa H \left|\ln \frac{\kappa}{H}\right| + 1\right), \quad (1.8)$$

which is valid under the relaxed assumption that

$$\Lambda_1 \kappa^{1/3} \leq H \leq \Lambda_2 \kappa, \quad (1.9)$$

Λ_1 and Λ_2 being positive constants. In particular, the assumption in (1.9) covers the situation in (1.7). The function $g(b)$ appears in the analysis of the two and three dimensional Ginzburg-Landau functional with constant magnetic field, [25, 8]. It is associated with some effective model energy. The function $g(b)$ will play a central role in this paper and its definition will be recalled later in this text (see (3.53)).

One purpose of this paper is to give a precise description of the aforementioned concentration of the order parameter and the energy when $\sigma \gg 1$, thereby leading to the assumption in (1.6).

The leading order term of the ground state energy in (1.5) is expressed via the quantity $E(\cdot)$ introduced in Theorem 3.8 below. The function $(0, \infty) \ni L \mapsto E(L)$ is a continuous function satisfying the following properties:

- $E(L)$ is defined via a reduced Ginzburg-Landau energy in the strip (this energy is introduced in (3.14)).
- $E(L) = 0$ iff $L \geq \lambda_0^{-3/2}$, where λ_0 is a universal constant defined as the bottom of the spectrum of a Montgomery operator, see (3.4).
- As $L \rightarrow 0_+$, the expected asymptotic behavior of $E(L)$ is like $L^{-4/3}$.

Through this text, we use the following notation. If A and B are two positive quantities, then

- $A \ll B$ means $A = \delta(\kappa)B$ and $\delta(\kappa) \rightarrow 0$ as $\kappa \rightarrow \infty$;
- $A \lesssim B$ means $A \leq CB$ and $C > 0$ is a constant independent of κ ;
- $A \gg B$ means $B \ll A$, and $A \gtrsim B$ means $B \lesssim A$;
- $A \approx B$ means $c_1 B \leq A \leq c_2 B$, $c_1 > 0$ and $c_2 > 0$ are constants independent of κ .

The main result in this paper is:

Theorem 1.1. *Suppose that the function B_0 satisfies Assumptions (1.2) and (1.3). Let $b : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a function satisfying*

$$\lim_{\kappa \rightarrow \infty} b(\kappa) = \infty \quad \text{and} \quad \limsup_{\kappa \rightarrow \infty} \kappa^{-1} b(\kappa) < \infty. \quad (1.10)$$

Suppose that

$$H = b(\kappa)\kappa.$$

As $\kappa \rightarrow \infty$, the ground state energy in (1.5) satisfies:

- (1) *If $b(\kappa) \gg \kappa^{1/2}$, then*

$$E_{\text{gs}}(\kappa, H) = \kappa \left(\int_{\Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) + \frac{\kappa^3}{H} o(1), \quad (1.11)$$

where ds is the arc-length measure in Γ .

- (2) *If $b(\kappa) \lesssim \kappa^{1/2}$, then,*

$$E_{\text{gs}}(\kappa, H) = \kappa^2 \int_{\Omega} g\left(\frac{H}{\kappa}|B_0(x)|\right) dx + \frac{\kappa^3}{H} o(1). \quad (1.12)$$

Remark 1.2. There is a small gap between the two regimes considered above. Hence it would be interesting to show that the two asymptotics match in this intermediate zone.

Remark 1.3. As we shall see in Section 2, Pan and Kwek [20] prove that if H is larger than a critical value $H_{c_3}(\kappa)$, then the minimizers of the functional in (1.1) are trivial and the ground state energy is $E_{\text{gs}}(\kappa, H) = 0$. Furthermore, the value of $H_{c_3}(\kappa)$ as given in [20] admits, as $\kappa \rightarrow \infty$, the following asymptotics

$$H_{c_3}(\kappa) \sim c_0 \kappa^2,$$

where c_0 is a universal explicit constant. As such, the assumption on the magnetic field in Theorem 1.1 is significant when $b(\kappa)\kappa \leq H \leq M\kappa^2$ and $M \in (0, c_0]$ is a constant. Note also that our theorem gives a bridge between the situations studied by Attar in [1, 2] and Pan-Kwek in [20].

Remark 1.4. As long as the intensity of the external magnetic field satisfies $\kappa \ll H \leq M\kappa^2$ and $M \in (0, c_0)$, the remainder term appearing in Theorem 1.1 is of lower order compared with the principal term. The function $g(b)$ is bounded and vanishes when $b \geq 1$. Accordingly,

$$\int_{\Omega} g\left(\frac{H}{\kappa}|B_0(x)|\right) dx = \int_{\{|B_0(x)| < \frac{\kappa}{H}\}} g\left(\frac{H}{\kappa}|B_0(x)|\right) dx \approx \frac{\kappa}{H}.$$

We shall see in Theorem 3.12 that,

$$\left(|\nabla B_0(x)|\frac{H}{\kappa^2}\right)^{1/3} E\left(|\nabla B_0(x)|\frac{H}{\kappa^2}\right) \approx \frac{\kappa^2}{H}.$$

Along the proof of Theorem 1.1, we obtain:

Theorem 1.5. *Suppose that the function B_0 satisfies Assumptions (1.2) and (1.3). Let $b : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a function satisfying (1.10). Suppose that*

$$H = b(\kappa)\kappa$$

and that (ψ, \mathbf{A}) is a minimizer of the functional in (1.1).

As $\kappa \rightarrow \infty$, there holds:

- (1) **Estimate of the magnetic energy.**

$$\kappa^2 H^2 \int_{\Omega} |\text{curl } \mathbf{A} - B_0|^2 dx = \frac{\kappa^3}{H} o(1).$$

- (2) **Estimate of the local energy.**

Let $D \subset \Omega$ be an open set with a smooth boundary such that $\partial D \cap \Gamma$ is a finite set.

- (a) *If $b(\kappa) \gg \kappa^{1/2}$, then*

$$\begin{aligned} \mathcal{E}_0(\psi, \mathbf{A}; D) &:= \int_D \left(|(\nabla - i\kappa H \mathbf{A})\psi|^2 - \kappa^2 |\psi|^2 + \frac{\kappa^2}{2} |\psi|^4 \right) dx \\ &= \kappa \left(\int_{D \cap \Gamma} \left(|\nabla B_0(x)|\frac{H}{\kappa^2} \right)^{1/3} E\left(|\nabla B_0(x)|\frac{H}{\kappa^2}\right) ds(x) \right) + \frac{\kappa^3}{H} o(1). \end{aligned} \quad (1.13)$$

- (b) *If $1 \ll b(\kappa) \lesssim \kappa^{1/2}$, then*

$$\mathcal{E}_0(\psi, \mathbf{A}; D) = \kappa^2 \int_D g\left(\frac{H}{\kappa}|B_0(x)|\right) dx + \frac{\kappa^3}{H} o(1).$$

- (3) **Concentration of the order parameter.**

Let $D \subset \Omega$ be an open set with a smooth boundary such that $\partial D \cap \Gamma$ is a finite set.

- (a) *If $b(\kappa) \gg \kappa^{1/2}$, then*

$$\int_D |\psi(x)|^4 dx = -\frac{2}{\kappa} \left(\int_{D \cap \Gamma} \left(|\nabla B_0(x)|\frac{H}{\kappa^2} \right)^{1/3} E\left(|\nabla B_0(x)|\frac{H}{\kappa^2}\right) ds(x) \right) + \frac{\kappa}{H} o(1).$$

(b) If $1 \ll b(\kappa) \lesssim \kappa^{1/2}$, then

$$\int_D |\psi(x)|^4 dx = - \int_D g \left(\frac{H}{\kappa} |B_0(x)| \right) dx + \frac{\kappa}{H} o(1).$$

Remark 1.6. It could be interesting to improve the second term when $D \cap \Gamma = \emptyset$. In Theorem 6.3, we will prove that the L^2 -norm of the order parameter ψ is concentrated near the set Γ , and that ψ exponentially decays as $\kappa \rightarrow \infty$, away from Γ .

Remark 1.7. In Theorem 1.5, the function $o(1)$ is bounded independently of the choice of the minimizer (ψ, \mathbf{A}) . In the first assertion, one can find a bound of $o(1)$ which depends only on the domain Ω and the function B_0 , while in the second and third assertions, the bound depends additionally on the domain D .

2. CRITICAL FIELDS

The identification of the critical *magnetic* fields is an important question regarding the functional in (1.1). This question has an early appearance in physics (see e.g. [11]) and was the subject of a vast mathematical literature in the past two decades. The two monographs [4, 24] contain an extensive review of many important results. In this section, we give a brief *informal* description of critical fields and highlight the importance of the case of a vanishing applied magnetic field.

2.1. Reminder: The constant field case. When the magnetic field B_0 is a (non-zero) constant, three critical values are assigned to the magnetic field H , namely H_{c_1} , H_{c_2} and H_{c_3} . The behavior of minimizers (and critical points) of the functional in (1.1) changes as the parameter H (i.e. magnetic field) crosses the values H_{c_1} , H_{c_2} and H_{c_3} . The identification of these critical values is not easy, especially the value H_{c_2} which is still *loosely* defined.

Let us recall that a critical point (ψ, \mathbf{A}) of the functional in (1.1) is said to be *normal* if $\psi = 0$ everywhere. The critical field $H_{c_3}(\kappa)$ is then defined as the value at which the transition from *normal* to *non-normal* critical points takes place.

The identification of the critical value $H_{c_3}(\kappa)$ is strongly related to the spectral analysis of the magnetic Schrödinger operator with a constant magnetic field and Neumann boundary condition. Suppose that $\Omega \subset \mathbb{R}^2$ is connected, open, has a smooth boundary and the boundary consists of a finite number of connected components, A_0 a vector field satisfying $\text{curl} A_0 = B_0$, the function B_0 is constant and positive, and $\lambda(H\kappa A_0)$ the lowest eigenvalue of the magnetic Schrödinger operator

$$-\Delta_{\kappa H A_0} = -(\nabla - i\kappa H A_0)^2 \quad \text{in } L^2(\Omega), \quad (2.1)$$

with Neumann boundary conditions. It is proved that the function $t \mapsto \lambda(tA_0)$ is monotonic for large values of t , see [4] and the references therein. Grosso modo¹ the critical field H_{c_3} is the unique solution of the equation,

$$\lambda(H_{c_3}(\kappa)\kappa A_0) = \kappa^2. \quad (2.2)$$

In this case, it was shown by Lu-Pan [16] that,

$$\lambda(H\kappa A_0) \sim (H\kappa)B_0\Theta_0, \quad \text{when } H\kappa \gg 1. \quad (2.3)$$

Further improvements of (2.3) are available, see [4] for the state of the art in 2009 and references therein.

As a consequence of (2.2) and (2.3), we get for κ sufficiently large,

$$H_{c_3}(\kappa) \sim \kappa / (\Theta_0 B_0). \quad (2.4)$$

¹Initially (see [16]), one should start by defining four critical values according to locally or globally minimizing solutions. Following the terminology of [4], these are upper or lower, global or local fields. The four fields are proved to be equal in [4].

The second critical field $H_{c_2}(\kappa)$ is usually defined as follows

$$H_{c_2}(\kappa) = \kappa/B_0. \quad (2.5)$$

Notice that this definition of H_{c_2} is asymptotically matching with the following definition²,

$$\lambda^D(H_{c_2}(\kappa)\kappa A_0) = \kappa^2, \quad (2.6)$$

where λ^D is the first eigenvalue of the operator in (2.1), but with Dirichlet boundary condition.

Near $H_{c_2}(\kappa)$, a transition takes place between surface and bulk superconductivity. At the level of the energy, this transition is described in [9].

We recall that $\Theta_0 < 1$. Hence, as expected, $H_{c_2}(\kappa) < H_{c_3}(\kappa)$ for κ sufficiently large. For the identification of the critical field $H_{c_1}(\kappa)$, we refer to Sandier-Serfaty [24]. A natural question is to extend this discussion in the variable magnetic field case (i.e. B_0 is a non-constant function).

2.2. The case of a non vanishing exterior magnetic field. Here we discuss the situation where the magnetic field B_0 is a non-constant function such that $B_0(x) \neq 0$ everywhere in $\overline{\Omega}$. In this case, it is proved by Lu-Pan [17, Theorem 1] that,

$$\lambda(H\kappa A_0) \sim (H\kappa) \min \left(\inf_{x \in \overline{\Omega}} |B_0(x)|, \Theta_0 \inf_{x \in \partial\Omega} |B_0(x)| \right), \quad (2.7)$$

as $H\kappa \rightarrow \infty$. Basically, this leads to consider two cases as follows.

Surface superconductivity. First, we assume that

$$\inf_{x \in \overline{\Omega}} |B_0(x)| > \Theta_0 \inf_{x \in \partial\Omega} |B_0(x)|. \quad (2.8)$$

In this case, the phenomenon of surface superconductivity observed in the constant magnetic field case is preserved. More precisely, the superconductivity starts to appear at the points where $(B_0)_{/\partial\Omega}$ is minimal. The critical value $H_{c_3}(\kappa)$ is still defined by (2.2). If the minima of $(B_0)_{/\partial\Omega}$ are non-degenerate, then the monotonicity of the eigenvalue $\lambda(tA_0)$ for large values of t is established in [22, Section 6]. Consequently, we get when κ is sufficiently large,

$$H_{c_3}(\kappa) \sim \frac{\kappa}{\Theta_0 \inf_{x \in \partial\Omega} |B_0(x)|}. \quad (2.9)$$

Tentatively, one could think to define $H_{c_2}(\kappa)$ either by

$$H_{c_2}(\kappa) = \frac{\kappa}{\inf_{x \in \overline{\Omega}} |B_0(x)|}, \quad (2.10)$$

or by

$$\lambda^D(H_{c_2}(\kappa)\kappa A_0) = \kappa^2, \quad (2.11)$$

where λ^D is the first eigenvalue of the operator in (2.1) with Dirichlet boundary condition. Notice that both formulas agree with their analogues in the constant magnetic field case (see (2.5) and (2.6)). Also, the values of $H_{c_2}(\kappa)$ given in (2.10) or (2.11) asymptotically match as $\kappa \rightarrow \infty$.

In order that the definition of $H_{c_2}(\kappa)$ in (2.11) be consistent, one should prove monotonicity of $t \mapsto \lambda^D(tA_0)$ for large values of t . This will ensure that (2.11) assigns a unique value of $H_{c_2}(\kappa)$. However, such a monotonicity is not proved yet. The definition in (2.10) was proposed in [4].

²Assuming the monotonicity of $t \mapsto \lambda^D(tA_0)$ for t large.

Interior onset of superconductivity. Here we assume that

$$\inf_{x \in \overline{\Omega}} |B_0(x)| < \Theta_0 \inf_{x \in \partial\Omega} |B_0(x)|. \quad (2.12)$$

In this case, the onset of superconductivity near the surface of the domain disappears. If one decreases gradually the intensity of the magnetic field H from ∞ , then superconductivity will start to appear near the minima of the function $|B_0|$, i.e. inside a compact subset of Ω .

In this situation, we have not to distinguish between the critical fields $H_{c_2}(\kappa)$ and $H_{c_3}(\kappa)$, since surface superconductivity is absent here. Consequently, we expect that,

$$H_{c_2}(\kappa) = H_{c_3}(\kappa) \sim \frac{\kappa}{\inf_{x \in \overline{\Omega}} |B_0(x)|}. \quad (2.13)$$

A partial justification of this fact can be done using the linearized Ginzburg-Landau equation near a normal solution. Actually, we may also define $H_{c_3}(\kappa)$ and $H_{c_2}(\kappa)$ as the values verifying (2.2) and (2.6). It should be noticed here that the vector field A_0 satisfies $\text{curl} A_0 = B_0$ and B_0 cannot be constant. Under the assumption (2.12), the known spectral asymptotics (which are actually the same in this case) of the Dirichlet and Neumann eigenvalues will lead us to the asymptotics given in the righthand side of (2.13). Under the additional assumption that $\inf_{x \in \overline{\Omega}} |B_0(x)|$ is attained at a unique minimum in Ω and that this minimum is non degenerate, a complete asymptotics of $\lambda^N(tA_0)$ can be given (see Helffer-Mohamed [15], Helffer-Kordyukov [13, 14], Raymond-Vu Ngoc [23]) and the monotonicity/strong diamagnetism property holds for large values of t (see Chapter 3 in [4]). Hence the definition of $H_{c_3}(\kappa)$ is clear in this case.

Besides the aforementioned linearized calculations, the results of [1] are useful to justify the equality of the critical fields $H_{c_2}(\kappa)$ and $H_{c_3}(\kappa)$ as well as their definition in (2.13).

First, we observe that if C is a positive constant such that $C < \frac{1}{\inf_{x \in \overline{\Omega}} |B_0(x)|}$, and if $H \leq C\kappa$, then the open set $D = \{x \in \Omega : |B_0(x)| < \frac{1}{C}\} \neq \emptyset$ is non-empty. Now, Theorem 1.4 of [1] asserts that,

$$\exists \kappa_0 > 0, \exists \epsilon_D > 0, \int_D |\psi(x)|^4 dx \geq \epsilon_D > 0,$$

for any $\kappa \geq \kappa_0$ and any *minimizer* (ψ, \mathbf{A}) of the functional in (1.1).

Consequently, a minimizer can not be a *normal solution*.

Now, suppose that the constant C satisfies

$$C > \frac{1}{\inf_{x \in \overline{\Omega}} |B_0(x)|}.$$

If $H \geq C\kappa$, then Theorem 1.4 of [1] asserts that any critical point (ψ, \mathbf{A}) of (1.1) satisfies,

$$\lim_{\kappa \rightarrow \infty} \int_{\Omega} |\psi(x)|^4 dx = 0,$$

hence, loosely speaking, critical points are *nearly* normal solutions. However, repeating the proof given in [4, Section 10.4] and using the asymptotics of the first eigenvalue in (2.7), one can get that such critical points are indeed normal solutions.

This discussion shows that the value appearing in the right hand side of (2.13) is indeed critical.

2.3. The case of a vanishing exterior magnetic field. We now discuss the case when B_0 vanishes along a curve, first considered in [20] and then in [1]. We assume that

$$|B_0| + |\nabla B_0| \neq 0 \text{ in } \overline{\Omega}, \quad (2.14)$$

which ensures that B_0 vanishes *non-degenerately*.

At each point of $B_0^{-1}(0) \cap \Omega$, Pan-Kwek [20] introduce a toy model (a Montgomery operator parameterized by the intensity of the magnetic field at this point) whose ground state energy, denoted by λ_0 , captures the ‘local’ ground state energy of the Schrödinger operator in (2.1).

Similarly, at every point x of $B_0^{-1}(0) \cap \partial\Omega$, a toy operator is defined on \mathbb{R}_+^2 parameterized (up to unitary equivalence) by the intensity of $B_0(x)$ and the angle $\theta(x) \in [0, \pi/2)$ between the unit normal of the boundary and $\nabla B_0(x)$. The ground state energy of this toy operator is denoted by $\lambda_0(\mathbb{R}_+, \theta(x))$.

The leading order behavior of the ground state energy of the operator in (2.1) is now described as follows [20],

$$\lambda(H\kappa A_0) \sim (H\kappa)^{2/3} \alpha_1^{2/3}, \quad (2.15)$$

as $H\kappa \rightarrow \infty$.

Here

$$\alpha_1 = \min \left(\lambda_0^{3/2} \min_{x \in \Gamma_{\text{blk}}} |\nabla B_0(x)|, \min_{x \in \Gamma_{\text{bnd}}} \lambda_0(\mathbb{R}_+, \theta(x)) |\nabla B_0(x)| \right), \quad (2.16)$$

$$\Gamma_{\text{blk}} = \{x \in \Omega : B_0(x) = 0\} \quad (2.17)$$

and

$$\Gamma_{\text{bnd}} = \{x \in \partial\Omega : B_0(x) = 0\}. \quad (2.18)$$

The critical value $H_{c_3}(\kappa)$ could tentatively be defined as the solution of the equation in (2.2). However, when $B_0 = \text{curl } A_0$ vanishes, monotonicity of $t \mapsto \lambda(tA_0)$ is not a direct application of Chapter 3 in [4] (see the discussion below). Nevertheless, for the various definitions of $H_{c_3}(\kappa)$ proposed in [20], one always get that, for large values of κ ,

$$H_{c_3}(\kappa) \sim \frac{\kappa^2}{\alpha_1}. \quad (2.19)$$

Surface superconductivity is absent if

$$\lambda_0^{3/2} \min_{x \in \Gamma_{\text{blk}}} |\nabla B_0(x)| < \min_{x \in \Gamma_{\text{bnd}}} \lambda_0(\mathbb{R}_+, \theta(x)) |\nabla B_0(x)|,$$

and in this case, we do not distinguish between H_{c_2} and H_{c_3} . However, if

$$\lambda_0^{3/2} \min_{x \in \Gamma_{\text{blk}}} |\nabla B_0(x)| > \min_{x \in \Gamma_{\text{bnd}}} \lambda_0(\mathbb{R}_+, \theta(x)) |\nabla B_0(x)|, \quad (2.20)$$

the phenomenon of surface superconductivity is observed in decreasing magnetic fields. Superconductivity will nucleate near the minima of the function

$$\Gamma_{\text{bnd}} \ni x \mapsto \lambda_0(\mathbb{R}_+, \theta(x)) |\nabla B_0(x)|.$$

In this case, a natural definition of $H_{c_2}(\kappa)$ can be,

$$H_{c_2}(\kappa) := \frac{\kappa^2}{\alpha_2}, \quad (2.21)$$

for large values of κ . Here

$$\alpha_2 = \lambda_0^{3/2} \min_{x \in \Gamma_{\text{blk}}} |\nabla B_0(x)|.$$

Recently, we learn from N. Raymond that his student J-P. Miqueu [18] is working on the case when (2.20) is satisfied. The project in [18] is to give a complete asymptotics of the first eigenvalue under the additional assumption that Γ touches the boundary transversally. If successful, then a modification of the proof³ given in [6] will yield the monotonicity of $\lambda^N(tA_0)$ for large values of t .

Clearly, the condition in (1.9) is violated when the intensity of the magnetic field H is comparable with the critical value $H_{c_3}(\kappa) \approx \kappa^2$, thereby preventing the application of the results of Attar [1].

³Personal communication with S. Fournais.

3. THE LIMITING PROBLEM

3.1. The Montgomery operator.

Consider the self-adjoint operator in $L^2(\mathbb{R}^2)$

$$P = - \left(\partial_{x_1} - i \frac{x_2^2}{2} \right)^2 - \partial_{x_2}^2. \quad (3.1)$$

The ground state energy

$$\lambda_0 = \inf \sigma(P) \quad (3.2)$$

of the operator P is described using the Montgomery operator as follows.

If $\tau \in \mathbb{R}$, let $\lambda(\tau)$ be the first eigenvalue of the Montgomery operator [19],

$$P(\tau) = - \frac{d^2}{dx_2^2} + \left(\frac{x_2^2}{2} + \tau \right)^2, \quad \text{in } L^2(\mathbb{R}). \quad (3.3)$$

Notice that the eigenvalue $\lambda(\tau)$ is positive, simple and has a unique positive eigenfunction φ^τ of L^2 norm 1. There exists a **unique** $\tau_0 \in \mathbb{R}$ such that

$$\lambda_0 = \lambda(\tau_0). \quad (3.4)$$

Hence $\lambda_0 > 0$. We write

$$\varphi_0 = \varphi^{\tau_0}$$

Clearly, the function

$$\psi_0(x_1, x_2) = e^{-i\tau_0 x_1} \varphi_0(x_2), \quad (3.5)$$

is a bounded (generalized) eigenfunction of the operator P with eigenvalue λ_0 . Moreover (see [12] and references therein) the minimum of λ at τ_0 is non-degenerate.

We collect some important properties of the family of operators $P(\tau)$.

Theorem 3.1. ([12])

- (1) $\tau_0 < 0$.
- (2) $\lim_{\tau \rightarrow \pm\infty} \lambda(\tau) = \infty$.
- (3) The function $\lambda(\tau)$ is increasing on the interval $[0, \infty)$.

3.2. A one dimensional energy.

Let $b > 0$ and $\alpha \in \mathbb{R}$. Consider the functional

$$\mathcal{E}_{\alpha,b}^{1D}(f) = \int_{-\infty}^{\infty} \left(|f'(t)|^2 + \left(\frac{t^2}{2} + \alpha \right)^2 |f(t)|^2 - b |f(t)|^2 + \frac{b}{2} |f(t)|^4 \right) dt, \quad (3.6)$$

defined over configurations in the space

$$B^1(\mathbb{R}) = \{f \in H^1(\mathbb{R}; \mathbb{R}) : t^2 f \in L^2(\mathbb{R})\}.$$

In light of Theorem 3.1, we may define two functions $z_1(b)$ and $z_2(b)$ satisfying,

$$z_1(b) < \tau_0 < z_2(b), \quad \lambda^{-1}([\tau_0, b]) = (z_1(b), z_2(b)). \quad (3.7)$$

Notice that, if $b < \lambda(0)$, then $z_2(b) < 0$. This follows from (3) in Theorem 3.1.

Theorem 3.2. ([4, Sec. 14.2])

- (1) The functional $\mathcal{E}_{\alpha,b}^{1D}$ has a non-trivial minimizer in the space $B^1(\mathbb{R})$ if and only if $\lambda(\alpha) < b$. Furthermore, a non-trivial minimizer f_α can be found which is a positive function and $\pm f_\alpha$ are the only real-valued minimizers.

(2) Let

$$\mathfrak{b}(\alpha, b) = \inf\{\mathcal{E}_{\alpha, b}^{1D}(f) : f \in B^1(\mathbb{R})\}. \quad (3.8)$$

There exists $\alpha_0 \in (z_1(b), z_2(b))$ such that,

$$\mathfrak{b}(\alpha_0, b) = \inf_{\alpha \in \mathbb{R}} \mathfrak{b}(\alpha, b). \quad (3.9)$$

(3) If $b < \lambda(0)$, then $\alpha_0 < 0$.

(4) (Feynman-Hellmann)

$$\int_{-\infty}^{\infty} \left(\frac{t^2}{2} + \alpha_0 \right) |f_{\alpha_0}(t)|^2 dt = 0. \quad (3.10)$$

The proof of this theorem can be obtained by adapting the analysis of [4, Sec. 14.2] devoted to the functional

$$\mathcal{F}_{\alpha, b}^{1D}(f) = \int_0^{\infty} \left(|f'(t)|^2 + (t + \alpha)^2 |f(t)|^2 - b|f(t)|^2 + \frac{b}{2}|f(t)|^4 \right) dt. \quad (3.11)$$

We note for future use that a minimizer of $\mathcal{E}_{\alpha, b}^{1D}$ satisfies the Euler-Lagrange equation:

$$-f''(t) + \left(\frac{t^2}{2} + \alpha \right)^2 f(t) - bf(t) + bf(t)^3 = 0, \quad (3.12)$$

and that $f \in \mathcal{S}(\mathbb{R})$.

According to Theorem 3.2, we observe that the functional $\mathcal{E}_{\alpha, b}^{1D}$ has non-trivial minimizers if and only if $\alpha \in (z_1(b), z_2(b))$.

3.3. Reduced Ginzburg-Landau functional.

Let $L > 0$, $R > 0$, $\mathcal{S}_R = (-R, R) \times \mathbb{R}$ and

$$\mathbf{A}_{\text{app}}(x) = \left(-\frac{x_2^2}{2}, 0 \right), \quad (x = (x_1, x_2) \in \mathbb{R}^2). \quad (3.13)$$

Consider the functional

$$\mathcal{E}_{L, R}(u) = \int_{\mathcal{S}_R} \left(|(\nabla - i\mathbf{A}_{\text{app}})u|^2 - L^{-2/3}|u|^2 + \frac{L^{-2/3}}{2}|u|^4 \right) dx, \quad (3.14)$$

and the ground state energy

$$\mathfrak{e}_{\text{gs}}(L; R) = \inf\{\mathcal{E}_{L, R}(u) : u \in H_0^1(\mathcal{S}_R)\}. \quad (3.15)$$

The functional in (3.14) has a minimizer (see [10, Theorem 3.6] or [21]). Useful properties of the minimizers are collected in the next theorem.

Theorem 3.3. *Let $L > 0$, $R > 0$ and $\varphi_{L, R} \in H_0^1(\mathcal{S}_R)$ be a minimizer of the functional $\mathcal{E}_{L, R}$ in (3.14). There holds,*

$$\|\varphi_{L, R}\|_{\infty} \leq 1. \quad (3.16)$$

Furthermore, there exist universal positive constants C_1 and C_2 such that the following is true:

(1) If $L < 2^{-\frac{3}{2}}$ and $R > 0$, then

$$\int_{\mathcal{S}_R \cap \{|x_2| \geq 4L^{-2/3}\}} \frac{|x_2|^3}{(\ln |x_2|)^2} |(\nabla - i\mathbf{A}_{\text{app}})\varphi_{L, R}|^2 dx \leq C_1 L^{-8/3} |\ln L|^{-2} R, \quad (3.17)$$

$$\int_{\mathcal{S}_R \cap \{|x_2| \geq 4L^{-2/3}\}} \frac{|x_2|}{(\ln |x_2|)^2} |\varphi_{L, R}(x)|^2 \leq C_1 L^{-1/3} |\ln L|^{-\frac{3}{2}} R. \quad (3.18)$$

(2) If $L \geq 2^{-\frac{3}{2}}$ and $R > 0$, then

$$\int_{\mathcal{S}_R \cap \{|x_2| \geq 8\}} \frac{|x_2|^3}{(\ln |x_2|)^2} |(\nabla - i\mathbf{A}_{\text{app}})\varphi_{L,R}|^2 dx \leq C_2 L^{2/3} R, \quad (3.19)$$

$$\int_{\mathcal{S}_R \cap \{|x_2| \geq 8\}} \frac{|x_2|}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^2 \leq C_2 L^{2/3} R. \quad (3.20)$$

Proof.

The minimizer $\varphi_{L,R}$ satisfies the Ginzburg-Landau equation,

$$-(\nabla - i\mathbf{A}_{\text{app}})^2 \varphi_{L,R} = L^{-2/3} (1 - |\varphi_{L,R}|^2) \varphi_{L,R}. \quad (3.21)$$

Hence (3.16) results from the strong maximum principle.

Let

$$\ell = \max(2, L^{-2/3}) \quad (3.22)$$

and $\chi(x_2)$ be a nonnegative even smooth function satisfying

$$\chi(x_2) = 0 \quad \text{if } |x_2| \leq \ell, \quad \chi(x_2) = \frac{|x_2|^{3/2}}{\ln |x_2|} \quad \text{if } |x_2| \geq 4\ell.$$

Clearly, χ' satisfies with our choice of ℓ ,

$$0 < |\chi'(x_2)| \leq \frac{3\sqrt{|x_2|}}{2 \ln |x_2|} \quad \text{if } |x_2| \geq 4\ell,$$

and one can choose χ on $[\ell, 4\ell]$ such that χ' satisfies,

$$|\chi'(x_2)| \leq \widehat{C} \ell^{1/2} |\ln \ell|^{-1} \quad \text{if } |x_2| \leq 4\ell,$$

for some constant \widehat{C} independent of ℓ .

Multiplying both sides of (3.21) by $\chi^2 \varphi_{L,R}$ and integrating by parts yield,

$$\int_{\mathcal{S}_R} \left(|(\nabla - i\mathbf{A}_{\text{app}})\chi\varphi_{L,R}|^2 - L^{-2/3} |\chi\varphi_{L,R}|^2 + L^{-2/3} \chi^2 |\varphi_{L,R}|^4 \right) dx = \int_{\mathcal{S}_R} |\chi'(x_2) \varphi_{L,R}(x)|^2 dx. \quad (3.23)$$

The following inequality is standard in the spectral analysis of the Schrödinger operators with magnetic fields [4, Lemma 1.4.1],

$$\int_{\mathcal{S}_R} |(\nabla - i\mathbf{A}_{\text{app}})\chi\varphi_{L,R}|^2 dx \geq \int_{\mathcal{S}_R} |\text{curl } \mathbf{A}_{\text{app}}(x)| |\chi(x_2)\varphi_{L,R}(x)|^2 dx = \int_{\mathcal{S}_R} |x_2| |\chi\varphi_{L,R}|^2 dx. \quad (3.24)$$

Since the function $x \mapsto \chi(x_2)\varphi_{L,R}(x)$ is supported in $\mathcal{S}_R \cap \{|x_2| \geq \ell\}$ and

$$\ell = \max(2, L^{-2/3}) \geq L^{-2/3},$$

we infer from the previous inequality that,

$$\int_{\mathcal{S}_R} |(\nabla - i\mathbf{A}_{\text{app}})\chi\varphi_{L,R}|^2 dx \geq L^{-2/3} \int_{\mathcal{S}_R} |\chi(x_2)\varphi_{L,R}(x)|^2 dx.$$

This inequality allows us to deduce from (3.23) that,

$$L^{-2/3} \int_{\mathcal{S}_R} \chi(x_2)^2 |\varphi_{L,R}(x)|^4 dx \leq \int_{\mathcal{S}_R} |\chi'(x_2) \varphi_{L,R}(x)|^2 dx. \quad (3.25)$$

We obtain an upper bound of the term on the right side as follows. First, we use the Cauchy-Schwarz inequality to write,

$$\begin{aligned} & \int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^2 dx \\ & \leq \left(\int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{1}{|x_2|(\ln |x_2|)^2} dx \right)^{1/2} \left(\int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|^3}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^4 dx \right)^{1/2} \\ & \leq C |\ln \ell|^{-\frac{1}{2}} R^{1/2} \left(\int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|^3}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^4 dx \right)^{1/2}. \end{aligned} \quad (3.26)$$

Next we use the assumptions on χ , (3.16) and (3.26),

$$\begin{aligned} & \int_{\mathcal{S}_R} |\chi'(x_2) \varphi_{L,R}(x)|^2 dx \\ & \leq \int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{9|x_2|}{4(\ln |x_2|)^2} |\varphi_{L,R}(x)|^2 dx + \int_{\mathcal{S}_R \cap \{|x_2| \leq 4\ell\}} |\chi'(x_2)|^2 |\varphi_{L,R}(x)|^2 dx \\ & \leq C |\ln \ell|^{-\frac{1}{2}} R^{1/2} \left(\int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|^3}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^4 dx \right)^{1/2} + C \ell^2 (\ln \ell)^{-2} R. \end{aligned} \quad (3.27)$$

Now we use the assumption on χ , (3.25) and (3.27) to get,

$$\begin{aligned} & L^{-2/3} \int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|^3}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^4 dx \\ & \leq L^{-2/3} \int_{\mathcal{S}_R} |\chi(x_2) \varphi_{L,R}(x)|^4 dx \\ & \leq C |\ln \ell|^{-\frac{1}{2}} R^{1/2} \left(\int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|^3}{4(\ln |x_2|)^2} |\varphi_{L,R}(x)|^4 dx \right)^{1/2} + C \ell^2 (\ln \ell)^{-2} R. \end{aligned}$$

As a consequence we obtain with a new constant C the following inequality,

$$\int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|^3}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^4 dx \leq C \left(L^{\frac{2}{3}} |\ln \ell|^{-1} + \ell^2 (\ln \ell)^{-2} \right) L^{\frac{2}{3}} R. \quad (3.28)$$

Next, we insert (3.28) into (3.26) and get,

$$\int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^2 dx \leq C |\ln \ell|^{-\frac{1}{2}} \left(L^{\frac{2}{3}} |\ln \ell|^{-1} + \ell^2 (\ln \ell)^{-2} \right)^{\frac{1}{2}} L^{\frac{1}{3}} R. \quad (3.29)$$

Again, the choice of ℓ allows us to deduce (3.18) and (3.20) from (3.26).

Now, we show how to get the two inequalities in (3.17) and (3.19). A simple decomposition of the integral below and the inequality in (3.16) yield,

$$\int_{\mathcal{S}_R} |\chi(x_2) \varphi_{L,R}(x)|^2 dx \leq \int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} |\chi(x_2) \varphi_{L,R}(x)|^2 dx + C \ell^4 |\ln \ell|^{-2} R.$$

The next inequality is an easy consequence of (3.24),

$$\int_{\mathcal{S}_R} |(\nabla - i\mathbf{A}_{\text{app}})\chi \varphi_{L,R}|^2 dx \geq 4\ell \int_{\mathcal{S}_R \cap \{|x_2| \geq 4\ell\}} |\chi(x_2) \varphi_{L,R}(x)|^2 dx.$$

We insert these two inequalities into (3.23). We get,

$$\frac{1}{2} \int_{\mathcal{S}_R} |(\nabla - i\mathbf{A}_{\text{app}})\chi \varphi_{L,R}|^2 dx \leq \int_{\mathcal{S}_R} |\chi'(x_2) \varphi_{L,R}(x)|^2 dx + C L^{-2/3} \ell^4 |\ln \ell|^{-2} R. \quad (3.30)$$

Now we use a simple commutator and get the following inequality,

$$\int_{S_R} |(\nabla - i\mathbf{A}_{\text{app}})\chi\varphi_{L,R}|^2 dx \geq \frac{1}{2} \int_{S_R} |\chi(x_2)(\nabla - i\mathbf{A}_{\text{app}})\varphi_{L,R}|^2 dx - 4 \int_{S_R} |\chi'(x_2)\varphi_{L,R}(x)|^2 dx.$$

As a consequence, we infer from (3.30),

$$\frac{1}{4} \int_{S_R} |\chi(\nabla - i\mathbf{A}_{\text{app}})\varphi_{L,R}|^2 dx \leq 3 \int_{S_R} |\chi'(x_2)\varphi_{L,R}(x)|^2 dx + CL^{-2/3}\ell^4 |\ln \ell|^{-2} R. \quad (3.31)$$

The term on the right side is controlled using (3.29) and (3.28). In that way we get,

$$\frac{1}{2} \int_{S_R} |\chi(\nabla - i\mathbf{A}_{\text{app}})\varphi_{L,R}|^2 dx \leq C\sqrt{L^{4/3} + L^{2/3}\ell^2(\ln \ell)^{-2}} R + C\ell^4 |\ln \ell|^{-2} R.$$

Thanks to the choice of χ and ℓ , we deduce the inequalities in (3.17) and (3.19). \square

Remark 3.4. Let $\Lambda > 0$. The inequalities in (3.17)-(3.20) remain true under the assumption that $L \leq \Lambda$ with the constant C_1 replaced by another constant C_Λ .

As a consequence of Theorem 3.3, we can obtain a uniform estimate of the energy of a minimizer $\varphi_{L,R}$.

Proposition 3.5. *Let $\Lambda > 0$. There exists a positive constant C_Λ such that, for all $L \in (0, \Lambda)$ and $R > 0$,*

$$\int_{S_R} |\varphi_{L,R}(x)|^2 dx \leq C_\Lambda L^{-2/3} R \quad \text{and} \quad \int_{S_R} |(\nabla - i\mathbf{A}_{\text{app}})\varphi_{L,R}|^2 dx \leq C_\Lambda L^{-4/3} R.$$

Proof. Let $\ell = \max(2, L^{-2/3})$. As a consequence of the inequalities in Theorem 3.3, we have,

$$\int_{S_R \cap \{|x_2| \geq 4\ell\}} \frac{|x_2|}{(\ln |x_2|)^2} |\varphi_{L,R}(x)|^2 dx \leq C \bar{\lambda} R,$$

where

$$\bar{\lambda} = L^{-2/3}(1 + |\ln L|)^{-2} + L^{4/3}.$$

Since $\ell \geq 2$, then there exists a universal constant a such that

$$|x_2| \geq 4\ell \implies \frac{|x_2|}{(\ln |x_2|)^2} \geq a > 0.$$

Consequently, we get,

$$\int_{S_R \cap \{|x_2| \geq 4\ell\}} |\varphi_{L,R}(x)|^2 dx \leq \frac{C\bar{\lambda}}{a} R.$$

On the other hand, using (3.16), we see that,

$$\int_{S_R \cap \{|x_2| \leq 4\ell\}} |\varphi_{L,R}(x)|^2 dx \leq C\ell R.$$

A combination of both inequalities lead us to

$$\int_{S_R} |\varphi_{L,R}(x)|^2 dx \leq C\ell R + \frac{C\bar{\lambda}}{a} R.$$

We have $\ell \sim L^{-2/3}$ and $\bar{\lambda} \sim L^{-2/3} |\ln L|^{-2}$ as $L \rightarrow 0_+$. Hence, we get

$$\int_{S_R} |\varphi_{L,R}(x)|^2 dx \leq C_\Lambda L^{-2/3} R.$$

To finish the proof of the theorem, we multiply both sides of (3.21) by $\overline{\varphi_{L,R}}$ and integrate by parts. In that way we obtain

$$\int_{S_R} |(\nabla - i\mathbf{A}_{\text{app}})\varphi_{L,R}|^2 dx \leq L^{-2/3} \int_{S_R} |\varphi_{L,R}(x)|^2 dx \leq C_\Lambda L^{-4/3} R.$$

\square

The value of the ground state energy defined in (3.15) is connected to the eigenvalue λ_0 in (3.4).

Proposition 3.6. *For $R > 0$, $L > 0$, if $\mathfrak{e}_{\text{gs}}(L; R)$ denotes the ground state energy in (3.15), there holds:*

- (1) *If $L \geq \lambda_0^{-3/2}$, then $\mathfrak{e}_{\text{gs}}(L; R) = 0$.*
- (2) *There exist positive constants C_1 , C_2 and C_3 such that, if $L < \lambda_0^{-3/2}$ and $R > 0$, then*

$$-C_1 L^{-4/3} R \leq \frac{\mathfrak{e}_{\text{gs}}(L; R)}{(1 - \lambda_0 L^{2/3})} \leq -C_2 L^{-2/3} R + \frac{C_3}{R}. \quad (3.32)$$

Proof.

Suppose that $L \geq \lambda_0^{-3/2}$. Let $u \in H_0^1(\mathcal{S}_R)$. The min-max principle and the condition on L tell us that $\mathcal{E}_{L,R}(u) \geq 0$ and consequently $\mathfrak{e}_{\text{gs}}(L; R) \geq 0$. But $\mathfrak{e}_{\text{gs}}(L; R) \leq \mathcal{E}_{L,R}(0) = 0$. This proves the statement in (1).

Now, suppose that $L < \lambda_0^{-3/2}$. Let $\theta \in C_c^\infty(\mathbb{R})$ be a function satisfying,

$$\text{supp } \theta \subset (-1, 1), \quad 0 \leq \theta \leq 1, \quad \theta = 1 \text{ in } (-1/2, 1/2),$$

and let

$$\theta_R(x) := \theta(x/R).$$

Let $t > 0$ and

$$u(x_1, x_2) = t \theta_R(x_1) \psi_0(x_1, x_2),$$

where ψ_0 is the function in (3.5).

Recall that ψ_0 satisfies $-(\nabla - i\mathbf{A}_{\text{app}})^2 \psi_0 = \lambda_0 \psi_0$. An integration by parts yields,

$$\begin{aligned} \int_{\mathcal{S}_R} |(\nabla - i\mathbf{A}_{\text{app}})u|^2 dx &= t^2 \left(\left\langle \theta_R(x_1)^2 \psi_0, -(\nabla - i\mathbf{A}_{\text{app}})^2 \psi_0 \right\rangle + \int_{\mathcal{S}_R} |\phi_0(x_2) \theta'_R(x_1)|^2 dx \right) \\ &= t^2 \left(\left\langle \theta_R(x_1)^2 \psi_0, -(\nabla - i\mathbf{A}_{\text{app}})^2 \psi_0 \right\rangle + \frac{1}{R} \int \theta'(x_1)^2 dx_1 \right) \\ &= t^2 \left(\lambda_0 \int_{\mathcal{S}_R} |\theta_R(x_1) \psi_0(x)|^2 dx + \frac{C}{R} \right). \end{aligned}$$

As a consequence, we get that,

$$\begin{aligned} \mathfrak{e}_{\text{gs}}(L, R) \leq \mathcal{E}_{L,R}(u) &\leq t^2 \left(\lambda_0 - L^{-2/3} \right) \int_{\mathcal{S}_R} |\theta_R \psi_0|^2 dx + t^2 \frac{C}{R} + t^4 \frac{L^{-2/3}}{2} \int_{\mathcal{S}_R} |\theta_R(x_1) \psi_0(x)|^4 dx \\ &\leq t^2 \left(R \left(\lambda_0 - L^{-2/3} \right) + \frac{C}{R} \right) + R \nu L^{-2/3} t^4. \end{aligned}$$

Here

$$\nu = \int_{\mathbb{R}} |\varphi_0(x_2)|^4 dx_2$$

and φ_0 is the L^2 -normalized function introduced in (3.16).

Selecting t such that

$$\left(\lambda_0 - L^{-2/3} \right) + \nu L^{-2/3} t^2 = \frac{1}{2} \left(\lambda_0 - L^{-2/3} \right)$$

finishes the proof of the upper bound.

The lower bound is obtained as follows. Let $\varphi_{L,R}$ be the minimizer in Theorem 3.3. It follows from the min-max principle that,

$$\mathfrak{e}_{\text{gs}}(L; R) = \mathcal{E}_{L,R}(\varphi_{L,R}) \geq L^{-2/3} (\lambda_0 L^{2/3} - 1) \int_{\mathcal{S}_R} |\varphi_{L,R}(x)|^2 dx.$$

Under the assumption $L < \lambda_0^{-2/3}$, Proposition 3.5 tells us that

$$\int_{\mathcal{S}_R} |\varphi_{L,R}(x)|^2 dx \leq C_1 L^{-2/3} R.$$

As a consequence, we get the lower bound. \square

Remark 3.7.

In light of Propositions 3.5 and 3.6, we observe that:

- (1) If $L \geq \lambda_0^{-2/3}$, then $\varphi_{L,R} = 0$ is the minimizer of the functional in (3.14) realizing the ground state energy in (3.15).
- (2) If $L \leq \lambda_0^{-2/3}$, every minimizer $\varphi_{L,R}$ satisfies,

$$\int_{\mathcal{S}_R} |(\nabla - i\mathbf{A}_{\text{app}})\varphi_{L,R}(x)|^2 dx \leq CL^{-4/3}, \quad \int_{\mathcal{S}_R} |\varphi_{L,R}(x)|^2 dx \leq CL^{-2/3}R, \quad (3.33)$$

where C is a universal constant.

Notice that the energy $\mathcal{E}_{L,R}(u)$ in (3.14) is invariant under translation along the x_1 -axis. This allows us to follow the approach in [8, 21] and obtain that the limit of $\frac{\mathbf{e}_{\text{gs}}(L;R)}{R}$ as $R \rightarrow \infty$ exists. The precise statement is:

Theorem 3.8. *Given $L > 0$, there exists $E(L) \leq 0$ such that,*

$$\lim_{R \rightarrow \infty} \frac{\mathbf{e}_{\text{gs}}(L;R)}{2R} = E(L).$$

The function $(0, \infty) \ni L \mapsto E(L) \in (-\infty, 0]$ is continuous, monotone increasing and

$$E(L) = 0 \quad \text{if and only if } L \geq \lambda_0^{-3/2}.$$

Furthermore,

$$\forall R > 0, \forall L > 0, \quad E(L) \leq \frac{\mathbf{e}_{\text{gs}}(L;R)}{2R}, \quad (3.34)$$

and there exists a constant C such that

$$\forall R \geq 2, \forall L > 0, \quad \frac{\mathbf{e}_{\text{gs}}(L;R)}{2R} \leq E(L) + C \left(1 + L^{-2/3}\right) R^{-2/3}. \quad (3.35)$$

Proof.

There is nothing to prove when $L \geq \lambda_0^{-3/2}$, hence we assume that

$$0 < L < \lambda_0^{-3/2}.$$

Step 1. Let $n \geq 2$ be a natural number, $a \in (0, 1)$ and consider the family of strips

$$I_j = \left(-n^2 - 1 - a + (2j - 1)\left(1 + \frac{a}{2}\right), -n^2 - 1 + (2j + 1)\left(1 + \frac{a}{2}\right)\right) \times \mathbb{R}, \quad (j \in \mathbb{Z}).$$

Notice that the width of each strip in the family (I_j) is $2(1 + a)$, and if two strips in the family overlap, then the width of the overlapping region is a . Consider a partition of unity of \mathbb{R}^2 such that,

$$\sum_j |\chi_j|^2 = 1, \quad 0 \leq \chi_j \leq 1, \quad \sum_j |\nabla \chi_j|^2 \leq \frac{C}{a^2}, \quad \text{supp } \chi_j \subset I_j,$$

where C is a universal constant.

Define $\chi_{R,j}(x) = \chi_j(x/R)$. That way we obtain the new partition of unity,

$$\sum_j |\chi_{R,j}|^2 = 1, \quad 0 \leq \chi_{R,j} \leq 1, \quad \sum_j |\nabla \chi_{R,j}|^2 \leq \frac{C}{a^2 R^2}, \quad \text{supp } \chi_{R,j} \subset I_{R,j},$$

where $I_{R,j} = \{Rx : x \in I_j\}$.

Notice that $(I_{R,j})_{j \in \{1, 2, \dots, n^2\}}$ is a covering of $\mathcal{S}_{n^2 R} = (-n^2 R, n^2 R) \times \mathbb{R}$ by n^2 strips, each having side-length $2(1 + a)R$.

Let $\varphi_{L,n^2R} \in H_0^1(\mathcal{S}_{n^2R})$ be the minimizer in Theorem 3.3. There holds the decomposition,

$$\begin{aligned} \mathfrak{e}_{\text{gs}}(L; n^2R) &= \mathcal{E}_{L,n^2R}(\varphi_{L,n^2R}) \\ &\geq \sum_{j=1}^{n^2} \left(\mathcal{E}_{L,n^2R}(\chi_{R,j}\varphi_{L,n^2R}) - \left\| |\nabla\chi_{R,j}| \varphi_{L,n^2R} \right\|_{L^2(\mathcal{S}_{n^2R})}^2 \right) \\ &= \left(\sum_{j=1}^{n^2} \mathcal{E}_{L,n^2R}(\chi_{R,j}\varphi_{L,n^2R}) \right) - \int_{\mathcal{S}_{n^2R}} \left(\sum_{j=1}^{n^2} |\nabla\chi_{R,j}|^2 \right) |\varphi_{L,n^2R}|^2 dx \\ &\geq \left(\sum_{j=1}^{n^2} \mathcal{E}_{L,n^2R}(\chi_{R,j}\varphi_{L,n^2R}) \right) - \frac{Cn^2L^{-2/3}}{a^2R} \quad [\text{By Remark 3.7}]. \end{aligned}$$

The function $\chi_{R,j}\varphi_{L,n^2R}$ is supported in an infinite strip of width $2(1+a)R$. Since the energy $\mathcal{E}_{L,R}(u)$ is (magnetic) translation-invariant along the x_1 direction, we get

$$\forall j, \quad \mathcal{E}_{L,n^2R}(\chi_{R,j}\varphi_{L,n^2R}) \geq \mathfrak{e}_{\text{gs}}(L; (1+a)R)$$

and consequently,

$$\mathfrak{e}_{\text{gs}}(L; n^2R) \geq n^2 \mathfrak{e}_{\text{gs}}(L; (1+a)R) - C \frac{n^2L^{-2/3}}{a^2R}.$$

Dividing both sides of the above inequality by n^2R and using the estimate in Proposition 3.6, we get

$$\frac{\mathfrak{e}_{\text{gs}}(L; n^2R)}{n^2R} \geq \frac{\mathfrak{e}_{\text{gs}}(L; (1+a)R)}{R} - C \left(aL^{-2/3} + \frac{L^{-2/3}}{a^2R^2} \right).$$

Using the trivial inequality $(1+a) \leq (1+a)^2$, we finally obtain:

$$\frac{\mathfrak{e}_{\text{gs}}(L; n^2R)}{n^2R} \geq \frac{\mathfrak{e}_{\text{gs}}(L; (1+a)^2R)}{(1+a)^2R} - C \left(aL^{-2/3} + \frac{L^{-2/3}}{a^2R^2} \right). \quad (3.36)$$

Step 2. Let $\ell > 0$. Let us define,

$$d(\ell, L) = \frac{\mathfrak{e}_{\text{gs}}(L; \ell^2)}{2}, \quad f(\ell, L) = \frac{d(\ell, L)}{\ell^2}.$$

Clearly, the function $\ell \mapsto d(\ell, L)$ is decreasing. Thanks to Proposition 3.6, we observe that $d(\ell, L) \leq 0$ and $f(\ell, L)$ is bounded. Furthermore, (3.36) used with $R = \ell^2$ tells us that,

$$f(n\ell, L) \geq f((1+a)\ell, L) - C \left(aL^{-2/3} + \frac{1}{a^2\ell^2} \right).$$

By [10, Lemma 3.10], we get the existence of $E(L)$ such that

$$\lim_{\ell \rightarrow \infty} f(\ell, L) = E(L).$$

The simple change of variable $\ell = \sqrt{R}$ gives us,

$$\lim_{R \rightarrow \infty} \frac{\mathfrak{e}_{\text{gs}}(L; R)}{2R} = E(L).$$

Step 3. Using a comparison argument and the translation invariance of the energy $\mathcal{E}_{L,R}(u)$, we observe that,

$$\forall n \in \mathbb{N}, \quad \mathfrak{e}_{\text{gs}}(L; n^2R) \leq n^2 \mathfrak{e}_{\text{gs}}(L; R).$$

Dividing both sides of the above inequality by $2n^2R$ and taking $n \rightarrow \infty$, we get,

$$E(L) = \lim_{n \rightarrow \infty} \frac{\mathfrak{e}_{\text{gs}}(L; n^2R)}{2n^2R} \leq \frac{\mathfrak{e}_{\text{gs}}(L; R)}{2R}.$$

The matching lower bound for $E(L)$ is obtained by taking $n \rightarrow \infty$ in (3.36), selecting $a = R^{-2/3}$ and replacing R by $(1+a)^2 R$.

Step 4. Proposition 3.6 tells us that $E(L) = 0$ if and only if $L \geq \lambda_0^{-3/2}$. The continuity and monotonicity properties of $E(L)$ are easily obtained through the study of the energy $\mathcal{E}_{L,R}(u)$ as a function of L . The details can be found in [10, Thm 3.13]. \square

As in the case of a constant magnetic field, it would be desirable to establish a simpler expression of $E(L)$ when $L \in (\lambda(0)^{-\frac{3}{2}}, \lambda_0^{-\frac{3}{2}})$.

Conjecture 3.9. *Let λ be the function introduced in (5.3). If*

$$\lambda_0 < L^{-2/3} < \lambda(0), \quad (3.37)$$

then

$$E(L) = E^{1D}(L^{-2/3}).$$

Here, for $b > 0$, $E^{1D}(b) = \mathfrak{b}(\alpha_0, b)$ and $\mathfrak{b}(\alpha_0, b)$ is defined in (3.9).

Remark 3.10. In [12], the following numerical estimate is given: $\lambda_0 \approx 0.57$. Furthermore, the lower bound:

$$\lambda(0) \leq \left(\frac{3}{4}\right)^{\frac{4}{3}} < 1, \quad (3.38)$$

is proved. Finally the strict inequality $\lambda_0 < \lambda(0)$ is a consequence of the uniqueness of the point of minimum of the function $\lambda(\tau)$.

3.4. The approximate functional. Let $\nu \in [0, 2\pi)$ be a given angle. Define the magnetic potential:

$$\mathbf{A}_{\text{app},\nu}(x) = -\frac{|x|^2}{2}\mathbf{n}, \quad \mathbf{n} = (\cos \nu, \sin \nu), \quad (x \in \mathbb{R}^2). \quad (3.39)$$

Let $\kappa > 0$, $\ell \in (0, 1)$, $\mathcal{D}_\ell = D(0, \ell)$ the disc centered at 0 and of radius ℓ , and $L > 0$. Consider the functional:

$$\mathcal{G}(\psi) = \int_{\mathcal{D}_\ell} \left(|(\nabla - iL\kappa^3 \mathbf{A}_{\text{app},\nu})\psi|^2 - \frac{\kappa^2}{2}|\psi|^2 + \frac{\kappa^2}{2}|\psi|^4 \right) dx, \quad (3.40)$$

together with the ground state energy

$$E_{\text{gs},r}(\kappa, L, \nu; \ell) = \inf\{\mathcal{G}(\psi) : \psi \in H_0^1(\mathcal{D}_\ell)\}. \quad (3.41)$$

The change of variable $x \mapsto \sqrt{m}\kappa x$ yields

$$E_{\text{gs},r}(\kappa, L; \ell) = \mathfrak{e}_{\text{gs}}(\nu, L; R), \quad (3.42)$$

where $m = L^{2/3}$, $R = \sqrt{m}\kappa\ell$, $\mathcal{D}_R = D(0, R)$,

$$\mathcal{E}_{\nu,L,R}(u) = \int_{\mathcal{D}_R} \left(|(\nabla - i\mathbf{A}_{\text{app},\nu})u|^2 - L^{-2/3}|u|^2 + \frac{L^{-2/3}}{2}|u|^4 \right) dx, \quad (3.43)$$

and

$$\mathfrak{e}_{\text{gs}}(\nu, L; R) = \inf\{\mathcal{E}_{\nu,L,R}(u) : u \in H_0^1(\mathcal{D}_R)\}. \quad (3.44)$$

We now show that the ground state energy $\mathfrak{e}_{\text{gs}}(\nu, L; R)$ is independent of ν . Let u be a given function in $H_0^1(\mathcal{D}_R)$. We perform the rotation

$$(x_1, x_2) \mapsto (x_1 \cos \nu + x_2 \sin \nu, -x_1 \sin \nu + x_2 \cos \nu),$$

which transforms the function u to a new function \tilde{u} , then the gauge transformation $\tilde{u} \mapsto v = e^{ix_1^3/6}\tilde{u}$ and get

$$\mathcal{E}_{\nu,L,R}(u) = \int_{\mathcal{D}_R} \left(|(\nabla - i\mathbf{A}_{\text{app}})v|^2 - L^{-2/3}|v|^2 + \frac{L^{-2/3}}{2}|v|^4 \right) dx =: G_{L,R}(v),$$

where \mathbf{A}_{app} is introduced in (3.13).
Hence we get,

$$\mathbf{e}_{\text{gs}}(\nu, L; R) = \inf\{G_{L,R}(v) : v \in H_0^1(\mathcal{D}_R)\}. \quad (3.45)$$

This simple observation allows us to prove the following theorem:

Theorem 3.11. *For $\nu \in [0, 2\pi)$, $L > 0$ and $R > 0$, we have,*

$$\mathbf{e}_{\text{gs}}(\nu, L; R) = \mathbf{e}_{\text{gs}}(0, L; R) \geq \mathbf{e}_{\text{gs}}(L; R), \quad (3.46)$$

where $\mathbf{e}_{\text{gs}}(L; R)$ and $\mathbf{e}_{\text{gs}}(\nu, L; R)$ are the ground state energies introduced in (3.15) and (3.44).
Moreover, there exists a constant C such that, for $L > 0$, $\rho \in (0, 1)$, and

$$R \geq \max\left(2^{1/\rho}L^{1/(3\rho)}, 2^{1/(1-\rho)}L^{-1/(1-\rho)}, 2\right),$$

we have

$$\mathbf{e}_{\text{gs}}(0, L; R) \leq \mathbf{e}_{\text{gs}}\left(L; (1 - L^{2/3}R^{-2\rho})R\right) + CL^{-4/3}|\ln L|^{-1}|\ln(L^{1/3}R^{-\rho})|^2 R^\rho. \quad (3.47)$$

Proof.

The independence of ν was observed in (3.45). From now on we can take $\nu = 0$.

Lower bound. Let $u \in H_0^1(\mathcal{D}_R)$ be a minimizer of the functional $G_{L,R}$. The function u can be extended by 0 to a function in $H_0^1(\mathcal{S}_R)$. Thus,

$$\mathbf{e}_{\text{gs}}(0, L; R) = G_{L,R}(u) = \mathcal{E}_{L,R}(u) \geq \mathbf{e}_{\text{gs}}(L; R).$$

Upper bound. Let $a \in (0, \frac{1}{2})$, $\tilde{R} = (1 - a)R$ and

$$v = \varphi_{L,\tilde{R}} \in H_0^1(\mathcal{S}_{\tilde{R}})$$

a minimizer of $G_{L,\tilde{R}}$. Remember that $\varphi_{L,\tilde{R}} = 0$ when $L \geq \lambda_0^{-3/2}$ (Proposition 3.6). We impose the condition

$$\sqrt{a}R > 2L^{-2/3}. \quad (3.48)$$

Consider a test function $\chi \in C_c^\infty(\mathbb{R})$ such that,

$$\begin{cases} 0 \leq \chi \leq 1, & \text{supp}\chi \subset (-\sqrt{a(2-a)}R, \sqrt{a(2-a)}R), \\ \chi = 1 & \text{in } (-\sqrt{a(1-a)}R, \sqrt{a(1-a)}R), \\ |\chi'| \leq \frac{C}{\sqrt{a}R} & \text{and } |\chi''| \leq \frac{C}{aR^2}. \end{cases}$$

Let

$$u(x_1, x_2) = \chi(x_2)v(x_1, x_2), \quad (x_1, x_2) \in \mathbb{R}^2.$$

Clearly, $u \in H_0^1(\mathcal{D}_R)$. Thus,

$$\begin{aligned}
\mathfrak{e}_{\text{gs}}(0, L; R) &\leq G_{L,R}(u) = \mathcal{E}_{L,R}(u) \\
&= \int_{\mathcal{S}_R} \left(\chi(x_2)^2 |(\nabla - i\mathbf{A}_{\text{app}})v|^2 - \chi(x_2)\chi''(x_2)|v|^2 - L^{-2/3}|\chi(x_2)v|^2 + \frac{L^{-2/3}}{2}|\chi(x_2)v|^4 \right) dx \\
&\leq \int_{\mathcal{S}_R} \left(|(\nabla - i\mathbf{A}_{\text{app}})v|^2 - L^{-2/3}|v|^2 + \frac{L^{-2/3}}{2}|v|^4 \right) dx \\
&\quad + L^{-\frac{2}{3}} \int_{\mathcal{S}_R} (1 - \chi^2)|v|^2 dx + \frac{CL^{-1/3}|\ln L|^{-3/2}(\ln(\sqrt{a}R))^2}{a^{3/2}R^2} \\
&= \int_{\mathcal{S}_{\tilde{R}}} \left(|(\nabla - i\mathbf{A}_{\text{app}})v|^2 - L^{-2/3}|v|^2 + \frac{L^{-2/3}}{2}|v|^4 \right) dx \\
&\quad + L^{-2/3} \int_{\mathcal{S}_R} (1 - \chi^2)|v|^2 dx + \frac{CL^{-1/3}|\ln L|^{-\frac{3}{2}}(\ln(\sqrt{a}R))^2}{a^{3/2}R^2} \\
&= \mathfrak{e}_{\text{gs}}(L; \tilde{R}) + \frac{CL^{-1}|\ln L|^{-\frac{3}{2}}(\ln(\sqrt{a}R))^2}{a^{1/2}} + \frac{CL^{-1/3}|\ln L|^{-\frac{3}{2}}(\ln(\sqrt{a}R))^2}{a^{3/2}R^2}.
\end{aligned}$$

The two terms $L^{-\frac{2}{3}} \int_{\mathcal{S}_R} (1 - \chi(x_2)^2)|v(x_1, x_2)|^2 dx$ and $\int_{\mathcal{S}_R} \chi(x_2)\chi''(x_2)|v(x_1, x_2)|^2 dx$ have been controlled by using the decay of $v = \varphi_{L, \tilde{R}}$ established in Theorem 3.3 (Formula (3.18)). Here, we have used Assumption (3.48).

We select $a = L^{2/3}R^{-2\rho}$. Under the assumptions on R , L and ρ , we see that $0 < a < \frac{1}{2}$ and $\sqrt{a}R > 2L^{-2/3}$. Remembering that $\tilde{R} = R - a$, this achieves the proof of Theorem 3.11. \square

3.5. A useful function.

In this subsection, we recall the construction of a function that describes the energy of the Ginzburg-Landau model with constant magnetic field [8, 25]. Consider $b \in (0, \infty)$, $r > 0$, and $Q_r = (-r/2, r/2) \times (-r/2, r/2)$. Define the functional,

$$F_{b, Q_r}(u) = \int_{Q_r} \left(b|(\nabla - i\mathbf{A}_0)u|^2 - |u|^2 + \frac{1}{2}|u|^4 \right) dx, \quad \text{for } u \in H^1(Q_r). \quad (3.49)$$

Here, \mathbf{A}_0 is the magnetic potential,

$$\mathbf{A}_0(x) = \frac{1}{2}(-x_2, x_1), \quad (x = (x_1, x_2) \in \mathbb{R}^2). \quad (3.50)$$

Define the two ground state energies,

$$e_D(b, r) = \inf\{F_{b, Q_r}(u) : u \in H_0^1(Q_r)\}, \quad (3.51)$$

$$e_N(b, r) = \inf\{F_{b, Q_r}(u) : u \in H^1(Q_r)\}. \quad (3.52)$$

It is known [1, 8, 25] that,

$$\forall b > 0, \quad g(b) = \lim_{r \rightarrow \infty} \frac{e_D(b, r)}{|Q_r|} = \lim_{r \rightarrow \infty} \frac{e_N(b, r)}{|Q_r|}, \quad (3.53)$$

where $|Q_r|$ denotes the area of Q_r ($|Q_r| = r^2$) and g is a continuous function such that

$$g(0) = -\frac{1}{2} \text{ and } g(b) = 0 \text{ when } b \geq 1. \quad (3.54)$$

Furthermore, there exists a constant C such that, for all $r \geq 1$ and $b > 0$,

$$g(b) - C \frac{\sqrt{b}}{r} \leq \frac{e_N(b, r)}{|Q_r|} \leq \frac{e_D(b, r)}{|Q_r|} \leq g(b) + C \frac{\sqrt{b}}{r}. \quad (3.55)$$

We will use the function $g(\cdot)$ to prove the following important theorem:

Theorem 3.12. *There exist two positive constants C_1 and C_2 such that, if $L \in (0, \lambda_0^{-3/2}]$, then*

$$-\frac{C_1(1 - \lambda_0 L^{2/3})}{L^{4/3}} \leq E(L) \leq -\frac{C_2(1 - \lambda_0 L^{2/3})}{L^{4/3}}.$$

Proof. The lower bound follows immediately by sending R to ∞ in the lower bound in Proposition 3.6 (see also Theorem 3.8). The upper bound in the second item of Proposition 3.6 gives us the upper bound

$$E(L) \leq -\frac{C(1 - \lambda_0 L^{2/3})}{L^{2/3}}, \quad (3.56)$$

valid for all $L \in (0, \lambda_0^{-3/2})$. We have just to improve it as $L \rightarrow 0$.

The improved upper bound with order $L^{-4/3}$ follows from the construction of a test function as follows. Let us cover \mathbb{R}^2 by a lattice of squares $Q_{\ell, j}$, where $Q_{\ell, j} = (-\ell + a_j, \ell - a_j)$ and

$$\ell = mL^{1/3}. \quad (3.57)$$

The choice of the positive constant m will be specified later. Notice that the magnetic potential \mathbf{A}_{app} (cf (3.13)) satisfies

$$\mathbf{B}_{\text{app}} = \text{curl } \mathbf{A}_{\text{app}} = x_2,$$

and that the gradient of the magnetic field \mathbf{B}_{app} is bounded.

There exists a constant C such that, for any j , we can select a gauge ϕ_j , such that, in the square $Q_{\ell, j}$, we have,

$$\left| \mathbf{A}_{\text{app}}(x) - \left(a_{j,2} \mathbf{A}_0(x - a_j) - \nabla \phi_j \right) \right| \leq C \ell^2,$$

where $a_j = (a_{j,1}, a_{j,2})$.

Now, we define the test function as follows,

$$v(x) = \begin{cases} e^{i\phi_j(x)} u_r(\sqrt{a_{j,2}}(x - a_j)) & \text{if } a_{j,2} > 0 \text{ and} \\ & x \in Q_{\ell, j} \subset \{|x_1| < R \text{ and } \frac{\epsilon}{2} L^{-2/3} < |x_2| < \epsilon L^{-2/3}\}, \\ e^{i\phi_j(x)} \overline{u_r(\sqrt{|a_{j,2}|}(x - a_j))} & \text{if } a_{j,2} < 0 \text{ and} \\ & x \in Q_{\ell, j} \subset \{|x_1| < R \text{ and } \frac{\epsilon}{2} L^{-2/3} < |x_2| < \epsilon L^{-2/3}\}, \\ 0 & \text{otherwise,} \end{cases} \quad (3.58)$$

where the function $u_r \in H_0^1(Q_r)$ is a minimizer of the ground state energy F_{b, Q_r} introduced in (3.51) and $\epsilon \in (0, 1)$ is a positive constant (to be determined in (3.60)).

We impose the following condition on m and ϵ

$$m \sqrt{\frac{\epsilon}{2}} \geq 1. \quad (3.59)$$

We will use the notation

$$\mathcal{E}_{L, R}(v; Q_{\ell, j}) = \int_{Q_{\ell, j}} \left(|(\nabla - i\mathbf{A}_{\text{app}})v|^2 - L^{-2/3}|v|^2 + \frac{L^{-2/3}}{2}|v|^4 \right) dx.$$

Notice that, if $a_{j,2} > 0$ and $Q_{\ell,j} \subset \{|x_1| < R \text{ and } \frac{\epsilon}{2}L^{-2/3} < |x_2| < \epsilon L^{-2/3}\}$, then, for all $\eta > 0$,

$$\begin{aligned} & \mathcal{E}_{L,R}(v; Q_{\ell,j}) \\ & \leq L^{-2/3} \left(\int_{Q_{\ell,j}} \left(L^{2/3}(1+\eta)|(\nabla - ia_{j,2}\mathbf{A}_0(x-a_j))v|^2 - L^{-2/3}|v|^2 + \frac{L^{-2/3}}{2}|v|^4 \right) dx \right) \\ & \quad + C\eta^{-1}\ell^6 \\ & = \frac{L^{-2/3}}{a_{j,2}} \left(\int_{Q_{\sqrt{a_{j,2}}\ell}} \left(L^{2/3}a_{j,2}(1+\eta)|(\nabla - i\mathbf{A}_0(x))u_r(x)|^2 - L^{-2/3}|v|^2 + \frac{L^{-2/3}}{2}|v|^4 \right) dx \right) \\ & \quad + C\eta^{-1}\ell^6 \\ & \leq \frac{L^{-2/3}}{a_{j,2}} \left(g\left(L^{2/3}a_{j,2}(1+\eta)\right)|a_{j,2}|\ell^2 + C\sqrt{L^{2/3}a_{j,2}(1+\eta)}\sqrt{a_{j,2}}\ell \right) + C\eta^{-1}\ell^6. \end{aligned}$$

To write the last inequality, (3.55) is used with $b = L^{2/3}a_{j,2}(1+\eta)$ and $r = \sqrt{a_{j,2}}\ell$. (Thanks to the condition (3.59), we have $r \geq 1$).

Similarly, if $a_{j,2} < 0$ and $Q_{\ell,j} \subset \{|x_1| < R \text{ and } \frac{\epsilon}{2}L^{-1/3} < |x_2| < \epsilon L^{-2/3}\}$, then,

$$\mathcal{E}_{L,R}(v; Q_{\ell,j}) \leq \frac{L^{-2/3}}{|a_{j,2}|} \left(g\left(L^{2/3}|a_{j,2}|(1+\eta)\right)|a_{j,2}|\ell^2 + C\sqrt{L^{2/3}(1+\eta)}|a_{j,2}|\ell \right) + C\eta^{-1}\ell^6.$$

Notice the simple decomposition of the energy of v ,

$$\mathcal{E}_{L,R}(v) = \sum_{j \in \mathcal{J}} \mathcal{E}_{L,R}(v; Q_{\ell,j}),$$

where $\mathcal{J} = \{j : Q_{\ell,j} \subset \{|x_1| < R \text{ and } \frac{\epsilon}{2}L^{-2/3} < |x_2| < \epsilon L^{-2/3}\}\}$.

Let

$$n = \text{Card } \mathcal{J}.$$

The numbers L and ℓ are small enough such that,

$$\frac{\epsilon}{4}L^{-2/3}R \leq n\ell^2 \leq \frac{\epsilon}{2}L^{-2/3}R < L^{-2/3}R.$$

Now, we have the following upper bound on the energy of v ,

$$\mathcal{E}_{L,R}(v) \leq L^{-2/3} \left(\sum_j g\left(L^{2/3}|a_{j,2}|(1+\eta)\right)\ell^2 + Cn\sqrt{L^{2/3}(1+\eta)}\ell \right) + Cn\eta^{-1}\ell^6.$$

We select $\eta = \frac{1}{2}$. Having in mind (3.54) we can select ϵ sufficiently small such that

$$g(t) \leq -\frac{1}{4}, \quad \forall t \in [0, 2\epsilon]. \quad (3.60)$$

Observing

$$L^{2/3}|a_{j,2}|(1+\eta) \leq 2\epsilon,$$

we get, for $R \geq n\ell^2L^{\frac{2}{3}}$,

$$\frac{\mathbf{e}_{\text{gs}}(L; R)}{R} \leq \frac{\mathcal{E}_{L,R}(v)}{R} \leq L^{-2/3} \left(-\frac{\epsilon}{16}L^{-2/3} + 2CL^{1/3}\frac{L^{-2/3}}{\ell} \right) + 2CL^{-2/3}\ell^4.$$

Sending $R \rightarrow \infty$, we deduce that,

$$E(L) \leq -\frac{\epsilon}{32}L^{-4/3} + C\frac{L^{-1}}{\ell} + CL^{-2/3}\ell^4.$$

Having in mind (3.57), we get

$$E(L) \leq \left(-\frac{\epsilon}{32} + \frac{C}{m}\right) L^{-4/3} + Cm^4 L^{2/3}. \quad (3.61)$$

Recalling (3.59) and (3.60), we select m such that

$$-\frac{\epsilon}{32} + \frac{C}{m} < 0 \text{ and } m > \sqrt{\frac{2}{\epsilon}}.$$

In that way, (3.61) gives the claimed upper bound as $L \rightarrow 0$. \square

4. A PRIORI ESTIMATES AND GAUGE TRANSFORMATION

Let $\kappa > 0$, $H > 0$ and (ψ, \mathbf{A}) be a critical point of the functional in (1.1), i.e. (ψ, \mathbf{A}) satisfies,

$$-(\nabla - i\kappa H \mathbf{A})^2 \psi = \kappa^2(1 - |\psi|^2)\psi, \quad (4.1)$$

$$-\nabla^\perp \operatorname{curl}(\mathbf{A} - \mathbf{F}) = \frac{1}{\kappa H} (\overline{\psi} (\nabla - i\kappa H \mathbf{A})\psi) \quad \text{in } \Omega, \quad (4.2)$$

and the two boundary conditions

$$\begin{aligned} \nu \cdot (\nabla - i\kappa H \mathbf{A})\psi &= 0 \\ \operatorname{curl}(\mathbf{A} - \mathbf{F}) &= 0 \quad \text{on } \partial\Omega, \end{aligned}$$

where ν is the unit exterior normal vector of $\partial\Omega$.

We note for further use the following identity. Multiplying both the equation in (4.1) by $\overline{\psi}$ then integrating over Ω , we get,

$$\mathcal{E}_0(\psi, \mathbf{A}; \Omega) := \int_{\Omega} \left(|(\nabla - i\kappa H \mathbf{A})\psi|^2 - \kappa^2 |\psi|^2 + \frac{\kappa^2}{2} |\psi|^4 \right) dx = -\frac{\kappa^2}{2} \int_{\Omega} |\psi|^4 dx \leq 0. \quad (4.3)$$

We need the following estimates on ψ and \mathbf{A} that we take from [4]. Earlier versions of these estimates are given in [16] when the magnetic field is constant.

Proposition 4.1. *Let $\alpha \in (0, 1)$. There exists a constant $C = C(\alpha, \Omega) > 0$ such that, if $\kappa > 0$, $H > 0$ and (ψ, \mathbf{A}) a critical point of the functional in (1.1), then,*

$$\|\psi\|_{\infty} \leq 1, \quad (4.4)$$

$$\|\operatorname{curl}(\mathbf{A} - \mathbf{F})\|_2 \leq \frac{C}{H} \|\psi\|_2, \quad (4.5)$$

$$\|(\nabla - i\kappa H \mathbf{A})\psi\|_2 \leq \kappa \|\psi\|_2, \quad (4.6)$$

$$\|\mathbf{A} - \mathbf{F}\|_{C^{1,\alpha}(\overline{\Omega})} \leq C \frac{1 + \kappa H + \kappa^2}{\kappa H} \|\psi\|_{\infty} \|\psi\|_2. \quad (4.7)$$

Using the regularity of the curl-div system, we obtain the following improved estimates of $\mathbf{A} - \mathbf{F}$.

Proposition 4.2. *Let $\alpha \in (0, 1)$. There exists a constant $C = C(\alpha, \Omega) > 0$ such that, if $\kappa > 0$, $H > 0$ and (ψ, \mathbf{A}) a critical point of the functional in (1.1), then,*

$$\|\mathbf{A} - \mathbf{F}\|_{C^{0,\alpha}(\overline{\Omega})} \leq C \left(\|\operatorname{curl}(\mathbf{A} - \mathbf{F})\|_2 + \frac{1}{\kappa H} \|(\nabla - i\kappa H \mathbf{A})\psi\|_2 \|\psi\|_{\infty} \right).$$

Proof. Let $a = \mathbf{A} - \mathbf{F}$. Notice that a satisfies $\operatorname{div} a = 0$ in Ω and $\nu \cdot a = 0$ on $\partial\Omega$. Thus, there exists $C(\Omega) > 0$ such that for all a satisfying the previous condition

$$\|a\|_{H^2(\Omega)} \leq C(\Omega) \|\operatorname{curl} a\|_{H^1(\Omega)}.$$

Since (ψ, \mathbf{A}) is a critical point of the functional in (1.1), then

$$\nabla^\perp \operatorname{curl} a = \frac{1}{\kappa H} \operatorname{Im} (\overline{\psi} (\nabla - i\kappa H)\psi).$$

Consequently, we get

$$\|a\|_{H^2(\Omega)} \leq C \left(\|\operatorname{curl}(\mathbf{A} - \mathbf{F})\|_2 + \frac{1}{\kappa H} \|(\nabla - i\kappa H \mathbf{A})\psi\|_2 \|\psi\|_\infty \right).$$

This finishes the proof of the proposition in light of the continuous embedding of $H^2(\Omega)$ in $C^{0,\alpha}(\overline{\Omega})$. \square

The next proposition provides us with a useful gauge transformation.

Proposition 4.3. *Given Ω and B_0 as in the introduction, there exists a constant $C > 0$ such that the following is true.*

- (1) *Let $\ell > 0$, $a_j \in \Omega$, $D(a_j, \ell) \subset \Omega$ and $x_j \in D(a_j, \ell)$. There exists a function $\varphi_j \in C^1(\overline{D(a_j, \ell)})$ such that, for all $x \in D(a_j, \ell)$,*

$$|\mathbf{F}(x) - (B_0(x_j)\mathbf{A}_0(x - a_j) + \nabla\varphi_j)| \leq C\ell^2. \quad (4.8)$$

- (2) *Let $\ell > 0$, $a_j \in \Gamma$ and $x_j \in \overline{D(a_j, \ell)} \cap \Gamma$. There exist $\nu_j \in [0, 2\pi)$ and a function $\phi_j \in C^1(\overline{D(a_j, \ell)} \cap \Omega)$ such that, for all $x \in D(a_j, \ell) \cap \Omega$,*

$$|\mathbf{F}(x) - (|\nabla B_0(x_j)|\mathbf{A}_{\text{app}, \nu_j}(x - a_j) + \nabla\phi_j)| \leq C\ell^3. \quad (4.9)$$

Proof.

The function φ_j in (1) is constructed in [1].

We give the construction of the function ϕ_j announced in (2). The vector field \mathbf{F} and the function B_0 are defined in a neighborhood of $\overline{\Omega}$ (w.l.o.g. we can even assume that they are defined in \mathbb{R}^2). In particular, $\mathbf{F}(x)$ and $B_0(x)$ are defined in $D(a_j, \ell)$ even when $D(a_j, \ell) \not\subset \Omega$.

Select $\nu_j \in [0, 2\pi)$ such that,

$$\nabla B_0(a_j) = |\nabla B_0(a_j)|(\cos \nu_j, \sin \nu_j).$$

We apply Taylor's formula to the function B_0 near a_j . Since $a_j \in \Gamma$, we get,

$$B_0(x) = |\nabla B_0(a_j)|(\cos \nu_j, \sin \nu_j) \cdot (x - a_j) + f_j(x), \quad (4.10)$$

where

$$|f_j(x)| \leq C|x - a_j|^2 \leq C\ell^2, \quad (x \in D(a_j, \ell)).$$

Taylor's formula applied to the function $|\nabla B_0|$ near a_j yields,

$$|\nabla B_0(x_j)| = |\nabla B_0(a_j)| + e_j,$$

where

$$|e_j| \leq C|x_j - a_j| \leq C\ell.$$

In that way, (4.10) becomes,

$$B_0(x) = |\nabla B_0(x_j)|(\cos \nu_j, \sin \nu_j) \cdot (x - a_j) + g_j(x), \quad (4.11)$$

where $g_j(x) = f_j(x) + e_j(\cos \nu_j, \sin \nu_j) \cdot (x - a_j)$ and satisfies

$$|g_j(x)| \leq C\ell^2.$$

Define the vector field:

$$\mathbf{G}_j(y) = \left(\int_0^1 s g_j(sy + a_j) ds \right) (-y_2, y_1), \quad \text{for } y = (y_1, y_2).$$

Clearly, $|\mathbf{G}_j(y)| \leq C\ell^3$, when $y \in D(0, \ell)$ and $y + a_j \in \Omega$.

We perform the translation $y = x - a_j$ and define

$$\tilde{\mathbf{F}}(y) = \mathbf{F}(y + a_j), \quad \text{for } y \in D(0, \ell).$$

In that way, the formula in (4.11) reads as follows,

$$\operatorname{curl} \left(\tilde{\mathbf{F}} - |\nabla B_0(x_j)|\mathbf{A}_{\text{app}, \nu_j} \right) = \operatorname{curl} \mathbf{G}_j \quad \text{in } D(0, \ell),$$

where $\mathbf{A}_{\text{app}, \nu_j}$ is introduced in (3.39).

Consequently, we deduce the existence of a function $\tilde{\phi}_j \in C^1(\overline{D(0, \ell)})$ such that,

$$\tilde{\mathbf{F}} - |\nabla B_0(x_j)| \mathbf{A}_{\text{app}, \nu_j} = \mathbf{G}_j + \nabla \tilde{\phi}_j, \quad \text{in } D(0, \ell).$$

The function ϕ_j is defined by $\phi_j(x) = \tilde{\phi}_j(x - a_j)$, for $x \in D(a_j, \ell)$. \square

5. ENERGY UPPER BOUND

For the statement of the next proposition, we introduce the quantity $\psi(\mu_1, \mu_2, \mu_3, \rho)$ which is defined for $\mu_1 > 0$, $\mu_2 > 0$, $\mu_3 > 0$ and $\rho \in (0, 1)$ by

$$\psi(\mu_1, \mu_2, \mu_3, \rho) := \max \left(2^{1/\rho} \mu_1^{\frac{1-\rho}{3\rho}}, 2^{1/(1-\rho)} \mu_2^{-\frac{4-\rho}{3(1-\rho)}}, 2\mu_3^{-1/3} \right)$$

Proposition 5.1. *Let $\Lambda > 0$, $\eta \in (0, 1/2)$ and $b : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\lim_{\kappa \rightarrow \infty} b(\kappa) = \infty$ and $\lim_{\kappa \rightarrow \infty} \kappa^{-1/2} b(\kappa) = 0$. There exist positive constants C , κ_0 and ℓ_0 such that if $\rho \in (0, 1)$, $\ell \in (0, \ell_0)$, $\delta \in (0, 1)$, $\kappa \geq \kappa_0$,*

$$\kappa \ell \geq \psi \left(\frac{H}{\kappa^2} \sup_{x \in \Gamma} |\nabla B_0(x)|, \frac{H}{\kappa^2} \inf_{x \in \Gamma} |\nabla B_0(x)|, \frac{H}{\kappa^2} \inf_{x \in \Gamma} |\nabla B_0(x)|, \rho \right), \quad (5.1)$$

and

$$b(\kappa) \kappa^{3/2} \leq H \leq \Lambda \kappa^2,$$

then the ground state energy in (1.5) satisfies,

$$E_{\text{gs}}(\kappa, H) \leq \kappa \int_{\Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) + a + b, \quad (5.2)$$

where

- ds is the arc-length measure on Γ ,
-

$$\check{\lambda} = \left(\sup_{x \in \Gamma} |\nabla B_0(x)| \right) \frac{H}{\kappa^2}, \quad (5.3)$$

- $a = C \left(\check{\lambda}^{-5/9} \kappa^{1/3} \ell^{-2/3} + \check{\lambda}^{-4/3} |\ln \check{\lambda}|^{-1} |\ln(\check{\lambda} \kappa^{-1} \ell^{-1})|^2 \kappa^\rho \ell^{\rho-1} + (\delta \kappa^2 + \delta^{-1} \kappa^2 H^2 \ell^6) \ell \right)$
- $b = C \left(\eta + \delta + \check{\lambda}^{\frac{2(1-\rho)}{3}} \kappa^{-2\rho} \ell^{-2\rho} \right) \frac{\kappa^3}{H}$.

Proof.

Step 1. *Existence of ℓ_0 .*

Recall the assumption that Γ is the union of a finite number of simple smooth curves and $\Gamma \cap \partial\Omega$ is a finite set. Given $\eta > 0$, there exists a constant $\ell_1 \in (0, 1)$ such that, for all $a \in \Gamma$ and $\ell \in (0, \ell_1)$ with $\overline{D(a, \ell)} \subset \Omega$, then $D(a, \ell) \cap \Gamma$ is connected and

$$2\ell - \frac{\eta}{2}\ell \leq \int_{D(a, \ell) \cap \Gamma} ds(x) \leq 2\ell + \frac{\eta}{2}\ell. \quad (5.4)$$

Notice that $\int_{D(a, \ell) \cap \Gamma} ds(x)$ is the arc-length (along Γ) of $D(a, \ell) \cap \Gamma$. Thus, the choice of ℓ_1 is such that the arc-length of $D(a, \ell) \cap \Gamma$ is approximately 2ℓ , whenever $\ell \in (0, \ell_1)$.

The arc-length measure of Γ is denoted by $|\Gamma|$. By assumption, Γ consists of a finite number of simple smooth curves $(\Gamma_i)_{i=1}^k$. Let

$$\ell_0 = \min \left(\frac{\eta}{16} \min_{1 \leq i \leq k} |\Gamma_i|, \frac{\ell_1}{16} \left(1 + \frac{\eta}{4} \right)^{-1} \right).$$

If $\ell \in (0, \ell_0)$, then $\ell < \ell_1$, (5.4) is satisfied and

$$\frac{2\ell}{|\Gamma_i|} < \frac{\eta}{4}, \quad (1 + \eta)\ell < \frac{\ell_1}{4}. \quad (5.5)$$

Step 2. *A covering of Γ .*

In the sequel, we suppose that $\ell \in (0, \ell_0)$. Consider $i \in \{1, \dots, k\}$ and the curve Γ_i . Let $\mathbf{n}_i \in \mathbb{N}$ be the unique natural number satisfying

$$\frac{|\Gamma_i|}{2\ell} \left(1 + \frac{\eta}{4}\right)^{-1} - 1 < \mathbf{n}_i \leq \frac{|\Gamma_i|}{2\ell} \left(1 + \frac{\eta}{4}\right)^{-1}. \quad (5.6)$$

Select \mathbf{n}_i distinct points $(b_{j,i})_j$ on Γ_i such that,

$$\forall j, \quad \text{dist}_{\Gamma_i}(b_{j,i}, b_{j+1,i}) = \frac{|\Gamma_i|}{\mathbf{n}_i},$$

where dist_{Γ_i} is the arc-length measure on Γ_i .

Obviously, the Euclidean distance $e_j := |b_{j+1,i} - b_{j,i}|$ satisfies $e_j \leq \text{dist}_{\Gamma_i}(b_{j,i}, b_{j+1,i}) = \frac{|\Gamma_i|}{\mathbf{n}_i}$. Thanks to (5.6) and (5.5), we have,

$$e_j \leq 2\ell(1 + \eta) < \ell_1.$$

Thus, if $\overline{D(b_{j,i}, e_j)} \subset \Omega$, we can use (5.4) with $\ell = e_j$ and get,

$$2e_j \left(1 - \frac{\eta}{4}\right) \leq 2\frac{|\Gamma_i|}{\mathbf{n}_i} \leq 2e_j \left(1 + \frac{\eta}{4}\right).$$

Thanks to (5.6), this leads to

$$e_j \geq \frac{|\Gamma_i|}{\mathbf{n}_i} \left(1 - \frac{\eta}{4}\right) \geq 2\ell.$$

Now, define the index set

$$\mathcal{J}_i = \{j : \overline{D(b_{j,i}, e_j)} \subset \Omega\},$$

and $N_i = \text{Card } \mathcal{J}_i$. Notice that, if $j \in \mathcal{J}_i$, then $e_j \geq 2\ell$ and $\overline{D(b_{j,i}, \ell)} \subset \overline{D(b_{j,i}, e_j/2)} \subset \Omega$. The sets $(\overline{D(b_{j,i}, \ell)})_{j \in \mathcal{J}_i}$ are pairwise disjoint.

Since $\Gamma_i \cap \partial\Omega$ is a finite set, then there exists a constant c such that,

$$\text{If } a \in \Gamma_i \text{ and } \text{dist}(a, \partial\Omega) \geq c\ell, \text{ then } \overline{D(a, \ell)} \subset \Omega.$$

Consequently, the number N_i satisfies

$$\mathbf{n}_i - C \leq N_i \leq \mathbf{n}_i,$$

where $C > 0$ is a constant. Thus, thanks to (5.6) and (5.5),

$$|\Gamma_i| \left(1 + \frac{\eta}{4}\right)^{-1} - C\ell \leq N_i \times 2\ell \leq |\Gamma_i| \left(1 + \frac{\eta}{4}\right)^{-1}.$$

Now, collecting the points $(b_{j,i})_{j \in \mathcal{J}_i, i \in \{1, \dots, k\}}$, we get the collection of points on Γ ,

$$(a_j)_{j \in \mathcal{J}} = (b_{j,i})_{j \in \mathcal{J}_i, i \in \{1, \dots, k\}},$$

such that,

$$\forall j \in \mathcal{J}, \quad a_j \in \Gamma \quad \text{and} \quad \overline{D(a_j, \ell)} \subset \Omega,$$

$$N = \text{Card } \mathcal{J} = \sum_{i=1}^k N_i \quad \text{and} \quad |\Gamma| = \sum_{i=1}^k |\Gamma_i|,$$

$$|\Gamma| \left(1 + \frac{\eta}{4}\right)^{-1} - C\ell \leq N \times 2\ell \leq |\Gamma| \left(1 + \frac{\eta}{4}\right)^{-1}. \quad (5.7)$$

Notice that

$$\bigcup_j \left(\Gamma \cap \overline{D(a_j, \ell)}\right) \subset \Gamma.$$

and the arc-length measure

$$\left| \bigcup_j \left(\Gamma \cap \overline{D(a_j, \ell)} \right) \right| = \int_{\bigcup_j \left(\Gamma \cap \overline{D(a_j, \ell)} \right)} ds(x) = \sum_j \int_{\Gamma \cap \overline{D(a_j, \ell)}} ds(x)$$

satisfies

$$|\Gamma| - C\eta \leq \left| \bigcup_j \left(\Gamma \cap \overline{D(a_j, \ell)} \right) \right| \leq |\Gamma| + C\eta.$$

Thus, the arc-length measure of the set $\Gamma \setminus \bigcup_j \left(\Gamma \cap \overline{D(a_j, \ell)} \right)$ satisfies,

$$\left| \Gamma \setminus \bigcup_j \left(\Gamma \cap \overline{D(a_j, \ell)} \right) \right| \leq C\eta. \quad (5.8)$$

Step 3. *Construction of a test configuration.* For each j , select an arbitrary point $x_j \in \overline{D(a_j, \ell)} \cap \Gamma$ and write

$$\nabla B_0(x_j) = |\nabla B_0(x_j)| (\cos \nu_j, \sin \nu_j),$$

with $\nu_j \in [0, 2\pi)$.

Define

$$L = L_j = |\nabla B_0(x_j)| \frac{H}{\kappa^2}, \quad R = R_j = L^{1/3} \kappa \ell. \quad (5.9)$$

Thanks to the assumption in (5.1), there holds the condition,

$$R \geq \max \left(2^{1/\rho} L^{1/(3\rho)}, 2^{1/(1-\rho)} L^{-1/(1-\rho)}, 2 \right). \quad (5.10)$$

We define a function $w \in H^1(\Omega)$ as follows. Consider the set of indices $\mathcal{J} = \{j : D(a_j, \ell) \subset \Omega\}$. Let $x \in \Omega$ and $j \in \mathcal{J}$. If $x \in D(a_j, \ell)$, define,

$$w(x) = e^{i\kappa H \phi_j} u_{L, \ell, \nu_j}(x - a_j), \quad (5.11)$$

where $u_{R, L, \nu_j} \in H_0^1(D(0, \ell))$ is a minimizer of the functional in (3.40) with $\nu = \nu_j$, and ϕ_j is the function constructed in Proposition 4.3. If $x \notin \bigcup_{j \in \mathcal{J}} \overline{D(a_j, \ell)}$, we set $w(x) = 0$.

Clearly, $w \in H^1(\Omega)$.

Step 4. *Upper bound of $\mathcal{E}(w, \mathbf{F})$.*

Notice that $\text{curl } \mathbf{F} = B_0$ and that the magnetic energy term in (1.1) vanishes for $\mathbf{A} = \mathbf{F}$. Thus, we have,

$$\mathcal{E}(w, \mathbf{F}) = \mathcal{E}_0(w, \mathbf{F}; \Omega) = \sum_j \mathcal{E}_0(w, \mathbf{F}; D(a_j, \ell)), \quad (5.12)$$

where the functional \mathcal{E}_0 is defined in (1.13).

Recalling the definition of w , we observe that,

$$\mathcal{E}_0(w, \mathbf{F}; D(a_j, \ell)) = \mathcal{E}_0(u_{L, \ell, \nu_j}(x - a_j), \mathbf{F} - \nabla \phi_j; D(a_j, \ell)).$$

Thanks to the choice of ϕ_j , we infer from Proposition 4.3,

$$\left| |\nabla B_0(x_j)| \mathbf{A}_{\text{app}, \nu_j}(x - a_j) - (\mathbf{F} - \nabla \phi_j) \right| \leq C \ell^3. \quad (5.13)$$

As a consequence, applying the Cauchy-Schwarz inequality, we get that, for any $\delta > 0$,

$$\mathcal{E}_0(w, \mathbf{F}; D(a_j, \ell)) \leq (1 + \delta) \mathcal{E}_0(u_{L, \ell, \nu_j}(x - a_j), |\nabla B_0(x_j)| \mathbf{A}_{\text{app}, \nu_j}(x - a_j); D(a_j, \ell)) + r_1,$$

where

$$r_1 = C (\delta \kappa^2 + \delta^{-1} \kappa^2 H^2 \ell^6) \int_{D(a_j, \ell)} |u_{L, \ell, \nu_j}(x - a_j)|^2 dx.$$

Recall that u_{L,ℓ,ν_j} being a minimizer, it satisfies

$$|u_{L,\ell,\nu_j}| \leq 1.$$

Thus,

$$r_1 \leq C (\delta \kappa^2 + \delta^{-1} \kappa^2 H^2 \ell^6) \ell^2. \quad (5.14)$$

Now, performing the translation $x \mapsto x - a_j$, we observe that,

$$\mathcal{E}_0(w, \mathbf{F}; D(a_j, \ell)) \leq (1 + \delta) \mathcal{E}_0(u_{L,\ell,\nu_j}, |\nabla B_0(x_j)| \mathbf{A}_{\text{app},\nu_j}; D(0, \ell)) + r_1.$$

With $L = L_j$ and $R = R_j$ in (5.9), we get in light of Theorem 3.11:

$$\mathcal{E}_0(w, \mathbf{A}; D(a_j, \ell)) \leq (1 + \delta) \mathbf{e}_{\text{gs}} \left(L_j; (1 - L_j^{2/3} R_j^{-2\rho}) R_j \right) + C L_j^{-4/3} |\ln L_j|^{-1} |\ln(L_j^{1/3} R_j^{-\rho})|^2 R_j^\rho + r_1.$$

Thanks to Theorem 3.8, we deduce that,

$$\begin{aligned} \mathcal{E}_0(w, \mathbf{F}; D(a_j, \ell)) &\leq 2(1 + \delta) (1 - L_j^{2/3} R_j^{-2\rho}) R_j E(L_j) + C (1 + L_j^{-2/3}) R_j^{1/3} \\ &\quad + C L_j^{-4/3} |\ln L_j|^{-1} |\ln(L_j^{1/3} R_j^{-\rho})|^2 R_j^\rho + r_1. \end{aligned} \quad (5.15)$$

Recall the definition of L_j and R_j in (5.9), and that the number of disks $D(a_j, \ell)$ is inversely proportional to ℓ , i.e. of order ℓ^{-1} .

In the sequel, the following remark will be used. The two terms

$$\check{\lambda} = \left(\sup_{x \in \Gamma} |\nabla B_0(x)| \right) \frac{H}{\kappa^2} \quad \text{and} \quad \underline{\lambda} = \left(\inf_{x \in \Gamma} |\nabla B_0(x)| \right) \frac{H}{\kappa^2}$$

are of the same order, i.e. there exist constants $c_1 > 0$ and $c_2 > 0$ such that $c_1 \check{\lambda} \leq \underline{\lambda} \leq c_2 \check{\lambda}$. Thus, terms controlled by $\underline{\lambda}$ are controlled by $\check{\lambda}$ and vice versa. We will express all terms controlled by $\underline{\lambda}$ and $\check{\lambda}$ in the form $\mathcal{O}(\check{\lambda})$.

Substituting (5.15) into (5.12) yields,

$$\begin{aligned} \mathcal{E}(w, \mathbf{F}) &\leq 2\kappa\ell(1 + \delta) (1 - \check{\lambda}^{\frac{2(1-\rho)}{3}} \kappa^{-2\rho} \ell^{-2\rho}) \left(\sum_j \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \right) \\ &\quad + C \check{\lambda}^{-5/9} \kappa^{1/3} \ell^{-2/3} + C \check{\lambda}^{-4/3} |\ln \check{\lambda}|^{-1} |\ln(\check{\lambda} \kappa^{-1} \ell^{-1})|^2 \kappa^\rho \ell^{\rho-1} + C r_1 \ell^{-1}. \end{aligned} \quad (5.16)$$

Thanks to (5.7) and the upper bound on $E(\cdot)$ obtained in Theorem 3.12, the term

$$\sum_j 2\ell \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \quad (5.17)$$

is of order κ^2/H . Thus, (5.16) becomes,

$$\begin{aligned} \mathcal{E}(w, \mathbf{F}) &\leq \kappa \left(\sum_j 2\ell \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \right) + C \left(\delta + \check{\lambda}^{\frac{2(1-\rho)}{3}} \kappa^{-2\rho} \ell^{-2\rho} \right) \frac{\kappa^3}{H} \\ &\quad + C \check{\lambda}^{-5/9} \kappa^{1/3} \ell^{-2/3} + C \check{\lambda}^{-4/3} |\ln \check{\lambda}|^{-1} |\ln(\check{\lambda} \kappa^{-1} \ell^{-1})|^2 \kappa^\rho \ell^{\rho-1} + C r_1 \ell^{-1}. \end{aligned} \quad (5.18)$$

In (5.17), replacing 2ℓ by the arc-length measure of $D(a_j, \ell) \cap \Gamma$ produces an error $\eta\ell/2$ and the sum becomes a Riemann sum over $V_\ell = \bigcup_{j \in \mathcal{J}} (\Gamma \cap \overline{D(a_j, \ell)})$.

The points x_j can be selected such that the Riemann sum is a *lower* Riemann sum. Thus,

$$\begin{aligned} \sum_j 2\ell \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \\ \leq \int_{V_\ell} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) + C\eta \frac{\kappa^2}{H}. \end{aligned}$$

Inserting this into (5.18), we get,

$$\begin{aligned} \mathcal{E}(w, \mathbf{F}) \leq & \kappa \left(\int_{V_\ell} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) + C \left(\eta + \delta + \check{\lambda}^{\frac{2(1-\rho)}{3}} \kappa^{-2\rho} \ell^{-2\rho} \right) \frac{\kappa^3}{H} \\ & + C \check{\lambda}^{-5/9} \kappa^{1/3} \ell^{-2/3} + C \check{\lambda}^{-4/3} |\ln \check{\lambda}|^{-1} |\ln(\check{\lambda} \kappa^{-1} \ell^{-1})|^2 \kappa^\rho \ell^{\rho-1} + C r_1 \ell^{-1}. \end{aligned}$$

As pointed earlier, the arc-length measure of the set $\Gamma \setminus V_\ell$ does not exceed $C\eta$. Recall the upper bound on $E(\cdot)$ obtained in Theorem 3.12. In that way, we get,

$$\left| \int_{\Gamma \setminus V_\ell} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right| \leq C \eta \frac{\kappa^2}{H}.$$

Consequently, we deduce the following upper bound,

$$\begin{aligned} \mathcal{E}(w, \mathbf{F}) \leq & \kappa \left(\int_{\Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) + C \left(\eta + \delta + \check{\lambda}^{\frac{2(1-\rho)}{3}} \kappa^{-2\rho} \ell^{-2\rho} \right) \frac{\kappa^3}{H} \\ & + C \check{\lambda}^{-5/9} \kappa^{1/3} \ell^{-2/3} + C \check{\lambda}^{-4/3} |\ln \check{\lambda}|^{-1} |\ln(\check{\lambda} \kappa^{-1} \ell^{-1})|^2 \kappa^\rho \ell^{\rho-1} + C r_1 \ell^{-1} + C \eta \frac{\kappa^3}{H}. \end{aligned}$$

The definition of the ground state energy in (1.5) tells us that $E_{\text{gs}}(\kappa, H) \leq \mathcal{E}(w, \mathbf{F})$. Recalling the definition of r_1 in (5.14) finishes the proof of (5.2). \square

As a straightforward application of Proposition 5.1, we deduce the following upper bound on the ground state energy in (1.5), valid in the regime $\kappa^{3/2} \ll H \lesssim \kappa^2$.

Theorem 5.2. *Let $\Lambda > 0$ and $\epsilon : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\lim_{\kappa \rightarrow \infty} \kappa \epsilon(\kappa) = \infty$ and $\lim_{\kappa \rightarrow \infty} \epsilon(\kappa) = 0$. If $\epsilon(\kappa) \kappa^2 \leq H \leq \Lambda \kappa^2$, then the ground state energy in (1.5) satisfies,*

$$E_{\text{gs}}(\kappa, H) \leq \kappa \left(\int_{\Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) + \frac{\kappa^3}{H} o(1), \quad (\kappa \rightarrow \infty), \quad (5.19)$$

where ds is the arc-length measure on Γ .

Proof. We use the upper bound in Proposition 5.1 with the following choice of the parameters:

$$\begin{cases} 0 < \rho < \frac{1}{8}, & \ell = \epsilon_1^{1/8} \kappa^{-3/4} \check{\lambda}^{-1/2}, & \delta = \epsilon_1^{7/8} |\ln \kappa| \kappa^{-1/4} \check{\lambda}^{-1/2}, \\ \epsilon_1 = \kappa^{-2/7} |\ln \kappa|^{-8/7}. \end{cases} \quad (5.20)$$

Clearly, ϵ_1 satisfies,

$$\epsilon_1 \ll 1 \quad \text{as } \kappa \rightarrow \infty.$$

Recall that $\check{\lambda}$ (introduced in (5.3)) satisfies

$$\kappa^{-1/2} \ll \check{\lambda} \lesssim 1.$$

In that way, the parameters δ and ℓ satisfy,

$$\begin{aligned} \delta & \ll \epsilon_1^{7/8} |\ln \kappa| \kappa^{1/4} = 1, \\ \ell & \ll \epsilon_1^{1/8} \kappa^{-3/4} \kappa^{1/2} = \epsilon_1^{1/8} \kappa^{-1/4} \ll 1. \end{aligned}$$

Let us show that the three conditions in (5.1) are satisfied. We rewrite

$$\kappa \ell = \epsilon_1^{1/8} \kappa^{1/4} \check{\lambda}^{-1/2} = \kappa^{3/14} |\ln \kappa|^{-1/7} \check{\lambda}^{-1/2} = \kappa^{\varsigma/14} |\ln \kappa|^{-1/7} \kappa^{(3-\varsigma)/14} \check{\lambda}^{-1/2},$$

with $\varsigma := \frac{1-8\rho}{3(1-\rho)}$.

Thanks to the condition $0 < \rho < \frac{1}{8}$, we have $0 < \varsigma < 1$.

The conditions in (5.1) will follow from

$$\kappa \ell \gg \max \left\{ \check{\lambda}^{\frac{1-\rho}{2\rho}}, \check{\lambda}^{-\frac{4-\rho}{3(1-\rho)}}, \check{\lambda}^{-\frac{1}{3}} \right\}.$$

We first observe

$$\kappa\ell \gg \check{\lambda}^{-(3-\varsigma)/7} \check{\lambda}^{-1/2} = \check{\lambda}^{-\frac{37-23\rho}{14 \times 3(1-\rho)}} \gtrsim 1.$$

Then, we deduce,

$$\begin{aligned} \kappa\ell \check{\lambda}^{\frac{4-\rho}{3(1-\rho)}} &\gg \check{\lambda}^{\frac{-29+21\rho}{14 \times 3(1-\rho)}} \gtrsim 1, \\ \kappa\ell \check{\lambda}^{1/3} &\gg \check{\lambda}^{\frac{-23+9\rho}{14 \times 3(1-\rho)}} \gtrsim 1, \end{aligned}$$

and we can apply Proposition 5.1.

Now, the remainder a in Proposition 5.1 satisfies⁴

$$a \ll \kappa \check{\lambda}^{-1}.$$

Notice indeed that,

$$\begin{aligned} \delta^{-1} \kappa^2 H^2 \ell^7 &\approx |\ln \kappa|^{-1} \kappa \check{\lambda}^{-1} \ll \kappa \check{\lambda}^{-1}, \\ \delta \kappa^2 \ell &= \epsilon_1 |\ln \kappa| \kappa \check{\lambda}^{-1} \ll \kappa \check{\lambda}^{-1}, \\ |\ln(\check{\lambda} \kappa \ell)|^2 \kappa^\rho \ell^{\rho-1} &= |\ln(\check{\lambda} \kappa \ell)|^2 (\epsilon_1^{\frac{1}{8}} \kappa^{1/4})^{\rho-1} \check{\lambda}^{\frac{1-3\rho}{2}} \kappa \check{\lambda}^{-1} \lesssim |\ln(\check{\lambda} \kappa \ell)|^2 (\epsilon_1^{\frac{1}{8}} \kappa^{1/4})^{\rho-1} \kappa \check{\lambda}^{-1} \ll \kappa \check{\lambda}^{-1}, \\ \check{\lambda}^{-5/9} \kappa^{1/3} \ell^{-2/3} &= \check{\lambda}^{7/9} \kappa^{-\frac{5}{6 \times 7}} |\ln \kappa|^{\frac{2}{3 \times 7}} \kappa \check{\lambda}^{-1} \ll \kappa \check{\lambda}^{-1}. \end{aligned}$$

The last point to verify is that

$$\check{\lambda}^{\frac{2(1-\rho)}{3}} \kappa^{-2\rho} \ell^{-2\rho} \ll 1,$$

which follows easily since we know that

$$1 - \rho > 0, \check{\lambda} \lesssim 1 \text{ and } \kappa \ell \gg 1.$$

Now sending $\kappa \rightarrow \infty$, the upper bound in Proposition 5.1 becomes,

$$\limsup_{\kappa \rightarrow \infty} \frac{H}{\kappa^3} \left\{ E_{\text{gs}}(\kappa, H) - 2\kappa \left(\int_{\Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) \right\} \leq C\eta.$$

Since this is true for all $\eta \in (0, 1/2)$, we get by sending $\eta \rightarrow 0_+$,

$$\limsup_{\kappa \rightarrow \infty} \frac{H}{\kappa^3} \left\{ E_{\text{gs}}(\kappa, H) - 2\kappa \left(\int_{\Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) \right\} \leq 0,$$

and the conclusion in Theorem 5.2 follows. \square

In the next theorem, we derive an upper bound of the ground state energy in (1.5) valid in the regime $\kappa^{-1} \ll H \lesssim \kappa^{3/2}$.

Theorem 5.3. *Let $\Lambda > 0$ and $\epsilon : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a function satisfying $\lim_{\kappa \rightarrow \infty} \epsilon(\kappa) = 0$ and $\lim_{\kappa \rightarrow \infty} \kappa \epsilon(\kappa) = \infty$.*

If $\epsilon(\kappa)\kappa \leq H \leq \Lambda\kappa^{3/2}$, then the ground state energy in (1.5) satisfies,

$$E_{\text{gs}}(\kappa, H) \leq \kappa^2 \int_{\Omega} g \left(\frac{H}{\kappa} |B_0(x)| \right) dx + \frac{\kappa^3}{H} o(1), \quad (5.21)$$

where $g(\cdot)$ is the function introduced in (3.53).

Proof. Here, we construct a test function as in (5.11) below. Let $\zeta = \kappa^{-1/16} H^{-1/2}$ and $(\mathcal{Q}_{\kappa, \zeta})$ be the lattice of squares generated by the square

$$\mathcal{Q}_{\zeta} = \left(-\frac{\zeta}{2}, \frac{\zeta}{2} \right) \times \left(-\frac{\zeta}{2}, \frac{\zeta}{2} \right).$$

⁴We recall that $\kappa \check{\lambda}^{-1}$ is of the order of κ^3/H .

Notice that ζ satisfies $\zeta \ll \frac{\kappa}{H} \ll 1$. Define

$$\mathcal{I} = \left\{ k : \mathcal{Q}_{k,\zeta} \subset \{ \text{dist}(x, \Gamma) \leq M \frac{\kappa}{H} \} \setminus \left(\bigcup_{j \in \mathcal{J}} D(a_j, \ell) \right) \text{ and } \text{dist}(\mathcal{Q}_{k,\zeta}, \Gamma) \geq M\zeta \right\},$$

where $M > 0$ is a constant selected sufficiently large so that, if $\text{dist}(x, \Gamma) > M \frac{\kappa}{H}$, then $|B_0(x)| \geq \frac{\kappa}{H}$. Notice that, since B_0 vanishes non-degenerately on Γ , if $k \in \mathcal{I}$, then

$$|B_0(x)| \geq M'\zeta > 0 \text{ in } \overline{\mathcal{Q}_{k,\zeta}}.$$

For all $k \in \mathcal{I}$, let a_k be the center of the square $\mathcal{Q}_{k,\zeta}$ and select an arbitrary point $x_k \in \overline{\mathcal{Q}_{k,\zeta}}$.

If $r > 0$ and $b > 0$, let $u_r \in H_0^1(\mathcal{Q}_r)$ be the minimizer of the ground state energy F_{b,\mathcal{Q}_r} introduced in (3.51). For all $k \in \mathcal{I}$, let $r_k = \zeta \sqrt{\kappa H |B_0(x_k)|}$, $b_k = \frac{H}{\kappa} |B_0(x_k)|$, $u_k = u_{r_k}$ and φ_k be the gauge function satisfying (see Proposition 4.3),

$$|\mathbf{F}(x) - (B_0(x_k) \mathbf{A}_0(x - a_k) + \nabla \varphi_k)| \leq C\zeta^2, \quad \text{in } \overline{\mathcal{Q}_{k,\zeta}}.$$

Define the test function v as follows,

$$v(x) = \begin{cases} e^{-i\phi_k(x)} u_k\left(\frac{r_k}{\zeta}(x - a_k)\right) & \text{if } x \in \mathcal{Q}_{k,\zeta} \subset \{B_0(x) > 0\}, \\ e^{-i\phi_j(x)} \overline{u_k}\left(\frac{r_k}{\zeta}(x - a_k)\right) & \text{if } x \in \mathcal{Q}_{k,\zeta} \subset \{B_0(x) < 0\}, \\ 0 & \text{otherwise.} \end{cases} \quad (5.22)$$

We outline the computation of $\mathcal{E}_0(v, \mathbf{F})$. The details of the computations are given in [1]. In every square $\mathcal{Q}_{k,\zeta}$, we have,

$$\mathcal{E}_0(v, \mathbf{F}; \mathcal{Q}_{k,\zeta}) \leq (1 + \kappa^{-1/16}) \frac{F_{b_k, \mathcal{Q}_{r_k}}(u_k)}{b_k} + \kappa^{1/16} \kappa^2 H^2 \zeta^6.$$

Thanks to the assumption on H and the definition of $\zeta = \kappa^{-1/16} H^{-1/2}$, we have $r_k \gg 1$. Thus, we may use (3.55) and write,

$$\begin{aligned} \mathcal{E}_0(v, \mathbf{F}; \mathcal{Q}_{k,\zeta}) &\leq (1 + \kappa^{-1/16}) \frac{r_k^2}{b_k} \left(g(b_k) + C \frac{\sqrt{b_k}}{r_k} \right) + \kappa^{1/16} \kappa^2 H^2 \zeta^6 \\ &= (1 + \kappa^{-1/16}) \zeta^2 \kappa^2 \left(g\left(\frac{H}{\kappa} |B_0(x_k)|\right) + C \frac{1}{\zeta \kappa} \right) + \kappa^{1/16} \kappa^2 H^2 \zeta^6. \end{aligned}$$

We sum over k and select the points x_k as follows

$$|B_0(x_k)| = \min\{|B_0(x)| : x \in \overline{\mathcal{Q}_{k,\zeta}}\}.$$

In that way, we obtain,

$$\begin{aligned} \mathcal{E}_0(v, \mathbf{F}) &= \sum_{k \in \mathcal{I}} \mathcal{E}_0(v, \mathbf{F}; \mathcal{Q}_{k,\zeta}) \\ &\leq (1 + \kappa^{-1/16}) \kappa^2 \int_{\bigcup_{k \in \mathcal{I}} \overline{\mathcal{Q}_{k,\zeta}}} \left(g\left(\frac{H}{\kappa} |B_0(x)|\right) + C \frac{1}{\zeta \kappa} + \kappa^{1/16} H^2 \zeta^4 \right) dx \\ &\leq (1 + \kappa^{-1/16}) \kappa^2 \int_{\bigcup_{k \in \mathcal{I}} \overline{\mathcal{Q}_{k,\zeta}}} g\left(\frac{H}{\kappa} |B_0(x)|\right) dx + C \kappa^{1/16} \frac{\kappa^2}{H\zeta} + M \kappa^{-1/16} \frac{\kappa^3}{H}. \end{aligned}$$

Notice that, since $g(b) = 0$ for $b \geq 1$ and B_0 vanishes non-degenerately on Γ , then

$$\int_{\bigcup_{k \in \mathcal{I}} \overline{\mathcal{Q}_{k,\zeta}}} g\left(\frac{H}{\kappa} |B_0(x)|\right) dx \leq \int_{\Omega} g\left(\frac{H}{\kappa} |B_0(x)|\right) dx + \frac{\kappa}{H} o(1).$$

Thus,

$$\mathcal{E}_0(v, \mathbf{F}) \leq (1 + \kappa^{-1/16}) \kappa^2 \int_{\Omega} g\left(\frac{H}{\kappa} |B_0(x)|\right) dx + C \kappa^{1/16} \frac{\kappa^2}{H\zeta} + M \kappa^{-1/16} \frac{\kappa^3}{H} + \frac{\kappa^3}{H} o(1).$$

Since $H \lesssim \kappa^{3/2}$, then

$$\kappa^{1/16} \frac{\kappa^2}{H\zeta} = \kappa^{1/8} \frac{H^{1/2} \kappa^3}{\kappa H} \ll 1$$

and

$$\mathcal{E}_0(v, \mathbf{F}) \leq (1 + \kappa^{-1/16}) \kappa^2 \int_{\Omega} g \left(\frac{H}{\kappa} |B_0(x)| \right) dx + \frac{\kappa^3}{H} o(1).$$

Since $E_{\text{gs}}(\kappa, H) \leq \mathcal{E}(v, \mathbf{F}) = \mathcal{E}_0(v, \mathbf{F})$, then we get the upper bound in (5.21). \square

6. EXPONENTIAL DECAY OF THE ORDER PARAMETER

The aim of this section is to prove that the order parameter ψ is exponentially small (in the L^2 -norm) away from the points where the magnetic field vanishes. This bound is needed in Section 7 to obtain a lower bound of the ground state energy in (1.5).

6.1. A rough bound. In this subsection we give a rough bound valid for any order parameter ψ .

Theorem 6.1. *Let $\Lambda > 0$ and $\epsilon : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\lim_{\kappa \rightarrow \infty} \kappa \epsilon(\kappa) = \infty$ and $\lim_{\kappa \rightarrow \infty} \epsilon(\kappa) = 0$. There exist constants C and κ_0 such that, if (ψ, \mathbf{A}) is a critical point of the functional in (1.1), $\kappa \geq \kappa_0$ and*

$$\epsilon(\kappa) \kappa^2 \leq H \leq \Lambda \kappa^2, \quad (6.1)$$

then

$$\|\psi\|_2 \leq C \left(\frac{\kappa}{H} \right)^{1/6}, \quad (6.2)$$

$$\|\text{curl}(\mathbf{A} - \mathbf{F})\|_2 \leq \frac{C}{H} \left(\frac{\kappa}{H} \right)^{1/6}, \quad (6.3)$$

and

$$\|(\nabla - i\kappa H \mathbf{A})\psi\|_2 \leq C \kappa \left(\frac{\kappa}{H} \right)^{1/6}. \quad (6.4)$$

An important ingredient in the proof of this theorem is:

Proposition 6.2. *Let $\Lambda > 0$ and $\epsilon : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\lim_{\kappa \rightarrow \infty} \kappa \epsilon(\kappa) = \infty$ and $\lim_{\kappa \rightarrow \infty} \epsilon(\kappa) = 0$. There exist positive constants C , ℓ_0 and κ_0 such that the following is true: For $\ell \in (0, \ell_0)$, $a \in (0, 1]$ and $h \in C_c^\infty(\Omega)$ such that*

$$\text{supp } h \subset \{x \in \Omega : \text{dist}(x, \partial\Omega) > \ell \ \& \ \text{dist}(x, \Gamma) > \sqrt{a} \ell\} \text{ and } \|h\|_\infty \leq 1,$$

if (ψ, \mathbf{A}) is a critical point of the functional in (1.1), $\kappa \geq \kappa_0$ and $\epsilon(\kappa) \kappa^2 \leq H \leq \Lambda \kappa^2$, then

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})\psi|^2 dx \geq \frac{1}{C} \kappa (H \sqrt{a} \ell - C^2) \int_{\Omega} |h\psi|^2 dx - C \kappa \int_{\Omega} (1 - h^2) |\psi|^2 dx. \quad (6.5)$$

Proof.

The support of the function $h\psi$ does not meet the boundary of Ω and Γ . We can use the celebrated inequality

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})h\psi|^2 dx \geq \kappa H \int_{\Omega} |\text{curl } \mathbf{A}| |h\psi|^2 dx.$$

The simple decomposition $\text{curl } \mathbf{A} = \text{curl } \mathbf{F} + \text{curl}(\mathbf{A} - \mathbf{F})$ and the triangle inequality yield,

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})h\psi|^2 dx \geq \kappa H \int_{\Omega} |\text{curl } \mathbf{F}| |h\psi|^2 dx - \kappa H \int_{\Omega} |\text{curl } \mathbf{A} - \text{curl } \mathbf{F}| |h\psi|^2 dx. \quad (6.6)$$

By assumption ∇B_0 does not vanish on Γ , hence

$$|\text{curl } \mathbf{F}| = |B_0(x)| \geq \frac{1}{M} \sqrt{a} \ell \quad \text{in } \{\text{dist}(x, \Gamma) \geq \sqrt{a} \ell\}, \quad (6.7)$$

for some constant $M > 0$.

Thus,

$$\int_{\Omega} |\operatorname{curl} \mathbf{F}| |h\psi|^2 dx \geq \frac{1}{M} \sqrt{a} \ell \int_{\Omega} |h\psi|^2 dx. \quad (6.8)$$

Next we use the Cauchy-Schwarz inequality and the inequality in (4.5) as follows

$$\begin{aligned} \kappa H \int_{\Omega} |\operatorname{curl} \mathbf{A} - \operatorname{curl} \mathbf{F}| |h\psi|^2 dx &\leq \kappa H \|\operatorname{curl} \mathbf{A} - \operatorname{curl} \mathbf{F}\|_2 \left(\int_{\Omega} |h\psi|^4 dx \right)^{1/2} \\ &\leq C \kappa \|\psi\|_2 \left(\int_{\Omega} |h\psi|^4 dx \right)^{1/2}. \end{aligned}$$

Since $\|\psi\|_{\infty} \leq 1$ and $\|h\|_{\infty} \leq 1$, we get further,

$$\|\psi\|_2 \left(\int_{\Omega} |h\psi|^4 dx \right)^{1/2} \leq \int_{\Omega} |\psi|^2 dx = \int_{\Omega} |h\psi|^2 dx + \int_{\Omega} (1-h^2)|\psi|^2 dx.$$

Therefore, we have,

$$\kappa H \int_{\Omega} |\operatorname{curl} \mathbf{A} - \operatorname{curl} \mathbf{F}| |h\psi|^2 dx \leq C \kappa \int_{\Omega} |\psi|^2 dx + C \kappa \int_{\Omega} (1-h^2)|\psi|^2 dx. \quad (6.9)$$

Inserting (6.9) and (6.8) into (6.6) finishes the proof of the proposition. \square

Proof of Theorem 6.1. Let $\ell > 0$ and $\Omega_{\ell} = \{x \in \Omega : \operatorname{dist}(x, \partial\Omega) > \ell \ \& \ \operatorname{dist}(x, \Gamma) > \ell\}$. Select a function $h \in C_c^{\infty}(\Omega)$ satisfying

$$0 \leq h \leq 1 \text{ in } \Omega, \quad h = 1 \text{ in } \Omega_{2\ell}, \quad h = 0 \text{ in } \Omega_{\ell},$$

and

$$|\nabla h| \leq \frac{C}{\ell} \text{ in } \Omega,$$

where C is a constant.

Thanks to the bound $\|\psi\|_{\infty} \leq 1$ and the assumptions on h , we have,

$$\int_{\Omega} |\psi|^2 \leq \int_{\Omega} |h\psi|^2 + C\ell, \quad (6.10)$$

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})\psi|^2 dx \geq \int_{\Omega} |h(\nabla - i\kappa H \mathbf{A})\psi|^2 dx \quad (6.11)$$

$$\geq \frac{1}{2} \int_{\Omega} |(\nabla - i\kappa H \mathbf{A})h\psi|^2 dx - C \int_{\Omega} |\nabla h|^2 |\psi|^2 dx. \quad (6.12)$$

Thanks to the estimate on the gradient of h , we may write,

$$\frac{1}{2} \int_{\Omega} |(\nabla - i\kappa H \mathbf{A})h\psi|^2 dx - \kappa^2 \int_{\Omega} |h\psi|^2 dx - C(\ell + \ell^{-1})\kappa^2 \leq \mathcal{E}_0(\psi, \mathbf{A}; \Omega) \leq 0,$$

where $\mathcal{E}_0(\psi, \mathbf{A}; \Omega)$ is introduced in (4.3).

Now, we use Proposition 6.2 with $a = 1$ and get,

$$\left(\frac{\kappa}{2C} (H\ell - C^2) - \kappa^2 \right) \int_{\Omega} |h\psi|^2 dx \leq C(\ell + \ell^{-1})\kappa^2.$$

Selecting $\ell = (\kappa/H)^{1/3}$, we get for κ large and H satisfying (6.1)

$$\int_{\Omega} |h\psi|^2 dx \leq C \left(\frac{\kappa}{H} \right)^{1/3}.$$

Now, thanks to (6.10), the first inequality (6.2) in Theorem 6.1 is proved. Now, the inequality (6.3) (resp. (6.4)) is simply a consequence of (4.5) (resp. (4.6)). \square

6.2. Exponential bound. In the next theorem, we establish that any order parameter decays exponentially fast away from the set Γ where the magnetic field vanishes.

Theorem 6.3. *Let $\Lambda > 0$ and $\epsilon : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\lim_{\kappa \rightarrow \infty} \kappa \epsilon(\kappa) = \infty$ and $\lim_{\kappa \rightarrow \infty} \epsilon(\kappa) = 0$. There exist positive constants C , m_0 and κ_0 such that, if (ψ, \mathbf{A}) is a critical point of the functional in (1.1), $\kappa \geq \kappa_0$, $\epsilon(\kappa)\kappa^2 \leq H \leq \Lambda\kappa^2$, then*

$$\int_{\Omega} \exp\left(2m_0 \frac{H}{\kappa} t(x)\right) \left(\frac{1}{\kappa^2} |(\nabla - i\kappa H \mathbf{A})\psi|^2 + |\psi(x)|^2\right) dx \leq C \int_{\{t(x) \leq C \frac{\kappa}{H}\}} |\psi(x)|^2 dx,$$

where $t(x) = \text{dist}(x, \Gamma)$.

Proof. Let

$$\zeta = (\kappa H)^{-1/3}. \quad (6.13)$$

The assumption on κ and H ensures that

$$\kappa^{-1} \lesssim \zeta \ll 1. \quad (6.14)$$

We will prove Theorem 6.3 by establishing the following two estimates (away from the boundary or in a neighborhood of the boundary),

$$\int_{\{\text{dist}(x, \partial\Omega) \geq \zeta\}} e^{2m_0 \frac{H}{\kappa} t(x)} \left(\frac{1}{\kappa^2} |(\nabla - i\kappa H \mathbf{A})\psi|^2 + |\psi(x)|^2\right) dx \leq C_1 \int_{\{t(x) \leq C \frac{\kappa}{H}\}} |\psi(x)|^2 dx, \quad (6.15)$$

and

$$\int_{\{\text{dist}(x, \partial\Omega) \leq \zeta\}} e^{2m_0 \frac{H}{\kappa} t(x)} \left(\frac{1}{\kappa^2} |(\nabla - i\kappa H \mathbf{A})\psi|^2 + |\psi(x)|^2\right) dx \leq C_2 \int_{\{t(x) \leq C \frac{\kappa}{H}\}} |\psi(x)|^2 dx, \quad (6.16)$$

expressing the localization of the energy of ψ near Γ .

The proof of (6.15) and (6.16) is divided into several steps.

Step 1.

Consider the parameters

$$\xi \in (1, \infty), \quad \sigma = \frac{H}{\kappa^2}, \quad \ell = \frac{\xi}{\sigma\kappa}. \quad (6.17)$$

Let

$$f(x) = \chi(x) \exp(\ell^{-1} t(x)),$$

and

$$g(x) = \eta(x) \exp(\ell^{-1} t(x)).$$

The functions $\chi \in C_c^\infty(\Omega)$ and $\eta \in C^\infty(\overline{\Omega})$ satisfy,

$$\begin{cases} 0 \leq \chi \leq 1 & \text{in } \Omega, \\ \chi = 1 & \text{in } \{\text{dist}(x, \partial\Omega) \geq \zeta\} \cup \{t(x) \geq \ell\}, \\ \chi = 0 & \text{in } \{\text{dist}(x, \partial\Omega) \leq \frac{1}{2}\zeta\} \cup \{t(x) \leq \frac{1}{2}\ell\}, \\ |\nabla \chi| \leq C\kappa & \text{in } \Omega. \end{cases} \quad (6.18)$$

and

$$\begin{cases} 0 \leq \eta \leq 1 & \text{in } \Omega, \\ \eta = 1 & \text{in } \{\text{dist}(x, \partial\Omega) \leq \zeta\} \cup \{t(x) \geq \ell\}, \\ \eta = 0 & \text{in } \{\text{dist}(x, \partial\Omega) \geq 2\zeta\} \cup \{t(x) \leq \frac{1}{2}\ell\}, \\ |\nabla \eta| \leq C\kappa & \text{in } \Omega. \end{cases} \quad (6.19)$$

Here we have used for the control of the gradient (6.14) and that

$$\kappa^{-1} \ll \ell \lesssim 1.$$

Using the Ginzburg-Landau equation in (4.1), we write,

$$\int_{\Omega} \left(|(\nabla - i\kappa H \mathbf{A})f\psi|^2 - |\nabla f|^2 |\psi|^2 \right) dx = \kappa^2 \int_{\Omega} (|\psi|^2 - |\psi|^4) f^2 dx \leq \kappa^2 \int_{\Omega} |f\psi|^2 dx, \quad (6.20)$$

and

$$\int_{\Omega} \left(|(\nabla - i\kappa H \mathbf{A})g\psi|^2 - |\nabla g|^2 |\psi|^2 \right) dx = \kappa^2 \int_{\Omega} (|\psi|^2 - |\psi|^4) g^2 dx \leq \kappa^2 \int_{\Omega} |g\psi|^2 dx. \quad (6.21)$$

Step 2.

In this step, we determine a lower bound of $\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 dx$. Notice that $f\psi \in C_c^\infty(\Omega)$.

Consequently, we may write (see (6.6)),

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 dx \geq \kappa H \int_{\Omega} |\operatorname{curl} \mathbf{F}| |f\psi|^2 dx - \kappa H \int_{\Omega} |\operatorname{curl} \mathbf{A} - \operatorname{curl} \mathbf{F}| |f\psi|^2 dx.$$

We use the following estimates,

$$\int_{\Omega} |\operatorname{curl} \mathbf{F}| |f\psi|^2 dx \geq \frac{1}{M} \ell \int_{\Omega} |f\psi|^2 dx \quad [\text{by (6.7)}]$$

$$\int_{\Omega} |\operatorname{curl} \mathbf{A} - \operatorname{curl} \mathbf{F}| |f\psi|^2 dx \leq \frac{C}{H} \left(\frac{\kappa}{H} \right)^{1/6} \|f\psi\|_4^2 \quad [\text{by (6.3)}],$$

and obtain

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 dx \geq \frac{1}{M} \kappa H \ell \int_{\Omega} |f\psi|^2 dx - C \kappa \left(\frac{\kappa}{H} \right)^{1/6} \|f\psi\|_4^2.$$

Notice that $f\psi \in C_c^\infty(\Omega) \subset H^1(\mathbb{R}^2)$. By the continuous Sobolev embedding of $H^1(\mathbb{R}^2)$ in $L^4(\mathbb{R}^2)$ and a scaling, we get for all $\eta \in (0, 1)$,

$$\begin{aligned} \|f\psi\|_4^2 &= \| |f\psi| \|_4^2 \\ &\leq C_{\text{Sob}} \left(\eta \|\nabla |f\psi|\|_2^2 + \eta^{-1} \| |f\psi| \|_2^2 \right) \\ &\leq C_{\text{Sob}} \left(\eta \|(\nabla - i\kappa H \mathbf{A})f\psi\|_2^2 + \eta^{-1} \| |f\psi| \|_2^2 \right) \quad [\text{By the diamagnetic inequality}]. \end{aligned}$$

We select $\eta = \frac{1}{C C_{\text{Sob}}} \kappa^{-1} \left(\frac{\kappa}{H} \right)^{-\frac{1}{6}}$ and obtain,

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 dx \geq \left(\frac{\kappa H \ell}{2M} - \widehat{C} \kappa^2 \left(\frac{\kappa}{H} \right)^{1/3} \right) \int_{\Omega} |f\psi|^2 dx. \quad (6.22)$$

Thanks to the choice of the parameters in (6.17), the lower bound in (6.22) becomes,

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 dx \geq \left(\frac{\xi \kappa^2}{2M} - \widehat{C} \kappa^2 \left(\frac{\kappa}{H} \right)^{1/3} \right) \int_{\Omega} |f\psi|^2 dx. \quad (6.23)$$

Step 3.

We insert (6.23) into (6.20) and use that

$$\begin{aligned} \int_{\Omega} |\nabla f|^2 |\psi|^2 dx &\leq 2\ell^{-2} \int_{\Omega} |f\psi|^2 dx + 2 \int_{\Omega} |\nabla \chi|^2 \exp(2\ell^{-1}t(x)) |\psi|^2 dx \\ &\leq 2\ell^{-2} \int_{\Omega} |f\psi|^2 dx + C \kappa^2 \int_{\Omega} |g\psi|^2 dx + C \kappa^2 \int_{\{\ell^{-1}t(x) \leq 1\}} |\psi|^2 dx, \end{aligned} \quad (6.24)$$

to obtain,

$$\begin{aligned} \int_{\Omega} \left(\frac{1}{2} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 + \frac{1}{2} \left(\frac{\xi \kappa^2}{2M} - 2 \frac{\sigma^2}{\xi^2} \kappa^2 - \widehat{C} \kappa^2 \left(\frac{\kappa}{H} \right)^{1/3} \right) |f\psi|^2 dx \right) \\ \leq \widehat{C} \kappa^2 \int_{\Omega} |g\psi|^2 dx + \widehat{C} \kappa^2 \int_{\{\ell^{-1}t(x) \leq 1\}} |\psi|^2 dx. \end{aligned} \quad (6.25)$$

Step 4.

We will determine a lower bound of $\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})g\psi|^2 dx$. We cover the set

$$\Omega_{\zeta, \ell} = \{x \in \overline{\Omega} : \text{dist}(x, \partial\Omega) \leq 2\zeta, \text{dist}(x, \Gamma) \geq \ell\}$$

by a family of squares (in tubular coordinates),

$$\mathcal{K}(a_j, \zeta) = \{x \in \overline{\Omega} : \text{dist}(x, \partial\Omega) \leq 2\zeta, \text{dist}_{\partial\Omega}(p(x), a_j) \leq 2\zeta\},$$

where:

- $\text{dist}_{\partial\Omega}$ is the arc-length distance along $\partial\Omega$.
- If $x \in \Omega_{\zeta, \ell}$ and ζ is sufficiently small, $p(x)$ is the unique point on $\partial\Omega$ satisfying $\text{dist}(x, p(x)) = \text{dist}(x, \partial\Omega)$.
- For all j , $a_j \in \partial\Omega \cap \overline{\Omega_{\zeta, \ell}}$.

Let (χ_j) be a partition of unity such that

$$\sum_j \chi_j^2 = 1, \quad \sum_j |\nabla \chi_j|^2 \leq C\zeta^{-2}, \quad \text{supp } \chi_j \subset \mathcal{K}(a_j, 2\zeta).$$

There holds the decomposition formula

$$\begin{aligned} \int_{\Omega} |(\nabla - i\kappa H \mathbf{A})g\psi|^2 dx &= \sum_j \int_{\Omega} |(\nabla - i\kappa H \mathbf{A})\chi_j g\psi|^2 dx - \sum_j \int_{\Omega} |\nabla \chi_j|^2 |g\psi|^2 dx \\ &\geq \sum_j \int_{\Omega} |(\nabla - i\kappa H \mathbf{A})\chi_j g\psi|^2 dx - C\zeta^{-2} \int_{\Omega} |g\psi|^2 dx. \end{aligned} \quad (6.26)$$

Next we define the gauge function

$$\alpha_j = (\mathbf{A}(a_j) - \mathbf{F}(a_j)) \cdot (x - a_j).$$

Using the Cauchy-Schwarz inequality, Proposition 4.2 and Theorem 6.1, we may write,

$$\begin{aligned} \int_{\Omega} |(\nabla - i\kappa H \mathbf{A})\chi_j g\psi|^2 dx &= \int_{\Omega} |(\nabla - i\kappa H(\mathbf{A} - \nabla \alpha_j))e^{-i\kappa H \alpha_j} \chi_j g\psi|^2 dx \\ &\geq \frac{1}{2} \int_{\Omega} |(\nabla - i\kappa H \mathbf{F})e^{-i\kappa H \alpha_j} \chi_j g\psi|^2 dx - C\kappa^2 \left(\frac{\kappa}{H}\right)^{1/3} \zeta^{2\alpha} \int_{\Omega} |\chi_j g\psi|^2 dx. \end{aligned} \quad (6.27)$$

Next, we observe that there exists a gauge function φ_j satisfying

$$|\mathbf{F}(x) - (B_0(x_j)\mathbf{A}_0(x - a_j) + \nabla \varphi_j)| \leq C\zeta^2 \quad \text{in } \mathcal{K}_j(a_j, \zeta).$$

Again, using the Cauchy-Schwarz inequality, we may write,

$$\begin{aligned} &\int_{\Omega} |(\nabla - i\kappa H \mathbf{F})e^{-i\kappa H \alpha_j} \chi_j g\psi|^2 dx \\ &\geq \frac{1}{2} \int_{\Omega} |(\nabla - i\kappa H B_0(a_j)\mathbf{A}_0(x - a_j))e^{-i\kappa H \varphi_j} e^{-i\kappa H \alpha_j} \chi_j g\psi|^2 dx - \kappa^2 H^2 \zeta^4 \int_{\Omega} |\chi_j g\psi|^2 dx. \end{aligned} \quad (6.28)$$

Now, we are reduced to the analysis of the Neumann realization of the Schrödinger operator with a constant magnetic field equal to $\kappa H B_0(a_j)$ in our case. In the half-plane case, the ground state energy of this operator is $\Theta_0 \kappa H |B_0(a_j)|$, where the constant Θ_0 is universal and satisfies $\Theta_0 \in (\frac{1}{2}, 1)$. The result remains asymptotically true in general domains with smooth and compact boundary [15]. More precisely, there exists a function

$$\text{err} : \mathbb{R}_+ \rightarrow \mathbb{R}_+,$$

such that $\lim_{|b| \rightarrow \infty} \text{err}(b) = 0$ and

$$\forall b, \quad \lambda^N(b) \geq \Theta_0 |b| - |b| \text{err}(b),$$

where $\lambda^N(b)$ is the lowest eigenvalue of the operator $-(\nabla - ib\mathbf{A}_0)^2$ in $L^2(\Omega)$ with Neumann boundary condition.

Notice that by the assumptions on ℓ and the points (a_j) , we may use (6.7) with $x = a_j$ and get,

$$\forall j, \quad \kappa H |B_0(a_j)| \geq \frac{1}{M} \ell \kappa H \gg 1.$$

Moreover, the magnetic potentials $\mathbf{A}_0(x)$ and $\mathbf{A}_0(x - a_j)$ are gauge equivalent since

$$\mathbf{A}_0(x - a_j) = \mathbf{A}_0(x) - \mathbf{A}_0(a_j) = \mathbf{A}_0(x) - \nabla u_j(x),$$

with $u_j(x) = A_0(a_j) \cdot x$.

In that way, when κ is sufficiently large, we may write,

$$\begin{aligned} \int_{\Omega} |(\nabla - i\kappa H B_0(a_j))\mathbf{A}_0(x - a_j)e^{-i\kappa H \varphi_j} e^{-i\kappa H \alpha_j} \chi_j g \psi|^2 dx \\ \geq \frac{\Theta_0}{2} \kappa H |B_0(a_j)| \int_{\Omega} |\chi_j g \psi|^2 dx \geq \frac{1}{4M} \ell \kappa H \int_{\Omega} |\chi_j g \psi|^2 dx. \end{aligned} \quad (6.29)$$

Collecting the estimates in (6.26), (6.27), (6.28) and (6.29), we get,

$$\int_{\Omega} |(\nabla - i\kappa H \mathbf{A})g\psi|^2 dx \geq \kappa \left(\frac{H\ell}{4M} - C\kappa H^2 \zeta^4 - \frac{C}{\kappa \zeta^2} - C\kappa \left(\frac{\kappa}{H}\right)^{1/3} \zeta^{2\alpha} \right) \int_{\Omega} |g\psi|^2 dx. \quad (6.30)$$

Recall the definition of the parameters in (6.17) and (6.13):

$$\zeta = (H\kappa)^{-1/3} = \sigma^{-1/3} \kappa^{-1}.$$

We insert (6.30) into (6.21) and use that

$$\begin{aligned} \int_{\Omega} |\nabla g|^2 |\psi|^2 dx &\leq 2\ell^{-2} \int_{\Omega} |g\psi|^2 dx + 2 \int_{\Omega} |\nabla \eta|^2 \exp(2\ell^{-1}t(x)) |\psi|^2 dx \\ &\leq 2\ell^{-2} \int_{\Omega} |g\psi|^2 dx + C\kappa^2 \int_{\Omega} |f\psi|^2 dx + C\kappa^2 \int_{\{\ell^{-1}t(x) \leq 1\}} |\psi|^2 dx, \end{aligned}$$

to obtain,

$$\begin{aligned} \int_{\Omega} \left(\frac{1}{2} |(\nabla - i\kappa H \mathbf{A})g\psi|^2 + \frac{1}{2} \left(\frac{\xi \kappa^2}{4M} - 2\ell^{-2} - C\sigma^{2/3} \kappa - C\kappa \left(\frac{\kappa}{H}\right)^{1/3} \zeta^{2\alpha} \right) |g\psi|^2 dx \right) \\ \leq \kappa^2 \int_{\Omega} |f\psi|^2 dx + C\kappa^2 \int_{\{\kappa t(x) \leq 1\}} |\psi|^2 dx. \end{aligned} \quad (6.31)$$

Step 5.

Adding the two inequalities in (6.25) and (6.31), we get,

$$\begin{aligned} \frac{1}{2} \int_{\Omega} (|(\nabla - i\kappa H \mathbf{A})g\psi|^2 + |(\nabla - i\kappa H \mathbf{A})f\psi|^2) dx \\ + \frac{1}{2} \int_{\Omega} \left(\frac{\xi \kappa^2}{4M} - C\frac{\sigma^2}{\xi^2} \kappa^2 - C\kappa^2 - C\sigma^{2/3} \kappa - C\kappa \left(\frac{\kappa}{H}\right)^{1/3} \zeta^{2\alpha} \right) (|f\psi|^2 + |g\psi|^2) \\ \leq C\kappa^2 \int_{\{\kappa t(x) \leq 1\}} |\psi|^2 dx. \end{aligned} \quad (6.32)$$

Recall that σ satisfies $\kappa^{-1} \ll \sigma \leq \Lambda$. We select ξ sufficiently large such that

$$\frac{\xi}{4M} - C\frac{\Lambda^2}{\xi^2} - C > 2.$$

Since $(\frac{\kappa}{H})^{1/3} \ll 1$ and $\zeta \ll 1$, we get,

$$\int_{\Omega} \left(\frac{1}{2} |(\nabla - i\kappa H \mathbf{A})f\psi|^2 + \frac{\kappa^2}{2} |f\psi|^2 dx \right) \leq C\kappa^2 \int_{\{\ell^{-1}t(x) \leq 1\}} |\psi|^2 dx,$$

and

$$\int_{\Omega} \left(\frac{1}{2} |(\nabla - i\kappa H \mathbf{A})g\psi|^2 + \frac{\kappa^2}{2} |g\psi|^2 dx \right) \leq C\kappa^2 \int_{\{\ell^{-1}t(x) \leq 1\}} |\psi|^2 dx.$$

Thanks to the definitions of f and g , the two aforementioned inequalities yield the inequalities in (6.15) and (6.16) with $m_0 = 1/\xi$. \square

As a consequence of Theorem 6.3, we get an improvement of the bound given in Theorem 6.1.

Proposition 6.4. *Under the assumptions of Theorem 6.3, there holds,*

$$\|\psi\|_2 \leq C \sqrt{\frac{\kappa}{H}}.$$

Combining the results in Propositions 4.1, 4.2 and 6.4, we obtain the improved estimates:

Proposition 6.5. *Under the assumptions of Theorem 6.3 and Proposition 4.2, there holds,*

$$\begin{aligned} \|\operatorname{curl} \mathbf{A} - \operatorname{curl} \mathbf{F}\|_2 &\leq \frac{C}{H} \sqrt{\frac{\kappa}{H}}, \\ \|\mathbf{A} - \mathbf{F}\|_{C^{1,\alpha}(\overline{\Omega})} &\leq C_{\alpha} \sqrt{\frac{\kappa}{H}}, \\ \|\mathbf{A} - \mathbf{F}\|_{C^{0,\alpha}(\overline{\Omega})} &\leq \frac{\widehat{C}_{\alpha}}{H} \sqrt{\frac{\kappa}{H}}. \end{aligned}$$

7. ENERGY LOWER BOUND

In this section, we will derive lower bounds of the following energy,

$$\mathcal{E}_0(\psi, \mathbf{A}; U) = \int_U \left(|(\nabla - i\kappa H \mathbf{A})\psi|^2 - \kappa^2 |\psi|^2 + \frac{\kappa^2}{2} |\psi|^4 \right) dx, \quad (7.1)$$

where $U \subset \mathbb{R}^2$ is an open set such that $\overline{U} \subset \Omega$.

Proposition 7.1. *Let $\Lambda > 0$ and $\epsilon : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\lim_{\kappa \rightarrow \infty} \kappa \epsilon(\kappa) = \infty$ and $\lim_{\kappa \rightarrow \infty} \epsilon(\kappa) = 0$. For $\alpha \in (0, 1)$, there exist positive constants C and κ_0 such that, for $\ell \in (0, 1)$, $\delta \in (0, 1)$, $a_j \in \Gamma$, $D(a_j, \ell) \subset \Omega$, $x_j \in \overline{D(a_j, \ell)} \cap \Gamma$, $h \in C_c^{\infty}(D(a_j, \ell))$ a function satisfying $\|h\|_{\infty} \leq 1$, (ψ, \mathbf{A}) a critical point of the functional in (1.1), $\kappa \geq \kappa_0$, and*

$$\epsilon(\kappa)\kappa^2 \leq H \leq \Lambda\kappa^2,$$

the following holds:

$$\mathcal{E}_0(h\psi, \mathbf{A}; D(a_j, \ell)) \geq (1 - \delta) 2\ell \kappa \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) - r,$$

where

$$r = C \left(\delta \kappa^2 + \delta^{-1} \left(\frac{\kappa^3}{H} \ell^{2\alpha} + \kappa^2 H^2 \ell^6 \right) \right) \int_{D(a_j, \ell)} |h\psi|^2 dx.$$

Proof. Let $\alpha_j = (\mathbf{A}(a_j) - \mathbf{F}(a_j)) \cdot (x - a_j)$. Thanks to Proposition 6.5, we have

$$|\mathbf{A} - (\mathbf{F} + \nabla \alpha_j)| \leq C \|\mathbf{A} - \mathbf{F}\|_{C^{0,\alpha}(\overline{\Omega})} |x - a_j|^{\alpha} \leq \frac{C}{H} \sqrt{\frac{\kappa}{H}} \ell^{\alpha} \quad \text{in } D(a_j, \ell). \quad (7.2)$$

Notice that,

$$\begin{aligned} \mathcal{E}_0(h\psi, \mathbf{A}; D(a_j, \ell)) &= \mathcal{E}_0(h\psi e^{-i\kappa H\alpha_j}, \mathbf{A} - \nabla\alpha_j; D(a_j, \ell)) \\ &\geq (1 - \delta)\mathcal{E}_0(h\psi e^{-i\kappa H\alpha_j}, \mathbf{F}; D(a_j, \ell)) \\ &\quad - C \left(\delta\kappa^2 \int_{D(a_j, \ell)} |h\psi|^2 dx + \delta^{-1}\kappa^2 H^2 \int_{D(a_j, \ell)} |\mathbf{A} - (\mathbf{F} + \nabla\alpha_j)|^2 |h\psi|^2 dx \right) \end{aligned} \quad (7.3)$$

Using (7.2), we get,

$$\mathcal{E}_0(h\psi, \mathbf{A}; D(a_j, \ell)) \geq (1 - \delta)\mathcal{E}_0(h\psi e^{-i\kappa H\alpha_j}, \mathbf{F}; D(a_j, \ell)) - C \left(\delta\kappa^2 + \delta^{-1}\frac{\kappa^3}{H}\ell^{2\alpha} \right) \int_{D(a_j, \ell)} |h\psi|^2 dx. \quad (7.4)$$

Let

$$f_j = h\psi e^{-i\kappa H\alpha_j} e^{i\kappa H\phi_j},$$

where ϕ_j is defined in Proposition 4.3.

Notice that $f_j \in H_0^1(D(a_j, \ell))$, $\|f_j\|_\infty \leq 1$ and, using (4.9),

$$\begin{aligned} &\mathcal{E}_0(h\psi e^{-i\kappa H\alpha_j}, \mathbf{F}; D(a_j, \ell)) \\ &= \mathcal{E}_0(f_j, \mathbf{F} - \nabla\phi_j; D(a_j, \ell)) \\ &\geq (1 - \delta)\mathcal{E}_0(f_j, |\nabla B_0(x_j)|\mathbf{A}_{\text{app}, \nu_j}(x - a_j); D(a_j, \ell)) - C(\delta\kappa^2 + \delta^{-1}\kappa^2 H^2 \ell^6) \int_{D(a_j, \ell)} |f_j|^2 dx. \end{aligned} \quad (7.5)$$

We will use Theorem 3.11 to get a lower bound of the energy

$$\mathcal{E}_0(f_j, |\nabla B_0(x_j)|\mathbf{A}_{\text{app}, \nu_j}(x - a_j); D(a_j, \ell)).$$

Define

$$L = L_j = |\nabla B_0(x_j)| \frac{H}{\kappa^2}. \quad (7.6)$$

Performing the translation $x \mapsto x + a_j$, we get that

$$\mathcal{E}_0(f_j, |\nabla B_0(x_j)|\mathbf{A}_{\text{app}, \nu_j}(x - a_j); D(a_j, \ell)) = \mathcal{G}(f_j) \geq E_{\text{gs}, r}(\kappa, L; \ell). \quad (7.7)$$

Here \mathcal{G} is the functional in (3.40) and $E_{\text{gs}, r}(\kappa, L; \ell)$ is the ground state energy in (3.41).

Let

$$R = L^{1/3}\kappa\ell.$$

Now, Theorems 3.11 and 3.8 applied successively tell us that

$$\mathcal{E}_0(f_j, |\nabla B_0(x_j)|\mathbf{A}_{\text{app}, \nu_j}(x - a_j); D(a_j, \ell)) \geq \mathbf{e}_{\text{gs}}(\nu, L; R) \geq 2RE(L) = L^{1/3}\kappa\ell E(L).$$

Recall the definition of L in (7.6). We insert the aforementioned estimate into (7.7). In that way, we infer from (7.5) and (7.4) the lower bound of Proposition 7.1. \square

Proposition 7.2. *For $r > 0$, $h \in C^\infty(\mathbb{R}^2)$ satisfying $\|h\|_\infty \leq 1$, and (ψ, \mathbf{A}) a critical point of the functional in (1.1), the following lower bound holds,*

$$\mathcal{E}_0(h\psi, \mathbf{A}; D(a_j, r) \cap \Omega) \geq -\pi\kappa^2 r^2. \quad (7.8)$$

Proof. Notice that all terms in $\mathcal{E}_0(h\psi, \mathbf{A}; D(a_j, r) \cap \Omega)$ are positive except the integral of $|h\psi|^2$. Thus,

$$\mathcal{E}_0(h\psi, \mathbf{A}; D(a_j, r) \cap \Omega) \geq -\kappa^2 \int_{\Omega \cap D(a_j, r)} |h\psi|^2 dx.$$

This finishes the proof of the proposition upon using $\|h\psi\|_\infty \leq 1$ and $\|\psi\|_\infty \leq 1$. \square

Theorem 7.3. *Let $\Lambda > 0$ and $\epsilon : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\lim_{\kappa \rightarrow \infty} \kappa \epsilon(\kappa) = \infty$ and $\lim_{\kappa \rightarrow \infty} \epsilon(\kappa) = 0$.*

There exist $\kappa_0 > 0$ and a function $\text{err} : \mathbb{R} \rightarrow \mathbb{R}$ such that the following is true:

- (1) $\lim_{\kappa \rightarrow \infty} \text{err}(\kappa) = 0$.
- (2) Let $D \subset \Omega$ be a regular open set, $h \in C^\infty(\overline{D})$, $\|h\|_\infty \leq 1$, (ψ, \mathbf{A}) a critical point of the functional in (1.1), $\kappa \geq \kappa_0$ and $\epsilon(\kappa)\kappa^2 \leq H \leq \Lambda\kappa^2$.
- (a) If $H \gg \kappa^{3/2}$, then,

$$\mathcal{E}_0(h\psi, \mathbf{A}; D) \geq \kappa \left(\int_{\Gamma \cap D} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) + \frac{\kappa^3}{H} \text{err}(\kappa). \quad (7.9)$$

- (b) If $H \lesssim \kappa^{3/2}$, then,

$$\mathcal{E}_0(h\psi, \mathbf{A}; D) \geq \kappa^2 \int_D g \left(\frac{H}{\kappa} |B_0(x)| \right) dx + \frac{\kappa^3}{H} \text{err}(\kappa). \quad (7.10)$$

Proof. Consider three parameters

$$a \in (0, 1), \quad \ell \in (0, 1), \quad \delta \in (0, 1),$$

and define the following sets,

$$D_1 = \{x \in \Omega : \text{dist}(x, \Gamma) < 2\sqrt{a}\ell\},$$

$$D_2 = \{x \in \Omega : \text{dist}(x, \Gamma) > \sqrt{a}\ell\}.$$

Let (χ_j) be a partition of unity satisfying

$$\sum_{j=1}^2 \chi_j^2 = 1, \quad \sum_{j=1}^2 |\nabla \chi_j|^2 \leq C(a\ell^2)^{-1}, \quad \text{supp } \chi_j \subset D_j \quad (j \in \{1, 2\}).$$

There holds the following decomposition of the energy,

$$\mathcal{E}_0(h\psi, \mathbf{A}; D) \geq \mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1) + \mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; D_2) - \sum_{j=1}^2 \int_\Omega |\nabla \chi_j|^2 |h\psi|^2 dx.$$

The error terms are controlled using the pointwise bounds on $|h|$, $|\psi|$, $|\nabla \chi_j|$, and the conditions on the support of χ_j . We obtain the following lower bound,

$$\mathcal{E}_0(h\psi, \mathbf{A}; D) \geq \mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1) + \mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; D_2) - C(\sqrt{a}\ell)^{-1}. \quad (7.11)$$

The regime $H \gg \kappa^{3/2}$.

In this regime, we shall see that $\mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1)$ is the leading term and $\mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; D_2)$ is an error term.

Lower bound of the term $\mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1)$.

Consider a constant $c > 0$ and distinct points (a_j) in Γ such that,

$$\forall j, \quad \ell - a\ell \leq \text{dist}(a_j, a_{j+1}) \leq \ell - a\ell.$$

Choose the constant a sufficiently small so that

$$D_1 = \{x \in \Omega : \text{dist}(x, \Gamma) < 2\sqrt{a}\ell\} \subset \bigcup_j D(a_j, \ell).$$

Consider a partition of unity satisfying

$$\sum_j f_j^2 = 1 \text{ in } D_1, \quad \text{supp } f_j \subset D(a_j, \ell), \quad \sum_j |\nabla f_j|^2 \leq \frac{C}{a^2 \ell^2}.$$

Notice that the support of each ∇f_j is in $D(a_j, \ell) \cap D(a_{j+1}, \ell) \cap D(a_{j-1}, \ell)$ and that the points (a_j) are selected such that the last domain has an area proportional to $\sqrt{a}\ell \times a\ell = a\sqrt{a}\ell^2$.

The partition of unity (f_j) allows us to decompose the energy as follows,

$$\begin{aligned} \mathcal{E}_0(\chi_1 h \psi, \mathbf{A}; D_1) &\geq \sum_j \mathcal{E}_0(f_j \chi_1 h \psi, \mathbf{A}; D_1) - \sum_j \left\| |\nabla f_j| \chi_1 h \psi \right\|_2^2 \\ &\geq \sum_j \mathcal{E}_0(h_j \psi, \mathbf{A}; D_1) - \frac{C}{\sqrt{a} \ell}, \end{aligned} \quad (7.12)$$

where $h_j = f_j \chi_1 h \psi$ is supported in $D \cap D(a_j, \ell)$.

If $D(a_j, \ell) \cap \partial\Omega \neq \emptyset$, then we can apply Proposition 7.2. Since $\Gamma \cap \partial\Omega$ is a finite set, then we get,

$$\sum_{D(a_j, \ell) \cap \partial\Omega \neq \emptyset} \mathcal{E}_0(h_j \psi, \mathbf{A}; D_1) \geq -C\kappa^2 \ell^2.$$

We select the parameter ℓ as follows,

$$\ell = \delta H^{-1/3}. \quad (7.13)$$

In that way, we obtain,

$$\ell \ll 1, \quad \kappa^2 \ell^2 \ll \frac{\kappa^3}{H}, \quad \frac{1}{\sqrt{a} \ell} \ll \frac{\kappa^3}{H},$$

and

$$\sum_{D(a_j, \ell) \cap \partial\Omega \neq \emptyset} \mathcal{E}_0(h_j \psi, \mathbf{A}; D_1) \geq \frac{\kappa^3}{H} o(1) \quad (\kappa \rightarrow \infty). \quad (7.14)$$

If $D(a_j, \ell) \subset D^c$, then $h_j = 0$ and

$$\mathcal{E}_0(h_j \psi, \mathbf{A}; D_1) = 0.$$

Now, if $j \in \mathcal{I} = \{j : D(a_j, \ell) \subset \Omega \text{ and } D(a_j, \ell) \cap D \neq \emptyset\}$, then we can apply Proposition 7.1 and get,

$$\begin{aligned} \sum_{j \in \mathcal{I}} \mathcal{E}_0(h_j \psi, \mathbf{A}; D_1) &\geq (1 - \delta) 2\ell \kappa \sum_{j \in \mathcal{I}} \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \\ &\quad - C \left(\delta \kappa^2 + \delta^{-1} \left(\frac{\kappa^3}{H} \ell^{2\alpha} + \kappa^2 H^2 \ell^6 \right) \right) \int_{\Omega} |h \psi|^2 dx, \end{aligned}$$

where, for all j , x_j is an arbitrary point in $\overline{D(a_j, \ell)}$.

Thanks to Proposition 6.4 and the choice of ℓ in (7.13), we get further,

$$\begin{aligned} \sum_{j \in \mathcal{I}} \mathcal{E}_0(h_j \psi, \mathbf{A}; D_1) &\geq (1 - \delta) 2\ell \kappa \sum_{j \in \mathcal{I}} \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \\ &\quad - C \left(\delta + \delta^{2\alpha-1} \frac{\kappa}{H} H^{-2\alpha/3} \right) \frac{\kappa^3}{H}. \end{aligned}$$

Theorem 3.12 allows us to write,

$$\left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \leq C \frac{\kappa^2}{H}. \quad (7.15)$$

Consequently,

$$\begin{aligned} \sum_{j \in \mathcal{I}} \mathcal{E}_0(h_j \psi, \mathbf{A}; D_1) &\geq 2\ell \kappa \sum_{j \in \mathcal{I}} \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \\ &\quad - C \left(\delta + \delta^{2\alpha-1} \frac{\kappa}{H} H^{-2\alpha/3} \right) \frac{\kappa^3}{H}. \end{aligned}$$

Inserting this and (7.14) into (7.12), and using that $(\sqrt{a}\ell)^{-1} \ll \frac{\kappa^3}{H}$, we get,

$$\begin{aligned} \mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1) &\geq \kappa \sum_{j \in \mathcal{I}} 2\ell \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \\ &\quad - C \left(\delta + \delta^{2\alpha-1} \frac{\kappa}{H} H^{-2\alpha/3} \right) \frac{\kappa^3}{H}. \end{aligned} \quad (7.16)$$

Thanks to (7.15), the sum

$$\sum_{j \in \mathcal{I}} 2\ell \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \quad (7.17)$$

is of order κ^2/H . Let $\eta > 0$. Select ℓ_0 sufficiently small such that, for all $\ell \in (0, \ell_0)$, the arc-length measure of $D(a_j, \ell) \cap \Gamma$ along Γ satisfies,

$$2\ell - \ell \frac{\eta}{2} \leq |D(a_j, \ell) \cap \Gamma| \leq 2\ell + \ell \frac{\eta}{2}.$$

Thus, replacing 2ℓ by $|D(a_j, \ell) \cap \Gamma|$ in the sum in (7.17) produces an error of order $\eta\ell$. Now, select $x_j \in \overline{D(a_j, \ell)}$ such that

$$|\nabla B_0(x_j)|^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) = \max_{D(a_j, \ell)} |\nabla B_0(x)|^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right).$$

In that way, the sum in (7.17) satisfies,

$$\begin{aligned} &\sum_{j \in \mathcal{I}} 2\ell \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) \\ &\geq \sum_{j \in \mathcal{I}} \left(\int_{D(a_j, \ell) \cap \Gamma} \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) dx \right) - C\eta \frac{\kappa^2}{H}. \end{aligned} \quad (7.18)$$

Recall (7.15). Since the balls $(D(a_j, \ell))$ overlap in a region of length $\mathcal{O}(a\ell)$, and the number of these balls is inversely proportional to ℓ , then

$$\begin{aligned} &\sum_{j \in \mathcal{I}} \left(\int_{D(a_j, \ell) \cap \Gamma} \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) dx \right) \\ &\geq \int_{D \cap \Gamma} \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x_j)| \frac{H}{\kappa^2} \right) dx - Ca \frac{\kappa^2}{H}. \end{aligned}$$

Inserting this into (7.18), then inserting the resulting inequality into (7.16), we get,

$$\begin{aligned} \mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1) &\geq \kappa \int_{\Gamma \cap D} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \\ &\quad - C \left(a + \delta + \delta^{2\alpha-1} \frac{\kappa}{H} H^{-2\alpha/3} + \eta \right) \frac{\kappa^3}{H}. \end{aligned} \quad (7.19)$$

Recall that $\alpha > 0$. Taking $\kappa \rightarrow \infty$, we get,

$$\begin{aligned} \liminf_{\kappa \rightarrow \infty} \frac{H}{\kappa^3} \left\{ \mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1) - \kappa \int_{\Gamma \cap D} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right\} \\ \geq -C(a + \delta + \eta). \end{aligned}$$

Taking $\eta \rightarrow 0_+$, we obtain,

$$\liminf_{\kappa \rightarrow \infty} \frac{H}{\kappa^3} \left\{ \mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1) - \kappa \int_{\Gamma \cap D} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right\} \geq -C(a+\delta),$$

i.e. when κ is sufficiently large,

$$\mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1) \geq \kappa \int_{\Gamma \cap D} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) - 2C(\delta + a) \frac{\kappa^3}{H}. \quad (7.20)$$

Lower bound of the term $\mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; D_2)$.

Since $H \gg \kappa^{3/2}$, the parameter ℓ defined in (7.13) satisfies,

$$\ell = \delta H^{-1/3} = \delta \frac{\kappa}{H} \frac{H^{1/3}}{\kappa} \gg \frac{\kappa}{H}.$$

Thanks to the exponential decay in Theorem 6.3, there holds,

$$\kappa^2 \int_{D_2} |\psi|^2 dx \ll \frac{\kappa^3}{H},$$

and

$$\mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; D_2) \geq -\frac{\kappa^3}{H} o(1) \quad (\kappa \rightarrow \infty). \quad (7.21)$$

Inserting (7.21) and (7.20) into (7.11), we get,

$$\mathcal{E}_0(h\psi, \mathbf{A}; D) \geq \kappa \int_{\Gamma \cap D} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) - C(\delta + a + o(1)) \frac{\kappa^3}{H}.$$

Taking the successive limits

$$\liminf_{\kappa \rightarrow \infty}, \quad \lim_{\delta \rightarrow 0_+}, \quad \lim_{a \rightarrow 0_+},$$

we arrive at

$$\mathcal{E}_0(h\psi, \mathbf{A}; D) \geq \kappa \int_{\Gamma \cap D} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) - \frac{\kappa^3}{H} o(1). \quad (7.22)$$

This finishes the proof of Theorem 7.3 in the case $H \gg \kappa^{3/2}$.

The regime $H \lesssim \kappa^{3/2}$.

In this regime, we shall see that $\mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1)$ is an error term and $\mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; D_2)$ is the leading term.

Since $H \lesssim \kappa^{3/2}$, the parameter ℓ introduced in (7.13) satisfies

$$\ell \lesssim \delta \frac{\kappa}{H}.$$

Consequently, we have,

$$\mathcal{E}_0(\chi_1 h\psi, \mathbf{A}; D_1) \geq -\kappa^2 \int_{\Omega} |\chi_1 h\psi|^2 dx \geq C\kappa^2 \ell \geq -\delta \frac{\kappa^3}{H}. \quad (7.23)$$

Unlike the regime $H \gg \kappa^{3/2}$, we can no more ignore the energy in $\{\sqrt{a}\ell \leq \text{dist}(x, \Gamma) \leq \frac{\kappa}{H}\}$.

We introduce the two parameters,

$$m > 1 \quad \text{and} \quad \zeta \in (0, \ell), \quad (7.24)$$

and the domain,

$$U = \left\{ x \in D_2 : \text{dist}(x, \Gamma) \geq m \frac{\kappa}{H} \right\}. \quad (7.25)$$

Thanks to the exponential decay in Theorem 6.3, we get,

$$\kappa^2 \int_U |\psi|^2 dx \leq C e^{-2m m_0} \frac{\kappa^3}{H}.$$

and

$$\mathcal{E}_0(\chi_2 h \psi, \mathbf{A}; U) \geq -C e^{-2m m_0} \frac{\kappa^3}{H}. \quad (7.26)$$

Our next task is to determine a lower bound of the energy $\mathcal{E}_0(\chi_2 h \psi, \mathbf{A}; D_2 \setminus U)$. Consider for $\zeta \in (0, 1)$ the lattice of squares $(\mathcal{Q}_{k, \zeta})_k$ generated by the square

$$\mathcal{Q}_\zeta = \left(-\frac{\zeta}{2}, \frac{\zeta}{2}\right) \times \left(-\frac{\zeta}{2}, \frac{\zeta}{2}\right).$$

Let

$$\begin{aligned} \mathcal{J}_{\text{blk}} &= \{k : \mathcal{Q}_{k, \zeta} \subset D_2 \setminus U \text{ and } \mathcal{Q}_{k, \zeta} \cap \partial\Omega = \emptyset\}, \\ \mathcal{J}_{\text{bnd}, 1} &= \{k : \mathcal{Q}_{k, \zeta} \cap (D_2 \setminus U) \neq \emptyset \text{ and } \mathcal{Q}_{k, \zeta} \cap \partial\Omega = \emptyset\}, \\ \mathcal{J}_{\text{bnd}, 2} &= \{k : \mathcal{Q}_{k, \zeta} \subset D_2 \setminus U \text{ and } \mathcal{Q}_{k, \zeta} \cap \partial\Omega \neq \emptyset\}. \end{aligned}$$

We have the obvious decomposition,

$$\mathcal{E}_0(\chi_2 h \psi, \mathbf{A}; D_2 \setminus U) \geq \sum_{k \in \mathcal{J}_{\text{blk}}} \mathcal{E}_0(\chi_2 h \psi, \mathbf{A}; \mathcal{Q}_{k, \zeta}) + \sum_{j=1}^2 \sum_{k \in \mathcal{J}_{\text{bnd}, j}} \mathcal{E}_0(\chi_2 h \psi, \mathbf{A}; \mathcal{Q}_{k, \zeta} \cap (D_2 \setminus U)). \quad (7.27)$$

Since $\Gamma \cap \partial\Omega$ is a finite set, then $N = \text{Card} \mathcal{J}_{\text{bnd}, 2}$ is bounded independently of κ . Now, the terms corresponding to $k \in \mathcal{J}_{\text{bnd}, j}$ are easily estimated as follows,

$$\sum_{j=1}^2 \sum_{k \in \mathcal{J}_{\text{bnd}, j}} \mathcal{E}_0(\chi_2 h \psi, \mathbf{A}; \mathcal{Q}_{k, \zeta} \cap (D_2 \setminus U)) \geq -\kappa^2 \int_{\{\text{dist}(x, \partial\Gamma) \leq C\zeta\}} |\psi|^2 dx - N \kappa^2 \zeta^2 \geq -C \kappa^2 \zeta. \quad (7.28)$$

For all k , let x_k be the center of the square $\mathcal{Q}_{k, \zeta}$ and a_k an arbitrary point in $\overline{\mathcal{Q}_{k, \zeta}}$. Repeating the proof of Proposition 7.1, we get, for all $k \in \mathcal{J}$ and $\eta \in (0, 1)$,

$$\begin{aligned} \mathcal{E}_0(\chi_2 h \psi, \mathbf{A}; \mathcal{Q}_{k, \zeta}) &\geq (1 - \eta) \mathcal{E}_0(\chi_2 h \psi e^{-i\kappa H u_k}, B_0(a_k) \mathbf{A}_0(x - x_k); \mathcal{Q}_{k, \zeta}) \\ &\quad - C \left(\eta \kappa^2 + \eta^{-1} \left(\frac{\kappa^3}{H} \zeta^{2\alpha} + \kappa^2 H^2 \zeta^4 \right) \right) \|\psi\|_{L^2(\mathcal{Q}_{k, \zeta})}^2, \end{aligned} \quad (7.29)$$

where u_k is a gauge function.

We select the parameter ζ as follows

$$\zeta = \eta H^{-1/2}. \quad (7.30)$$

Clearly, ζ satisfies,

$$\zeta \ll \ell \lesssim \delta \frac{\kappa}{H} \ll 1, \quad \kappa^2 \zeta \ll \frac{\kappa}{H}, \quad H^2 \zeta^4 = \eta^4, \quad \zeta \sqrt{\kappa H |B_0(a_k)|} \gtrsim \zeta \sqrt{\kappa H \sqrt{a} \ell} \gtrsim \eta \delta^{1/2} a^{1/4}.$$

Applying a scaling and a translation, we may use (3.55) and get,

$$\begin{aligned} &\frac{1}{(\zeta \sqrt{\kappa H |B_0(a_k)|})^2} \mathcal{E}_0(\chi_2 h \psi e^{-i\kappa H u_k}, B_0(a_k) \mathbf{A}_0(x - x_k); \mathcal{Q}_{k, \zeta}) \\ &\geq \frac{\kappa}{H |B_0(a_k)|} \left(g \left(\frac{H}{\kappa} |B_0(a_k)| \right) - C \frac{\sqrt{\frac{H}{\kappa} |B_0(a_k)|}}{\zeta \sqrt{\kappa H |B_0(a_k)|}} \right). \end{aligned}$$

We insert this into (7.29), sum over $k \in \mathcal{J}_{\text{blk}}$ and use Proposition 6.4 to get,

$$\sum_{k \in \mathcal{J}_{\text{blk}}} \mathcal{E}_0(\chi_2 h \psi, \mathbf{A}; \mathcal{Q}_{k, \zeta}) \geq \zeta^2 \kappa^2 \sum_{k \in \mathcal{J}_{\text{blk}}} \left(g \left(\frac{H}{\kappa} |B_0(a_k)| \right) - \frac{C}{\zeta \kappa} \right) - (C\eta + o(1)) \frac{\kappa^3}{H}.$$

The sum in the inequality above becomes a lower Riemann sum if for each k the point (a_k) is selected in $\overline{Q_{k,\ell}}$ as follows,

$$|B_0(a_k)| = \max\{|B_0(x)| : x \in \overline{Q_{k,\ell}}\}.$$

Notice that $\mathcal{N}_{\text{blk}} = \text{Card } \mathcal{J}_{\text{blk}}$ satisfies $\mathcal{N}_{\text{blk}} \times \zeta^2 \approx |D_2 \setminus U|$ as $\zeta \rightarrow 0$ and

$$|D_2 \setminus U| = |\{\sqrt{a}\ell \leq \text{dist}(x, \Gamma) \leq m\frac{\kappa}{H}\}| \leq Cm\frac{\kappa}{H}.$$

Consequently, we have,

$$\sum_{k \in \mathcal{J}_{\text{blk}}} \mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; \mathcal{Q}_{k,\zeta}) \geq \kappa^2 \int_{\mathcal{D}_\zeta} g\left(\frac{H}{\kappa}|B_0(x)|\right) dx - m\frac{C\kappa}{\zeta}\frac{\kappa}{H} - (C\eta + o(1))\frac{\kappa^3}{H},$$

where

$$\mathcal{D}_\zeta = \bigcup_{k \in \mathcal{J}_{\text{blk}}} \overline{Q_{k,\zeta}} \subset D_2 \setminus U.$$

Since the function g is non-positive, then we get that,

$$\begin{aligned} \sum_{k \in \mathcal{J}_{\text{blk}}} \mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; \mathcal{Q}_{k,\zeta}) &\geq \kappa^2 \int_{\mathcal{D}_2 \setminus U} g\left(\frac{H}{\kappa}|B_0(x)|\right) dx - Cm\frac{\kappa}{\zeta}\frac{\kappa}{H} - (C\eta + o(1))\frac{\kappa^3}{H} \\ &\geq \kappa^2 \int_{\mathcal{D}_2 \setminus U} g\left(\frac{H}{\kappa}|B_0(x)|\right) dx - Cm\eta^{-1}\frac{H^{1/2}}{\kappa}\frac{\kappa^3}{H} - (C\eta + o(1))\frac{\kappa^3}{H}. \end{aligned} \quad (7.31)$$

We insert (7.31) and (7.28) into (7.27). Since $g\left(\frac{H}{\kappa}|B_0(x)|\right) = 0$ in $\{|B_0(x)| \geq \frac{H}{\kappa}\}$ and $H \lesssim \kappa^{3/2}$, it results the inequality,

$$\mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; D_2 \setminus U) \geq \kappa^2 \int_D g\left(\frac{H}{\kappa}|B_0(x)|\right) dx - (C\eta + o(1))\frac{\kappa^3}{H}, \quad (\kappa \rightarrow \infty). \quad (7.32)$$

Combining (7.32) and (7.26), we get

$$\mathcal{E}_0(\chi_2 h\psi, \mathbf{A}; D_2) \geq \kappa^2 \int_D g\left(\frac{H}{\kappa}|B_0(x)|\right) dx - Ce^{-2m m_0}\frac{\kappa^3}{H} - (C\eta + o(1))\frac{\kappa^3}{H}.$$

Now, we insert this inequality and (7.23) into (7.11) to get,

$$\mathcal{E}_0(h\psi, \mathbf{A}; D) \geq \kappa^2 \int_D g\left(\frac{H}{\kappa}|B_0(x)|\right) dx - C(a + \delta + \eta + e^{-2m m_0} + o(1))\frac{\kappa^3}{H}.$$

By taking the successive limits,

$$\liminf_{\kappa \rightarrow \infty}, \quad \lim_{a \rightarrow 0_+}, \quad \lim_{\delta \rightarrow 0_+}, \quad \lim_{\eta \rightarrow 0_+}, \quad \lim_{m \rightarrow \infty},$$

we get

$$\mathcal{E}_0(h\psi, \mathbf{A}; D) \geq \kappa^2 \int_D g\left(\frac{H}{\kappa}|B_0(x)|\right) dx - \frac{\kappa^3}{H} o(1),$$

which finishes the proof of Theorem 7.3 in the regime $H \lesssim \kappa^{3/2}$. \square

We get by applying Theorem 7.3 with $D = \Omega$:

Corollary 7.4. *Let $\Lambda > 0$ and $\epsilon : \mathbb{R} \rightarrow \mathbb{R}_+$ such that $\lim_{\kappa \rightarrow \infty} \kappa \epsilon(\kappa) = \infty$ and $\lim_{\kappa \rightarrow \infty} \epsilon(\kappa) = 0$.*

There exist $\kappa_0 > 0$ and a function $\text{err} : \mathbb{R} \rightarrow \mathbb{R}$ such that the following is true:

- (1) $\lim_{\kappa \rightarrow \infty} \text{err}(\kappa) = 0$.
- (2) *Let $\kappa \geq \kappa_0$, $\epsilon(\kappa)\kappa^2 \leq H \leq \Lambda\kappa^2$ and (ψ, \mathbf{A}) be a critical point of the functional in (1.1).*

(a) If $H \gg \kappa^{3/2}$, then,

$$\mathcal{E}_0(h\psi, \mathbf{A}) \geq \kappa \left(\int_{\Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) + \frac{\kappa^3}{H} \text{err}(\kappa). \quad (7.33)$$

(b) If $H \lesssim \kappa^{3/2}$, then,

$$\mathcal{E}_0(h\psi, \mathbf{A}) \geq \kappa^2 \int_{\Omega} g \left(\frac{H}{\kappa} |B_0(x)| \right) dx + \frac{\kappa^3}{H} \text{err}(\kappa). \quad (7.34)$$

We conclude this section with the

Proof of Theorem 1.1. We have just to combine the conclusions of Theorem 5.2 and Corollary 7.4. \square

8. LOCAL ENERGY ESTIMATES

8.1. Preliminaries. Let $D \subset \Omega$ be an open set with a smooth boundary such that $\partial D \cap \Gamma$ is a finite set. Let $\rho_0 \in (0, 1)$, $\rho \in (0, \rho_0)$ and

$$D_\rho = \{x \in \Omega : \text{dist}(x, D) < \rho\}.$$

We select ρ_0 sufficiently small so that the boundary of ∂D_ρ is smooth. Let $h_1 \in C_c^\infty(D_\rho)$ and $h_2 \in C^\infty(\mathbb{R}^2)$ be functions satisfying

$$0 \leq h_1 \leq 1, \quad |\nabla h_1| + |\nabla h_2| \leq \frac{C}{\rho} \quad \text{in } \mathbb{R}^2, \quad h_1 = 1 \quad \text{in } D_\rho, \quad \text{and } h_1^2 + h_2^2 = 1.$$

Notice that

$$\text{supp } h_2 \subset \overline{D}^c.$$

Let (ψ, \mathbf{A}) be a minimizer of (1.1). We will estimate the following energy

$$\mathcal{E}_0(\psi, \mathbf{A}; D) = \int_D \left(|\nabla - i\kappa H \mathbf{A} \psi|^2 - \kappa^2 |\psi|^2 + \frac{\kappa^2}{2} |\psi|^4 \right) dx.$$

Notice that we have the following decomposition of the energy

$$\mathcal{E}_0(\psi, \mathbf{A}; \Omega) \geq \mathcal{E}_0(h_1 \psi, \mathbf{A}; D_\rho) + \mathcal{E}_0(h_2 \psi, \mathbf{A}; \overline{D}^c) - \frac{C}{\rho^2} \int_{\Omega} |\psi|^2 dx.$$

Now we use the estimate in Proposition 6.4 and write,

$$\mathcal{E}_0(\psi, \mathbf{A}; \Omega) \geq \mathcal{E}_0(h_1 \psi, \mathbf{A}; D_\rho) + \mathcal{E}_0(h_2 \psi, \mathbf{A}; \overline{D}^c) - \frac{C}{\rho^2} \frac{\kappa}{H}. \quad (8.1)$$

Recall that we deal with two separate regimes:

$$\begin{cases} \mathbf{Regime I} : & \kappa \ll H \lesssim \kappa^{3/2}; \\ \mathbf{Regime II} : & \kappa^{3/2} \ll H \lesssim \kappa^2. \end{cases}$$

We define the quantity $C_0(\kappa, H; D)$ as follows:

$$C_0(\kappa, H; D) = \begin{cases} \kappa \left(\int_{D \cap \Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) & \text{in Regime I,} \\ \kappa^2 \int_D g \left(\frac{H}{\kappa} |B_0(x)| \right) dx & \text{in Regime II.} \end{cases} \quad (8.2)$$

Notice that, in Regimes I and II, the result of Theorem 1.1 reads as follows,

$$E_{\text{gs}}(\kappa, H) = C_0(\kappa, H; \Omega) + \frac{\kappa^3}{H} o(1), \quad (\kappa \rightarrow \infty),$$

8.2. Upper bound. The results in this section are valid under the assumption that (ψ, \mathbf{A}) is a minimizer of the functional in (1.1).

We have $\mathcal{E}_0(\psi, \mathbf{A}; \Omega) \leq E_{\text{gs}}(\kappa, H)$. Using $|\psi| \leq 1$ and the upper bound in Theorem 5.2, we get,

$$\mathcal{E}_0(h_1\psi, \mathbf{A}; D_\rho) + \mathcal{E}_0(h_2\psi, \mathbf{A}; \overline{D}^c) \leq C_0(\kappa, H; \Omega) + \frac{\kappa^3}{H} \text{err}(\kappa) + \frac{C}{\rho^2} \frac{\kappa}{H}.$$

Using Theorem 7.3, we may write,

$$\mathcal{E}_0(h_2\psi, \mathbf{A}; D^c) \geq C_0(\kappa, H; \overline{D}^c) + \frac{\kappa^3}{H} \text{err}(\kappa).$$

As a consequence, we have,

$$\mathcal{E}_0(h_1\psi, \mathbf{A}; D_\rho) \leq C_0(\kappa, H; D) + \frac{\kappa^3}{H} \text{err}(\kappa) + \frac{C}{\rho^2} \frac{\kappa}{H}.$$

Since $h_1 = 1$ in D , we get the simple decomposition of the energy,

$$\mathcal{E}_0(\psi, \mathbf{A}; D) = \mathcal{E}_0(h_1\psi, \mathbf{A}; D_\rho) - \mathcal{E}_0(h_1\psi, \mathbf{A}; D_\rho \setminus D).$$

Since $\|h_1\|_\infty \leq 1$ and the boundary of $D_\rho \setminus D$ is smooth, we get in light of Theorem 7.3,

$$\mathcal{E}_0(h_1\psi, \mathbf{A}; D_\rho \setminus D) \geq C_0(\kappa, H; D_\rho \setminus D) + \frac{\kappa^3}{H} \text{err}_\rho(\kappa).$$

In light of the upper bound in Theorem 3.12, we have,

$$\left(\int_{(D_\rho \setminus D) \cap \Gamma} \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right)^{1/3} E \left(|\nabla B_0(x)| \frac{H}{\kappa^2} \right) ds(x) \right) \leq C \frac{\kappa^2}{H} \rho.$$

In the same vein, since $g(b)$ is bounded and vanishes when $b \geq 1$, then,

$$\int_{D_\rho \setminus D} g \left(\frac{H}{\kappa} |B_0(x)| \right) dx \leq C \rho \frac{\kappa}{H}.$$

As a consequence, we get,

$$C_0(\kappa, H; D_\rho \setminus D) \leq C \frac{\kappa^3}{H} \rho,$$

and

$$\mathcal{E}_0(\psi, \mathbf{A}; D) \leq C_0(\kappa, H; D) + \frac{\kappa^3}{H} \left(C \rho + \text{err}_\rho(\kappa) \right) + \frac{C}{\rho^2} \frac{\kappa}{H}$$

Sending κ to infinity, we deduce that,

$$\limsup_{\kappa \rightarrow \infty} \frac{H}{\kappa^3} \{ \mathcal{E}_0(\psi, \mathbf{A}; D_\rho) - C_0(\kappa, H; D) \} \leq C \rho.$$

Next, we send ρ to 0_+ and get,

$$\limsup_{\kappa \rightarrow \infty} \frac{H}{\kappa^3} \{ \mathcal{E}_0(\psi, \mathbf{A}; D) - C_0(\kappa, H; D) \} \leq 0. \quad (8.3)$$

Notice that the upper bound in (8.3) is valid for any open set $D \subset \Omega$ with smooth boundary. In particular, it is true when D is replaced by $\overline{D}^c = \Omega \setminus \overline{D}$, i.e.

$$\limsup_{\kappa \rightarrow \infty} \frac{H}{\kappa^3} \{ \mathcal{E}_0(\psi, \mathbf{A}; \overline{D}^c) - C_0(\kappa, H; \overline{D}^c) \} \leq 0. \quad (8.4)$$

8.3. Lower bound. We continue to assume that (ψ, \mathbf{A}) is a minimizer of the functional in (1.1). We will give a lower bound of the energy $\mathcal{E}_0(\psi, \mathbf{A}; D)$. We plug the lower bound in Corollary 7.4 into the following simple decomposition of the energy,

$$\mathcal{E}_0(\psi, \mathbf{A}; D) + \mathcal{E}_0(\psi, \mathbf{A}; \overline{D}^c) = \mathcal{E}_0(\psi, \mathbf{A}; \Omega).$$

In that way, we get,

$$\mathcal{E}_0(\psi, \mathbf{A}; D) \geq C_0(\kappa, H; \Omega) - \mathcal{E}_0(\psi, \mathbf{A}; \overline{D}^c) + \frac{\kappa^3}{H} \text{err}(\kappa).$$

Notice the following simple decomposition of the term on the right hand side,

$$\mathcal{E}_0(\psi, \mathbf{A}; D) \geq C_0(\kappa, H; D) + \frac{\kappa^3}{H} \text{err}(\kappa) - \{ \mathcal{E}_0(\psi, \mathbf{A}; \overline{D}^c) - C_0(\kappa, H; \overline{D}^c) \}.$$

Now we send κ to ∞ and using (8.4), we get,

$$\liminf_{\kappa \rightarrow \infty} \frac{H}{\kappa^3} \{ \mathcal{E}_0(\psi, \mathbf{A}; D) - C_0(\kappa, H; D) \} \geq 0. \quad (8.5)$$

8.4. Conclusion for the local energy.

Combining (8.3) and (8.5), we get, in the two regimes we are considering, that the local energy in D of a minimizer (ψ, \mathbf{A}) satisfies,

$$\mathcal{E}_0(\psi, \mathbf{A}; D) = C_0(\kappa, H; D) + \frac{\kappa^3}{H} o(\kappa), \quad (8.6)$$

where $C_0(\kappa, H; D)$ is introduced in (8.2).

9. PROOF OF THEOREM 1.5

The proof of (1) in Theorem 1.5 is a simple combination of the upper bound in Theorem 5.2 and the lower bound in Theorem 7.3 (used with $D = \Omega$).

The assertion (2) in Theorem 1.5 is the conclusion of Section 8.4.

The rest of the section is devoted to the proof of statement (3) in Theorem 1.5. This will be done in three steps. Recall the definition of the quantity $C_0(\kappa, H; D)$ in (8.2) and that we work under the assumption on H described in Regimes I and II. It is sufficient to prove that the following formula is true in Regimes I and II,

$$\int_D |\psi(x)|^4 dx = -\frac{2}{\kappa^2} C_0(\kappa, H; D) + \frac{\kappa}{H} o(1), \quad (\kappa \rightarrow \infty),$$

where (ψ, \mathbf{A}) is a minimizer of the energy in (1.1).

Step 1: The case $D = \Omega$.

A minimizer (ψ, \mathbf{A}) satisfies the Ginzburg-Landau equation in (4.1). Recall the useful identity in (4.3),

$$-\frac{\kappa^2}{2} \int_{\Omega} |\psi(x)|^4 dx = \mathcal{E}_0(\psi, \mathbf{A}; \Omega).$$

Thanks to the formulas in Subsection 8.4 used with $D = \Omega$, we observe that (8.6) yields,

$$\int_{\Omega} |\psi(x)|^4 dx = -\frac{2}{\kappa^2} C_0(\kappa, H; \Omega) + \frac{\kappa}{H} o(1), \quad (9.1)$$

where the formula is valid in Regimes I and II.

Step 2: Upper bound.

Let

$$\ell = \kappa^{-1/4} \sqrt{\frac{\kappa}{H}} \quad \text{and} \quad D_\ell = \{x \in D : \text{dist}(x, \partial D) \geq \ell\}.$$

Consider a cut-off function $\chi_\ell \in C_c^\infty(D)$ such that,

$$\|\chi_\ell\|_\infty \leq 1, \quad \|\nabla \chi_\ell\| \leq \frac{C}{\ell}, \quad \chi_\ell = 1 \text{ in } D_\ell.$$

Multiplying both sides of the equation in (4.1) by $\chi_\ell^2 \bar{\psi}$ then integrating by parts and using the estimate in Proposition 6.4 yield,

$$\int_D (|\nabla - i\kappa H \mathbf{A}| \chi_\ell \psi|^2 - \kappa^2 \chi_\ell^2 |\psi|^2 + \kappa^2 \chi_\ell^2 |\psi|^4) dx = \int_D |\nabla \chi_\ell|^2 |\psi|^2 dx = \mathcal{O}\left(\frac{C}{\ell^2} \frac{\kappa}{H}\right) = \frac{\kappa^3}{H} o(1).$$

Since $1 \geq \chi_\ell^2 \geq \chi_\ell^4$, this formula implies,

$$-\frac{\kappa^2}{2} \int_D \chi_\ell^2 |\psi|^4 dx \geq \mathcal{E}_0(\chi_\ell \psi, \mathbf{A}; D) - \frac{\kappa^3}{H} o(1). \quad (9.2)$$

Using the bounds $\|\psi\|_\infty \leq 1$ and $\|\psi\|_2 \leq C \sqrt{\frac{\kappa}{H}}$, the fact that χ_ℓ is supported in D and $\chi_\ell = 1$ in D_ℓ , we get,

$$\begin{aligned} \int_D |\psi(x)|^4 dx &= \int_D \chi_\ell^2(x) |\psi(x)|^4 dx + \int_D (1 - \chi_\ell^2(x)) |\psi(x)|^4 dx \\ &= \int_D \chi_\ell^2(x) |\psi(x)|^4 dx + \mathcal{O}\left(\sqrt{\ell} \sqrt{\frac{\kappa}{H}}\right) \\ &= \int_D \chi_\ell^2(x) |\psi(x)|^4 dx + \frac{\kappa}{H} o(1). \end{aligned} \quad (9.3)$$

Now, we infer from (9.2) and Theorem 7.3,

$$\int_D |\psi(x)|^4 dx \leq -\frac{2}{\kappa^2} C_0(\kappa, H; D) + \frac{\kappa}{H} o(1). \quad (9.4)$$

Step 3: Lower bound.

Notice that (9.4) is valid for any open domain $D \subset \Omega$ with a smooth boundary, in particular, it is valid when D is replaced by the complementary of \bar{D} in Ω : \bar{D}^c . We have the simple decomposition,

$$\begin{aligned} \int_D |\psi(x)|^4 dx &= \int_\Omega |\psi(x)|^4 dx - \int_{\bar{D}^c} |\psi(x)|^4 dx \\ &\geq \int_\Omega |\psi(x)|^4 dx - \frac{2}{\kappa^2} C_0(\kappa, H; \bar{D}^c) + \frac{\kappa}{H} o(1). \end{aligned}$$

Using the asymptotics in (9.1) obtained in Step 1, we deduce that,

$$\int_D |\psi(x)|^4 dx \geq -\frac{2}{\kappa^2} C_0(\kappa, H; D) + \frac{\kappa}{H} o(1).$$

Combining this lower bound and the upper bound in (9.4), we obtain the asymptotics announced in the third assertion of Theorem 1.5.

Acknowledgements. The authors would like to thank S. Fournais and N. Raymond for useful discussions. B. Helffer is partially supported by the ANR program NOSEVOL. A. Kachmar is partially supported by a grant from Lebanese University.

REFERENCES

- [1] K. Attar. The ground state energy of the two dimensional Ginzburg-Landau functional with variable magnetic field. To appear in *Annales de l'institut Henri Poincaré - Analyse Non-Linéaire*.
- [2] K. Attar. Energy and vorticity of the Ginzburg-Landau model with variable magnetic field. In preparation.
- [3] M. Correggi, N. Rougerie. On the Ginzburg-Landau functional in the surface superconductivity regime. [arXiv:1309.2268](https://arxiv.org/abs/1309.2268) (2013).
- [4] S. Fournais, B. Helffer. *Spectral Methods in Surface Superconductivity*. Progress in Nonlinear Differential Equations and Their Applications. **77** Birkhäuser (2010).
- [5] S. Fournais, B. Helffer. Bulk superconductivity in type II superconductors near the second critical field. *J. Europ. Math. Soc.* **12** 461470 (2008).
- [6] S. Fournais, B. Helffer. Strong diamagnetism for general domains and application. *Ann. Inst. Fourier* **57** (7) 2389-2400 (2007).
- [7] S. Fournais, B. Helffer. Optimal uniform elliptic estimates for the Ginzburg-Landau system. *Adventures in Mathematical Physics. Contemp. Math.* **447** (2007), 83-102.
- [8] S. Fournais, A. Kachmar. The ground state energy of the three dimensional Ginzburg-Landau functional. Part I. Bulk regime. *Communications in Partial Differential Equations.* **38** 339-383 (2013).
- [9] S. Fournais, A. Kachmar. Nucleation of bulk superconductivity close to critical magnetic field. *Advances in Mathematics* **226** 1213-1258 (2011).
- [10] S. Fournais, A. Kachmar, and M. Persson. The ground state energy of the three dimensional Ginzburg-Landau functional. Part II. Surface regime. *J. Math. Pures Appl.* **99** 343-374 (2013).
- [11] P.G. de Gennes. Boundary effects in superconductors. *Rev. Mod. Phys.* January 1964.
- [12] B. Helffer. The Montgomery operator revisited. *Colloquium Mathematicum*, Vol. 118, No 2, 391-400 (2011).
- [13] B. Helffer, Yu. A. Kordyukov. Semiclassical spectral asymptotics for a two-dimensional magnetic Schrödinger operator: The case of discrete wells, *Spectral Theory and Geometric Analysis. Contemp. Math.* **535**, 55-78; AMS, Providence, RI, 2011.
- [14] B. Helffer, Yu. A. Kordyukov. Accurate semiclassical spectral asymptotics for a two-dimensional magnetic Schrödinger operator, to appear in *Ann. Henri Poincaré* (2014).
- [15] B. Helffer, A. Mohamed. Semiclassical analysis for the ground state energy of a Schrödinger operator with magnetic wells. *J. Funct. Anal.* **138** (1) (1996), 40-81.
- [16] K. Lu, X.B. Pan. Estimates of the upper critical field for the Ginzburg-Landau equations of superconductivity. *Physica D* **127** (1999), 73-104.
- [17] K. Lu, X.B. Pan. Eigenvalue problems of Ginzburg-Landau operator in bounded domains. *Journal of mathematical Physics*, Vol. 40 (6), 2647-2670 (1999).
- [18] J-P. Miqueu. Equation de Schrödinger avec un champ magnétique qui s'annule. Thèse de doctorat (in preparation).
- [19] R. Montgomery. Hearing the zero locus of a magnetic field. *Commun. Math. Phys.* **168** (3) 651-675 (1995).
- [20] X.B. Pan, K.H. Kwek. Schrödinger operators with non-degenerately vanishing magnetic fields in bounded domains. *Trans. Amer. Math. Soc.* **354** (10) 4201-4227 (2002).
- [21] X.B. Pan. Surface superconductivity in applied magnetic fields above H_{C_2} . *Commun. Math. Phys.* **228** (2002), 228-370.
- [22] N. Raymond. Sharp asymptotics for the Neumann Laplacian with variable magnetic field: case of dimension 2. *Ann. Henri Poincaré* **10**(1) (2009), 95-122.
- [23] N. Raymond, S. Vu Ngoc. *Geometry and spectrum in 2D magnetic wells*. Preprint [arXiv:1306.5054](https://arxiv.org/abs/1306.5054), 2013. To appear in *Annales Institut Fourier* (2014).
- [24] E. Sandier, S. Serfaty. *Vortices for the Magnetic Ginzburg-Landau Model*. Progress in Nonlinear Differential Equations and their Applications. **70** Birkhäuser (2007).
- [25] E. Sandier, S. Serfaty. The decrease of bulk superconductivity close to the second critical field in the Ginzburg-Landau model. *SIAM. J. Math. Anal.* **34** No. 4 (2003), 939-956.

(B. Helffer) LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ DE PARIS-SUD 11, BÂT 425, 91405 ORSAY, FRANCE AND LABORATOIRE JEAN LERAY (UNIVERSITÉ DE NANTES)

E-mail address: bernard.helffer@math.u-psud.fr

(A. Kachmar) DEPARTMENT OF MATHEMATICS, LEBANESE UNIVERSITY, HADAT, LEBANON; SCHOOL OF ARTS AND SCIENCES, LEBANESE INTERNATIONAL UNIVERSITY, BEIRUT, LEBANON

E-mail address: ayman.kashmar@liu.edu.lb