

# Multibump nodal solutions for an indefinite superlinear elliptic problem\*

Pedro M. Girão<sup>†</sup> and José Maria Gomes<sup>‡</sup>  
 Instituto Superior Técnico  
 Av. Rovisco Pais  
 1049-001 Lisbon, Portugal

## Abstract

We define some Nehari-type constraints using an orthogonal decomposition of the Sobolev space  $H_0^1$  and prove the existence of multibump nodal solutions for an indefinite superlinear elliptic problem.

## 1 Introduction

Consider a Lipschitz bounded domain  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 1$ , and a function  $a \in C(\bar{\Omega})$ , with  $a = a^+ - a^-$ , where  $a^+ = \max\{a, 0\}$  as usual. Assume the set  $a^+ > 0$  is the union of a finite number,  $L \geq 1$ , of open connected and disjoint Lipschitz components. We separate the components arbitrarily into three families

$$\begin{aligned}\Omega^+ = \{x \in \Omega : a^+(x) > 0\} &= (\cup_{i=1}^I \tilde{\omega}_i) \cup (\cup_{j=1}^J \hat{\omega}_j) \cup (\cup_{k=1}^K \bar{\omega}_k) \\ &= \tilde{\Omega} \cup \hat{\Omega} \cup \underline{\Omega},\end{aligned}$$

so that  $L = I + J + K$ ; we also assume

$$\Omega^- = \{x \in \Omega : a^-(x) > 0\} = \Omega \setminus \overline{\Omega^+}.$$

Let  $\mu > 0$  and  $p$  be a superquadratic and subcritical exponent,  $2 < p < 2^*$ , with  $2^* = 2N/(N-2)$  for  $N \geq 3$ , and  $2^* = +\infty$  for  $N = 1$  or  $2$ . Our main result is

**Theorem 1.1.** *For every large  $\mu$ , there exists an  $H_0^1(\Omega)$  weak solution  $u_\mu$  of*

$$-\Delta u = (a^+ - \mu a^-)|u|^{p-2}u \quad \text{in } \Omega. \quad (1)$$

---

\*2000 Mathematics Subject Classification: 35J65 (35J20)

Keywords: Multibump solutions, Nehari manifold, sign-changing solutions, elliptic equations

<sup>†</sup>Email: pgirao@math.ist.utl.pt. Partially supported by the Center for Mathematical Analysis, Geometry and Dynamical Systems through FCT Program POCTI/FEDER and by grant POCI/FEDER/MAT/55745/2004.

<sup>‡</sup>Email: jgomes@math.ist.utl.pt. Supported by FCT grant SFRH/BPD/29098/2006.

Furthermore, the family  $\{u_\mu\}$  has the property that (modulo a subsequence)

$$u_\mu \rightharpoonup u \quad \text{in } H_0^1(\Omega) \text{ as } \mu \rightarrow +\infty, \quad (2)$$

where

$$\begin{cases} -\Delta u = a^+|u|^{p-2}u & \text{in } \tilde{\omega}_i, \\ u^\pm \not\equiv 0 & \text{in } \tilde{\omega}_i, \end{cases} \quad i = 1, \dots, I,$$

$$\begin{cases} -\Delta u = a^+|u|^{p-2}u & \text{in } \hat{\omega}_j, \\ u^+ \not\equiv 0, u^- \equiv 0 & \text{in } \hat{\omega}_j, \end{cases} \quad j = 1, \dots, J,$$

$$u \equiv 0 \quad \text{in } \bar{\omega}_k, \quad k = 1, \dots, K,$$

and

$$u \equiv 0 \quad \text{in } \Omega^-.$$

The one-dimensional version of (1) was studied in [15] with topological shooting arguments and phase-plane analysis. Theorem 1.1 extends the main result in [7] where the case  $\tilde{\Omega} = \emptyset$  was considered, so that the function  $u$  in (2) was positive. The authors used a volume constrain regarding the  $L^p$  norm, rescaling and a min-max argument based on the Mountain Pass Lemma. A careful analysis allowed them to distinguish between the solutions that arise from the  $2^L$  different possible partitionings of  $\Omega^+ = \hat{\Omega} \cup \underline{\Omega}$ . However, the argument in [7] does not seem either to extend easily to the present situation or to be suited to non-homogeneous nonlinearities.

Our approach is adapted from the work [18] regarding a system of equations related to

$$\begin{cases} -\epsilon^2 \Delta u + V(x)u = f(u) & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \end{cases}$$

when  $\epsilon$  is small and the functions  $V$  and  $f$  satisfy appropriate conditions. The positive function  $V$  was assumed to have a finite number of minima. In particular, the authors proved the existence of multipeak positive solutions by defining a Nehari-type manifold which, roughly speaking, imposes that the derivative of the associated Euler-Lagrange functional at a function  $u$  should vanish when applied to a truncation of  $u$  around a minimum of the potential function  $V$ .

The perspective of [18] is related to the one of [16] which, using Nehari conditions and a cut-off operator, simplifies the original techniques for gluing together mountain-pass type solutions of [12], [13] and [20].

Our method consists in defining a Nehari-type set,  $\mathcal{N}_\mu$ , by imposing that the derivative of the associated Euler-Lagrange functional at a function  $u$  should vanish when applied to the positive and negative parts of some projections of  $u$ . The idea to use these projections is borrowed from [7], where they are also used, but in a different way.

We prove that the Euler-Lagrange functional associated to (1) has a minimum over the set  $\mathcal{N}_\mu$  using an argument similar to the one found in [8]. Since our set  $\mathcal{N}_\mu$  is not a manifold (see [5, Lemma 3.1]), one has to demonstrate, as in [9], that the minima are indeed critical points. As mentioned above, in the case that  $\tilde{\Omega} = \emptyset$  we recover the main result of [7], but with a simpler proof.

Our results are somewhat parallel to the ones of singular perturbation problems like in [14]. The large parameter  $\mu$  in (1) plays the role of the small parameter  $\epsilon$ . The solutions concentrate in the set  $\tilde{\Omega} \cup \hat{\Omega}$  and vanish in the set  $\underline{\Omega} \cup \Omega^-$  as  $\mu \rightarrow +\infty$ .

In [1] flow invariance properties together with a weak splitting condition proved the existence of infinitely many geometrically distinct two bump solutions of a periodic superlinear Schrödinger equation. The paper [4] is concerned with the singular perturbed equation above. As a special case, the authors observed the existence of multiple pairs of concentrating nodal solutions at an isolated minimum of the potential.

There has been much interest in elliptic problems with a sign changing weight. We refer to [2], [3], [6], [11], [17], [19], [21] and the references therein.

For simplicity we restrict the proof to the case where  $I = J = K = 1$ , but it extends to the other ones as well. The work is organized as follows. In Section 2 we provide estimates for minimizing sequences on the set  $\mathcal{N}_\mu$ . In Section 3 we prove the existence of a minimizer in the set  $\mathcal{N}_\mu$ . Finally, in Section 4 we prove that a minimizer in the set  $\mathcal{N}_\mu$  is a critical point using a local deformation and a degree argument similar to the one in [10].

## 2 Estimates for minimizing sequences on a Nehari-type set $\mathcal{N}_\mu$

As mentioned in the Introduction, we consider a Lipschitz bounded domain  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 1$ , and a function  $a \in C(\Omega)$ . We assume the set  $a^+ > 0$  is the union of three Lipschitz components,

$$\{x \in \Omega : a^+(x) > 0\} = \tilde{\omega} \cup \hat{\omega} \cup \bar{\omega},$$

and

$$\{x \in \Omega : a^-(x) > 0\} = \Omega \setminus \overline{(\tilde{\omega} \cup \hat{\omega} \cup \bar{\omega})}. \quad (3)$$

We introduce a positive parameter  $\mu$  and consider  $2 < p < 2^*$ .

We denote by  $\langle \cdot, \cdot \rangle$  the usual inner product on the Sobolev space  $H_0^1(\Omega)$ , i.e.  $\langle u, v \rangle = \int \nabla u \cdot \nabla v$  for  $u, v \in H_0^1(\Omega)$ . When the region of integration is

not specified it is understood that the integrals are over  $\Omega$ . We denote by  $\| \cdot \|$  the induced norm. We define the spaces

$$\begin{aligned}\underline{H}(\tilde{\omega}) &= \{u \in H_0^1(\Omega) : u = 0 \text{ in } \Omega \setminus \tilde{\omega}\}, \\ \underline{H}(\hat{\omega}) &= \{u \in H_0^1(\Omega) : u = 0 \text{ in } \Omega \setminus \hat{\omega}\}, \\ \underline{H}(\bar{\omega}) &= \{u \in H_0^1(\Omega) : u = 0 \text{ in } \Omega \setminus \bar{\omega}\},\end{aligned}$$

which can be obtained from the spaces  $H_0^1(\tilde{\omega})$ ,  $H_0^1(\hat{\omega})$ ,  $H_0^1(\bar{\omega})$  by extending functions as zero on  $\Omega \setminus \tilde{\omega}$ ,  $\Omega \setminus \hat{\omega}$ ,  $\Omega \setminus \bar{\omega}$ , respectively.

Each  $u \in H_0^1(\Omega)$  can be decomposed as

$$u = \tilde{u} + \hat{u} + \bar{u} + \underline{u},$$

with  $\tilde{u}$ ,  $\hat{u}$  and  $\bar{u}$  the projections of  $u$  on  $\underline{H}(\tilde{\omega})$ ,  $\underline{H}(\hat{\omega})$  and  $\underline{H}(\bar{\omega})$ , respectively. We recall the projections are defined by

$$\begin{aligned}\tilde{u} \in \underline{H}(\tilde{\omega}) &: \forall \varphi \in \underline{H}(\tilde{\omega}), \quad \langle u, \varphi \rangle = \langle \tilde{u}, \varphi \rangle, \\ \hat{u} \in \underline{H}(\hat{\omega}) &: \forall \varphi \in \underline{H}(\hat{\omega}), \quad \langle u, \varphi \rangle = \langle \hat{u}, \varphi \rangle, \\ \bar{u} \in \underline{H}(\bar{\omega}) &: \forall \varphi \in \underline{H}(\bar{\omega}), \quad \langle u, \varphi \rangle = \langle \bar{u}, \varphi \rangle.\end{aligned}$$

Clearly, these projections are orthogonal and continuous with respect to the weak topology. The function  $\underline{u}$  is harmonic in  $\tilde{\omega} \cup \hat{\omega} \cup \bar{\omega}$ .

The following is Theorem 1.1 in the case when  $I = J = K = 1$ .

**Proposition 2.1.** *For every large  $\mu$ , there exists an  $H_0^1(\Omega)$  weak solution  $u_\mu$  of*

$$-\Delta u = (a^+ - \mu a^-)|u|^{p-2}u \quad \text{in } \Omega. \quad (4)$$

Furthermore, the family  $\{u_\mu\}$  has the property that, modulo a subsequence,

$$u_\mu \rightharpoonup u \quad \text{in } H_0^1(\Omega) \text{ as } \mu \rightarrow +\infty, \quad (5)$$

where

$$u = \tilde{u} + \hat{u}, \quad (6)$$

$$\begin{cases} -\Delta \tilde{u} = a^+|\tilde{u}|^{p-2}\tilde{u} & \text{in } \tilde{\omega}, \\ \tilde{u}^\pm \neq 0, \end{cases} \quad (7)$$

and

$$\begin{cases} -\Delta \hat{u} = a^+|\hat{u}|^{p-2}\hat{u} & \text{in } \hat{\omega}, \\ \hat{u}^+ \neq 0, \hat{u}^- \equiv 0. \end{cases} \quad (8)$$

The solutions of (4) are the critical points of the  $C^2$  functional  $I_\mu : H_0^1(\Omega) \rightarrow \mathbb{R}$ , defined by

$$I_\mu(u) = \frac{1}{2} \|u\|^2 - \frac{1}{p} \int (a^+ - \mu a^-) |u|^p.$$

We fix a function  $v$  such that  $v = \tilde{v} + \hat{v}^+$ , with  $\tilde{v}^+, \tilde{v}^-, \hat{v} \not\equiv 0$  and

$$I'_\mu(v)(\tilde{v}^+) = I'_\mu(v)(\tilde{v}^-) = I'_\mu(v)(\hat{v}) = 0$$

for some (and hence all)  $\mu > 0$ .

The restriction of  $I_\mu$  to  $\underline{H}(\hat{\omega}) \oplus \underline{H}(\bar{\omega})$  is independent of  $\mu$  and has a strict local minimum at zero. We fix a small  $\rho_0 > 0$  such that zero is the unique minimizer of  $I_\mu$  in  $\{u \in \underline{H}(\hat{\omega}) \oplus \underline{H}(\bar{\omega}) : \max\{\|\hat{u}\|, \|\bar{u}\|\} \leq \rho_0\}$ . For  $0 < \rho \leq \rho_0$ , we denote by  $c_\rho$  the positive constant

$$c_\rho := \inf_{\substack{u \in \underline{H}(\hat{\omega}) \oplus \underline{H}(\bar{\omega}) \\ \rho \leq \max\{\|\hat{u}\|, \|\bar{u}\|\} \leq \rho_0}} I_\mu(u). \quad (9)$$

The solutions of (4) will be obtained by minimizing the functional  $I_\mu$  on the following Nehari-type set,  $\mathcal{N}_\mu$ . Let  $\rho_0$  be as above and  $R > \|v\|$ .

**Definition 2.2.**  $\mathcal{N}_\mu$  is the set of functions  $u = \tilde{u} + \hat{u} + \bar{u} + \underline{u} \in H_0^1(\Omega)$  such that

- ( $\mathcal{N}_i$ )  $\tilde{u}^+, \tilde{u}^-, \hat{u}^+ \not\equiv 0$ ,
- ( $\mathcal{N}_{ii}$ )  $I'_\mu(u)(\tilde{u}^+) = I'_\mu(u)(\tilde{u}^-) = I'_\mu(u)(\hat{u}^+) = 0$ ,
- ( $\mathcal{N}_{iii}$ )  $I_\mu(u) \leq I_\mu(v) + 1$ ,
- ( $\mathcal{N}_{iv}$ )  $\|\underline{u}\| \leq \min\{\|\tilde{u}^+\|, \|\tilde{u}^-\|, \|\hat{u}^+\|\} < \|\tilde{u} + \hat{u}^+\| \leq R$ ,
- ( $\mathcal{N}_v$ )  $\max\{\|\hat{u}^-\|, \|\bar{u}\|\} \leq \rho_0$ .

We remark that  $v \in \mathcal{N}_\mu$  for all  $\mu > 0$ .

The square of the  $H_0^1(\Omega)$  norm of  $u$  is equal to the sum of the squares of the  $H_0^1(\Omega)$  norms of the components of  $u$ , but the  $p$ -th power of the  $L^p(\Omega)$  norm of  $u$  does not have such a nice property. However, the next lemma says that this is almost the case when  $\mu$  is large.

**Lemma 2.3.** *Let  $\delta > 0$  be given. There exists  $\mu_\delta$  such that, if  $\mu > \mu_\delta$ ,*

$$\forall u \in \mathcal{N}_\mu, \quad \int |\underline{u}|^p < \delta.$$

*Proof.* Suppose, by contradiction, that for some  $\delta > 0$  there exists  $\mu_n \rightarrow +\infty$  and  $u_n \in \mathcal{N}_{\mu_n}$  with

$$\int |\underline{u}_n|^p \geq \delta. \quad (10)$$

As  $\|u_n\|$  is bounded, we may suppose  $u_n \rightharpoonup u$ . We have  $\underline{u}_n \rightharpoonup \underline{u}$  and  $\underline{u} \equiv 0$  in  $\Omega \setminus (\tilde{\omega} \cup \hat{\omega} \cup \bar{\omega})$ . Otherwise, by (3) and modulo a subsequence,

$$\int a^- |\underline{u}_n|^p \geq c > 0.$$

This would contradict  $(\mathcal{N}_{iii})$  for sufficiently large  $n$ :

$$\frac{1}{2} \|u_n\|^2 - \frac{1}{p} \int a^+ |u_n|^p + \frac{\mu_n}{p} \int a^- |\underline{u}_n|^p \leq I_\mu(v) + 1.$$

So the function  $\underline{u}$  belongs to  $\underline{H}(\tilde{\omega}) \oplus \underline{H}(\hat{\omega}) \oplus \underline{H}(\bar{\omega})$  and is harmonic in  $\tilde{\omega} \cup \hat{\omega} \cup \bar{\omega}$ . It follows that  $\underline{u}$  must be identically equal to zero in  $\Omega$ . This contradicts (10).  $\square$

Usually one may obtain a lower bound for the  $H_0^1(\Omega)$  norm of  $\tilde{u}^+$ ,  $\tilde{u}^-$  and  $\hat{u}^+$  from  $(\mathcal{N}_i)$  and a condition like  $(\mathcal{N}_{ii})$ . Here, in addition, we require the first inequality in  $(\mathcal{N}_{iv})$  to prove

**Lemma 2.4.** *There exists a constant  $\kappa$ , independent of  $\mu$ , such that*

$$\forall u \in \mathcal{N}_\mu, \quad \min \{ \|\tilde{u}^+\|, \|\tilde{u}^-\|, \|\hat{u}^+\| \} \geq \kappa > 0. \quad (11)$$

*Proof.* Let  $w$  be one of the three functions  $\tilde{u}^+$ ,  $-\tilde{u}^-$  or  $\hat{u}^+$ . Denote by  $\chi$  the characteristic function of the set  $\{x \in \Omega : w(x) \neq 0\}$  and let  $c$  be the Sobolev constant  $(\int |v|^p)^{1/p} \leq c \|v\|, \forall v \in H_0^1(\Omega)$ . From  $I'_\mu(u)w = 0$ ,

$$\begin{aligned} \|w\|^2 &= \int a^+ |u|^{p-2} u w \leq \|a\|_\infty \left( \int \chi |u|^p \right)^{\frac{p-1}{p}} \left( \int |w|^p \right)^{\frac{1}{p}} \\ &\leq \|a\|_\infty c^p (\|\underline{u}\| + \|w\|)^{p-1} \|w\| \leq 2^{p-1} \|a\|_\infty c^p \|w\|^p, \end{aligned}$$

because of the first inequality in  $(\mathcal{N}_{iv})$ . Since  $w \neq 0$ , due to  $(\mathcal{N}_i)$ , we may take

$$\kappa = (2^{p-1} \|a\|_\infty c^p)^{-1/(p-2)}.$$

$\square$

Now we fix a  $\mu$  and turn to minimizing sequences  $(u_n)$  for  $I_\mu$  restricted to  $\mathcal{N}_\mu$ . Later it will be important that the limit of such a sequence has a neighborhood whose points satisfy  $(\mathcal{N}_i)$ ,  $(\mathcal{N}_{iii})$ ,  $(\mathcal{N}_{iv})$  and  $(\mathcal{N}_v)$ . This follows from

**Lemma 2.5.** *Let  $\bar{R}$  be fixed,  $\|v\| < \bar{R} < R$ , and  $\delta$  be given,  $0 < \delta < \rho_0$ . There exists  $\mu_\delta > 0$  such that for every  $\mu > \mu_\delta$  and every minimizing sequence  $(u_n)$  for  $I_\mu$  restricted to  $\mathcal{N}_\mu$ , we have, for large  $n$ ,*

- (a)  $I_\mu(u_n) \leq I_\mu(v) + \frac{1}{2}$ ,
- (b)  $\|\tilde{u}_n + \hat{u}_n^+\| < \bar{R}$ ,
- (c)  $\max\{\|\hat{u}_n^-\|, \|\bar{u}_n\|\} < \delta$ ,
- (d)  $\|\underline{u}_n\| < \delta$ ;

also

- (e)  $\frac{\mu}{p} \int a^- |\underline{u}_n|^p < \delta$ .

*Proof.* (a) Immediate since  $(u_n)$  is minimizing and  $v \in \mathcal{N}_\mu$  for all  $\mu$ .

(b) Suppose

$$\|\tilde{u}_n + \hat{u}_n^+\| \geq \bar{R} \quad (12)$$

for large  $n$ .

$$\begin{aligned} I_\mu(u_n) &= \frac{1}{2} \|\tilde{u}_n + \hat{u}_n^+\|^2 + \frac{1}{2} \|\hat{u}_n^-\|^2 + \frac{1}{2} \|\bar{u}_n\|^2 + \frac{1}{2} \|\underline{u}_n\|^2 \\ &\quad - \frac{1}{p} \int a^+ |u_n|^{p-2} u_n (\tilde{u}_n + \hat{u}_n^+) + \frac{1}{p} \int a^+ |u_n|^{p-2} u_n \hat{u}_n^- \\ &\quad - \frac{1}{p} \int a^+ |u_n|^{p-2} u_n \bar{u}_n - \frac{1}{p} \int a^+ |u_n|^{p-2} u_n \underline{u}_n + \frac{\mu}{p} \int a^- |\underline{u}_n|^p \\ &\geq \left( \frac{1}{2} - \frac{1}{p} \right) \|\tilde{u}_n + \hat{u}_n^+\|^2 + o(1). \end{aligned}$$

Here and henceforth  $o(1)$  denotes a value, independent of  $u \in \mathcal{N}_\mu$ , that can be made arbitrarily small by choosing  $\mu$  sufficiently large. For the proof of the last inequality we used  $(\mathcal{N}_{ii})$ ,

$$\frac{1}{2} \|\hat{u}_n^-\|^2 + \frac{1}{p} \int a^+ |u_n|^{p-2} u_n \hat{u}_n^- \geq o(1)$$

and

$$\frac{1}{2} \|\bar{u}_n\|^2 - \frac{1}{p} \int a^+ |u_n|^{p-2} u_n \bar{u}_n \geq o(1)$$

(consequences of  $(\mathcal{N}_v)$  and Lemma 2.3),

$$-\frac{1}{p} \int a^+ |u_n|^{p-2} u_n \underline{u}_n = o(1)$$

(consequence of  $(\mathcal{N}_{iv})$ ,  $(\mathcal{N}_v)$  and Lemma 2.3), and

$$\frac{1}{2} \|\underline{u}_n\|^2 + \frac{\mu}{p} \int a^- |\underline{u}_n|^p \geq 0.$$

We now use (12) and the definition of  $\bar{R}$ . For sufficiently large  $\mu$ ,

$$I_\mu(u_n) \geq \left(\frac{1}{2} - \frac{1}{p}\right) \bar{R}^2 + o(1) > \left(\frac{1}{2} - \frac{1}{p}\right) \|v\|^2 + c = I_\mu(v) + c,$$

for some  $c > 0$ . This contradicts the fact that  $(u_n)$  is minimizing.

(c) Suppose  $\|\hat{u}_n^-\| \geq \delta$  for large  $n$ . As in (b), we have

$$\begin{aligned} I_\mu(u_n) &= I_\mu(u_n + \hat{u}_n^-) + \frac{1}{2} \|\hat{u}_n^-\|^2 - \frac{1}{p} \int a^- |\hat{u}_n^-|^p + o(1) \\ &\geq I_\mu(u_n + \hat{u}_n^-) + c_\delta + o(1), \end{aligned}$$

due to Lemma 2.3 and then (9). This implies that

$$\lim I_\mu(u_n) > \liminf I_\mu(u_n + \hat{u}_n^-),$$

for sufficiently large  $\mu$ , and contradicts the assumption that  $(u_n)$  is minimizing, because  $u_n + \hat{u}_n^- \in \mathcal{N}_\mu$ . Similarly, one proves that  $\|\bar{u}_n\| \geq \delta$  for large  $n$  leads to a contradiction, for sufficiently large  $\mu$ , because  $u_n - \bar{u}_n \in \mathcal{N}_\mu$ .

(d) Suppose  $\|\underline{u}_n\| \geq \delta$  for large  $n$ . From  $(\mathcal{N}_{ii})$  and Lemma 2.3, we know

$$\begin{aligned} \|\tilde{u}_n^+\|^2 &= \int a^+ |\tilde{u}_n^+|^p + o(1), \\ \|\tilde{u}_n^-\|^2 &= \int a^+ |\tilde{u}_n^-|^p + o(1), \\ \|\hat{u}_n^+\|^2 &= \int a^+ |\hat{u}_n^+|^p + o(1). \end{aligned}$$

We define  $\tilde{r}_n$ ,  $\tilde{s}_n$  and  $\hat{t}_n$  by

$$\tilde{r}_n = \left( \frac{\|\tilde{u}_n^+\|^2}{\int a^+ |\tilde{u}_n^+|^p} \right)^{\frac{1}{p-2}}, \quad \tilde{s}_n = \left( \frac{\|\tilde{u}_n^-\|^2}{\int a^+ |\tilde{u}_n^-|^p} \right)^{\frac{1}{p-2}}, \quad \hat{t}_n = \left( \frac{\|\hat{u}_n^+\|^2}{\int a^+ |\hat{u}_n^+|^p} \right)^{\frac{1}{p-2}},$$

so that  $\tilde{r}_n, \tilde{s}_n, \hat{t}_n = 1 + o(1)$  by Lemma 2.4, and

$$v_n := \tilde{r}_n \tilde{u}_n^+ - \tilde{s}_n \tilde{u}_n^- + \hat{t}_n \hat{u}_n^+ - \hat{u}_n^- + \bar{u}_n.$$

Provided  $\mu$  is large, we can guarantee  $v_n \in \mathcal{N}_\mu$  for large  $n$  due to (a), (b), (c) and Lemma 2.4. We now obtain an upper bound for  $I_\mu(v_n)$ :

$$\begin{aligned}
I_\mu(v_n) &= I_\mu(\tilde{u}_n + \hat{u}_n + \bar{u}_n) + o(1) \\
&\leq I_\mu(u_n) + o(1) \\
&\quad - \left( \frac{1}{2} \|\underline{u}_n\|^2 - \frac{1}{p} \int a^+ (|u_n|^p - |u_n - \underline{u}_n|^p) + \frac{\mu}{p} \int a^- |\underline{u}_n|^p \right) \\
&\leq I_\mu(u_n) + o(1) - \frac{1}{2} \|\underline{u}_n\|^2 \\
&\leq I_\mu(u_n) + o(1) - \frac{1}{2} \delta^2.
\end{aligned} \tag{13}$$

This implies that  $\liminf I_\mu(v_n) < \lim I_\mu(u_n)$  for sufficiently large  $\mu$ , which is impossible.

(e) Follows from inequality (13).  $\square$

### 3 Existence of a minimizer in $\mathcal{N}_\mu$

For each  $u \in \mathcal{N}_\mu$ , we consider the 3-dimensional manifold with boundary in  $H_0^1(\Omega)$  parametrized on  $[0, 2]^3$  by

$$\zeta(\tilde{r}, \tilde{s}, \hat{t}) = \tilde{r}\tilde{u}^+ - \tilde{s}\tilde{u}^- + \hat{t}\hat{u}^+ - \hat{u}^- + \bar{u} + \underline{u}. \tag{14}$$

We call  $f$  the function  $I_\mu \circ \zeta$ , so that

$$\begin{aligned}
f(\tilde{r}, \tilde{s}, \hat{t}) &= \frac{\tilde{r}^2}{2} \|\tilde{u}^+\|^2 + \frac{\tilde{s}^2}{2} \|\tilde{u}^-\|^2 + \frac{\hat{t}^2}{2} \|\hat{u}^+\|^2 + K \\
&\quad - \frac{1}{p} \int a^+ |\tilde{r}\tilde{u}^+ + \underline{u}|^p - \frac{1}{p} \int a^+ |\underline{u} - \tilde{s}\tilde{u}^-|^p - \frac{1}{p} \int a^+ |\hat{t}\hat{u}^+ + \underline{u}|^p,
\end{aligned}$$

with

$$\begin{aligned}
K &= \frac{1}{2} \|\hat{u}^-\|^2 + \frac{1}{2} \|\bar{u}\|^2 + \frac{1}{2} \|\underline{u}\|^2 \\
&\quad - \frac{1}{p} \int a^+ |\underline{u} - \hat{u}^-|^p - \frac{1}{p} \int a^+ |\bar{u} + \underline{u}|^p + \frac{\mu}{p} \int a^- |\underline{u}|^p.
\end{aligned}$$

Two properties of  $f$  are immediate, namely  $f(1, 1, 1) = I_\mu(u)$  and  $\nabla f(1, 1, 1) = 0$  by  $(\mathcal{N}_{ii})$ . The critical point  $(1, 1, 1)$  is characterized in

**Lemma 3.1.** *For  $\mu$  sufficiently large, independent of  $u \in \mathcal{N}_\mu$ , the point  $(1, 1, 1)$  is an absolute maximum of  $f$ . Furthermore, if*

$$|(\tilde{r}, \tilde{s}, \hat{t}) - (1, 1, 1)| \geq \theta > 0,$$

then

$$f(\tilde{r}, \tilde{s}, \hat{t}) \leq f(1, 1, 1) - d_\theta. \quad (15)$$

The constant  $d_\theta > 0$  may be chosen independent of  $u$  and  $\mu$ .

*Proof.* We define an auxiliary function  $g : [0, 2]^3 \rightarrow \mathbb{R}$  by

$$\begin{aligned} g(\tilde{r}, \tilde{s}, \hat{t}) &:= \left( \frac{\tilde{r}^2}{2} - \frac{\tilde{r}^p}{p} \right) \|\tilde{u}^+\|^2 + \left( \frac{\tilde{s}^2}{2} - \frac{\tilde{s}^p}{p} \right) \|\tilde{u}^-\|^2 \\ &\quad + \left( \frac{\hat{t}^2}{2} - \frac{\hat{t}^p}{p} \right) \|\hat{u}^+\|^2 + K, \end{aligned}$$

which satisfies  $\nabla g(1, 1, 1) = 0$  and

$$D^2g(1, 1, 1) = -(p-2) \operatorname{diag} \left\{ \|\tilde{u}^+\|^2, \|\tilde{u}^-\|^2, \|\hat{u}^+\|^2 \right\} \leq -(p-2)\kappa I,$$

where  $\kappa$  was defined in Lemma 2.4. One easily checks that in a small neighborhood of  $(1, 1, 1)$  the second derivative  $D^2g$  is below a negative definite matrix which is independent of  $u \in \mathcal{N}_\mu$ . We also have that, for any derivative  $D^\alpha$  with  $|\alpha| \leq 2$ ,

$$|D^\alpha f - D^\alpha g| = o(1), \quad (16)$$

by Lemma 2.3; notice that the right-hand-side is uniform in  $u$  and  $\mu$ . Thus, by (16) with  $|\alpha| = 2$ ,  $f$  has a strict local maximum at  $(1, 1, 1)$ . We take  $\alpha = 0$  to conclude this maximum is absolute. Of course, the previous two statements hold provided  $\mu$  is sufficiently large.  $\square$

Let  $\mu$  be fixed and  $(u_n)$  be a minimizing sequence for  $I_\mu$  restricted to  $\mathcal{N}_\mu$ . Since  $\mathcal{N}_\mu$  is bounded in  $H_0^1(\Omega)$ , we may assume

$$u_n \rightharpoonup u \quad \text{in } H_0^1(\Omega).$$

**Lemma 3.2.** *If  $\mu$  is sufficiently large, the function  $u$  belongs to  $\mathcal{N}_\mu$ . Therefore (by the lower semi-continuity of the norm) the function  $u$  is a minimizer of  $I_\mu$  restricted to  $\mathcal{N}_\mu$ .*

*Proof.* We may assume  $\tilde{u}_n^+ \rightharpoonup \tilde{u}^+$ ,  $\tilde{u}_n^- \rightharpoonup \tilde{u}^-$ ,  $\hat{u}_n^+ \rightharpoonup \hat{u}^+$  in  $H_0^1(\Omega)$ , since  $w_n \rightharpoonup w$  in  $H_0^1(\Omega)$  implies a subsequence of  $w_n$  converges pointwise a.e. to  $w$ . From  $(\mathcal{N}_{ii})$  and Lemma 2.4,

$$\min \left\{ \int a^+ |u|^{p-2} u \tilde{u}^+, - \int a^+ |u|^{p-2} u \tilde{u}^-, \int a^+ |u|^{p-2} u \hat{u}^+ \right\} \geq \kappa.$$

These three integrals are also bounded above by a constant independent of  $\mu$  because  $\mathcal{N}_\mu$  is bounded. It follows from Lemma 2.3 that the integrals

$$\int a^+ |\tilde{u}^+|^p, \quad \int a^+ |\tilde{u}^-|^p, \quad \int a^+ |\hat{u}^+|^p$$

are bounded below by a positive constant independent of  $\mu$ . The Sobolev inequality now implies that the norms

$$\|\tilde{u}^+\|, \quad \|\tilde{u}^-\|, \quad \|\hat{u}^+\|$$

are bounded below by a positive constant independent of  $\mu$ . From the lower-semicontinuity of the norm,

$$\|\tilde{u}^+\| \leq \liminf \|\tilde{u}_n^+\|, \quad \|\tilde{u}^-\| \leq \liminf \|\tilde{u}_n^-\|, \quad \|\hat{u}^+\| \leq \liminf \|\hat{u}_n^+\|. \quad (17)$$

We wish to prove that equalities hold. Otherwise, choose  $(\tilde{r}, \tilde{s}, \hat{t})$ , defined by

$$\begin{aligned} \tilde{r} &= \left( \frac{\|\tilde{u}^+\|^2}{\int a^+ |u|^{p-2} u \tilde{u}^+} \right)^{\frac{1}{p-2}}, \quad \tilde{s} = \left( \frac{\|\tilde{u}^-\|^2}{-\int a^+ |u|^{p-2} u \tilde{u}^-} \right)^{\frac{1}{p-2}}, \\ \hat{t} &= \left( \frac{\|\hat{u}^+\|^2}{\int a^+ |u|^{p-2} u \hat{u}^+} \right)^{\frac{1}{p-2}}, \end{aligned}$$

so that the function

$$w := \tilde{r} \tilde{u}^+ - \tilde{s} \tilde{u}^- + \hat{t} \hat{u}^+ - \hat{u}^- + \bar{u} + \underline{u}$$

satisfies  $(\mathcal{N}_{ii})$ . By (17), the strong convergence in  $L^p(\Omega)$ , and what we have just seen,

$$(\tilde{r}, \tilde{s}, \hat{t}) \in [c, 1]^3 \setminus \{(1, 1, 1)\},$$

for some  $c > 0$  independent of  $\mu$ . The function  $w$  clearly satisfies  $(\mathcal{N}_i)$  and  $(\mathcal{N}_v)$ . Lemma 2.3 guarantees that  $(\mathcal{N}_{iv})$  is satisfied for sufficiently large  $\mu$ . Consider the estimate

$$\begin{aligned} I_\mu(\tilde{r} \tilde{u}^+ - \tilde{s} \tilde{u}^- + \hat{t} \hat{u}^+ - \hat{u}^- + \bar{u} + \underline{u}) \\ &< \liminf I_\mu(\tilde{r} \tilde{u}_n^+ - \tilde{s} \tilde{u}_n^- + \hat{t} \hat{u}_n^+ - \hat{u}_n^- + \bar{u}_n + \underline{u}_n) \\ &\leq \lim I_\mu(u_n), \end{aligned}$$

where the last inequality is due to Lemma 3.1. It shows that  $w$  satisfies  $(\mathcal{N}_{iii})$ . Therefore  $w \in \mathcal{N}_\mu$  and  $I_\mu(w) < \lim I_\mu(u_n)$ . This is a contradiction. We have established that equality holds in all three of (17). Therefore  $u \in \mathcal{N}_\mu$  for large  $\mu$ .  $\square$

## 4 A minimizer in $\mathcal{N}_\mu$ is a critical point

In the previous section we obtained a minimizer  $u$  of  $I_\mu$  on  $\mathcal{N}_\mu$ . We will now prove that this minimizer is indeed a critical point of  $I_\mu$ . This will be done by using a deformation argument on the manifold introduced above. Let  $\sigma$  be the restriction to the interval  $[1/2, 2]^3$  of the  $\varsigma$  corresponding to the minimizer  $u$ . Recall  $\varsigma$  was defined in (14). We define a negative gradient flow in a neighborhood of  $u$  in the following way. Let  $B_\rho(u) := \{w \in H_0^1(\Omega) : \|w - u\| < \rho\}$ , where  $\rho$  is chosen small enough so that

$$\sigma(\tilde{r}, \tilde{s}, \hat{t}) \in B_\rho(u) \Rightarrow \frac{1}{2} < \tilde{r}, \tilde{s}, \hat{t} < 2 \quad (18)$$

and  $w \in B_\rho(u)$  implies that  $w$  satisfies  $(\mathcal{N}_i)$ ,  $(\mathcal{N}_{iii})$ ,  $(\mathcal{N}_{iv})$  and  $(\mathcal{N}_v)$ , for sufficiently large  $\mu$ . Such a  $\rho$  exists because the function  $u$  satisfies (11) and (a), (b), (c) and (d) of Lemma 2.5. Let  $\varphi$  be a Lipschitz function,  $\varphi : H_0^1(\Omega) \rightarrow [0, 1]$ , such that  $\varphi = 1$  on  $B_{\rho/2}(u)$  and  $\varphi = 0$  on the complement of  $B_\rho(u)$ . Consider the Cauchy problem

$$\begin{cases} \frac{d\eta}{d\tau} = -\varphi(\eta)\nabla I_\mu(\eta), \\ \eta(0) = w, \end{cases} \quad (19)$$

whose solution we denote by  $\eta(\tau; w)$ . For  $\tau \geq 0$ , let

$$\sigma_\tau(\tilde{r}, \tilde{s}, \hat{t}) = \eta(\tau; \sigma(\tilde{r}, \tilde{s}, \hat{t})).$$

**Lemma 4.1.** *The set  $\sigma_\tau([1/2, 2]^3)$  intersects  $\mathcal{N}_\mu$  in a nonempty set.*

*Proof.* Consider the maps  $\tilde{\phi}^\pm, \hat{\phi}, \tilde{\psi}^\pm, \hat{\psi}$  from  $\{w \in H_0^1(\Omega) : \tilde{w}^\pm \neq 0, \hat{w}^+ \neq 0\}$  to  $\mathbb{R}$ , defined by

$$\begin{aligned} \tilde{\phi}^\pm(w) &= \frac{\pm \int a^+ |w|^{p-2} w \tilde{w}^\pm}{\|\tilde{w}^\pm\|^2}, & \hat{\phi}(w) &= \frac{\int a^+ |w|^{p-2} w \hat{w}^+}{\|\hat{w}^+\|^2}, \\ \tilde{\psi}^\pm(w) &= \frac{\int a^+ |\tilde{w}^\pm|^p}{\|\tilde{w}^\pm\|^2}, & \hat{\psi}(w) &= \frac{\int a^+ |\hat{w}^+|^p}{\|\hat{w}^+\|^2}. \end{aligned}$$

These maps are well defined on  $\sigma_\tau([1/2, 2]^3)$ , because if  $w \in B_\rho(u)$ , then  $w$  satisfies  $(\mathcal{N}_i)$ . We finally define

$$\Phi_\tau := (\tilde{\phi}^+, \tilde{\phi}^-, \hat{\phi}) \circ \sigma_\tau$$

and

$$\Psi := (\tilde{\psi}^+, \tilde{\psi}^-, \hat{\psi}) \circ \sigma,$$

from  $([1/2, 2]^3)$  to  $\mathbb{R}^3$ . Since  $\int |\underline{u}|^p = o(1)$  uniformly in  $u$  and  $\mu$  and the value of  $\kappa$  in Lemma 2.4 is independent of  $\mu$ ,

$$\begin{aligned}\Psi(\tilde{r}, \tilde{s}, \hat{t}) &= \left( \tilde{r}^{p-2} \tilde{\psi}^+(u), \tilde{s}^{p-2} \tilde{\psi}^-(u), \hat{t}^{p-2} \hat{\psi}(u) \right) \\ &= \left( (1 + o(1)) \tilde{r}^{p-2}, (1 + o(1)) \tilde{s}^{p-2}, (1 + o(1)) \hat{t}^{p-2} \right),\end{aligned}\quad (20)$$

with the last three  $o(1)$  independent of  $u$  and  $\mu$ . As a consequence,

$$\text{dist} \left( \Psi \left( \partial[1/2, 2]^3 \right), (1, 1, 1) \right) \geq c > 0,$$

the constant  $c$  being independent of  $u$  and  $\mu$ . We deduce from (20) that for large  $\mu$ ,

$$\text{deg} \left( \Psi, [1/2, 2]^3, (1, 1, 1) \right) = 1.$$

Notice that condition (18) and the definition of the flux (19) guarantee

$$\Phi_\tau|_{\partial[1/2, 2]^3} = \Phi_0|_{\partial[1/2, 2]^3} = \Psi|_{\partial[1/2, 2]^3} + o(1)$$

and therefore

$$\text{deg} \left( \Phi_\tau, [1/2, 2]^3, (1, 1, 1) \right) = 1.$$

for  $\mu$  large enough. This proves that

$$\sigma_\tau \left( [1/2, 2]^3 \right) \cap \mathcal{N}_\mu \neq \emptyset.$$

□

We are ready to give the

*Proof of Proposition 2.1.* Let  $\mu$  be large and  $u_\mu$  be a minimizer of  $I_\mu$  restricted to  $\mathcal{N}_\mu$ . The existence of such a  $u_\mu$  was proven in Lemma 3.2. Suppose that  $I'_\mu(u_\mu) \neq 0$ . By Lemma 3.1, with  $u = u_\mu$ ,  $\max I_\mu \circ \sigma \left( [1/2, 2]^3 \right) = I_\mu(u_\mu)$ , and so for any small  $\tau > 0$ ,

$$\max I_\mu \circ \sigma_\tau \left( [1/2, 2]^3 \right) < I_\mu(u_\mu).$$

This contradicts Lemma 4.1. So  $I'_\mu(u_\mu) = 0$ , and the minimizer of  $I_\mu$  on  $\mathcal{N}_\mu$  is a weak solution of (4).

Consider now  $u$  as in (5). Properties (6), (7) and (8) follow from Lemma 2.4 and Lemma 2.5 (c), (d), as

$$\min \left\{ \int a^+ |u_\mu|^{p-2} u_\mu \tilde{u}_\mu^+, - \int a^+ |u_\mu|^{p-2} u_\mu \tilde{u}_\mu^-, \int a^+ |u_\mu|^{p-2} u_\mu \hat{u}_\mu^+ \right\} \geq \kappa.$$

□

Theorem 1.1 can be proved as Proposition 2.1 with obvious adaptations.

## References

- [1] Ackermann, N.; Weth, T.. Multibump solutions of nonlinear periodic Schrödinger equations in a degenerate setting. *Commun. Contemp. Math.* 7 (2005), no. 3, 269–298.
- [2] Alama, S.; Del Pino, M.. Solutions of elliptic equations with indefinite nonlinearities via Morse theory and linking. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 13 (1996), no. 1, 95–115.
- [3] Alama, S.; Tarantello, G.. On semilinear elliptic equations with indefinite nonlinearities. *Calc. Var. Partial Differential Equations* 1 (1993), no. 4, 439–475.
- [4] Bartsch, T.; Clapp, M.; Weth, T.. Configuration spaces, transfer, and 2-nodal solutions of a semiclassical nonlinear Schrödinger equation. *Math. Ann.* to appear.
- [5] Bartsch, T.; Weth, T.. A note on additional properties of sign changing solutions to superlinear elliptic equations. *Topol. Methods Nonlinear Anal.* 22 (2003), no. 1, 1–14.
- [6] Berestycki, H.; Capuzzo-Dolcetta, I.; Nirenberg, L.. Variational methods for indefinite superlinear homogeneous elliptic problems. *NoDEA Nonlinear Differential Equations Appl.* 2 (1995), no. 4, 553–572.
- [7] Bonheure, D.; Gomes, J.M.; Habets, P.. Multiple positive solutions of superlinear elliptic problems with sign-changing weight. *J. Differential Equations* 214 (2005), no. 1, 36–64.
- [8] Castro, A.; Cossio, J.; Neuberger, J.M.. A sign-changing solution for a superlinear Dirichlet problem. *Rocky Mountain J. Math.* 27 (1997), no. 4, 1041–1053.
- [9] Cerami, G.; Solimini, S.; Struwe, M.. Some existence results for superlinear elliptic boundary value problems involving critical exponents. *J. Funct. Anal.* 69 (1986), no. 3, 289–306.
- [10] Clapp, M.; Weth, T.. Minimal nodal solutions of the pure critical exponent problem on a symmetric domain. *Calc. Var. Partial Differential Equations* 21 (2004), no. 1, 1–14.
- [11] Costa, D.G.; Ramos, M.; Tehrani, H.. Non-zero solutions for a Schrödinger equation with indefinite linear and nonlinear terms. *Proc. Roy. Soc. Edinburgh Sect. A* 134 (2004), no. 2, 249–258.

- [12] Coti Zelati, V.; Rabinowitz, P.H.. Homoclinic orbits for second order Hamiltonian systems possessing superquadratic potentials. *J. Amer. Math. Soc.* 4 (1991), no. 4, 693–727.
- [13] Coti Zelati, V.; Rabinowitz, P.H.. Homoclinic type solutions for a semi-linear elliptic PDE on  $\mathbb{R}^n$ . *Comm. Pure Appl. Math.* 45 (1992), no. 10, 1217–1269.
- [14] Del Pino, M.; Felmer, P.L.. Multi-peak bound states for nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 15 (1998), no. 2, 127–149.
- [15] Gaudenzi, M.; Habets, P.; Zanolin, F.. A seven-positive-solutions theorem for a superlinear problem. *Adv. Nonlinear Stud.* 4 (2004), no. 2, 149–164.
- [16] Li, Y.; Wang, Z.Q.. Gluing approximate solutions of minimum type on the Nehari manifold. *Proceedings of the USA-Chile Workshop on Nonlinear Analysis (Viña del Mar-Valparaiso, 2000)*, 215–223, *Electron. J. Differ. Equ. Conf.*, 6, Southwest Texas State Univ., San Marcos, TX, 2001.
- [17] Ramos, M.. Remarks on a priori estimates for superlinear elliptic problems. *Topological methods, variational methods and their applications (Taiyuan, 2002)*, 193–200, *World Sci. Publ.*, River Edge, NJ, 2003.
- [18] Ramos, M.; Tavares, H.. Solutions with multiple spike patterns for an elliptic system. Preprint.
- [19] Ramos, M.; Terracini, S.; Troestler, C.. Superlinear indefinite elliptic problems and Pohožaev type identities. *J. Funct. Anal.* 159 (1998), no. 2, 596–628.
- [20] Séré, É.. Existence of infinitely many homoclinic orbits in Hamiltonian systems. *Math. Z.* 209 (1992), no. 1, 27–42.
- [21] Tehrani, H.. On indefinite superlinear elliptic equations. *Calc. Var. Partial Differential Equations* 4 (1996), no. 2, 139–153.