

REALIZATION OF QUANTUM GROUP POISSON BOUNDARIES AS CROSSED PRODUCTS

MEHRDAD KALANTAR, MATTHIAS NEUFANG AND ZHONG-JIN RUAN

ABSTRACT. For a locally compact quantum group \mathbb{G} , consider the convolution action of a quantum probability measure μ on $L_\infty(\mathbb{G})$. As shown by Junge–Neufang–Ruan, this action has a natural extension to a Markov map on $\mathcal{B}(L_2(\mathbb{G}))$. We prove that the Poisson boundary of the latter can be realized concretely as the von Neumann crossed product of the Poisson boundary associated with μ under the action of \mathbb{G} induced by the coproduct. This yields an affirmative answer, for general locally compact quantum groups, to a problem raised by Izumi (2004) in the commutative situation, in which he settled the discrete case, and unifies earlier results of Jaworski, Neufang and Runde.

1. INTRODUCTION AND PRELIMINARIES

Poisson boundaries and harmonic functions have played a very important role in the study of random walks on discrete groups, and more generally in harmonic analysis and ergodic theory on locally compact groups (see for instance Furstenberg’s seminal work [6]). The noncommutative version of this concept can be considered in two different directions: one replaces $L_\infty(G)$ by certain quantum groups, such as its dual (quantum) group von Neumann algebra $VN(G)$ or quantum groups arising from mathematical physics; in another direction, one can perform the quantum mechanical passage from $L_\infty(G)$ to $\mathcal{B}(L_2(G))$.

In the first direction, Chu–Lau [2] studied the case of $VN(G)$, and Poisson boundaries over (discrete) quantum groups were first investigated by Izumi [9], in particular for the dual of Woronowicz’s compact quantum group $SU_q(2)$. Izumi’s results were further generalized to other discrete quantum groups by Izumi–Neshveyev–Tuset, Tomatsu, Vaes, Vander Vennet and Vergnioux ([12], [25], [26], [27]).

The second way of quantization has been studied by Neufang–Ruan–Spronk in [19], based on a natural isometric representation Θ of the measure algebra $M(G)$ of a locally compact group G , as completely bounded maps on $\mathcal{B}(L_2(G))$, extending the convolution action of measures on $L_\infty(G)$ (cf. [7], [24]). The structure of the Poisson boundary at the level of $\mathcal{B}(L_2(G))$, associated with a probability measure, has been analyzed by Izumi in [10, 11] for discrete groups, and by Jaworski–Neufang for locally compact groups in [13], answering a question raised in [10]. Dually, [19] investigates the extension of the pointwise action of the completely bounded Fourier algebra multipliers $M_{cb}A(G)$, from $VN(G)$ to the level of $\mathcal{B}(L_2(G))$; the structure of the corresponding Poisson boundary associated with positive definite functions has been described in [20], providing a noncommutative analogue of the situation considered in [2].

2000 *Mathematics Subject Classification.* Primary 46L55, 46L89; Secondary 46L07, 60J50.

The second and third author were partially supported by NSERC and the Simons Foundation, respectively.

In this paper, we combine these two quantization methods: for a locally compact quantum group \mathbb{G} , we consider the natural extension of the action of a quantum probability measure μ on $L_\infty(\mathbb{G})$ to $\mathcal{B}(L_2(\mathbb{G}))$, as introduced and studied by Junge–Neufang–Ruan in [14]. We completely describe the structure of the Poisson boundary $\mathcal{H}_{\Theta(\mu)}$ of the induced Markov operator $\Theta(\mu)$. The main result (Theorem 2.4) of the paper gives a concrete realization of $\mathcal{H}_{\Theta(\mu)}$ as the (von Neumann) crossed product of \mathcal{H}_μ with the quantum group \mathbb{G} under a natural action.

To obtain an action of the quantum group \mathbb{G} on the Poisson boundary, it is natural to consider the restriction of the co-multiplication to the latter. In our setting, a difficulty lies in the fact that in general $L_\infty(\mathbb{G})$ need not have the slice map property (S_σ); for instance, given a discrete group G , $VN(G)$ has S_σ if and only if G has the approximation property (AP), cf. [8]. Note that this problem does not arise in the case of commutative or discrete quantum groups \mathbb{G} since $L_\infty(\mathbb{G})$ then has S_σ . We solve this problem by using the Fubini product, a tool from operator space theory.

In [10, Theorem 4.1] Izumi proved a crossed product formula for Poisson boundaries associated with $\ell_\infty(G)$, where G is a countable discrete group, and as a direct consequence he concluded that the natural action of G on its Poisson boundary is amenable. Hence, he provided a new proof for this well-known result of Zimmer [28], and therefore raised the question if such an identification result extends to all second countable locally compact groups [10, Problem 4.3]. This problem was answered affirmatively in [13]. The present paper answers Izumi’s question even in the setting of locally compact quantum groups \mathbb{G} . As a corollary, we conclude that the crossed product von Neumann algebra of the natural action of \mathbb{G} on its (noncommutative) Poisson boundary is injective.

We recall some definitions and preliminary results that will be used in the sequel. For more details on locally compact quantum groups we refer the reader to [17], [18].

A *locally compact quantum group* \mathbb{G} is a quadruple $(L_\infty(\mathbb{G}), \Gamma, \varphi, \psi)$, where $L_\infty(\mathbb{G})$ is a von Neumann algebra, $\Gamma : L_\infty(\mathbb{G}) \rightarrow L_\infty(\mathbb{G}) \bar{\otimes} L_\infty(\mathbb{G})$ is a co-associative co-multiplication, and φ and ψ are (normal faithful semi-finite) left, respectively, right invariant weights on $L_\infty(\mathbb{G})$, called the *Haar weights*. The weight ψ determines a Hilbert space $L_2(\mathbb{G}) = L_2(\mathbb{G}, \psi)$, and we obtain the *right fundamental unitary operator* V on $L_2(\mathbb{G}) \otimes L_2(\mathbb{G})$, which satisfies the *pentagonal relation* $V_{12}V_{13}V_{23} = V_{23}V_{12}$. Here we used the leg notation $V_{12} = V \otimes 1$, $V_{23} = 1 \otimes V$, and $V_{13} = (\iota \otimes \chi)V_{12}$, where $\chi(x \otimes y) = y \otimes x$ is the flip map on $\mathcal{B}(H \otimes K)$. We will also use the flip map σ defined on $H \otimes K$. The fundamental unitary operator induces a coassociative comultiplication

$$(1.1) \quad \tilde{\Gamma} : x \in \mathcal{B}(L_2(\mathbb{G})) \rightarrow \tilde{\Gamma}(x) = V(x \otimes 1)V^* \in \mathcal{B}(L_2(\mathbb{G}) \otimes L_2(\mathbb{G}))$$

on $\mathcal{B}(L_2(\mathbb{G}))$, for which we have $\tilde{\Gamma}|_{L_\infty(\mathbb{G})} = \Gamma$.

Let $L_1(\mathbb{G})$ be the predual of $L_\infty(\mathbb{G})$. Then the pre-adjoint of Γ induces an associative completely contractive multiplication

$$(1.2) \quad \star : f \otimes g \in L_1(\mathbb{G}) \hat{\otimes} L_1(\mathbb{G}) \rightarrow f \star g = (f \otimes g)\Gamma \in L_1(\mathbb{G})$$

on $L_1(\mathbb{G})$. The convolution actions $x \star \mu := (\iota \otimes \mu)\Gamma(x)$ and $\mu \star x := (\mu \otimes \iota)\Gamma(x)$ are normal completely bounded maps on $L_\infty(\mathbb{G})$.

We denote by $C_0(\mathbb{G})$ the *reduced quantum group C^* -algebra*, which is a weak* dense C^* -subalgebra of $L_\infty(\mathbb{G})$. Let $M(\mathbb{G})$ denote the operator dual $C_0(\mathbb{G})^*$. There exists a completely contractive multiplication on $M(\mathbb{G})$ given by the convolution

$$\star : \mu \otimes \nu \in M(\mathbb{G}) \widehat{\otimes} M(\mathbb{G}) \mapsto \mu \star \nu = \mu(\iota \otimes \nu)\Gamma = \nu(\mu \otimes \iota)\Gamma \in M(\mathbb{G})$$

such that $M(\mathbb{G})$ contains $L_1(\mathbb{G})$ as a norm closed two-sided ideal.

We denote by $C_u(\mathbb{G})$ the *universal quantum group C^* -algebra* of \mathbb{G} (see [16] for details). There is a comultiplication

$$\Gamma_u : C_u(\mathbb{G}) \rightarrow M(C_u(\mathbb{G}) \otimes_{\min} C_u(\mathbb{G})),$$

and the operator dual $M_u(\mathbb{G}) = C_u(\mathbb{G})^*$, which can be regarded as the space of all *quantum measures* on \mathbb{G} , is a unital completely contractive Banach algebra with multiplication given by

$$\omega \star_u \mu = \omega(\iota \otimes \mu)\Gamma_u = \mu(\omega \otimes \iota)\Gamma_u.$$

Moreover, the convolution algebras $M(\mathbb{G})$ and $L_1(\mathbb{G})$ can be canonically identified with norm closed two-sided ideals in $M_u(\mathbb{G})$. Therefore, for each $\mu \in M_u(\mathbb{G})$, we obtain a pair of completely bounded maps

$$(1.3) \quad \mathfrak{m}_\mu^l(f) = \mu \star f \quad \text{and} \quad \mathfrak{m}_\mu^r(f) = f \star \mu$$

on $L_1(\mathbb{G})$ with $\max\{\|\mathfrak{m}_\mu^l\|_{cb}, \|\mathfrak{m}_\mu^r\|_{cb}\} \leq \|\mu\|$. The adjoint map $\Phi_\mu = (\mathfrak{m}_\mu^r)^*$ is a normal completely bounded map on $L_\infty(\mathbb{G})$ satisfying the covariance condition

$$(1.4) \quad \Gamma \circ \Phi_\mu = (\iota \otimes \Phi_\mu) \circ \Gamma,$$

or equivalently, $\Phi_\mu(f \star x) = f \star \Phi_\mu(x)$ for all $x \in L_\infty(\mathbb{G})$ and $f \in L_1(\mathbb{G})$. We are particularly interested in the case when μ is a state in $M_u(\mathbb{G})$. In this case, Φ_μ is unital completely positive, i.e., a Markov operator, on $L_\infty(\mathbb{G})$.

2. CONCRETE REALIZATION OF THE POISSON BOUNDARY IN $\mathcal{B}(L_2(\mathbb{G}))$

In the following, \mathbb{G} denotes a general locally compact quantum group. We write $\mathcal{P}_u(\mathbb{G})$ for the set of all states on $C_u(\mathbb{G})$ (i.e., the ‘quantum probability measures’). We consider the space of fixed points $\mathcal{H}^\mu = \{x \in L_\infty(\mathbb{G}) : \Phi_\mu(x) = x\}$. It is easy to see that \mathcal{H}^μ is a weak* closed operator system in $L_\infty(\mathbb{G})$. In fact, we obtain a natural von Neumann algebra product on this space. Let us recall this construction for the convenience of the reader (cf. [9, Section 2.5]).

We first define a projection $\mathcal{E}_\mu : L_\infty(\mathbb{G}) \rightarrow L_\infty(\mathbb{G})$ of norm one by

$$(2.1) \quad \mathcal{E}_\mu(x) = w^* - \lim_{\mathcal{U}} \frac{1}{n} \sum_{k=1}^n \Phi_\mu^k(x)$$

with respect to a free ultrafilter \mathcal{U} on \mathbb{N} . It is easy to see that $\mathcal{H}^\mu = \mathcal{E}_\mu(L_\infty(\mathbb{G}))$, and that the Choi-Effros product

$$(2.2) \quad x \circ y = \mathcal{E}_\mu(xy)$$

defines a von Neumann algebra product on \mathcal{H}^μ . We note that this product is independent of the choice of the free ultrafilter \mathcal{U} since every completely positive isometric linear isomorphism between two von Neumann algebras is a *-isomorphism. To avoid confusion, we denote by $\mathcal{H}_\mu = (\mathcal{H}^\mu, \circ)$ this von Neumann algebra, and we call \mathcal{H}_μ the *Poisson boundary* of μ .

It follows from [14, Theorem 4.5] that the Markov operator Φ_μ has a unique weak* continuous (unital completely positive) extension $\Theta(\mu)$ to $\mathcal{B}(L_2(\mathbb{G}))$ such that

$$(2.3) \quad \tilde{\Gamma} \circ \Theta(\mu) = (\iota \otimes \Phi_\mu) \circ \tilde{\Gamma}.$$

Similarly to (2.1) we obtain a projection $\mathcal{E}_{\Theta(\mu)}$ of norm one on $\mathcal{B}(L_2(\mathbb{G}))$ and a von Neumann algebra product on $\mathcal{H}^{\Theta(\mu)}$. We denote this von Neumann algebra by $\mathcal{H}_{\Theta(\mu)}$. Our main result (Theorem 2.4) shows that there is a left \mathbb{G} action on \mathcal{H}_μ such that $\mathcal{H}_{\Theta(\mu)}$ is *-isomorphic to the von Neumann algebra crossed product of \mathcal{H}_μ by \mathbb{G} .

Unlike in the classical and the discrete settings, in the case of general locally compact quantum groups, the fact that the coproduct induces an action of \mathbb{G} on a Poisson boundary is not trivial. So, first we need to prove this result. For this purpose, we need to recall the Fubini product for weak* closed operator spaces on Hilbert spaces. Suppose that V and W are weak* closed subspaces of $\mathcal{B}(H)$ and $\mathcal{B}(K)$, respectively. We define the *Fubini product* of V and W to be the space

$$V \overline{\otimes}_{\mathcal{F}} W = \{X \in \mathcal{B}(H) \overline{\otimes} \mathcal{B}(K) : (\omega \otimes \iota)(X) \in W \text{ and } (\iota \otimes \phi)(X) \in V \text{ for all } \omega \in \mathcal{B}(H)_*, \phi \in \mathcal{B}(K)_*\}.$$

In this case, $V_* = \mathcal{B}(H)_*/V_\perp$ and $W_* = \mathcal{B}(K)_*/W_\perp$ are operator preduals of V and W , respectively. It is known from [23, Proposition 3.3] (see also [3] and [4, §7.2]) that the Fubini product $V \overline{\otimes}_{\mathcal{F}} W$ is a weak* closed subspace of $\mathcal{B}(H \otimes K)$ such that we have the weak* homeomorphic completely isometric isomorphism

$$(2.4) \quad V \overline{\otimes}_{\mathcal{F}} W = (V_* \hat{\otimes} W_*)^*,$$

where $V_* \hat{\otimes} W_*$ is the operator space projective tensor product of V_* and W_* . In particular, if M and N are von Neumann algebras, the Fubini product coincides with the von Neumann algebra tensor product, i.e., we have

$$M \overline{\otimes}_{\mathcal{F}} N = M \overline{\otimes} N.$$

It is also known from operator space theory that there is a canonical completely isometrically identification

$$(2.5) \quad (V_* \hat{\otimes} W_*)^* = \mathcal{CB}(V_*, W)$$

given by the left slice maps. Now, if W_1 and W_2 are dual operator spaces and $\Psi : W_1 \rightarrow W_2$ is a (not necessarily weak* continuous) completely bounded map, we can apply (2.4) and (2.5) to obtain a completely bounded map

$$\iota \otimes \Psi : V \overline{\otimes}_{\mathcal{F}} W_1 \rightarrow V \overline{\otimes}_{\mathcal{F}} W_2$$

such that

$$(\omega \otimes \iota)(\iota \otimes \Psi)(X) = \Psi((\omega \otimes \iota)(X))$$

for all $X \in V \overline{\otimes}_{\mathcal{F}} W_1$ and $\omega \in V_*$. It is easy to see that we have $\|\iota \otimes \Psi\|_{cb} = \|\Psi\|_{cb}$, and if Ψ is a completely isometric isomorphism (respectively, completely contractive projection) then so is $\iota \otimes \Psi$.

Now since \mathcal{H}^μ and $\mathcal{H}_\mu = (\mathcal{H}^\mu, \circ)$ have the same predual, the identity map ι_μ is a weak* homeomorphic and completely isometric isomorphism from the weak* closed operator system \mathcal{H}^μ onto the von Neumann algebra \mathcal{H}_μ . So we obtain the weak* homeomorphic and completely isometric isomorphism

$$\iota \otimes \iota_\mu : L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}^\mu \rightarrow L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}_\mu.$$

Since both $L_\infty(\mathbb{G})$ and \mathcal{H}_μ are von Neumann algebras, we can identify $L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}_\mu$ with the von Neumann algebra $L_\infty(\mathbb{G}) \overline{\otimes} \mathcal{H}_\mu$. We note that since \mathcal{E}_μ is a (not necessarily normal) projection of norm one from $L_\infty(\mathbb{G})$ onto $\mathcal{H}^\mu \subseteq L_\infty(\mathbb{G})$, the map $\iota \otimes \mathcal{E}_\mu$ defines a projection of norm one from $L_\infty(\mathbb{G}) \overline{\otimes} L_\infty(\mathbb{G}) = L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} L_\infty(\mathbb{G})$ onto $L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}^\mu \subseteq L_\infty(\mathbb{G}) \overline{\otimes} L_\infty(\mathbb{G})$ and thus induces a Choi-Effros product such that $L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}^\mu$ becomes a von Neumann algebra. It turns out that (up to the above identification) this von Neumann algebra is exactly equal to $L_\infty(\mathbb{G}) \overline{\otimes} \mathcal{H}_\mu$.

Proposition 2.1. *For any $\mu \in \mathcal{P}_u(\mathbb{G})$ the restriction of Γ to \mathcal{H}_μ induces a left action Γ_μ of \mathbb{G} on the von Neumann algebra \mathcal{H}_μ .*

Proof. Let us first show that $\Gamma(\mathcal{H}^\mu) \subseteq L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}^\mu$. Given $x \in \mathcal{H}^\mu$ and $f \in L_1(\mathbb{G})$, we have

$$\Phi_\mu((f \otimes \iota)\Gamma(x)) = (f \otimes \iota)(\iota \otimes \Phi_\mu)\Gamma(x) = (f \otimes \iota)\Gamma(\Phi_\mu(x)) = (f \otimes \iota)\Gamma(x).$$

This shows that $(f \otimes \iota)\Gamma(x)$ is contained in \mathcal{H}^μ for all $f \in L_1(\mathbb{G})$. On the other hand, we clearly have $(\iota \otimes g)\Gamma(x) \in L_\infty(\mathbb{G})$ for all $g \in (\mathcal{H}_\mu)_*$. Hence, $\Gamma(x) \in L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}^\mu$.

It is clear that $\Gamma|_{\mathcal{H}^\mu} : \mathcal{H}^\mu \rightarrow L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}^\mu$ is a normal injective unital completely positive isometry. This induces the map $\Gamma_\mu : \mathcal{H}_\mu \rightarrow L_\infty(\mathbb{G}) \overline{\otimes} \mathcal{H}_\mu$ given by $\Gamma_\mu = (\iota \otimes \iota_\mu) \circ \Gamma \circ \iota_\mu^{-1}$. It suffices to show that Γ_μ is an algebra homomorphism with respect to the corresponding Choi-Effros products on \mathcal{H}_μ and $L_\infty(\mathbb{G}) \overline{\otimes}_{\mathcal{F}} \mathcal{H}_\mu$. Given $x, y \in \mathcal{H}_\mu$, we now use (2.1) and (2.2) to obtain that

$$\begin{aligned} \Gamma_\mu(x \circ y) &= \Gamma(\mathcal{E}_\mu(xy)) = \Gamma(w^* - \lim_{\mathcal{U}} \frac{1}{n} \sum_{k=1}^n \Phi_\mu^k(xy)) = w^* - \lim_{\mathcal{U}} \frac{1}{n} \sum_{k=1}^n \Gamma(\Phi_\mu^k(xy)) \\ &= w^* - \lim_{\mathcal{U}} \frac{1}{n} \sum_{k=1}^n (\iota \otimes \Phi_\mu^k)(\Gamma(xy)) = (\iota \otimes \mathcal{E}_\mu)(\Gamma(x)\Gamma(y)) \\ &= \Gamma_\mu(x) \circ \Gamma_\mu(y) \in L_\infty(\mathbb{G}) \overline{\otimes} \mathcal{H}_\mu. \end{aligned}$$

Since Γ is a comultiplication on $L_\infty(\mathbb{G})$, it is clear that Γ_μ satisfies

$$(\iota \otimes \Gamma_\mu) \circ \Gamma_\mu = (\Gamma \otimes \iota) \circ \Gamma_\mu.$$

So Γ_μ defines a left action. □

As we discussed above, we have a projection $\mathcal{E}_{\Theta(\mu)}$ of norm one from $\mathcal{B}(L_2(\mathbb{G}))$ onto $\mathcal{H}^{\Theta(\mu)}$ and obtain the Choi-Effros von Neumann algebra product on $\mathcal{H}_{\Theta(\mu)}$ given by

$$X \circ Y = \mathcal{E}_{\Theta(\mu)}(XY)$$

for $X, Y \in \mathcal{H}_{\Theta(\mu)} = (\mathcal{H}^{\Theta(\mu)}, \circ)$. It is easy to see that the restriction of $\mathcal{E}_{\Theta(\mu)}$ to $L_\infty(\mathbb{G})$ is equal to \mathcal{E}_μ and \mathcal{H}_μ is a von Neumann subalgebra of $\mathcal{H}_{\Theta(\mu)}$. Recall that $\tilde{\Gamma}$ denotes the comultiplication on $\mathcal{B}(L_2(\mathbb{G}))$ defined in (1.1).

Proposition 2.2. *For any $\mu \in \mathcal{P}_u(\mathbb{G})$ the restriction of $\tilde{\Gamma}$ to $\mathcal{H}_{\Theta(\mu)}$ induces a normal injective unital $*$ -homomorphism*

$$\tilde{\Gamma}_{\Theta(\mu)} : \mathcal{H}_{\Theta(\mu)} \rightarrow \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu$$

between von Neumann algebras. Moreover, the restriction of $\tilde{\Gamma}_{\Theta(\mu)}$ to \mathcal{H}_μ is equal to Γ_μ .

Proof. For any $x \in \mathcal{H}_{\Theta(\mu)}$, we have $\tilde{\Gamma}(x) \in \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} L_\infty(\mathbb{G})$. We can apply (2.3) to get

$$\Phi_\mu \left((\omega \otimes \iota) \tilde{\Gamma}(x) \right) = (\omega \otimes \iota) (\iota \otimes \Phi_\mu) \tilde{\Gamma}(x) = (\omega \otimes \iota) (\tilde{\Gamma}(\Theta(\mu)(x))) = (\omega \otimes \iota) (\tilde{\Gamma}(x))$$

for all $\omega \in \mathcal{B}(L_2(\mathbb{G}))_*$. This shows that $\tilde{\Gamma}(\mathcal{H}_{\Theta(\mu)}) \subseteq \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes}_{\mathcal{F}} \mathcal{H}_\mu = \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu$. The rest of proof is similar to that given in the proof of Proposition 2.1. □

For $\mu \in \mathcal{P}_u(\mathbb{G})$, let Γ_μ be the left action of \mathbb{G} on the von Neumann algebra \mathcal{H}_μ given in Proposition 2.1. The crossed product $\mathbb{G} \rtimes_{\Gamma_\mu} \mathcal{H}_\mu$ is defined to be the von Neumann algebra

$$\mathbb{G} \rtimes_{\Gamma_\mu} \mathcal{H}_\mu = [\Gamma_\mu(\mathcal{H}_\mu) \cup (L_\infty(\hat{\mathbb{G}}) \otimes 1)]''$$

in $\mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu$. The following result, which is crucial for us, even holds in the setting of measured quantum groupoids (cf. [5, Theorem 11.6]).

Theorem 2.3. *Let $\mu \in \mathcal{P}_u(\mathbb{G})$. Denote by χ the flip map $\chi(a \otimes b) = b \otimes a$. Then*

$$\beta : x \in \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu \rightarrow (\sigma V^* \sigma \otimes 1) ((\chi \otimes \iota) (\iota \otimes \Gamma_\mu)(x)) (\sigma V \sigma \otimes 1) \in L_\infty(\mathbb{G}) \overline{\otimes} \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu$$

defines a left action of \mathbb{G} on the von Neumann algebra $\mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu$, and we have

$$\mathbb{G} \rtimes_{\Gamma_\mu} \mathcal{H}_\mu = (\mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu)^\beta,$$

where $(\mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu)^\beta = \{y \in \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu : \beta(y) = 1 \otimes y\}$ is the fixed point algebra of β .

Now we can prove the main theorem of this paper. Applied to the case $\mathbb{G} = L_\infty(G)$, even without the assumption of second countability, our theorem yields [13, Proposition 6.3] which provided the answer to Izumi's original question. Specializing to the case $\mathbb{G} = VN(G)$, our result implies the main theorem of [20] where it was assumed either that G has the approximation property, or μ belongs to the Fourier algebra $A(G) = L_1(\mathbb{G})$.

Theorem 2.4. *Let \mathbb{G} be a locally compact quantum group and let $\mu \in \mathcal{P}_u(\mathbb{G})$. The induced map*

$$\tilde{\Gamma}_{\Theta(\mu)} : \mathcal{H}_{\Theta(\mu)} \rightarrow \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu$$

defines a von Neumann algebra isomorphism between $\mathcal{H}_{\Theta(\mu)}$ and $\mathbb{G} \rtimes_{\Gamma_\mu} \mathcal{H}_\mu$.

Proof. Since $V \in L_\infty(\hat{\mathbb{G}})' \overline{\otimes} L_\infty(\mathbb{G})$, it is easy to see from (1.1) and (2.3) that we have

$$\Theta(\mu)(\hat{x}) \otimes 1 = V^*(\tilde{\Gamma}(\Theta(\mu)(\hat{x})))V = V^*(\iota \otimes \Phi_\mu)(V(\hat{x} \otimes 1)V^*)V = \hat{x} \otimes 1$$

for all $\hat{x} \in L_\infty(\hat{\mathbb{G}})$. This shows that $\hat{x} \in \mathcal{H}_{\Theta(\mu)}$ and

$$\hat{x} \otimes 1 = \tilde{\Gamma}(\hat{x}) = \tilde{\Gamma}_{\Theta(\mu)}(\hat{x}) \in \tilde{\Gamma}_{\Theta(\mu)}(\mathcal{H}_{\Theta(\mu)})$$

for all $\hat{x} \in L_\infty(\hat{\mathbb{G}})$. Moreover, since $\mathcal{H}_\mu \subseteq \mathcal{H}_{\Theta(\mu)}$ and the restriction of $\tilde{\Gamma}_{\Theta(\mu)}$ to \mathcal{H}_μ is equal to Γ_μ , we obtain

$$\Gamma_\mu(\mathcal{H}_\mu) = \tilde{\Gamma}_{\Theta(\mu)}(\mathcal{H}_\mu) \subseteq \tilde{\Gamma}_{\Theta(\mu)}(\mathcal{H}_{\Theta(\mu)}).$$

Since $\tilde{\Gamma}_{\Theta(\mu)}$ is a unital $*$ -homomorphism from $\mathcal{H}_{\Theta(\mu)}$ into $\mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu$, this implies that

$$\mathbb{G} \rtimes_{\Gamma_\mu} \mathcal{H}_\mu = (\Gamma_\mu(\mathcal{H}_\mu) \cup (L_\infty(\hat{\mathbb{G}}) \otimes 1))'' = \tilde{\Gamma}_{\Theta(\mu)}((\mathcal{H}_\mu \cup L_\infty(\hat{\mathbb{G}}))'') \subseteq \tilde{\Gamma}_{\Theta(\mu)}(\mathcal{H}_{\Theta(\mu)}).$$

Conversely, let $x \in \mathcal{H}_{\Theta(\mu)}$. Then $\tilde{\Gamma}_{\Theta(\mu)}(x) \in \mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu$ and

$$\begin{aligned} \beta(\tilde{\Gamma}_{\Theta(\mu)}(x)) &= (\sigma V^* \sigma \otimes 1)((\chi \otimes \iota)(\iota \otimes \tilde{\Gamma})(\tilde{\Gamma}_{\Theta(\mu)}(x)))(\sigma V \sigma \otimes 1) \\ &= (\sigma V^* \sigma \otimes 1)((\sigma \otimes \iota)(\tilde{\Gamma} \otimes \iota)(\tilde{\Gamma}_{\Theta(\mu)}(x))(\sigma \otimes \iota))(\sigma V \sigma \otimes 1) \\ &= (\sigma V^* \otimes 1)((V \otimes 1)(\tilde{\Gamma}_{\Theta(\mu)}(x)_{13})(V^* \otimes 1))(V \sigma \otimes 1) \\ &= (\sigma \otimes 1)(\tilde{\Gamma}_{\Theta(\mu)}(x)_{13})(\sigma \otimes 1) = 1 \otimes \tilde{\Gamma}_{\Theta(\mu)}(x). \end{aligned}$$

Therefore, we have

$$\tilde{\Gamma}_{\Theta(\mu)}(x) \in (\mathcal{B}(L_2(\mathbb{G})) \overline{\otimes} \mathcal{H}_\mu)^\beta = \mathbb{G} \rtimes_{\Gamma_\mu} \mathcal{H}_\mu$$

for all $x \in \mathcal{H}_{\Theta(\mu)}$. This completes the proof. \square

The following result is an immediate consequence of Theorem 2.4.

Corollary 2.5. *The crossed product von Neumann algebra $\mathbb{G} \rtimes_{\Gamma_\mu} \mathcal{H}_\mu$ is injective.*

As pointed out by Izumi in [10] as a concrete example for the crossed product formula he obtained for countable discrete groups, in the case of random walks on the free group \mathbb{F}_n with respect to the uniform distribution on the generators, the result yields an identification of the Poisson boundary on the level of $\mathcal{B}(\ell_2(\mathbb{F}_n))$ ($n \geq 2$) with the Powers factor of type $III_{1/(2n-1)}$. Moreover, as noted in [10], the crossed product realization immediately implies amenability of the natural G -action on the Poisson boundary in the sense of Zimmer. The concept of an amenable action was first introduced by Zimmer [28] in the context of a measure class preserving action of a locally compact second countable group on a standard Borel space. It was subsequently shown that if G is a second countable locally compact group and $\alpha : G \curvearrowright X$ is a measure class preserving action of G on a standard probability space (X, ν) , then α is an amenable action if and only if the crossed product $G \rtimes_{\alpha} L_{\infty}(X, \nu)$ is injective. This was generalized to the case of actions of locally compact groups on von Neumann algebras by Anantharaman-Delaroche [1]. From this perspective, our result suggests that injectivity of the crossed product may provide a notion of ‘Zimmer amenability’ of a quantum group action. We note that a concept of (topologically) amenable actions for discrete quantum groups \mathbb{G} in the C^* -algebra framework was defined by Vaes–Vergnioux in [27]. It is shown in [27, Proposition 4.4] that topological amenability of the action on a unital nuclear C^* -algebra entails nuclearity of the (reduced) C^* -crossed product.

Our proof of Theorem 2.4 gives a simple way to obtain the following.

Proposition 2.6. *Let G be a discrete group. Then the following are equivalent:*

- (i) G is exact;
- (ii) for every probability measure μ on G , the reduced C^* -algebra crossed product $G \rtimes_{\Gamma_{\mu}} \mathcal{H}_{\mu}$ is nuclear;
- (iii) for some probability measure μ on G , the crossed product $G \rtimes_{\Gamma_{\mu}} \mathcal{H}_{\mu}$ is nuclear.

Proof. (i) \Rightarrow (ii): Denote $\mathcal{R}_{\mu} := \mathcal{E}_{\Theta(\mu)}(UC^*(G))$. Since $\mathcal{E}_{\Theta(\mu)}(a) = \mathcal{E}_{\mu}(a)$ for all $a \in \ell_{\infty}(G)$, and $\mathcal{E}_{\Theta(\mu)}(\hat{a}) = \hat{a}$ for all $\hat{a} \in C_r^*(G)$, it follows from the proof of Theorem 2.4 that

$$(2.6) \quad \mathcal{R}_{\mu} = \tilde{\Gamma}_{\Theta(\mu)}^{-1}(G \rtimes_{\Gamma_{\mu}} \mathcal{H}_{\mu}).$$

Moreover, we have $\mathcal{R}_{\mu} \subseteq UC^*(G)$, and hence $\mathcal{E}_{\Theta(\mu)}$ restricts to a unital completely positive idempotent on $UC^*(G)$. Since G is exact, the uniform Roe algebra $UC^*(G)$ is nuclear [21]. Now it is easy to see from the completely positive factorization property

$$\mathcal{R}_{\mu} \hookrightarrow UC^*(G) \rightarrow M_{n(\alpha)} \rightarrow UC^*(G) \rightarrow \mathcal{R}_{\mu}$$

that \mathcal{R}_{μ} (with its Choi–Effros product) is also nuclear. But by (2.6), $\tilde{\Gamma}_{\Theta(\mu)}$ induces a completely isometric order isomorphism of the operator systems $\mathcal{R}_{\mu} \cong G \rtimes_{\Gamma_{\mu}} \mathcal{H}_{\mu}$. Hence the latter is nuclear.

(ii) \Rightarrow (iii): Obvious.

(iii) \Rightarrow (i): This follows from the fact that $C_r^*(G) \subseteq G \rtimes_{\Gamma_{\mu}} \mathcal{H}_{\mu}$. □

Ozawa proved in [21] that a discrete group G is exact if and only if its action on the Stone–Ćech compactification βG is amenable; equivalently, G admits an amenable action on some compact space.

However, if G acts amenably on βG , of course it does not need to do so on an arbitrary compact space (indeed, amenability of the trivial action on a one-point set is equivalent to amenability of the group itself). Since βG is often too large to be useful in applications, it is interesting to find “smaller” amenable G -spaces, when they exist. This motivated the above result.

Finally, one may compare Proposition 2.6 with the following characterization of the amenability, due to Kaimanovich–Vershik [15] (see also [22] for non-discrete groups); note that the implication (ii) \Rightarrow (i) follows immediately from Proposition 2.6.

Theorem 2.7. *Let G be a countable discrete group. Then the following are equivalent:*

- (i) G is amenable;
- (ii) $\mathcal{H}_\mu = \mathbb{C}1$ for some probability measure μ on G .

We remark that requiring condition (ii) to hold for all adapted probability measures μ on G (i.e., such that the subgroup generated by the support of μ is dense in G) is a much stronger property than amenability, called the Liouville property; there exist even solvable groups without the Liouville property [15, Proposition 6.1].

REFERENCES

- [1] C. Anantharaman-Delaroche, *Action moyennable d’un groupe localement compact sur une algèbre de von Neumann*, Math. Scand. **45** (1979), no. 2, 289–304.
- [2] C.-H. Chu and A. T.-M. Lau, *Harmonic functions on groups and Fourier algebras*, Lecture Notes in Mathematics, **1782**, Springer-Verlag, Berlin, 2002.
- [3] E. G. Effros, J. Kraus and Z.-J. Ruan, *On two quantized tensor norms*, Operator Algebras, Mathematical Physics, and Low Dimensional Topology (Istanbul 1991), Res. Notes Math. 5, A K Peters, Wellesley, MA 1993, 125–145.
- [4] E. G. Effros and Z.-J. Ruan, *Operator spaces*, London Math. Soc. Monographs, New Series **23**, Oxford University Press, New York, 2000.
- [5] M. Enock, *Measured quantum groupoids in action*, Mém. Soc. Math. Fr. (N.S.), no. 114 (2008).
- [6] H. Furstenberg, *Boundary theory and stochastic processes on homogeneous spaces*, Harmonic analysis on homogeneous spaces (Proc. Sympos. Pure Math., Vol. XXVI, Williams Coll., Williamstown, Mass., 1972), Amer. Math. Soc., Providence, R.I., 1973, 193–229.
- [7] F. Ghahramani, *Isometric representation of $M(G)$ on $B(H)$* , Glasgow Math. J. **23** (1982), 119–122.
- [8] U. Haagerup and J. Kraus, *Approximation properties for group C^* -algebras and group von Neumann algebras*, Trans. Amer. Math. Soc. **344** (1994), 667–699.
- [9] M. Izumi, *Non-commutative Poisson boundaries and compact quantum group actions*, Adv. Math. **169** (2002), no. 1, 1–57.
- [10] M. Izumi, *Non-commutative Poisson boundaries*, in: *Discrete geometric analysis*, Contemp. Math., 347, Amer. Math. Soc., Providence, RI, 2004, 69–81.
- [11] M. Izumi, *E_0 -semigroups: around and beyond Arveson’s work*, J. Operator Theory **68** (2012), no. 2, 335–363.
- [12] M. Izumi, S. Neshveyev and L. Tuset, *Poisson boundary of the dual of $SU_q(n)$* , Comm. Math. Phys. **262** (2006), no. 2, 505–531.
- [13] W. Jaworski and M. Neufang, *The Choquet–Deny equation in a Banach space*, Canad. J. Math. **59** (2007), no. 4, 795–827.

- [14] M. Junge, M. Neufang and Z.-J. Ruan, *A representation theorem for locally compact quantum groups*, Int. J. Math. **20** (2009), 377–400.
- [15] V. A. Kaimanovich and A. M. Vershik, *Random walks on discrete groups: boundary and entropy*, Ann. Probability, **11** (1983), 457–490.
- [16] J. Kustermans, *Locally compact quantum groups in the universal setting*, International J. Math. **12** (2001), 289–338.
- [17] J. Kustermans and S. Vaes, *Locally compact quantum groups*, Ann. Sci. Ecole Norm. Sup. **33** (2000), 837–934.
- [18] J. Kustermans and S. Vaes, *Locally compact quantum groups in the von Neumann algebraic setting*, Math. Scand. **92** (2003), 68–92.
- [19] M. Neufang, Z.-J. Ruan and N. Spronk, *Completely isometric representations of $M_{cb}A(G)$ and $UCB(\hat{G})^*$* , Trans. Amer. Math. Soc. **360** (2008), 1133–1161.
- [20] M. Neufang and V. Runde, *Harmonic operators: the dual perspective*, Math. Z. **255** (2007), 669–690.
- [21] N. Ozawa, *Amenable actions and exactness for discrete groups*, C. R. Acad. Sci. Paris Sr. I Math. **330** (2000), no. 8, 691–695.
- [22] J. Rosenblatt, *Ergodic and mixing random walks on locally compact groups*, Math. Ann. **257** (1981), no. 1, 31–42.
- [23] Z.-J. Ruan, *On the predual of dual algebras*, J. Operator Theory **27** (1992), 179–192.
- [24] E. Størmer, *Regular abelian Banach algebras of linear maps of operator algebras*, J. Funct. Anal. **37** (1980), 331–373.
- [25] R. Tomatsu, *A characterization of right coideals of quotient type and its application to classification of Poisson boundaries*, Comm. Math. Phys. **275** (2007), no. 1, 271–296.
- [26] S. Vaes and N. Vander Venet, *Poisson boundary of the discrete quantum group $\widehat{A_u(F)}$* , Compos. Math. **146** (2010), no. 4, 1073–1095.
- [27] S. Vaes and R. Vergnioux, *The boundary of universal discrete quantum groups, exactness, and factoriality*, Duke Math. J. **140** (2007), no. 1, 35–84.
- [28] R. Zimmer, *Amenable ergodic group actions and an application to Poisson boundaries of Random walks*, J. Funct. Anal. **27** (1978), 350–372.

SCHOOL OF MATHEMATICS AND STATISTICS, CARLETON UNIVERSITY, OTTAWA, ONTARIO, CANADA K1S 5B6

E-mail address: `mkalanta@math.carleton.ca`

SCHOOL OF MATHEMATICS AND STATISTICS, CARLETON UNIVERSITY, OTTAWA, ONTARIO, CANADA K1S 5B6

E-mail address: `mneufang@math.carleton.ca`

UNIVERSITÉ LILLE 1 - SCIENCES ET TECHNOLOGIES, U.F.R. DE MATHÉMATIQUES, LABORATOIRE DE MATHÉMATIQUES PAUL PAINLEVÉ - UMR CNRS 8524, 59655 VILLENEUVE D'ASCQ CÉDEX, FRANCE

E-mail address: `matthias.neufang@math.univ-lille1.fr`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA, IL 61801, USA

E-mail address: `ruan@math.uiuc.edu`