

# An efficient curing policy for epidemics on graphs\*

Kimon Drakopoulos, Asuman Ozdaglar, John N. Tsitsiklis

Laboratory of Information and Decision Systems, Massachusetts Institute of Technology, Cambridge, MA 02139,  
{kimondr,asuman,jnt}@mit.edu.

August 21, 2018

We provide a dynamic policy for the rapid containment of a contagion process modeled as an SIS epidemic on a bounded degree undirected graph with  $n$  nodes. We show that if the budget  $r$  of curing resources available at each time is  $\Omega(W)$ , where  $W$  is the CutWidth of the graph, and also of order  $\Omega(\log n)$ , then the expected time until the extinction of the epidemic is of order  $O(n/r)$ , which is within a constant factor from optimal, as well as sublinear in the number of nodes. Furthermore, if the CutWidth increases only sublinearly with  $n$ , a sublinear expected time to extinction is possible with a sublinearly increasing budget  $r$ .

**1. Introduction** Many contagion processes over large networks can lead to costly cascades unless controlled by outside intervention. Examples include epidemics spreading over a population of individuals, viruses attacking a network of connected computers, or financial contagion in a network of banks. In this paper we study how this type of contagion can be prevented or contained by dynamically curing some of the infected nodes under a budget constraint on the amount of curing resources that can be deployed at each time.

More specifically, we consider a canonical SIS epidemic model on an undirected graph<sup>1</sup> with  $n$  nodes, with a common infection rate along any edge that connects an infected and a healthy node, and node-specific curing rates  $\rho_v(t)$  at each node  $v$ . The curing rates are to be chosen according to a curing policy which is based on the past history of the process and the network structure, subject to an upper bound on the total curing rate  $\sum_v \rho_v(t)$ .

In a companion paper, [6] we characterize the cases for which a contagion process can be rapidly contained, i.e., the expected time to extinction can be made sublinear in the number  $n$  of nodes using a sublinear curing budget. Our characterization involves the CutWidth of the underlying graph. Intuitively, the CutWidth measures the required budget of curing resources in a simpler deterministic curing problem, in which infected nodes are cured one at a time, subject to the constraint that the number of edges between healthy and infected nodes is at all times less than or equal to the budget of curing resources. In [6], we establish that if the CutWidth increases at least linearly with  $n$ , then a sublinear (in  $n$ ) expected time to extinction is impossible with a sublinear budget  $r$ . On the other hand, [6] provides a *nonconstructive proof* that for graphs with sublinear CutWidth and bounded degree, there exists a dynamic policy that achieves sublinear expected time to extinction using only a sublinear budget, for any set of initially infected nodes. The main contribution of the present paper is the *construction of a specific* policy with the latter desirable properties.

Our policy is based on a combinatorial result which states the following. Given an initial set of infected nodes, nodes can be removed from that set, one at a time, in way that the maximum cut (number of edges) between healthy and infected nodes encountered during this process is upper bounded by the sum of the CutWidth of the graph and the cut associated with the initial set. Let

\*Research partially supported by the Draper Laboratories and NSF grant CMMI-1234062. A preliminary version of some of the results in this paper are included in a paper submitted to the 53rd IEEE Conference on Decision and Control, December 2014.

<sup>1</sup>Our results actually are easily generalized to the case of directed graphs.

us refer to the sequence of subsets encountered during this process as a *target path*. The main idea underlying our policy is to allocate the entire curing budget to appropriate nodes so that we stay most of the time, with high probability, on or near the target path. We show that this is indeed possible, as long as the curing resource budget scales in proportion to the CutWidth. We also show that the policy is optimal (within a multiplicative constant) if the available budget is also  $\Omega(\log n)$ .<sup>2</sup>

A similar model, but in which the curing rate allocation is done statically (open-loop) has been studied in [5, 7, 3, 11], but the proposed methods were either heuristic or based on mean-field approximations of the evolution process. Closer to our work, the authors of [2] let the curing rates be proportional to the degree of each node — independent of the current state of the network, which means that curing resources may be wasted on healthy nodes. On a graph with bounded degree, the policy in [2] achieves sublinear time to extinction, but requires a curing budget that is proportional to the number of nodes. In contrast, our policy achieves the same performance (sublinear time to extinction) for all bounded degree graphs with small CutWidth by properly focusing the curing resources. As an extreme example, consider a line graph with  $n$  nodes, and assume that the  $n/2$  leftmost nodes are initially infected. The degree-based policy of [2] requires a total budget proportional to  $n$  and allocates it proportional to the degree. In contrast, our policy can achieve sublinear expected time to extinction with a  $\Omega(\log n)$  but sublinear budget. This is because, instead of allocating the available budget to all nodes, our policy focuses on specific nodes on the boundary between healthy and infected nodes, in this instance on the rightmost infected node. By extending this idea, our policy achieves a similar improvement for all graphs with sublinear CutWidth.

The rest of the paper is organized as follows. In Section 2 we present the details of the model that we are considering. In Section 3 we introduce the CutWidth and establish the combinatorial result mentioned earlier. In Section 4 we present the policy and analyze its performance. In Section 5 we develop some corollaries that demonstrate the possibility of fast extinction using a sublinear budget and the approximate optimality of our policy in a certain regime. We also mention some examples. Finally, in Section 6 we offer some closing remarks.

**2. The Model** We consider a network, represented by a connected undirected graph  $G = (V, E)$ , where  $V$  denotes the set of nodes and  $E$  denotes the set of edges. We use  $n$  to denote the number of nodes. Two nodes  $u, v \in V$  are *neighbors* if  $(u, v) \in E$ . We denote by  $\Delta$  the maximum of the node degrees.

We assume that the nodes in a set  $I_0 \subseteq V$  are initially infected and that the infection spreads according to a controlled contact process where the rate at which infected nodes get cured is determined by a network controller. Specifically, each node can be in one of two states: *infected* or *healthy*. The controlled contact process — also known as the SIS epidemic model — on  $G$  is a right-continuous, continuous-time Markov process  $\{I_t\}_{t \geq 0}$  on the state space  $\{0, 1\}^V$ , where  $I_t$  stands for the set of infected nodes at time  $t$ . We refer to  $I_t$  as the *infection process*.

State transitions at each node occur independently according to the following dynamics.

- a) The process is initialized at the given initial state  $I_0$ .
- b) If a node  $v$  is healthy, i.e., if  $v \notin I_t$ , the transition rate associated with a change of the state of that node to being infected is equal to an infection rate  $\beta$  times the number of infected neighbors of  $v$ , that is,

$$\beta \cdot |\{(u, v) \in E : u \in I_t\}|,$$

where we use  $|\cdot|$  to denote the cardinality of a set. By rescaling time, we can and will assume throughout the paper that  $\beta = 1$ .

<sup>2</sup> We write  $f(n) = o(g(n))$  if  $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$ . We write  $f(n) = \Omega(g(n))$  if  $\liminf_{n \rightarrow \infty} f(n)/g(n) > 0$ . Finally, we write  $f(n) = O(g(n))$  if  $\limsup_{n \rightarrow \infty} f(n)/g(n) < \infty$ .

- c) If a node  $v$  is infected, i.e., if  $v \in I_t$ , the transition rate associated with a change of the state of that node to being healthy is equal to a curing rate  $\rho_v(t)$  that is determined by the network controller, as a function of the current and past states of the process. We are assuming here that the network controller has access to the entire past evolution of the process.

We assume a *budget constraint* of the form

$$\sum_{v \in V} \rho_v(t) \leq r, \quad (1)$$

for each time instant  $t$ , reflecting the fact that curing is costly. A *curing policy* is a mapping which at any time  $t$  maps the past history of the process to a curing vector  $\rho(t) = \{\rho_v(t)\}_{v \in V}$  that satisfies (1).

We define the *time to extinction* as the time until the process reaches the absorbing state where all nodes are healthy:

$$\tau = \min\{t \geq 0 : I_t = \emptyset\}.$$

The *expected time to extinction* (the expected value of  $\tau$ ) is the performance measure that we will be focusing on.

**3. Graph theoretic preliminaries** In this section we introduce the notions of a cut and of the CutWidth that will be used in the description of our policy. We state some of their properties and then proceed to develop a key combinatorial result that will play a critical role in the analysis of our policy's performance. Throughout, we assume that we are dealing with a particular given graph  $G$ .

**3.1. CutWidth** For convenience, we will be using the shorthand term “bag” to refer to “a subset of  $V$ .” We also use the following notation. For any two bags  $A$  and  $B$ , and any  $v \in V$ , we let

$$A \setminus B = \{v \in A : v \notin B\},$$

and

$$A - v = A \setminus \{v\}.$$

We also use  $A^c$  to denote the complement,  $V \setminus A$  of  $A$ .

We next define the concept of a *monotone crusade*. A monotone crusade from a bag  $A$  to another bag  $B$ , where  $B \subseteq A$ , is a finite sequence of bags that starts with  $A$  and ends with  $B$ , so that at each step of the sequence no nodes are added (cf. Part (iii) of Definition 1), and exactly one node is removed (cf. Part (iv) of Definition d:crus).

**DEFINITION 1.** For any two bags  $A$  and  $B$ , with  $B \subseteq A$ , a (monotone) crusade from  $A$  to  $B$ , or  $(A \downarrow B)$ -**crusade** for short, is a sequence  $\omega = (\omega_0, \omega_1, \dots, \omega_k)$  of bags of length  $|\omega| = k+1$ , with the following properties:

- (i)  $\omega_0 = A$ ,
- (ii)  $\omega_k = B$ ,
- (iii)  $\omega_{i+1} \subset \omega_i$ , for  $i = 0, 1, \dots, k-1$ , and
- (iv)  $|\omega_i \setminus \omega_{i+1}| = 1$ , for  $i = 0, 1, \dots, k-1$ .

We denote by  $\mathcal{C}(A \downarrow B)$  the set of all  $(A \downarrow B)$ -crusades.

The number of edges connecting a bag  $A$  with its complement is called the cut of the bag. Its importance lies in that it is equal to the total rate at which new infections occur, when the set of currently infected nodes is  $A$ .

**DEFINITION 2.** For any bag  $A$ , its **cut**,  $c(A)$ , is defined as the cardinality of the set of edges

$$\{(u, v) : u \in A, v \in A^c\}.$$

In Proposition 1 below, we record, without proof, an elementary property of cuts.

PROPOSITION 1. *For any two bags  $A$  and  $B$ , we have*

$$c(A \cup B) \leq c(A) + c(B) \leq c(A) + \Delta \cdot |B|.$$

We define the width of a monotone crusade  $\omega$  as the maximum cut encountered during the crusade. Intuitively, this is the largest infection rate to be encountered if the nodes were to be cured according to the sequence prescribed by the crusade deterministically, if no new infections happen in between.

DEFINITION 3. Given an  $(A \downarrow B)$ -crusade  $\omega = (\omega_0, \dots, \omega_k)$ , its **width**  $z(\omega)$  is defined by

$$z(\omega) = \max_{0 \leq i \leq k} \{c(\omega_i)\}.$$

We next define what we call the impedance of a bag  $A$ , as the minimum possible width among the  $\mathcal{C}(A \downarrow \emptyset)$ -crusades. This minimization captures the objective of finding a crusade along which the total infection rate is always small.

DEFINITION 4. The **impedance**  $\delta(A)$  of a bag  $A$  is defined by

$$\delta(A) \doteq \min_{\omega \in \mathcal{C}(A \downarrow \emptyset)} z(\omega). \quad (2)$$

For the special case where  $A = V$ , the impedance is known as the **CutWidth** [10, 1, 4, 8], and will be denoted by  $W$ .

We say that a (monotone) crusade  $(A \downarrow B)$ -crusade  $\omega = (\omega_0, \dots, \omega_{k-1})$  is *optimal* if it attains the minimum in Eq. (2). It can be seen that the impedances satisfy the Bellman equation:

$$\delta(A) = \max \{c(A), \min\{\delta(B) : B \subseteq A, |A \setminus B| = 1\}\}. \quad (3)$$

Furthermore, along an optimal crusade, we have  $\delta(\omega_{i+1}) \leq \delta(\omega_i)$ , for  $i = 0, 1, \dots, k-1$ . Finally, we note that  $c(A) \leq \delta(A)$ .

**3.2. Impedance and CutWidth** In this subsection we discuss the relation between the impedance of an arbitrary bag and the CutWidth. The impedance of a bag  $A$  is at least  $c(A)$ , which in general may be much larger than the CutWidth.<sup>3</sup> This is a concern because the stochastic nature of the infections can always bring the process to a bag with high impedance, and therefore high subsequent infection rates. The next lemma provides an upper bound on the impedance of a bag  $A$  in terms of the CutWidth  $W$  of the graph and the cut of  $A$ . Its proof is given in the Appendix.

LEMMA 1. *For any bag  $A$ , we have*

$$\delta(A) \leq W + c(A).$$

**4. The CURE policy** In this section, we present our curing policy and we study the resulting expected time to extinction, starting from an arbitrary initial set of infected nodes. Loosely speaking, the policy, at any time, tries to follow a certain desirable (monotone) crusade, called a target path, by allocating all of the curing resources to a single node, namely, the node that should be removed in order to obtain the next bag along the target path. On the other hand, this ideal scenario may be interrupted by infections, at which point the policy shifts its attention to newly infected nodes, and attempts to return to a bag on the target path. It turns out that under certain

<sup>3</sup> As an example, consider a line graph, and let  $A$  be the set of even-numbered nodes. Then,  $c(A)$  is approximately  $n$ , whereas the CutWidth of the line graph is equal to 1.

assumptions, this is successful with high probability and does not take too much time. However, with small probability, the process veers far off from the target path; in that case the policy “restarts” in a manner that we will make precise in the sequel.

It is quite intuitive (and formally established in [6]) that a fast (sublinear) time to extinction may not be possible if the curing budget is smaller than the CutWidth. For this reason, we focus on the regime where the curing rate is at least proportional to the CutWidth, and more concretely, on the regime where  $r \geq 4W$ , which we henceforth assume.

Under the above assumptions on the budget  $r$ , and the additional assumptions that  $r = \Omega(\log n)$  and  $r \geq 8\Delta$ , we will construct a policy whose expected time to extinction is  $O(n/r)$ ; cf. Theorem 1 and Corollary 1.

The CURE policy is defined hierarchically: it consists of a sequence of attempts; each attempt consists of a waiting period, followed by a sequence of segments; finally, each segment consists of a path-following phase and an excursion. Each attempt, at the end of its waiting period, determines a path that its segments will try to follow. If an attempt fails to cure all nodes, a new attempt is initiated. We now proceed to describe in detail a typical attempt.

**Waiting period.** A typical attempt starts at some bag  $A$ , with a waiting period. (If this is the first attempt, then  $A = I_0$ . Otherwise, it is the bag at the end of the preceding attempt.) During the waiting period, all curing rates  $\rho_v(t)$  are kept at zero. The waiting period ends at the first subsequent time that<sup>4</sup>

$$c(I_t) \leq r/8.$$

Let  $B$  be the bag  $I_t$  right at the end of the waiting period, and let  $\omega^B = (\omega_0^B, \dots, \omega_{|B|}^B)$  the corresponding optimal crusade, which we refer to as the *target path*.

**Segments.** Each segment of an attempt starts either at the end of the waiting period or at the end of a preceding segment of the same attempt. In all cases, the segment starts with a bag on the target path. For the first segment, this is guaranteed by the definition of the target path. For subsequent segments, it will be guaranteed by our specifications of what happens at the end of the preceding segment.

**Path-following phase.** Let  $v_1, \dots, v_m$  be the nodes in the bag at the beginning of a segment, arranged in the order according to which they are to be removed along the target path. For example, the bag at the beginning of the segment is  $\{v_1, \dots, v_m\}$ , the next bag is  $\{v_2, \dots, v_m\}$ , etc. During the path-following phase, the entire curing budget is first allocated to node  $v_1$  until it gets cured, then to node  $v_2$ , etc. The phase ends when either:

(i) all nodes have been cured, i.e.,  $I_t = \emptyset$ ; in this case, the attempt is considered successful and the process is over.

(ii) an infection occurs; in this case,  $I_t$  is outside the target path, and the segment continues with an excursion phase.

**Excursion.** If a path-following phase ended with an infection, the policy enters an excursion phase. Let  $C$  be the next bag (on the target path), namely the bag that would have been reached if a node was cured before the infection happened. We define  $D_t = I_t \setminus C$ ; this is the set of infected nodes that do not belong to the next bag on the target path. The goal during the excursion is to cure the nodes in  $D_t$ , so as to reach the bag  $C$  on the target path, and then start a subsequent segment. With this purpose in mind, during an excursion, we allocate all the available budget on an arbitrary node in  $D_t$ . The excursion ends when  $|D_t|$  becomes either zero or at least  $r/8\Delta$ .

(i) If the excursion ends with  $|D_t| = 0$ , we say that we have a *short* excursion. At that time, we are back on the target path, with  $I_t = C$ , and we are ready to start with the next segment.

<sup>4</sup> Note that the waiting period is guaranteed to terminate in finite time, with probability 1. This is because if it were infinite, then healthy nodes would keep getting infected until eventually  $I_t = V$ . But  $c(V) = 0$ , which means that at some point the condition  $c(I_t) \leq r/8$  would be satisfied and the waiting period would be finite, a contradiction.

(ii) If on the other hand  $|D_t| \geq r/8\Delta$ , we say that the excursion was *long*, and that the attempt has failed. In this case, the attempt has no more segments, and a new attempt will be initiated, starting with a waiting period.

**4.2. Performance analysis — Outline** We now proceed to establish an upper bound on the expected time to extinction, under the assumption that  $r \geq 4W$ , for any set of initially infected nodes. If the process always stayed on the target path, that is if we had no infections, the expected time to extinction would be the time until all nodes (at most  $n$  of them) were cured. Given that nodes are cured at a rate of  $r$ , the expected time to extinction would have been  $O(n/r)$ . On the other hand, infections do delay the curing process, by initiating excursions, and we need to show that these do not have a major impact.

There are two kinds of excursions to consider, *short* ones, at the end of which  $|D_t| = 0$ , and *long* ones, at the end of which  $|D_t| \geq r/8\Delta$ . During an excursion, the size of  $D_t$  (the “distance” from the target path) is at most  $r/8\Delta$ . Using also an upper bound on the size of the cut along the target path, we can show that the infection rate during an excursion is smaller than the curing rate. For this reason, during an excursion, the process  $|D_t|$  has a downward drift. As a consequence, using a standard argument, the expected duration of an excursion is small and there is high probability that the excursion ends with  $|D_t| = 0$ , so that the excursion is short and we continue with the next segment. As a result, the expected duration of an attempt behaves similar to the case of no excursions and is also of order  $O(n/r)$ . Finally, by viewing each attempt as an independent trial, we can establish an upper bound for the overall policy. A formal version of this argument is the content of the rest of this section.

**4.3. Excursion analysis** Let us focus on a particular excursion, and let, for simplicity,  $M_t = |D_t|$ . The process  $M_t$  evolves on the finite set  $\{0, 1, \dots, r/8\Delta\}$ . (For simplicity, and without loss of generality, we assume that  $r/8\Delta$  is an integer.) Recall that  $C$  was defined as the bag on the target path that we were trying to reach at the end of the excursion. The difference  $D_t$  at the time that the excursion starts consists of exactly two nodes: the node that we were trying to remove just before the excursion started and the node outside the target path that got infected. Thus, the process  $M_t$  is initialized at 2, at the beginning of the excursion. The process  $M_t$  is stopped as soon one of the two boundary points, 0 or  $r/8\Delta$ , is reached. At each time before the process is stopped, there is a rate equal to  $r$  of downward transitions. Furthermore, there is a rate  $c(I_t)$  of upward transitions, corresponding to new infections.

LEMMA 2. *The rate  $c(I_t)$  of upward transitions during an excursion satisfies  $c(I_t) \leq r/2$ .*

*Proof:* The definition  $D_t = I_t \setminus C$  implies that  $I_t \subseteq C \cup D_t$ . Consequently,

$$\begin{aligned} c(I_t) &\leq c(C) + c(D_t) \leq c(C) + \Delta \cdot |D_t| \\ &= c(C) + \Delta \cdot M_t \leq c(C) + \frac{r}{8}. \end{aligned} \tag{4}$$

We have used here Proposition 1, in the first and second inequality, together with the fact  $M_t \leq r/8\Delta$ .

On the other hand,  $C$  is on the target path associated with  $B$ , the bag obtained at the end of the waiting period. As remarked at the end of Section 3.1, the impedance does not increase along an optimal crusade, and therefore,  $\delta(C) \leq \delta(B)$ . Using also Lemma 1, we have

$$\delta(C) \leq \delta(B) \leq W + c(B).$$

Recall now that a waiting period ends with a bag whose cut is at most  $r/8$ . Therefore,  $c(B) \leq r/8$ . It follows that  $c(C) \leq W + r/8$ . Using this fact, together with the assumption  $r \geq 4W$  and Eq. (4), we obtain

$$c(I_t) \leq c(C) + \frac{r}{8} \leq \left(W + \frac{r}{8}\right) + \frac{r}{8} \leq \frac{r}{4} + \frac{r}{8} + \frac{r}{8} = \frac{r}{2}.$$

We now establish the properties of the excursions that we have claimed; namely, that excursions are short, with high probability, and do not last too long.

LEMMA 3. a) *The probability that the excursion is long is at most*

$$p = \frac{2^2 - 1}{2^{r/8\Delta} - 1}.$$

b) *The expected length of an excursion is upper bounded by  $4/r$ .*

*Proof:* a) Using Lemma 2, the process  $M_t$  is stochastically dominated by a process  $N_t$  on the same space  $\{0, 1, \dots, r/8\Delta\}$ , which is initialized to be equal to the value of  $M_t$  at the beginning of the excursion (which is 2), has a rate  $r$  of downward transitions, a rate  $r/2$  of upward transitions, and stops at the first time that it reaches one of the two boundary values. Note that the ratio of the downward to the upward drift is equal to 2. The probability, denoted by  $p$ , that the process  $N_t$  will first reach the upper boundary is a well-studied quantity and is given by the expression in part (a) of the lemma. The proof is standard and can be found in Section 2.1 of [9] (for a non-martingale based proof) or Section 2.3 of [12] (for a martingale based proof). Since  $M_t$  is stochastically dominated by  $N_t$ , the probability that  $M_t$  will first reach the upper boundary is no larger.

b) For simplicity, let us suppose that the excursion starts at time  $t = 0$ . We define the process

$$H_t = M_t + \frac{r}{2}t$$

and the stopped version,  $\hat{H}_t$  which stops at the time  $T$  that the excursion ends. It is straightforward to verify that  $\hat{H}_t$  is a supermartingale, because the upward drift of the process is  $\beta c(I_t) \leq r/2$  and the downward drift is  $r$ , so that the total downward drift is at least  $r/2$ . Furthermore,  $H_0 = M_0 = 2$ . Using Doob's optional stopping theorem we obtain

$$2 = \mathbb{E}[M_0] = \mathbb{E}[H_0] \geq \mathbb{E}[H_T] + \frac{r}{2} \cdot T \geq \frac{r}{2} \cdot T,$$

from which we conclude that

$$\mathbb{E}[T] \leq \frac{4}{r}.$$

Note that if  $r \geq \alpha \log n$ , where  $\alpha$  is a sufficiently large constant, then  $p$  can be made smaller than  $1/n^2$ , so that  $np$  tends to zero. We will be using this observation later on. We will now bound the length of a waiting period.

LEMMA 4. *The expected length of a waiting period is bounded above by  $8n/r$ .*

*Proof:* A waiting period involves at most  $n$  infections. The waiting period ends as soon as  $c(I_t) \leq r/8$ . Therefore, during the waiting period, infections happen at a rate of at least  $r/8$ . In particular, during the waiting period, the expected time between consecutive infections is at most  $8/r$ . For a maximum of  $n$  infections, the expected time is upper bounded by  $8n/r$ .

We can now combine the various bounds we have derived so far in order to bound the expected time to extinction under our policy.

THEOREM 1. *Suppose that  $r \geq 4W$  and that  $r$  is large enough so that  $np < 1$ , where  $p$  is as defined in Lemma 3. For any initial bag, the expected time to extinction under the CURE policy is upper bounded by*

$$\frac{1}{1 - np} \cdot \frac{13n}{r}.$$

*Proof:* We start by upper bounding the expected duration of an attempt. The length of the waiting period of an attempt is upper bounded by  $8n/r$ , by Lemma 4.

We now consider the total length of the path-following phases of an attempt. At a typical time in the path-following phase, the current bag is on the target path, of the form  $\{v_k, \dots, v_m\}$ , where we are using the notation introduced in Section 4.1. At some point, a transition occurs and either:

(i) the new bag is  $\{v_{k+1}, \dots, v_m\}$  and the path-following phase continues, or

(ii) an infection occurs; in this case, and unless the attempt fails, the next path-following phase will again start from the bag  $\{v_{k+1}, \dots, v_m\}$ .

In both cases, the expected time spent in the path-following phase until we move to the next bag on the target path is at most  $1/r$ . Since the target path has length at most  $n$ , the expected total duration of the path-following phases of an attempt is at most  $n/r$ .

Similarly, the number of excursions during an attempt is at most  $n$ . By Lemma 3, the expected length of an excursion is at most  $4/r$ . Therefore, the expected total time spent on excursions, during the same attempt, is upper bounded by  $4n/r$ .

Putting everything together, the expected duration of an attempt is at most  $(8n/r) + (n/r) + (4n/r) = 13n/r$ .

Each attempt involves  $n$  segments. During each segment, there is probability at most  $p$  that the excursion is long and that the attempt fails. Therefore, the overall probability that the attempt will fail is at most  $np$ . Given that the process regenerates at the beginning of each attempt, the expected number of attempts is at most  $1/(1 - np)$ , and the desired result follows.

**5. Corollaries and near-optimality of the CURE policy** Theorem 1 has a number of interesting consequences, which we collect in the corollary that follows. We argue that if all nodes are initially infected, then the expected time to extinction under any policy is at least  $n/r$ . Furthermore, in a certain regime of parameters, our policy achieves  $O(n/r)$  expected time to extinction and is therefore optimal within a multiplicative constant. Finally, if the CutWidth increases sublinearly with the number of nodes, then the expected time to extinction can be made sublinear in  $n$ , using only a sublinear budget. This last result is also proved in [6], using a different, nonconstructive argument.

**COROLLARY 1.** *a) For any graph with  $n$  nodes and with all nodes initially infected, the expected time to extinction is at least  $n/r$ , under any policy.*

*b) Suppose that the budget  $r$  satisfies*

$$r \geq 4W, \quad r \geq 16 \cdot \log_2 n \cdot \Delta.$$

*Then, for large enough  $n$ , and for any initial set of infected nodes, the expected time to extinction under the CURE policy is at most  $26n/r$ , which is sublinear in  $n$ .*

*c) Suppose that the budget  $r$  satisfies the conditions in part (b), together with the condition*

$$r = \Omega(n/\log n).$$

*Then, the expected time to extinction under the CURE policy is of order  $O(\log n)$ .*

*d) If the CutWidth increases sublinearly with  $n$ , then it is possible to have sublinear time to extinction with a sublinear budget.*

*Proof:* a) Since nodes are cured at a rate of at most  $r$ , and there are  $n$  nodes to be cured, the expected time to extinction must be at least  $n/r$ , even in the absence of infections.

b) When  $r \geq 16 \cdot \log_2 n \cdot \Delta$ , we have  $r/8\Delta \geq 2 \log_2 n$ , and  $2^{r/8\Delta} \geq n^2$ . Thus, the probability  $p$  in Lemma 3 is of order  $O(1/n^2)$ , and  $np$  is of order  $O(1/n)$ . In particular, for large enough  $n$ , the factor  $1/(1 - np)$  is less than 2. By Theorem 1, the expected time to extinction is at most  $26n/r$ . This is sublinear in  $n$ , because  $r$  tends to infinity.

- c) This is an immediate consequence of part (b).  
d) If  $W$  increase sublinearly with  $n$ , we can satisfy the conditions in parts (b) and (c) while keeping  $r$  sublinear in  $n$ , and still achieve sublinear, e.g.,  $O(\log n)$  expected time to extinction.

We continue by mentioning some examples. For a line graph with  $n$  nodes, the CutWidth is equal to 1 and  $\Delta = 2$ . Therefore, by part (b) of Corollary 1 we can guarantee an approximately optimal expected time to extinction, of order  $O(n/r)$ , as long as  $r \geq 16 \cdot \log_2 n \cdot \Delta = 32 \log_2 n$  in part (b) of Corollary. We note, however, that for this example, our analysis is not tight, and the requirement  $r \geq 32 \log_2 n$  is stronger than necessary.

For a square grid-graph with  $n$  nodes, the Cut-Width is approximately  $\sqrt{n}$  and  $\Delta = 4$ . In this case, the requirement  $r \geq 4W \approx 4\sqrt{n}$  is the dominant one, and suffices to guarantee an approximately optimal expected time to extinction, of order  $O(n/r)$ .

In both of these examples, we can of course let  $r$  be much larger than the minimum required, namely,  $O(\log n)$  and  $O(\sqrt{n})$ , respectively, in order to obtain a smaller expected time to extinction, e.g., the  $O(\log n)$  expected time to extinction in part (c) of the corollary.

**6. Conclusions** We have presented a dynamic curing policy with desirable performance characteristics. For example, if the CutWidth is sublinear in the number of nodes, the policy achieves sublinear expected time to extinction, using a sublinear curing budget. This policy applies to any subset of initially infected nodes and the resulting expected time to extinction is order-optimal when the available budget is sufficiently large.

Our analysis brings up a number of open problems of both practical and theoretical interest. Specifically, a major drawback of the CURE policy is computational complexity because calculating the impedance of a bag or finding a target path is computationally hard. Therefore, one possible direction is the design of computationally efficient policies with some performance guarantees, perhaps for special cases.

Alternatively, In the same spirit, certain combinatorial optimization tools have been developed for the approximation of the CutWidth of a graph. Similar tools can perhaps be employed to approximate the impedance of a bag. An interesting direction is the analysis of the performance of the CURE policy when instead of optimal crusades, approximately optimal crusades are used.

Finally, we have argued in this paper that the CURE policy is efficient in the sense of attaining near-optimal,  $O(n/r)$  expected time to extinction, in a certain parameter regime. It is an interesting problem to look for approximately optimal policies over a wider set of regimes, as well as for the case where the initial set of infected nodes is small.

## Appendix. Proof of Lemma 1

Consider a monotone crusade  $\omega \in \mathcal{C}(V \downarrow \emptyset)$  whose width is equal to the CutWidth  $W$ . This crusade starts with  $V$  and removes nodes one at a time, until the empty set is obtained. Let  $v_1, v_2, \dots, v_n$  be the nodes in  $V$ , arranged in the order in which they are removed.

Let us now fix a bag  $A$ . We construct a monotone crusade  $\hat{\omega} \in \mathcal{C}(A \downarrow \emptyset)$  as follows. We start with  $A$  and remove its nodes one at a time, according to the order prescribed by  $\omega$ . For example, if  $n = 4$ , and  $A = \{v_2, v_4\}$ , the monotone crusade that starts from  $A$  first removes node  $v_2$  and then removes node  $v_4$ .

At any intermediate step during the crusade  $\hat{\omega}$ , the current bag is of the form  $A \cap \{v_k, \dots, v_n\}$ , for some  $k$ . It only remains to show that the cut of this bag is upper bounded by  $c(A) + W$ . Let  $R = \{v_1, \dots, v_{k-1}\}$ . Note that

$$c(R) \leq W,$$

because of the definition of the width and the assumption that the width of  $\omega$  is  $W$ . Note also that the current bag is simply  $A \cap R^c$ .

For any two sets  $S_1$  and  $S_2$ , let  $e(S_1, S_2)$  be the number of edges that join them. We have that

$$\begin{aligned}
 c(A \cap R^c) &= e(A \cap R^c, (A \cap R^c)^c) \\
 &= e(A \cap R^c, A^c \cup R) \\
 &\leq e(A \cap R^c, A^c) + e(A \cap R^c, R) \\
 &\leq e(A, A^c) + e(R^c, R) \\
 &= c(A) + c(R) \\
 &\leq c(A) + W.
 \end{aligned}$$

We conclude that the cut associated with any intermediate bag in the crusade  $\hat{\omega}$  is upper bounded by  $c(A) + W$ . It follows that the width of  $\hat{\omega}$ , and therefore  $\delta(A)$  as well, is also upper bounded by that same quantity.

## References

- [1] D. Bienstock and P. Seymour. Monotonicity in graph searching. *Journal of Algorithms*, 12(2):239 – 245, 1991.
- [2] C. Borgs, J. Chayes, A. Ganesh, and A. Saberi. How to distribute antidote to control epidemics. *Random Structures and Algorithms*, 37(2):204–222, 2010.
- [3] F. R. K. Chung, P. Horn, and A. Tsiatas. Distributing antidote using pagerank vectors. *Internet Mathematics*, 6(2):237–254, 2009.
- [4] M. Chung, F. Makedon, I. Sudborough, and J. Turner. Polynomial time algorithms for the min cut problem on degree restricted trees. *SIAM Journal on Computing*, 14(1):158–177, 1985.
- [5] R. Cohen, S. Havlin, and D. ben-Avraham. Efficient immunization strategies for computer networks and populations. *Physical Review Letters*, 91:247901, 2003.
- [6] K. Drakopoulos, A. Ozdaglar, and J. N. Tsitsiklis. Network resilience against epidemic spread. *working paper*, 2014.
- [7] E. Gourdin, J. Omic, and P. Van Mieghem. Optimization of network protection against virus spread. In *Design of Reliable Communication Networks (DRCN), 2011 8th International Workshop on the*, pages 86–93. IEEE, October 2011.
- [8] E. Korach and N. Solel. Tree-width, path-width, and cutwidth. *Discrete Applied Mathematics*, 43(1):97 – 101, 1993.
- [9] D. A. Levin, Y. Peres, and E. L. Wilmer. *Markov Chains and Mixing Times*. American Mathematical Society, 2006.
- [10] F. Makedon and I. Sudborough. On minimizing width in linear layouts. *Discrete Applied Mathematics*, 23(3):243 – 265, 1989.
- [11] V. M. Preciado, M. Zargham, C. Enyioha, A. Jadbabaie, and G. J. Pappas. Optimal vaccine allocation to control epidemic outbreaks in arbitrary networks. *CoRR*, abs/1303.3984, 2013.
- [12] J. M. Steele. *Stochastic Calculus and Financial Applications*. Springer, 2000.