

# Many-body characterization of topological superconductivity: The Richardson-Gaudin-Kitaev chain

Gerardo Ortiz,<sup>1</sup> Jorge Dukelsky,<sup>2</sup> Emilio Cobanera,<sup>3</sup> Carlos Eсеbbag,<sup>4</sup> and Carlo Beenakker<sup>3</sup>

<sup>1</sup>*Department of Physics, Indiana University, Bloomington, IN 47405, USA\**

<sup>2</sup>*Instituto de Estructura de la Materia, C.S.I.C., Serrano 123, E-28006 Madrid, Spain*

<sup>3</sup>*Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands*

<sup>4</sup>*Departamento de Física y Matemáticas, Universidad de Alcalá, 28871 Alcalá de Henares, Spain*

(Dated: July 2014)

What distinguishes trivial from topological superfluids in interacting many-body systems where the number of particles is conserved? Building on a class of integrable pairing Hamiltonians, we present a number-conserving, interacting variation of the Kitaev model, the Richardson-Gaudin-Kitaev chain, that remains exactly solvable for periodic and antiperiodic boundary conditions. Our model allows us to identify fermion parity switches that distinctively characterize topological superconductivity in interacting many-body systems. Although the Majorana zero-modes in this model have only a power-law confinement, we may still define many-body Majorana operators by tuning the flux to a fermion parity switch. We derive a closed-form expression for an interacting topological invariant and show that the transition away from the topological phase is of third order.

In recent years, the physics of Majorana zero-energy modes has become a key subfield of condensed matter physics [1–4]. On the theory side, a central result is the bulk-boundary correspondence [5] that associates Majorana zero-modes to the boundary of (or defects in) a topologically non-trivial superconductor, with the Kitaev chain as a prototypical example [6]. The mathematical formalism underlying this correspondence relies on the symmetries and topological invariants of the Bogoliubov-de Gennes equation [7], a mean-field description of the superconducting state in which the conservation of the number of fermions (a continuous symmetry) is broken down to a discrete symmetry, the conservation of fermion-number parity. Majorana zero-modes are directly linked to the spontaneous breaking of this residual discrete symmetry [8].

As the experimental side of Majorana physics continues to develop [9–14], it becomes crucial to unveil how much of the mean-field picture survives beyond its natural limits. This has motivated recent studies [15–21], with the focus on the anomalous  $2\Phi_0 = h/e$  flux-periodicity of the Josephson effect — the hallmark of a topological superconductor [6].

A main thrust of this paper is the characterization of interacting many-body, number-conserving, topological superconductors, or superfluids, and the fate of Majorana zero-modes beyond mean-field. The theoretical study of any interacting quantum system is hampered by the exponential growth of the Hilbert space with the number of particles. An additional complication of superconducting systems is the lack of simple principles to guide the design of particle-number conserving models, in which the phase of the order parameter is not a good quantum number. To overcome both obstacles, we have constructed an exactly solvable, number-conserving variation of the Kitaev

chain. Because our model belongs to a class of integrable pairing models [22–24] based on the  $s$ -wave reduced BCS Hamiltonian first solved by Richardson [25], and on a generalization of the Gaudin magnet [26], we will refer to it as the Richardson-Gaudin-Kitaev (RGK) chain.

The RGK model is integrable for periodic and antiperiodic boundary conditions. It allows us to obtain precise answers for the characterization problem posed here, including the very existence of a topologically nontrivial phase, in an interacting number-conserving system, the order of the phase transition into the trivial phase, the definition of a topological invariant beyond mean-field, and the emergence of many-body Majorana zero-modes.

*Richardson-Gaudin-Kitaev chain.*— Our model is defined by the Hamiltonian

$$H_{\text{RGK}} = \sum_{k \in \mathcal{S}_k^\phi} \varepsilon_k \hat{c}_k^\dagger \hat{c}_k - 8G \sum_{k, k' \in \mathcal{S}_{k+}^\phi} \eta_k \eta_{k'} \hat{c}_k^\dagger \hat{c}_{-k}^\dagger \hat{c}_{-k'} \hat{c}_{k'} \quad (1)$$

in terms of fermion creation operators  $\hat{c}_k^\dagger$ , with momentum  $k$ -dependent single-particle spectrum  $\varepsilon_k = -2t_1 \cos k - 2t_2 \cos 2k$  and interaction strength  $G > 0$ . The interaction is modulated by the potential

$$\eta_k = \sin(k/2) \sqrt{t_1 + 4t_2 \cos^2(k/2)}, \quad (2)$$

which displays the odd-parity behavior  $\eta_k = -\eta_{-k}$  characteristic of  $p$ -wave superconductivity. The pair potential is related to the single-particle spectrum by  $4\eta_k^2 = \varepsilon_k + 2t_+$ ,  $t_+ = t_1 + t_2$ . This relation is the key to achieve exact solvability.

In real space, we define  $c_j = L^{-1/2} \sum_{k \in \mathcal{S}_k^\phi} e^{ijk} \hat{c}_k$  for a chain of length  $L$ , measured in units of the lattice constant. We take  $\phi$ -dependent boundary conditions  $c_{i+L} = e^{i\phi/2} c_i$ . In a ring geometry, periodic boundary conditions ( $\phi = 0$ ) correspond to enclosed flux  $\Phi = 0$  and antiperiodic boundary conditions ( $\phi = 2\pi$ ) correspond to  $\Phi = \Phi_0$ . The resulting sets of allowed momenta  $\mathcal{S}_k^\phi$  are  $\mathcal{S}_k^0 = \mathcal{S}_{k+}^0 \oplus \mathcal{S}_{k-}^0 \oplus \{0, -\pi\}$  and  $\mathcal{S}_k^{2\pi} = \mathcal{S}_{k+}^{2\pi} \oplus \mathcal{S}_{k-}^{2\pi}$ ,

\*Electronic address: ortizg@indiana.edu

with  $\mathcal{S}_{k\pm}^0 = L^{-1}\{\pm 2\pi, \pm 4\pi, \dots, \pm(\pi L - 2\pi)\}$  and  $\mathcal{S}_{k\pm}^{2\pi} = L^{-1}\{\pm\pi, \pm 3\pi, \dots, \pm(\pi L - \pi)\}$ . The RGK Hamiltonian in real space reads

$$H_{\text{RGK}} = - \sum_{i=1}^L \sum_{r=1}^2 \left( t_r c_i^\dagger c_{i+r} + \text{H.c.} \right) - 2G I_\phi^\dagger I_\phi, \quad (3)$$

$$I_\phi \equiv 2i \sum_{k \in \mathcal{S}_{k+}^\phi} \eta_k \hat{c}_k \hat{c}_{-k} = \sum_{i>j}^L \eta(i-j) c_i c_j. \quad (4)$$

There are at least two cases where the pairing function  $\eta(m)$  can be determined in closed form by Fourier transformation of Eq. (2). For  $t_1 = 0$  and  $t_2 \neq 0$ ,  $\eta(m) = \sqrt{t_2} \delta_{m1}$ , i.e. we have nearest-neighbor pairing only. For  $t_1 \neq 0$  and  $t_2 = 0$  we obtain

$$\eta(m) = \frac{(-1)^m 8\sqrt{t_1}}{\pi} \frac{m}{1-4m^2}, \quad \text{for } L \rightarrow \infty, \quad (5)$$

so a long-range pairing interaction with a slow  $1/m$  decay with distance  $m = i - j$ . In general,  $\eta(m)$  is a monotonous, decaying function of  $m$ .

This long-range pairing interaction is the difference with the original Kitaev model [6] that allows for the exact solution beyond the mean-field approximation. As we shall see in a moment, the long-range coupling still allows for a topologically nontrivial phase. It may also be physically relevant for chains of magnetic nanoparticles on a superconducting substrate [27, 28], which have recently been shown to support topologically protected Majorana zero-modes in the presence of a long-range coupling [29].

*Mean-field approximation.*— Before we work out the exact solution of the RGK chain, we would like to establish first whether it displays a non-trivial topological phase in the mean-field approximation.

We set  $t_2 = 0$  for simplicity and take the pairing interaction (5). The mean-field approximation to the RGK chain is obtained from the substitution  $2GI^\dagger I \rightarrow \Delta^* I + \Delta I^\dagger$ , with gap function  $\Delta = 2G \langle I_L \rangle = e^{i\theta} |\Delta|$ . We define Majorana fermion operators  $a_i = e^{-i\theta/2} c_i + e^{i\theta/2} c_i^\dagger$ ,  $b_i = e^{-i\theta/2} c_i - e^{i\theta/2} c_i^\dagger$ . The mean-field Hamiltonian is

$$H_{\text{mf}} = \frac{it_1}{2} \sum_{i=1}^{L-1} (b_i a_{i+1} - a_i b_{i+1}) - \frac{i}{2} \sum_{i>j}^L \Delta_{i-j} (b_i a_j + a_i b_j), \quad (6)$$

where  $\Delta_{i-j} = |\Delta| \eta(i-j)$ , and displays a topological phase characterized by power-law Majorana edge modes and an associated  $4\pi$ -periodic Josephson effect. For clarity of presentation, let us compute approximate edge modes to leading order in  $\delta_2 = \Delta_2 / (\Delta_1 + t_1)$ . The Majorana mode localized at the left end of the chain is  $\tilde{a}_1 = a_1 + \sum_{i=2}^{L-1} \delta_i \sum_{j=2}^L (-\delta_2)^{L-j} a_j$ , so that  $i[H_b^{\text{mf}}, \tilde{a}_1] = \Delta_{L-1} b_L + \mathcal{O}(\delta_2)$ . We see that  $\Delta_1 + t_1$  controls the localization of the Majoranas, while  $\Delta_{L-1}$  controls the vanishing of the commutator with the Hamiltonian.

Due to the long-range pairing interaction (5), the wave function of the Majorana modes decays algebraically

rather than exponentially in the bulk, and their energy approaches zero as a power law in  $1/L$ , similarly to what has been found in other mean-field models based on the Kitaev chain with long-range coupling [29–31].

*Exact solution.*— To show that the RGK chain is exactly solvable, let us rewrite

$$H_{\text{RGK}} = 8H_\phi + \delta_{\phi,0} (\varepsilon_0 \hat{c}_0^\dagger \hat{c}_0 + \varepsilon_{-\pi} \hat{c}_{-\pi}^\dagger \hat{c}_{-\pi}) - 4t_+ S^z + C_\phi, \quad (7)$$

$$H_\phi = \sum_{k \in \mathcal{S}_{k+}^\phi} \eta_k^2 S_k^z - G \sum_{k,k' \in \mathcal{S}_{k+}^\phi} \eta_k \eta_{k'} S_k^+ S_{k'}^-, \quad (8)$$

with  $C_\phi = 2t_2 \delta_{\phi,0}$ . We introduced the operators  $S_k^z = \frac{1}{2} (\hat{c}_k^\dagger \hat{c}_k + \hat{c}_{-k}^\dagger \hat{c}_{-k} - 1)$  and  $S_k^+ = \hat{c}_k^\dagger \hat{c}_{-k}^\dagger$  for each pair  $(k, -k)$  of pairing-active momenta. They satisfy the algebra of  $\text{SU}(2)$ . Hence  $S^z = \sum_{k \in \mathcal{S}_{k+}^\phi} S_k^z$  defines a conserved quantity,  $[H_\phi, S^z] = 0$ .

The Hamiltonian  $H_\phi$  belongs to the hyperbolic family of exactly solvable pairing Hamiltonians [24, 32], whose best known representative is the chiral  $p$ -wave superfluid [33–36]. The rational family includes  $s$ -wave pairing and has been used in the study of the BCS-BEC crossover phenomenon [37]. Eigenstates for  $2M + N_\nu$  fermions are

$$|\Phi_{M,\nu}\rangle = \prod_{\alpha=1}^M \left( \sum_{k \in \mathcal{S}_{k+}^\phi} \frac{\eta_k}{\eta_k^2 - E_\alpha} \hat{c}_k^\dagger \hat{c}_{-k}^\dagger \right) |\nu\rangle. \quad (9)$$

The state  $|\nu\rangle$  with  $N_\nu$  unpaired fermions satisfies  $S_k^- |\nu\rangle = 0$  for all  $k$ . Moreover,  $S_k^z |\nu\rangle = -s_k |\nu\rangle$ , with  $s_k = 0$  if the level  $k$  is singly-occupied or  $s_k = 1/2$  if it is empty. The corresponding energy levels are  $E_{M,\nu} = \langle \nu | H_\phi | \nu \rangle + \sum_{\alpha=1}^M E_\alpha$ , with spectral parameters  $E_\alpha$  determined by the Richardson-Gaudin equations

$$\sum_{k \in \mathcal{S}_{k+}^\phi} \frac{s_k}{\eta_k^2 - E_\alpha} - \sum_{\beta(\neq\alpha)} \frac{1}{E_\beta - E_\alpha} = \frac{Q_\phi}{E_\alpha}, \quad (10)$$

where  $Q_\phi = 1/2G - \sum_{k \in \mathcal{S}_{k+}^\phi} s_k + M - 1$ .

In the case of periodic boundary conditions ( $\phi = 0$ ) the two momenta  $k = 0, -\pi$  are not affected by the interactions and need to be included separately. The eigenvectors then are  $|\Psi_N\rangle = |n_0 n_{-\pi}\rangle \otimes |\Phi_{M,\nu}\rangle$ , where  $n_0, n_{-\pi} \in \{0, 1\}$  and the total number of fermions is  $N = 2M + N_\nu + n_0 + n_{-\pi}$ .

The Richardson-Gaudin equations (10) become singular when two or more  $E_\alpha$ 's approach the same single-particle energy  $\eta_k^2$  and also around  $E_\alpha = 0$ . At specific values of the interaction strength  $G$

$$G_n^\phi = \frac{2}{L} g_n^\phi = \frac{2}{L - 2(2M + N_\nu) + 2(n + 1 - \delta_{\phi,0})}, \quad (11)$$

there are  $n$  solutions  $E_\alpha$ ,  $1 \leq n \leq M$ , that vanish identically [34, 36]. In particular, at  $n = M$ , where (9) becomes a pair condensate,  $G_M^\phi$  is precisely the Moore-Read coupling. And, at  $n = 1$  where  $Q_\phi = 0$ ,  $G_1^\phi = G_c$  is the

Read-Green coupling, associated to the non-analytic behavior of observables in the thermodynamic limit [34, 36].

The numerical solution of Eq. (10) is particularly simple for values of  $G > G_1^\phi$ , when all  $E_\alpha$ 's are real and negative, and for  $G = G_n^\phi$  when  $n$  of the  $E_\alpha$ 's vanish while  $M - n$  are real and negative. We carried out computations with systems of up to  $L \approx 2000$  sites at quarter filling. Without the integrability condition, this would have required diagonalization of a Hamiltonian matrix with the unwieldy dimension  $5 \cdot 10^{242}$ . The exact solvability reduces the complexity of the problem to the solution of the  $M \simeq 250$  nonlinear coupled equations (10).

We were also able to perform extrapolations to the thermodynamic limit  $N, L \rightarrow \infty$  at finite density  $\rho = N/L$  and rescaled interaction strength  $g = GL/2$ . In that limit, Eqs. (10) relate to the mean-field gap and number equations for  $H_\phi$  [34]

$$\frac{2\pi}{g} = \int_0^\pi \frac{\eta_k^2}{E_k} dk, \quad \rho = \frac{1}{\pi} \int_0^\pi v_k^2 dk, \quad (12)$$

$$E_k = \sqrt{\left(\frac{\eta_k^2}{2} - \mu\right)^2 + \eta_k^2 \Delta^2}, \quad v_k^2 = \frac{1}{2} - \frac{\eta_k^2 - 2\mu}{4E_k}, \quad (13)$$

with quasi-energies  $E_k$  and occupation probabilities  $v_k^2$ .

*Phase diagram and topological transition.*— To establish the quantum phase diagram of the RGK chain, one needs the ground state energy  $\mathcal{E}_0(\rho, g)$  of  $H_{\text{RGK}}$ . Depending on the boundary condition and fermion-number parity, one has to consider either  $N_\nu = 0$  or 1. For periodic boundary conditions, since the levels  $k = 0, -\pi$  decouple from the rest,  $N_\nu = 0$  for both even and odd  $N$ . If  $N$  is odd, the unpaired particle occupies the  $k = 0$  level without blocking an active level. For antiperiodic boundary conditions the ground state has  $N_\nu = 0$  for  $N$  even, while for  $N$  odd it has  $N_\nu = 1$  with blocked level  $k_0$ . The resulting ground state energy is given by

$$\mathcal{E}_0^\phi(N) = 8 \sum_{\alpha=1}^M E_\alpha - 4t_+ M + J_{\phi,0} + \delta_{N_\nu,1}(4\eta_{k_0}^2 - 2t_+), \quad (14)$$

where  $J_{\phi,0} = \delta_{\phi,0}(\varepsilon_0 \delta_{n_0,1} + \varepsilon_{-\pi} \delta_{n_{-\pi},1})$ . In the thermodynamic limit the energy density reduces to

$$e_0 \equiv \lim_{L \rightarrow \infty} \mathcal{E}_0^\phi/L = -2t_+\rho - \frac{4}{g}\Delta^2 + \frac{4}{\pi} \int_0^\pi \eta_k^2 v_k^2 dk. \quad (15)$$

The resulting phase diagram is shown in Fig. 1. The RGK chain is gapped for all  $g > 0$ , except for the Read-Green coupling  $g_c = G_{n=1}^\phi L/2$  where it becomes critical in the thermodynamic limit (without any dependence on the choice of boundary conditions). This critical line defines the phase boundary separating weak (topological) from strong (trivial) pairing phases, and thus is a line of non-analyticities. At  $g_c$  a cusp develops in the second derivative of  $e_0$ , that leads to a singular discontinuous behavior of the third-order derivative. Hence the transition from a weakly-paired to a strongly-paired superconductor is of third order, just like for the two-dimensional chiral  $p$ -wave superconductor [34, 36].




FIG. 1: Third-order phase transition between the topological (weak-pairing) and trivial (strong-pairing) superconducting phases. The continuous line denotes the second order derivative of the ground-state energy density  $e_0$ , evaluated in the thermodynamic limit. Circles are the exact solution for  $L = 2048$ ,  $N = 512$ , and antiperiodic boundary conditions. Top inset: Discontinuity in the third-order derivative. Bottom inset: Quantum phase diagram in the  $(\rho, g)$ -plane. Dashed and full lines represent, respectively, the Moore-Read ( $g_M^{-1} = 1 - \rho$ ) and Read-Green ( $g_c^{-1} = 1 - 2\rho$ ) boundaries.

Open circles in Fig. 1 correspond to the second-order derivative of the exact  $e_0$  for the antiperiodic RGK chain with  $L = 2048$ ,  $N = 512$ , and  $t_1 = 1, t_2 = 0$ , obtained by solving Eqs. (10) for some selected pairing strength values, and illustrates how close to the thermodynamic limit these systems sizes are. We will demonstrate shortly that the weak-pairing phase of the RGK chain is indeed topologically nontrivial.

*Fermion parity switches.*— We next introduce a quantitative criterion to establish emergence of topological superconductivity in particle-number conserving, many-body systems. The criterion exploits the behavior of the ground state energy of a system of  $N$ , and  $N \pm 1$  particles, for both periodic and antiperiodic boundary conditions. The emergence of topological order in a superconducting wire, closed in a ring and described in mean-field, is associated with switches in the ground-state fermion parity  $\mathcal{P}(\phi)$  upon increasing the enclosed flux  $\Phi = (\phi/2\pi) \times \Phi_0$  [6, 38–43]. Any spin-active superconductor, topologically trivial or not, may experience a crossing of the ground state energies for even and odd number of electrons [44–47]. Regardless of spin, what matters is the number of crossings  $N_X$  between  $\Phi = 0 = \phi$  and  $\Phi = \Phi_0$ ,  $\phi = 2\pi$ . The superconductor is topologically nontrivial if  $N_X$  is odd, otherwise it is trivial.

In the many-body, number conserving, case we need to identify the relevant parity switches signaling the emergence of a topological superconducting phase. Our exact solution gives us access to  $\mathcal{P}(\phi)$  only at  $\phi = 0$  and  $\phi = 2\pi$ , but this is sufficient to determine whether  $N_X$  is even or odd. Notice that odd  $N_X$  means that the flux  $\Phi$  should




FIG. 2: Ground state energies for the RGK chain (in units of  $t_1 \equiv 1$ , for  $t_2 = 0$ ) for even ( $N = 2M$ ) and odd ( $N = 2M \pm 1$ ) number of fermions, and with periodic ( $\phi = 0$ ) or antiperiodic ( $\phi = 2\pi$ ) boundary conditions. The main plot shows the odd-even difference as a function of the interaction strength  $g$  for a finite system (data points, for  $N = 512$ ,  $L = 2048$ ) and in the thermodynamic limit (continuous lines). The topologically nontrivial state is entered for  $g < g_c = 2$ . The insets show the even and odd energies themselves, for the finite system (lower two insets) and in the thermodynamic limit (upper two insets), illustrating the fermion parity switches  $\mathcal{P}_N(\phi)$ .

be advanced by  $2\Phi_0$  — rather than  $\Phi_0$  — in order to return to the initial ground state, which is the essence of the  $4\pi$ -periodic Josephson effect [6, 48].

To identify the fermion parity switches we calculate the ground state energy  $\mathcal{E}_0^\phi(N)$  for a given number  $N$  of fermions in the chain of length  $L$ , with periodic ( $\phi = 0$ ) or antiperiodic ( $\phi = 2\pi$ ) boundary conditions, and compare  $\mathcal{E}_0^{\text{odd}}(\phi) = \frac{1}{2}\mathcal{E}_0^\phi(N+1) + \frac{1}{2}\mathcal{E}_0^\phi(N-1)$  and  $\mathcal{E}_0^{\text{even}}(\phi) = \mathcal{E}_0^\phi(N)$ , where we assumed  $N$  even. The difference (inverse compressibility)  $\chi(\phi) = \mathcal{E}_0^{\text{odd}}(\phi) - \mathcal{E}_0^{\text{even}}(\phi)$  determines  $\mathcal{P}_N(\phi) = \text{sign } \chi(\phi)$ , so it has the opposite sign at  $\phi = 0$  and  $\phi = 2\pi$  in the topologically nontrivial phase. We also find that  $\mathcal{P}_{N \in \text{even}}(\phi) = -\mathcal{P}_{N \in \text{odd}}(\phi)$  in the topologically nontrivial phase. The results, shown in Fig. 2, unambiguously demonstrate the topologically nontrivial nature of the superconductor for  $g < g_c$  — both in a finite system and in the thermodynamic limit, and without relying on any mean-field approximation.

The ground state of the odd ( $2M \pm 1$ ) system strongly depends on the boundary conditions. For periodic boundary conditions the unpaired particle always occupies the  $k_0 = 0$  level, while for the antiperiodic case it starts blocking the Fermi momentum  $k_F = k_0$  at  $g = 0$ , continuously decreasing its modulus with increasing  $g$ , up to  $k_0 = \pi/L$  at  $g_0 \sim 1.1936$  ( $\rho = 1/4$ ), corresponding to  $\mu = \Delta^2$  in the thermodynamic limit. In that limit  $\chi(\phi)$  has a particularly simple form:  $\chi(0) = -8\mu$ , and  $\chi(2\pi) = 8|\mu|$  for  $g > g_0$ .

*Topological invariant and zero-modes*— The occupa-

tion number  $\mathcal{N}_{k=0}$  of the  $k = 0$  single-particle state is a topological invariant [34, 49], being the one-dimensional analogue of the winding number in two space dimensions [50, 51]. By combining the integrals of motion [24, 34] with the Hellmann-Feynman theorem we find

$$\mathcal{N}_k = \frac{1}{2} - s_k - 4s_k\gamma^2 \sum_{\alpha=1}^M \frac{\eta_k^2}{(\eta_k^2 - E_\alpha)^2} \frac{\partial E_\alpha}{\partial \gamma}, \quad (16)$$

$$\gamma = \frac{G_c W(G)}{3W(G) - 2G_c}, \quad W(G) = \frac{G_c G}{2G - G_c}. \quad (17)$$

The mapping  $W(G)$  is an involution,  $W(W(G)) = G$ , with fixed point  $W(G_c) = G_c$ , typical of self-dual transformations. In the thermodynamic limit,  $\lim_{L \rightarrow \infty} \mathcal{N}_{k=0} = \Theta[G_c - G]$  with  $\Theta[x]$  the unit step function, thus signaling the topological phase transition at  $G = G_c$ .

We saw in the mean-field calculation that the Majorana zero-modes have a power-law decay, so for any finite  $L$  they overlap and are displaced from precisely zero energy. It is known for the mean-field Kitaev chain that it is still possible to construct exact zero-energy modes by tuning the flux to a fermion parity switch [52]. We may follow a similar strategy for our many-body system. By varying the boundary conditions in the topologically nontrivial phase the ground state energies  $\mathcal{E}_0^{\text{odd}}(\phi)$  and  $\mathcal{E}_0^{\text{even}}(\phi)$  cross at some critical  $\phi^*$ . (They must cross, because their order is inverted at  $\phi = 0$  and  $\phi = 2\pi$ .) At this value of  $\phi = \phi^*$  one can identify many-body Majorana zero-modes as follows.

We define the normalized ground states  $|\Psi_0^{\text{even}}\rangle = |\Psi_0^N\rangle$ ,  $|\Psi_0^{\text{odd}}\rangle = (|\Psi_0^{N+1}\rangle + |\Psi_0^{N-1}\rangle)/\sqrt{2}$ , with the transition operator  $\hat{T} = |\Psi_0^{\text{even}}\rangle\langle\Psi_0^{\text{odd}}|$ . Clearly,  $\hat{T}^2 = 0$ , and  $\{\hat{T}, \hat{T}^\dagger\} = \hat{P}_0$ . Here,  $\hat{P}_0 = |\Psi_0^{\text{even}}\rangle\langle\Psi_0^{\text{even}}| + |\Psi_0^{\text{odd}}\rangle\langle\Psi_0^{\text{odd}}|$  is the projector onto the ground-state subspace. The corresponding Majorana operators are

$$\Gamma_1 = \hat{T} + \hat{T}^\dagger, \quad \Gamma_2 = \hat{T} - \hat{T}^\dagger, \quad \{\Gamma_1, \Gamma_2\} = 0. \quad (18)$$

We expect these zero-modes to be localized at the edges of the RGK chain due to the energy gap in the bulk, and it would be interesting to check this numerically.

In conclusion, we have constructed a variation of the Kitaev Hamiltonian that is both number conserving and interacting, but still exactly solvable. This allowed us to identify the fermion parity switches needed for characterizing topological superconductivity in interacting many-body systems. We have shown that our Richardson-Gaudin-Kitaev model shares the features of the mean-field Kitaev model that have made it a paradigm of topological superconductivity. There is one difference, the long-range nature of the pairing interaction, but in view of recent experimental developments [28, 29], this may be a welcome feature of the model rather than a drawback. Irrespective of the potential application of our model to a real physical system, we anticipate that its integrability will make it a fruitful starting point for theoretical studies of interacting models with topological order.

JD and CE are supported by grant FIS2012-34479 of the Spanish Ministry of Economy and Competitiveness. EC and CB acknowledge support by the Netherlands Or-

ganization for Scientific Research (OCW/NWO/FOM) and an ERC Synergy Grant.

- 
- [1] M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [2] X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).
- [3] J. Alicea, *Rep. Prog. Phys.* **75**, 076501 (2012).
- [4] C. W. J. Beenakker, *Annu. Rev. Con. Mat. Phys.* **4**, 113 (2013).
- [5] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, *New J. Phys.* **12**, 065010 (2010).
- [6] A. Yu. Kitaev, *Phys. Usp.* **44** (suppl.), 131 (2001).
- [7] P.-G. de Gennes, *Superconductivity of Metals and Alloys* (Benjamin, New York, 1966).
- [8] B. van Heck, E. Cobanera, J. Ulrich, and F. Hassler, *Phys. Rev. B* **89**, 165416 (2014).
- [9] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, *Science* **336**, 1003 (2012).
- [10] A. Das, Y. Ronen, Y. Most, Y. Oreg, M. Heiblum, and H. Shtrikman, *Nature Phys.* **8**, 887 (2012).
- [11] M. T. Deng, C. L. Yu, G. Y. Huang, M. Larsson, P. Caroff, and H. Q. Xu, *Nano Lett.* **12**, 6414 (2012).
- [12] A. D. K. Finck, D. J. Van Harlingen, P. K. Mohseni, K. Jung, and X. Li, *Phys. Rev. Lett.* **110**, 126406 (2013).
- [13] H. O. H. Churchill, V. Fatemi, K. Grove-Rasmussen, M. T. Deng, P. Caroff, H. Q. Xu, and C. M. Marcus, *Phys. Rev. B* **87**, 241401 (2013).
- [14] T. D. Stanescu and S. Tewari, *J. Phys. Cond. Matt.* **25**, 233201 (2013).
- [15] A. M. Tselik, arXiv:1106.2996.
- [16] M. Cheng and H.-H. Tu, *Phys. Rev. B* **84**, 094503 (2011).
- [17] F. Hassler and D. Schuricht, *New J. Phys.* **14**, 125018 (2012).
- [18] A. Manolescu, D. C. Marinescu, and T. D. Stanescu, *J. Phys. Condens. Matter* **26**, 172203 (2014).
- [19] J. D. Sau, B. I. Halperin, K. Flensberg, and S. Das Sarma, *Phys. Rev. B* **84**, 144509 (2011).
- [20] L. Fidkowski, R. M. Lutchyn, C. Nayak, and M. P. A. Fisher, *Phys. Rev. B* **84**, 195436 (2011).
- [21] B. van Heck, F. Hassler, A. R. Akhmerov, and C. W. J. Beenakker, *Phys. Rev. B* **84**, 180502(R) (2011).
- [22] L. Amico, A. Di Lorenzo, and A. Osterloh, *Phys. Rev. Lett.* **86**, 5759 (2001).
- [23] J. Dukelsky, C. Esebbag, and P. Schuck, *Phys. Rev. Lett.* **87**, 066403 (2001).
- [24] G. Ortiz, R. Somma, J. Dukelsky, and S. M. A. Rombouts, *Nuc. Phys. B* **707**, 421 (2005).
- [25] R. W. Richardson, *Phys. Lett.* **3**, 277 (1963).
- [26] M. Gaudin, *J. Phys. (Paris)* **37**, 1087 (1976).
- [27] T.-P. Choy, J. M. Edge, A. R. Akhmerov, and C. W. J. Beenakker, *Phys. Rev. B* **84**, 195442 (2011).
- [28] S. Nadj-Perge, I. K. Drozdov, B. A. Bernevig, and A. Yazdani, *Phys. Rev. B* **88**, 020407 (2013).
- [29] F. Pientka, L. I. Glazman, and F. von Oppen, *Phys. Rev. B* **88**, 155420 (2013).
- [30] D. Vodola, L. Lepori, E. Ercolessi, A. V. Gorshkov, and G. Pupillo, arXiv:1405.5440.
- [31] W. DeGottardi, M. Thakurathi, S. Vishveshwara, and D. Sen, *Phys. Rev. B* **88**, 165111 (2013).
- [32] J. Dukelsky, S. Pittel, and G. Sierra, *Rev. Mod. Phys.* **76**, 643 (2004).
- [33] M. I. Ibañez, J. Links, G. Sierra, and S. Y. Zhao, *Phys. Rev. B* **79**, 180501(R) (2009).
- [34] S. M. A. Rombouts, J. Dukelsky, and G. Ortiz, *Phys. Rev. B* **82**, 224510 (2010).
- [35] C. Dunning, M. Ibanez, J. Links, G. Sierra, and S.-Y. Zhao, *J. Stat. Mech.* P08025 (2010).
- [36] S. Lerma H., S. M. A. Rombouts, J. Dukelsky, and G. Ortiz, *Phys. Rev. B* **84**, 100503(R) (2011).
- [37] G. Ortiz and J. Dukelsky, *Phys. Rev. A* **72**, 043611 (2005).
- [38] A. Keselman, L. Fu, A. Stern, and E. Berg, *Phys. Rev. Lett.* **111**, 116402 (2013).
- [39] C. W. J. Beenakker, J. M. Edge, J. P. Dahlhaus, D. I. Pikulin, S. Mi, and M. Wimmer, *Phys. Rev. Lett.* **111**, 037001 (2013).
- [40] J. D. Sau and E. Demler, *Phys. Rev. B* **88**, 205402 (2013).
- [41] A. Haim, A. Keselman, E. Berg, and Y. Oreg, *Phys. Rev. B* **89**, 220504 (2014).
- [42] F. Crépin and B. Trauzettel, *Phys. Rev. Lett.* **112**, 077002 (2014).
- [43] C. W. J. Beenakker, arXiv:1407.2131.
- [44] A. Sakurai, *Prog. Theor. Phys.* **44**, 1472 (1970).
- [45] A. V. Balatsky, I. Vekhter, and J.-X. Zhu, *Rev. Mod. Phys.* **78**, 373 (2006).
- [46] W. Chang, V. E. Manucharyan, T. S. Jespersen, J. Nygård, and C. M. Marcus, *Phys. Rev. Lett.* **110**, 217005 (2013).
- [47] E. J. H. Lee, X. Jiang, M. Houzet, R. Aguado, C. M. Lieber, and S. De Franceschi, *Nature Nanotech.* **9**, 79 (2014).
- [48] K. Sengupta, I. Žutić, H.-J. Kwon, V. M. Yakovenko, and S. Das Sarma, *Phys. Rev. B* **63**, 144531 (2001).
- [49] S. S. Botelho and C. A. R. Sà de Melo, *J. Low Temp. Phys.* **140**, 409 (2005).
- [50] G. E. Volovik, *Sov. Phys. JETP* **67**, 1804 (1988).
- [51] M. S. Foster, M. Dzero, V. Gurarie, and E. A. Yuzbashyan, *Phys. Rev. B* **88**, 104511 (2013).
- [52] A. R. Akhmerov, J. P. Dahlhaus, F. Hassler, M. Wimmer, and C. W. J. Beenakker, *Phys. Rev. Lett.* **106**, 057001 (2011). [Supplemental material.]