Maxima of the signless Laplacian spectral radius for planar graphs^{*}

Guanglong Yu^{a^{\dagger}} Jianyong Wang^b Shu-Guang Guo^a

^aDepartment of Mathematics, Yancheng Teachers University, Yancheng, 224002, Jiangsu, P.R. China
^bDepartment of Mathematics, Linyi University, Linyi, 276005, Shandong, P.R. China

Abstract

The signless Laplacian spectral radius of a graph is the largest eigenvalue of its signless Laplacian. In this paper, we prove that the graph $K_2 \nabla P_{n-2}$ has the maximal signless Laplacian spectral radius among all planar graphs of order $n \ge 456$.

AMS Classification: 05C50 **Keywords:** Signless Laplacian; Spectral radius; Planar graph

1 Introduction

The (adjacency) spectral radius of a graph is used in many fields, including chemistry, physics and computer science [4], [10], [22]. It arises a broad research now. Spectral radius of planar graphs is of great interest in harmonic analysis, and bounds on it can be used in the design and analysis of certain Monte Carlo algorithms [19]. Spectral radius of planar graphs have applications not only in the theory of algorithms but also in theoretical physics. Boots and Royle investigated the spectral radius of planar graphs motivated by an application in geography networks [2]. This makes the research about the spectral radius of planar graphs interesting and dynamic, and many interesting results have emerged (see [19], [3], [15, 16], [20], for example). Very recently the signless Laplacian has attracted the attention of researchers. Some results on the signless Laplacian spectrum have been reported since 2005 and a new spectral theory called the *Q*-theory is being developed by many researchers [6]-[9], [13]. A nature question is that how about the application of the *Q*-theory. By computer investigations of graphs [11], E. van Dam and W. Haemers found that the signless Laplacian spectrum or Laplacian

^{*}Supported by NSFC (Nos. 11271315, 11171290, 11201417) and Jiangsu Qing Lan Project (2014A).

[†]E-mail addresses: yglong01@163.com.

spectrum in distinguishing non-isomorphic graphs. In computer science, signless Laplacian spectral radius can also be used to study the properties of a network [10], [24, 25]. Schwenk and Wilson initiated the study of the eigenvalues of planar graphs [20]. In [3], D. Cao and A. Vince conjectured that $K_2 \nabla P_{n-2}$ has the maximum spectral radius among all planar graphs of order n, where ∇ denotes the *join* of two graphs obtained from the union of these two graphs by joining each vertex of the first graph to each vertex of the second graph. The conjuncture is still open now. With the development of the Q-theory, a natural question is that what about the maximum signless Laplacian spectral radius of planar graphs. By some comparisons in [13], it seems plausible that $K_2 \nabla P_{n-2}$ also has the maximal signless Laplacian spectral radius among planar graphs. In this paper, we confirm that among planar graphs with order $n \ge 456$, $K_2 \nabla P_{n-2}$ has the maximal signless Laplacian spectral radius.

The layout of this paper is as follows. Section 2 gives some notations and some working lemmas. In section 3, our results are presented.

2 Preliminary

All graphs considered in this paper are undirected and simple, i.e. no loops or multiple edges are allowed. Denote by G = G[V(G), E(G)] a graph with vertex set V(G)and edge set E(G). |V(G)| = n is the order. m(G) = |E(G)| is edge number. Recall that given a graph G, Q(G) = D(G) + A(G) is called the signless Laplacian matrix of G, where $D(G) = \text{diag}(d_1, d_2, \ldots, d_n)$ with $d_i = d_G(v_i)$ being the degree of vertex v_i $(1 \leq i \leq n)$, and A(G) is the adjacency matrix of G. The signless Laplacian spectral radius of G is the largest eigenvalue q(G) of its signless Laplacian Q(G). From spectral graph theory, we know that if graph G is connected, then there is a positive real eigenvector corresponding to q(G), and the unit positive eigenvector corresponding to q(G) is always called *Perron eigenvector*. For a graph G of order n, if $X = (x_1, x_2, \ldots, x_n)^T \in \mathbb{R}^n$ is a positive eigenvector corresponding to q(G) satisfying that $\sum_{i=1}^n x_i = 1$, then X is called a standard eigenvector corresponding to q(G).

Denote by K_n , C_n , P_n a complete graph, a cycle and a path of order n respectively. For a graph G, we let V(G), E(G) denote the vertex set and the edge set respectively. If there is no ambiguity, we use d(v) instead of $d_G(v)$. We use δ or $\delta(G)$ to denote the minimum vertex degree of a graph, use Δ or $\Delta(G)$ to denote the largest vertex degree of a graph, and use Δ' or $\Delta'(G)$ to denote the second largest vertex degree in a graph. In a graph, the notation $v_i \sim v_j$ denotes that vertex v_i is adjacent to v_j . Denote by $K_{s,t}$ a complete bipartite graph with one part of size s and another part of size t. In a graph G of order $n \geq 4$, for a vertex $u \in V(G)$, let $N_G(u)$ denote the neighbor set of u, and let $N_G[u] = \{u\} \cup N_G(u)$. $G(u) = G[N_G[u]], G^{\circ}(u) = G[N_G(u)]$ denote the subgraphs induced by $N_G[u], N_G(u)$ respectively.

The reader is referred to [1, 14] for the facts about planar and outer-planar graphs. A graph which can be drawn in the plane in such a way that edges meet only at points corresponding to their common ends is called a planar graph, and such a drawing is called a planar embedding of the graph. A simple planar graph is (edge) maximal if no edge can be added to the graph without violating planarity. In the planar embedding of a maximal planar graph G of order $n \geq 3$, each face is triangle. For a planar graph G, we have $m(G) \leq 3n - 6$ with equality if and only if it is maximal. In a maximal planar graph G of order $n \ge 4$, $\delta(G) \ge 3$. A graph G is *outer-planar* if it has a planar embedding, called *standard embedding*, in which all vertices lie on the boundary of its outer face. A simple outer-planar graph is (edge) maximal if no edge can be added to the graph without violating outer-planarity. In the standard embedding of a maximal outer-planar graph Gof order $n \geq 3$, the boundary of the outer face is a Hamiltonian cycle (a cycle contains all vertices) of G, and each of the other faces is triangle. For an outer-planar graph G, we have $m(G) \leq 2n-3$ with equality if and only if it is maximal. In a maximal planar graph G of order $n \ge 4$, for a vertex $u \in V(G)$, $G^{\circ}(u)$ is an outer-planar graph, and $G(u) = u \nabla G^{\circ}(u)$. From a nonmaximal planar graph G, by adding edges to G, a maximal planar graph G' can be obtained. From spectral graph theory, for a graph G, it is known that q(G+e) > q(G) if $e \notin E(G)$. Consequently, when we consider the maxima of the signless Laplacian spectral radius among planar graphs, it suffices to consider the maximal planar graphs directly.

Next we introduce some working lemmas.

Lemma 2.1 [21] Let u be a vertex of a maximal outer-planar graph on $n \ge 2$ vertices. Then $\sum_{v \sim u} d(v) \le n + 3d(u) - 4$.

Lemma 2.2 [17] Let G be a graph. Then $q(G) \leq \max_{u \in V(G)} \{ d_G(u) + \frac{1}{d_G(u)} \sum_{v \sim u} d_G(v) \}.$

Lemma 2.3 [8] Let G be a connected graph containing at least one edge. Then $q(G) \ge \Delta + 1$ with equality if and only if $G \cong K_{1,n-1}$.

3 Main results

Lemma 3.1 Let G be a maximal planar graph with order $n \ge 6$ and largest degree $\Delta(G)$. Then $q(G) \le \max_{u \in V(G)} \{ d_G(u) + 2 + \frac{3n - 9}{d_G(u)} \}$. Moreover, (i) if $\Delta(G) = n - 1$, then $q(G) \le n + 4 - \frac{6}{n-1}$; (ii) if $\Delta(G) = n - 2$, then $q(G) \le n + 3 - \frac{3}{n-2}$; (iii) if $\Delta(G) \le n - 3$, then $q(G) \le n + 2$. **Proof.** For any vertex $u \in V(G)$, let $N_G(u) = \{v_1, v_2, \dots, v_t\}$, $V_1 = V(G) \setminus N_G[u]$. For $1 \leq i \leq t$, let $\alpha_i = d_{G^\circ(u)}(v_i)$. Note that $m(G^\circ(u)) = |E(G^\circ(u))| = \frac{1}{2} \sum_{i=1}^t \alpha_i$. Between

 $N_G(u)$ and V_1 , there are $3n - 6 - \frac{1}{2} \sum_{i=1}^{l} \alpha_i - d_G(u)$ edges. Consequently,

$$\sum_{v \sim u} d_G(v) = d_G(u) + [3n - 6 - \frac{1}{2}\sum_{i=1}^t \alpha_i - d_G(u)] + \sum_{i=1}^t \alpha_i = 3n - 6 + \frac{1}{2}\sum_{i=1}^t \alpha_i.$$

From the narration in Section 2, we know that $G^{\circ}(u)$ is an outer-planar graph. Then $m(G^{\circ}(u)) \leq 2d_G(u) - 3$. As a result, $\sum_{v \sim u} d_G(v) \leq 3n - 9 + 2d_G(u)$, and

$$d_G(u) + \frac{1}{d_G(u)} \sum_{v \sim u} d_G(v) \le d_G(u) + 2 + \frac{3n - 9}{d_G(u)}$$

By Lemma 2.2, $q(G) \le \max_{u \in V(G)} \{ d_G(u) + 2 + \frac{3n - 9}{d_G(u)} \}.$

Let $f(x) = x + 2 + \frac{3n-9}{x}$. It can be checked that f(x) is convex. Note that $3 \le d_G(u) \le \Delta$. Then

$$d_G(u) + \frac{1}{d_G(u)} \sum_{v \sim u} d_G(v) \le \max\{5 + \frac{3n - 9}{3}, \Delta + 2 + \frac{3n - 9}{\Delta}\}.$$

By Lemma 2.2, then (i)-(iii) follow. This completes the proof. \Box

Let $\mathcal{H} = K_2 \nabla P_{n-2}$ for $n \ge 4$ (see Fig. 3.1) and let $\mathcal{H} = K_n$ for n = 1, 2, 3.



Fig 3.1. \mathcal{H}

Lemma 3.2 Suppose that the order n of \mathcal{H} is at least 5 (see Fig. 3.1). Then $q(\mathcal{H}) > n+2$.

Proof. Let $X = (x_1, x_2, \ldots, x_n)^T \in \mathbb{R}^n$ be a standard eigenvector corresponding to $q(\mathcal{H})$, where x_i corresponds to vertex v_i and $\sum_{i=1}^n x_i = 1$. By symmetry, $x_1 = x_2$, $x_3 = x_n$. By Lemma 2.3, $q(G) \ge n$.

Note that

$$q(G)x_1 = (n-1)x_1 + x_2 + \sum_{i=3}^n x_i = (n-2)x_1 + x_1 + x_2 + \sum_{i=3}^n x_i = (n-2)x_1 + 1.$$

Then $(q(G) - n + 2)x_1 = 1$, and

$$x_1 = \frac{1}{q(G) - n + 2}.$$
 (1)

Note that

$$q(G)\sum_{i=3}^{n} x_i = 3x_3 + 3x_n + 4\sum_{i=4}^{n-1} x_i + x_3 + x_n + 2\sum_{i=4}^{n-1} x_i + (n-2)x_1 + (n-2)x_2$$
$$= 3x_3 + 3x_n + 4\sum_{i=4}^{n-1} x_i + x_3 + x_n + 2\sum_{i=4}^{n-1} x_i + 2(n-2)x_1$$
$$= 6\sum_{i=3}^{n} x_n + 2(n-2)x_1 - 2(x_3 + x_n).$$

Then

$$(q(G) - 6) \sum_{i=3}^{n} x_i = 2(n-2)x_1 - 2(x_3 + x_n) = 2(n-2)x_1 - 4x_3,$$
$$\sum_{i=3}^{n} x_i = \frac{2(n-2)x_1 - 4x_3}{q(G) - 6},$$

and

$$1 = \sum_{i=1}^{n} x_i = \frac{2(n-2)x_1 - 4x_3}{q(G) - 6} + x_1 + x_2$$
$$= \frac{2(n-2)x_1 - 4x_3}{q(G) - 6} + \frac{2}{q(G) - n + 2}$$
$$= \frac{\frac{2(n-2)}{q(G) - n + 2} - 4x_3}{q(G) - 6} + \frac{2}{q(G) - n + 2}.$$

Then

$$x_3 = \frac{2(n-2) - (q(G) - n)(q(G) - 6)}{4(q(G) - n + 2)}.$$
 (2)

Note that $q(G)x_3 = 3x_3 + x_1 + x_2 + x_4$. Then

$$q(G)x_1 - q(G)x_3 = (n-2)x_1 - 2x_3 + \sum_{i=5}^n x_i$$
$$= (n-4)x_1 + 2x_1 - 2x_3 + \sum_{i=5}^n x_i,$$

and then

$$(q(G) - 2)(x_1 - x_3) = (n - 4)x_1 + \sum_{i=5}^n x_i.$$

This implies that $x_1 > x_3$. Combining with (1) and (2), we get

$$\frac{2(n-2) - (q(G) - n)(q(G) - 6)}{4(q(G) - n + 2)} < \frac{1}{q(G) - n + 2}.$$
 (3)

Simplifying (3), we get $q^{2}(G) - (6+n)q(G) + 4n + 8 > 0$. Then

$$q(G) > \frac{6+n+\sqrt{(6+n)^2-16(n+2)}}{2} = n+2.$$

This completes the proof. $\hfill \Box$

By the narration in Section 2 and Lemmas 3.1, 3.2, to consider the maxima of the signless Laplacian spectral radius among planar graphs of order $n \ge 5$, it suffices to consider only the graphs with $\Delta = n - 1$ and $\Delta = n - 2$.

Lemma 3.3 [18] Let A be an irreducible nonnegative square real matrix of order n and $X = (x_1, x_2, x_3, \ldots, x_n)^T$ be a real vector. For $1 \le i \le n$, let A_i be the *i*th row of A. If for any $1 \le i \le n$, $A_iX \le rx_i$, we say that $AX \le rX$.

Lemma 3.4 [23] Let A be an irreducible nonnegative square real matrix of order n and spectral radius ρ . If there exists a nonnegative real vector $y \neq 0$ and a real coefficient polynomial function f such that $f(A)y \leq ry$ $(r \in \mathbf{R})$, then $f(\rho) \leq r$.

Lemma 3.5 Let G be a maximal planar graph with order $n \ge 115$ and $d_G(v_1) = \Delta(G) = n-2$. In G, there are exactly $1 \le k \le 12$ vertices $v_2, v_3, \ldots, v_{k+1}$ such that $\frac{n}{6} + 1 \le d_G(v_i) \le n-61$, and for $k+2 \le i \le n$, $d_G(v_i) < \frac{n}{6} + 1$. Then $q(G) \le n-2$.

Proof. Note that in G, $d_G(v_i) \ge 3$ for $1 \le i \le n$, $\sum_{i=1}^n d_G(v_i) = 2(3n-6)$ and

$$k(\frac{n}{6}+1) + n - 2 + 3(n-k-1) = (\frac{k}{6}+4)n - 2k - 5 \qquad \text{(if } k \ge 13\text{)}$$
$$\ge 6n + \frac{n}{6} - 31 > 6n - 12.$$

Hence, $k \leq 12$. Let $X = (x_1, x_2, x_3, \ldots, x_n)^T$ be a positive vector satisfying that x_i corresponds to vertex v_i and

$$x_{i} = \begin{cases} 1, & i = 1; \\ \frac{1}{k}, & 2 \le i \le k+1; \\ \frac{3}{n-k-1}, & k+2 \le i \le n \end{cases}$$

For x_1 ,

$$\frac{(n-2)x_1 + \sum_{v_j \sim v_1} x_j}{x_1} \le n-2 + 1 + \frac{3(n-k-2)}{n-k-1} < n+2.$$

For x_i $(k+2 \le i \le n)$, when $d(v_i) \ge k+1$,

$$\frac{d_G(v_i)x_i + \sum_{v_j \sim v_i} x_j}{x_i} \le d_G(v_i) + \frac{\sum_{j=1}^{k+1} x_j + \frac{3(d_G(v_i) - k - 1)}{n - k - 1}}{\frac{3}{n - k - 1}}$$

$$\leq d_G(v_i) + \frac{2 + \frac{3(d_G(v_i) - k - 1)}{n - k - 1}}{\frac{3}{n - k - 1}}$$
$$\leq d_G(v_i) + \frac{2}{3}(n - k - 1) + d_G(v_i) - k - 1$$
$$= 2d_G(v_i) + \frac{2}{3}n - \frac{5}{3}k - \frac{5}{3} \leq n + 2;$$

when $d_G(v_i) \leq k$,

$$\frac{d_G(v_i)x_i + \sum_{v_j \sim v_i} x_j}{x_i} \le d_G(v_i) + \frac{\sum_{j=1}^{d_G(v_i)} x_j}{\frac{3}{n-k-1}}$$
$$\le d_G(v_i) + \frac{1 + \frac{d_G(v_i) - 1}{k}}{\frac{3}{n-k-1}}$$
$$= d_G(v_i) + \frac{n-k-1}{3} + \frac{(d_G(v_i) - 1)(n-k-1)}{3k}$$
$$< k + \frac{n-k-1}{3} + \frac{n-k-1}{3}$$
$$= \frac{2(n-1)+k}{3} < n+2.$$

For x_i $(2 \le i \le k+1)$, noting that $d_G(v_i) \ge \frac{n}{6} + 1 > k$, we get

$$\frac{d_G(v_i)x_i + \sum_{v_j \sim v_i} x_j}{x_i} = \frac{(d_G(v_i) - 1)x_i + x_i + \sum_{v_j \sim v_i} x_j}{x_i} \\
\leq d_G(v_i) - 1 + \frac{\sum_{j=1}^{k+1} x_j + \frac{3(d_G(v_i) - k)}{n-k-1}}{\frac{1}{k}} \\
= d_G(v_i) - 1 + \frac{2 + \frac{3(d_G(v_i) - k)}{n-k-1}}{\frac{1}{k}} \\
= (1 + \frac{3k}{n-k-1})d_G(v_i) - \frac{3k^2}{n-k-1} + 2k - 1. \quad (4)$$

Let $f(k) = (1 + \frac{3k}{n-k-1})d_G(v_i) - \frac{3k^2}{n-k-1} + 2k - 1$. Taking derivation of f(k) with respect to k, we get $f'(k) = \frac{(n-k-1)(2n-2+3d_G(v_i)-8k)+3kd_G(v_i)-3k^2}{(n-k-1)^2}$. Note that $n \ge 115$. As a result, if $d_G(v_i) \ge k$, then f'(k) > 0. This implies that f(k) is monotone increasing with respect to k. Note that $d_G(v_i) \le n - 61$. Then (4) < n + 2.

Then $Q(G)X \leq (n+2)X$. By Lemma 3.4, $q(G) \leq n+2$. This completes the proof.

Lemma 3.6 Let G be a maximal planar graph with order $n \ge 380$, $d_G(v_1) = \Delta(G) = n-2$, and $\Delta'(G) \ge n-62$. Then $q(G) \le n+2$. **Proof.** Suppose $d_G(v_2) = \Delta'(G)$. Let $X = (x_1, x_2, x_3, \ldots, x_n)^T$ be a positive vector satisfying that x_i corresponds to vertex v_i and

$$x_i = \begin{cases} 1, & i = 1; \\ 1, & i = 2; \\ \frac{3}{n-2}, & 3 \le i \le n. \end{cases}$$

For v_1 ,

$$\frac{(n-2)x_1 + \sum_{v_j \sim v_1} x_j}{x_1} \le n - 2 + 1 + \frac{3(n-3)}{n-2} < n+2$$

Next, there are two cases to consider.

Case 1 $v_2 \in N_G(v_1)$. Suppose $N_G(v_1) = \{v_2, v_3, \ldots, v_{n-2}, v_{n-1}\}$. Suppose v_1 is in the outer face of $G_{v_1}^{\circ}$. Then v_n is in one of the inner faces of $G_{v_1}^{\circ}$ (see Fig. 3.2).

For
$$v_2$$
,

$$\frac{d_G(v_2)x_2 + \sum_{v_j \sim v_2} x_j}{x_2} \le d_G(v_2) + 1 + \frac{3(d_G(v_2) - 1)}{n - 2}$$
$$= d_G(v_2) + 1 + \frac{3d_G(v_2)}{n - 2} - \frac{3}{n - 2}$$
$$\le n + 2 - \frac{3}{n - 2}. \qquad (d_G(v_2) \le n - 2)$$

Denote by $C_{v_1} = v_2 v_3 \cdots v_{n-2} v_{n-1} v_2$ the Hamiltonian cycle in $G_{v_1}^{\circ}$. Suppose that $v_i s$ $(2 \le i \le n-1)$ are distributed along clockwise direction on C_{v_1} and suppose $N_{G_{v_1}^{\circ}}(v_2) = \{v_{2_1}, v_{2_2}, \ldots, v_{2_t}\}$, where for $1 \le i \le t-1$, $2_i < 2_{i+1}$, $v_{2_1} = v_3$, $v_{2_t} = v_{n-1}$ (see Fig. 3.2). For $1 \le j \le t$, suppose there are l_{j-1} vertices between $v_{2_{j-1}}$ and v_{2_j} along clockwise direction on C_{v_1} , where if j = 1, we let $v_{2_0} = v_2$. Along clockwise direction on C_{v_1} , suppose there are l_t vertices between v_{2_t} and v_2 .



For each v_{2_j} $(1 \le j \le t$, see Fig. 3.2), noting that

$$l_{j-1} + l_j \le n - 3 - d_G(v_2)$$

and $d_G(v_2) \ge n - 62$, then

$$d_G(v_{2_j}) \le l_{j-1} + l_j + 5 \le n + 2 - d_G(v_2) \le 64,$$

and then

$$\frac{d_G(v_{2_j})x_{2_j} + \sum_{v_k \sim v_{2_j}} x_k}{x_{2_j}} \le d_G(v_{2_j}) + \frac{2 + \frac{3(d_G(v_{2_j}) - 2)}{n - 2}}{\frac{3}{n - 2}}$$
$$= 2d_G(v_{2_j}) - 2 + \frac{2}{3}(n - 2)$$
$$\le 126 + \frac{2}{3}(n - 2) \le n + 2. \qquad (n \ge 380)$$

For each $v_f \in (N_G(v_1) \setminus \{v_2, v_{2_1}, v_{2_2}, \ldots, v_{2_t}\})$, then along clockwise direction on C_{v_1} , there exists $0 \leq s \leq t$ such that v_f is between v_{2_s} and $v_{2_{s+1}}$, where $v_{2_{t+1}} = v_2$ (see Fig. 3.3). Note that

$$l_s \le n - 3 - d_G(v_2).$$

Then $d_G(v_f) \leq l_s + 3 \leq n - d_G(v_2) \leq 62$. As a result,

$$\frac{d_G(v_f)x_f + \sum_{v_k \sim v_f} x_k}{x_f} \le d_G(v_f) + \frac{2 + \frac{3(d_G(v_f) - 2)}{n - 2}}{\frac{3}{n - 2}}$$
$$= 2d_G(v_f) - 2 + \frac{2}{3}(n - 2)$$
$$\le 122 + \frac{2}{3}(n - 2) \le n + 2. \qquad (n \ge 380)$$



For v_n , note that v_n is in one of the inner faces of $G_{v_1}^{\circ}$. Suppose that in $G_{v_1}^{\circ}$, v_n is in a face $v_2v_{2_z}v_{2_z+1}v_{2_z+2}\cdots v_{2_{z+1}}v_2$ (see Fig. 3.4). Note that $l_z \leq n-3-d_G(v_2)$ and $d_G(v_n) \leq l_z + 3$. Then $d_G(v_n) \leq n-d_G(v_2) \leq 62$, and

$$\frac{d_G(v_n)x_n + \sum_{v_k \sim v_n} x_k}{x_n} \le d_G(v_n) + \frac{1 + \frac{3(d_G(v_n) - 1)}{n - 2}}{\frac{3}{n - 2}}$$
$$= 2d_G(v_n) - 1 + \frac{n - 2}{3}$$
$$\le 123 + \frac{1}{3}(n - 2) < n + 2 \qquad (n \ge 380).$$

Case 2 $v_2 \notin N_G(v_1)$. Suppose that v_1 is in the outer face of $G_{v_1}^{\circ}$. Then v_2 is in one of the inner faces of $G_{v_1}^{\circ}$. Then $N_G(v_1) = \{v_3, v_4, v_5, \ldots, v_{n-1}, v_n\}$. Suppose that

 $C_{v_1} = v_3 v_4 \cdots v_{n-1} v_n v_3$ is the Hamiltonian cycle in $G_{v_1}^{\circ}$, $v_i s$ $(3 \leq i \leq n)$ are distributed along clockwise direction on C_{v_1} , and suppose $N_{G_{v_1}^{\circ}}(v_2) = \{v_{2_1}, v_{2_2}, \ldots, v_{2_t}\}$, where for $1 \leq i \leq t-1, 2_i < 2_{i+1}$ (see Fig. 3.5). For $2 \leq j \leq t$, along clockwise direction on C_{v_1} , suppose there are l_{j-1} vertices between $v_{2_{j-1}}$ and v_{2_j} . Suppose that there are l_t vertices between v_{2_t} and v_{2_1} .

For each v_{2_j} $(2 \le j \le t)$, noting that $l_{j-1} + l_j \le n - 2 - d_G(v_2)$ and $d_G(v_2) \ge n - 62$, then $d_G(v_{2_j}) \le l_{j-1} + l_j + 4 \le n + 2 - d_G(v_2) \le 64$; for v_{2_1} , noting that $l_1 + l_t \le n - 2 - d_G(v_2)$, then $d_G(v_{2_1}) \le l_1 + l_t + 4 \le n + 2 - d_G(v_2) \le 64$. Then for each v_{2_j} $(1 \le j \le t)$,

$$\frac{d_G(v_{2j})x_{2j} + \sum_{v_k \sim v_{2j}} x_k}{x_{2j}} \le d_G(v_{2j}) + \frac{2 + \frac{3(d_G(v_{2j}) - 2)}{n - 2}}{\frac{3}{n - 2}}$$
$$= 2d_G(v_{2j}) - 2 + \frac{2}{3}(n - 2)$$
$$\le 126 + \frac{2}{3}(n - 2) \le n + 2 \qquad (n \ge 380).$$

For each $v_i \in (N_G(v_1) \setminus \{v_{2_1}, v_{2_2}, \ldots, v_{2_t}\})$, along clockwise direction on C_{v_1}, v_i is between v_{2_k} and $v_{2_{k+1}}$ for some $1 \le k \le t-1$, or v_i is between v_{2_t} and v_{2_1} . Then for some $1 \le k \le t$, $d_G(v_i) \le l_k + 2$. Note that $l_k \le n-2 - d_G(v_2)$. Then $d_G(v_i) \le n - d_G(v_2) \le 62$, and

$$\frac{d_G(v_i)x_i + \sum_{v_k \sim v_i} x_k}{x_i} \le d_G(v_i) + \frac{1 + \frac{3(d_G(v_i) - 1)}{n - 2}}{\frac{3}{n - 2}}$$
$$= 2d_G(v_i) - 1 + \frac{1}{3}(n - 2)$$
$$\le 123 + \frac{1}{3}(n - 2) < n + 2 \qquad (n \ge 380)$$

For v_2 ,

$$\frac{d_G(v_2)x_2 + \sum_{v_k \sim v_2} x_k}{x_2} \le d_G(v_2) + \frac{3d_G(v_2)}{n-2} \le n+1. \qquad (d_G(v_2) \le n-2)$$

By above discussion, we get $Q(G)X \leq (n+2)X$. By Lemma 3.4, we get that $q(G) \leq n+2$. This completes the proof. \Box

Lemma 3.7 Let G be a maximal planar graph with order $n \ge 4$ and $d(v_1) = \Delta(G) = n-2$. If for $2 \le i \le n$, $d_G(v_i) < 1 + \frac{n}{6}$. Then $q(G) \le n-2$.

Proof. Let $X = (x_1, x_2, x_3, ..., x_n)^T$ be a positive vector satisfying that x_i corresponds to vertex v_i and

$$x_i = \begin{cases} 1, & i = 1; \\ \frac{4}{n-1}, & 2 \le i \le n \end{cases}$$

For v_1 ,

$$\frac{(n-2)x_1 + \sum_{v_j \sim v_1} x_j}{x_1} \le n - 2 + \frac{4(n-2)}{n-1} < n+2.$$

For $v_i \ (2 \le i \le n)$,

$$\frac{d_G(v_i)x_i + \sum_{v_j \sim v_i} x_j}{x_i} \le d_G(v_i) + \frac{1 + \frac{4(d_G(v_i) - 1)}{n - 1}}{\frac{4}{n - 1}} \le 2d_G(v_i) - 1 + \frac{n - 1}{4} \le \frac{7n}{12} + \frac{3}{4} < n + 2.$$

As a result, $Q(G)X \leq (n+2)X$. By Lemma 3.4, we get that $q(G) \leq n+2$. This completes the proof. \Box

Theorem 3.8 Let G be a maximal planar graph with order $n \ge 380$ and $\Delta(G) = n - 2$. Then $q(G) \le n + 2$.

Proof. This theorem follows from Lemmas 3.5-3.7.

Lemma 3.9 Let G be a maximal planar graph with order $n \ge 91$ and $d_G(v_1) = \Delta(G) = n-1$. There are exactly $1 \le k \le 13$ vertices $v_2, v_3, \ldots, v_{k+1}$ in G such that

$$\frac{n}{7} + \frac{19}{7} \le d(v_i) \le n - 75.$$

For $k+2 \le i \le n$, $d_G(v_i) < \frac{n}{7} + \frac{19}{7}$. Then $q(G) \le n+2$.

Proof. Note that in *G*, $d_G(v_i) \ge 3$ for $1 \le i \le n$, $\sum_{i=1}^n d_G(v_i) = 2(3n-6)$, and note that if $k \ge 14$,

$$k(\frac{n}{7} + \frac{19}{7}) + n - 1 + 3(n - k - 1) \ge 14(\frac{n}{7} + \frac{19}{7}) + n - 1 + 3(n - 15) > 6n - 12.$$

Hence, $k \leq 13$. Let $X = (x_1, x_2, x_3, \ldots, x_n)^T$ be a positive vector satisfying that x_i corresponds to vertex v_i and

$$x_{i} = \begin{cases} 1, & i = 1; \\ \frac{2}{3k}, & 2 \le i \le k+1; \\ \frac{7}{3(n-k-1)}, & k+2 \le i \le n. \end{cases}$$

For v_1 ,

$$\frac{(n-1)x_1 + \sum_{v_j \sim v_1} x_j}{x_1} = n - 1 + 3 = n + 2.$$

For v_i $(k+2 \le i \le n)$, if $d_G(v_i) \ge k+1$, then

$$\frac{d_G(v_i)x_i + \sum_{v_j \sim v_i} x_j}{x_i} \le d_G(v_i) + \frac{\frac{5}{3} + \frac{7(d_G(v_i) - k - 1)}{3(n - k - 1)}}{\frac{7}{3(n - k - 1)}} \le 2d_G(v_i) - k - 1 + \frac{5}{7}(n - k - 1) \le 2d_G(v_i) - \frac{12k}{7} - \frac{12}{7} + \frac{5}{7}n \le n + 2;$$

if $d_G(v_i) \leq k$, then

$$\frac{d_G(v_i)x_i + \sum_{v_j \sim v_i} x_j}{x_i} \le d_G(v_i) + \frac{\frac{5}{3}}{\frac{7}{3(n-k-1)}} \le d_G(v_i) + \frac{5}{7}(n-k-1) < n+2.$$

For v_i $(2 \le i \le k+1)$, $d_G(v_i) \ge k$, then

$$\frac{d_G(v_i)x_i + \sum_{v_j \sim v_i} x_j}{x_i} = \frac{(d_G(v_i) - 1)x_i + x_i + \sum_{v_j \sim v_i} x_j}{x_i}$$

$$\leq d_G(v_i) - 1 + \frac{\sum_{j=1}^{k+1} x_j + \frac{7(d_G(v_i) - k)}{3(n-k-1)}}{\frac{2}{3k}}$$

$$\leq d_G(v_i) - 1 + \frac{\frac{5}{3} + \frac{7(d_G(v_i) - k)}{3(n-k-1)}}{\frac{2}{3k}}$$

$$\leq d_G(v_i) - 1 + \frac{5}{2}k + \frac{\frac{7}{2}k(d_G(v_i) - k)}{n-k-1}.$$
(5)

As the proof of Lemma 3.5, noting that $d_G(v_i) \leq n - 75$, we can prove that $(5) \leq n + 2$. Then $Q(G)X \leq (n+2)X$. By Lemma 3.4, we get that $q(G) \leq n + 2$. This completes the proof. \Box

Lemma 3.10 Let G be a maximal planar graph with order $n \ge 6$ and $d(v_1) = \Delta(G) = n-1$. If for $2 \le i \le n$, $d_G(v_i) < \frac{n}{7} + \frac{19}{7}$. Then $q(G) \le n+2$.

Proof. Let $X = (x_1, x_2, x_3, ..., x_n)^T$ be a positive vector satisfying that x_i corresponds to vertex v_i and

$$x_{i} = \begin{cases} 1, & i = 1; \\ \\ \frac{3}{n-1}, & 2 \le i \le n \end{cases}$$

For v_1 ,

$$\frac{(n-1)x_1 + \sum_{v_j \sim v_1} x_j}{x_1} = n - 1 + 3 = n + 2.$$

For $v_i \ (2 \le i \le n)$,

$$\frac{d_G(v_i)x_i + \sum_{v_j \sim v_i} x_j}{x_i} \le d_G(v_i) + \frac{1 + \frac{3(d_G(v_i) - 1)}{n - 1}}{\frac{3}{n - 1}}$$

$$= 2d_G(v_i) - 1 + \frac{n-1}{3}$$

$$< \frac{2n}{7} + \frac{31}{7} + \frac{n-1}{3}$$

$$\le n+2. \qquad (n \ge 6)$$

As a result, $Q(G)X \leq (n+2)X$. By Lemma 3.4, we get that $q(G) \leq n+2$. This completes the proof. \Box

Lemma 3.11 Let G be a maximal planar graph with order $n \ge 461$, $d_G(v_1) = \Delta(G) = n-1$ and $n - 81 \le \Delta'(G) \le n - 4$. Then $q(G) \le n + 2$.

Proof. Suppose $d_G(v_2) = \Delta'(G)$. Let $X = (x_1, x_2, x_3, \ldots, x_n)^T$ be a positive vector satisfying that x_i corresponds to vertex v_i and

$$x_{i} = \begin{cases} 1, & i = 1; \\ \frac{4}{7}, & i = 2; \\ \frac{17}{7(n-2)}, & 3 \le i \le n \end{cases}$$

For v_1 ,

$$\frac{(n-1)x_1 + \sum_{v_j \sim v_1} x_j}{x_1} = n - 1 + 3 = n + 2.$$

For v_2 ,

$$\frac{d_G(v_2)x_2 + \sum_{v_j \sim v_2} x_j}{x_2} \le d_G(v_2) + \frac{1 + \frac{17(d_G(v_2) - 1)}{7(n - 2)}}{\frac{4}{7}}$$
$$= d_G(v_2) + \frac{7}{4} + \frac{17(d_G(v_2) - 1)}{4(n - 2)}$$
$$< n + 2. \qquad (d_G(v_2) \le n - 4)$$

Suppose that v_1 is in the outer face of $G_{v_1}^{\circ}$, $C_{v_1} = v_2 v_3 \cdots v_{n-1} v_n v_2$ is the Hamiltonian cycle in $G_{v_1}^{\circ}$, $v_i s$ $(2 \le i \le n)$ are distributed along clockwise direction on C_{v_1} , and suppose $N_{G_{v_1}^{\circ}}(v_2) = \{v_{2_1}, v_{2_2}, \ldots, v_{2_t}\}$, where for $1 \le i \le t-1$, $2_i < 2_{i+1}, v_{2_1} = v_3, v_{2_t} = v_n$. On C_{v_1} , along clockwise direction, for $1 \le j \le t$, suppose that there are l_{j-1} vertices between $v_{2_{j-1}}$ and v_{2_j} , where if j = 1, we let $v_{2_0} = v_2$. Along clockwise direction on C_{v_1} , suppose that there are l_t vertices between v_{2_t} and v_2 . For each v_{2_j} $(1 \le j \le t)$, noting that $l_{j-1} + l_j \le n-2 - d_G(v_2)$ and $d_G(v_2) \ge n-81$, then $d_G(v_{2_j}) \le l_{j-1} + l_j + 4 \le n+2 - d_G(v_2) \le 83$, and then

$$\frac{d_G(v_{2_j})x_{2_j} + \sum_{v_k \sim v_{2_j}} x_k}{x_{2_j}} \le d_G(v_{2_j}) + \frac{\frac{11}{7} + \frac{17(d_G(v_{2_j}) - 2)}{7(n-2)}}{\frac{17}{7(n-2)}} = 2d_G(v_{2_j}) - 2 + \frac{11}{17}(n-2)$$

$$\leq 164 + \frac{11}{17}(n-2)$$

 $\leq n+2. \qquad (n \geq 456)$

For each $v_f \in (N_G(v_1) \setminus \{v_2, v_{2_1}, v_{2_2}, \ldots, v_{2_t}\})$, along clockwise direction, then there exists $0 \leq s \leq t$ such that v_f is between v_{2_s} and $v_{2_{s+1}}$ on C_{v_1} , and then $d_G(v_f) \leq l_s + 2 \leq n - d_G(v_2) \leq 81$. As a result,

$$\frac{d_G(v_f)x_f + \sum_{v_k \sim v_f} x_k}{x_f} \le d_G(v_f) + \frac{\frac{11}{7} + \frac{17(d_G(v_f) - 2)}{7(n - 2)}}{\frac{17}{7(n - 2)}}$$
$$= 2d_G(v_f) - 2 + \frac{11}{17}(n - 2)$$
$$\le 160 + \frac{11}{17}(n - 2)$$
$$\le n + 2. \qquad (n \ge 456)$$

As a result, $Q(G)X \leq (n+2)X$. By Lemma 3.4, we get that $q(G) \leq n+2$. This completes the proof. \Box

Lemma 3.12 Let G be a maximal planar graph with order $n \ge 15$, $d_G(v_1) = \Delta(G) = n-1$. (i) if $\Delta'(G) = n - 2$, then $q(G) < q(\mathcal{H})$; (ii) if $\Delta'(G) = n - 3$, then $q(G) < q(\mathcal{H})$.



Fig. 3.6. $D_1 - D_4$

Proof. Suppose $d_G(v_2) = \Delta'(G)$, v_1 is in the outer face of $G_{v_1}^{\circ}$, and suppose that $C_{v_1} = v_2 v_3 \cdots v_{n-1} v_n v_2$ is the Hamiltonian cycle in $G_{v_1}^{\circ}$ (see Fig. 3.6).

(i) Suppose $d_G(v_2) = \Delta'(G) = n - 2$ and $v_k \notin N_G(v_2)$ $(4 \le k \le n - 1)$. Then $G_{v_1}^{\circ} \cong D_1$ (see Fig. 3.6). For convenience, we suppose $G_{v_1}^{\circ} = D_1$. By Lemma 2.3, we get that q(G) > 15. Let $X = (x_1, x_2, \ldots, x_n)^T \in \mathbb{R}^n$ be the Perron eigenvector corresponding to q(G), where x_i corresponds to vertex v_i .

Note that

$$q(G)x_k = 3x_k + x_{k-1} + x_{k+1} + x_1, (6)$$

$$q(G)x_2 = (n-2)x_2 + x_1 + \sum_{3 \le i \le n, i \ne k} x_i.$$
 (7)

(6), (7) tell us that

$$q(G)x_2 - q(G)x_k = (n-2)x_2 - 3x_k + \sum_{3 \le i \le k-2} x_i + \sum_{k+2 \le i \le n} x_i,$$
$$(q(G) - 3)(x_2 - x_k) = (n-5)x_2 + \sum_{3 \le i \le k-2} x_i + \sum_{k+2 \le i \le n} x_i.$$

Because $n \ge 15$, it follows immediately that $x_2 > x_k$.

Note that

$$q(G)x_{k-1} = 5x_{k-1} + x_1 + x_2 + x_{k-2} + x_k + x_{k+1},$$

$$q(G)x_{k+1} = 5x_{k+1} + x_1 + x_2 + x_{k-1} + x_k + x_{k+2}.$$

Then

$$q(G)(x_{k-1} + x_{k+1}) = 6(x_{k-1} + x_{k+1}) + 2(x_1 + x_k + x_2) + x_{k-2} + x_{k+2}.$$
(8)

From (6) and (7), we also get that

$$q(G)(x_2 + x_k) = (n-2)x_2 + 3x_k + 2x_1 + 2x_{k-1} + 2x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+2 \le i \le n} x_i.$$
 (9)

By (9)-(8), we get that

$$q(G)(x_2 + x_k) - q(G)(x_{k-1} + x_{k+1}) = (n - 11)x_2 + 4(x_2 + x_k) - 4(x_{k-1} + x_{k+1}) + 3(x_2 - x_k) + \sum_{3 \le i \le k-3} x_i + \sum_{k+3 \le i \le n} x_i.$$

It follows that

$$(q(G)-4)[x_2+x_k-(x_{k-1}+x_{k+1})] = (n-11)x_2+3(x_2-x_k) + \sum_{3 \le i \le k-3} x_i + \sum_{k+3 \le i \le n} x_i.$$
 (10)

Because $n \ge 15$, (10) tells us that $x_2 + x_k > x_{k-1} + x_{k+1}$.

Let $F = G - v_{k-1}v_{k+1} + v_2v_k$. Note the relation between the Rayleigh quotient and the largest eigenvalue of a non-negative real symmetric matrix, and note that

$$X^{T}Q(F)X - X^{T}Q(G)X = (x_{2} + x_{k})^{2} - (x_{k-1} + x_{k+1})^{2}.$$

It follows that when $n \ge 15$, then $q(F) > X^T Q(F) X > X^T Q(G) X = q(G)$. Because $F \cong \mathcal{H}, q(\mathcal{H}) > q(G)$. Then (i) follows.

(ii) Suppose $d_G(v_2) = \Delta'(G) = n-3$. Then there are three cases for G, that is, $G \cong D_2$, $G \cong D_3$ or $G \cong D_4$ (see Fig. 3.6). By Lemma 2.3, we know that q(G) > 15.

Case 1 $G \cong D_2$. For convenience, we suppose that $G = D_2$. Because $d_G(v_2) = \Delta'(G) = n - 3, 4 \le k \le n - 2$. Let $X = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$ be the Perron eigenvector corresponding to q(G), where x_i corresponds to vertex v_i .

Note that

$$q(G)x_{k+1} = 3x_{k+1} + x_1 + x_k + x_{k+2},$$
(11)

$$q(G)x_k = 4x_k + x_1 + x_{k-1} + x_{k+1} + x_{k+2}.$$
(12)

By (12)-(11), we get

$$q(G)(x_k - x_{k+1}) = 3x_k - 2x_{k+1} + x_{k-1}.$$

Then

$$(q(G) - 2)(x_k - x_{k+1}) = x_k + x_{k-1}.$$
(13)

Because $n \ge 15$, (13) implies $x_k > x_{k+1}$.

Note that

$$q(G)x_2 = (n-3)x_2 + x_1 + \sum_{3 \le i \le k-1} x_i + \sum_{k+2 \le i \le n} x_i.$$
 (14)

By (14)+(12), we get

$$q(G)(x_2+x_k) = (n-3)x_2+2x_1+4x_k+2x_{k-1}+x_{k+1}+2x_{k+2}+\sum_{3\le i\le k-2}x_i+\sum_{k+3\le i\le n}x_i.$$
 (15)

Note that

$$q(G)x_{k-1} = 5x_{k-1} + x_1 + x_2 + x_{k-2} + x_k + x_{k+2}.$$
(16)

By (16)+(11), we get that

$$q(G)(x_{k-1} + x_{k+1}) = 5x_{k-1} + x_{k-2} + 2x_1 + x_2 + 2x_k + 3x_{k+1} + 2x_{k+2}.$$
 (17)

By (17)-(12), we get that

$$q(G)(x_{k-1} + x_{k+1} - x_k) = x_1 + x_2 + x_{k-2} + 4x_{k-1} + 2x_{k+1} + x_{k+2} - 2x_k.$$

Then

$$(q(G) - 2)(x_{k-1} + x_{k+1} - x_k) = x_1 + x_2 + x_{k-2} + 2x_{k-1} + x_{k+2} > 0.$$
(18)

Because $n \ge 15$, (18) implies that $x_{k-1} + x_{k+1} > x_k$.

By (14)-(11), we get that

$$q(G)(x_2 - x_{k+1}) = (n-3)x_2 + x_{k-1} - x_k - 3x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+3 \le i \le n} x_i > 0.$$

Then

$$(q(G) - 4)(x_2 - x_{k+1}) = (n - 7)x_2 + x_{k-1} + x_{k+1} - x_k + \sum_{3 \le i \le k-2} x_i + \sum_{k+3 \le i \le n} x_i > 0.$$
(19)

Because $n \ge 15$, (19) implies $x_2 > x_{k+1}$.

By (14)-(12), we get that

$$q(G)(x_2 - x_k) = (n - 3)x_2 - 4x_k - x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+3 \le i \le n} x_i$$

Then

$$(q(G) - 4)(x_2 - x_k) = (n - 8)x_2 + x_2 - x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+3 \le i \le n} x_i > 0.$$
(20)

Because $n \ge 15$, (20) implies that $x_2 > x_k$.

By (14)-(16), we get that

$$q(G)(x_2 - x_{k-1}) = (n-4)x_2 - 4x_{k-1} - x_k + \sum_{3 \le i \le k-3} x_i + \sum_{k+3 \le i \le n} x_i.$$

Then

$$(q(G) - 4)(x_2 - x_{k-1}) = (n - 9)x_2 + x_2 - x_k + \sum_{3 \le i \le k-3} x_i + \sum_{k+3 \le i \le n} x_i.$$
 (21)

Because $n \ge 15$, (21) implies that $x_2 > x_{k-1}$.

Note that

$$q(G)x_{k+2} = 6x_{k+2} + x_{k+1} + x_k + x_{k-1} + x_{k+3} + x_2 + x_1.$$
(22)

By (14)-(22), we get that

$$q(G)(x_2 - x_{k+2}) = (n-4)x_2 - 5x_{k+2} - x_k - x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+4 \le i \le n} x_i.$$

Then

$$(q(G) - 5)(x_2 - x_{k+2}) = (n - 11)x_2 + x_2 - x_k + x_2 - x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+4 \le i \le n} x_i.$$
 (23)

Because $n \ge 15$, (23) implies that $x_2 > x_{k+2}$.

By (16)+(22), we get that

$$q(G)(x_{k-1} + x_{k+2}) = 2x_1 + 2x_2 + x_{k-2} + 6x_{k-1} + 2x_k + x_{k+1} + 7x_{k+2} + x_{k+3}.$$
 (24)

By (15)-(24), we get that

$$q(G)(x_{2} + x_{k}) - q(G)(x_{k-1} + x_{k+2})$$

$$= (n-5)x_{2} - 4x_{k-1} + 2x_{k} - 5x_{k+2} + \sum_{3 \le i \le k-3} x_{i} + \sum_{k+4 \le i \le n} x_{i}$$

$$= (n-14)x_{2} + 4x_{2} - 4x_{k-1} + 2x_{k} + 5x_{2} - 5x_{k+2} + \sum_{3 \le i \le k-3} x_{i} + \sum_{k+4 \le i \le n} x_{i}.$$
(25)

Because $n \ge 15$, (25) implies that $x_2 + x_k > x_{k-1} + x_{k+2}$.

Let $\mathbb{F} = G - v_{k-1}v_{k+2} + v_2v_k$. Note that $X^TQ(\mathbb{F})X - X^TQ(G)X = (x_2 + x_k)^2 - (x_{k-1} + x_{k+2})^2$. It follows that when $n \ge 15$, then $q(\mathbb{F}) > X^TQ(\mathbb{F})X > X^TQ(G)X = q(G)$. By (i), it follows immediately that $q(\mathcal{H}) > q(\mathbb{F}) > q(G)$.

Case 2 $G \cong D_3$.

For convenience, we suppose that $G = D_3$. Because $d_G(v_2) = \Delta'(G) = n - 3, 5 \le k \le n - 2$. Let $X = (x_1, x_2, \ldots, x_n)^T \in \mathbb{R}^n$ be the Perron eigenvector corresponding to q(G), where x_i corresponds to vertex v_i .

Note that

$$q(G)x_{k+1} = 3x_{k+1} + x_1 + x_k + x_{k+2},$$
(26)

$$q(G)x_{k-1} = 3x_{k-1} + x_1 + x_k + x_{k-2},$$
(27)

$$q(G)x_2 = (n-3)x_2 + x_1 + x_k + \sum_{3 \le i \le k-2} x_i + \sum_{k+2 \le i \le n} x_i.$$
 (28)

Then

$$q(G)x_2 - q(G)x_{k+1} = (n-3)x_2 - 3x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+3 \le i \le n} x_i,$$
$$(q(G) - 3)(x_2 - x_{k+1}) = (n-6)x_2 + \sum_{3 \le i \le k-2} x_i + \sum_{k+3 \le i \le n} x_i > 0.$$

This implies that $x_2 > x_{k+1}$. By (28)-(27), we get that

$$q(G)x_2 - q(G)x_{k-1} = (n-3)x_2 - 3x_{k-1} + \sum_{3 \le i \le k-3} x_i + \sum_{k+2 \le i \le n} x_i$$

Then

$$(q(G) - 3)(x_2 - x_{k-1}) = (n - 6)x_2 + \sum_{3 \le i \le k-3} x_i + \sum_{k+2 \le i \le n} x_i > 0.$$

This implies that $x_2 > x_{k-1}$. Note that

$$q(G)x_k = 6x_k + x_1 + x_2 + x_{k-2} + x_{k-1} + x_{k+1} + x_{k+2},$$
(29)

$$q(G)x_{k+2} = 5x_{k+2} + x_1 + x_2 + x_{k+1} + x_k + x_{k+3}.$$
(30)

By (28)-(29), we get

$$q(G)x_2 - q(G)x_k = (n-4)x_2 - 5x_k - x_{k-1} - x_{k+1} + \sum_{3 \le i \le k-3} x_i + \sum_{k+3 \le i \le n} x_i.$$

Then

$$(q(G) - 5)(x_2 - x_k) = (n - 11)x_2 + 2x_2 - x_{k-1} - x_{k+1} + \sum_{3 \le i \le k-3} x_i + \sum_{k+3 \le i \le n} x_i.$$

This implies that $x_2 > x_k$. By (28)-(30), we get

$$q(G)x_2 - q(G)x_{k+2} = (n-4)x_2 - 4x_{k+2} - x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+4 \le i \le n} x_i.$$

Then

$$(q(G) - 4)(x_2 - x_{k+2}) = (n - 9)x_2 + x_2 - x_{k+1} + \sum_{3 \le i \le k-2} x_i + \sum_{k+4 \le i \le n} x_i.$$

This implies that $x_2 > x_{k+2}$. By (26)+(28), we get that

$$q(G)(x_2 + x_{k+1}) = (n-3)x_2 + 3x_{k+1} + 2x_1 + 2x_k + 2x_{k+2} + \sum_{3 \le i \le k-2} x_i + \sum_{k+3 \le i \le n} x_i.$$
 (31)

By (29)+(30), we get that

$$q(G)(x_k + x_{k+2}) = 2x_1 + 2x_2 + 2x_{k+1} + 7x_k + 6x_{k+2} + x_{k-1} + x_{k-2} + x_{k+3}.$$
(32)

By (31)-(32), we get that

$$q(G)(x_{2} + x_{k+1}) - q(G)(x_{k} + x_{k+2})$$

$$= (n-5)x_{2} + x_{k+1} - 5x_{k} - 4x_{k+2} - x_{k-1} + \sum_{3 \le i \le k-3} x_{i} + \sum_{k+4 \le i \le n} x_{i}$$

$$= (n-15)x_{2} + x_{k+1} + 5x_{2} - 5x_{k} + 4x_{2} - 4x_{k+2} + x_{2} - x_{k-1} + \sum_{3 \le i \le k-3} x_{i} + \sum_{k+4 \le i \le n} x_{i}.$$
(33)

Because $n \ge 15$, (33) implies that $x_2 + x_{k+1} > x_k + x_{k+2}$.

Let $\mathbb{F} = G - v_k v_{k+2} + v_2 v_{k+1}$. Note that $X^T Q(\mathbb{F}) X - X^T Q(G) X = (x_2 + x_{k+1})^2 - (x_k + x_{k+2})^2$. It follows that when $n \ge 15$, then $q(\mathbb{F}) > X^T Q(\mathbb{F}) X > X^T Q(G) X = q(G)$. Because $n \ge 15$, by (i), it follows immediately that $q(\mathcal{H}) > q(\mathbb{F}) > q(G)$.

Case 3 $G \cong D_4$.

For convenience, we suppose that $G = D_4$. Because $d_G(v_2) = \Delta'(G) = n - 3$, $4 \le k \le l - 2$, $l \le n - 1$. Let $X = (x_1, x_2, ..., x_n)^T \in \mathbb{R}^n$ be the Perron eigenvector corresponding to q(G), where x_i corresponds to vertex v_i . Let $\mathbb{F} = G - v_{l-1}v_{l+1} + v_2v_l$. As (i), it can be proved that $q(G) < q(\mathbb{F})$. By (i), we get that $q(\mathbb{F}) < q(\mathcal{H})$. Then $q(G) < q(\mathcal{H})$.

From above three cases, (ii) follows. This completes the proof. \Box

Theorem 3.13 Let G be a maximal planar graph with order $n \ge 456$, $d(v_1) = \Delta(G) = n - 1$ and $\Delta'(G) \le n - 2$. Then $q(G) < q(\mathcal{H})$.

Proof. This theorem follows from Lemmas 3.2 and 3.9-3.12. \Box

Theorem 3.14 Let G be a maximal planar graph with order $n \ge 456$. Then $q(G) \le q(\mathcal{H})$ with equality if and only if $G \cong \mathcal{H}$.

Proof. This theorem follows from Lemmas 3.1, 3.2, Theorems 3.8 and 3.13. \Box

Theorem 3.15 Let G be a planar graph with order $n \ge 456$. Then $q(G) \le q(\mathcal{H})$ with equality if and only if $G \cong \mathcal{H}$.

Proof. This theorem follows from the narrations in Section 2 and Theorem 3.14. \Box

Remark By computations with computer, we can check that among all planar graphs of order $n \leq 6$, the graph \mathcal{H} has the maximal signless Laplacian spectral radius. And by computation with computer, we can check and find which graph has the maximal signless Laplacian spectral radius among all planar graphs with not large order. But it seems obviously it is not a good way by computation to check and find which graph has the maximal signless Laplacian spectral radius among all planar graphs with large order because magnitudes of work need do. As for the planar graphs of order $n \leq 455$, we think that it need a smart way to check and find which graph has the maximal signless Laplacian spectral radius. By some computations with computer, we conjuncture that among all planar graphs of order $n \leq 455$, the graph \mathcal{H} still has the maximal signless Laplacian spectral radius.

References

- [1] J. Bondy, U. Murty, Graph theory, Springer Press, 2008.
- [2] B. Boots, G. Royle, A conjecture on the maximum value of the principal eigenvalue of a planar graph, Geographical Anal. 23 (1991) 276-282.
- [3] D. Cao, A. Vince, The spectral radius of a planar graph, Linear Algebra Appl. 187 (1993) 251-257.
- [4] D. Cvetkovic, M. Doob, H. Sachs, Spectra of Graphs: Theory and Application, Academic Press, New York, 1979.
- [5] D. Cvetković, P. Rowlinson, The largest eigenvalue of a graph: A survey, Linear Multilinear Algebra 28 (1990) 3-33.
- [6] D. Cvetković, P. Rowlinson, S. Simić, Signless Laplacians of finite graphs, Linear Algebra Appl. 423 (2007) 155-171.

- [7] D. Cvetković, S. Simić, Towards a spectral theory of graphs based on the signless Laplacian,
 I, Publications De l'institut Mathématique Nouvelle série, tome 85 (99) (2009) 19-33.
- [8] D. Cvetković, S. Simić, Towards a spectral theory of graphs based on the signless Laplacian, II, Linear Algebra Appl. 432 (2010) 2257-2272.
- [9] D. Cvetković, S. Simić, Towards a spectral theory of graphs based on the signless Laplacian, III, Discrete Math. 4 (2010) 156-166.
- [10] D. Cvetković, S. Simić, Graph spectra in Computer Science, Linear Algebra Appl. 434 (2011) 1545-1562.
- [11] E. van Dam, W. Haemers, Which graphs are determined by their spectrum, Linear Algebra Appl. 373 (2003) 241-272.
- [12] Z. Dvořák, B. Mohar, Spectral radius of finite and infinite planar graphs and of graphs of bounded genus, J. Combin. Theory Ser. B 100 (2010) 729-739.
- [13] L. Feng, G. Yu, A. Ilić, D. Stevanović, The signless Laplacian spectral radius of graphs on surfaces, Linear Multilinear Algebra 61 (2013) 573-581.
- [14] F. Harary, Graph theory, Addison-Wesley, New York, 1969.
- [15] Y. Hong, On the spectral radius and the genus of graphs, J. Combin. Theory Ser. B 65 (1995), 262-268.
- [16] Y. Hong, Sharp upper bounds of the spectral radius of graphs in terms of genus, J. Combin. Theory Ser. B 74 (1998), 153-159.
- [17] R. Merris, A note on Laplacian graph eigenvalues, Linear Algebra Appl. 285 (1998) 33-35.
- [18] H. Minc, Nonnegative Matrices, John Wiley & Sons Inc., New York, 1988
- [19] B. Mohar, W. Woess, A survey on spectra of infinite graphs, Bull. Lond. Math. Soc. 21 (1989) 209-234.
- [20] A. Schwenk, R. Wilson, On the eigenvalues of a graph, in Selected Topics in Graph Theory, L.W. Beineke and R.J. Wilson, eds., Academic Press, New York, 1978, pp. 307-336.
- [21] J. Shu, Y. Hong, Upper bounds of the spectral radius for outer-planar graphs and Halin graphs, Chin. Ann. Math. Ser. A 21 (2001) 677-682.
- [22] Y. Wang, D. Chakrabarti, C. Wang, C. Faloutsos, Epidemic spreading in real networks: an eigenvalue viewpoint, in: 22nd Symposium in Reliable Distributed Computing, Florence Italy, 6-8 October, 2003.
- [23] G. Yu, J. Shu, Y. Hong, Bounds of the spectral radii of $K_{2,3}$ -minor free graphs, Electron. J. Linear Algebra, 23 (2012), 171-179.
- [24] L. Zhang, H. Jiang, Impulsive cluster anticonsensus of discretemultiagent linear dynamic systems, Discrete Dyn. Nat. Soc., 2012 (2012), Article ID857561, 11 pages.
- [25] L. Zhang, H. Jiang, S. Guo, ICluster Anticonsensus of Multiagent Systems Based on the Q-Theory, Discrete Dyn. Nat. Soc., 2014 (2014), Article ID 254749, 7 pages.