

BURCHNALL-CHAUNDY POLYNOMIALS AND THE LAURENT PHENOMENON

A.P. VESELOV AND R. WILLOX

ABSTRACT. The Burchnell-Chaundy polynomials $P_n(z)$ are determined by the differential recurrence relation

$$P'_{n+1}(z)P_{n-1}(z) - P_{n+1}(z)P'_{n-1}(z) = P_n(z)^2$$

with $P_{-1} = P_0(z) = 1$. The fact that this recurrence relation has all solutions polynomial is not obvious and is similar to the integrality of Somos sequences and the Laurent phenomenon. We discuss this parallel in more detail and extend it to two difference equations

$$Q_{n+1}(z+1)Q_{n-1}(z) - Q_{n+1}(z)Q_{n-1}(z+1) = Q_n(z)Q_n(z+1)$$

and

$$R_{n+1}(z+1)R_{n-1}(z-1) - R_{n+1}(z-1)R_{n-1}(z+1) = R_n^2(z)$$

related to two different KdV-type reductions of the Hirota-Miwa and Dodgson octahedral equations. As a corollary we have a new form of the Burchnell-Chaundy polynomials in terms of the initial data $P_n(0)$, which is shown to be Laurent.

1. INTRODUCTION

In the 1920s Burchnell and Chaundy [5] discovered a remarkable sequence of polynomials satisfying the recurrence relation

$$(1) \quad P'_{n+1}(z)P_{n-1}(z) - P_{n+1}(z)P'_{n-1}(z) = P_n(z)^2$$

with $P_{-1} = P_0(z) = 1$, where $'$ means differentiation in z :

$$P_1 = z, \quad P_2 = \frac{1}{3}(z^3 + \tau_2), \quad P_3 = \frac{1}{45}(z^5 + 5\tau_2z^3 + \tau_3z - 5\tau_2^2),$$

$$P_4 = \frac{1}{4725}(z^{10} + 15\tau_2z^7 + 7\tau_3z^5 - 35\tau_2\tau_3z^2 + 175\tau_2^3z - \frac{7}{3}\tau_3^2 + \tau_4z^3 + \tau_4\tau_2), \dots$$

Note that at each step we have an additional integration constant because of the freedom in the solution of the differential equation

$$P_{n+1}(z) \rightarrow P_{n+1}(z) + cP_{n-1}(z)$$

(we are using here Adler and Moser's notation with $\tau_1 = 0$, see [2]).

These polynomials have been rediscovered several times, most notably by Stelmacher and Lagnese in the theory of Huygens' principle [19] and by Adler and Moser in the theory of rational solutions of the Korteweg-de Vries equation [2]. They appeared also in the bispectral theory of Duistermaat and Grünbaum [8], who explained their important role in the theory of monodromy-free Schrödinger operators and the relation to Schur polynomials with triangular Young diagrams. The Burchnell-Chaundy polynomials $P_n(z)$ are special cases of the Schur-Weierstrass polynomials introduced by Buchstaber, Enolskii and Leykin in the theory of Klein sigma-functions [3] (see also Nakayashiki [16] for the relation to KP tau functions and Sato theory).

Note that the very existence of these polynomials looks like a miracle since the relation (1) can be rewritten as

$$\frac{d}{dz} \frac{P_{n+1}}{P_{n-1}} = \frac{P_n^2}{P_{n-1}^2},$$

which means that all the residues of the right-hand side must be zero (see the discussion of this in [5] and [2]).

We would like to make the parallel with the sequence

$$p_{n+1}p_{n-1} = p_n^2 + 1, \quad p_{-1} = p_0 = 1,$$

which surprisingly is integer for all n : 1, 2, 5, 13, 34, 89, ... (these in fact are nothing else but every other Fibonacci number). This sequence is related to the cluster algebra of type $A_1^{(1)}$ and gives a nice example of the so-called Laurent phenomenon studied by Fomin and Zelevinsky: for general initial data p_{-1} and p_0 , the solution p_n is a Laurent polynomial in p_{-1} and p_0 with integer coefficients (see [6, 9]). In particular, if $p_{-1} = p_0 = 1$ this implies the integrality of the p_n (which, of course, can be proven in an elementary way as well). We see that the Burchnell-Chaundy sequence is a functional analogue of the same phenomenon with the role of integers \mathbb{Z} played by the polynomial ring $\mathbb{Q}[z]$.

This work started as an attempt to see if this conceptual parallel with the Laurent phenomenon can be made a real connection. This led us naturally to the following *difference Burchnell-Chaundy equation*

$$(2) \quad Q_{n+1}(z+1)Q_{n-1}(z) - Q_{n+1}(z)Q_{n-1}(z+1) = Q_n(z)Q_n(z+1),$$

with $Q_{-1}(z) = Q_0(z) = 1$.

We prove that this equation also has polynomial solutions $Q_n(z)$ with coefficients that are Laurent polynomials of the initial data $q_k = Q_k(0)$, such that

$$A_n Q_n(z) \in \mathbb{Z}[z; q_1^{\pm 1}, \dots, q_{n-2}^{\pm 1}, q_{n-1}, q_n], \quad A_n = \prod_{j=1}^n (2j-1)!!,$$

where $(2k+1)!! = 1 \times 3 \times 5 \times \dots \times (2k+1)$. The first three polynomials have the form

$$\begin{aligned} Q_1 &= z + q_1, & Q_2 &= \frac{z(z^2 - 1)}{3} + q_1 z^2 + q_1^2 z + q_2, \\ Q_3 &= \frac{z^2(z^2 - 1)(z^2 - 4)}{45} + \frac{2q_1 z^5}{15} + \frac{q_1^2 z^4}{3} + \frac{(q_1^3 - q_1 + q_2)z^3}{3} + \frac{(3q_1 q_2 - q_1^2)z^2}{3} \\ &\quad + \left(\frac{q_3}{q_1} + \frac{q_2^2}{q_1} + \frac{2q_2}{3} - \frac{q_1^3}{3} + \frac{q_1}{5} \right) z + q_3. \end{aligned}$$

If we set $z = m \in \mathbb{Z}$, the difference Burchnell-Chaundy equation simply becomes a certain KdV-type reduction of the Hirota-Miwa equation, which is known to have the Laurent property [9]. This explains the Laurent property of the coefficients, but the proof of polynomiality of $Q_n(z)$ needs additional arguments and is related to specific properties of the Cauchy problem we consider. We follow here the classical approach [5], [2] by using an explicit description of solutions in terms of Casorati determinants, which are standard in the theory of integrable systems. As a result we obtain an independent proof of the Laurent property for the coefficients and a recursive way for their calculation.

We show also that these polynomials, after rescaling $R_n(z) = 2^{-n(n+1)/2} Q_n(z)$, satisfy the difference equation

$$(3) \quad R_{n+1}(z+1)R_{n-1}(z-1) - R_{n+1}(z-1)R_{n-1}(z+1) = R_n^2(z)$$

with the same initial data $R_{-1}(z) = R_0(z) = 1$. This equation, which we call *difference Dodgson equation*, for $z = m \in \mathbb{Z}$ is a reduction of the 3D Dodgson

octahedral equation, which is formally equivalent to the Hirota-Miwa equation but has different geometry.

We show that under a natural continuum limit the polynomials $Q_n(z)$ become the usual Burchnall-Chaundy polynomials $P_n(z)$, thus giving one more proof for their existence.

A corollary of our results is the proof that in terms of the initial data $c_n = P_n(0)$,

$$A_n P_n \in \mathbb{Z}[z; c_1^{\pm 1}, \dots, c_{n-2}^{\pm 1}, c_{n-1}, c_n], \quad A_n = \prod_{j=1}^n (2j-1)!!,$$

have coefficients which are Laurent polynomials in c_k with integer coefficients:

$$\begin{aligned} P_1 &= z + c_1, \quad P_2 = \frac{1}{3}(z^3 + 3c_1 z^2 + 3c_1^2 z + 3c_2), \\ P_3 &= \frac{1}{45}(z^6 + 6c_1 z^5 + 15c_1^2 z^4 + 15(c_1^3 + c_2)z^3 + 45c_1 c_2 z^2 + 45(\frac{c_2^2}{c_1} + \frac{c_3}{c_1})z + 45c_3), \\ P_4 &= \frac{1}{4725}(z^{10} + 10c_1 z^9 + 45c_1^2 z^8 + 15(7c_1^3 + 3c_2)z^7 + 105(c_1^4 + 3c_2 c_1)z^6 \\ &+ 315(\frac{c_2^2}{c_1} + \frac{c_3}{c_1} + 2c_1^2 c_2)z^5 + 1575(c_2^2 + c_3)z^4 + 1575(\frac{c_3^2}{c_1^2} + c_1 c_3 + \frac{c_4}{c_2} + \frac{c_3^2}{c_1^2 c_2} + 2\frac{c_3 c_2}{c_1^2})z^3 \\ &+ 4725(\frac{c_3^2}{c_1 c_2} + \frac{c_2 c_3}{c_1} + \frac{c_1 c_4}{c_2})z^2 + 4725(\frac{c_3^2}{c_2} + \frac{c_1^2 c_4}{c_2})z + 4725c_4), \dots \end{aligned}$$

This form of the Burchnall-Chaundy polynomials seems to be new and demonstrates one more feature of these remarkable polynomials.

2. DODGSON, HIROTA-MIWA AND BURCHNALL-CHAUNDY

In 1866 Charles Dodgson (known to the world under the name of Lewis Carroll) proposed a very original method of computing determinants called ‘‘condensation method’’ [7]. One can view this method as the solution of a very special Cauchy problem for the discrete *Dodgson octahedral equation*

$$(4) \quad u_{l,m+1,n+1} u_{l,m-1,n-1} - u_{l,m+1,n-1} u_{l,m-1,n+1} = u_{l-1,m,n} u_{l+1,m,n},$$

where $m, n, l \in \mathbb{Z}$, $m \equiv n \equiv l \pmod{2}$. Let A be an $N \times N$ matrix for which the determinant is to be computed. Assume that N is odd and consider the following Cauchy data: $u_{-1,m,n} = 1$, for all $m, n \in \mathbb{Z}$, and

$$u_{0,-N-1+2i,-N-1+2j} = A_{ij},$$

when $i, j = 1, \dots, N$ and $u_{0,-N-1+2i,-N-1+2j} = 0$ otherwise. For even N we should consider the same initial data for $l = 0$ and $l = 1$ respectively. The Dodgson condensation can be viewed as the evolution, according to (4), along the positive l -direction inside the characteristic cone: $\max(|m|, |n|) \leq N - l - (N \bmod 2)$. The support of such a solution for a generic matrix will therefore be a pyramid with matrix A in the base and the determinant $\det A$ at the top. Outside this pyramid we can assume all $u_{l,m,n}$ for $l > 0$ to be zero such that the equation is trivially satisfied.

In the 1980s Hirota [11] discovered a discrete version of the KP equation on a standard cubic lattice, which was subsequently studied by Miwa [15] in relation to Sato theory, and which is now known as the *Hirota-Miwa equation*

$$(5) \quad a v_{l+1,m,n} v_{l,m+1,n+1} + b v_{l,m+1,n} v_{l+1,m,n+1} + c v_{l,m,n+1} v_{l+1,m+1,n} = 0,$$

where $l, m, n \in \mathbb{Z}$ and a, b, c are arbitrary non-zero parameters. Note that the exact values of the parameters are not very important and can be changed by the gauge




FIGURE 1. Dodgson's Cauchy pyramid for computing 3×3 determinants

transformation

$$v_{l,m,n} = \left(\frac{\tilde{a}}{a}\right)^{mn} \left(\frac{\tilde{b}}{b}\right)^{ln} \left(\frac{\tilde{c}}{c}\right)^{lm} \tilde{v}_{l,m,n},$$

which however will affect the Cauchy data.

Formally the Hirota-Miwa equation may be considered as a version of the Dodgson equation if one interprets these six vertices of the cube as the vertices of the octahedron, see Figure 2.




FIGURE 2. Support of the Hirota-Miwa equation in the cube

It is however clear that from a geometric point of view this is unnatural. Indeed, the typical initial value problems for the Hirota-Miwa equation are very different from the Dodgson initial value problem. In particular, typical explicit solutions for the Hirota-Miwa equation are given by determinants of the same size, while for Dodgson it is crucial that the size of the matrices is shrinking when moving up in the pyramid.

This difference is clear if we compare two natural reductions of these equations. For the octahedral equation we consider the reduction $u_{l+1,m,n} = u_{l-1,m,n}$ leading to the discrete Dodgson equation

$$(6) \quad u_{m+1,n+1}u_{m-1,n-1} - u_{m+1,n-1}u_{m-1,n+1} = u_{m,n}^2,$$

where $m \equiv n \pmod{2}$. If we extend this equation to all integers m, n and replace m by z we come to the difference Dodgson equation (3).

For Hirota-Miwa a natural reduction is $v_{l+1,m,n+1} = v_{l,m,n}$ which, for $a = 1$, $b = c = -1$, leads to the *discrete KdV equation* in Hirota bilinear form [10]

$$(7) \quad Q_{m+1,n+1}Q_{m,n-1} - Q_{m,n+1}Q_{m+1,n-1} = Q_{m,n}Q_{m+1,n},$$

the functional version of which is the Burchnell-Chaundy equation (2). Note that the support of the equation has a domino shape, which is a cube squashed along one of the face diagonals, see Figure 3.




FIGURE 3. Domino-type support for the discrete KdV equation.

Because of their common origin it is natural to expect that the equations (2) and (3) are closely related, which is indeed the case.

Theorem 1. *The difference Burchnell-Chaundy (2) and Dodgson (3) equations are equivalent on the set of initial data satisfying $\Phi_0 = 0$, where*

$$(8) \quad \Phi_n(z) := Q_n(z+1)Q_{n-1}(z-1) + Q_{n-1}(z+1)Q_n(z-1) - 2Q_{n-1}(z)Q_n(z).$$

More precisely, if the initial data $Q_{-1}(z), Q_0(z)$ of the Cauchy problem for the dBCh equation (2) satisfy the constraint

$$(9) \quad Q_0(z+1)Q_{-1}(z-1) + Q_{-1}(z+1)Q_0(z-1) - 2Q_{-1}(z)Q_0(z) = 0,$$

then $R_n(z) = 2^{-\frac{n(n+1)}{2}} Q_n(z)$ satisfy the Dodgson relation (3).

Proof. If we modify the Dodgson equation as

$$(10) \quad R_{n+1}(z+1)R_{n-1}(z-1) - R_{n+1}(z-1)R_{n-1}(z+1) = 2R_n^2(z),$$

then we simply have to show that under the constraint $\Phi_0 = 0$ we have $Q_n(z) = R_n(z)$. Note that the constraint is preserved by the Burchnell-Chaundy dynamics. Indeed, assume that

$$(11) \quad \Phi_n(z) = Q_n(z+1)Q_{n-1}(z-1) + Q_{n-1}(z+1)Q_n(z-1) - 2Q_{n-1}(z)Q_n(z) = 0$$

and we have two dBCh relations

$$(12) \quad Q_{n+1}(z)Q_{n-1}(z-1) - Q_{n+1}(z-1)Q_{n-1}(z) = Q_n(z-1)Q_n(z),$$

$$(13) \quad Q_{n+1}(z+1)Q_{n-1}(z) - Q_{n+1}(z)Q_{n-1}(z+1) = Q_n(z)Q_n(z+1).$$

Multiplying (12) by $Q_n(z+1)$ and (13) by $Q_n(z-1)$ we have

$$\begin{aligned} Q_{n+1}(z+1)Q_n(z-1)Q_{n-1}(z) - Q_{n+1}(z)Q_n(z-1)Q_{n-1}(z+1) \\ = Q_n(z-1)Q_n(z)Q_n(z+1), \\ Q_{n+1}(z)Q_n(z+1)Q_{n-1}(z-1) - Q_{n+1}(z-1)Q_n(z+1)Q_{n-1}(z) \\ = Q_n(z-1)Q_n(z)Q_n(z+1), \end{aligned}$$

so

$$(14) \quad \begin{aligned} & Q_{n+1}(z+1)Q_n(z-1)Q_{n-1}(z) - Q_{n+1}(z)Q_n(z-1)Q_{n-1}(z+1) \\ & = Q_{n+1}(z)Q_n(z+1)Q_{n-1}(z-1) - Q_{n+1}(z-1)Q_n(z+1)Q_{n-1}(z). \end{aligned}$$

On the other hand $Q_{n-1}(z)\Phi_{n+1}(z) - Q_{n+1}(z)\Phi_n(z)$ can be rewritten as

$$\begin{aligned} & Q_{n-1}(z)Q_{n+1}(z+1)Q_n(z-1) + Q_{n-1}(z)Q_n(z+1)Q_{n+1}(z-1) \\ & \quad - Q_{n+1}(z)Q_n(z+1)Q_{n-1}(z-1) - Q_{n+1}(z)Q_{n-1}(z+1)Q_n(z-1), \end{aligned}$$

which is zero due to (14).

Now in order to prove that the $R_n(z) = Q_n(z)$ satisfy (10), we multiply relation (11) by $Q_n(z)$, (12) by $Q_{n-1}(z+1)$, (13) by $Q_{n-1}(z-1)$ and add all of them to obtain

$$Q_{n-1}(z)Q_{n+1}(z+1)Q_{n-1}(z-1) - Q_{n-1}(z)Q_{n+1}(z-1)Q_{n-1}(z+1) = 2Q_{n-1}(z)Q_n^2(z).$$

Dividing now by $Q_{n-1}(z)$ proves the claim. \square

This theorem can be reformulated as follows. Consider a two by two square formed by two adjacent KdV dominos (see Figure 3). It can be alternatively described as two horizontal dominos with relations $\Phi_n = 0$ and $\Phi_{n+1} = 0$ on them. Then the claim is that any three of these domino relations implies the fourth one and the Dodgson relation (10).

3. CAUCHY PROBLEM FOR DISCRETE EQUATIONS AND LAURENT PROPERTY

Fomin and Zelevinsky [9] have shown that the Hirota-Miwa and the octahedral Dodgson equations have the Laurent property for suitable initial data. They also considered some reductions, including the discrete KdV equation

$$(15) \quad \alpha Q_{m+1,n+1}Q_{m,n-1} + \beta Q_{m,n+1}Q_{m+1,n-1} = Q_{m,n}Q_{m+1,n}$$

(which they called "the knight recurrence" and attributed to Noam Elkies).

In the case of a particular Cauchy problem for this equation with $\alpha = 1, \beta = -1$ and

$$Q_{m,-1} = Q_{m,0} = 1, \quad Q_{0,n} = q_n, \quad m, n \in \mathbb{Z},$$

this implies that the general solution $Q_{m,n}$ is a Laurent polynomial in q_i with integer coefficients. In particular, this implies that when all $q_i = 1$, all the $Q_{m,n}$ are integers as can be seen in Figure 4. Note that on the line $m = 1$ in this figure we have the Fibonacci numbers $Q_{1,n} = F_n$,

$$F_{n+1} = F_n + F_{n-1}, \quad F_{-1} = F_0 = 1,$$

which grow like φ^n , where $\varphi = \frac{1+\sqrt{5}}{2}$ is the golden ratio. On the next line we have the sequence $Q_{2,n} = G_n$ satisfying the linear recurrence,

$$F_{n-1}G_{n+1} = F_{n+1}G_{n-1} + F_nG_n, \quad G_{-1} = G_0 = 1,$$

with Fibonacci numbers as the coefficients, which grows like φ^{2n} . Similarly, $|Q_{m,n}|$ for fixed m grows exponentially like φ^{mn} as $n \rightarrow \infty$.

In the m direction the growth is much slower: for fixed n the $|Q_{m,n}|$ grow like $m^{\frac{n(n+1)}{2}}$. This indicates that the dependence on m is polynomial, but to prove this for arbitrary initial data q_n seems to be not easy.

Instead we prefer to use (like Burchnell and Chaundy [5] and Adler and Moser [2]) alternative tools, standard in the theory of integrable systems, by presenting explicit determinantal expressions for the solutions. The Wronskians from [2, 5] are naturally replaced by Casoratians in our case, which is usual in the discrete setting, see e.g. [17]. Note however that in soliton KP theory, the matrices that yield determinant solutions usually have the same size, which for example in the case of

12181	-507	-455	-91	21	5	1	21	397	6469	104145	1332565	15181325
377	13	13	21	9	-3	1	13	149	1629	14001	115245	908245
615	-26	-23	-4	3	2	1	8	59	350	2109	11492	52375
249	51	5	1	1	-1	1	5	21	91	329	977	2477
-39	-19	-7	-1	1	1	1	3	9	21	41	71	113
-5	-4	-3	-2	-1	0	1	2	3	4	5	6	7
1	1	1	1	1	1	1	1	1	1	1	1	1
1	1	1	1	1	1	1	1	1	1	1	1	1
7	6	5	4	3	2	1	0	-1	-2	-3	-4	-5
113	71	41	21	9	3	1	1	1	-1	-7	-19	-39
2477	977	329	91	21	5	1	-1	1	1	5	51	249
52375	11492	2109	350	59	8	1	2	3	-4	-23	-26	615
908245	115245	14001	1629	149	13	1	-3	9	21	13	13	377
15181325	1332565	104145	6469	397	21	1	5	21	-91	-455	-507	12181

FIGURE 4. Solution to the discrete KdV equation with Cauchy data $Q_{0,n} = Q_{m,-1} = Q_{m,0} = 1$. The axes are given by the column of 1's (n axis) and the top row of 1's (m axis).

the discrete KdV equation (15) would require the sum of the coefficients α and β to be equal to 1. This is not true in our case (7), for which it is known [2, 5] that it is essential to have matrices of growing size.

4. CASORATI DETERMINANTAL FORMULAE

Let $Q_n(z)$ be a solution of the Cauchy problem for the difference Burchnall-Chaundy equation (2) with Cauchy data $Q_{-1}(z) = Q_0(z) = 1$ and arbitrary values for $Q_n(0)$. Since the Cauchy data $Q_{-1}(z) = Q_0(z) = 1$ satisfy the constraint (9) the corresponding polynomials $R_n(z)$ that satisfy (3) are simply

$$R_n(z) = 2^{-\frac{n(n+1)}{2}} Q_n(z).$$

The following procedure is a natural difference analogue of the Adler and Moser description of Burchnall-Chaundy polynomials [2]. Define first the sequence of polynomials $x_n(z)$ by

$$(16) \quad \Delta x_n(z) = x_{n-1}(z), \quad x_0 = 1,$$

where

$$\Delta f(z) := f(z+1) - f(z).$$

They depend on some additional parameters, which are not unique. The generating function

$$F(z, u) := \sum_{k=0}^{\infty} x_k(z) u^k,$$

satisfies the relation

$$\Delta F = uF$$

and has the general form

$$F(z, u) = A(t)(1+u)^z = A(t)e^{\sum_{k=1}^{\infty} (-1)^{k+1} \frac{u^k}{k} z},$$

where $A(t)$ is an arbitrary function of the parameters. We fix the choice of these parameters, which will turn out to be related to the KdV flows, by choosing

$$A(t) = e^{\sum_{k=1}^{\infty} (-1)^{k+1} t_k \frac{u^k}{k}},$$

so that

$$(17) \quad F(z, t, u) = e^{\sum_{k=1}^{\infty} (-1)^{k+1} (z+t_k) \frac{u^k}{k}},$$

leading to

$$x_0 = 1, \quad x_1 = z + t_1, \quad x_2 = \frac{1}{2}[(z + t_1)^2 - (z + t_2)], \quad \dots$$

These polynomials can be given also as the determinants (see e.g. [12], page 28)

$$(18) \quad x_k(z) = \frac{1}{k!} \begin{vmatrix} z_1 & -1 & 0 & \dots & \dots & 0 \\ z_2 & z_1 & -2 & \dots & \dots & 0 \\ z_3 & z_2 & z_1 & -3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & 0 \\ z_{k-1} & z_{k-2} & \dots & \dots & z_1 & 1-k \\ z_k & z_{k-1} & \dots & \dots & z_2 & z_1 \end{vmatrix}, \quad z_k = (-1)^{k+1} (z + t_k).$$

Let now $y_k = x_{2k-1}$ and consider the Casoratians $Q_n(z) = C(y_1, \dots, y_n)$, where the Casoratian $C(f_1, \dots, f_n)$ of the functions $f_1(z), \dots, f_n(z)$ is defined as

$$(19) \quad C(f_1, \dots, f_n) = \det \|f_i(z + j - 1)\|, \quad i, j = 1, \dots, n.$$

The Casoratians

$$(20) \quad Q_k = C(y_1, \dots, y_k)$$

can be written as the determinants

$$(21) \quad Q_k = \begin{vmatrix} x_1 & x_3 & x_5 & \dots & \dots & x_{2k-1} \\ 1 & x_2 & x_4 & \dots & \dots & x_{2k-2} \\ 0 & x_1 & x_3 & x_5 & \dots & x_{2k-3} \\ 0 & 1 & x_2 & x_4 & \dots & x_{2k-4} \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & \dots & x_{k-2} & x_k \end{vmatrix}$$

Theorem 2. *The Casoratians $Q_k(z)$ given by (21) with $x_j(z)$ defined by (18) satisfy the Burchall-Chaundy relations (2).*

Proof. We essentially follow the arguments of Adler and Moser [2].

Let ϕ_1, \dots, ϕ_k be arbitrary functions of z and consider the difference operator

$$C_k(\chi) = C(\phi_1, \dots, \phi_k, \chi).$$

Then we have what Adler and Moser call *Jacobi identity*:

$$[C_k(\chi), C_{k+1}] = -(TC_k) C_{k+1}(\chi),$$

where

$$C_k = C(\phi_1, \dots, \phi_k),$$

T is the shift operator:

$$Tf(z) = f(z + 1),$$

and by $[A, B] = C(A, B)$ we mean the Casoratian of A and B :

$$[A, B] = A(TB) - B(TA).$$

The proof is simple: both $[C_k(\chi), C_{k+1}]$ and $C_{k+1}(\chi)$ are difference operators of order $k + 1$ with the same kernel generated by $\phi_1, \dots, \phi_{k+1}$. Thus they can only differ by a factor, which is easily seen to be $-(TC_k)$.

In our case we have $\Delta^2 \phi_k = \phi_{k-1}$ and $\phi_1 = z$. It is then easy to see that this implies that if we take $\chi = 1$ then

$$C_k(1) = (-1)^k C(\Delta \phi_1, \dots, \Delta \phi_k).$$

However,

$$C(\Delta\phi_1, \dots, \Delta\phi_k) = C(\Delta^2\phi_2, \dots, \Delta^2\phi_k) = C(\phi_1, \dots, \phi_{k-1}) = C_{k-1},$$

and similarly: $C_{k+1}(1) = (-1)^{k+1}C_k$. Substituting this into the Jacobi identity we have

$$C_{k-1}(TC_{k+1}) - C_{k+1}(TC_{k-1}) = C_k(TC_k),$$

which means that $Q_n(z) = C_n$ satisfies (2). \square

Corollary 1. *The coefficients of the Burchnall-Chaundy polynomials $Q_k(z)$ are polynomial in the parameters t_j .*

Here are the first few polynomials expressed in KdV parameters:

$$\begin{aligned} Q_1 &= z + t_1, & Q_2 &= \frac{1}{3}(z(z^2 - 1) + 3t_1z^2 + 3t_1^2z + t_1^3 - t_3) \\ Q_3 &= \frac{1}{45}(z^2(z^2 - 1)(z^2 - 4) + 6t_1z^5 + 15t_1^2z^4 + (20t_1^3 - 5t_3 - 15t_1)z^3 + 15t_1(t_1^3 - t_1 - t_3)z^2 \\ &\quad + (9t_1 - 10t_3 + 9t_5 - 15t_1^2t_3 - 5t_1^3 + 6t_1^5)z + t_1^6 - 5t_3^2 - 5t_1^3t_3 + 9t_1t_5). \end{aligned}$$

Now we need to find the relation between the KdV parameters t_j and the Cauchy data $q_k = Q_k(0)$.

Substituting $z = 0$ in formula (18) we have the homogeneous polynomials

$$x_k = \frac{(-1)^{k+1}}{k}t_k + \phi_k(t_1, \dots, t_{k-1}),$$

where the weight of x_k and t_k is k :

$$\begin{aligned} x_1 &= t_1, & x_2 &= -\frac{1}{2}t_2 + \frac{1}{2}t_1^2, & x_3 &= \frac{1}{3}t_3 - \frac{1}{2}t_1t_2 + \frac{1}{6}t_1^3, \\ x_4 &= -\frac{1}{4}t_4 + \frac{1}{3}t_1t_3 + \frac{1}{8}t_2^2 - \frac{1}{4}t_1^2t_2 + \frac{1}{24}t_1^4, \\ x_5 &= \frac{1}{5}t_5 - \frac{1}{4}t_1t_4 - \frac{1}{6}t_2t_3 + \frac{1}{6}t_1^2t_3 + \frac{1}{8}t_1t_2^2 - \frac{1}{12}t_1^3t_2 + \frac{1}{120}t_1^5, \dots \end{aligned}$$

These relations can be inverted as the determinants (see [12], page 28)

$$(22) \quad t_k = \begin{vmatrix} x_1 & 1 & 0 & \dots & \dots & 0 \\ 2x_2 & x_1 & 1 & \dots & \dots & 0 \\ 3x_3 & x_2 & x_1 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & 0 \\ \cdot & \cdot & \dots & \dots & x_1 & 1 \\ kx_k & x_{k-1} & \dots & \dots & x_2 & x_1 \end{vmatrix}.$$

Note that $t_k \in \mathbb{Z}[x_1, \dots, x_k]$ are polynomials with integer coefficients.

Substituting $z = 0$ to (20) we have the relations

$$\begin{aligned} q_1 &= t_1, & q_2 &= -\frac{1}{3}t_3 + \frac{1}{3}t_1^3, & q_3 &= \frac{1}{5}t_1t_5 - \frac{1}{9}t_3^2 - \frac{1}{9}t_1^3t_3 + \frac{1}{45}t_1^6, \\ q_4 &= \frac{1}{21}(t_3 - t_1^3)t_7 - \frac{1}{25}t_5^2 + \frac{1}{15}t_1^2t_3t_5 + \frac{1}{75}t_1^5t_5 - \frac{1}{27}t_1t_3^3 - \frac{1}{315}t_1^7t_3 + \frac{1}{4725}t_1^{10}, \dots \end{aligned}$$

We can also define $q_{-1} = q_0 = 1$.

Note that the initial conditions q_k are homogeneous in t_j of weight $k(k+1)/2$.

Theorem 3. *The polynomial q_k depends only on odd parameters t_{2i-1} , ($i = 1, \dots, k$) and has the form*

$$q_k = \frac{(-1)^{k+1}}{2k-1}q_{k-2}t_{2k-1} + \psi_k(t_1, t_3, \dots, t_{2k-3}),$$

for some polynomials ψ_k with rational coefficients.

The parameter t_{2k-1} can be expressed in terms of q_j as a Laurent polynomial with integer coefficients

$$t_{2k-1} = (2k-1) \frac{(-1)^{k+1} q_k}{q_{k-2}} + \varphi_k(q_1, \dots, q_{k-1}) \in \mathbb{Z}[q_1^\pm, q_2^\pm, \dots, q_{k-2}^\pm, q_{k-1}, q_k].$$

Proof. The generating function (17) satisfies the differential relations

$$\frac{\partial}{\partial t_j} F = (-1)^{j+1} \frac{w^j}{j} F,$$

which imply that

$$\frac{\partial x_k}{\partial t_j} = \frac{(-1)^{j+1}}{j} x_{k-j}.$$

Differentiating the determinantal expression (21) with respect to t_2 we see that the derivative of the j -th column is precisely the preceding column and consequently, as the derivative of the first column is zero, that Q_k does not depend on t_2 . Similarly we see that the same is true for all even times (cf. [3]).

This means that we can choose the even times as we want. Let us choose them in such a way that the corresponding $x_{2p}(0) = 0$ for all $p \in \mathbb{Z}_+$. This then defines the t_{2p} as certain polynomials of odd times with rational coefficients:

$$t_2 = t_1^2, \quad t_4 = \frac{4}{3} t_1 t_3 + \frac{1}{2} t_2^2 - t_1^2 t_2 + \frac{1}{6} t_1^4 = \frac{4}{3} t_1 t_3 - \frac{1}{3} t_1^4, \quad \dots$$

From the determinant (21) we see that in this case the $q_k = Q_k(0)$ satisfy the recurrence relation

$$q_k = (-1)^{k+1} x_{2k-1} q_{k-2} + g_k(x_1, x_3, \dots, x_{2k-3}),$$

for some polynomial g_k with integer coefficients, which implies the first claim. This relation also allows us to express, recursively, x_{2k-1} (and hence t_{2k-1} using (22)) as a Laurent polynomial of q_1, \dots, q_k with integer coefficients. \square

The first four expressions for the t_{2k-1} have the form:

$$t_1 = q_1, \quad t_3 = -3q_2 + q_1^3, \quad t_5 = \frac{5q_3 - 5q_1^3 q_2 + 5q_2^2 + q_1^6}{q_1},$$

$$t_7 = -\frac{7q_1^2 q_4 + 14q_2^2 q_3 - q_1^9 q_2 + 7q_3^2 + 7q_2^4 - 14q_1^3 q_2^3 + 7q_1^6 q_2^2 - 7q_1^3 q_2 q_3}{q_1^2 q_2}.$$

Combining the expression (21) for Q_n with the previous theorem we have

Corollary 2. *Laurent phenomenon for the Burchnall-Chaundy sequence (2):*

$$A_n Q_n(z) \in \mathbb{Z}[z; q_1^\pm, \dots, q_{n-2}^\pm, q_{n-1}, q_n], \quad q_k = Q_k(0).$$

The first three polynomials are given explicitly in the Introduction.

The polynomials $Q_n^0(z)$ corresponding to zero Cauchy data $q_n = Q_n^0(0) = 0$ have to be dealt with as a special case because of the Laurent nature of the solution. Instead we can consider their expressions in terms of the t_j , which are polynomial, and therefore allow us to set $t_j = 0$. The corresponding polynomials admit the following explicit description.

Define the polynomials $Q_n^0(z)$ by the following recurrence relation

$$(23) \quad Q_n^0(z) = \frac{1}{(2n-1)!!} \prod_{j=1}^n (z+n+1-2j) Q_{n-1}^0(z),$$

or explicitly as

$$(24) \quad Q_n^0(z) = A_n^{-1} z^{\lfloor \frac{n+1}{2} \rfloor} \prod_{j=1}^{n-1} (z^2 - j^2)^{\lfloor \frac{n+1-i}{2} \rfloor}, \quad A_n = \prod_{j=1}^n (2j-1)!!,$$

where $(2k+1)!! = 1 \times 3 \times 5 \times \dots \times (2k+1)$ and $[x]$ denotes the integer part of x .

Theorem 4. *The polynomials $Q_n^0(z)$ satisfy the difference Burchnall-Chaundy relations with zero initial data $Q_n^0(0) = 0$.*

Proof is by a direct check.

The polynomials $Q_n^0(z)$ correspond to all $t_k = 0$ and thus have Casoratian form (20) with binomial

$$x_k = \frac{z(z-1)\dots(z-k+1)}{k!}.$$

It is interesting that they also can be given as symmetric Casoratians of simple monomials.

The *symmetric Casoratian* $C^*(f_1, \dots, f_n)$ of the functions $f_1(z), \dots, f_n(z)$ is defined as the determinant

$$(25) \quad C^*(f_1, \dots, f_n) = \det \|f_i(z+n+1-2j)\|, \quad i, j = 1, \dots, n.$$

Define the functions

$$(26) \quad f_j(x) = \frac{1}{(2j-1)!} z^{2j-1}, \quad j = 1, \dots, n.$$

Theorem 5. *The polynomials $Q_n^0(z)$ can be given as the symmetric Casoratians*

$$(27) \quad Q_n^0(z) = 2^{-n(n+1)/2} C^*(f_1, \dots, f_n)$$

of the functions (26).

The proof easily follows from the Vandermonde formula.

5. CONTINUUM LIMIT: USUAL BURCHNALL-CHAUNDY POLYNOMIALS

It is well-known [1, 2] that the Burchnall-Chaundy polynomials P_n are the τ -functions of the rational solutions of the Korteweg-de Vries equation

$$u_{T_1} = D^3 u - 6uDu, \quad D = \frac{d}{dx}$$

and its higher analogues

$$u_{T_k} = D^{2k+1} u + \dots$$

(see [14], where these equations are precisely defined for scaled u). Namely, in a proper parametrisation

$$u(x, T_1, \dots, T_n) = -2D^2 \log P_n(x, T_1, \dots, T_n)$$

are the general rational solutions of the KdV hierarchy [1, 2].

Adler and Moser realised that the parameters τ_k they used (see the formulae in the Introduction) are different from the KdV times T_k but are related to them by a non-trivial invertible polynomial transformation.

We claim that our parameters t_{2k+1} are simply related to the KdV times by the scaling

$$t_{2k+1} = 4^k (2k+1) T_k.$$

Theorem 6. *The continuum limit*

$$P_n(x, t_1, t_3, \dots, t_{2n-1}) = \lim_{\varepsilon \rightarrow 0} \varepsilon^{\frac{n(n+1)}{2}} Q_n\left(\frac{x}{\varepsilon}, \frac{t_3}{\varepsilon^3}, \dots, \frac{t_{2n-1}}{\varepsilon^{2n-1}}\right),$$

yields the usual Burchnall-Chaundy polynomials parametrized by the scaled KdV times t_3, \dots, t_{2n-1} .

Note that $\varepsilon^{\frac{n(n+1)}{2}} Q_n(\frac{x}{\varepsilon}, \frac{t_3}{\varepsilon^3}, \dots, \frac{t_{2n-1}}{\varepsilon^{2n-1}})$ is polynomial in ε because of the homogeneity property of the q 's. The proof then follows from Miwa's results on the relation between the Hirota-Miwa equation and KP/KdV hierarchy [15] and by comparison with the normalisation of the KdV flows in [14].

Since the initial data q_k are homogeneous in t_k they do not change during the continuum limit and we have $q_k = P_k(0) = c_k$.

Corollary 3. *Laurent phenomenon for the usual Burchnell-Chaundy relation (1):*

$$A_n P_n(z) \in \mathbb{Z}[z; c_1^\pm, \dots, c_{n-2}^\pm, c_{n-1}, c_n], \quad c_k = P_k(0).$$

The explicit form of the first 4 polynomials is given in the Introduction.

6. CONCLUDING REMARKS

The main question we are dealing with in this paper is basically what happens with the Laurent property when we replace a discrete equation by its functional difference version. Our point is that the answer depends very much on the type of the corresponding Cauchy problem.

It is known to the experts in integrable systems that the equation itself does not guarantee integrability, which holds only for particular initial value problems and functional class of the initial data. For example, for the KdV equation the initial value problem is integrable for decaying or periodic initial data [18], but for general initial data not much can be said. At the discrete level this fact is less visible and the importance of the choice of Cauchy problem is probably not well recognised. A discussion of the role the Cauchy problem plays with respect to the Laurent phenomenon for integrable equations can be found in [13].

To illustrate our point let us consider the same discrete KdV dynamics (7), but in the m direction, and its functional version by replacing n by x :

$$F_{m+1}(x+1)F_m(x-1) - F_{m+1}(x)F_m(x) - F_{m+1}(x-1)F_m(x+1) = 0$$

with $F_0(x) = 1$. It has the explicit solutions $F_m(x) = C_m \varphi^{mx}$, $\varphi = \frac{1+\sqrt{5}}{2}$, but these correspond to particular Cauchy data satisfying $F_m(1) = \varphi^m F_m(0)$. What can one say about the analytic structure of the solutions for general Cauchy data (in particular, for $F_m(1) = F_m(0) = 1$)? So, in other words, what are the analytic functions interpolating integers on the vertical lines in Figure 4?

It is clear that the case of the Burchnell-Chaundy polynomials is special, but the question is how special it is. It would be interesting therefore to study other reductions, in particular period 3 reductions which correspond to the Boussinesq equation and which should be related to Schur-Weierstrass polynomials for trigonal curves [3]. It is natural to link this with the theory of periodic Darboux/dressing chains [20, 21].

It would also be interesting to know if the polynomials Q_n play any special role in the theory of hyperelliptic sigma-functions [4].

7. ACKNOWLEDGEMENTS

A.P.V. is grateful to the Graduate School of Mathematical Sciences of the University of Tokyo for the support of his visit in April-July 2014, during which this work was done. R.W. would like to acknowledge support from the Japan Society for the Promotion of Science, through the JSPS grant: KAKENHI 24540204.

REFERENCES

- [1] H. Airault, H.P. McKean and J. Moser *Rational and elliptic solutions of the Korteweg-de Vries equation and a related many-body problem*. Comm. Pure Appl. Math. **30** (1977), 95–178.

- [2] M. Adler and J. Moser *On a class of polynomials connected with the Korteweg-de Vries equation*. Commun. Math. Phys., **61** (1978), 1–30.
- [3] V.M. Buchstaber, V.Z. Enolskii and D.V. Leykin *Rational analogs of abelian functions*. Funct. Anal. Appl., **33** (1999), 83–94.
- [4] V.M. Buchstaber, V.Z. Enolskii and D.V. Leykin *Kleinian functions, hyperelliptic Jacobians and applications*. Reviews in Math. and Math. Phys. **10** (1997), 1–125.
- [5] J.L. Burchnall and T.W. Chaundy *A set of differential equations which can be solved by polynomials*. Proc. London Math. Soc. **30** (1929-30), 401–414.
- [6] P. Caldero and A. Zelevinsky *Laurent expansions in cluster algebras via quiver representations*. Moscow Mathematical Journal **6**, no. 3 (2006), 411–429.
- [7] C.L. Dodgson *Condensation of determinants, being a new and brief method for computing their arithmetical values*. Proc. Royal Soc. London **15** (1886-67), 150–155.
- [8] J.J. Duistermaat and F.A. Grünbaum *Differential equations in the spectral parameter*. Comm. Math. Phys. **103**, no. 2 (1986), 177–240.
- [9] S. Fomin and A. Zelevinsky *The Laurent Phenomenon*. Adv. Appl. Math. **28** (2002), 119–144.
- [10] R. Hirota *Nonlinear partial difference equations. I. A difference analogue of the Korteweg-de Vries equation*. J. Phys. Soc. Japan **43** (1977), 1424–1433.
- [11] R. Hirota *Discrete analogue of a generalised Toda equation*. J. Phys. Soc. Japan **50** (1981), 3787–3791.
- [12] I.G. Macdonald *Symmetric functions and Hall polynomials*. 2nd edition, Oxford Univ. Press, 1995.
- [13] T. Mase *The Laurent phenomenon and discrete integrable systems*. RIMS Kokyuroku Bessatsu **B41** (2013), 43–64.
- [14] R.M. Miura, C.S. Gardner and M.D. Kruskal *Korteweg-de-Vries equation and generalizations V*. J. Math. Phys. **11** (3) (1970), 952–953.
- [15] T. Miwa *On Hirota’s difference equation*. Proc. Japan Acad. A **58** (1982), 9–12.
- [16] A. Nakayashiki *Sigma function as a tau function*. IMRN **2010** (2009), 383–394.
- [17] Y. Ohta, R. Hirota, S. Tsujimoto and T. Imai *Casorati and discrete Gram type determinant representations of solutions to the discrete KP hierarchy*. J. Phys. Soc. Japan **62** (1993), 1872–1886.
- [18] S.P. Novikov, S.V. Manakov, L.P. Pitaevskii and V.E. Zakharov. *Theory of solitons: The inverse scattering method*. Plenum, New York, 1984.
- [19] J.E. Lagnese and K.L. Stellmacher *A method of generating classes of Huygens’ operators*. J. Math. & Mech. **17** (1967), 461–472.
- [20] A.P. Veselov and A.B. Shabat *Dressing chains and the spectral theory of the Schrödinger operator* Funct. Anal. Appl. **27** (1993), 81–96.
- [21] R. Willox and J. Satsuma *Sato Theory and Transformations Groups. A Unified Approach to Integrable Systems in Discrete Integrable Systems*, B. Grammaticos, Y. Kosmann-Schwarzbach, Tamizhmani, Tamizharasi (Eds.), Springer-Verlag Berlin (2004), 17–55.

DEPARTMENT OF MATHEMATICAL SCIENCES, LOUGHBOROUGH UNIVERSITY, LOUGHBOROUGH LE11 3TU, UK AND MOSCOW STATE UNIVERSITY, MOSCOW 119899, RUSSIA
E-mail address: A.P.Veselov@lboro.ac.uk

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO 153-8914
E-mail address: willox@ms.u-tokyo.ac.jp