Fast Genome-Wide QTL Association Mapping on Pedigree and Population Data

Hua Zhou, 1John Blangero, 2Thomas D. Dyer,2Kei-hang K. Chan, 3 4Kenneth Lange,3 5 6Eric M. Sobel3

December 23, 2014

¹Department of Statistics, North Carolina State University, Raleigh, NC 27695-8203

- ²Department of Genetics, Texas Biomedical Research Institute, San Antonio, TX 78245-0549
- ³Department of Human Genetics, University of California, Los Angeles, CA 90095-1766
- ⁴Department of Epidemiology, University of California, Los Angeles, CA 90095-1766
- ⁵Department of Biomathematics, University of California, Los Angeles, CA 90095-1766

⁶Department of Statistics, University of California, Los Angeles, CA 90095-1766

Abstract

Since most analysis software for genome-wide association studies (GWAS) currently exploit only unrelated individuals, there is a need for efficient applications that can handle general pedigree data or mixtures of both population and pedigree data. Even data sets thought to consist of only unrelated individuals may include cryptic relationships that can lead to false positives if not discovered and controlled for. In addition, family designs possess compelling advantages. They are better equipped to detect rare variants, control for population stratification, and facilitate the study of parent-of-origin effects. Pedigrees selected for extreme trait values often segregate a single gene with strong effect. Finally, many pedigrees are available as an important legacy from the era of linkage analysis. Unfortunately, pedigree likelihoods are notoriously hard to compute. In this paper we re-examine the computational bottlenecks and implement ultra-fast pedigree-based GWAS analysis. Kinship coefficients can either be based on explicitly provided pedigrees or automatically estimated from dense markers. Our strategy (a) works for random sample data, pedigree data, or a mix of both; (b) entails no loss of power; (c) allows for any number of covariate adjustments, including correction for population stratification; (d) allows for testing SNPs under additive, dominant, and recessive models; and (e) accommodates both univariate and multivariate quantitative traits. On a typical personal computer (6 CPU cores at 2.67 GHz), analyzing a univariate HDL (high-density lipoprotein) trait from the San Antonio Family Heart Study (935,392 SNPs on 1357 individuals in 124 pedigrees) takes less than 2 minutes and 1.5 GB of memory. Complete multivariate QTL analysis of the three timepoints of the longitudinal HDL multivariate trait takes less than 5 minutes and 1.5 GB of memory. The algorithm is implemented as the Ped-GWAS Analysis (Option 29) in the MENDEL statistical genetics software package, which is freely available for Macintosh, Linux, and Windows platforms from http://genetics.ucla.edu/software/mendel.

Key words: genome-wide association study; pedigree; kinship; score test; fixed-effects models, multivariate traits

1 Introduction

Genome-wide association studies (GWAS) are now at a crossroads. After the discovery of thousands of genes influencing hundreds of common traits⁹, much of the low-hanging fruit has been plucked ^{13,36}. Because of the enormous sample sizes of current studies, new trait genes are still being uncovered. Unfortunately, most entail small effects. Is it possible that inheritance is predominantly polygenic, and a law of diminishing returns has set in? The push to exploit rare variants is one response to this dilemma. The previous generation of geneticists relied on linkage to map rare variants. Linkage mapping fell from grace because of its poor resolution. Reducing a genome search to a one or two megabase region leaves too large an expanse of DNA to sift. The real gold of linkage mapping may well be its legacy pedigrees²⁶. Pedigree data is particularly attractive in association studies because it permits control of population substructure and study of parent-of-origin effects. Related affecteds are also more likely to share the same disease predisposing gene than unrelated affecteds. Even in population-based association studies, taking into account estimated identity-by-descent (IBD) information is apt to reduce false positives and increases power. The recent availability of dense marker data from genotyping chips enables quick and accurate estimation of global and even local IBD⁷.

Geneticists turned to random sample and case-control data because of the relative ease of collecting population data and the computational challenges posed by pedigrees. The tide of computational complexity is now beginning to turn. To handle pedigree data in association testing, statistical geneticists have proposed semiparametric methods such as the generalized linear mixed model (GLMM)^{1,3} and generalized estimating equations (GEE)^{5,4}. Although such methods work for both quantitative and binary traits, they are compromised by current restrictions that reduce power. The GEE approach requires input of a working correlation structure for each pedigree. The kinship coefficient matrix is a natural candidate. However, current implementations require the same working correlation matrix across all clusters, which implicitly requires all pedigrees to have the same structure⁴. This is a dubious and restrictive assumption. In the limited context of case-control studies, recent methods such as M_{QLS} ³², ROADTRIPS³³, and FPCA⁴⁶ correct for pedigree and ethnically induced

	Mendel	FaST-LMM	GEMMA
Multi-threaded operation	Yes	Yes	No
Can estimate kinships via SNPs	Yes	Yes	Yes
Imports & exports kinship estimates	Yes	Yes	Yes
Allows retained co-variates	Yes	Yes	Yes
Allows linear constraints on co-variates	Yes	No	No
Can use either LRT or score test	Yes	No	Yes*
Allows multivariate analysis	Yes	No	Yes
Can perform multiple univariate analyses	Yes	No	No
Allows > 2 variance components	Yes	No	No
Analyzes X-linked loci	Yes	No	No
Automatic SNP filtering on MAF	Yes	No	Yes
Allows non-additive SNP models	Yes	No	No
Detects outlier pedigrees	Yes	No	No
Detects outlier individuals	Yes	No	No
Can simulate genotype/phenotype data	Yes	No	No
Reads in fractional genotype values	No	Yes	Yes

Table 1: Comparison of features in MENDEL, FAST-LMM, and GEMMA for GWAS of QTLs. *GEMMA can use the likelihood ratio, score, or Wald test.

correlations by exploiting dense marker data. Other authors attack the same issues more broadly from the GLMM perspective^{10,41,23}. Korte et al.¹² generalizes GLMM to multivariate traits. Models based on the transmission-disequilibrium test (TDT)³⁰ and its generalization, the family-based association test (FBAT)^{15,16,35,37,38}, are promising but ignore covariates and polygenic background. See Van Steen³⁴ for a recent overview of FBAT methods for GWAS. We treat all of these extensions in a unified framework consistent with exceptionally fast computing.

The present paper re-examines the computational bottlenecks encountered in association mapping with pedigree data. It turns out that the previous objections to pedigree GWAS can be overcome. Kinship coefficients can be based on explicitly provided pedigree structure or estimated from dense markers when genealogies are missing or dubious. Frequentist hypothesis testing usually operates by comparing maximum likelihoods under the null and alternative hypotheses. Maximization of the alternative likelihood must be conducted for each and every marker. Score tests constitute a more efficient strategy than likelihood ratio tests. This is the point of departure taken by Chen and Abecasis⁶, but they use approximations that we avoid. The GLOGS program³¹ makes similar approximations in the case-control setting. Here we consider arbitrary pedigrees and multivariate quantitative traits. Score tests require no additional iteration under the alternative model. All that is needed is evaluation of a quadratic form combining the score vector and the expected information matrix at the maximum likelihood estimates under the null model. Although it takes work to assemble these quantities, a careful analysis of the algorithm shows that fast testing is perfectly feasible.

In our implementation of score testing, the few SNPs with the most significant score-test p-values are automatically re-analyzed by the slightly more powerful, but much slower, likelihood ratio test (LRT). Our fixed effects (mean component) model assumes Gaussian variation of the trait; the two alleles of a SNP shift trait means. There is no confounding of association and linkage. This framework carries with it several advantages. First, it applies to random sample data, pedigree data, or a mix of both. Second, it enables covariate adjustment, including correction for population stratification. Third, it accommodates additive, dominant, and recessive SNP models. Fourth, it also accommodates both univariate and multivariate traits. And fifth, as just mentioned, it fosters both likelihood ratio tests and score tests. The mean component model is now implemented in our software package MENDEL for easy use by the genetics community. In addition, MENDEL provides a complete suite of tools for pedigree analysis, including GWAS data preparation and manipulation, pedigree genotype simulation (gene dropping), trait simulation, genotype imputation, local and global kinship coefficient estimation, and pedigree-based GWAS (ped-GWAS)^{21,20}.

The competing software packages EMMAX¹¹, MMM²⁷, FAST-LMM^{23,24}, and GEMMA^{44,45} already implement variance component models for quantitative trait locus (QTL) analysis. Exhaustive comparison of MENDEL to each of these programs is beyond the scope of the current paper. We limit our comparisons of MENDEL to the state-of-art packages FAST-LMM and GEMMA, arguably the fastest and most sophisticated of the competition. Table 1 summarizes some of the qualitative features of these packages. Our numerical examples also demonstrate an order of magnitude advantage in speed of MENDEL over FAST-LMM and GEMMA. This advantage stems from our careful formulation of the score test and our exploitation of the multicore processors resident in almost all personal computers and computational clusters.

3

2 Methods

2.1 QTL Association Mapping with Pedigrees

QTL association mapping typically invokes the multivariate Gaussian distribution to model the trait values $y = (y_i)$ over a pedigree. The observed trait value y_i of person *i* can be either univariate or multivariate. For simplicity we first assume y_i is univariate and later indicate the necessary changes for multivariate y_i . The standard model¹⁷ collects the corresponding trait means into a vector ν and the corresponding covariances into a matrix Ω and represents the loglikelihood of a pedigree as

$$L = -\frac{1}{2} \ln \det \mathbf{\Omega} - \frac{1}{2} (\mathbf{y} - \mathbf{\nu})^t \mathbf{\Omega}^{-1} (\mathbf{y} - \mathbf{\nu}), \qquad (1)$$

where det denotes the determinant function and the covariance matrix is typically parametrized as

$$\boldsymbol{\Omega} = 2\sigma_a^2 \boldsymbol{\Phi} + \sigma_d^2 \boldsymbol{\Delta}_7 + \sigma_h^2 \boldsymbol{H} + \sigma_e^2 \boldsymbol{I}.$$
(2)

Here the variance component Φ is the global kinship coefficient matrix capturing additive polygenic effects, and Δ_7 is a condensed identity coefficient matrix capturing dominance genetic effects. When pedigree structure is explicitly given, these genetic identity coefficients are easily calculated¹⁷. With unknown or dubious genealogies, the global kinship coefficient can be accurately estimated from dense markers⁷. The household effect matrix H has entries $h_{ij} = 1$ if individuals i and j belong to the same household and 0 otherwise. Individual environmental contributions and trait measurement errors are incorporated via the identity matrix I. MENDEL's implementation of this model can include both the two standard variance classes, additive and environmental, as well as the two extra variances classes, dominance and household. Inclusion of additional variance classes has no significant effect on MENDEL's speed of computation.

In general, a mixed model for QTL association mapping captures polygenic and other random effects through Ω and captures QTL fixed effects through ν . Let β denote the full vector of regression coefficients parameterizing ν . In a linear model one postulates that $\nu = A\beta$ for some predictor matrix

Genotype	Additive	Dominant	Recessive
1/1	_1	-1	-1
1/2	0	-1	+1
2/2	+1	+1	+1

Table 2: Genotype encodings for the major gene models. The additive model is the default choice. In the genotype column, "1" and "2" represent the first and second alleles for each SNP. An effect size estimate reflects the change in trait values due to each positive unit change in the encodings. For example, the default additive model estimates the mean trait difference in moving from a 1/2 genotype to a 2/2 genotype.

A incorporating relevant covariates such as age, gender, and diet. In testing association against a given SNP, *A* is augmented by an extra column whose entries encode genotypes according to one of the models (additive, dominant, and recessive) shown in see Table 2. To accommodate imprecise imputation in an additive model, these encodings can be made fractional. The corresponding component of β , β_{SNP} , is the SNP effect size. In likelihood ratio association testing one contrasts the null hypothesis $\beta_{SNP} = 0$ with the alternative hypothesis $\beta_{SNP} \neq 0$. In testing a univariate trait, the likelihood ratio statistic asymptotically follows a χ_1^2 distribution. In testing a multivariate trait with T > 1 components, each row of *A* must be replicated *T* times. The likelihood ratio statistic then asymptotically follows a χ_T^2 distribution. To implement likelihood ratio testing, iterative maximum likelihood estimation must be undertaken for each and every SNP under the alternative hypothesis. This unfortunate requirement is the major stumbling block retarding pedigree analysis.

Score tests serve as convenient substitutes for likelihood ratio tests. The current paper describes how to implement ultra-fast score tests for screening SNPs. Only SNPs with the most significant score test p-values are further subjected to the more accurate likelihood ratio test. An advantage of the likelihood ratio method is that it estimates effect sizes. In contrast, the score test only requires parameter estimates under the null hypothesis and involves no iteration beyond fitting the null model. The score vector is the gradient $\nabla L(\theta)$ of the loglikelihood $L(\theta)$, where the full parameter vector θ includes variance components such as the additive genetic variance in addition to the regression coefficient vector β . The transpose $dL(\theta)$ of the score is a row vector called the first differential of $L(\theta)$. The expected information $J(\theta)$ is the covariance matrix of the score vector. It is well known that the expected value of the observed information matrix (negative second differential) $-d^2L(\theta)$ coincides with $J(\theta)^{28}$. The score statistic

$$S(\boldsymbol{\theta}) = dL(\boldsymbol{\theta})J(\boldsymbol{\theta})^{-1}\nabla L(\boldsymbol{\theta}) \approx dL(\boldsymbol{\theta})[-d^2L(\boldsymbol{\theta})]^{-1}\nabla L(\boldsymbol{\theta})$$

is evaluated at the maximum likelihood estimates under the null hypothesis with the parameter β_{SNP} of the alternative hypothesis set to 0.

2.2 Fast Score Test for Individual SNPs

Under the multivariate model, the expected information matrix $J(\theta)$ for a single pedigree can be written in the block diagonal form

$$J(\boldsymbol{\theta}) = \begin{pmatrix} \mathrm{E}[-d_{\boldsymbol{\beta}}^{2}L(\boldsymbol{\theta})] & 0\\ 0 & \mathrm{E}[-d_{\boldsymbol{\sigma}}^{2}L(\boldsymbol{\theta})] \end{pmatrix},$$
(3)

where σ denotes the vector of variance parameters¹⁷. For independent pedigrees, the loglikelihoods (1) and corresponding score vectors and expected information matrices add. Hence, the block diagonal form of $J(\theta)$ is preserved. Because the inverse of a block diagonal matrix is block diagonal, the score statistic splits into a piece contributed by the variance components plus a piece contributed by the mean components. The maximum likelihood estimate $\hat{\theta} = (\hat{\beta}, \hat{\sigma})$ under the null model is a stationary point of the loglikelihood. Thus, the variance components segment $\nabla_{\sigma} L(\hat{\theta})$ of the score vector vanishes. We therefore focus on the mean components segment of the score vector.

If the pedigrees are labeled $1, \ldots, n$, then the pertinent quantities for implementing the score test are

$$\sum_{i=1}^{n} \nabla_{\boldsymbol{\beta}} L_{i}(\boldsymbol{\theta}) = \sum_{i=1}^{n} \boldsymbol{A}_{i}^{t} \boldsymbol{\Omega}_{i}^{-1} \boldsymbol{r}_{i}$$
$$\sum_{i=1}^{n} \mathrm{E}[-d_{\boldsymbol{\beta}}^{2} L_{i}(\boldsymbol{\theta})] = \sum_{i=1}^{n} \boldsymbol{A}_{i}^{t} \boldsymbol{\Omega}_{i}^{-1} \boldsymbol{A}_{i},$$

where $\mathbf{r}_i = \mathbf{y}_i - \mathbf{A}_i \hat{\boldsymbol{\beta}}$ is the residual for pedigree *i* and the covariance matrix Ω_i for pedigree *i* is determined by equation (2). See Chapter 8 of Lange¹⁷ for a detailed derivation of the score and expected information. Since the score statistic is calculated from estimated parameters under the null model, residuals do not change when we expand the null model to the alternative model keeping $\beta_{\text{SNP}} = 0$. Calculation of the maximum likelihood estimate $\hat{\theta}$ under the null is accomplished by a quasi-Newton algorithm whose initial step reduces to Fisher scoring^{22,17}.

For pedigree *i* under the alternative hypothesis, the design matrix A_i can be written as (a_i, N_i) , where N_i is the design matrix under the null hypothesis and a_i conveys the genotypes at the current SNP. In testing a univariate trait, the entries of a_i are taken from Table 2. If allele counts are imputed under the additive model, then the entries of a_i may be fractional numbers drawn from the interval [-1,1]. In testing a multivariate trait with T > 1 components, each row of $A_i = (a_i, N_i)$ must be replicated T times. The only exceptions to this rule occur for people missing some but not all component traits; otherwise, the covariance matrix Ω_i for pedigree *i* decomposes into a sum of Kronecker products¹⁷. Regardless of whether the trait is univariate or multivariate, one must compute the quantities

$$\sum_{i=1}^{n} \nabla_{\boldsymbol{\beta}} L_{i}(\boldsymbol{\theta}) = \begin{pmatrix} \sum_{i=1}^{n} \boldsymbol{a}_{i}^{t} \boldsymbol{\Omega}_{i}^{-1} \boldsymbol{r}_{i} \\ \sum_{i=1}^{n} \boldsymbol{N}_{i}^{t} \boldsymbol{\Omega}_{i}^{-1} \boldsymbol{r}_{i} \end{pmatrix}$$
$$\sum_{i=1}^{n} \mathrm{E}[-d_{\boldsymbol{\beta}}^{2} L_{i}(\boldsymbol{\theta})] = \begin{pmatrix} \sum_{i=1}^{n} \boldsymbol{a}_{i}^{t} \boldsymbol{\Omega}_{i}^{-1} \boldsymbol{a}_{i} & \sum_{i=1}^{n} \boldsymbol{a}_{i}^{t} \boldsymbol{\Omega}_{i}^{-1} \boldsymbol{N}_{i} \\ \sum_{i=1}^{n} \boldsymbol{N}_{i}^{t} \boldsymbol{\Omega}_{i}^{-1} \boldsymbol{a}_{i} & \sum_{i=1}^{n} \boldsymbol{N}_{i}^{t} \boldsymbol{\Omega}_{i}^{-1} \boldsymbol{N}_{i} \end{pmatrix}.$$

At the maximum likelihood estimates under the null model, the partial score vector $\sum_{i=1}^{n} N_i^t \Omega_i^{-1} r_i$ vanishes. Hence, the score statistic for testing a SNP can be expressed as

$$S = \mathbf{R}^{t} \left[\mathbf{Q} - \mathbf{W}^{t} \left(\sum_{i=1}^{n} \mathbf{N}_{i}^{t} \mathbf{\Omega}_{i}^{-1} \mathbf{N}_{i} \right)^{-1} \mathbf{W} \right]^{-1} \mathbf{R},$$

$$egin{array}{rcl} m{Q}&=&\sum_{i=1}^nm{a}_i^tm{\Omega}_i^{-1}m{a}_i, &m{R}&=&\sum_{i=1}^nm{a}_i^tm{\Omega}_i^{-1}m{r}_i, \ m{W}&=&\sum_{i=1}^nm{N}_i^tm{\Omega}_i^{-1}m{a}_i. \end{array}$$

In the score statistic *S*, the covariance matrices Ω_i^{-1} and residual vectors r_i are evaluated at the maximum likelihood estimates under the null model. Large sample theory says that *S* asymptotically follows a χ_T^2 distribution.

These formulas suggest that we precompute and store the quantities Ω_i^{-1} , $\Omega_i^{-1}N_i$, and $\Omega_i^{-1}r_i$ for each pedigree *i* and the overall sum $\sum_{i=1}^n N_i^t \Omega_i^{-1} N_i$ at the maximum likelihood estimates under the null hypothesis. From these parts, the basic elements of the score statistic can be quickly assembled. The most onerous quantity that must be computed on the fly as each new SNP is encountered is $\sum_{i=1}^n a_i^t \Omega_i^{-1} a_i$. If there are p_i people in pedigree *i*, then computation of the quadratic form $a_i^t \Omega_i^{-1} a_i$ requires $O(p_i^2)$ arithmetic operations. This looks worse than it is in practice since the entries of a_i are integers (-1, 0, and 1) in the absence of fractional imputation. This simplification allows one to avoid a fair amount of arithmetic. Assembling the remaining parts of the score statistic requires $O(p_i)$ arithmetic operations.

Individuals missing univariate trait values are omitted from analysis. Individuals missing some but not all components of a multivariate trait are retained in analysis. The proper adjustments for missing data are made automatically in the score statistic because sections of Gaussian random vectors are Gaussian.

SNPs with minor allele counts below a user-designated threshold are also omitted from analysis. Note that if the minor allele count across a study is 0, then the given SNP is mono-allelic and worthless in association testing. MENDEL's default threshold of 3 is motivated by the rule of thumb in contingency table testing that all cells have an expected count of at least 3. For a multivariate trait, a SNP may fall below the threshold for some component traits but not for others. This situation can occur when each trait displays a different pattern of missing data across individuals. MENDEL re-

where

tains such anomalous SNPs only for those component traits with a sufficient number of minor alleles. Again, proper adjustments are made automatically within the score test statistic to account for partial data.

MENDEL's analysis yields a score test p-value for each SNP. For the user-designated most significant SNPs, MENDEL's subsequent likelihood ratio test outputs an estimated SNP effect size, a standard error of that estimate, and the fraction of the total variance explained by that SNP. For a multivariate trait, MENDEL outputs a SNP effect size and associated standard error for each component trait. In the initial analysis under the null model with no SNPs, MENDEL provides estimates with standard errors of all mean and variance components included in the model. Finally, an estimate of heritability with standard error is also provided.

The extension of the score test to the multivariate *t*-distribution is straightforward¹⁸. Suppose η equals the degrees of freedom of the *t*-distribution and m_i equals the number of observed persontrait combinations for pedigree *i*. The sections of the score and expected information pertinent to the mean components for the pedigree reduce to

$$egin{array}{rcl}
abla_{oldsymbol{eta}} L_i(oldsymbol{ heta}) &=& rac{\eta+m_i}{\eta+s_i}oldsymbol{A}_i^toldsymbol{\Omega}_i^{-1}oldsymbol{r}_i \ &=& rac{\eta+m_i}{\eta+m_i+2}oldsymbol{A}_i^toldsymbol{\Omega}_i^{-1}oldsymbol{A}_i, \end{array}$$

where r_i is the residual and $s_i = r_i^t \Omega_i^{-1} r_i$ is the associated Mahalanobis distance. A sensible choice for η is its estimate under the null model.

2.3 Kinship Estimation From SNPs

MENDEL can either calculate the global kinship coefficient matrix Φ from the provided pedigree structures or estimate it from dense genotypes. In global kinship estimation MENDEL's default uses an evenly spaced 20% of the available SNPs, and only compares pairs of individuals within defined pedigrees. Hence, Φ is block diagonal. Users can trivially elect to exploit a larger fraction of the available SNPs or estimate kinship for *all* pairs of individuals. Given *S* selected SNPs, MENDEL estimates the global kinship coefficient of individuals *i* and *j* based on either the genetic relation matrix (GRM) method

$$\hat{\Phi}_{ij} = \frac{1}{2S} \sum_{k=1}^{S} \frac{(x_{ik} - 2p_k)(x_{jk} - 2p_k)}{2p_k(1 - p_k)}$$

or the method of moments (MoM)^{7,19}

$$\hat{\Phi}_{ij} = \frac{e_{ij} - \sum_{k=1}^{S} \left[p_k^2 + (1 - p_k)^2 \right]}{S - \sum_{k=1}^{S} \left[p_k^2 + (1 - p_k)^2 \right]},$$

where p_k is the minor allele frequency at SNP k, x_{ik} is the number of minor alleles in *i*'s genotype at SNP k, and

$$e_{ij} = \frac{1}{4} \sum_{k=1}^{S} \left[x_{ik} x_{jk} + (2 - x_{ik})(2 - x_{jk}) \right]$$

is the observed fraction of alleles identical by state (IBS) between i and j. The GRM method is MENDEL's default. In general, one can think of the GRM method centering and scaling each genotype, while the MoM method uses the raw genotypes and then centers and scales the final result.

2.4 Other Utilities for Handling Pedigree Data

To encourage thorough testing of new statistical methods, such as the current Ped-GWAS score test, we have implemented both genotype and trait simulation in our genetic analysis program MENDEL²⁰. MENDEL does genotype simulation (gene dropping) subject to prescribed allele frequencies, a given genetic map, and Hardy-Weinberg and linkage equilibrium. If one fixes founder haplotypes and simulates conditional on these, then the unrealistic assumption of linkage equilibrium can be relaxed. Missing data patterns are respected or imposed by the user. It is also possible to set the rate for randomly deleting data and to simulate genotypes for people of mixed ethnicity by defining different ancestral populations, each with its own allele frequencies. If this feature is invoked, then each pedigree founder should be assigned to a population.

Trait simulation can be layered on top of genotype simulation. MENDEL simulates either univariate

traits determined by generalized linear models or multivariate Gaussian traits determined by variance component models. The biggest limitations are the restriction to a single major locus and the generalized linear model assumption that trait correlations are driven solely by this locus. Variance component models enable inclusion of environmental effects and more complicated correlations among relatives. In the variance component setting, univariate as well as multivariate Gaussian traits can be simulated. Most variance component models are built on Gaussian distributions, but MENDEL allows one to replace these by multivariate *t*-distributions. Thus, users can investigate robust statistics less prone to distortion by outliers. More theoretical and implementation details appear in the MENDEL documentation²⁰.

3 Results

3.1 The San Antonio Family Heart Study

We analyzed a real data set collected by the San Antonio Family Heart Study (SAFHS)²⁵. The data consist of 3637 individuals in 211 Mexican American families. High-density lipoprotein (HDL) levels were measured at up to three time points for each of the 1429 phenotyped individuals. These traits are denoted HDL₁, HDL₂, and HDL₃, measured at corresponding ages AGE₁, AGE₂, and AGE₃. Some of the phenotyped individuals have HDL measurements at only one or two of the time points. Of the 1429 phenotyped individuals, 1413 were genotyped at 944,427 genome-wide SNPs. The genotyping success rate exceeded 98% in 1388 of these individuals over 124 pedigrees. The largest family contains 247 individuals; five others also contain more than 90 individuals. The smallest pedigree was a singleton. Genotyping success rates were above 98% for 935,392 SNPs.

3.2 Comparison with FAST-LMM and GEMMA

For fair comparisons, we directed MENDEL to estimate SNP-based global kinship coefficients for all pairs of individuals ignoring the input pedigrees. This is the default in FAST-LMM and GEMMA. In addition, we ran MENDEL's default in which the coefficients are estimated only for pairs of individuals

within the same input pedigree. We also slightly adjusted some of the default quality control thresholds so the programs would be analyzing roughly the same set of SNPs and individuals. For example, by default MENDEL filters SNPs with fewer than three occurrences of the minor allele in the data; in contrast, FAST-LMM only filters SNPs with zero occurrences of the minor allele, and GEMMA filters SNPs with minor allele frequency (MAF) < 0.01. All other defaults were observed throughout. Users can easily adjust the MENDEL analysis parameters via its control file and the FAST-LMM and GEMMA analysis parameters via their command line.

We first carried out three univariate QTL analyses of HDL₁, HDL₂, and HDL₃, using SEX and AGE₁, AGE₂, or AGE₃ as covariates. We then ran a multivariate QTL analysis of HDL₁, HDL₂, and HDL₃ jointly, which we refer to as HDL_{Joint}. For the multivariate analysis, the most appropriate configuration is to constrain the effects of the SEX and AGE covariates to be the same on all three HDL measurements. Such linear constraints are imposed in MENDEL via a few simple lines in its control file. FAST-LMM and GEMMA do not allow constraints on covariates. Therefore, we also ran a multivariate analysis with only the SEX covariate and no constraints. With no constraints, SEX will have a slightly different effect on each component phenotype in the multivariate analysis. For instance, MENDEL's default run estimated a female effect of 2.5 ± 0.3 on HDL₁, 2.1 ± 0.4 on HDL₂, and 2.7 ± 0.4 on HDL₃. FAST-LMM cannot perform any multivariate analyses.

Table 3 reports all SNPs with MAF > 0.01 that achieve genome-wide significance (p-values less than 5×10^{-8}) as reported by at least one software package. For the univariate analyses, each software package found the same set of significant SNPs, except that one of GEMMA's p-values was slightly short of the significance threshold. Figure 1 shows a Manhattan plot and a Q-Q plot from the HDL₁ analysis by MENDEL given kinship estimates for all pairs of individuals. The results for the other analyses, both univariate and multivariate, were similar. Each MENDEL all-pairs univariate analysis had genomic control λ in the range 1.002 to 1.006; in default mode, λ was in the range 0.992 to 1.022. The various Q-Q plots and associated λ values show there is no systematic biases in the data or analysis. In the all-pairs MENDEL HDL₁ analysis, the grand mean (intercept) was 49.0 \pm 0.8. The SEX covariate was significant in all null models. For example, in the all-pairs MENDEL HDL_{Joint} analysis with constrained covariates, the SEX effect was 2.4 \pm 0.3 for females and, by design, the

opposite for males. The AGE covariate was not significant in any run. For example, again in the allpairs HDL_{Joint} analysis with parameter constraints, the AGE effect was 0.04 ± 0.02 . In the null model for the all-pairs MENDEL HDL₁ analysis, the additive variance was estimated as 78.8 \pm 9.9, and the environmental variance was estimated as 78.1 \pm 7.2. This gives an overall heritability estimate for HDL₁ of 0.50 ± 0.04 . Similar variance estimates were seen in other null models.

For the multivariate analysis without parameter constraints, MENDEL is able to include almost twice as many individuals in the analysis as GEMMA (see Table 4). GEMMA only includes individuals phenotyped at all component traits and covariates. This probably explains why MENDEL finds several more SNPs with significant p-values than GEMMA.

Table 4 tallies the run times and memory footprints from each analysis on a typical personal computer with adequate RAM to accommodate FAST-LMM (6 CPU cores at 2.67 GHz, with 48 GB total RAM). Even when estimating the global kinship coefficients for all pairs of individuals, each univariate QTL run took MENDEL less than 8 minutes to read, quality check, and analyze the data for kinship estimates and association tests, roughly 10% of the time required for FAST-LMM and 5% of the time required by GEMMA. (For GEMMA, the kinship estimation and association tests are run separately. The run times reported here are their total.)

The three programs use different association test strategies: MENDEL performs score tests for all SNPs and LRTs for the top SNPs; FAST-LMM performs LRTs; and GEMMA by default performs Wald tests, but the user can change this to LRTs or score tests. For the univariate analyses on a six-core computer, excluding estimation of kinship coefficients, GEMMA's run times under the Wald test and LRT options were roughly similar to FAST-LMM's; GEMMA's run time under the score test option was roughly double MENDEL's in all-pairs mode. This is impressive given GEMMA's lack of multithreading. It is kinship estimation, which in practice can be done once per data set, that is substantially slower in GEMMA (running roughly 135 minutes) than in FAST-LMM or MENDEL (less than 1 minute).

Each trivariate QTL run took MENDEL less than 90 minutes. MENDEL required roughly one-eighth the time of GEMMA while analyzing almost twice as many individuals. MENDEL is also memory efficient. The univariate and multivariate runs each required less than 1.5 GB of memory, which is well below the amount of RAM in a typical computer. FAST-LMM's memory usage is more than 15 times larger than MENDEL's. GEMMA uses even less memory than MENDEL but is considerably slower.

4 Discussion

We have implemented an ultra-fast algorithm for QTL analysis of pedigree data or mix of population and pedigree data. In our opinion MENDEL's comprehensive environment for genetic data analysis is a decided advantage. In addition to its exceptional speed and memory efficiency, MENDEL can handle multivariate quantitative traits and detect outlier trait values and pedigrees. Most competing programs ignore multivariate traits and outliers altogether.

A recent review of univariate QTL analysis packages for family data⁸ shows that all the explored packages obtain similar results, leaving speed, features, and ease of use as the important factors in choosing between them. Once the current version of MENDEL came out, the authors of the review were kind enough to add a comment (http://www.plosgenetics.org/annotation/listThread. action?root=81847www.plosgenetics.org/annotation/listThread.action?root=81847) to their article observing that MENDEL was now the fastest and one of the easiest to use packages they reviewed.

In the SAFHS example data set we used with HDL phenotypes, all the significant SNPs we found had MAF < 0.01. Due to these low MAFs, we do not claim these SNPs are strong candidates for further study. However, the key point here is that all four methods found the same SNPs, at least for the univariate analyses. We also note that the p-values are quite similar regardless of whether one uses kinship estimates between all individuals (MENDEL's all-pairs mode) or only between individuals within the same input pedigrees (MENDEL's default mode). This suggests that the input pedigree structures for this data set are substantially correct and complete, with few mistaken or hidden relationships. Obviously, this may not be true for other data sets. By supplying good kinship estimates ignoring pedigree structures, the currently reviewed packages make the hard fieldwork of relationship discovery superfluous.

A future version of MENDEL will address its failure to read fractional genotype values. This is

simply a logistical issue, as all MENDEL's internal genotype computations are already handled as floating point operations. Another imminent feature is a fourth style of kinship coefficient estimation that allows the user to force theoretical kinship coefficients for pairs of individuals within the same pedigree and estimated kinships for all other pairs.

By supplying a comprehensive, fast, and easy to use package for GWAS on quantitative traits in general pedigrees, we hope to encourage exploitation of family-based data sets for gene mapping. A gene mapping study should collect as large a sample as possible consistent with economic constraints and uniform trait phenotyping. If the sample includes pedigrees, all the better. One should *not* let the choice of statistical test determine the data collected; on the contrary, the data should determine the test. Here we have argued that score tests can efficiently handle unrelated individuals, pedigrees, or a mixture of both. For human studies, where controlled breeding is forbidden, nature has provided pedigrees segregating every genetic trait. Many of these pedigrees are known from earlier linkage era studies and should be treasured as valuable resources.

Let us suggest a few directions for future work. The current method works marker by marker and is ill equipped to perform model selection. Lasso penalized regression is available to handle model selection for case-control and random sample data 40,39,43,42 and can be generalized to variance component models. Although we have generalized the score test to distributions such as the multivariate t, extending it to discrete traits may be out of reach. For likelihood based methods, there simply are no discrete analogues of the Gaussian distribution that lend themselves to graceful evaluation of pedigree likelihoods. Treating case/control data as a 0/1 quantitative variable is a possibility that has been explored by Pirinen et al.²⁷. The GEE method is another fallback option because it does not depend on precise distributional assumptions.

In rare variant mapping, grouping related SNPs in a variance component may be a good alternative to the mean component models used here. Each variant may be too rare to achieve significance in hypothesis testing. Fortunately, aggregating genotype information within biological units such as genes or pathways offer better power than marginal testing of individual SNPs. See Asimit and Zeggini² for a recent review of aggregation strategies. Kwee et al.¹⁴ have successfully applied a variance component model for association testing of SNP sets in a sample of unrelated subjects. Rönnegård et al.²⁹ consider score tests for random effects models in the context of experimental line crosses. Score tests may well be the key to implementing random effect models in pedigrees. However, the computational demands are apt to be more formidable than those encountered here with fixed effects models. In particular, if tests are based simply on local identity-by-descent (IBD) sharing, then the boundaries between pedigrees disappear, and the entire sample collapses to one large pedigree. The required local kinship coefficients can again be well estimated from dense markers, but this demands more computation than the estimation of global kinship coefficients under the mean components model advocated here⁷. Since inversion of a pedigree covariance matrix scales as the cube of the number of individuals in the pedigree, treating the entire sample as a single pedigree will put a practical upper limit on sample size. There are other issues in implementing variance component models such as assigning p-values and dealing with multivariate traits that are best left to a separate paper.

Acknowledgement

Funding: NIH (GM053275, HG006139, MH059490) and NSF (DMS-1310319). Burroughs Wellcome Fund Inter-school Training Program in Metabolic Diseases (KHKC).

References

- Amin, N., van Duijn, C. M., and Aulchenko, Y. S. (2007). A genomic background based method for association analysis in related individuals. *PLoS ONE*, 2(12):e1274.
- [2] Asimit, J. and Zeggini, E. (2010). Rare variant association analysis methods for complex traits. *Annual Review of Genetics*, 44(1):293–308.
- [3] Aulchenko, Y. S., de Koning, D.-J., and Haley, C. (2007). Genomewide rapid association using mixed model and regression: A fast and simple method for genomewide pedigree-based quantitative trait loci association analysis. *Genetics*, 177(1):577–585.

- [4] Chen, M.-H., Liu, X., Wei, F., Larson, M. G., Fox, C. S., Vasan, R. S., and Yang, Q. (2011). A comparison of strategies for analyzing dichotomous outcomes in genome-wide association studies with general pedigrees. *Genetic Epidemiology*, 35(7):650–657.
- [5] Chen, M.-H. and Yang, Q. (2010). GWAF: an R package for genome-wide association analyses with family data. *Bioinformatics*, 26(4):580–581.
- [6] Chen, W.-M. and Abecasis, G. R. (2007). Family-based association tests for genome-wide association scans. *American Journal of Human Genetics*, 81(5):913–926.
- [7] Day-Williams, A. G., Blangero, J., Dyer, T. D., Lange, K., and Sobel, E. M. (2011). Linkage analysis without defined pedigrees. *Genetic Epidemiology*, 35(5):360–370.
- [8] Eu-ahsunthornwattana, J., Miller, E. N., Fakiola, M., Wellcome Trust Case Control Consortium 2, Jeronimo, S. M. B., Blackwell, J. M., and Cordell, H. J. (2014). Comparison of methods to account for relatedness in genome-wide association studies with family-based data. *PLoS Genetics*, 10(7):e1004445.
- [9] Hindorff, L. A., Sethupathy, P., Junkins, H. A., Ramos, E. M., Mehta, J. P., Collins, F. S., and Manolio, T. A. (2009). Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proceedings of the National Academy of Sciences*, 106(23):9362– 9367.
- [10] Kang, H. M., Sul, J. H., Service, S. K., Zaitlen, N. A., Kong, S.-Y., Freimer, N. B., Sabatti, C., and Eskin, E. (2010). Variance component model to account for sample structure in genome-wide association studies. *Nature Genetics*, 42(4):348–354.
- [11] Kang, H. M., Zaitlen, N. A., Wade, C. M., Kirby, A., Heckerman, D., Daly, M. J., and Eskin, E. (2008). Efficient control of population structure in model organism association mapping. *Genetics*, 178:1709–1723.
- [12] Korte, A., Vilhjalmsson, B. J., Segura, V., Platt, A., Long, Q., and Nordborg, M. (2012). A mixed-

model approach for genome-wide association studies of correlated traits in structured populations. *Nature Genetics*, 44(9):1066–1071.

- [13] Ku, C. S., Loy, E. Y., Pawitan, Y., and Chia, K. S. (2010). The pursuit of genome-wide association studies: where are we now? *Journal of Human Genetics*, 55(4):195–206.
- [14] Kwee, L. C., Liu, D., Lin, X., Ghosh, D., and Epstein, M. P. (2008). A powerful and flexible multilocus association test for quantitative traits. *American Journal of Human Genetics*, 82(2):386– 397.
- [15] Laird, N. M., Horvath, S., and Xu, X. (2000). Implementing a unified approach to family-based tests of association. *Genetic Epidemiology*, 19(Suppl 1):S36–S42.
- [16] Lange, C. and Laird, N. M. (2002). On a general class of conditional tests for family-based association studies in genetics: the asymptotic distribution, the conditional power, and optimality considerations. *Genetic Epidemiology*, 23(2):165–180.
- [17] Lange, K. (2002). Mathematical and Statistical Methods for Genetic Analysis. Statistics for Biology and Health. Springer-Verlag, New York, 2nd edition.
- [18] Lange, K., Little, R. J. A., and Taylor, J. M. G. (1989). Robust statistical modeling using the t distribution. *Journal of the American Statistical Association*, 84(408):881–896.
- [19] Lange, K., Papp, J. C., Sinsheimer, J. S., and Sobel, E. M. (2014). Next-generation statistical genetics: Modeling, penalization, and optimization in high-dimensional data. *Annual Review of Statistics and Its Application*, 1(1):279–300.
- [20] Lange, K., Papp, J. C., Sinsheimer, J. S., Sripracha, R., Zhou, H., and Sobel, E. M. (2013).Mendel: The Swiss army knife of genetic analysis programs. *Bioinformatics*, 29(12):1568–1570.
- [21] Lange, K., Sinsheimer, J. S., and Sobel, E. (2005). Association testing with Mendel. *Genetic Epidemiology*, 29(1):36–50.
- [22] Lange, K., Westlake, J., and Spence, M. A. (1976). Extensions to pedigree analysis iii. variance components by the scoring method. *Annals of Human Genetics*, 39(4):485–491.

- [23] Lippert, C., Listgarten, J., Liu, Y., Kadie, C. M., Davidson, R. I., and Heckerman, D. (2011). FaST linear mixed models for genome-wide association studies. *Nature Methods*, 8(10):833–835.
- [24] Listgarten, J., Lippert, C., Kadie, C. M., Davidson, R. I., Eskin, E., and Heckerman, D. (2012). Improved linear mixed models for genome-wide association studies. *Nature Methods*, 9(6):525– 526.
- [25] Mitchell, B. D., Kammerer, C. M., Blangero, J., Mahaney, M. C., Rainwater, D. L., Dyke, B., Hixson, J. E., Henkel, R. D., Sharp, R. M., Comuzzie, A. G., VandeBerg, J. L., Stern, M. P., and MacCluer, J. W. (1996). Genetic and environmental contributions to cardiovascular risk factors in Mexican Americans: The San Antonio Family Heart Study. *Circulation*, 94(9):2159–2170.
- [26] Ott, J., Kamatani, Y., and Lathrop, M. (2011). Family-based designs for genome-wide association studies. *Nature Reviews Genetics*, 12(7):465–474.
- [27] Pirinen, M., Donnelly, P., and Spencer, C. C. A. (2013). Efficient computation with a linear mixed model on large-scale data sets with applications to genetic studies. *Annals of Applied Statistics*, 7(1):369–390.
- [28] Rao, C. (2009). Linear Statistical Inference And Its Applications. Wiley, 2nd edition.
- [29] Rönnegård, L., Besnier, F., and Carlborg, O. (2008). An improved method for quantitative trait loci detection and identification of within-line segregation in f2 intercross designs. *Genetics*, 178(4):2315–2326.
- [30] Spielman, R. S. and Ewens, W. J. (1998). A sibship test for linkage in the presence of association: the sib transmission/disequilibrium test. *American Journal of Human Genetics*, 62(2):450–458.
- [31] Stanhope, S. A. and Abney, M. (2012). GLOGS: a fast and powerful method for GWAS of binary traits with risk covariates in related populations. *Bioinformatics*, 28(11):1553–1554.
- [32] Thornton, T. and McPeek, M. S. (2007). Case-control association testing with related individuals: A more powerful quasi-likelihood score test. *American Journal of Human Genetics*, 81(2):321–337.

- [33] Thornton, T. and McPeek, M. S. (2010). ROADTRIPS: Case-control association testing with partially or completely unknown population and pedigree structure. *American Journal of Human Genetics*, 86(2):172–184.
- [34] Van Steen, K. (2011). Perspectives on genome-wide multi-stage family-based association studies. Statistics in Medicine, 30(18):2201–2221.
- [35] Van Steen, K. and Lange, C. (2005). PBAT: a comprehensive software package for genome-wide association analysis of complex family-based studies. *Human Genomics*, 2(1):67–69.
- [36] Visscher, P. M., Brown, M. A., McCarthy, M. I., and Yang, J. (2012). Five years of GWAS discovery. American Journal of Human Genetics, 90(1):7–24.
- [37] Won, S., Bertram, L., Becker, D., Tanzi, R., and Lange, C. (2009a). Maximizing the power of genome-wide association studies: A novel class of powerful family-based association tests. *Statistics in Biosciences*, 1(2):125–143.
- [38] Won, S., Wilk, J. B., Mathias, R. A., O'Donnell, C. J., Silverman, E. K., Barnes, K., O'Connor, G. T., Weiss, S. T., and Lange, C. (2009b). On the analysis of genome-wide association studies in family-based designs: A universal, robust analysis approach and an application to four genome-wide association studies. *PLoS Genetics*, 5(11):e1000741.
- [39] Wu, T. T., Chen, Y., Hastie, T., Sobel, E. M., and Lange, K. (2009). Genome-wide association analysis by lasso penalized logistic regression. *Bioinformatics*, 25(6):714–721.
- [40] Wu, T. T. and Lange, K. (2008). Coordinate descent algorithms for lasso penalized regression. Annals of Applied Statistics, 2(1):224–244.
- [41] Zhang, Z., Ersoz, E., Lai, C.-Q., Todhunter, R. J., Tiwari, H. K., Gore, M. A., Bradbury, P. J., Yu, J., Arnett, D. K., Ordovas, J. M., and Buckler, E. S. (2010). Mixed linear model approach adapted for genome-wide association studies. *Nature Genetics*, 42(4):355–360.

- [42] Zhou, H., Alexander, D., Sehl, M., Sinsheimer, J. S., Sobel, E. M., and Lange, K. (2011). Penalized regression for genome-wide association screening of sequence data. *Pacific Symposium on Biocomputing*, 2011:106–117.
- [43] Zhou, H., Sehl, M. E., Sinsheimer, J. S., and Lange, K. (2010). Association screening of common and rare genetic variants by penalized regression. *Bioinformatics*, 26(19):2375–2382.
- [44] Zhou, X. and Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association studies. *Nature Genetics*, 44:821–824.
- [45] Zhou, X. and Stephens, M. (2014). Efficient multivariate linear mixed model algorithms for genome-wide association studies. *Nature Methods*, 11(4):407–409.
- [46] Zhu, Y. and Xiong, M. (2012). Family-based association studies for next-generation sequencing. *American Journal of Human Genetics*, 90(6):1028–1045.

Trait	SNP	Chr.	Base Pair Position	MAF	$-\log_{10}(\text{p-val})$ MENDEL default	$-\log_{10}(p\text{-val})$ Mendel all-pairs	$-\log_{10}(p-val)$ – FAST-LMM
	rs7303112	12	97,596,023	0.00455	10.21	10.71	7.63
HDL_1	rs8040647	15	32,304,988	0.00147*	7.44	7.56	7.35
	rs9972594	15	32,421,102	0.00147*	7.44	7.56	7.37
	rs7167103	15	32,830,477	0.00147*	7.44	7.56	7.35
HDL_2	rs7100957	10	28,207,332	0.00183*	8.84	8.95	8.88
HDL_3	rs17060933	8	22,510,029	0.00382	8.23	8.28	8.61
	rs7303112	12	97,596,023	0.00644	9.89	9.94	
HDL _{Joint}	rs16925210	10	25,308,103	0.00217	8.15	8.33	
with	rs7091416	10	25,318,381	0.00217	8.15	8.33	Not
constrained	rs10075658	5	148,911,957	0.00144*	8.16	8.21	Available
covariates	rs7733139	5	145,977,990	0.00217	7.36	7.34	
	rs7100957	10	28,207,332	0.00870	7.20	7.30	
	rs7303112	12	97,596,023	0.00644	9.82	9.88	
HDL _{Joint}	rs16925210	10	25,308,103	0.00217	8.04	8.23	
without	rs7091416	10	25,318,381	0.00217	8.04	8.23	Not
constrained	rs10075658	5	148,911,957	0.00144*	8.12	8.17	Available
covariates	rs7733139	5	145,977,990	0.00217	7.41	7.40	
	rs7100957	10	28,207,332	0.00870	7.19	7.30	
	rs10083226	13	104,434,452	0.00219	7.10	7.31	

Table 3: All SNPs with minor allele frequency (MAF) above 0.001 that reach genome-wide significance in any of the analyses of the HDL traits from the San Antonio Family Heart Study (SAFHS). All default parameters were used except for minor changes to the quality control thresholds (see text). Also, MENDEL was run in both default and all-pairs modes. MENDEL's default mode estimates non-zero global kinship coefficients only for pairs of individuals within the same input pedigree; MENDEL in all-pairs mode, FAST-LMM, and GEMMA estimate coefficients for all pairs of individuals. Genomewide significance was declared for p-values $< 5 \times 10^{-8} \implies -\log_{10}(p-value) > 7.3$. The SAFHS has 1413 genotyped and phenotyped individuals in 124 pedigrees. The genotypes include roughly 1 million SNPs. The phenotypes include the subjects' high-density lipoprotein (HDL) level and age at three time points. The HDL_{Joint} runs are multivariate analyses of HDL₁, HDL₂, and HDL₃ jointly; all other runs are univariate analyses. See the text for a list of the covariates used in each analysis. Note that in the multivariate analysis, MENDEL is able to use roughly twice as many individuals as GEMMA (see text and Table 4), which may explain the less significant findings for GEMMA. Each MAF is based on the pedigree founders, except where marked by an asterisk (*). In these cases the minor allele did not appear in the genotyped founders, and its frequency was estimated from all genotyped individuals.



Figure 1: The results of MENDEL'S HDL₁ univariate analysis in the SAFHS data set with global kinship coefficients estimated for all pairs of individuals. Upper: The Manhattan plot graphs roughly one million SNPs against their $-\log_{10}(p\text{-value})$. The horizontal line is the genome-wide significance threshold, $7.3 = -\log_{10}(5 \times 10^{-8})$. Lower: The Q-Q plot graphs the observed $-\log_{10}(p\text{-value})$ quantiles versus their expectations. The genomic control value of $\hat{\lambda} = 1.006$ derived from this comparison suggests no systematic biases in the data or analysis.

Program	Trait Ana	alyzed	Analyzed	RunTime	RAM
	S	Samples	SNPs	(min:sec)	(GB)
MENDEL default MENDEL all-pairs FAST-LMM GEMMA	HDL_1	1357 1357 1397 1397	935,392 935,392 941,546 919,050	1:51 7:49 76:11 206:54	1.2 1.2 30.0
MENDEL default	HDL_2	818	935,392	1:33	1.1
MENDEL all-pairs		818	935,392	3:25	1.1
FAST-LMM		840	934,216	49:44	18.0
GEMMA		840	914,051	180:21	0.3
MENDEL default	HDL ₃	914	935,392	1:38	1.1
MENDEL all-pairs		914	935,392	3:54	1.1
FAST-LMM		939	937,208	54:58	20.0
GEMMA		939	918,626	182:26	0.3
MENDEL default MENDEL all-pairs FAST-LMM GEMMA	HDL _{Joint} with constrained covariates	1388 935,392 4:08 1388 935,392 83:24 ied Not Available iss Not Available		4:08 83:24 ot Available ot Available	1.2 1.2
MENDEL default MENDEL all-pairs FAST-LMM GEMMA	HDL _{Joint} without constrained covariates	1388 1388 1 712	935,392 935,392 No 912,318	3:49 80:04 ot Available 630:37	1.2 1.2 0.6

Table 4: Comparison of run times and memory (RAM) usage on a typical computer but with adequate RAM to accommodate FAST-LMM (6 CPU cores at 2.67 GHz, with 48 GB total RAM). The listed run times include reading the data set, performing quality checks, estimating the kinship coefficients, and calculating the association test p-values. All default parameters were used except for minor changes to the quality control thresholds (see text). Also, MENDEL was run in both default and all-pairs modes. MENDEL's default mode estimates non-zero global kinship coefficients only for pairs of individuals within the same input pedigree; MENDEL in all-pairs mode, FAST-LMM, and GEMMA estimate coefficients for all pairs of individuals. For the multivariate analysis, MENDEL includes roughly twice as many individuals as GEMMA because GEMMA only analyzes individuals phenotyped at all component traits and covariates. MENDEL performs score tests for all SNPs and LRTs for the top SNPs; FAST-LMM performs LRTs; and GEMMA by default performs Wald tests, but the user can change this to LRTs or score tests. Using score tests in GEMMA would make it faster (see text).