A note on gauge-fixing in the electroweak sector of nmUED

Anindya Datta¹ *, Avirup Shaw² [†]

¹Department of Physics, University of Calcutta, 92 Acharya Prafulla Chandra Road, Kolkata 700009, India

²Regional Centre for Accelerator-based Particle Physics, Harish-Chandra Research Institute, Chhatnag Road, Jhusi, Allahabad 211019, India

Abstract

Electroweak observables are highly sensitive to the loop corrections. Therefore, a proper gauge-fixing mechanism is always needed to define the propagators which are involved in Feynman loop amplitude. With this spirit we compute gauge-fixing mechanism in five dimensional (5-D) Universal Extra-Dimensional (UED) model with boundary localised terms (BLTs). These BLTs are not 5-D operators in four-dimensional (4-D) effective theory but some sort of boundary conditions on the respective fields at the fixed points of S^1/Z_2 orbifold. Furthermore, these BLTs non-trivially modify the Kaluza-Klein (KK) spectra and some of the interactions among the KK-excitations compared to the minimal UED (mUED), in which, these BLTs are absent. In this note we calculate the gauge-fixing mechanism in the electroweak sector of such non-trivial UED scenario. Moreover, we discuss the composition and masses of Goldstone and any physical scalar that emerge after the symmetry breaking in this set up with different choices of gauge.

PACS No: 11.10 Kk, 11.15.-q, 12.60.-i

Extra dimensional models in which all the Standard Model (SM) fields propagate in one or more compactified space-like extra dimensions offer viable solutions to several long standing problems of Particle Physics like Dark Matter [1, 2, 3, 4, 5], Unification [6, 7, 8] etc. We are interested in a particular incarnation of such models namely UED [9] where all the SM fields can feel a single compactified extra space-like dimension y. The effective 4-D action of UED, consists of SM particles and towers of their KK-excitations. Members of a tower is specified by so called KK-number (n), which is nothing but the discretised momentum in direction of the extra dimension. Masses of the KK-excitations in n^{th} KK-state is typically nR^{-1} , where R is the radius of compactification. The interactions among the fields are governed by the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge symmetry of the SM. To accommodate the chiral fermions of the SM, one has to demand for some extra Z_2 symmetry of the action after orbifolding the extra direction. These results into two special points y = 0 and $y = \pi R$ called the *fixed points*, along the y direction.

Radiative corrections to the masses of the KK-states are important from phenomenological view point and has been considered in [10, 11]. UED being an effective theory, lacks the ultraviolet

^{*}email: adphys@caluniv.ac.in

[†]email: avirup.cu@gmail.com

completion in radiative corrections. So assumptions have been made in Ref.[11], such that at the cut-off scale, radiative corrections to all KK-masses identically vanish. In principle, one can take into account the contributions (to the radiative corrections) beyond the cut-off, by adding BLTs at the fixed points (y = 0 and $y = \pi R$) with their coefficients as free parameters. These models are collectively called non-minimal UED (nmUED). Theoretical as well as different possibilities of phenomenological aspects of nmUED have been investigated to some extent in existing literature [12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27].

In this note, we discuss the gauge-fixing procedure in the context of nmUED model. In literature there exist several articles [28, 29, 30, 31] in which the issue of gauge-fixing procedure has been discussed in the context of UED, but in case of nmUED it would not be a straight forward job due to the presence of boundary terms. To our knowledge, the first time we are going to analyse this issue in electroweak sector of this non-trivial extension of UED scenario. Furthermore, we will discuss the masses and composition of Goldstone and other physical scalars emerging after the electroweak symmetry breaking. At this point we would like to make a comment that the gauge-fixing mechanism for 5-D QED in presence of BLTs, has been discussed in [32]. We extend the prescription in Ref.[32] to the case when the symmetry is spontaneously broken in a *non-abelian gauge theory*.

Let us start with the free action (governed by $SU(2)_L \times U(1)_Y$ gauge group) of 5-D gauge and scalar fields with their respective boundary localised kinetic terms (BLKTs) [18, 19, 20, 21, 22, 23, 24, 25, 26]:

$$S_{V} = -\frac{1}{4} \int d^{5}x \bigg[W_{MN} W^{MN} + r_{W} \{ \delta(y) + \delta(y - \pi R) \} W_{\mu\nu} W^{\mu\nu} + B_{MN} B^{MN} + r_{B} \{ \delta(y) + \delta(y - \pi R) \} B_{\mu\nu} B^{\mu\nu} \bigg], \qquad (1)$$

$$S_{H} = \int d^{5}x \bigg[(D_{M}H)^{\dagger} (D^{M}H) + r_{H} \{ \delta(y) + \delta(y - \pi R) \} (D_{\mu}H)^{\dagger} (D^{\mu}H) \bigg].$$
(2)

Here, r_W , r_B and r_H parametrise the strength of the BLKTs for the respective fields. 5-D field strength tensors are given below:

$$W^{a}_{MN} \equiv (\partial_{M}W^{a}_{N} - \partial_{N}W^{a}_{M} - \tilde{g}_{2}\epsilon^{abc}W^{b}_{M}W^{c}_{N}), \qquad (3)$$
$$B_{MN} \equiv (\partial_{M}B_{N} - \partial_{N}B_{M}).$$

 W_M^a and B_M (M = 0, 1...4) are the 5-D gauge fields corresponding to $SU(2)_L$ and $U(1)_Y$ gauge group respectively. 5-D covariant derivative is defined as $D_M \equiv \partial_M + i\tilde{g}_2 \frac{\sigma^a}{2} W_M^a + i\tilde{g}_1 \frac{Y}{2} B_M$, with the 5-D gauge coupling constants \tilde{g}_2 and \tilde{g}_1 . $\frac{\sigma^a}{2}$ and $\frac{Y}{2}$ are the corresponding generators of the gauge groups. $H = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$ is the 5-D Higgs doublet with $\phi^0 = \frac{(\tilde{v}+h+i\chi)}{\sqrt{2}}$, \tilde{v} being the 5-D vacuum expectation value (VEV). Before going to discuss gauge-fixing mechanism we would like to make some clarifying remarks which could help the reader to understand the following gauge-fixing action. First of all, generally KK-decomposition of neutral gauge bosons become very complicated by the fact that Band W^3 mix in the bulk as well as on the boundary. So, unless $r_W = r_B$, it would not be possible to diagonalise the bulk and boundary actions simultaneously by the same 5-D field redefinition. In the following we will stick to the $r_W = r_B$ equality condition. However, in general one can deal with $r_W \neq r_B$, but in this case the mixing term between B and W^3 in the bulk and on the boundary points generate off-diagonal terms in the neutral gauge boson mass matrix. In this situation Unitary gauge (in which it is difficult to calculate loop corrections) choice is the only one favourable option where we can play with this nmUED scenario [19, 20, 21, 22, 23, 24]. The reason is due to the fact that in UED or nmUED like scenario one can argue about the Unitary gauge choice condition which is equivalent to large R^{-1} limit [19, 20, 21, 22, 23, 24]. As in this case contribution to the KK-masses from spontaneous breaking of the electroweak symmetry are negligible with respect to extra-dimensional contribution. Later we will discuss this issue elaborately.

Eq.1 and 2 must be supplemented by the gauge-fixing action. One of the aims for fixing the gauge is to get rid of the terms in which V_{μ} couples to V_4 (arise in Eq.1) and/or to $\eta (\equiv \phi^+ \text{ or } \chi)$ (in Eq.2). Here we use V to generically denote vector fields and also we use r_V as the generic BLKT parameter for gauge bosons. Further, we use η as the generic notation for 5-D Goldstone fields. To ensure the (second) cancellation, Higgs BLKT parameter r_H should be included in the gauge-fixing action. We have adapted the gauge-fixing action from [33, 34], where a similar kind of gauge-fixing action can be found in presence of boundary loclaised terms. However, we have considered the following gauge-fixing action appropriate for nmUED model¹ (for the purpose of writing gauge-fixing action we use the conventional notations W^{\pm} , Z (for weak gauge bosons) and A (for photon), however in rest of our analysis we will use generic notation V),

$$S_{GF} = -\frac{1}{\xi_y} \int d^5 x \Big| \partial_\mu W^{\mu +} + \xi_y (\partial_y W^{4+} + i M_W \phi^+ \{1 + r_H (\delta(y) + \delta(y - \pi R))\}) \Big|^2 - \frac{1}{2\xi_y} \int d^5 x [\partial_\mu Z^\mu + \xi_y (\partial_y Z^4 - M_Z \chi \{1 + r_H (\delta(y) + \delta(y - \pi R))\})]^2 - \frac{1}{2\xi_y} \int d^5 x [\partial_\mu A^\mu + \xi_y \partial_y A^4]^2.$$
(4)

The above gauge-fixing action is somewhat very special and at the same time very crucial for this nmUED scenario. Due to the presence of the BLKTs in the Lagrangian lead to a non-homogeneous weight function for the fields with respect to the extra dimension. This inhomogeneity forces us to choose a y-dependent gauge-fixing parameter ξ_y as [32],

$$\xi = \xi_y \left(1 + r_H \{ \delta(y) + \delta(y - \pi R) \} \right), \tag{5}$$

¹From now and rest of our analysis we use $\partial_4(D_4) \equiv \partial_y(D_y)$.

here ξ is independent of y. This relation can be seen as *renormalisation* of the gauge-fixing parameter as the BLKTs are in some sense counter terms taking into account the unknown ultraviolet contribution in loop calculations. In this sense, ξ_y is the bare gauge-fixing parameter while ξ can be viewed as the renormalized gauge-fixing parameter taking the values 0 (Landau gauge), 1 (Feynman gauge) or ∞ (Unitary gauge).

Appropriate KK-expansion of the fields which are involved in the above actions can be schematically written as:

$$V_{\mu}(x,y) = \sum_{n} V_{\mu}^{(n)}(x)a^{n}(y)$$
(6)

$$V_4(x,y) = \sum_n V_4^{(n)}(x)b^n(y),$$
(7)

and

$$H(x,y) = \sum_{n} H^{(n)}(x) f^{n}(y).$$
 (8)

Variation of the action functional equations (Eq.1 and 2) and utilizing Eq.6, 7 and 8 we obtain equation for the y-dependent wave functions of V_{μ} as follows [18, 19]:

$$\left[\partial_y^2 + m_{V^n}^2 + \{\delta(y) + \delta(y - \pi R)\}\{r_V m_{V^n}^2 - (r_H - r_V) M_V^2\}\right] a^n(y) = 0.$$
(9)

The above equation can be used for both W and Z bosons. M_V is the electroweak symmetry breaking mass. For, $f^n(y)$ and $b^n(y)$ we have [16, 19]:

$$\left[\partial_y^2 + m_{H^n}^2 (1 + r_H \{\delta(y) + \delta(y - \pi R)\})\right] F^n(y) = 0,$$
(10)

where $F^n(y) \equiv f^n(y)$ or $b^n(y)$. Here, $m_{V^n}(m_{H^n})$ are the KK-mass solutions of gauge (Higgs) fields obtained from transcendental equation which we will discuss in the following.

For the sake of completeness in Eq.9 we have shown the equation of motion for $a^n(y)$. However, the Eq.9 or its solution will not play any major role in our analysis. One can see that in case of $r_V = r_H$, Eq.9 has the identical form as that of Eq.10 and they have the same solutions. In the following we will show without loss of any generality that this *equality* condition helps us to fix the gauge properly in this nmUED scenario.

We must keep in mind that the zero mode fields must correspond to respective SM particles. However, some of the fields in 5-D theory do not have any SM counterpart. Consequently application of appropriate boundary conditions, while solving the differential equation (Eq.10), is very crucial. As for example, $V_{\mu}(x, y)$ and H(x, y) should have zero modes, while $V_4(x, y)$ cannot have a zero mode.

Using the proper boundary conditions [16, 19], solutions for $f^n(y)$ can be written in a compact form[18, 25, 26] as $N_n \cos[m_{H^n}(y - \frac{\pi R}{2})]$ for even (including the zero mode) KK-modes and $N_n \sin[m_{H^n}(y - \frac{\pi R}{2})]$ for odd KK-modes. Here N's are the normalisation constant. Note that these functions have a definite parity of $(-1)^n$ (called KK-parity) under $y \leftrightarrow y - \pi R$ transformation².

The eigenvalues m_{H^n} 's are the solutions of the transcendental equation [16, 22, 25, 26]:

$$\frac{r_H m_{H^n}}{2} = \begin{cases} -\tan\left(\frac{m_{H^n} \pi R}{2}\right) & \text{for } n \text{ even,} \\ \cot\left(\frac{m_{H^n} \pi R}{2}\right) & \text{for } n \text{ odd.} \end{cases}$$
(11)

It should be noted that zero mode of V_4 is not desirable, one can easily write the solutions of y-dependent part of $V_4(x, y)$ i.e., $b^n(y)$ as $N_n \sin[m_{H^n}(y - \frac{\pi R}{2})]$ and $N_n \cos[m_{H^n}(y - \frac{\pi R}{2})]$ for even and odd KK-modes respectively.

From the above discussions it is evident that y-dependent wave functions $b^n(y)$ and $f^n(y)$ are transformed in opposite way under Z_2 parity, consequently $b^n(y)$ and $f^n(y)$ should satisfy the relation $\partial_y f^n(y) = m_{H^n} b^n(y)$. In fact, f^n 's are so called *primitive* of b^n 's [32]. In general a function $\tilde{b}^n(y)$ would be called the primitive of $b^n(y)$, if

$$\partial_y \tilde{b}^n(y) = m_{H^n} b^n(y). \tag{12}$$

Thus, $b^n(y)$ and $\tilde{b}^n(y)$ obey the following condition [32]:

$$\frac{m_{H^n}\partial_y b^n(y)}{[1+r_H\{\delta(y)+\delta(y-\pi R)\}]} = -m_{H^n}^2 \tilde{b}^n(y),$$
(13)

or equivalently we can write it as (using Eq.12),

$$[\partial_y^2 + m_{H^n}^2 (1 + r_H \{\delta(y) + \delta(y - \pi R)\})]\tilde{b}^n(y) = 0.$$
(14)

This equation (equivalent to Eq.10) immediately gives the following orthogonality relations [19]:

$$\int_{0}^{\pi R} dy \, \left[1 + r_f \{\delta(y) + \delta(y - \pi R)\}\right] \tilde{b}^n(y) \tilde{b}^m(y) = \delta^{nm},\tag{15}$$

$$\int_{0}^{\pi R} dy \, \partial_y \tilde{b}^n(y) \partial_y \tilde{b}^m(y) = m_{H^n}^2 \delta^{nm}.$$
(16)

In principle, $a^n(y)$ could also be a primitive of $b^n(y)$ when r_H would equal to r_V . So in our analysis $\tilde{b}^n \equiv f^n$, a^n .

Now let us come to the main point of our interest. What are the composition and masses of any physical scalar mode that emerges after the symmetry breaking? To find this out one has to look into the mass-mixing matrix constructed out of the $V_4^{(n)}$ ($\equiv W_4^{+(n)}$ or $Z_4^{(n)}$) and the $\eta^{(n)}$ ($\equiv \phi^{+(n)}$ or $\chi^{(n)}$). All the elements of this matrix are obtained from Eq.2 and Eq.4. In Eq.2 we have a term like $(D_y H)^{\dagger} (D^y H)$ from which the relevant terms are as follows $\frac{1}{2}M_V^2 V_4 V^4$, $\frac{1}{2}(\partial_y \eta)(\partial^y \eta)$ and $M_V(\partial_y \eta)V^4$. Using the KK-decompositions (Eq.7 and Eq.8),

²This is equivalent to a reflection symmetry along a line $y = \frac{\pi R}{2}$ in the y-space.

primitive condition (Eq.12) and orthogonality relation (Eq.16), we can express the first two terms as $-\frac{1}{2}M_V^2 V_4^{(n)} V_4^{(n)}$, $-\frac{1}{2}m_{H^n}^2 \eta^{(n)} \eta^{(n)}$ and these two terms contribute to diagonal elements of the mass-mixing matrix. Again using Eq.8, Eq.12 and Eq.16 we can express the remaining term³ as $-M_V m_{H^n} \eta^{(n)} V_4^{(n)}$ which contributes to off-diagonal elements. The complementary parts of the matrix elements (for both diagonal and off-diagonal) are generated from the Eq.4 (gauge-fixing action) and the relevant terms (written in terms of generic notation V and η) are given by $-\frac{\xi(\partial_y V^4)^2}{2(1+r_H\{\delta(y)+\delta(y-\pi R)\})}$, $-\frac{\xi}{2}M_V^2(1+r_H\{\delta(y)+\delta(y-\pi R)\})\eta^2$ and $\xi M_V(\partial_y V^4)\eta$. Then again using KK-decompositions (Eq.7 and Eq.8), Eq.13 and Eq.15 we can express these terms as $-\frac{\xi}{2}m_{H^n}^2 V_4^{(n)} V_4^{(n)}$, $-\frac{\xi}{2}M_V^2 \eta^{(n)} \eta^{(n)}$ and $\xi M_V m_{H^n} \eta^{(n)} V_4^{(n)}$ respectively. Note that, the construction of mass-mixing matrix elements of the above forms are possible only in the limit $r_V = r_H$, as otherwise we can not identify $\tilde{b}^n \equiv f^n$. Using above expressions we can write the mass-mixing matrix in $V_4^{(n)}(x) - \eta^{(n)}(x)$ basis in R_ξ gauge as:

$$M_{\xi}^{n} = \begin{pmatrix} M_{V}^{2} + \xi m_{H^{n}}^{2} & (1 - \xi) M_{V} m_{H^{n}} \\ (1 - \xi) M_{V} m_{H^{n}} & \xi M_{V}^{2} + m_{H^{n}}^{2} \end{pmatrix}.$$
 (17)

Let us now pay some attention to the eigenvalues of this matrix M_{ξ}^n . A straight forward diagonalisation would results into two eigenvalues $\lambda_1 = (M_V^2 + m_{H^n}^2)$ and $\lambda_2 = \xi (M_V^2 + m_{H^n}^2)$ and corresponding eigenstates are given by:

$$\mathcal{H}^{(n)} = \frac{\left(m_{H^n} \eta^{(n)} - M_V V_4^{(n)}\right)}{M_{V^{(n)}}},\tag{18}$$

$$\mathcal{G}^{(n)} = \frac{\left(M_V \eta^{(n)} + m_{H^n} V_4^{(n)}\right)}{M_{V^{(n)}}},\tag{19}$$

where, $M_{V^{(n)}} = \sqrt{M_V^2 + m_{H^n}^2}$. It is thus very simple to identify the state with mass(-square) λ_1 with the physical scalar (as this is independent of gauge choice) and the remaining one with the Goldstone. Note that, in the Feynman gauge (with $\xi = 1$) these two states are mass degenerate. Unitary gauge is equivalent to choosing $\xi \to \infty$, when this second state becomes infinitely heavy and decouples from the theory. And, finally in the Landau gauge (with $\xi = 0$), the ξ dependent eigenvalue becomes zero, resulting in vanishing determinant of the mass-matrix and a massless Goldstone mode characteristic of this gauge. The mixing angle (θ) is independent of the gauge choice as, $\tan 2\theta = \frac{2M_V m_{H^n}}{m_{H^n}^2 - M_V^2}$. The orthogonal combination to the Goldstone, with eigenvalue ($M_A^2 + m_{H^n}^2$) is a physical scalar which remains in the theory. Mass of this physical mode is independent of the gauge choice. To this end let us remind that, though the structure of the orthogonal combination of the physical scalar and Goldstone are identical with UED scenario, but the extra dimensional contributions are obtained from the solution of transcendental equation (Eq.11). Furthermore, without the equality condition $r_V = r_H$, such simplified version (identical with UED scenario) of gauge-fixing procedure would not be possible in muUED scenario.

³As we have already mentioned that we generically use η instead of ϕ^+ and χ , so for η we can use the KK-decomposition given in Eq.8.

In literature [16, 19, 23, 24], taking $V_4 \rightarrow 0$ is synonymous with the Unitary gauge. Here, we would like to shed light on this issue in context of our earlier discussion. Extremising the gauge fixed action of V_M (Eq.1 and Eq.4) would lead to an equation of motion for $V_\mu(x, y)$ (under the assumption of $r_V = r_H$):

$$\left[\left[\partial^{\mu} \partial_{\mu} g^{\mu\nu} + M_V^2 g^{\mu\nu} - \left(1 - \frac{1}{\xi}\right) \partial^{\nu} \partial^{\mu} \right] \left[1 + r_V \{ \delta(y) + \delta(y - \pi R) \} \right] - \partial_y^2 g^{\mu\nu} \right] V_{\mu}(x, y) = 0.$$
 (20)

Using $\xi \to \infty$ in the above equation would lead us to the equation of motion for V_{μ} in the Unitary gauge. However one can also achieve this without gauge-fixing and simply by setting $V_4 \rightarrow 0$ in Eq.1 followed by extermisation of the action. In fact, setting V_4 to zero in Eq.1 would straightforwardly remove the undesirable terms in which V_{μ} couples to V_4 via derivative, which was the main purpose of gauge-fixing. And, as long as we are interested in V_{μ} and its interactions with other physical particles, taking $V_4 \rightarrow 0$ is as good as Unitary gauge⁴. However we must keep in mind that the *Goldstone* is a linear combination of $V_4^{(n)}$ and $\eta^{(n)}$ (see Eq.19). And the coefficients of this linear combination is independent of choice of gauge. Whenever, one needs to have the interactions involving physical scalar (like Charged Higgs) and/or Goldstones (in some gauge other than the Unitary gauge), one should be more careful and should use the proper definitions of these scalars. For example, the Refs. [25, 26, 27] deal with the electroweak radiative corrections in nmUED scenario and the entire calculations have been performed in Feynman gauge where Goldstone bosons appear in the loop diagrams. The prescription of gauge-fixing mechanism in the above Refs. [25, 26, 27] has been adopted from the formalism presented in this letter. A more careful look at the expression for the mixing angle in the physical scalar/Goldstone sector would reveal that in the large R^{-1} limit $(R^{-1} >> v(\text{SM VEV}))$, physical scalar can be identified with η (see Eq.18) while V_4 becomes the Goldstone (see Eq.19).

In summary, we have discussed, the gauge-fixing procedure in a model of Universal Extra Dimensions with boundary localised kinetic terms where a gauge symmetry is spontaneously broken by Higgs mechanism. Finally we have discussed the composition and masses of the Goldstone and physical scalar in this model.

Acknowledgements AD is partially supported by Department of Atomic Energy-Board of Research in Nuclear Sciences (DAE-BRNS) research project. AS acknowledges financial support Department of Atomic Energy, Government of India, for the Regional Centre for Acceleratorbased Particle Physics (RECAPP), Harish-Chandra Research Institute. Authors are grateful to Amitava Raychaudhuri for carefully reading the manuscript and several illuminating discussions.

 $^{^{4}}$ An equivalent statement can also be found in the Ref [33] in which all fermions are localised on the 4-D brane.

References

- [1] G. Servant and T. M. P. Tait, Nucl. Phys. B 650 (2003) 391, [arXiv:hep-ph/0206071].
- [2] D. Majumdar, Mod. Phys. Lett. A 18 (2003) 1705.
- [3] K. Kong and K. T. Matchev, JHEP 0601 (2006) 038 [arXiv:hep-ph/0509119].
- [4] F. Burnell and G. D. Kribs, Phys. Rev. D 73 (2006) 015001, [arXiv:hep-ph/0509118].
- [5] G. Belanger, M. Kakizaki and A. Pukhov, JCAP **1102** (2011) 009, [arXiv:1012.2577 [hep-ph]].
- [6] K. R. Dienes, E. Dudas and T. Gherghetta, Phys. Lett. B 436 (1998) 55, [arXiv:hep-ph/9803466].
- [7] K. Dienes, E. Dudas, and T. Gherghetta, Nucl. Phys. B 537 (1999) 47, [arXiv:hep-ph/9806292].
- [8] G. Bhattacharyya, A. Datta, S. K. Majee, A. Raychaudhuri, Nucl.Phys. B 760 (2007) 117, [arXiv:hep-ph/0608208].
- [9] T. Appelquist, H. C. Cheng and B. A. Dobrescu, Phys. Rev. D 64 (2001) 035002, [arXiv:hep-ph/0012100].
- [10] H. Georgi, A. K. Grant and G. Hailu, Phys. Lett. B 506 (2001) 207, [arXiv:hep-ph/0012379].
- [11] H. C. Cheng, K. T. Matchev and M. Schmaltz, Phys. Rev. D 66 (2002) 036005, [arXiv:hep-ph/0204342].
- [12] G. R. Dvali, G. Gabadadze, M. Kolanovic and F. Nitti, Phys. Rev. D 64 (2001) 084004, [arXiv:hep-ph/0102216].
- [13] F. del Aguila, M. Perez-Victoria and J. Santiago, JHEP 0302 (2003) 051, [arXiv:hep-th/0302023].
- [14] F. del Aguila, M. Perez-Victoria and J. Santiago, Acta Phys. Polon. B 34 (2003) 5511, [arXiv:hep-ph/0310353].
- [15] T. Flacke, K. Kong and S.C. Park, JHEP **05** (2013) 111 [arXiv:1303.0872 [hep-ph]].
- [16] M. S. Carena, T. M. P. Tait and C. E. M. Wagner, Acta Phys. Polon. B 33 (2002) 2355 [arXiv:hep-ph/0207056].
- [17] C. Schwinn, Phys. Rev. D 69 (2004) 116005, [hep-ph/0402118].
- [18] T. Flacke, A. Menon and D. J. Phalen, Phys. Rev. D 79 (2009) 056009 [arXiv:0811.1598 [hep-ph]].

- [19] A. Datta, U. K. Dey, A. Shaw, and A. Raychaudhuri, Phys. Rev. D 87 (2013) 076002 [arXiv:1205.4334 [hep-ph]].
- [20] A. Datta, K. Nishiwaki and S. Niyogi, JHEP **1211** (2012) 154, [arXiv:1206.3987 [hep-ph]].
- [21] A. Datta, K. Nishiwaki and S. Niyogi, JHEP **1401** (2014) 104, [arXiv:1310.6994 [hep-ph]].
- [22] A. Datta, U. K. Dey, A. Raychaudhuri, and A. Shaw, Phys. Rev. D 88 (2013) 016011, [arXiv:1305.4507 [hep-ph]].
- [23] A. Datta, A. Raychaudhuri, and A. Shaw, Phys. Lett. B 730 (2014) 42, [arXiv:1310.2021 [hep-ph]].
- [24] A. Shaw, Eur. Phys. J. C **75** (2015) 33, [arXiv:1405.3139 [hep-ph]].
- [25] T. Jha and A. Datta, JHEP **1503** (2015) 012, [arXiv:1410.5098 [hep-ph]].
- [26] A. Datta and A. Shaw, Phys. Rev. D 93 (2016) 055048, [arXiv:1506.08024 [hep-ph]].
- [27] U. K. Dey and T. Jha, [arXiv:1602.03286 [hep-ph]].
- [28] J. Papavassiliou and A. Santamaria, Phys. Rev. D 63 (2001) 125014, [arXiv:hep-ph/0102019].
- [29] F. J. Petriello, JHEP **0205** (2002) 003, [arXiv:hep-ph/0204067].
- [30] A. J. Buras, M. Spranger and A. Weiler, Nucl. Phys. B 660 (2003) 225, [arXiv:hep-ph/0212143].
- [31] C. Csaki, J. Hubisz and P. Meade [arXiv:hep-ph/0510275].
- [32] A. Mück, Ph. D dissertation, University of Würzburg, 2004.[http://www.physik.uni-wuerzburg.de/fileadmin/11030200/Dissertationen/mueck-diss.pdf]
- [33] A. Muck, A. Pilaftsis and R. Ruckl, Phys. Rev. D **65** (2002) 085037, [arXiv:hep-ph/0110391].
- [34] A. Muck, A. Pilaftsis and R. Ruckl, [arXiv:hep-ph/0203032].