Supporting Online Material to the manuscript: "Water isotope diffusion rates from the NorthGRIP ice core for the last 16,000 years - glaciological and paleoclimatic implications."

V. Gkinis^{1,2}, S. B. Simonsen^{1,3}, S. L. Buchardt¹, J. W. C. White², and B. M. Vinther¹

¹Centre for Ice and Climate, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark

²Institute for Alpine and Arctic Research, University of Colorado, Boulder, 1560 30th Street Boulder, CO 80303 USA

³Div. of Geodynamics, DTU space National Space Institute, Elektrovej, Build. 327, Kgs. Lyngby, Denmark

I. Derivation of the integration equations for σ^2

In this section we show how we derive the numerical expressions for the diffusion length (Eq. (8) in main text (MT)). Starting from the differential equation for the diffusion length we have:

$$\frac{\mathrm{d}\sigma^2}{\mathrm{d}t} - 2\,\dot{\varepsilon}_z(t)\,\sigma^2 = 2D(t)\,. \tag{1}$$

We consider the vertical strain rate due to densification:

$$\dot{\varepsilon}_z(t) = \frac{-\partial \rho}{\partial t} \frac{1}{\rho} \tag{2}$$

By substituting t with ρ and combining Eq.(1), (2) we get:

$$\frac{\mathrm{d}\sigma^2}{\mathrm{d}\rho} + \frac{2\sigma^2}{\rho} = 2\left(\frac{\mathrm{d}\rho}{\mathrm{d}t}\right)^{-1}D(\rho) \tag{3}$$

Multiplication of both sides of Eq.3 with the integrating factor

$$F(\rho) = e^{\int \frac{2}{\rho} d\rho} = \rho^2, \tag{4}$$

gives:

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\rho^2 \sigma^2 \right) = 2\rho^2 \left(\frac{\mathrm{d}\rho}{\mathrm{d}t} \right)^{-1} D(\rho), \tag{5}$$

from which we get the result:

$$\sigma^{2}(\rho) = \frac{1}{\rho^{2}} \int_{\rho_{o}}^{\rho} 2\rho^{2} \left(\frac{\mathrm{d}\rho}{\mathrm{d}t}\right)^{-1} D(\rho) \,\mathrm{d}\rho. \tag{6}$$

In a similar way for the ice diffusion length (Eq. (12) in the MT) we have:

$$\frac{\mathrm{d}\sigma^2}{\mathrm{d}t} - 2\,\dot{\varepsilon}_z(t)\,\sigma^2 = 2D(t)\,,\tag{7}$$

where the total thinning is given by:

$$S(t') = e^{\int_0^{t'} \dot{\varepsilon}_z(t) dt} . \tag{8}$$

We multiply both sides of Eq. (7) with the integrating factor

$$F(\rho) = e^{\int_0^{t'} -2\dot{\varepsilon}_z(t)dt},\tag{9}$$

which results in

$$\frac{\mathrm{d}}{\mathrm{d}t} \left[\sigma^2 e^{\int_0^{t'} -2\dot{\varepsilon}_z(t)\mathrm{d}t} \right] = 2D(t)e^{\int_0^{t'} -2\dot{\varepsilon}_z(t)\mathrm{d}t}, \quad (10)$$

From this, we get the expression for the ice diffusion length

$$\sigma_{ice}^2(t') = S(t')^2 \int_0^{t'} 2D_{ice}(t) S(t)^{-2} dt.$$
 (11)

II. THE DIFFUSIVITY PARAMETRIZATION

A. The firn diffusivity

We use the diffusivity parametrization as introduced by Johnsen et al. (2000).

$$D(\rho) = \frac{m \, p \, D_{ai}}{R \, T \, \alpha_i \, \tau} \left(\frac{1}{\rho} - \frac{1}{\rho_{ice}} \right). \tag{12}$$

The terms used in Eq. (12) and their parameterizations used are described below:

- m: molar weight (kg)
- p: saturation vapor pressure over ice (Pa). We use (Murphy and Koop, 2005):

$$p = \exp\left(9.5504 - \frac{5723.265}{T} + 3.530 \ln(T) - 0.0073 T\right).$$
(13)

• D_a : diffusivity of water vapor in air (m²s⁻¹). We use (Hall and Pruppacher, 1976):

$$D_a = 2.1 \cdot 10^{-5} \left(\frac{T}{T_o}\right)^{1.94} \left(\frac{P_o}{P}\right) \tag{14}$$

with $P_o=1$ Atm, $T_o=273.15$ K and P,T the ambient pressure (Atm) and temperature (K). Additionally from Merlivat (1978) $D_{a^2\mathrm{H}}=\frac{D_a}{1.0251}$ and $D_{a^{18}\mathrm{O}}=\frac{D_a}{1.0285}$.

- R: molar gas constant R = $8.314478 \left(\text{m}^3\text{Pa}\left(\text{K mol}\right)^{-1}\right)$
- T: Ambient temperature (K)

• α_i : Ice – Vapor fractionation factor, we use the formulations by Majoube (1971) and Merlivat and Nief (1967) for $\alpha_{s/v}^2$ and $\alpha_{s/v}^{18}$ respectively.

$$\ln \alpha_{Ice/Vapor} (^{2}\text{H}/^{1}\text{H}) = 16288/T^{2} - 9.34 \times 10^{-2}$$

$$(15)$$

$$\ln \alpha_{Ice/Vapor} (^{18}\text{O}/^{16}\text{O}) = 11.839/T - 28.224 \times 10^{-3}$$

$$(16)$$

• τ : The firn tortuosity. We use (Schwander et al., 1988; Johnsen et al., 2000):

$$\frac{1}{\tau} = \begin{cases} 1 - b_{\tau} \left(\frac{\rho}{\rho_{ice}}\right)^2 & \text{, for } \rho \le \frac{\rho_{ice}}{\sqrt{b}} \\ 0 & \text{, for } \rho > \frac{\rho_{ice}}{\sqrt{b}} \end{cases} , \quad (17)$$

where $b_{\tau} = 1.30$, implying a close-off density of $\rho_{\rm co} = 804.3~{\rm kgm}^{-3}$.

B. The ice diffusivity

Ice diffusion is believed to occur via a vacancy mechanism with transport of molecules within the ice lattice. Based on isotopic probe experiments, there is a strong consensus that the ice diffusivity coefficient is the same for H₂ ¹⁸O, D₂O and T₂O (Ramseier, 1967; Blicks et al., 1966; Itagaki, 1967; Delibaltas et al., 1966) The dependence of the ice diffusivity parameter to temperature is described by an Arrhenius type equation

$$D = D_0 \exp\left(-Q/RT\right),\tag{18}$$

where Q is the activation energy and D_0 a pre-exponential factor. The results of the studies mentioned above agree well with each other. Here we plot the diffusivity parametrization coefficients suggested by those studies (Fig. 1). In this work we follow Ramseier (1967) and use $Q=0.62~\rm eV$ and $D_o=9.2\cdot 10^{-4}~\rm m^2~s^{-1}$. Note that the results of Ramseier (1967) are based on measurements of both artificially as well as naturally grown ice collected at Mendenhall glacier, Alaska.

Enhanced ice diffusion rates have been proposed to be the cause of the excess diffusion observed in the Holocene section of the GRIP ice core (Johnsen et al., 2000). For the early Holocene part of the GRIP core, the authors of that study observed higher diffusion rates than expected by the theory. In order to diminish the discrepancy between modeled and observed diffusion rates Johnsen et al. (2000) introduced the term "excess ice diffusion", referring to a possibly higher diffusivity coefficient due to isotopic exchange in the liquid phase on thin water films and ice crystal veins. However, the thickness of the water films and the diameter of the ice crystal veins required, are unrealistically high.

Although the existence of an "excess ice diffusion" mechanism cannot be excluded based on the findings of our study, it should be mentioned that the diffusion model used in Johnsen et al. (2000) assumes an accumulation rate and temperature signal that is based on the δ^{18} O record of the GRIP core. The GRIP δ^{18} O signal is characterized by a rather flat curve throughout the Holocene showing no indication of an early Holocene optimum, a feature that is mostly due to ice sheet elevation effects "masking" the temperature change

(Vinther et al., 2009). As a result, it is expected that the diffusion length calculations in Johnsen et al. (2000) would underestimate the diffusion signal throughout the Holocene. We conclude that the "excess ice diffusion" issue requires more work in the future. Considering that the ice diffusivity coefficient is very similar for all the isotopologues of water, a study focusing on the differential diffusion signal between $\delta^{18}O$ and δD would provide a better insight in the problem. An effort will be undertaken as soon as an adequately long Holocene section of the NEEM ice core is analyzed for both $\delta^{18}O$ and δD using dual isotope laser spectroscopy.

III. EXAMPLES OF DIFFUSION LENGTHS FOR DIFFERENT ICE CORE SITES

In this section we present an ensemble of implementations of the diffusion–densification model for various combinations of surface forcings that represent typical modern day conditions for a number of ice core sites on Greenland and Antarctica. The contours in the plot are generated by integration of Eq. (6) and expressed in m ice eq. The forcing for each ice core site is given in Table III and the results are shown in Fig. 2

IV. Estimation of σ^2 from the high resolution data set

In order to estimate the diffusion length value from high resolution water isotope data we minimize the 2-norm $\|P_s - \hat{P}_s\|$ where \hat{P}_s is an estimate of the power spectral density of a high resolution $\delta^{18}{\rm O}$ data section and P_s is a model description of the power spectral density.

 P_s is obtained by the use of the Burg's spectral estimation method. The method fits an autoregressive model of order μ (AR- μ) by minimizing the forward–backward prediction error filter (Hayes, 1996; Press et al., 2007; Andersen, 1974). For the theoretical model we have:

$$P_s = P_\sigma + |\hat{\eta}(k)|^2, \tag{19}$$

where $P_{\sigma} = P_0 e^{-k^2 \sigma_i^2}$ is the effect of the firn diffusion process with squared diffusion length σ^2 . Regarding the noise, we find red noise described by an AR-1 process with an autoregressive coefficient $q_1 = 0.15$ to provide a good description of the noise signal we observe. The spectrum of this signal is (Kay and Marple, 1981):

$$|\hat{\eta}(k)|^2 = \frac{\sigma_{\eta}^2 \Delta}{|1 + q_1 \exp(-ik\Delta)|^2},$$
 (20)

where σ_{η}^2 is the variance of the noise. The angular frequency $k=2\pi f$ is in the range $f\in\left[0,\frac{1}{2\Delta}\right]$ defined by the Nyquist frequency and thus the sampling resolution Δ . We vary the parameters σ^2 , P_0 , and σ_{η}^2 of the spectral model in order to minimize the misfit between P_s and \hat{P}_s in a least squares sense.

As it can be seen in Fig. 3, it is the characterization of the full spectrum that yields information on σ^2 . It is thus not necessary to specifically study the relative attenuation of individual spectral peaks as for example the annual signal. This approach allows for a study of the diffusion signal even after the spectral signature of the annual signal diminishes.

V. AR ORDER SELECTION

An interesting feature of the Burg estimation method is that the order μ of the AR filter affects the spectral resolution of \hat{P}_s (Hayes, 1996; Press et al., 2007). Low μ values result in smoother spectra with inferior spectral resolution, while higher order spectra show better performance in resolving neighboring spectral peaks. This can be seen in the spectral estimates presented in Fig.3 where we plot spectral estimates with $\mu=30$ and $\mu=40$. As described above, the goal of the σ^2 estimation is to characterize the overall shape of the spectrum. As a result, relatively low values of μ produce smooth spectra of relatively low spectral resolution and can be adequate for the purpose of our application.

We look into both the influence of the value of μ on the σ^2 estimate by performing 41 power spectrum estimates with $\mu \in [40,80]$. Possible interferences of spectral features due to longer scale climate variability that could have an effect on the estimation of σ^2 are also investigated with this test. In Fig. 4 we show the mean value of the 41 spectral estimates before and after strain correction. The standard deviation of the estimated σ for every depth is presented on the top subplot of the figure. It can be seen that on average the standard deviation is about 2 orders of magnitude lower than the absolute values of $\sqrt{\sigma^2}$ thus approximately in the 1% range. The low standard deviation of the 41 estimates suggests that a possible effect of spectral features due to low frequency climate variability is of second order. The same applies for the selection of the AR order μ , in Burg's spectral estimation.

VI. TEST WITH SYNTHETIC DATA

Additional to the MEM order selection sensitivity test shown in section V, we investigate the precision and accuracy of the σ^2 estimation using synthetic data. We perform two tests which we describe below.

A. Synthetic data test 1

For the first test we investigate the influence of short or long memory of the $\delta^{18}{\rm O}$ time series due to climate variability. We generate high resolution synthetic $\delta^{18}O$ data by assuming an AR-1 process with the AR-1 coefficient ϕ_1 of the process being equal to 0.2 and 0.9995. The process is applied on Gaussian noise with a mean of zero and a standard deviation of 10 %. The time series generated have a spacing of $\Delta x = 10^{-3} \,\mathrm{m}$ and a total length $L = 20 \,\mathrm{m}$. A ϕ_1 equal to 0.9995 with $\Delta x = 10^{-3} \,\mathrm{m}$ corresponds to a time constant for the memory of the process equal to $\tau_{0.9995} = -\Delta x / \ln \phi_1 = 2 \,\mathrm{m}$ (Percival, 1993). For the part of the record we study, this is equivalent to climate variability at decadal time scales. The high resolution AR-1 process is then convolved with the Gaussian filter of predefined variance, simulating the effect of firn diffusion. Measurement white noise is then added to the result of the convolution. We then sample the high resolution diffused time series at a resolution of $\Delta = 0.05 \,\mathrm{m}$ representing the width of a discrete ice core sample and perform the σ^2 estimation as described in section IV.

The test is run using 3 different values for the diffusion length; 0.05, 0.035 and 0.025 m. We perform the procedure 100 times for every diffusion length generating a new AR-1 process for every repetition. This results in a total of $2\times3\times100=600$ experiments. We compare the estimated values for σ^2 with the target values and calculate the mean and RMS value for the estimation. In Fig. 5 we show one example for each of the three sets of different diffusion length values used presenting the raw time series for both values of ϕ_1 and the respective power spectral densities. The results for the 6 sets of experiments are presented in Table I.

B. Synthetic data test 2

For the second test we follow a similar procedure as in test 1 generating an AR-1 process that we then diffuse using a fixed value for the diffusion length. We choose $\sigma=0.08\,\mathrm{m}$. The synthetic data sets are then sampled with 4 different resolutions with $\Delta=0.05,0.08,0.10\,\mathrm{and}\,0.12\,\mathrm{m}$ The diffusion length is estimated and corrected for discrete sampling as described in section 4 of the MT. From the spectral estimation point of view, the effect of the coarser sampling resolution is equivalent to the ice flow thinning. The lower Nyquist frequency caused by the coarser sampling scheme results in an inferior estimation of the noise signal (Fig. 6). The diffusion length estimates that we present in Table II indicate that the estimation scheme is accurate and insensitive to the memory of the AR-1 process as well as the sampling scheme.

An RMS value of 0.5 cm based on the synthetic data tests is taken into account when calculating the confidence intervals in Fig. 12. The equivalent temperature uncertainty of $\pm 0.5 \, \mathrm{cm}$ is approximately 1 K for the NorthGRIP site.

VII. ICE FLOW THINNING EFFECTS

The layer thinning induced by the ice flow, impacts the diffusion length estimation mainly in two ways. First, due to the discrete sampling scheme as the diffusion length estimation moves towards the deeper parts of the core, a single sample averages more years of climate information. This effect is essentially taken care of by means of the discrete sampling correction described in section 4 of the MT. The term σ_{dis}^2 is constant with depth. However based on Eq. (20) of the MT one can see that the effective correction for discrete sampling scales with the total thinning function S(z). In Fig. 9, we illustrate the effect of this correction with depth.

Second, the ice flow thinning will result in the diffusion length value decreasing with depth. With lower σ^2 values, a spectrum estimate up to the Nyquist frequency $1/2\Delta$ will contain a decreasing part of the noise signal $|\hat{\eta}(k)|^2$. After a certain depth, the sampling resolution is not high enough to resolve $|\hat{\eta}(k)|^2$. The result of this effect is that the estimation of the P_σ signal requires an assumption about $|\hat{\eta}(k)|^2$ and thus it can limit the extend to which the diffusion technique can be applied to the deeper parts of the core.

In Fig. 7 we plot the expected diffusion length value assuming a simple case of constant temperature and accumulation rate at the surface and a certain ice layer thinning

history. Then, based on this modeled diffusion length profile, in Fig. 8, we calculate six power spectral densities for $z = 200,600,900,1200,1400 \,\mathrm{and}\, 1600 \,\mathrm{m}$. For the spectral calculations we use 2 different sampling schemes, Δ_1 = $5\,\mathrm{cm}$ and $\Delta_2=2.5\,\mathrm{cm}$. The plots illustrate the effect of the ice layer thinning as well as the sampling resolution on the shape of the power spectral density. As depth increases, a progressively smaller portion of the noise signal is resolved. Conclusively, for the deeper parts of the core where the ice layer thinning has reduced the diffusion length, a higher sampling resolution ($\Delta < 2.5\,\mathrm{cm}$) is preferable for an even more accurate estimation of P_s and subsequently σ^2 to be possible. For the NorthGRIP reconstruction we present here, we are able resolve the noise signal down to the depth of approximately 1450 m. For depths higher than 1450 m we make the simplest possible assumption that the noise level is equal to the average values we have observed in the Holocene section.

The accuracy of the ice flow model in inferring the ice thinning function has an influence on the uncertainty of our temperature reconstruction. The value of the diffusion length of a layer at depth z, estimated from the spectral properties of a set of $\delta^{18}\mathrm{O}$ data needs to be corrected for ice flow thinning. An inaccurately estimated thinning function affects the inferred values of σ^2_{firm} in a linear way as we show in Eq. (13) and (20) of the MT. As far as the inferred temperatures are concerned, the ice thinning function impacts the slope of the signal, but has no influence on its variability.

In Fig. 10 we performed the temperature calculation using six different scenarios for the ice thinning function S(z). We assume the simple scenario of a thinning function that varies linearly with depth and a value of S(z=2100 m) equal to 0.22, 0.24, 0.26, 0.28, 0.30, 0.32. It is apparent that the change in temperature due to thinning function uncertainties can be significant. However this type of uncertainty only affects the slope of the temperature signal. So, unrealistic thinning function scenarios are relatively straightforward to rule out. Estimates of temperature from other proxies for any point of the record, can be useful in order to select a plausible scenario for the thinning function. This allows for a more accurate determination of slope of the temperature signal inferred by means of the firn diffusion method.

A direct consequence of this is that the method can potentially be useful in providing combined paleotemperature and glaciological information. In this study the unrealistically high temperature values we inferred for the Holocene climatic optimum pointed to possible inaccuracies of the ice thinning function used for the estimation. When fixing the temperature gradient between the Holocene optimum and present conditions to be approximately 3 K, we were able to propose a more likely scenario for the ice thinning function and hence the accumulation rate history. The temperature reconstruction using the proposed ice thinning function for NorthGRIP is presented in red color in Fig. 10.

VIII. FIRN DENSIFICATION UNCERTAINTIES

Uncertainties related to the densification model affect the estimation of the diffusion length $\sigma_{\rm firm}^2$. Hereby we examine

the influence of four parameters involved in the densification—diffusion model. We run a set of sensitivity experiments where the four firn densification parameters are perturbed in order to create a family of 1000 implementations of the diffusion-densification model for each experiment. For all the following sensitivity tests we also consider the standard deviation of the diffusion length spectral estimate as calculated in sections V and VI.

The first two parameters we consider are the surface and close-off densities ρ_0 and ρ_{co} . We perform two sensitivity experiments where the values of ρ_0 and ρ_{co} are drawn from a Gaussian distribution with a defined mean and standard deviation (Table IV). For the close-off density a value of $804~\pm20~{\rm kgm^{-3}}~(1\sigma)$ is used (Schwander et al., 1988; Jean-Baptiste et al., 1998; Johnsen et al., 2000). The range of values we choose for ρ_{co} brackets within 2σ the more extreme estimates of 775 and 840 kgm⁻³ shown in Scher and Zallen (1970) and Stauffer et al. (1985) respectively. For the surface density we use a value of $320 \pm 40 \text{ kgm}^{-3}$ (1σ) , based on modern observations of the firn column density at NothGRIP. Previous high resolution density observations by Albert and Shultz (2002) for Summit, Greenland indicate that the surface density can vary within $\pm 50 \text{kgm}^{-3}$ of its mean value. As a result, with a 1σ of 40 kgm^{-3} the Gaussian distribution of ρ_0 covers this range adequately in our sensitivity experiments.

Additionally, we include two parameters that describe the dependence of the densification rate to temperature. Based on Herron and Langway (1980)

$$\frac{\mathrm{d}\rho(z)}{\mathrm{d}t} = K(T)A^{\vartheta}\left(\rho_{\mathrm{ice}} - \rho\left(z\right)\right),\tag{21}$$

where K(T) is a temperature dependent Arrhenius-type densification rate coefficient described by:

$$K(T) = 11 \exp\left(-\frac{10160}{RT}\right) \quad \rho < 550 \text{ kgm}^{-3},$$
 (22)

and

$$K(T) = 575 \exp\left(-\frac{21400}{RT}\right) \quad \rho \ge 550 \text{ kgm}^{-3}.$$
 (23)

In order to perturb the model we use the term K'(t) in Eq. (21) where K'(T) = fK(T) and $f = 1 \pm 0.2$ (1σ). This results in a family of density profiles that are used for the diffusion length calculation. In Fig. 11 1σ and 2σ intervals are illustrated together with firn density measurements from NorthGRIP.

The results of these sensitivity experiments are illustrated in Fig.12. Based on these results we conclude that using a fixed value for the surface and close–off densities is a plausible approach. The combined uncertainty of the ρ_o and ρ_{co} parameters is in the order of 1 K and thus the temperature history we infer is consistent over a wide range of densification parameter values. Combining all densification parameters the uncertainty of the estimation is equal to $\pm 2.5 \, \mathrm{K} \, (1\sigma)$. Combining this in a Gaussian sense with the spectral estimation uncertainty from section VI we get a total of $\pm 2.7 \, \mathrm{K} \, (1\sigma)$.

REFERENCES

- Albert, M. R., Shultz, E. F., May 2002. Snow and firn properties and air-snow transport processes at summit, greenland. Atmospheric Environment 36 (15-16), 2789–2797.
- Andersen, N., 1974. Calculation of filter coefficients for maximum entropy spectral analysis. Geophysics 39 (1), 69–72.
- Blicks, H., Dengel, O., Riehl, N., 1966. Diffusion von protonen (tritonen) in reinen und dotierten eis-einkristallen. Physik Der Kondensiterten Materie 4 (5), 375–381.
- Dahl Jensen, D., Mosegaard, K., Gundestrup, N., Clow, G. D., Johnsen, S. J., Hansen, A. W., Balling, N., Oct. 1998. Past temperatures directly from the Greenland ice sheet. Science 282 (5387), 268–271.
- Delibaltas, P., Dengel, O., Helmreich, D., Riehl, N., Simon, H., 1966. Diffusion von ¹⁸O in eis-einkristallen. Physik Der Kondensiterten Materie 5 (3), 166–170.
- Hall, W. D., Pruppacher, H. R., 1976. Survival of ice particles falling from cirrus clouds in subsaturated air. Journal of the Atmospheric Sciences 33 (10), 1995–2006.
- Hayes, M. H., 1996. Statistical digital signal processing and modeling. John Wiley & Sons.
- Herron, M. M., Langway, C., 1980. Firn densification an empirical-model. Journal Of Glaciology 25 (93), 373–385.
- Itagaki, K., Feb. 1967. Self-diffusion in single crystal ice. J. Phys. Soc. Jpn. 22 (2), 427–431.
- Jean-Baptiste, P., Jouzel, J., Stievenard, M., Ciais, P., May 1998. Experimental determination of the diffusion rate of deuterated water vapor in ice and application to the stable isotopes smoothing of ice cores. Earth And Planetary Science Letters 158 (1-2), 81–90.
- Johnsen, S. J., Clausen, H. B., Cuffey, K. M., Hoffmann, G., Schwander, J., Creyts, T., 2000. Diffusion of stable isotopes in polar firn and ice. the isotope effect in firn diffusion. In: Hondoh, T. (Ed.), Physics of Ice Core Records. Hokkaido University Press, Sapporo, pp. 121–140.
- Johnsen, S. J., Dahl Jensen, D., Dansgaard, W., Gundestrup, N., Nov. 1995. Greenland paleotemperatures derived from GRIP bore hole temperature and ice core isotope profiles. Tellus Series B-Chemical And Physical Meteorology 47 (5), 624–629.
- Johnsen, S. J., DahlJensen, D., Gundestrup, N., Steffensen, J. P., Clausen, H. B., Miller, H., Masson-Delmotte, V., Sveinbjornsdottir, A. E., White, J., 2001. Oxygen isotope and palaeotemperature records from six greenland ice-core stations: Camp Century, Dye-3, GRIP, GISP2, Renland and NorthGRIP. Journal Of Quaternary Science 16 (4), 299–307.
- Kay, S. M., Marple, S. L., 1981. Spectrum analysis a modern perspective. Proceedings Of The Ieee 69 (11), 1380–1419.
- Majoube, M., 1971. Fractionation in o-18 between ice and water vapor. Journal De Chimie Physique Et De Physico-Chimie Biologique 68 (4), 625–&.
- Merlivat, L., 1978. Molecular diffusivities of $\mathrm{H_2^{16}O}$, $\mathrm{HD^{16}O}$, and $\mathrm{H_2^{18}O}$ gases. Journal of Chemical Physics 69 (6), 2864–2871.
- Merlivat, L., Nief, G., 1967. Fractionnement isotopique lors des changements detat solide-vapeur et liquide-vapeur de leau a des temperatures inferieures a 0 degrees c. Tellus

- 19 (1), 122–127.
- Murphy, D. M., Koop, T., Apr. 2005. Review of the vapour pressures of ice and supercooled water for atmospheric applications. Quarterly Journal Of The Royal Meteorological Society 131 (608), 1539–1565.
- Percival, D. B., 1993. Spectral analysis for physical applications. Cambridge University Press.
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., Flannery, B. P., August 2007. Numerical Recipes: The Art of Scientific Computing. Cambridge University Press.
- Ramseier, R. O., 1967. Self-diffusion of tritium in natural and synthetic ice monocrystals. Journal Of Applied Physics 38 (6), 2553–2556.
- Scher, H., Zallen, R., 1970. Critical density in percolation processes. Journal of Chemical Physics 53 (9), 3759–3761.
- Schwander, J., Stauffer, B., Sigg, A., 1988. Air mixing in firm and the age of air at pore close-off. Annals Of Glaciology 10, 141–145.
- Stauffer, B., Schwander, J., H., O., 1985. Enclosure of air during metamorphosis of dry firn to ice. Annals of Glaciology 6, 108–112.
- Vinther, B. M., Buchardt, S. L., Clausen, H. B., Dahl-Jensen,
 D., Johnsen, S. J., Fisher, D. A., Koerner, R. M., Raynaud,
 D., Lipenkov, V., Andersen, K. K., Blunier, T., Rasmussen,
 S. O., Steffensen, J. P., Svensson, A. M., 2009. Holocene thinning of the greenland ice sheet. Nature 461 (7262), 385–388.

$\sigma [{ m cm}]$	5	3.5	2.5
$\phi_1 = 0.2$	4.99 ± 0.20	3.47 ± 0.19	2.30 ± 0.25
$\phi_1 = 0.9995$	5.06 ± 0.15	3.50 ± 0.13	2.56 ± 0.39

TABLE I: Summary of the synthetic data test 1. The mean and RMS value of the diffusion length estimation is given in cm for 3 different target values of the diffusion length. Statistics are based on 100 realizations for each experiment.

	Sampling Interval [cm]			
	5	8	10	12
$\phi_1 = 0.2$	8.1 ± 0.3	8.0 ± 0.4	8.1 ± 0.3	8.2 ± 0.7
$\phi_1 = 0.9995$	8.2 ± 0.2	8.1 ± 0.3	8.1 ± 0.2	8.1 ± 0.5

TABLE II: Summary of the synthetic data test 2. The mean and RMS value of the diffusion length estimation given in cm, is based on 100 realizations of the experiment for each set of AR-1 coefficient ϕ_1 and sampling interval Δx .

Site	Location	Accum. Rate [myr1]	Temperature [C]	σ_{18}^2 [m]
Dome C	75°06′S 123°21′E	0.027	-54.5	0.067
GISP2	72°36′N 38°30′W	0.24	-31.4	0.079
GRIP	72°35′N 37°38′W	0.23	-31.7	0.0795
NEEM	77°45′S 51°06′W	0.2	-30	0.088
NorthGRIP	75°10′N 42°32′W	0.207	-32	0.081
SipleDome	81°40′S 148°46′W	0.087	-25	0.145
South Pole	90°S 00°	0.076	-51	0.054
Vostoc	78°27′S 10°51′E	0.024	-55.5	0.067

TABLE III: Surface forcing used for the diffusion length calculations in Fig. 2.

	f_o	f_1	$ ho_0$	$ ho_{ m co}$
Experiment 1	1	1	320 kgm^{-3}	$804 \pm 20 { m kgm}^{-3}$
Experiment 2	1	1	$320 \pm 40 \; { m kgm}^{-3}$	$804~\mathrm{kgm}^{-3}$
Experiment 3	1 ± 0.2	1 ± 0.2	320 kgm^{-3}	$804 \; {\rm kgm^{-3}}$
Experiment 4	1 ± 0.2	1 ± 0.2	$320 \pm 40 \; { m kgm}^{-3}$	$804 \pm 20 \ { m kgm^{-3}}$

TABLE IV: Summary of the sensitivity experiments run




Fig. 1: Ice diffusivity parametrizations based on isotopic probe experiments for the temperature range 200 - 270 K




Fig. 2: Calculation of σ_{18}^2 for the close–off density of $\rho_{co}=804.3~{\rm kgm}^{-3}$ in m of ice equivalent. $\rho_o=330~{\rm kgm}^{-3}$ for all sites.




Fig. 3: Examples of raw $\delta^{18}{\rm O}$ data and estimated power spectral densities \hat{P}_s for four depth intervals using $\mu=30$ (red spectra) and $\mu=40$ (black spectra). The power spectral model is illustrated; P_s in green, P_σ in cyan and $|\hat{\eta}(k)|^2$ in pink. The red vertical line in the top three plots indicates the approximate position of the frequency representing the annual layer thickness. For the bottom plot this frequency is $\approx 40\,{\rm m}^{-1}$ and omitted from the plot.




Fig. 4: Result of the AR order selection test (section V). The black curve is the mean value of the diffusion length calculated using a μ in the range [40,80]. The blue curve is the diffusion length curve corrected for ice flow thinning (red curve) effects. The mean P_0 and σ_{η}^2 values are given in the two middle plots. In the top plot the standard deviation of the 41 estimates of the diffusion length for every depth in m ice eq. is shown.




Fig. 5: Synthetic data test 1. Time series after diffusion and discrete sampling for $\phi=0.9995$ and 0.2. The power spectral densities and the estimated \hat{P}_s are also presented. (a) $\sigma=0.05\,\mathrm{m}$. (b) $\sigma=0.035\,\mathrm{m}$. (c) $\sigma=0.025\,\mathrm{m}$.




Fig. 6: Synthetic data test 2. $\sigma = 0.08\,\mathrm{m}$ for all 4 sets of experiments. Power spectral densities (column 1) and time series after convolution with the diffusion filter and discrete sampling (columns 2 and 3) for both values of ϕ_1 are presented. Here we show the 100th realization of the experiment for every set. (a) $\Delta = 0.05\,\mathrm{m}$. (b) $\Delta = 0.08\,\mathrm{m}$. (c) $\Delta = 0.10\,\mathrm{m}$. (d) $\Delta = 0.12\,\mathrm{m}$.




Fig. 7: Effect of the ice layer thinning on the value of the diffusion length. For the calculation of $\sigma_{\rm firn}$ the parameters we used for the H–L model were typical of Holocene conditions for the NorthGRIP site: P=0.7 Atm, $\rho_0=330~{\rm kgm^{-3}}$, $\rho_{\rm CO}=804.3~{\rm kgm^{-3}}$, T=242.15 K, and $A=0.2~{\rm myr^{-1}}$ ice eq.




Fig. 8: Modeled power spectral densities for 6 different depths using diffusion length values from the calculation of Fig. 7 (Note that all plots represent identical accumulation rate and temperature forcing at the surface). We use the spectral model as in Eq. 19 with $q_1=0.15$, $P_{ini}=0.2$. The blue highlighted area represents the expected spectra for the case of $\Delta=5 \mathrm{cm} \left(f_{Nyq}=10 \, \mathrm{m}^{-1}\right)$ while the full range of the modeled spectra represents the case with $\Delta=2.5 \, \mathrm{cm} \left(f_{Nyq}=20 \, \mathrm{m}^{-1}\right)$.




Fig. 9: Discrete sampling correction




Fig. 10: Ice thinning uncertainty. Temperature reconstructions for NorthGRIP using (from top to bottom) S(2100m) = 0.22, 0.24, 0.26, 0.28, 0.30, 0.32. Records filtered with a 500y low-pass filter.




Fig. 11: Firn density measurements from NorthGRIP (red) compared to implementations of the H-L densification model with varying values of f. Solid and dashed curves represent 1σ and 2σ respectively.




Fig. 12: Sensitivity tests results. The top panel illustrates the mean temperature history as calculated from Experiment 4 (all parameters varied) bracketed by the 95% confidence interval estimated with the sensitivity test. For the 95% confidence interval we have also taken into account the uncertainty of the spectral estimation based on the synthetic data tests and using an RMS value of $\pm 0.5 \, \mathrm{cm}$ that is equivalent to $\approx \pm 1 \, \mathrm{K}$ in temperature. In the bottom panel we present the value of the standard deviation (1σ) for each sensitivity test.