

Available online at www.sciencedirect.com

Nuclear Physics B **Proceedings** Supplement

Nuclear Physics B Proceedings Supplement 00 (2021) 1[–4](#page-2-0)

Generation of cosmic magnetic fields in electroweak plasma

Maxim Dvornikov^{a,b}

^a Institute of Physics, University of São Paulo, CP 66318, CEP 05315-970 São Paulo, SP, Brazil *^bPushkov Institute of Terrestrial Magnetism, Ionosphere and Radiowave Propagation (IZMIRAN), 142190 Troitsk, Moscow, Russia*

Abstract

We study the generation of strong magnetic fields in magnetars and in the early universe. For this purpose we calculate the antisymmetric contribution to the photon polarization tensor in a medium consisting of an electronpositron plasma and a gas of neutrinos and antineutrinos, interacting within the Standard Model. Such a contribution exactly takes into account the temperature and the chemical potential of plasma as well as the photon dispersion law in this background matter. It is shown that a nonvanishing Chern-Simons parameter, which appears if there is a nonzero asymmetry between neutrinos and antineutrinos, leads to the instability of a magnetic field resulting to its growth. We apply our result to the description of the magnetic field amplification in the first second of a supernova explosion. It is suggested that this mechanism can explain strong magnetic fields of magnetars. Then we use our approach to study the cosmological magnetic field evolution. We find a lower bound on the neutrino asymmetries consistent with the well-known Big Bang nucleosynthesis bound in a hot universe plasma. Finally we examine the issue of whether a magnetic field can be amplified in a background matter consisting of self-interacting electrons and positrons.

Keywords: magnetic field, Chern-Simons theory, magnetar, early universe

The origin of magnetic fields (*B* fields) in some astrophysical and cosmological media is still a puzzle for the modern physics and astrophysics. There are multiple models for the generation of strong *B* fields in magnetars [\[1\]](#page-2-1). The observable galactic *B* field can be a remnant of a strong primordial *B* field existed in the early universe [\[2\]](#page-2-2). Recently the indication on the existence of the inflationary *B* field was claimed basing on the analysis of BICEP2 data [\[3\]](#page-2-3). In the present work we analyze the possibility for the strong *B* field generation in an electroweak plasma. First, we study the *B* field generation driven by neutrino asymmetries. Then, we apply our results for the description of strong *B* fields in magnetars and in the early universe. Finally, we analyze the evolution of a *B* field in a self-interacting electronpositron plasma.

To study the *B* field evolution we start with the anal-

ysis of the electromagnetic properties of an electroweak plasma consisting of electrons e^- , positrons e^+ , neutrinos *ν*, and antineutrinos $\bar{ν}$ of all types. These particles are supposed to interact in frames of the Fermi theory. This interaction is parity violating. Thus the photon polarization tensor $\Pi_{\mu\nu}$ acquires a contribution [\[4\]](#page-2-4),

$$
\Pi_{ij}(k) = i\varepsilon_{ijn} k^n \Pi_2 + \dots,
$$
 (1)

where $\Pi_2 = \Pi_2(k)$ is the new form factor, or the Chern-Simons (CS) parameter, we will be looking for and $k^{\mu} = (k_0, \mathbf{k})$ is the photon momentum. Here we adopt
the potation of [5] the notation of [\[5\]](#page-2-5)

First, we will be interested in the contribution to Π_2 arising from the interaction of a e^-e^+ plasma with a $\nu\bar{\nu}$
as a In this case the most general analytical expresgas. In this case the most general analytical expression for Π_2 can be obtained on the basis of the Feyn-man diagram shown in Fig. [1.](#page-1-0) We shall represent Π_2 as $\Pi_2 = \Pi_2^{(\nu)} + \Pi_2^{(\nu e)}$, where $\Pi_2^{(\nu)}$ is the contribution of only the neutrino gas and $\Pi_2^{(ve)}$ is the contribution of the e^-e^+ plasma with the nonzero temperature *T* and the chemi-

Email address: maxim.dvornikov@usp.br (Maxim Dvornikov)

Figure 1: The Feynman diagram for the one loop contribution to the photon polarization tensor in case of a e^-e^+ plasma interacting with a $v\bar{v}$ gas. The electron propagators are shown as broad straight lines since they account for the densities of background ν and $\bar{\nu}$ [\[6\]](#page-2-6).

cal potential μ .

The expression for $\Pi_2^{(v)}$ can be obtained using the standard quantum field theory technique [\[6\]](#page-2-6),

$$
\Pi_2^{(\nu)} = V_5 \frac{e^2}{2\pi^2} \frac{k^2}{m^2} \int_0^1 dx \frac{x(1-x)}{1 - \frac{k^2}{m^2} x(1-x)}.
$$
 (2)

where e is the electron charge, m is the electron mass, $V_5 = (V_R - V_L)/2$, and $V_{R,L}$ are the potentials of the interaction of right and left chiral projections of the *e*[−]e⁺ field with the *ν*ν̄ background. The explicit form of *V*_{R,L} can be found in [\[6\]](#page-2-6).

The expression for $\Pi_2^{(ve)}$ can be obtained using the technique for the summation over the Matsubara frequencies [\[6\]](#page-2-6),

$$
\Pi_2^{(ve)} = V_5 e^2 \int_0^1 dx \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\mathcal{E}_{\mathbf{p}}^3}
$$

$$
\times \left\{ I_0^+ - (1 - x) \left[\frac{1}{\mathcal{E}_{\mathbf{p}}^2} (\mathbf{p}^2 [3 - 5x] - 3x \left[k^2 x (1 - x) + m^2 \right] \right) \left(J_0^+ + J_0^- \right) - \frac{\beta k_0}{2} x (1 - 2x) \left(J_1^+ - J_1^- \right) - x \left(J_2^+ + J_2^- \right) \right\}, \quad (3)
$$

where

$$
I_0^+ = \frac{1}{\exp[\beta(\mathcal{E}_{\mathbf{p}} + \mu_+)] + 1} + \frac{\beta \mathcal{E}_{\mathbf{p}}}{2} \frac{1}{\cosh[\beta(\mathcal{E}_{\mathbf{p}} + \mu_+)] + 1} + (\mu_+ \to -\mu_+),
$$

$$
J_0^+ = \frac{1}{\exp[\beta(\mathcal{E}_{\mathbf{p}} + \mu_+)] + 1} + \frac{\beta \mathcal{E}_{\mathbf{p}}}{2} \frac{1 + \frac{\beta \mathcal{E}_{\mathbf{p}}}{3} \tanh\left[\frac{\beta}{2}(\mathcal{E}_{\mathbf{p}} + \mu_+)\right]}{1 + (\mu_+ \to -\mu_+),}
$$

$$
J_1^+ = \frac{1 + \beta \mathcal{E}_{\mathbf{p}} \tanh\left[\frac{\beta}{2}(\mathcal{E}_{\mathbf{p}} + \mu_+)\right]}{\cosh[\beta(\mathcal{E}_{\mathbf{p}} + \mu_+)] + 1} - (\mu_+ \to -\mu_+),
$$

$$
J_2^+ = \frac{1}{\exp[\beta(\mathcal{E}_{\mathbf{p}} + \mu_+)] + 1} + \frac{\beta \mathcal{E}_{\mathbf{p}}}{2} \frac{1 - \beta \mathcal{E}_{\mathbf{p}} \tanh\left[\frac{\beta}{2}(\mathcal{E}_{\mathbf{p}} + \mu_+)\right]}{1 + (\mu_+ \to -\mu_+)}.
$$
 (4)

Figure 2: The function *F* versus k_0 for a e^-e^+ plasma interacting with a $v\bar{v}$ gas. (a) Degenerate relativistic plasma. (b) Hot relativistic plasma.

Here $\mathcal{E}_{\mathbf{p}} = \sqrt{\mathbf{p}^2 + M^2}$, $\beta = 1/T$, $\mu_+ = \mu + k_0 x$ and $M^2 = m^2 - k^2 x (1 - x)$. To obtain $J_{0,1,2}^-$ in Eq. [\(3\)](#page-1-1) we should replace $\mu_+ \rightarrow \mu_- = \mu - k_0 x$ in I^+ in Eq. (4) should replace $\mu_+ \to \mu_- = \mu - k_0 x$ in $J_{0,1,2}^+$ in Eq. [\(4\)](#page-1-2). Should replace μ_+ μ_- = μ κ_0 x in $J_{0,1,2}$ in Eq. (7).
Note that in Eqs. [\(3\)](#page-1-1) and [\(4\)](#page-1-2) we assume that $k^2 < 4m^2$, i.e. no creation of e^-e^+ pairs occurs [\[7\]](#page-2-7).

It is convenient to represent Π_2 as $\Pi_2 = 2 \frac{\alpha_{em}}{\pi} V_5 F$, where *F* is the dimensionless function and $\alpha_{em} = \frac{e^2}{4t}$
is the fine structure constant. Using Eqs. (2)-(4) is where \overline{t} is the dimensionless random and $a_{em} = 4\pi$
is the fine structure constant. Using Eqs. [\(2\)](#page-1-3)-[\(4\)](#page-1-2), in Fig. [2](#page-1-4) we show the behavior of F versus k_0 in relativistic plasmas. It should be noted that in the static limit $F(k_0 = 0) \neq 0$. To plot Fig. [2](#page-1-4) we take into account the dispersion law of long electromagnetic waves in plasma $k^2 = k^2(T, \mu)$ [\[6\]](#page-2-6) and the fact that an electron
acquires the effective mass $m^2 = e^2 (r^2 + \pi^2 T^2)$ in a acquires the effective mass $m_{\text{eff}}^2 = \frac{e^2}{8\pi^2}(\mu^2 + \pi^2 T^2)$ in a
hot and dense matter [7]. As shown in [6], the nonzero hot and dense matter [\[7\]](#page-2-7). As shown in [\[6\]](#page-2-6), the nonzero $\Pi_2(0) = \Pi_2(k_0 = 0)$ results in the instability of a *B* field leading to the exponential growth of a seed field.

We can apply our results for the description of the *B* field evolution in a dense relativistic electron gas in a supernova explosion. It is known that, just after the core collapse, a supernova is a powerful source of ^ν*^e* whereas the fluxes of $v_{\mu,\tau}$ and $\bar{v}_{e,\mu,\tau}$ are negligible [\[8\]](#page-2-8). Thus $V_5 \neq 0$ and we get that $\Pi_2(0) = \frac{\sqrt{2}}{\pi}$ $V_5 \neq 0$ and we get that $\Pi_2(0) = \frac{\sqrt{2}}{\pi} \alpha_{\text{em}} G_{\text{F}} n_{\nu_e} F(0) \neq 0$, where G_F is the Fermi constant and $|F(0)| \approx 2$, see Fig. [2\(a\),](#page-1-5) since electrons are degenerate. The magnetic diffusion time $t_{diff} = \sigma \Pi_2^{-2}(0) \approx 2.3 \times 10^{-2}$ s for $n = 3.7 \times 10^{37}$ cm⁻³ and $n = 1.9 \times 10^{37}$ cm⁻³ in the $n_e = 3.7 \times 10^{37} \text{ cm}^{-3}$ and $n_{\nu_e} = 1.9 \times 10^{37} \text{ cm}^{-3}$ in the superpoval core [6]. Here σ is the electron gas conduc-supernova core [\[6\]](#page-2-6). Here σ is the electron gas conductivity. Thus at $t \sim 10^{-3}$ s $\ll t_{\text{diff}}$, when the flux of v_e is maximal, no seed magnetic field dissinates. Therefore maximal, no seed magnetic field dissipates. Therefore the neutrino driven instability can result in the growth of the *B* field. It should be noted that the scale of the *B* field turns out to be small $\Lambda \sim 10^{-3}$ cm. However, at later stages of the star evolution V_5 diminishes and Λ can be comparable with the magnetar radius. Thus our mechanism can be used to explain strong *B* fields of magnetars.

Figure 3: The Feynman diagrams contributing to the photon polarization tensor in case of a e^-e^+ self-interacting plasma. Here A_μ is the potential of the electromagnetic field.

Now let us apply out results to study the *B* field evolution in the primordial plasma. At the stages of the early universe evolution before the neutrino decoupling at $T > (2 - 3)$ MeV, the e^-e^+ plasma is hot and relativistic Assuming the causal scenario in which relativistic. Assuming the causal scenario, in which $\Lambda < H^{-1}$, where *H* is the Hubble constant, we get that $|E| = E = E + 1.11 \times 10^{-6} \sqrt{a^* / 106.75} \times (T/\text{MeV})^{-1}$ $|\xi_{v_e} - \xi_{v_\mu} - \xi_{v_\tau}| > 1.1 \times 10^{-6} \sqrt{g^*/106.75} \times (T/\text{MeV})^{-1}$,
see [6] where $\xi = u/T$ a^{*} is the number of relativis-see [\[6\]](#page-2-6), where $\xi_{\alpha} = \mu_{\alpha}/T$, g^* is the number of relativistic degrees of freedom, and μ_{α} is the chemical potential tic degrees of freedom, and μ_{α} is the chemical potential of neutrinos of the type $\alpha = v_e, v_\mu, v_\tau$. Here we use that $|F(0)| \approx 0.2$, see Fig. [2\(b\).](#page-1-6) Assuming that before the Big Bang nucleosynthesis at $T \sim (2-3)$ MeV all neutrino flavors equilibrate owing to neutrino oscillations $\xi_{v_e} \sim \xi_{v_\mu} \sim \xi_{v_\tau}$, we get the lower bound on the neutrino
asymmetries, which is consistent with the well-known asymmetries, which is consistent with the well-known Big Bang nucleosynthesis upper bound on $|\xi_{\alpha}|$, see [\[9\]](#page-2-9).

Finally, let us examine the issue of whether a *B* field can be amplified in a e^-e^+ plasma self-interacting within the Fermi model, i.e. when a $v\bar{v}$ gas is not present. In this case the contributions to Π_2 are schematically de-picted in Fig. [3.](#page-2-10) The analytical expression for $\Pi_2^{(ee)}$ $\frac{1}{2}$ can be obtained analogously to the previous case $[10]$,

$$
\Pi_{2}^{(ee)} = \frac{\left(1 - 4\sin^{2}\theta_{W}\right)}{2\sqrt{2}} e^{2} G_{F} \left(n_{e} - n_{\bar{e}}\right) \int_{0}^{1} (1 - x) dx
$$
\n
$$
\times \int \frac{d^{3}p}{(2\pi)^{3}} \frac{1}{\mathcal{E}_{p}^{3}} \left\{ \left[J_{2}^{\prime} - J_{2}^{\prime\prime}\right] - \left[J_{0}^{\prime} - J_{0}^{\prime\prime}\right] \frac{1}{\mathcal{E}_{p}^{2}} \right\}
$$
\n
$$
\times \left(\mathbf{p}^{2} \left[3 - 2x\right] - 3\left[m^{2}(1 + x) + k^{2}x^{2}\right]\right)\right\}, \quad (5)
$$

where $n_{e,\bar{e}}$ are the electron and positron densities, θ_W is the Weinberg angle, and $J'_{0,2} = J_{0,2}^+$ in Eq. [\(4\)](#page-1-2), with $U' = U_0$. The expressions for I'' can be obtained from $J'_{0,2}$ if we make the replacement $\mu' \to \mu'' = \mu + k_0(1-x)$
there As in deriving of Eqs. (3) and (4) here we also $\mathcal{U} = \mu_+$. The expressions for $J_{0,1}^{\prime\prime}$ can be obtained from
 \mathcal{U} if we make the replacement $u' \rightarrow u'' - u + k_0(1 - x)$ $\mu_{0,2}$ in we make the replacement $\mu_{0,1}$ $\mu_{0,2}$ in $\mu_{0,1}$ $\mu_{0,2}$
there. As in deriving of Eqs. [\(3\)](#page-1-1) and [\(4\)](#page-1-2), here we also assume that $k^2 < 4m^2$.

Let us express

Let us express Π_2 in Eq. [\(5\)](#page-2-11) as Π_2 $\frac{\alpha_{\text{em}}}{\sqrt{2\pi}}\left(1-4\sin^2\theta_W\right)G_F(n_e-n_{\bar{e}})F$, where *F* is the $\sqrt{2\pi}$ (1) on $\sqrt{9}$ p or \sqrt{e} and $\sqrt{2}$ and $\sqrt{2\pi}$ compared the state dimensionless function. We shall analyze this function

Figure 4: The function *F* versus k_0 for a e^-e^+ self-interacting plasma. (a) Degenerate relativistic plasma. (b) Hot relativistic plasma.

in the static limit $k_0 \rightarrow 0$. We mention that, if we neglect k_0 in Eq. [\(5\)](#page-2-11), then $J'_{0,2} = J''_{0,2}$ and $\Pi_2 \to 0$. The behavior of *F* for relativistic plasmas is shown in Fig. [4,](#page-2-12) where one can see that $\Pi_2(0) = 0$. In Fig. [4](#page-2-12) we also account for the thermal corrections to the photon dispersion and to the electron mass. It means that a e^-e^+ plasma does not reveal the instability of a *B* field leading to its growth. Therefore, contrary to the claim of [\[5\]](#page-2-5), one can use this mechanism for neither the explanation of strong *B* fields of magnetars nor the *B* field amplification in the early universe.

In conclusion we mention that we have derived the CS term Π_2 in an electroweak plasma consisting of $e^$ and e^+ as well as *v* and $\bar{\nu}$ of all flavors. These particles are involved in the parity violating interaction. It makes are involved in the parity violating interaction. It makes possible the existence of a nonzero CS term. In case of a e^-e^+ plasma interacting with a $v\bar{v}$ background, the CS term is nonvanishing in the static limit when $k_0 = 0$. CS term is nonvanishing in the static limit when $k_0 = 0$. Therefore, a *B* field becomes unstable in this system. We have shown that a seed field can be exponentially amplified. This feature of an electroweak plasma in question can be used to explain strong *B* fields of magnetars and to study the evolution of a primordial *B* field. We have also demonstrated that there is no *B* field instability in a self-interacting e^-e^+ plasma.

I am thankful to the organizers of $37th$ ICHEP for the invitation and a financial support, to V.B. Semikoz for helpful discussions, and to FAPESP (Brazil) for a grant.

References

- [1] R.C. Duncan and C. Thompson, Asprophys. J. **392**, L9 (1992).
- [2] A. Neronov and I. Vovk, Science 328, 73 (2010).
- [3] C. Bonvin, *et al.*, Phys. Rev. Lett. 112, 191303 (2014).
- [4] S. Mohanty, *et al.*, Phys. Rev. D 58, 093007 (1998).
- [5] A. Boyarsky, et al., Phys. Rev. Lett. **109**, 111602 (2012).
- M. Dvornikov and V.B. Semikoz, JCAP 1405, 002 (2014) [\[arXiv:1311.5267\]](http://arxiv.org/abs/1311.5267).
- [7] E. Braaten, Astrophys. J. 392, 70 (1992).
- [8] H.-Th. Janka, *et al.*, Phys. Rept. 442, 38 (2007).
- [9] G. Mangano, *et al.*, Phys. Lett. B 708, 1 (2012).

[10] M. Dvornikov, Phys. Rev. D 90, 041702 (2014) [\[arXiv:1405.3059\]](http://arxiv.org/abs/1405.3059).