

DECOUPLINGS FOR CURVES AND HYPERSURFACES WITH NONZERO GAUSSIAN CURVATURE

JEAN BOURGAIN AND CIPRIAN DEMETER

ABSTRACT. We prove two types of results. First we develop the decoupling theory for hypersurfaces with nonzero Gaussian curvature, which extends our earlier work from [3]. As a consequence of this we obtain sharp (up to ϵ losses) Strichartz estimates for the hyperbolic Schrödinger equation on the torus.

Our second main result is an l^2 decoupling for non degenerate curves which has implications for Vinogradov's mean value theorem.

1. STATEMENTS OF RESULTS

Let S be a compact C^2 hypersurface in \mathbb{R}^n with nonzero Gaussian curvature. The typical example to have in mind is the truncated paraboloid defined for $v = (v_1, \dots, v_{n-1}) \in (\mathbb{R} \setminus \{0\})^{n-1}$ as

$$H_v^{n-1} := \{(\xi_1, \dots, \xi_{n-1}, v_1 \xi_1^2 + \dots + v_{n-1} \xi_{n-1}^2) : |\xi_i| \leq 1/2\}.$$

H_v^{n-1} is called elliptic when all v_i have the same sign and hyperbolic otherwise.

Let $\mathcal{N}_\delta = \mathcal{N}_\delta(S)$ be the δ neighborhood of S and let \mathcal{P}_δ be a finitely overlapping cover of \mathcal{N}_δ with $\sim \delta^{1/2} \times \dots \times \delta^{1/2} \times \delta$ rectangular boxes θ . We will denote by f_θ the Fourier restriction of f to θ .

We will write $A \sim B$ if $A \lesssim B$ and $B \lesssim A$. The implicit constants hidden inside the symbols \lesssim and \sim will in general depend on fixed parameters such as p, n, α and sometimes on variable parameters such as ϵ . We will in general not record the dependence on the fixed parameters.

Our first result is the following l^p decoupling theorem¹.

Theorem 1.1. *Let $n \geq 2$. If $\text{supp}(\hat{f}) \subset \mathcal{N}_\delta$ then for $p \geq \frac{2(n+1)}{n-1}$ and $\epsilon > 0$*

$$\|f\|_p \lesssim_\epsilon \delta^{\frac{n}{p} - \frac{n-1}{2} - \epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_p^p \right)^{1/p}. \quad (1)$$

This is a close cousin of the following l^2 decoupling proved in [3] in the case when S has definite second fundamental form

$$\|f\|_p \lesssim_\epsilon \delta^{-\frac{n-1}{4} + \frac{n+1}{2p} - \epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_p^2 \right)^{1/2}, \quad p \geq \frac{2(n+1)}{n-1}. \quad (2)$$

Key words and phrases. decouplings, mean value theorems, Strichartz estimates.

The first author is partially supported by the NSF grant DMS-1301619. The second author is partially supported by the NSF Grant DMS-1161752.

¹Perhaps a more appropriate nomenclature for the l^p and l^2 decouplings we consider in this paper would be $l^p(L^p)$ and $l^2(L^p)$ decouplings. For simplicity of notation, we prefer the former notation.

We point out that (1) is slightly weaker than (2) as it follows from (2) via Hölder's inequality. In particular, Theorem 1.1 for $n = 2$ is contained in [3]. We mention that sharp l^p decouplings were first considered by Wolff in the case of the cone, see [10].

As briefly explained in [3], (2) is false for the hyperbolic paraboloid due to the fact that it contains lines. On the other hand, apart from the dependence on ϵ , inequality (1) is sharp, as is (2). This can be easily seen by considering the case of the sphere $S = S^{n-1}$ and f with $\hat{f} = 1_{\mathcal{N}_\delta}$. The main new difficulty in proving (1) as compared to (2) is the fact that intersections of hyperbolic paraboloids with hyperplanes do not always have nonzero Gaussian curvature. It will however be crucial to our argument the fact that at most one of the principal curvatures of these sections can be small. An application of Theorem 1.1 to curves is discussed in Section 6.

A modification of our proof of Theorem 1.1 leads to the following related l^2 decoupling result for H_v^{n-1} . Let us denote by $d(v)$ the minimum between the number of positive and negative entries of v .

Theorem 1.2. *Let $n \geq 2$. If $\text{supp}(\hat{f}) \subset \mathcal{N}_\delta(H_v^{n-1})$ then for $p \geq 2$ and $\epsilon > 0$ we have*

$$\|f\|_p \lesssim_\epsilon \delta^{-\epsilon} K_{n,p,v}^{(2)}(\delta) \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_p^2 \right)^{1/2}, \quad (3)$$

where $K_{n,p,v}^{(2)}(\delta) = \delta^{-\frac{n-1}{4} + \frac{n+1}{2p}}$ when $p \geq \frac{2(n+1-d(v))}{n-1-d(v)}$ and $K_{n,p,v}^{(2)}(\delta) = \delta^{d(v)(-\frac{1}{4} + \frac{1}{2p})}$ when $2 \leq p \leq \frac{2(n+1-d(v))}{n-1-d(v)}$.

Note that Theorems 1.1 and 1.2 are independent, neither of them implies the other one. Of course, (3) generalizes (2) (which corresponds to $d(v) = 0$), and leads to Strichartz estimates for the hyperbolic Schrödinger equation. More precisely, fix $v_1, \dots, v_{n-1} \in \mathbb{R} \setminus \{0\}$. For $\phi \in L^2(\mathbb{T}^{n-1})$ consider the "generalized Laplacian" operator

$$T\phi(x_1, \dots, x_{n-1}) = \sum_{(\xi_1, \dots, \xi_{n-1}) \in \mathbb{Z}^{n-1}} (\xi_1^2 v_1 + \dots + \xi_{n-1}^2 v_{n-1}) \hat{\phi}(\xi_1, \dots, \xi_{n-1}) e(i\xi_1 x_1 + \dots + \xi_{n-1} x_{n-1})$$

on the irrational torus $\prod_{i=1}^{n-1} \mathbb{R}/(|v_i|\mathbb{Z})$. Let also

$$e^{itT} \phi(x_1, \dots, x_{n-1}, t) = \sum_{(\xi_1, \dots, \xi_{n-1}) \in \mathbb{Z}^{n-1}} \hat{\phi}(\xi_1, \dots, \xi_{n-1}) e(x_1 \xi_1 + \dots + x_{n-1} \xi_{n-1} + t(\xi_1^2 v_1 + \dots + \xi_{n-1}^2 v_{n-1})).$$

Following the approach described in [3], Theorem 1.2 implies the next corollary.

Corollary 1.3 (Strichartz estimates for irrational tori: the hyperbolic case). *Let $\phi \in L^2(\mathbb{T}^{n-1})$ with $\text{supp}(\hat{\phi}) \subset [-N, N]^{n-1}$. Then for each $\epsilon > 0$, $p \geq 2$ and each interval $I \subset \mathbb{R}$ with $|I| \gtrsim 1$ we have*

$$\|e^{itT} \phi\|_{L^p(\mathbb{T}^{n-1} \times I)} \lesssim_\epsilon N^\epsilon K_{n,p,v}^{(2)}(N^{-2}) |I|^{1/p} \|\phi\|_2, \quad (4)$$

where $K_{n,p,v}^{(2)}$ is as in Theorem 1.2 and the implicit constant does not depend on I and N .

Estimates of this type have been considered recently, see for example [7] and [9]. It is interesting to note that in the non elliptic case ($d(v) \geq 1$), the exponent of N in $K_{n,p,v}^{(2)}(N^{-2})$ is always nonzero, in contrast with the continuous case (when \mathbb{T}^n is replaced with \mathbb{R}^n) and the elliptic case (on either \mathbb{T}^n or \mathbb{R}^n). This exponent is sharp in the case when $|v_i| = 1$ for all i . To see this, use ϕ Fourier supported on the lattice points of a vector subspace of dimension $d(v)$ of H_v^{n-1} . This example does not exist in the case when v has rationally independent entries. Our method here does not seem to shed any light on the issue of whether one can improve the constant $K_{n,p,v}^{(2)}(N^{-2})$ in (4) in the irrational case.

It is possible that the term N^ϵ is not necessary in (4) when $p > \frac{2(n+1-d(v))}{n-1-d(v)}$. As observed in [3], this is indeed the case for the elliptic Schrödinger equation on the rational torus. See also [7] and [9] where similar sharp results are proved in the range $2 \leq p \leq 4$, when $n = 3$.

The proof of Theorem 1.2 will be sketched in Section 3.

In the second part of the paper we consider curves $\Phi : [0, 1] \rightarrow \mathbb{R}^n$,

$$\Phi(t) = (\phi_1(t), \dots, \phi_n(t))$$

with $\phi_i \in C^n([0, 1])$ and such that the Wronskian

$$W(\phi'_1, \dots, \phi'_n)(t)$$

is nonzero on $[0, 1]$. These are usually referred to in literature as *nondegenerate* curves.

Abusing earlier notation, let \mathcal{N}_δ be the δ neighborhood of $\Phi([0, 1])$ and let \mathcal{P}_δ be the partition of \mathcal{N}_δ with δ neighborhoods θ of $\Phi(I)$, with I dyadic interval of length $\delta^{1/n}$. We will as before denote by f_θ the Fourier restriction of f to θ . One important aspect to note at this point is the fact that θ is a curved tube, in particular not a cylinder. To make θ a cylinder, one needs to consider intervals I of length $\delta^{1/2}$.

Theorem 1.4. *For each such curve Φ and each $f : \mathbb{R}^n \rightarrow \mathbb{C}$ with Fourier support in \mathcal{N}_δ we have*

$$\|f\|_{L^p(\mathbb{R}^n)} \lesssim_\epsilon \delta^{-\epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_{L^p(\mathbb{R}^n)}^2 \right)^{\frac{1}{2}},$$

for each $\epsilon > 0$ and $2 \leq p \leq 4n - 2$.

By using a limiting procedure (see for example the discussion in [3] and (34) below) one can replace f with a sum of Dirac deltas and derive the following corollary.

Corollary 1.5. *Fix Φ as above and let $p \leq 4n - 2$. Then for each δ -separated set Λ of points on the curve Φ and each coefficients $a_\xi \in \mathbb{C}$ we have*

$$\left(\frac{1}{|B_R|} \int_{B_R} \left| \sum_{\xi \in \Lambda} a_\xi e(\xi \cdot x) \right|^p dx \right)^{1/p} \lesssim_\epsilon \delta^{-\epsilon} \|a_\xi\|_{l^2(\Lambda)}, \quad (5)$$

for each ϵ and each ball $B_R \subset \mathbb{R}^n$ of radius $R \gtrsim \delta^{-n}$.

In particular, by applying this to (a rescaled version of) the curve

$$\Phi(t) = (t, t^2, \dots, t^n)$$

we get

$$\left(\frac{1}{|B_R|} \int_{B_R} \left| \sum_{\xi \in \Lambda} a_\xi e(\xi \cdot x) \right|^p dx\right)^{1/p} \lesssim_\epsilon N^{-\epsilon} \|a_\xi\|_{l^2(\Lambda)}, \quad (6)$$

for $p \leq 4n - 2$, each 1-separated set Λ of points on the curve Φ ($0 \leq t \leq \infty$) with size N and each $R \gtrsim N^n$.

Let us now explore an immediate consequence. Given an integer $k \geq 2$ we define the k -energy of Λ as

$$\mathbb{E}_k(\Lambda) = |\{(\lambda_1, \dots, \lambda_{2k}) \in \Lambda^{2k} : \lambda_1 + \dots + \lambda_k = \lambda_{k+1} + \dots + \lambda_{2k}\}|.$$

By letting $R \rightarrow \infty$ in (6) with $p = 4n - 2$ and $a_\xi = 1$, we immediately get that

$$\mathbb{E}_{2n-1}(\Lambda) \lesssim_\epsilon \Lambda^{2n-1+\epsilon}$$

for each Λ as above. In particular, we have

$$\mathbb{E}_{2n-1}(\{(l, l^2, \dots, l^n) : l = 1, 2, \dots, N\}) \lesssim_\epsilon N^{2n-1+\epsilon}.$$

This is a special case of Vinogradov's mean value theorem, due to (and significantly improved by the recent work of) Wooley, see for example [11] and the references therein. Our method however shows that the integer case is not special, but is rather a particular case of a larger phenomenon. Note also that our method allows for arbitrary coefficients a_ξ in (6).

Vinogradov's mean value theorem conjectures that (6) should hold in the integer case, with $a_\xi = 1$, for p as large as $n(n+1)$. It is possible that (6) also holds for $p \leq n(n+1)$, for arbitrary coefficients and frequency points Λ . However, our method does not currently seem to shed any light on this issue. Further applications of variants of inequality (6) to number theory are presented in [4].

Acknowledgment. The second author is indebted to Andrea Nahmod for bringing the question about the hyperbolic Schrödinger equation to his attention and for stimulating discussions on the topic.

2. l^p DECOUPLINGS FOR HYPERSURFACES

In this section we present the proof of Theorem 1.1. We start by observing that the induction on scales argument from the last section in [3] allows us to focus on the hypersurfaces H_ν^{n-1} .

The proof of Theorem 1.1 for H_ν^{n-1} will be done in two separate stages. First, we develop the multilinear decoupling theory and show that it is essentially equivalent to its linear counterpart.

Let $g : H_\nu^{n-1} \rightarrow \mathbb{C}$. For a cap τ on H_ν^{n-1} we let $g_\tau = g|_\tau$ be the (spatial) restriction of g to τ . We denote by $\pi : H_\nu^{n-1} \rightarrow [-1/2, 1/2]^{n-1}$ the projection map and by $d\sigma$ the natural surface measure on H_ν^{n-1} .

Definition 2.1. *We say that the caps τ_1, \dots, τ_n on H_ν^{n-1} are ν -transverse if the volume of the parallelepiped spanned by any unit normals n_i at τ_i is greater than ν .*

In the following, the norm $\|f\|_{L^p(w_{B_R})}$ will refer to the weighted L^p integral

$$\left(\int_{\mathbb{R}^n} |f(x)|^p w_{B_R}(x) dx \right)^{1/p}$$

for some weight w_{B_R} which is Fourier supported in $B(0, \frac{1}{R})$ and satisfies

$$1_{B_R}(x) \lesssim w_{B_R}(x) \leq \left(1 + \frac{|x - c(B_R)|}{R}\right)^{-10n}. \quad (7)$$

We denote by $C_{p,n,\nu}(\delta, \nu)$ the smallest constant such that

$$\left\| \left(\prod_{i=1}^n |g_{\tau_i} \widehat{d\sigma}| \right)^{1/n} \right\|_{L^p(B_{\delta^{-1}})} \leq C_{p,n,\nu}(\delta, \nu) \left[\prod_{i=1}^n \left(\sum_{\substack{\theta: \delta^{1/2}\text{-cap} \\ \theta \subset \tau_i}} \|g_{\theta} \widehat{d\sigma}\|_{L^p(w_{B_{\delta^{-1}}})}^p \right)^{1/p} \right]^{1/n},$$

for each ν -transverse caps $\tau_i \subset H_v^{n-1}$, each δ^{-1} ball $B_{\delta^{-1}}$ and each $g : H_v^{n-1} \rightarrow \mathbb{C}$.

Let also $K_{p,n,\nu}(\delta)$ be the smallest constant such that

$$\|\widehat{gd\sigma}\|_{L^p(B_{\delta^{-1}})} \leq K_{p,n,\nu}(\delta) \left(\sum_{\theta: \delta^{1/2}\text{-cap}} \|g_{\theta} \widehat{d\sigma}\|_{L^p(w_{B_{\delta^{-1}}})}^p \right)^{1/p},$$

for each $g : H_v^{n-1} \rightarrow \mathbb{C}$ and each δ^{-1} ball $B_{\delta^{-1}}$.

Hölder's inequality gives

$$C_{p,n,\nu}(\delta, \nu) \leq K_{p,n,\nu}(\delta).$$

We will show that the reverse inequality essentially holds true. This will be an important intermediate step that will help us prove that for $2 \leq p \leq \frac{2(n+1)}{n-1}$

$$K_{p,n,\nu}(\delta) \lesssim_{\epsilon} \delta^{-\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})-\epsilon}. \quad (8)$$

Note that this estimate is the expected one, if the trivial bound for $p = 2$ is interpolated with the bound at the critical index $p = \frac{2(n+1)}{n-1}$. In other words, (8) is an equivalent reformulation of Theorem 1.1.

Theorem 2.2. *Fix $n \geq 3$, $\nu \in \{-1, 1\}^{n-1}$ and let $p \geq 2$. Assume one of the following holds*

- (i) $n = 3$
- (ii) $n \geq 4$ and

$$K_{p,n-2,\nu'}(\delta') \lesssim_{\epsilon} \delta'^{-\frac{n-3}{2}(\frac{1}{2}-\frac{1}{p})-\epsilon} \quad (9)$$

for each $\delta' > 0$, $\nu' \in \{-1, 1\}^{n-3}$ and each $\epsilon > 0$.

Then for each $0 < \nu \leq 1$ there is $\epsilon(\nu)$ with $\lim_{\nu \rightarrow 0} \epsilon(\nu) = 0$ and C_{ν} such that

$$K_{p,n,\nu}(R^{-1}) \leq C_{\nu} R^{\epsilon(\nu)} C_{p,n,\nu}(R^{-1}, \nu)$$

for each $R > 1$.

We first prove a lemma for paraboloids that are allowed to have one small (possibly zero) principal curvature. The motivation behind this consideration will be explained in the end of the proof of Proposition 2.6.

Lemma 2.3. *Let $n \geq 3$. Fix $v_1, \dots, v_{n-2} \in \{-1, 1\}$ and let $|a| \lesssim 1$ be arbitrary, possibly zero. Let \mathcal{P}_δ be a partition of the neighborhood \mathcal{N}_δ associated with the hypersurface $H_{(v_1, \dots, v_{n-2}, a)}^{n-1}$.*

If $\text{supp}(\hat{f}) \subset \mathcal{N}_\delta$ then for $p \geq 2$ we have, uniformly over the parameter $|a| \lesssim 1$

$$\|f\|_p \lesssim \delta^{-\frac{1}{2} + \frac{1}{p}} K_{p, n-1, (v_1, \dots, v_{n-2})}(\delta) \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_p^p \right)^{1/p}.$$

Before we embark in the proof of the lemma, we give some heuristics on numerology. A simple L^p orthogonality principle asserts that given any pairwise disjoint subsets S_1, \dots, S_M in \mathbb{R}^n we have

$$\|f\|_p \leq M^{1 - \frac{2}{p}} \left(\sum_{i=1}^M \|f_{S_i}\|_p^p \right)^{1/p} \quad (10)$$

for each $2 \leq p \leq \infty$ and each f Fourier supported in the union of the S_i . We may refer to this as being *trivial L^p decoupling*. If the sets S_i are arbitrary, the universal exponent $1 - \frac{2}{p}$ of M is sharp. To see, it suffices to consider the case when S_i are equidistant unit balls with collinear centers. However, this exponent becomes smaller when geometry is favorable. For example, the L^p decoupling inequality (8) corresponds to $M \sim \delta^{-\frac{n-1}{2}}$, and the exponent there is $\frac{1}{2} - \frac{1}{p}$, half of the universal one. The absence of curvature is an enemy, and one expects a penalty of $\delta^{\frac{1}{2p} - \frac{1}{4}}$ for each zero principal curvature. For example, when a is small (possibly zero), $H_{(v_1, \dots, v_{n-2}, a)}^{n-1}$ has a decoupling constant $\delta^{\frac{1}{2p} - \frac{1}{4}}$ larger than in the case $a \sim 1$.

Proof The proof is rather standard, we sketch it briefly. The case $n = 3$ is entirely typical, we prefer it only to simplify the notation. We can of course also assume $v_1 = 1$. By first performing the trivial decoupling (10) in the direction of e_2 (which corresponds to the entry a), it suffices to prove that for each $\alpha \in [-\frac{1}{2}, \frac{1}{2}]$ we have

$$\left\| \sum_{\theta \in \mathcal{P}_{\delta, \alpha}} f_\theta \right\|_p \lesssim K_{p, 2, 1}(\delta) \left(\sum_{\theta \in \mathcal{P}_{\delta, \alpha}} \|f_\theta\|_p^p \right)^{1/p}, \quad (11)$$

where $\mathcal{P}_{\delta, \alpha}$ consists of those $\theta \in \mathcal{P}_\delta$ that intersect the parabola

$$\{(\xi_1, \alpha, \xi_3) : \xi_3 = \xi_1^2 + a\alpha^2\}.$$

We next show how a standard parabolic change of coordinates will allow us to assume $\alpha = 0$. It is easy to see (the reader is again referred to [3]) that (11) is equivalent with

$$\int_{B_{\delta^{-1}}} \left| \int_{[-1/2, 1/2] \times [\alpha, \alpha + \delta^{1/2}]} f(\xi_1, \xi_2) e(x_1 \xi_1 + x_2 \xi_2 + x_3(\xi_1^2 + a\xi_2^2)) d\xi_1 d\xi_2 \right|^p dx_1 dx_2 dx_3 \lesssim K_{p, 2, 1}(\delta)^p \sum_{I: \delta^{1/2}\text{-interval}} \int_{B_{\delta^{-1}}} \left| \int_{I \times [\alpha, \alpha + \delta^{1/2}]} f(\xi_1, \xi_2) e(x_1 \xi_1 + x_2 \xi_2 + x_3(\xi_1^2 + a\xi_2^2)) d\xi_1 d\xi_2 \right|^p dx_1 dx_2 dx_3$$

for each $B_{\delta^{-1}}$. It is now rather immediate that we can assume $\alpha = 0$, since the image of $B_{\delta^{-1}}$ under the transformation

$$(x_1, x_2, x_3) \mapsto (x_1, x_2 + 2\alpha a x_3, x_3)$$

is roughly $B_{\delta^{-1}}$.

Note however that all $\theta \in \mathcal{P}_{\delta,0}$ lie in the δ neighborhood of the cylinder

$$\{(\xi_1, \xi_2, \xi_3) : \xi_3 = \xi_1^2\},$$

and (11) follows immediately from Fubini. \blacksquare

A simple induction on scales similar to the one in Section 7 in [3] allows us to extend the previous lemma to arbitrary hypersurfaces with (at least) $n - 2$ principal curvatures bounded away from zero.

Lemma 2.4. *Let $n \geq 3$ and $p \geq 2$. Let S be a C^2 compact hypersurface in \mathbb{R}^n which at any given point has at least $n - 2$ principal curvatures with magnitudes ~ 1 , while the remaining one is $\lesssim 1$. Let as usual \mathcal{P}_δ be a partition of the neighborhood \mathcal{N}_δ associated S . Assume that for each $\delta' > 0$*

$$\max_{v \in \{-1,1\}^{n-2}} K_{p,n-1,v}(\delta') \lesssim_\epsilon \delta'^{-\frac{n-2}{2}(\frac{1}{2}-\frac{1}{p})-\epsilon}.$$

Then for each δ and each $\text{supp}(\hat{f}) \subset \mathcal{N}_\delta$ we have

$$\|f\|_p \lesssim_\epsilon \delta^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{p})-\epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_p^p \right)^{1/p}.$$

Proof For $\delta < 1$, let as before $K_{p,n,S}(\delta)$ be the smallest constant such that for each f with Fourier support in \mathcal{N}_δ we have

$$\|f\|_p \leq K_{p,n,S}(\delta) \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_p^p \right)^{1/p}.$$

First note that for each such f

$$\|f\|_p \leq K_{p,n,S}(\delta^{\frac{2}{3}}) \left(\sum_{\tau \in \mathcal{P}_{\delta^{\frac{2}{3}}}} \|f_\tau\|_p^p \right)^{1/p}.$$

Second, our assumption on the principal curvatures of S combined with Taylor's formula shows that on each $\tau \in \mathcal{P}_{\delta^{\frac{2}{3}}}$, S is within δ from a paraboloid H_v^{n-1} with at least $n - 2$ of the entries of v having magnitudes of order 1. By invoking Lemma 2.3 for this paraboloid (via a simple rescaling), combined with parabolic rescaling we get

$$\|f_\tau\|_p \lesssim (\delta^{1/3})^{\frac{1}{p}-\frac{1}{2}} \max_{v \in \{-1,1\}^{n-2}} K_{p,n-1,v}(\delta^{1/3}) \left(\sum_{\theta \in \mathcal{P}_\delta: \theta \subset \tau} \|f_\theta\|_p^p \right)^{1/p}.$$

For each $\epsilon > 0$, we conclude the existence of C_ϵ such that for each $\delta < 1$

$$K_{p,n,S}(\delta) \leq C_\epsilon [\delta^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{p})-\epsilon}]^{1/3} K_{p,n,S}(\delta^{\frac{2}{3}}).$$

By iteration this immediately leads to the desired conclusion. \blacksquare

We next present a lemma that will play a key role in the proof of Proposition 2.6 below.

Lemma 2.5. *Let A be an invertible symmetric $n \times n$ matrix and let S be an m dimensional affine subspace of \mathbb{R}^n . There exists $\delta = \delta(A)$ such that if the m dimensional quadratic hypersurface*

$$x_{m+1} = \langle Ax, x \rangle, \quad x \in S$$

has l principal curvatures in the interval $[-\delta, \delta]$ then

$$l \leq n - m.$$

Proof We may assume S contains the origin. Choose δ small enough so that the hypothesis forces the existence of an l dimensional subspace S_1 of S such that

$$\|P_S Ax\| \leq \frac{1}{2} \|A^{-1}\|^{-1} \|x\| \quad (12)$$

for each $x \in S_1$. Here P_S is the orthogonal projection onto S . We claim that $S_1 \cap A^{-1}S = \{0\}$, which will easily imply the desired conclusion. Indeed, otherwise there is $x \in S_1$ with $\|x\| = 1$ and $Ax \in S$, and (12) forces the contradiction. \blacksquare

Here is the basic step in the Bourgain-Guth-type induction on scales that relates the linear and the multilinear decoupling.

Proposition 2.6. *Fix $n \geq 3$, $v \in \{-1, 1\}^{n-1}$ and let $p \geq 2$. Assume one of the following holds*

(i) $n = 3$

(ii) $n \geq 4$ and $K_{p,n-2,v'}(\delta') \lesssim_\epsilon \delta'^{-\frac{n-3}{2}(\frac{1}{2}-\frac{1}{p})-\epsilon}$ for each $\delta' > 0$, $v' \in \{-1, 1\}^{n-3}$ and each $\epsilon > 0$.

Then for each ϵ there exist constants C_ϵ, C_n such that for each $R > 1$ and $K \geq 1$

$$\begin{aligned} \|\widehat{gd\sigma}\|_{L^p(w_{B_R})} &\leq C_\epsilon \left[\left(\sum_{\substack{\alpha \subset H_v^{n-1} \\ \alpha: \frac{1}{K} \text{- cap}}} \|\widehat{g_\alpha d\sigma}\|_{L^p(w_{B_R})}^p \right)^{1/p} + K^{\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})+\epsilon} \left(\sum_{\substack{\beta \subset H_v^{n-1} \\ \beta: \frac{1}{K^{1/2}} \text{- cap}}} \|\widehat{g_\beta d\sigma}\|_{L^p(w_{B_R})}^p \right)^{1/p} \right] + \\ &+ K^{C_n} C_{p,n,v}(R^{-1}, K^{-n}) \left(\sum_{\substack{\Delta \subset H_v^{n-1} \\ \Delta: \frac{1}{R^{1/2}} \text{- cap}}} \|\widehat{g_\Delta d\sigma}\|_{L^p(w_{B_R})}^p \right)^{1/p} \end{aligned}$$

Proof

We first prove the case $n = 3$ and then indicate the modifications needed for $n \geq 4$.

It is rather immediate that if $Q_1, Q_2, Q_3 \subset [-1/2, 1/2]^2$, the volume of the parallelepiped spanned by the unit normals to H_v^2 at $\pi^{-1}(Q_i)$ is comparable to the area of the triangle $\Delta Q_1 Q_2 Q_3$.

As in [2], we may think of $|\widehat{g_\alpha d\sigma}|$ as being essentially constant on each ball B_K . Denote by $c_\alpha(B_K)$ this value and let α^* be the cap that maximizes it.

The starting point in the argument is the observation in [2] that for each B_K there exists a line $L = L(B_K)$ in the (ξ_1, ξ_2) plane such that if

$$S_L = \{(\xi_1, \xi_2) : \text{dist}((\xi_1, \xi_2), L) \leq \frac{C}{K}\}$$

then for $x \in B_K$

$$\begin{aligned} |\widehat{gd\sigma}(x)| &\leq \\ &C \max_\alpha |\widehat{g_\alpha d\sigma}(x)| + \end{aligned} \quad (13)$$

$$K^4 \max_{\substack{\alpha_1, \alpha_2, \alpha_3 \\ K^{-2}\text{-transverse}}} \left(\prod_{i=1}^3 |\widehat{g_{\alpha_i} d\sigma}(x)| \right)^{1/3} + \quad (14)$$

$$\left| \sum_{\alpha \subset \pi^{-1}(S_L) \cap H_v^2} \widehat{g_{\alpha} d\sigma}(x) \right|. \quad (15)$$

To see this, we distinguish three scenarios. First, if $c_{\alpha}(B_K) \leq K^{-2}c_{\alpha^*}(B_K)$ for each α with $\text{dist}(\pi(\alpha), \pi(\alpha^*)) \geq \frac{10}{K}$, then (13) suffices, as

$$|\widehat{gd\sigma}(x)| \leq \sum_{\alpha} c_{\alpha}(B_K).$$

Otherwise, there is α^{**} with $\text{dist}(\pi(\alpha^{**}), \pi(\alpha^*)) \geq \frac{10}{K}$ and $c_{\alpha^{**}}(B_K) \geq K^{-2}c_{\alpha^*}(B_K)$. The line L is determined by α^*, α^{**} .

Second, if there is α^{***} such that $\pi(\alpha^{***})$ intersects the complement of S_L and $c_{\alpha^{***}}(B_K) \geq K^{-2}c_{\alpha^*}(B_K)$ then (14) suffices. Indeed, note that $\alpha^*, \alpha^{**}, \alpha^{***}$ are K^{-2} transverse.

Otherwise the sum of (13) and (15) will suffice.

The only nontrivial case to address is the one corresponding to this latter scenario. Cover $\pi^{-1}(S_L) \cap H_v^2$ by pairwise disjoint strips U of length $\sim \frac{1}{K^{1/2}}$. An application of the trivial l^p decoupling (10) shows that

$$\left\| \sum_{\alpha: \pi(\alpha) \subset S_L} \widehat{g_{\alpha} d\sigma} \right\|_{L^p(B_K)} \lesssim K^{\frac{1}{2} - \frac{1}{p}} \left(\sum_U \|\widehat{g_U d\sigma}\|_{L^p(w_{B_K})}^p \right)^{1/p}.$$

This is the best we can say in general. Indeed, in the case of the hyperbolic paraboloid $v = (1, -1)$, if the line L happens to be $\xi_2 = \pm \xi_1$ then $\pi^{-1}(L)$ is itself a line. The absence of curvature prevents any non-trivial estimate to hold.

Note however that since we are dealing with the third scenario,

$$\left(\sum_U \|\widehat{g_U d\sigma}\|_{L^p(B_K)}^p \right)^{1/p} \lesssim \left(\sum_{\substack{\beta: \frac{1}{K^{1/2}} \text{ cap} \\ \pi(\beta) \subset S_L}} \|\widehat{g_{\beta} d\sigma}\|_{L^p(w_{B_K})}^p \right)^{1/p} + \|\widehat{g_{\alpha^*} d\sigma}\|_{L^p(w_{B_K})}.$$

We conclude that in either case

$$\begin{aligned} \|\widehat{gd\sigma}\|_{L^p(B_K)} &\lesssim \left[\left(\sum_{\substack{\alpha \subset H_v^2 \\ \alpha: \frac{1}{K} \text{ cap}}} \|\widehat{g_{\alpha} d\sigma}\|_{L^p(w_{B_K})}^p \right)^{1/p} + K^{\frac{1}{2} - \frac{1}{p}} \left(\sum_{\substack{\beta \subset H_v^2 \\ \beta: \frac{1}{K^{1/2}} \text{ cap}}} \|\widehat{g_{\beta} d\sigma}\|_{L^p(w_{B_K})}^p \right)^{1/p} \right] + \\ &+ K^{10} C_{p,3,v}(R^{-1}, K^{-2}) \left(\sum_{\substack{\Delta \subset H_v^2 \\ \Delta: \frac{1}{R^{1/2}} \text{ cap}}} \|\widehat{g_{\Delta} d\sigma}\|_{L^p(w_{B_K})}^p \right)^{1/p}. \end{aligned}$$

Finally, raise to the p^{th} power and sum over $B_K \subset B_R$. Also, the norm $\|\widehat{gd\sigma}\|_{L^p(B_R)}$ can be replaced by the weighted norm $\|\widehat{gd\sigma}\|_{L^p(w_{B_R})}$ via the standard localization argument.

One may repeat this argument in the case $n \geq 4$ as follows. For each B_K there exists a hyperplane $\mathcal{H} = \mathcal{H}(B_K)$ in the $(\xi_1, \dots, \xi_{n-2})$ space such that for $x \in B_K$

$$\begin{aligned} |\widehat{gd\sigma}(x)| &\leq \\ &C \max_{\alpha} |\widehat{g_{\alpha} d\sigma}(x)| + \end{aligned}$$

$$K^{C_n} \max_{\substack{\alpha_1, \dots, \alpha_n \\ K^{-n}\text{-transverse}}} \left(\prod_{i=1}^n |\widehat{g_{\alpha_i} d\sigma}(x)| \right)^{1/n} + \\ \left| \sum_{\alpha \subset \pi^{-1}(S_{\mathcal{H}}) \cap H_v^{n-1}} \widehat{g_{\alpha} d\sigma}(x) \right|.$$

Here

$$S_{\mathcal{H}} = \{(\xi_1, \dots, \xi_{n-1}) : \text{dist}((\xi_1, \dots, \xi_{n-1}), \mathcal{H}) \lesssim \frac{1}{K}\}$$

We only need to explain how to accommodate the previous argument to control the last term. Cover $\pi^{-1}(S_{\mathcal{H}}) \cap H_v^{n-1}$ by pairwise disjoint strips U of dimension $\sim \frac{1}{K^{1/2}} \times \dots \times \frac{1}{K^{1/2}} \times \frac{1}{K}$. These strips are inside the $\frac{1}{K}$ neighborhood of the $n-2$ dimensional manifold

$$S_{\mathcal{H},v} = \{(\xi_1, \dots, \xi_n) \in H_v^{n-1} : (\xi_1, \dots, \xi_{n-1}) \in \mathcal{H}\},$$

and correspond to a tiling of this manifold by $\frac{1}{K^{1/2}}$ -caps. The important new observation is that $S_{\mathcal{H},v}$, regarded as an $n-2$ dimensional hypersurface in the hyperplane

$$\{(\xi_1, \dots, \xi_n) : (\xi_1, \dots, \xi_{n-1}) \in \mathcal{H}\}$$

has at least $n-3$ of its $n-2$ principal curvatures bounded away from zero, at any given point. This is of course a consequence of Lemma 2.5. The case $n=3$ discussed earlier shows that one (in this case the only) principal curvature may indeed happen to be zero. More generally, consider any hyperbolic paraboloid H_v^{n-1} . Fix any A_1, \dots, A_{n-1} such that

$$\sum_{i=1}^{n-1} v_i A_i^2 = 0.$$

Let \mathcal{H} be the hyperplane

$$\sum_{i=1}^{n-1} v_i A_i \xi_i = 0$$

It is easy to check that for each point $(\xi_1^*, \dots, \xi_n^*)$ in the corresponding manifold $S_{\mathcal{H},v}$, the (appropriate part of the) line

$$\frac{\xi_1 - \xi_1^*}{A_1} = \dots = \frac{\xi_{n-1} - \xi_{n-1}^*}{A_{n-1}} = \frac{\xi_n - \xi_n^*}{0}$$

is inside $S_{\mathcal{H},v}$. In other words $S_{\mathcal{H},v}$ is a cylinder, and one of its principal curvatures will be zero.

Using our hypothesis and Lemma 2.4 we can write

$$\left\| \sum_{\alpha \subset \pi^{-1}(S_{\mathcal{H}}) \cap H_v^{n-1}} \widehat{g_{\alpha} d\sigma} \right\|_{L^p(B_K)} \lesssim_{\epsilon} K^{\frac{n-1}{2}(\frac{1}{2} - \frac{1}{p}) + \epsilon} \left(\sum_U \|\widehat{g_U d\sigma}\|_{L^p(w_{B_K})}^p \right)^{1/p}.$$

As before, this can be further bounded by

$$\lesssim K^{\frac{n-1}{2}(\frac{1}{2} - \frac{1}{p}) + \epsilon} \left[\left(\sum_{\beta: \frac{1}{K^{1/2}}\text{-cap}} \|\widehat{g_{\beta} d\sigma}\|_{L^p(w_{B_K})}^p \right)^{1/p} + \left(\sum_{\alpha: \frac{1}{K}\text{-cap}} \|\widehat{g_{\alpha} d\sigma}\|_{L^p(w_{B_K})}^p \right)^{1/p} \right].$$

The argument is now complete. ■

Simple parabolic rescaling leads to the following more general version. The interested reader should consult the proof of the analogous result in [3] for details.

Proposition 2.7. *Fix $n \geq 3$, $v \in \{-1, 1\}^{n-1}$ and let $p \geq 2$. Assume one of the following holds*

(i) $n = 3$

(ii) $n \geq 4$ and $K_{p,n-2,v'}(\delta') \lesssim_\epsilon \delta'^{-\frac{n-3}{2}(\frac{1}{2}-\frac{1}{p})-\epsilon}$ for each $\delta' > 0$, $v' \in \{-1, 1\}^{n-3}$ and each $\epsilon > 0$.

Then for each ϵ there exist constants C_ϵ, C_n such that for each $R > 1$ and $K \geq 1$ and for each δ -cap τ on H_v^{n-1} we have

$$\begin{aligned} \|\widehat{g_\tau d\sigma}\|_{L^p(w_{B_R})} &\leq C_\epsilon \left[\left(\sum_{\substack{\alpha: \frac{\delta}{K} \text{- cap} \\ \alpha: \frac{\delta}{K} \text{- cap}}} \|\widehat{g_\alpha d\sigma}\|_{L^p(w_{B_R})}^p \right)^{1/p} + K^{-\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})+\epsilon} \left(\sum_{\substack{\beta: \frac{\delta}{K^{1/2}} \text{- cap} \\ \beta: \frac{\delta}{K^{1/2}} \text{- cap}}} \|\widehat{g_\beta d\sigma}\|_{L^p(w_{B_R})}^p \right)^{1/p} \right] + \\ &+ K^{C_n} C_{p,n,v}((R\delta^2)^{-1}, K^{-n}) \left(\sum_{\substack{\Delta: \frac{1}{R^{1/2}} \text{- cap} \\ \Delta: \frac{1}{R^{1/2}} \text{- cap}}} \|\widehat{g_\Delta d\sigma}\|_{L^p(w_{B_R})}^p \right)^{1/p} \end{aligned}$$

We are now ready to prove Theorem 2.2. Let $K = \nu^{-1/n}$. We can certainly assume that the quantity $\delta^{\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})} C_{p,n,v}(\delta, \nu)$ is an essentially decreasing function of $\delta > 0$, otherwise the results we aim to prove become trivial. This can be worked out rigorously as in [3] using parabolic rescaling, we leave the details to the reader. In particular, we may assume that

$$\delta^{\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})} C_{p,n,v}(\delta, \nu) \lesssim_\epsilon R^{-\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})+\epsilon} C_{p,n,v}(R^{-1}, \nu) \quad (16)$$

for each $\delta > R^{-1}$ and $\epsilon > 0$.

Iterate Proposition 2.7 starting with caps of scale 1 until all resulting caps have scale $R^{-1/2}$. Each iteration lowers the scale of the caps from δ to at least $\frac{\delta}{K^{1/2}}$. When iteration is over, we end up with a sum of terms of the form

$$T_\Gamma = \Gamma K^{C_n} \left(\sum_{\Delta: \frac{1}{R^{1/2}} \text{- cap}} \|\widehat{g_\Delta d\sigma}\|_{L^p(w_{B_R})}^p \right)^{1/p},$$

with various coefficients Γ . Each such term arises via $\leq \log_K R$ iterations. Also, a crude estimate shows that we end up with at most $3^{\log_K R} = R^{O(\log_{\nu^{-1}} 3)}$ such terms.

It remains to get a uniform upper bound on Γ . Tracing back the iteration history of T_Γ , assume it went through m_1 steps where scale was lowered by K and m_2 steps where scale was lowered by $K^{1/2}$. Then obviously, for each ϵ

$$\Gamma \leq (C_\epsilon)^{m_1+m_2} K^{[\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})+\epsilon]m_2} C_{p,n,v}((RK^{-m_2-2m_1})^{-1}, \nu).$$

Using the bound $m_1 + m_2 \leq \log_K R$ this is further bounded by

$$R^{\log_{\nu^{-1}} C_\epsilon} K^{\epsilon \log_K R} C_{p,n,v}((RK^{-m_2-2m_1})^{-1}, \nu) (RK^{-m_2-2m_1})^{-\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})} R^{\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})}$$

Finally, (16) shows that

$$\Gamma \lesssim_\epsilon R^{\epsilon + \log_{\nu^{-1}} C_\epsilon} C_\epsilon C_{p,n,v}(R^{-1}, \nu).$$

The proof is now complete, by carefully letting ϵ approach zero at slower rate than ν .

We now enter the second and final stage of the argument for Theorem 1.1. For the remainder of the section we fix $v \in \{-1, 1\}^{n-1}$ and $p > \frac{2(n+1)}{n-1}$. To simplify notation we let $K(\delta) = \delta^{\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})} K_{p,n,v}(\delta)$.

As in [3], let

$$\gamma = \liminf_{\delta \rightarrow 0} \frac{\log K(\delta)}{\log(\delta^{-1})}.$$

It follows that for each ϵ

$$\delta^{-\gamma} \lesssim K(\delta) \lesssim_{\epsilon} \delta^{-\gamma-\epsilon}.$$

Write $\gamma = \frac{n-1}{4} - \frac{n+1}{2p} + \alpha$. Recall that we need to prove that $\alpha = 0$.

Define

$$\xi = \frac{2}{(p-2)(n-1)}, \quad \eta = \frac{n(np-2n-p-2)}{2p(n-1)^2(p-2)}. \quad (17)$$

Since $p > \frac{2(n+1)}{n-1}$ we have that $\xi < \frac{1}{2}$. A simple computation reveals that the assumption $\alpha > 0$ is equivalent with

$$\gamma \frac{1-\xi}{1-2\xi} > \frac{n-1}{4} - \frac{n^2+n}{2p(n-1)} + \frac{2\eta}{1-2\xi}.$$

Under this assumption it follows that we can choose $s_0 \in \mathbb{N}$ large enough and ϵ_0 small enough so that

$$\gamma \left(\frac{1-\xi}{1-2\xi} - \frac{\xi(2\xi)^{s_0}}{1-2\xi} \right) > \frac{n-1}{4} - \frac{n^2+n}{2p(n-1)} + 2^{s_0}\epsilon_0 + \frac{2\eta}{1-2\xi}(1-(2\xi)^{s_0}) + \frac{n}{(n-1)p}(2\xi)^{s_0}. \quad (18)$$

Choose $\nu > 0$ small enough such that $\epsilon_0 > \epsilon(\nu)$, with $\epsilon(\nu)$ as in Theorem 2.2. Note that s_0 , ϵ_0 and ν depend only on the fixed parameters p, n, α . As a result, we follow our convention and do not record the dependence on them when using the symbol \lesssim .

Throughout the rest of the section ν , s_0 and ϵ_0 will always refer to these values. Introduce the following semi-norms

$$\begin{aligned} \|f\|_{p,\delta,B} &= \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_{L^p(w_B)}^2 \right)^{1/2}, \\ |||f|||_{p,\delta,B} &= \delta^{-\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})} \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_{L^p(w_B)}^p \right)^{1/p} \end{aligned}$$

and note that

$$\|f\|_{p,\delta,B} \leq |||f|||_{p,\delta,B} \quad (19)$$

For a fixed $0 \leq \beta \leq 1$ consider the inequality

$$\left\| \left(\prod_{i=1}^n |g_i \widehat{d\sigma}| \right)^{1/n} \right\|_{L^p(B_N)} \lesssim_{\epsilon} A_\beta(N) N^\epsilon X(B_N)^{1-\beta} Y(B_N)^\beta, \quad (20)$$

for arbitrary $\epsilon > 0$, N , g_i and B_N as before. Here

$$X(B_N) = \left(\prod_{i=1}^n |||g_i \widehat{d\sigma}|||_{p,\delta,B_N} \right)^{\frac{1}{n}},$$

$$Y(B_N) = \left(\prod_{i=1}^n \|\widehat{g_i d\sigma}\|_{\frac{p(n-1)}{n}, \delta, B_N} \right)^{\frac{1}{n}}.$$

The following holds.

Proposition 2.8. (a) Inequality (20) holds true for $\beta = 1$ with $A_1(N) = N^{\frac{n-1}{4} - \frac{n^2+n}{2p(n-1)}}$.
 (b) Moreover, if we assume (20) for some $\beta \in (0, 1]$, then we also have (20) for $\frac{2\beta}{(p-2)(n-1)}$ with

$$A_{\frac{2\beta}{(p-2)(n-1)}}(N) = A_\beta(N^{1/2}) \delta^{-\frac{\gamma}{2}(1 - \frac{2\beta}{(p-2)(n-1)})} N^{\frac{n(np-2n-p-2)}{2p(n-1)^2(p-2)}\beta}.$$

The proof follows line by line the proof of the analogous Proposition 6.3 in [3]. More precisely, part (i) here follows right away from Proposition 6.3 (i) by simply invoking (19). Also, the only modification needed to prove part (ii) is to notice that the following consequence of Hölder's inequality used in [3]

$$\|\widehat{g_i d\sigma}\|_{\frac{p(n-1)}{n}, \delta, B_N} \leq \|\widehat{g_i d\sigma}\|_{p, \delta, B_N}^{1 - \frac{2}{(p-2)(n-1)}} \|\widehat{g_i d\sigma}\|_{2, \delta, B_N}^{\frac{2}{(p-2)(n-1)}}$$

continues to hold if $\|\cdot\|$ is replaced with $\|\|\cdot\|\|$.

Proposition 2.8 implies that for each $s \geq 0$

$$A_{\xi^s}(N) = N^{\psi(\xi^s)}$$

with

$$\psi(\xi^{s+1}) = \frac{1}{2}\psi(\xi^s) + \frac{\gamma}{2}(1 - \xi^{s+1}) + \eta\xi^s. \quad (21)$$

Recall that $\xi < \frac{1}{2}$. Iterating (21) gives

$$\psi(\xi^s) = \frac{1}{2^s}\psi(1) + \gamma(1 - 2^{-s}) + 2\left(\frac{\eta}{\xi} - \frac{\gamma}{2}\right)\frac{2^{-s} - \xi^s}{\xi^{-1} - 2} \quad (22)$$

Note that $Y(B_N) \lesssim X(B_N)N^{\frac{n}{(n-1)p}}$. As (20) holds for $\beta = \xi^s$ and arbitrary ν -transverse caps τ_i we get

$$\delta^{\frac{n-1}{2}(\frac{1}{2} - \frac{1}{p})} C_{p,n,\nu}(\delta) \lesssim_{\epsilon,s} \delta^{-\epsilon} A_{\xi^s}(N) N^{\frac{n\xi^s}{(n-1)p}}. \quad (23)$$

To finish the proof of Theorem 1.1 for H_v^{n-1} , we will argue using induction on n that $\alpha = 0$. As observed earlier, the case $n = 2$ is covered by the main theorem in [3]. So the first case to consider is $n = 3$. Since (23) (with $s = s_0$) holds for arbitrarily small δ and ϵ , using Theorem 2.2 we get

$$\gamma - \epsilon_0 \leq \psi(\xi^{s_0}) + \frac{n\xi^{s_0}}{(n-1)p}. \quad (24)$$

Combining (22) and (24) we find

$$\gamma\left(\frac{1-\xi}{1-2\xi} - \frac{\xi(2\xi)^{s_0}}{1-2\xi}\right) \leq \psi(1) + 2^{s_0}\epsilon_0 + \frac{2\eta}{1-2\xi}(1 - (2\xi)^{s_0}) + \frac{n}{(n-1)p}(2\xi)^{s_0},$$

which contradicts (18). Thus $\alpha = 0$ and Theorem 1.1 is proved for $n = 3$ and $p > 4$.

Assume now that $n \geq 4$ and that Theorem 1.1 was proved in all dimensions $d \leq n - 1$. To prove Theorem 1.1 in \mathbb{R}^n for $p > \frac{2(n+1)}{n-1}$, it suffices to prove it for $\frac{2(n+1)}{n-1} < p < \frac{2(n-1)}{n-3}$. Note that in this range we have $p < \frac{2(d+1)}{d-1}$ for $d = n - 2$, in particular (9) holds. Thus Theorem 2.2 is applicable due to our induction hypothesis and we reach a contradiction as in the case $n = 3$ discussed above.

It remains to see why Theorem 1.1 holds for the endpoint $p = p_n = \frac{2(n+1)}{n-1}$. Via a localization argument, $K_{p,n,v}(\delta)$ is comparable to the best constant $K_{p,n,v}^*(\delta)$ that makes the following inequality true for each N -ball B_N and each f Fourier supported in \mathcal{N}_δ

$$\|f\|_{L^p(B_N)} \leq K_{p,n,v}^*(\delta) \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_{L^p(\mathbb{R}^n)}^p \right)^{1/p}. \quad (25)$$

It suffices now to invoke Theorem 1.1 for $p > \frac{2(n+1)}{n-1}$ together with

$$\|f\|_{L^{pn}(B_N)} \lesssim \|f\|_{L^p(B_N)} N^{\frac{n}{pn} - \frac{n}{p}} \quad (\text{by Hölder's inequality})$$

$$\|f_\theta\|_{L^p(\mathbb{R}^n)} \lesssim N^{\frac{n+1}{2p} - \frac{n+1}{2pn}} \|f_\theta\|_{L^{pn}(\mathbb{R}^n)} \quad (\text{by Bernstein's inequality}),$$

and then to let $p \rightarrow p_n$.

3. THE PROOF OF THEOREM 1.2

We start this section by explaining the numerology in Theorem 1.2, in particular the origin of the critical index $\frac{2(n+1-d(v))}{n-1-d(v)}$. There are two examples to consider. Example 1 relates to the fact that the essentially sharp estimate

$$|\widehat{d\sigma_{H_v^{n-1}}}(x)| \lesssim (1 + |x|)^{\frac{1-n}{2}}$$

is universal, it does not depend on the signature of H_v^{n-1} . Using f such that \widehat{f} is (a smooth approximation of) the characteristic function of the δ neighborhood of H_v^{n-1} leads to the lower bound

$$K_{n,p,v}^{(2)}(\delta) \gtrsim \delta^{-\frac{n-1}{4} + \frac{n+1}{2p}}, \quad p \geq \frac{2(n+1)}{n-1}, \quad (26)$$

which was shown to be sharp in [3] in the elliptic case.

In the non elliptic case we have that $d(v) \geq 1$ and H_v^{n-1} will contain a compact subset V of an affine subspace of dimension $d(v)$. Example 2 is concerned with the case when \widehat{f} is the characteristic function of the δ neighborhood of V . A standard computation shows that for this f we have

$$\|f\|_p \sim \delta^{d(v)(\frac{1}{2p} - \frac{1}{4})} \left(\sum_{\theta} \|f_\theta\|_p^2 \right)^{1/2}, \quad p \geq 2$$

which leads to

$$K_{n,p,v}^{(2)}(\delta) \gtrsim \delta^{d(v)(-\frac{1}{4} + \frac{1}{2p})}, \quad p \geq 2. \quad (27)$$

Now (26) and (27) suggest that

$$K_{n,p,v}^{(2)}(\delta) \sim \max\left\{ \delta^{-\frac{n-1}{4} + \frac{n+1}{2p}}, \delta^{d(v)(-\frac{1}{4} + \frac{1}{2p})} \right\}, \quad p \geq 2.$$

We will prove that this is indeed correct. Note that there is a regime change precisely at $p = \frac{2(n+1-d(v))}{n-1-d(v)}$.

It is worth mentioning that Example 1 and 2 above also apply (with the correct change in numerology) to the case of l^p decouplings in Theorem 1.1. However, one may check that when $p \geq 2\frac{n+1}{n-1}$ the lower bound coming from Example 1 always dominates the one produced by Example 2.

The proof of Theorem 1.2 follows via induction on the dimension n . The case $n = 2$ was proved in [3]. Assume we have proved the theorem for all H_v^{n-2} with $v \in (\mathbb{R} \setminus \{0\})^{n-2}$, for some fixed $n \geq 3$. Fix now $v \in (\mathbb{R} \setminus \{0\})^{n-1}$. By invoking interpolation with L^2 and L^∞ , it suffices to prove the theorem for H_v^{n-1} when p is greater than but arbitrarily close to the critical index $\frac{2(n+1-d(v))}{n-1-d(v)}$. In particular, it suffices to consider $\frac{2(n+1-d(v))}{n-1-d(v)} < p < \frac{2(n-d(v))}{n-2-d(v)}$.

The guiding principle is that whenever curvature is absent one uses the trivial l^2 decoupling for $2 \leq p < \infty$, which amounts to the following. Given any pairwise disjoint parallelepipeds S_1, \dots, S_M in \mathbb{R}^n we have

$$\|f\|_p \leq M^{\frac{1}{2} - \frac{1}{p}} \left(\sum_{i=1}^M \|f_{S_i}\|_p^2 \right)^{1/2} \quad (28)$$

for each $2 \leq p \leq \infty$ and each f Fourier supported in the union of the S_i . We will refer to this as being *trivial l^2 decoupling*. Note that this is an analogue of (10), which follows again by interpolation.

We now present the main steps in the argument. First, our induction hypothesis implies the following result for hypersurfaces with small (possibly zero) principal curvatures, in the style of Lemma 2.4. More precisely, for a given hypersurface S in \mathbb{R}^{n-1} ($n \geq 3$) with $p(S)$ principal curvatures ≥ 1 , $q(S)$ principal curvatures ≤ -1 and $r(S)$ principal curvatures in $(-1, 1)$, define

$$d(S) = r(S) + \min(p(S), q(S)).$$

Lemma 3.1. *If $\text{supp}(\hat{f}) \subset \mathcal{N}_\delta(S)$ and $2 \leq p \leq \frac{2(n-d(S))}{n-2-d(S)}$ we have*

$$\|f\|_p \lesssim_\epsilon \delta^{-\epsilon} K_{S,p}^{(2)}(\delta) \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_p^2 \right)^{1/2}, \quad (29)$$

where $K_{S,p}^{(2)}(\delta) = \delta^{d(S)(-\frac{1}{4} + \frac{1}{2p})}$.

Proof

Of course, there is nothing special about 1 and -1. This result is about “large” versus “small” principal curvatures. As before, we may assume $S = H_{v'}^{n-2}$ where v' has $p(S)$ entries equal to 1, $q(S)$ entries equal to -1 and $r(S)$ entries in $(-1, 1)$. The case $r(S) = 0$ is an immediate consequence of our induction hypothesis. If $r(S) \geq 1$, perform first a trivial l^2 decoupling in the direction of each of the $r(S)$ “small” principal curvatures. This will contribute a factor of $\delta^{r(S)(-\frac{1}{4} + \frac{1}{2p})}$ to $K_{S,p}^{(2)}(\delta)$, according to (28). Then use the induction hypothesis for the cross sections S' corresponding to the remaining $p(S) + q(S)$ curvatures. Note that $d(S') = \min(p(S), q(S))$ and that S' is a hypersurface with nonzero

Gaussian curvature in $\mathbb{R}^{n-1-r(S)}$. As a result, the critical index for S' is $\frac{2(n-d(S))}{n-2-d(S)}$, and the induction hypothesis is indeed applicable. Thus the cross sections contribute $\delta^{d(S')(-\frac{1}{4}+\frac{1}{2p})}$ to $K_{S,p}^{(2)}(\delta)$. Finally, the two contributions to $K_{S,p}^{(2)}(\delta)$ can be pieced together by invoking Fubini, as explained in the proof of Lemma 2.3. \blacksquare

Next, we record the following analogue of both Proposition 2.6 from here and Proposition 5.5 from [3].

Proposition 3.2. $2 \leq p \leq \frac{2(n-d(v))}{n-2-d(v)}$. For each ϵ there exist constants C_ϵ, C_n such that for each $R > 1$ and $K \geq 1$

$$\begin{aligned} \|\widehat{gd\sigma}\|_{L^p(w_{B_R})} &\leq C_\epsilon \left[\left(\sum_{\substack{\alpha \subset H_v^{n-1} \\ \alpha: \frac{1}{K} - \text{cap}}} \|\widehat{g_\alpha d\sigma}\|_{L^p(w_{B_R})}^2 \right)^{1/2} + K^{d(v)(\frac{1}{2p}-\frac{1}{4})+\epsilon} \left(\sum_{\substack{\beta \subset H_v^{n-1} \\ \beta: \frac{1}{K^{1/2}} - \text{cap}}} \|\widehat{g_\beta d\sigma}\|_{L^p(w_{B_R})}^2 \right)^{1/2} \right] + \\ &+ K^{C_n} C_{p,n,v}^{(2)}(R^{-1}, K^{-n}) \left(\sum_{\substack{\Delta \subset H_v^{n-1} \\ \Delta: \frac{1}{R^{1/2}} - \text{cap}}} \|\widehat{g_\Delta d\sigma}\|_{L^p(w_{B_R})}^2 \right)^{1/2}. \end{aligned}$$

Here, in analogy with earlier notation, $C_{p,n,v}^{(2)}$ is the multilinear version of $K_{p,n,v}^{(2)}$.

Proof This follows as before by using the Bourgain-Guth induction on scales, and our induction hypothesis for arbitrary cross sections S' of H_v^{n-1} with a "vertical" hyperplane. It suffices to prove that $d(S') \leq d(v)$. This follows via an argument similar to the one in Lemma 2.5, we sketch it briefly. Assume for simplicity that $v_i = 1$ for $1 \leq i \leq p(v)$ and $v_i = -1$ for $p(v) + 1 \leq i \leq n - 1$. Assume also that the hyperplane contains the origin, and let H' be its intersection with the hyperplane $x_n = 0$. Call $A : \mathbb{R}^{n-1} \rightarrow \mathbb{R}^{n-1}$ the linear transformation so that H_v^{n-1} is (part of) the graph of $x \mapsto \langle Ax, x \rangle$, $x \in \mathbb{R}^{n-1}$. Obviously v_i are its eigenvalues and $\langle e_i \rangle$ are the corresponding eigenspaces. Define

$$X_+ = \langle e_i : 1 \leq i \leq p(v) \rangle, \quad X_- = \langle e_i : p(v) + 1 \leq i \leq n - 1 \rangle.$$

Similarly, let $A' : H' \rightarrow H'$ be the linear transformation so that S' is (part of) the graph of $x \mapsto \langle A'x, x \rangle$ over $x \in H'$. Let e'_1, \dots, e'_{n-2} be an appropriate orthonormal basis of H' consisting of eigenvectors of A' with eigenvalues v'_i . Assume that $v'_i \geq 1$ for $1 \leq i \leq p(S')$, $v'_i \leq -1$ for $p(S') + 1 \leq i \leq p(S') + q(S')$ and $|v'_i| < 1$ for $p(S') + q(S') + 1 \leq i \leq p(S') + q(S') + r(S') = n - 2$. Define

$$X'_+ = \langle e'_i : 1 \leq i \leq p(S') \rangle, \quad X'_- = \langle e'_i : p(S') + 1 \leq i \leq p(S') + q(S') \rangle,$$

$$X'_0 = \langle e'_i : p(S') + q(S') + 1 \leq i \leq n - 2 \rangle.$$

It suffices now to prove that $p(S') + r(S') \leq p(v)$ and $q(S') + r(S') \leq q(v)$. Assume for contradiction that $p(S') + r(S') > p(v)$. Then $(X'_+ \oplus X'_0) \cap X_-$ must contain a unit vector x . Note first that since $x \in H'$, we have $\langle Ax, x \rangle = \langle A'x, x \rangle$. On the other hand, since $x \in X_-$ we must have $\langle Ax, x \rangle = -1$, while $x \in X'_+ \oplus X'_0$ implies that $\langle Ax, x \rangle > -1$. The contradiction is now obvious. \blacksquare

Iterations of Proposition 3.2 show the equivalence between $C_{p,n,v}^{(2)}$ and $K_{p,n,v}^{(2)}$. Since we work with $p > \frac{2(n+1-d(v))}{n-1-d(v)}$, we also have that $p > \frac{2(n+1)}{n-1}$. In particular, the parameter ξ from (17) is $< \frac{1}{2}$, as desired. The rest of the argument continues exactly as in the elliptic

case from [3], see also the l^p decouplings from the previous section. This is due to the fact that when $p > \frac{2(n+1-d(v))}{n-1-d(v)}$, (2) and (3) are identical.

4. MULTILINEAR VERSUS LINEAR DECOUPLING FOR CURVES

In this section we start the proof of Theorem 1.4. It is easy to see that for each $t_0 \in [0, 1]$, there is an affine transformation L_{t_0} of \mathbb{R}^n , more precisely a rotation followed by a translation, such that

$$\begin{cases} L_{t_0}(\Phi(t_0)) = \mathbf{0} \\ L_{t_0}(\Phi'(t_0)) \perp \langle e_2, \dots, e_n \rangle \\ L_{t_0}(\Phi''(t_0)) \perp \langle e_3, \dots, e_n \rangle \\ \dots\dots\dots \\ L_{t_0}(\Phi^{(n-1)}(t_0)) \perp \langle e_n \rangle \end{cases}$$

In this new local system of coordinates, the equation of the curve near $t = 0$ becomes $\tilde{\Phi}(t) = (C_{1,1}t + C_{1,2}t^2 + \dots + C_{1,n}t^n, C_{2,2}t^2 + \dots + C_{2,n}t^n, \dots, C_{n,n}t^n) + O(t^{n+1}, t^{n+1}, \dots, t^{n+1})$. The coefficients $C_{i,j}$ depend on t_0 but satisfy $\kappa \leq |C_{i,i}| \leq \kappa^{-1}$ and $|C_{i,j}| \leq \kappa^{-1}$ for $i < j$, with $\kappa > 0$ independent of t_0 , due to the Wronskian condition.

By invoking a simple induction on scales argument, it suffices to prove Theorem 1.4 for the special curves $\Phi_{\mathbf{C}}$, $\mathbf{C} = (C_{i,j})_{1 \leq i \leq j \leq n}$

$$\Phi_{\mathbf{C}}(t) = (C_{1,1}t + C_{1,2}t^2 + \dots + C_{1,n}t^n, C_{2,2}t^2 + \dots + C_{2,n}t^n, \dots, C_{n,n}t^n), \quad (30)$$

with $C_{i,j}$ as above. The estimates will of course depend only on κ . To see this, for $\delta < 1$ and $p \leq 4n - 2$, let $K_p^*(\delta)$ be the smallest constant such that for each f with Fourier support in the neighborhood \mathcal{N}_δ of Φ we have

$$\|f\|_{2(n+1)} \leq K_p^*(\delta) \left(\sum_{\theta \in \mathcal{P}_\delta} \|f_\theta\|_p^2 \right)^{\frac{1}{2}}.$$

First, for each such f

$$\|f\|_{2(n+1)} \leq K_p^*(\delta^{\frac{n}{n+1}}) \left(\sum_{\tau \in \mathcal{P}_{\delta^{\frac{n}{n+1}}}} \|f_\tau\|_p^2 \right)^{\frac{1}{2}}. \quad (31)$$

The previous discussion shows that the portion of Φ inside a given $\tau \in \mathcal{P}_{\delta^{\frac{n}{n+1}}}$ is within δ from a curve (30). Let a be the left endpoint of the interval of length $\delta^{\frac{1}{n+1}}$ corresponding to τ . We will perform generalized parabolic rescaling as follows. Consider the linear transformation

$$L_\tau(\xi_1, \dots, \xi_n) = (\xi'_1, \dots, \xi'_n) = \left(\frac{\xi_1 - a}{\delta^{\frac{1}{n+1}}}, \frac{\xi_2 - 2a\xi_1 + a^2}{\delta^{\frac{2}{n+1}}}, \frac{\xi_3 - 3a\xi_2 + 3a^2\xi_1 - a^3}{\delta^{\frac{3}{n+1}}}, \dots \right). \quad (32)$$

It maps $\tau \cap \mathcal{N}_\delta$ into $\mathcal{N}_{\delta^{\frac{1}{n+1}}}$ and each $\theta \in \mathcal{N}_\delta$ with $\theta \subset \tau$ into some $\theta' \in \mathcal{N}_{\delta^{\frac{1}{n+1}}}$. Using this change of variables and Theorem 1.4 with δ replaced with $\delta^{\frac{1}{n+1}}$ we get

$$\|f_\tau\|_p \lesssim_\epsilon \delta^{-\epsilon} \left(\sum_{\theta \in \mathcal{P}_\delta: \theta \subset \tau} \|f_\theta\|_p^2 \right)^{\frac{1}{2}}. \quad (33)$$

For each $\epsilon > 0$, using (31) and (33) we conclude the existence of C_ϵ such that for each $\delta < 1$

$$K_p^*(\delta) \leq C_\epsilon \delta^{-\epsilon} K_p^*(\delta^{\frac{n}{n+1}}).$$

By iteration this immediately leads to $K_p^*(\delta) \lesssim_\epsilon \delta^{-\epsilon}$.

We further observe that it suffices to consider curves $\Phi_{\mathbf{C}}$ with \mathbf{C} equal to the identity matrix. This is because the decoupling inequality is preserved under linear transformations. For the remainder of the section we let

$$\Phi(t) = (t, t^2, \dots, t^n).$$

For each dyadic interval $I \subset [0, 1]$ define the extension operator

$$E_I g(x) = \int_I g(t) e(itx_1 + t^2 x_2 + \dots + t^n x_n) dt.$$

Unless specified otherwise, all intervals will be implicitly assumed to be dyadic intervals in $[0, 1]$.

We will denote by $K_p(R)$ the smallest constant such that

$$\|E_{[0,1]} g\|_{L^p(B_R)} \leq K_p(R) \left(\sum_{|U|=R^{-1/n}} \|E_U g\|_{L^p(w_{B_R})}^2 \right)^{\frac{1}{2}}$$

holds true for each $g : [0, 1] \rightarrow \mathbb{C}$ and B_R .

It is easy to see that $K_p(R) \sim K_p^*(R^{-1})$. In particular, Theorem 1.4 for some p is in fact equivalent with the following inequality

$$\|E_{[0,1]} g\|_{L^p(B_N)} \lesssim_\epsilon N^\epsilon \left(\sum_{|U|=N^{-1/n}} \|E_U g\|_{L^p(w_{B_N})}^2 \right)^{\frac{1}{2}}. \quad (34)$$

For dyadic $0 < \nu < 1$, denote by $C_p(R, \nu)$ the smallest constant such that

$$\left\| \left(\prod_{i=1}^n |E_{I_i} g| \right)^{1/n} \right\|_{L^p(B_R)} \leq C_p(R, \nu) \left(\prod_{i=1}^n \sum_{\substack{U \subset I_i \\ |U|=R^{-1/n}}} \|E_U g\|_{L^p(w_{B_R})}^2 \right)^{\frac{1}{2n}},$$

holds for each nonadjacent dyadic intervals $I_1, \dots, I_n \subset [0, 1]$ of size ν , each $g : [0, 1] \rightarrow \mathbb{C}$ and B_R . We will refer to such intervals as being ν -transverse.

It is immediate that $C_p(R, \nu) \leq K_p(R)$. We now show that the reverse inequality is also essentially true.

Theorem 4.1. *For each ν there exists $\epsilon(\nu)$ with $\lim_{\nu \rightarrow 0} \epsilon(\nu) = 0$ and C_ν such that for each R*

$$K_p(R) \leq C_\nu R^{\epsilon(\nu)} C_p(R, \nu).$$

The proof of the theorem is a simple version of the Bourgain-Guth induction on scales [2]. We will prove a few preliminary results.

Proposition 4.2. *For each R -ball B_R we have*

$$\begin{aligned} \|E_{[0,1]} g\|_{L^p(B_R)} &\leq C_p(R, \nu) C_\nu \left(\sum_{|U|=R^{-1/n}} \|E_U g\|_{L^p(w_{B_R})}^2 \right)^{1/2} + \\ &+ C \left(\sum_{|I|=\nu} \|E_I g\|_{L^p(B_R)}^2 \right)^{1/2}. \end{aligned}$$

The constant C is independent of ν , it only depends on n and p .

Proof We start by writing

$$E_{[0,1]}g = \sum_{I:\nu\text{-interval}} E_I g.$$

It is rather immediate that for each $x \in B_R$

$$\begin{aligned} |E_{[0,1]}g(x)| &\leq C \max_{I:\nu\text{-interval}} |E_I g(x)| + C_\nu \sum_{I_1, \dots, I_n: \nu\text{-transverse}} \left(\prod_{j=1}^n |E_{I_j} g(x)| \right)^{\frac{1}{n}} \leq \\ &C \left(\sum_{I:\nu\text{-interval}} |E_I g(x)|^2 \right)^{1/2} + C_\nu \sum_{I_1, \dots, I_n: \nu\text{-transverse}} \left(\prod_{j=1}^n |E_{I_j} g(x)| \right)^{\frac{1}{n}}. \end{aligned}$$

The result now follows by integrating the p -th power. \blacksquare

We now rescale to get the following version.

Proposition 4.3. *Let H be a δ -interval in $[0, 1]$. For each $R \gtrsim \delta^{-n}$ and each R -ball B_R we have*

$$\begin{aligned} \|E_H g\|_{L^p(B_R)} &\leq C_\nu C_p(R\delta^n, \nu) \left(\sum_{|U|=R^{-1/n}} \|E_U g\|_{L^p(w_{B_R})}^2 \right)^{1/2} + \\ &+ C \left(\sum_{\substack{J:\delta\nu\text{-interval} \\ J \subset H}} \|E_J g\|_{L^p(B_R)}^2 \right)^{1/2}. \end{aligned}$$

Proof Note that if $H = [a, a + \delta]$ then the change of variables $s = \frac{t-a}{\delta}$ shows that

$$|E_H g(x)| = \delta |E_{[0,1]} g^{a,\delta}(x')|$$

where $g^{a,\delta}(s) = g(s\delta + a)$ and $x' = (x'_1, \dots, x'_n)$ with

$$\begin{cases} x'_1 = \delta(x_1 + 2ax_2 + 3a^2x_3 + \dots) \\ x'_2 = \delta^2(x_2 + 3ax_3 + \dots) \\ \dots\dots\dots \\ x'_n = \delta^n x_n \end{cases}.$$

In particular

$$\|E_H g\|_{L^p(B_R)} = \delta^{1 - \frac{n(n+1)}{2p}} \|E_{[0,1]} g^{a,\delta}\|_{L^p(C_R)}$$

where C_R is a $\sim \delta R \times \delta^2 R \times \dots \times \delta^n R$ -cylinder. Cover C_R by a family \mathcal{F} of $O(1)$ -overlapping $\delta^n R$ -balls $B_{\delta^n R}$ and write using Proposition 4.2 and Minkowski's inequality

$$\begin{aligned} \|E_H g\|_{L^p(B_R)} &\lesssim \delta^{1 - \frac{n(n+1)}{2p}} \left(\sum_{B_{\delta^n R} \in \mathcal{F}} \|E_{[0,1]} g^{a,\delta}\|_{L^p(B_{\delta^n R})}^p \right)^{1/p} \leq \\ &C_\nu C_p(\delta^n R, \nu) \delta^{1 - \frac{n(n+1)}{2p}} \left(\sum_{B_{\delta^n R} \in \mathcal{F}} \left(\sum_{|U|=(\delta^n R)^{-1/n}} \|E_U g^{a,\delta}\|_{L^p(w_{B_{\delta^n R}})}^2 \right)^{p/2} \right)^{1/p} + \\ &+ C \delta^{1 - \frac{n(n+1)}{2p}} \left(\sum_{B_{\delta^n R} \in \mathcal{F}} \left(\sum_{|I|=\nu} \|E_I g^{a,\delta}\|_{L^p(B_{\delta^n R})}^2 \right)^{p/2} \right)^{1/p} \lesssim \\ &C_\nu C_p(\delta^n R, \nu) \delta^{1 - \frac{n(n+1)}{2p}} \left(\sum_{|U|=(\delta^n R)^{-1/n}} \|E_U g^{a,\delta}\|_{L^p(w_{C_R})}^2 \right)^{1/2} + \end{aligned}$$

$$+C\delta^{1-\frac{n(n+1)}{2p}}\left(\sum_{|I|=\nu}\|E_I g^{a,\delta}\|_{L^p(w_{C_R})}^2\right)^{1/2}.$$

Changing back to the original variables gives us the desired estimate. \blacksquare

We are now ready to prove Theorem 4.1. Given $\epsilon > 0$, choose ν so that $\log_{\nu^{-1}} C \ll n\epsilon$, where C is the constant in Proposition 4.3. Iterate Proposition 4.3 starting with scale $\delta = 1$ until we reach scale $\delta = R^{-1/n}$. Each iteration lowers the scale of the intervals from δ to $\delta\nu$. We thus have to iterate $\frac{\log_{\nu^{-1}} R}{n}$ times. In particular

$$\begin{aligned}\|E_{[0,1]}g\|_{L^p(B_R)} &\lesssim \sum_{s=0}^{\frac{\log_{\nu^{-1}} R}{n}} C_\nu C^s C_p(R\nu^{ns}, \nu) \left(\sum_{|U|=R^{-1/n}} \|E_U g\|_{L^p(w_{B_R})}^2\right)^{1/2} \\ &\leq R^\epsilon C_\nu C_p(R, \nu) \left(\sum_{|U|=R^{-1/n}} \|E_U g\|_{L^p(w_{B_R})}^2\right)^{1/2},\end{aligned}$$

since $C_p(\cdot, \nu)$ is an essentially nondecreasing function.

5. THE l^2 DECOUPLING FOR CURVES

To prove (34) we develop a variant of the argument for hypersurfaces from [3]. That argument relied on the deep multilinear theorem of Bennett, Carbery and Tao [1]. Our proof for curves will use the following analogous, but much simpler result, see for example Lemma 2.5 in [8].

Proposition 5.1. *Let $I_1, \dots, I_n \subset [0, 1]$ be ν -transverse and let $g : [0, 1] \rightarrow \mathbb{C}$. Then*

$$\left\|\left(\prod_{i=1}^n |E_{I_i} g|\right)^{1/n}\right\|_{L^{2n}(\mathbb{R}^n)} \lesssim_\nu \left(\prod_{i=1}^n \|g\|_{L^2(I_i)}\right)^{1/n} \quad (35)$$

The whole argument consists of changing variables

$$(t_1, \dots, t_n) \in I_1 \times \dots \times I_n \mapsto (\Phi(t_1) + \dots + \Phi(t_n)) \in \mathbb{R}^n$$

followed by Plancherel's inequality $\|\widehat{G}\|_2 \lesssim \|G\|_2$. An immediate consequence is the fact that given any f_i Fourier supported in a δ -neighborhood of $\Phi(I_i)$, we have

$$\left\|\left(\prod_{i=1}^n |f_i|\right)^{1/n}\right\|_{L^{2n}(\mathbb{R}^n)} \lesssim_\nu N^{-\frac{n-1}{2}} \left(\prod_{i=1}^n \|f_i\|_{L^2(\mathbb{R}^n)}\right)^{1/n}. \quad (36)$$

Indeed, foliate the δ neighborhood \mathcal{N}_δ of $\Phi([0, 1])$ into translates of the curve $\Phi([0, 1])$, apply (35) to each of them and use Hölder's inequality.

Another immediate corollary of (35) is the following multilinear version of the decoupling inequality (34), that follows by simply invoking L^2 orthogonality

$$\left\|\left(\prod_{i=1}^n |E_{I_i} g|\right)^{1/n}\right\|_{L^{2n}(B_N)} \lesssim_\nu \left\|\left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=N^{-1}}} |E_J g|^2\right)^{\frac{1}{2n}}\right\|_{L^{2n}(w_{B_N})} \quad (37)$$

There are two ways in which (37) is stronger than (34). First, note that multilinear decoupling holds at frequency scales as small as N^{-1} . Second, note that by Minkowski's inequality

$$\left\| \left(\sum_J |h_J|^2 \right)^{1/2} \right\|_p \leq \left(\sum_J \|h_J\|_p^2 \right)^{1/2},$$

which makes the right hand side of (37) smaller than the right hand side of (34).

An application of the Bourgain-Guth induction on scales described in the end of the previous section shows that (37) implies (34) for $2 \leq p \leq 2n$. In the remaining part of this section we will show how to bridge the gap between $p = 2n$ and $p = 4n - 2$.

The first step in our iteration scheme is the following lemma. We will consider a partition of \mathcal{N}_δ into tubular regions τ , each of which is a δ -neighborhood of $\Phi(J)$, for some interval J of length $\delta^{1/2}$. The scale $\delta^{1/2}$ we use here is much smaller than $\delta^{1/n}$, and in particular ensures that τ is essentially a $\delta^{1/2} \times \delta \times \dots \times \delta$ cylinder. This will in turn allow for wave packet decompositions to come into play.

We will use the notation $\delta = N^{-1}$.

Lemma 5.2. *Let $I_1, \dots, I_n \subset [0, 1]$ be ν -transverse intervals, and assume f_i are Fourier supported in δ -neighborhoods of $\Phi(I_i)$. Then for each $2n \leq p \leq \infty$*

$$\left\| \left(\prod_{i=1}^n \left(\sum_{\tau} |f_{i,\tau}|^2 \right)^{1/2} \right)^{1/n} \right\|_{L^p(\mathbb{R}^n)} \lesssim_\nu N^{-\frac{n(n-1)}{p}} \left(\prod_{i=1}^n \sum_{\tau} \|f_{i,\tau}\|_{L^{p/n}(\mathbb{R}^n)}^2 \right)^{\frac{1}{2n}}. \quad (38)$$

Proof Note first that (36) implies via a standard randomization argument that

$$\left\| \left(\prod_{i=1}^n \sum_{\tau} |f_{i,\tau}|^2 \right)^{\frac{1}{2n}} \right\|_{L^{2n}(\mathbb{R}^n)} \lesssim_\nu N^{-\frac{n-1}{2}} \left(\prod_{i=1}^n \sum_{\tau} \|f_{i,\tau}\|_{L^2(\mathbb{R}^n)}^2 \right)^{\frac{1}{2n}}. \quad (39)$$

We also have the trivial inequality

$$\left\| \left(\prod_{i=1}^n \sum_{\tau} |f_{i,\tau}|^2 \right)^{\frac{1}{2n}} \right\|_{L^\infty(\mathbb{R}^n)} \lesssim \left(\prod_{i=1}^n \sum_{\tau} \|f_{i,\tau}\|_{L^\infty(\mathbb{R}^n)}^2 \right)^{\frac{1}{2n}}. \quad (40)$$

The result now follows using interpolation via wave packet decompositions, see for example the proof of Proposition 6.2 in [3]. \blacksquare

Given $g : [0, 1] \rightarrow \mathbb{C}$, there is an immediate reformulation of the lemma: for each $2n \leq p \leq \infty$ and each ball B_N of radius $N = \delta^{-1}$ we have

$$\left\| \left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=\delta^{1/2}}} |E_J g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_{B_N})} \lesssim_\nu N^{-\frac{n(n-1)}{p}} \left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=\delta^{1/2}}} \|E_J g\|_{L^{p/n}(w_{B_N})}^2 \right)^{\frac{1}{2n}}. \quad (41)$$

Let now Δ be an arbitrary ball of radius $N^{1/2}$. Write using Hölder's inequality

$$\left(\sum_{|J'|=\delta^{1/4}} \|E_{J'} g\|_{L^{p/n}(w_\Delta)}^2 \right)^{\frac{1}{2}} \leq \left(\sum_{|J'|=\delta^{1/4}} \|E_{J'} g\|_{L^2(w_\Delta)}^2 \right)^{\frac{1-\kappa}{2}} \left(\sum_{|J'|=\delta^{1/4}} \|E_{J'} g\|_{L^p(w_\Delta)}^2 \right)^{\frac{\kappa}{2}}. \quad (42)$$

where

$$\frac{n}{p} = \frac{1-\kappa}{2} + \frac{\kappa}{p}.$$

The next key element in our argument is the almost orthogonality specific to L^2 , which will allow us to pass from scale $\delta^{1/4}$ to scale $\delta^{1/2}$. Indeed, since $(E_J g)w_\Delta$ are almost orthogonal for $|J| = \delta^{1/2}$, we have

$$\left(\sum_{|J'|=\delta^{1/4}} \|E_{J'} g\|_{L^2(w_\Delta)}^2 \right)^{1/2} \lesssim \left(\sum_{|J|=\delta^{1/2}} \|E_J g\|_{L^2(w_\Delta)}^2 \right)^{1/2}.$$

We can now rely on the fact that $|E_J g|$ is essentially constant on balls Δ' of radius $N^{1/2}$ to argue that

$$\left(\sum_{\substack{J \subset I_i \\ |J|=\delta^{1/2}}} \|E_J g\|_{L^2(\Delta')}^2 \right)^{\frac{1}{2}} \sim |\Delta'|^{1/2} \left(\sum_{\substack{J \subset I_i \\ |J|=\delta^{1/2}}} |E_J g|^2 \right)^{\frac{1}{2}}_{|\Delta'}$$

and thus

$$\left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=\delta^{1/2}}} \|E_J g\|_{L^2(w_\Delta)}^2 \right)^{\frac{1}{2n}} \lesssim |\Delta|^{\frac{1}{2} - \frac{1}{p}} \left\| \left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=\delta^{1/2}}} |E_J g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_\Delta)}. \quad (43)$$

Combining (41), (42) and (43) we get

$$\left\| \left(\prod_{i=1}^n \sum_{\substack{J' \subset I_i \\ |J'|=\delta^{1/4}}} |E_{J'} g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_\Delta)} \lesssim_\nu \left\| \left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=\delta^{1/2}}} |E_J g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_\Delta)}^{1-\kappa} \left(\prod_{i=1}^n \sum_{\substack{J' \subset I_i \\ |J'|=\delta^{1/4}}} \|E_{J'} g\|_{L^p(w_\Delta)}^2 \right)^{\frac{\kappa}{2n}}.$$

Summing this up over $\Delta \subset B_N$ we get

$$\begin{aligned} & \left\| \left(\prod_{i=1}^n \sum_{\substack{J' \subset I_i \\ |J'|=\delta^{1/4}}} |E_{J'} g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_{B_N})} \leq \\ & \leq C \left\| \left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=\delta^{1/2}}} |E_J g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_{B_N})}^{1-\kappa} \left(\prod_{i=1}^n \sum_{\substack{J' \subset I_i \\ |J'|=\delta^{1/4}}} \|E_{J'} g\|_{L^p(w_{B_N})}^2 \right)^{\frac{\kappa}{2n}}, \end{aligned} \quad (44)$$

where C depends only on n , ν and p . By iterating this inequality we get for integers $s \geq 2$

$$\begin{aligned} & \left\| \left(\prod_{i=1}^n \sum_{\substack{J_s \subset I_i \\ |J_s|=\delta^{2-s}}} |E_{J_s} g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_{B_N})} \leq \\ & \leq C^{s-1} \left\| \left(\prod_{i=1}^n \sum_{\substack{J_1 \subset I_i \\ |J_1|=\delta^{1/2}}} |E_{J_1} g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_{B_N})}^{(1-\kappa)^{s-1}} \prod_{j=2}^s \left(\prod_{i=1}^n \sum_{\substack{J_j \subset I_i \\ |J_j|=\delta^{2-j}}} \|E_{J_j} g\|_{L^p(w_{B_N})}^2 \right)^{\frac{\kappa(1-\kappa)^{s-j}}{2n}}. \end{aligned} \quad (45)$$

We next observe the following result.

Lemma 5.3. *For $2n \leq p \leq \infty$ and $s \geq 2$*

$$\left\| \left(\prod_{i=1}^n |E_{I_i} g| \right)^{1/n} \right\|_{L^p(B_N)} \lesssim_\nu N^{2-s(\frac{1}{2} - \frac{n}{p})} \left\| \left(\prod_{i=1}^n \sum_{\substack{J_s \subset I_i \\ |J_s|=\delta^{2-s}}} |E_{J_s} g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_{B_N})},$$

with the implicit constant independent of N and s .

Proof Let Δ_{s-1} be any ball with radius N^{2-s+1} . By interpolating the inequality (consequence of (37))

$$\left\| \left(\prod_{i=1}^n |E_{I_i} g| \right)^{1/n} \right\|_{L^{2n}(\Delta_{s-1})} \lesssim_{\nu} \left\| \left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=N^{-s}}} |E_J g|^2 \right)^{\frac{1}{2n}} \right\|_{L^{2n}(w_{\Delta_{s-1}})}$$

with the trivial inequality

$$\left\| \left(\prod_{i=1}^n |E_{I_i} g| \right)^{1/n} \right\|_{L^{\infty}(\Delta_{s-1})} \lesssim N^{2-s-1} \left\| \left(\prod_{i=1}^n \sum_{\substack{J \subset I_i \\ |J|=N^{-s}}} |E_J g|^2 \right)^{\frac{1}{2n}} \right\|_{L^{\infty}(w_{\Delta_{s-1}})}$$

we get that

$$\left\| \left(\prod_{i=1}^n |E_{I_i} g| \right)^{1/n} \right\|_{L^p(\Delta_{s-1})} \lesssim_{\nu} N^{2-s(\frac{1}{2}-\frac{n}{p})} \left\| \left(\prod_{i=1}^n \sum_{\substack{J_s \subset I_i \\ |J_s|=\delta^{2-s}}} |E_{J_s} g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_{\Delta_{s-1}})}.$$

Interpolation is again possible due to the availability of wave packet decompositions in this context. Finally, it suffices to sum over $\Delta_{s-1} \subset B_N$. \blacksquare

Generalized parabolic rescaling (32) shows that for each $N^{-\rho}$ interval J with $\rho \leq \frac{1}{n}$ we have

$$\|E_J g\|_{L^p(B_N)} \leq K_p(N^{1-\rho n}) \left(\sum_{\substack{I \subset J \\ |I|=N^{-1/n}}} \|E_I g\|_{L^p(w_{B_N})} \right)^{1/2}.$$

We will apply this rescaling to the terms in (45) with $2^{-j} \leq \frac{1}{n}$. There will be $O_n(1)$ terms in (45) for which this procedure is not applicable. However, we will be content with the trivial estimates

$$\left(\prod_{i=1}^n \sum_{\substack{J_j \subset I_i \\ |J_j|=\delta^{2-j}}} \|E_{J_j} g\|_{L^p(w_{B_N})}^2 \right)^{\frac{1}{2n}} \leq N^A \left(\prod_{i=1}^n \sum_{\substack{U \subset I_i \\ |U|=\delta^{1/n}}} \|E_U g\|_{L^p(w_{B_N})}^2 \right)^{\frac{1}{2n}},$$

for $2^{-j} > \frac{1}{n}$, and

$$\left\| \left(\prod_{i=1}^n \sum_{\substack{J_1 \subset I_i \\ |J_1|=\delta^{1/2}}} |E_{J_1} g|^2 \right)^{\frac{1}{2n}} \right\|_{L^p(w_{B_N})} \leq N^A \left(\prod_{i=1}^n \sum_{\substack{U \subset I_i \\ |U|=\delta^{1/n}}} \|E_U g\|_{L^p(w_{B_N})}^2 \right)^{\frac{1}{2n}}.$$

The value of A will not be important for our computations.

Combining these estimates with (45) and Lemma 5.3 we have for ν -transverse I_i

$$\left\| \left(\prod_{i=1}^n |E_{I_i} g| \right)^{1/n} \right\|_{L^p(B_N)} \lesssim_{\nu} C^{s-1} \left(\prod_{i=1}^n \sum_{\substack{U \subset I_i \\ |U|=\delta^{1/n}}} \|E_U g\|_{L^p(w_{B_N})}^2 \right)^{\frac{1}{2n}} \times$$

$$\times N^{2-s(\frac{1}{2}-\frac{n}{p})} K_p(N^{1-2^{-s}n})^{\kappa} K_p(N^{1-2^{-s+1}n})^{\kappa(1-\kappa)} \dots K_p(N^{1-2^{-j_0}n})^{\kappa(1-\kappa)^{s-j_0}} N^{O_n((1-\kappa)^s)},$$

where $2^{-j_0} \leq \frac{1}{n} < 2^{-j_0+1}$.

This in turn implies that

$$C_p(N, \nu) \lesssim_{\nu}$$

$$C^{s-1} N^{2^{-s}(\frac{1}{2} - \frac{n}{p})} K_p(N^{1-2^{-s}n})^\kappa K_p(N^{1-2^{-s+1}n})^{\kappa(1-\kappa)} \dots K_p(N^{1-2^{-j_0}n})^{\kappa(1-\kappa)^{s-j_0}} N^{O_n((1-\kappa)^s)}.$$

Assume $K_p(N) \sim N^{\gamma_p}$. This can be made rigorous, see for example Section 6 in [3]. In light of Theorem 4.1 we have

$$\gamma_p \leq 2^{-s} \left(\frac{1}{2} - \frac{n}{p} \right) + \kappa \gamma_p \left(\frac{1 - (1 - \kappa)^{s-j_0+1}}{\kappa} - n 2^{-s} \frac{1 - (2(1 - \kappa))^{s-j_0+1}}{2\kappa - 1} \right) + O_n((1 - \kappa)^s).$$

If $p > 4n - 2$ then $2(1 - \kappa) = \frac{4(n-1)}{p-2} < 1$ and we can let $s \rightarrow \infty$ to get

$$\gamma_p \leq \frac{p - 4n + 2}{2np}.$$

This is of course good enough to conclude the proof of Theorem 1.4 for $p = 4n - 2$. To get the range $2 \leq p < 4n - 2$, we caution that there is no interpolation argument available in this context (see the counterexample in the last section). As explained earlier, this is due to the fact that the neighborhoods $\theta \in \mathcal{P}_\delta$ do not have cylindrical shape. There is however at least one way around this. Namely, note that (44) holds with $\kappa = 1$ (Minkowski's inequality), hence for each $\frac{p-2n}{p-2} = \kappa_p \leq \kappa \leq 1$. Thus, if $2n < p < p_0 = 4n - 2$, one may replace κ_p with $\kappa_{p_0+\epsilon}$, and get Theorem 1.4 in this range, too. Finally, recall that the result for $2 \leq p \leq 2n$ follows from (37).

6. A MORE GENERAL PERSPECTIVE ON DECOUPLINGS FOR CURVES

Recall the definition of the extension operator

$$E_I g(x) = \int_I g(t) e(itx_1 + t^2 x_2 + \dots + t^n x_n) dt.$$

We denote by $\|F\|_{L^p_\sharp(B)} = (\frac{1}{|B|} \int_{\mathbb{R}^n} |F|^p w_B)^{1/p}$ the normalized integral. Given $2 \leq q \leq p < \infty$, $1 \leq m \leq n$ and $\frac{1}{n} \leq \alpha \leq 1$, we will say that we have (p, q, α, m) decoupling if

$$\|(\prod_{i=1}^m |E_{I_i} g|)^{1/m}\|_{L^p_\sharp(B_N)} \lesssim_\epsilon N^{\alpha(\frac{1}{2} - \frac{1}{q}) + \epsilon} \left(\prod_{i=1}^m \sum_{\substack{|U|=N^{-\alpha} \\ U \subset I_i}} \|E_U g\|_{L^q_\sharp(B_N)}^q \right)^{\frac{1}{mq}}$$

holds true for each $g : [0, 1] \rightarrow \mathbb{C}$, each N -ball $B_N \subset \mathbb{R}^n$ and each ~ 1 -transverse intervals $I_i \subset [0, 1]$.

A few comments are appropriate. First, (p, q, α, m) decoupling implies (p', q', α, m') decoupling when $p' \leq p$, $q \leq q'$, $m' \geq m$. Also, (p, p, α, m) decoupling implies (p, p, α', m) decoupling when $\alpha' \leq \alpha$. It thus follows that (p, q, α, m) decoupling is a bit stronger than (p, p, α, m) decoupling, which was referred to in the earlier part of this paper as m -linear l^p decoupling at frequency scale $N^{-\alpha}$. In particular, (p, q, α, m) decoupling implies the following estimate for exponential sums: for each δ -separated set Λ of points on the curve $\Phi([0, 1])$ and each coefficients $a_\xi \in \mathbb{C}$ we have

$$\left(\frac{1}{|B_R|} \int_{B_R} \left| \prod_{i=1}^m \sum_{\xi \in \Lambda \cap \Phi(I_i)} a_\xi e(\xi \cdot x) \right|^{p/m} dx \right)^{1/p} \lesssim_\epsilon \delta^{\frac{1}{2} - \frac{1}{p} - \epsilon} \|a_\xi\|_{l^p(\Lambda)}, \quad (46)$$

for each ϵ and each ball $B_R \subset \mathbb{R}^n$ of radius $R \gtrsim \delta^{-\frac{1}{\alpha}}$.

We have repeatedly used throughout the paper the fact that one can interpolate decouplings when $\alpha = \frac{1}{2}$. Let us now see why interpolation fails when $\alpha = \frac{1}{n}$. Recall that we have both $(4n - 2, 4n - 2, \frac{1}{n}, n)$ and $(2n, 2, \frac{1}{n}, n)$ decoupling. If interpolation held true, this would give $(3n, 6, \frac{1}{n}, n)$ decoupling, namely

$$\left\| \left(\prod_{i=1}^n |E_{I_i} g| \right)^{1/n} \right\|_{L_{\sharp}^{3n}(B_N)} \lesssim_{\epsilon} N^{\frac{1}{3n} + \epsilon} \left(\prod_{i=1}^n \sum_{\substack{|U|=N-\frac{1}{n} \\ U \subset I_i}} \|E_U g\|_{L_{\sharp}^6(B_N)} \right)^{\frac{1}{6n}}.$$

Note however that this is false for n large enough. Indeed, when $g = 1_{[0,1]}$ the left hand side is greater than

$$N^{-1/3} \left\| \left(\prod_{i=1}^n |E_{I_i} g| \right)^{1/n} \right\|_{L^{3n}(B_1)} \gtrsim N^{-1/3}.$$

On the other hand, a stationary phase computation (see for example [6]) shows that if $n \geq 3$ then

$$\|E_U g\|_{L_{\sharp}^6(B_N)} \lesssim \|E_{[0,1]} g\|_{L_{\sharp}^6(B_N)} \lesssim_{\epsilon} N^{-\frac{4}{9} + \epsilon}$$

Given $\frac{1}{n} \leq \alpha \leq 1$, let $l = l(\alpha)$ be such that $\frac{1}{l+1} < \alpha \leq \frac{1}{l}$. If $\alpha < \frac{1}{n}$ we set $l(\alpha) = n$. The following simple example shows that there can be no (p, q, α, m) decoupling for $p > 2\frac{n-l}{\alpha} + l(l+1)$. Indeed, such a decoupling would imply that

$$\left(\frac{1}{N^n} \int_{[0, N]^n} \left| \prod_{i=1}^m \sum_{k \in I_i} e\left(\frac{k}{N^\alpha} x_1 + \dots + \left(\frac{k}{N^\alpha}\right)^n x_n\right) \right|^{p/m} dx_1 \dots dx_n \right)^{\frac{1}{p}} \lesssim_{\epsilon} N^{\frac{\alpha}{2} + \epsilon}.$$

To see why this leads to a contradiction, consider integers $M_1, \dots, M_l \geq 1$ such that $M_j N^{j\alpha} \leq N < (M_j + 1)N^{j\alpha}$ for $1 \leq j \leq l$.

By periodicity, the left hand side is greater than

$$\begin{aligned} & \left(\frac{1}{N^n} \int_{\prod_{j=1}^l [0, M_j N^{j\alpha}] \times [0, N]^{n-l}} \left| \prod_{i=1}^m \sum_{\frac{k}{N^\alpha} \in I_i} e\left(\frac{k}{N^\alpha} x_1 + \dots + \left(\frac{k}{N^\alpha}\right)^n x_n\right) \right|^{p/m} dx_1 \dots dx_n \right)^{\frac{1}{p}} = \\ & \left(\frac{\prod_{j=1}^l M_j}{N^n} \int_{\prod_{j=1}^l [0, N^{j\alpha}] \times [0, N]^{n-l}} \left| \prod_{i=1}^m \sum_{\frac{k}{N^\alpha} \in I_i} e\left(\frac{k}{N^\alpha} x_1 + \dots + \left(\frac{k}{N^\alpha}\right)^n x_n\right) \right|^{p/m} dx_1 \dots dx_n \right)^{\frac{1}{p}} \gtrsim \\ & \left(\frac{N^l}{N^{n+\alpha+\dots+l\alpha}} \min_{(x_1, \dots, x_n) \in [0, \frac{1}{100}]^n} \left| \prod_{i=1}^m \sum_{\frac{k}{N^\alpha} \in I_i} e\left(\frac{k}{N^\alpha} x_1 + \dots + \left(\frac{k}{N^\alpha}\right)^n x_n\right) \right|^{p/m} \right)^{\frac{1}{p}} \gtrsim \\ & N^{\alpha + \frac{1}{p} \left(l - \frac{\alpha l(l+1)}{2} - n \right)}. \end{aligned}$$

Let us now take a look at some examples. First, the case $\alpha = 1$. Inequality (37) shows that we have $(p, 2, 1, n)$ decoupling for $2 \leq p \leq 2n$. The example above shows that there can be no $(p, q, 1, m)$ decoupling for $p > 2n$.

The value $\alpha = \frac{1}{n}$ corresponds to the natural scaling of the curve, and it is thus naturally associated with linear decoupling ($m = 1$). Our Theorem 1.4 here implies that we have

$(4n-2, 4n-2, \frac{1}{n}, 1)$ decoupling. It seems possible that $(p, p, \frac{1}{n}, 1)$ decoupling would hold for p as large as $n(n+1)$ and the example above shows that one can not hope for a larger p . If true, this would imply Vinogradov's mean value theorem.

The case $\alpha = \frac{1}{2}$ is particularly interesting. It seems reasonable to expect $(4n-2, 4n-2, \frac{1}{2}, m)$ decoupling for some $m > 1$, at least for $m = n$. The exponent $4n-2$ is again suggested by the example above but also by interpolation heuristics. In [5], the first author proves the $(3n, 6, \frac{1}{2}, \frac{n}{2})$ decoupling when n is even. This will be the key step in improving the upper bound on the Riemann zeta function on the critical line.

As a last example for $\alpha = \frac{1}{2}$, we will show below how our Theorem 1.1 here implies $(2(n+1), \frac{2(n+1)}{n-1}, \frac{1}{2}, n-1)$ decoupling. It is interesting to observe that the reciprocals of $(2n, 2)$, $(3n, 6)$, $(2(n+1), \frac{2(n+1)}{n-1})$, $(4n-2, 4n-2)$ are collinear. This is consistent with the fact that interpolation is available in the case $\alpha = \frac{1}{2}$.

Proposition 6.1. *Let I_1, \dots, I_{n-1} be ~ 1 -transverse intervals. Then we have*

$$\left\| \left(\prod_{j=1}^{n-1} |E_{I_j} g| \right)^{\frac{1}{n-1}} \right\|_{L^2_{\sharp}(B_N)^{2(n+1)}} \lesssim_{\epsilon} N^{\frac{1}{2(n+1)} + \epsilon} \left(\prod_{i=1}^{n-1} \sum_{\substack{\Delta: \frac{1}{N^{1/2}}\text{-interval} \\ \Delta \subset I_i}} \|E_{\Delta} g\|_{L^2_{\sharp} \left(\frac{2(n+1)}{n-1} \right) (B_N)} \right)^{\frac{1}{2(n+1)}}.$$

Proof The first step is to construct a hypersurface S from the curve Φ , and invoke Theorem 1.1 for it. That leads to the following inequality proved in [4]

$$\left\| \left(\prod_{j=1}^{n-1} |f_j| \right)^{\frac{1}{n-1}} \right\|_{L^2(B_N)^{2(n+1)}} \lesssim_{\epsilon} N^{\frac{1}{2(n+1)} + \epsilon} \left(\sum_{\substack{\Delta_i: \frac{1}{N^{1/2}}\text{-interval} \\ \Delta_i \subset I_i}} \left\| \prod_{i=1}^{n-1} f_{i, \Delta_i} \right\|_{L^2 \left(\frac{2(n+1)}{n-1} \right) (w_{B_N})} \right)^{\frac{1}{2(n+1)}}, \quad (47)$$

whenever f_i is Fourier supported in the $\frac{1}{N}$ -neighborhood of $\Phi(I_i)$ and f_{i, Δ_i} denotes the Fourier restriction of f_i to the $\frac{1}{N}$ -neighborhood of $\Phi(\Delta_i)$. It is worth noting that when $n = 3$, the hypersurface S is

$$\xi_3 = -\frac{\xi_1^3}{2} + 3\xi_1\xi_2.$$

The principal curvatures are $-\frac{3}{2}(\xi_1 \pm \sqrt{\xi_1^2 + 1})$ and the second fundamental form of S is not definite. This example explains the need for our main Theorem 1.1 in this paper.

The second step of the argument consists of exploiting multilinearity on the right hand side of (47). To avoid technicalities we will provide an informal argument. Using wave packet decompositions, we note that for each q

$$|f_{\Delta}|^q \sim \sum_{T \cap B_N \neq \emptyset} c_T N^{-\frac{1}{2} - n} 1_T \text{ on } B_N,$$

where T are parallel $N \times N \dots \times N \times N^{1/2}$ -plates dual to the $\frac{1}{N}$ -neighborhood of $\Phi(\Delta)$ and

$$\sum c_T \sim \int_{B_N} |f_{\Delta}|^q.$$

Using the key estimate

$$\int_{B_N} 1_{T_1} \dots 1_{T_{n-1}} \lesssim N^{\frac{n+1}{2}}$$

for transverse plates T_1, \dots, T_{n-1} , we obtain (with $q = 2\frac{n+1}{n-1}$)

$$\int_{B_N} \prod_{i=1}^{n-1} f_{i,\Delta_i} \lesssim N^{-(n-\frac{1}{2})(n-1)+\frac{n+1}{2}} \prod_{i=1}^{n-1} \|f_{i,\Delta_i}\|_{L^{2\frac{n+1}{n-1}}(w_{B_N})}^{2\frac{n+1}{n-1}}.$$

This finishes the proof. ■

REFERENCES

- [1] Bennett, J., Carbery, A. and Tao, T. *On the multilinear restriction and Kakeya conjectures*, Acta Math. 196 (2006), no. 2, 261-302
- [2] Bourgain, J. and Guth, L. *Bounds on oscillatory integral operators based on multilinear estimates*, Geom. Funct. Anal. 21 (2011), no. 6, 1239-1295
- [3] Bourgain, J. and Demeter, C. *The proof of the l^2 Decoupling Conjecture*, preprint available on arxiv.
- [4] Bourgain, J. *Decoupling inequalities and some mean-value theorems*, preprint available on arxiv.
- [5] Bourgain, J. *Decoupling, exponential sums and the Riemann-zeta function*, preprint available on arxiv.
- [6] Brandolini, L.; Gigante, G.; Greenleaf, A.; Iosevich, A.; Seeger, A.; Travaglini, G. *Average decay estimates for Fourier transforms of measures supported on curves*. J. Geom. Anal. 17 (2007), no. 1, 15-40.
- [7] Godet, N. and Tzvetkov, N. *Strichartz estimates for the periodic non-elliptic Schrödinger equation*, C. R. Math. Acad. Sci. Paris 350 (2012), no. 21-22, 955-958.
- [8] Ham, S and Lee, S. *Restriction estimates for space curves with respect to general measure*, available at <http://arxiv.org/pdf/1303.6562.pdf>
- [9] Wang, Y. *Periodic cubic hyperbolic Schrödinger equation on \mathbb{T}^2* , J. Funct. Anal. 265 (2013), no. 3, 424-434.
- [10] Wolff, T. *Local smoothing type estimates on L^p for large p* , Geom. Funct. Anal. 10 (2000), no. 5, 1237-1288
- [11] Wooley, T. *Approximating the main conjecture in Vinogradov's mean value theorem*, available at <http://arxiv.org/abs/1401.2932>

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ 08540
E-mail address: bourgain@math.ias.edu

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, 831 EAST 3RD ST., BLOOMINGTON IN 47405
E-mail address: demeterc@indiana.edu