

Global Existence and Regularity Results for Strongly Coupled Nonregular Parabolic Systems via Iterative Methods.

DUNG LE¹

Abstract

The global existence of classical solutions to strongly coupled parabolic systems is shown to be equivalent to the availability of an iterative scheme producing a sequence of solutions with uniform continuity in the BMO norms. Amann's results on global existence of classical solutions still hold under much weaker condition that their BMO norms do not blow up in finite time. The proof makes use of some new global and local weighted Gagliardo-Nirenberg inequalities involving BMO norms.

1 Introduction

Among the long standing questions in the theory of strongly coupled parabolic systems and its applications are the global existence and regularity properties of their solutions. We consider in this paper the following system

$$\begin{cases} u_t = \operatorname{div}(A(u)Du) + f(u) & (x, t) \in Q = \Omega \times (0, T), \\ u(x, 0) = U_0(x) & x \in \Omega \\ \text{Boundary conditions for } u \text{ on } \partial\Omega \times (0, T). \end{cases} \quad (1.1)$$

Here, Ω is a bounded domain with smooth boundary $\partial\Omega$ in \mathbb{R}^n , $n > 1$, and $u : \Omega \rightarrow \mathbb{R}^m$, $f : \mathbb{R}^m \rightarrow \mathbb{R}^m$ are vector valued functions. $A(u)$ is a full matrix $m \times m$. Thus, the above is a system of m equations. The vector valued solution u satisfies either Dirichlet or Neumann boundary condition on $\partial\Omega \times (0, T)$.

The system (1.1) arises in many mathematical biology and ecology applications as well as in differential geometry theory. In the last few decades, papers concerning such strongly coupled parabolic systems usually assumed that the solutions under consideration were bounded, a very hard property to check as maximum principles had been unavailable for systems in general. In addition, past results usually relied on the following local existence result of Amann.

Theorem 1.1 ([1, 2]) *Suppose $\Omega \subset \mathbb{R}^n$, $n \geq 2$, with $\partial\Omega$ being smooth. Assume that (1.1) is normally elliptic. Let $p_0 \in (n, \infty)$ and U_0 be in $W^{1,p_0}(\Omega)$. Then there exists a maximal time $T_0 \in (0, \infty]$ such that the system (1.1) has a unique classical solution in $(0, T_0)$ with*

$$u \in C([0, T_0), W^{1,p_0}(\Omega)) \cap C^{1,2}((0, T_0) \times \bar{\Omega})$$

Moreover, if $T_0 < \infty$ then

$$\lim_{t \rightarrow T_0^-} \|u(\cdot, t)\|_{W^{1,p_0}(\Omega)} = \infty. \quad (1.2)$$

¹Department of Mathematics, University of Texas at San Antonio, One UTSA Circle, San Antonio, TX 78249. Email: Dung.Le@utsa.edu

Mathematics Subject Classifications: 35J70, 35B65, 42B37.

Key words: Parabolic systems, Hölder regularity, A_p classes, BMO weak solutions.

The proof of the above result worked directly with the system and based on semigroup and interpolation of functional spaces theories. We refer the readers to [1] for the definition of normal ellipticity. The checking of (1.2) is the most difficult one as known techniques for the regularity of solutions to scalar equations could not be extended to systems and counterexamples were available.

In this paper, we propose a different approach using iterative techniques and depart from the boundedness assumptions. Namely, we consider the following schemes

$$(u_k)_t = \operatorname{div}(A(u_{k-1})Du_k) + f(u_{k-1}, Du_k) \quad k \geq 1.$$

Under very weak assumptions on the uniform boundedness and continuity of the BMO norms of the solutions to the above systems, we will show the global existence of a classical solution to (1.1). Thus, global existence and regularity of solutions are established at once. Furthermore, without the boundedness assumptions the systems are no longer regular elliptic, we will only assume that the matrix $A(u)$ in (1.1) is *uniformly elliptic*.

We also improve Theorem 1.1 by replacing the condition (1.2) with a weaker ones using the BMO or W^{1,p_0} norms of u with $p_0 = n$. In a forthcoming work, we will show that the results in this paper can apply to a class of generalized Shigesada-Kawasaki-Teramoto models ([13]) consisting of more than 2 equations. Namely, we will establish the global existence of classical solutions to the following system

$$u_t = \Delta(P(u)) + f(u), \tag{1.3}$$

where $P(u), f(u)$ are vector valued functions whose components have quadratic (or even polynomial) growth in u .

In the proof we make use of some new local Gagliardo-Nirenberg inequalities involving BMO norms and weights in A_p classes. These are the generalizations of the inequalities by Strzelecki and Rivière in [12].

2 Preliminaries and Main Results

Throughout this paper Ω is a bounded domain with smooth boundary in \mathbb{R}^n , $n > 1$. To describe our assumptions we recall the definitions of BMO spaces and A_p classes.

For any locally integrable vector valued function $u \in L^1_{loc}(\Omega, \mathbb{R}^m)$ and measurable set $B \subset \Omega$ with its Lebesgue measure $|B| \neq 0$, we denote

$$\mathop{\int}\limits_B u \, dx = \frac{1}{|B|} \int_B u \, dx.$$

For a smooth function u defined on $\Omega \times (0, T)$, $T > 0$, its temporal and spatial derivatives are denoted by u_t, Du respectively. If A is a function in u then we also abbreviate $\frac{\partial A}{\partial u}$ by A_u .

We will frequently work with a ball $B(z, R)$ centered at $z \in \mathbb{R}^n$ with radius R . If z is understood we will write B_R for $B(z, R)$ For any $x_0 \in \Omega$ and $R > 0$, we also denote $\Omega(x_0, R) = \Omega \cap B(x_0, R)$. For any locally measurable function u on Ω and $x_0 \in \Omega$ and

$R > 0$ we write $u_{x_0,R}$ for the average of u over $\Omega(x_0, R)$. If x_0 is understood, we simply write u_R for $u_{x_0,R}$.

We say that a locally integrable vector valued function $u : \Omega \rightarrow \mathbb{R}^m$ is BMO (Bounded Mean Oscillation) if the seminorm

$$[u]_{BMO(\Omega)} = \sup_{x_0 \in \Omega, R > 0} \int_{\Omega(x_0, R)} |u - u_{\Omega(x_0, R)}| dx < \infty,$$

where the supremum is taken over all balls $B \subset \Omega$. The space $BMO(\Omega)$ is the Banach space of BMO functions on Ω with norm

$$\|u\|_{BMO(\Omega)} = \|u\|_{L^1(\Omega)} + [u]_{BMO(\Omega)}.$$

We recall the following well known fact (e.g., see [7]).

Lemma 2.1 *If $\lambda : \mathbb{R}^m \rightarrow \mathbb{R}$ is Hölder function then $\lambda(u)$ is BMO if u is BMO.*

There is a connection between BMO functions and the so called A_γ weights, which are defined as follows. Let Ψ be a measurable nonnegative function on Ω and $\gamma > 1$. We say that Ψ belongs to the class A_γ or Ψ is an A_γ weight if the quantity

$$[\Psi]_\gamma = \sup_{B(x, R) \subset \Omega} \left(\int_{B(x, R)} \Psi dx \right) \left(\int_{B(x, R)} \Psi^{1-\gamma'} dx \right)^{\gamma-1} < \infty. \quad (2.1)$$

Here, $\gamma' = \gamma/(\gamma - 1)$. The A_∞ class is defined by $A_\infty = \cup_{\gamma > 1} A_\gamma$. For more details on these classes we refer the readers to [5, 11, 14].

We also recall the following result from [7] on the connection between BMO functions and weights.

Lemma 2.2 *Let Ψ be a positive function and μ is a nonnegative Radon measure on Ω such that Ψ, Ψ^{-1} are in $BMO(\Omega)$ then Ψ belongs to $\cap_{\gamma > 1} A_\gamma$ and $[\Psi]_\gamma$ is bounded by a constant depending on $[\Psi]_{BMO(\Omega)}$ and $[\Psi^{-1}]_{BMO(\Omega)}$.*

We also recall the definition of the Campanato spaces $\mathcal{L}^{p,\gamma}(\Omega, \mathbb{R}^m)$. For any $p \geq 1$ and $\gamma > 0$ and $u \in L^p(\Omega, \mathbb{R}^m)$, we define

$$[u]_{p,\gamma}^p = \sup_{x_0 \in \Omega, \rho > 0} \rho^{-\gamma} \int_{\Omega(x_0, \rho)} |u - u_{x_0, \rho}|^p dx.$$

Then $\mathcal{L}^{p,\gamma}(\Omega, \mathbb{R}^m)$ is the Banach space of such functions with finite norm

$$\|u\|_{p,\gamma} = \|u\|_p + [u]_{p,\gamma}.$$

Clearly, $\mathcal{L}^{p,n}(\Omega, \mathbb{R}^m) = BMO(\Omega, \mathbb{R}^m)$. Moreover, it is well known that ([6, Theorem 2.9, p.52]) $\mathcal{L}^{p,\gamma}(\Omega, \mathbb{R}^m)$ is isomorphic to $C^{0,\alpha}(\Omega)$ if $\alpha = \frac{\gamma-n}{p} > 0$.

As usual, $W^{1,p}(\Omega, \mathbb{R}^m)$, $p \geq 1$, will denote the standard Sobolev spaces whose elements are vector valued functions $u : \Omega \rightarrow \mathbb{R}^m$ with finite norm

$$\|u\|_{W^{1,p}(\Omega, \mathbb{R}^m)} = \|u\|_{L^p(\Omega)} + \|Du\|_{L^p(\Omega)},$$

where Du is the the derivative of u .

We now state our structural conditions on the system (1.1).

A.1) (Uniform ellipticity) There are positive constants C, λ_0 and smooth functions $\lambda(u), \Lambda(u)$ such that $\lambda(u) \geq \lambda_0$ and $\Lambda(u) \leq C\lambda(u)$ for all $u \in \mathbb{R}^m$ and

$$\lambda(u)|\xi|^2 \leq \langle A(u)\xi, \xi \rangle \leq \Lambda(u)|\xi|^2 \quad \forall u \in \mathbb{R}^m, \xi \in \mathbb{R}^{nm}.$$

A.2) Assume that $A \in C^1(\mathbb{R}^m)$. Let Φ_0, Φ be defined as

$$\Phi_0(u) = \lambda^{\frac{1}{2}}(u) \text{ and } \Phi(u) = \frac{|A_u(u)|}{\lambda^{\frac{1}{2}}(u)} \quad u \in \mathbb{R}^m.$$

Assume that the quantities

$$k_1 := \sup_{u \in \mathbb{R}^m} \frac{|\Phi_u|}{\Phi}, \quad k_2 := \sup_{u \in \mathbb{R}^m} \frac{\Phi}{\Phi_0} \quad (2.2)$$

are finite.

A.3) (Weights) If $u \in BMO(\Omega)$ then $\Phi^{\frac{2}{3}}(u)$ belongs to the $A_{\frac{4}{3}}$ class and the quantity $[\Phi(u)^{\frac{2}{3}}]_{\frac{4}{3}}$ can be controlled by the norm $\|u\|_{BMO(\Omega)}$.

In applications, if $A(u)$ has a polynomial growth in u then we can assume that $|A(u)| \sim \lambda(u)$ and $|A_u| \sim |\lambda_u|$. In this case

$$\Phi \sim \frac{|\lambda_u|}{\lambda^{\frac{1}{2}}}, \quad |\Phi_u| \sim \frac{|\lambda_{uu}|}{\lambda^{\frac{1}{2}}} + \frac{|\lambda_u|^2}{\lambda^{\frac{3}{2}}}.$$

Therefore, if $\sup_u \frac{|\lambda_u|}{\lambda}$ and $\sup_u \frac{|\lambda_{uu}|}{|\lambda_u|}$ are bounded then (2.2) of A.2) is verified. It is clear that this is the case if $\lambda(u)$ is a polynomial in $|u|$, say $\lambda(u) \sim (1 + |u|)^k$ for some $k \geq 0$. Similarly, concerning A.3), we see that $\Phi^\alpha(u) \sim (1 + |u|)^{\alpha(k/2-1)}$. Thus, if $2 \leq k < 5$ then $\alpha(k/2 - 1) \leq 1$ for some $\alpha > 2/3$. In this case, $\Phi^\alpha(u)$ is Hölder in u so that it is BMO if u is. Of course, $\Phi^{-1}(u)$ is bounded so that it is also BMO. Lemma 2.2 shows that $\Phi^\alpha(u)$ is then an A_γ weight for all $\gamma > 1$ and A.3) is satisfied.

On the other hand, it is clear from the definition of weights that $[\Phi^\alpha]_{\beta+1} = [\Phi^{-\frac{\alpha}{\beta}}]_{\frac{\beta+1}{\beta}}^\beta$ so that if $1 < k < 2$ then $\Phi^{\frac{\alpha}{\beta}}$ is bounded from above and $\Phi^{-\frac{\alpha}{\beta}} \sim (1 + |u|)^{\frac{\alpha}{\beta}(1-k/2)}$. Thus, we can find $\alpha > 2/3$ and $\beta < 1/3$ such that $\Phi^{-\frac{\alpha}{\beta}}$ is Hölder in u and therefore BMO if u is. Again, this gives that $\Phi^{-\frac{\alpha}{\beta}}$ is a weight and belongs to $\cap_{\gamma>1} A_\gamma$.

Concerning global existence of classical solutions, we also assume that the ellipticity constants λ, Λ in A.1) are not too far apart.

R) (The ratio condition) There is $\delta \in [0, 1)$ such that

$$\frac{n-2}{n} = \delta \sup_{u \in \mathbb{R}^m} \frac{\lambda(u)}{\Lambda(u)}. \quad (2.3)$$

One should note that there are examples in [3] of blow up solutions to (1.1) if the condition R) is violated.

We assume the following growth conditions on the nonlinearity f .

F) There are positive constants C, b such that for any vector valued functions $u \in C^1(\Omega, \mathbb{R}^m)$ and $p \in C^1(\Omega, \mathbb{R}^{mn})$

$$|f(u, p)| \leq C|p| + C|u|^b + C. \quad (2.4)$$

$$|Df(u, p)| \leq C|Dp| + C|u|^{b-1}|Du| + C. \quad (2.5)$$

To solve (1.1), we can make use of the following iterative scheme. We start with any smooth vector valued function u_0 on Q and define a sequence $\{u_k\}$ of solutions to the following linear systems

$$(u_k)_t = \operatorname{div}(A(u_{k-1})Du_k) + f(u_{k-1}, Du_k) \quad k \geq 1. \quad (2.6)$$

The initial and boundary conditions for the above systems are those of u in (1.1). Note that the global existence of the strong solutions to the above systems is not generally available by standard theories (see [4]) because of the presence of Du_k in f . However, this is the case if we assume R) and the linear growth of f in Du of F) and make use of the results in [3].

Concerning the approximation sequence $\{u_k\}$, we assume the following uniform bound and continuity of their BMO norms.

V) Let $\{u_k\}$ and Φ be defined by (2.6) and A.2). There exists a continuous function C on $(0, \infty)$ such that for any $T > 0$

$$\|u_0(\cdot, t)\|_{C^1(\Omega)}, \|u_k(\cdot, t)\|_{BMO(\Omega)} \leq C(T) \quad \forall t \in (0, T), k = 1, 2, \dots$$

Moreover, for any $\varepsilon > 0$ and $(x, t) \in Q$, there exists $R = R(\varepsilon, T) > 0$ such that

$$\|u_k(\cdot, t)\|_{BMO(B_R(x))} < \varepsilon \quad \forall t \in (0, T), k = 0, 1, \dots \quad (2.7)$$

In addition, for all $(x, t) \in Q$ and integer $k \geq 1$ we assume that

$$\text{either } \frac{\Phi(u_k(x, t))}{\Phi(u_{k-1}(x, t))} \leq C(T) \text{ or } \frac{\lambda(u_k(x, t))}{\lambda(u_{k-1}(x, t))} \leq C(T). \quad (2.8)$$

The uniform boundedness assumption on the BMO norm of u_k is of course much weaker than the L^∞ boundedness assumptions in literature. Moreover, the uniform continuity assumption (2.7) on the BMO norms is somehow necessary for the regularity of the limit solution u .

The assumption (2.8) seems to be technical at first glance but it is clearly necessary if we would like to produce a the sequence $\{u_k\}$ that converges in $L^\infty(Q)$ to a solution of (1.1). In particular, if $\lambda(u)$ behaves like $\lambda(u) \sim (1 + |u|)^k$ for some $k \geq 0$ then, as discussed earlier, we see that $\Phi(u) \sim (1 + |u|)^{(k/2-1)}$. Thus,

$$\frac{\Phi(u_k(x, t))}{\Phi(u_{k-1}(x, t))} \sim \left(\frac{1 + |u_k(x, t)|}{1 + |u_{k-1}(x, t)|} \right)^{k/2-1}, \quad \frac{\lambda(u_k(x, t))}{\lambda(u_{k-1}(x, t))} \sim \left(\frac{1 + |u_k(x, t)|}{1 + |u_{k-1}(x, t)|} \right)^k.$$

Hence, if $k \in [0, 2]$ then (2.8) is clearly verified. The generalized Shigesada-Kawasaki-Teramoto model (1.3) clearly is a typical example.

We then have our main result of this paper as follows.

Theorem 2.3 *Assume A.1)-A.3), R), F) and V). Then the sequence $\{u_k\}$ has a subsequence that converges strongly to a classical solution u in $C^0(\Omega \times (0, \infty))$.*

A simple consequence of the above theorem is the following.

Corollary 2.4 *Assume A.1)-A.3), R), F) and V). Then the system (1.1) has a classical solution u that exists globally on $\Omega \times (0, \infty)$ if and only if there is u_0 and a sequence $\{u_k\}$ satisfying V) such that $\{u_k\}$ has a subsequence that converges weakly to u in $L^2(\Omega \times (0, T))$ for each $T > 0$.*

The necessary part is trivial as we can take $u_k = u$ for all $k \geq 0$, the solution sequence is then a constant one. The sufficient part comes from Theorem 2.3 and the uniqueness of weak limits in $L^2(\Omega \times (0, T))$.

In the next theorem we discuss the global existence of classical solutions when their local existence can be achieved by other methods (e.g., Theorem 1.1).

Theorem 2.5 *Assume A.1)-A.3), R) and F). Let $p_0 \in (n, \infty)$ and U_0 be in $W^{1,p_0}(\Omega)$. Suppose that $T_0 \in (0, \infty]$ is the maximal existence time for a classical solution*

$$u \in C([0, T_0], W^{1,p_0}(\Omega)) \cap C^{1,2}((0, T_0) \times \bar{\Omega})$$

for the system (1.1). Suppose that there is a function C in $C^0((0, T_0])$ such that

$$\|u(\cdot, t)\|_{BMO(\Omega)} \leq C(t) \quad \forall t \in (0, T_0).$$

Moreover, for any $\varepsilon > 0$ and $(x, t) \in Q$, there exists $R = R(\varepsilon) > 0$ such that

$$\|u(\cdot, t)\|_{BMO(B_R(x))} < \varepsilon \quad \forall t \in (0, T_0). \quad (2.9)$$

Then $T_0 = \infty$.

By Poincaré's inequality,

$$\int_{B_R} |u - u_R|^n dx \leq C(n) \int_{B_R} |Du|^n dx,$$

it is easy to see that if $Du(\cdot, t) \in L^n(\Omega)$ then $u(\cdot, t)$ is BMO and $\|u(\cdot, t)\|_{BMO(B_R)}$ is small if R is small. Therefore, as a simple consequence of the above theorem, we have the following improvisation of Theorem 1.1.

Corollary 2.6 *In addition to the assumptions of Theorem 1.1, we assume R). Then there exists a maximal time $T_0 \in (0, \infty]$ such that the system (1.1) has a unique classical solution in $(0, T_0)$ with*

$$u \in C([0, T_0], W^{1,p_0}(\Omega)) \cap C^{1,2}((0, T_0) \times \bar{\Omega})$$

Moreover, if $T_0 < \infty$ then

$$\lim_{t \rightarrow T_0^-} \|u(\cdot, t)\|_{W^{1,n}(\Omega)} = \infty. \quad (2.10)$$

Again, we remark that if the condition R) is violated then there are counterexamples for finite time blow up solutions to (1.1).

3 Weighted Gagliardo-Nirenberg inequalities

In this section we will establish global and local weighted Gagliardo-Nirenberg interpolation inequalities which allow us to control the L^{2p+2} norm of Du_k in the proof of our main theorems.

Since the inequalities can be useful for other applications, we will prove them under a set of the following independent assumptions.

GN.1) Let Φ_0, Φ be positive functions on \mathbb{R}^m with $\Phi \in C^1(\mathbb{R}^m)$. Assume that the following quantities are finite

$$k_1 := \sup_{u \in \mathbb{R}^m} \frac{|\Phi_u|}{\Phi}, \quad k_2 := \sup_{u \in \mathbb{R}^m} \frac{\Phi}{\Phi_0}. \quad (3.1)$$

GN.2) For some $p \geq 1$ suppose that $\Phi(u)^{\frac{2}{p+2}}$ belongs to the $A_{\frac{p}{p+2}+1}$ class if u is BMO and the quantity $[\Phi^{\frac{2}{p+2}}]_{\frac{p}{p+2}+1}$ can be controlled by the norm $\|u\|_{BMO(\Omega)}$.

In the rest of this paper we will slightly abuse our notations and write the dot product $\langle u, v \rangle$ as uv for any two vectors u, v because its meaning should be clear in the context. Similarly, when there is no ambiguity C will denote a universal constant that can change from line to line in our argument. Furthermore, $C(\cdot \cdot \cdot)$ is used to denote quantities which are bounded in terms of their parameters.

Lemma 3.1 *Assume GN.1) and GN.2). Let $u, U : \Omega \rightarrow \mathbb{R}^m$ be vector valued functions with $U \in C^2(\Omega)$, $u \in C^1(\Omega)$. Suppose further that either U or $\Phi^2(u) \frac{\partial U}{\partial \nu}$ vanish on the boundary $\partial\Omega$ of Ω . We set*

$$I_1 := \int_{\Omega} \Phi^2(u) |DU|^{2p+2} dx, \quad \hat{I}_1 := \int_{\Omega} \Phi^2(u) |Du|^{2p+2} dx, \quad (3.2)$$

and

$$I_2 := \int_{\Omega} \Phi_0^2(u) |DU|^{2p-2} |D^2U|^2 dx. \quad (3.3)$$

Then there is a constant C_{Φ} depending on $[\Phi^{\frac{2}{p+2}}(u)]_{\frac{p}{p+2}+1}$ for which

$$I_1 \leq C_{\Phi} \|U\|_{BMO(\Omega)} \left[k_1 (I_1 + \hat{I}_1) + k_2 I_1^{\frac{1}{2}} I_2^{\frac{1}{2}} \right]. \quad (3.4)$$

Before we go to the proof of this lemma, let us recall the following facts from Harmonic Analysis. We first recall the definition of the maximal function of a function $F \in L^1_{loc}(\Omega)$

$$M(F)(x) = \sup_{\varepsilon} \left\{ \int_{B_{\varepsilon}(x)} F(x) dx : \varepsilon > 0 \text{ and } B_{\varepsilon}(x) \subset \Omega \right\}.$$

We also note here the Hardy-Littlewood theorem for any $F \in L^q(\Omega)$

$$\int_{\Omega} M(F)^q dx \leq C(q) \int_{\Omega} F^q dx, \quad q > 1. \quad (3.5)$$

In addition, let us recall the definition of the Hardy space \mathcal{H}^1 . For any $y \in \Omega$ and $\varepsilon > 0$. Let ϕ be any function in $C_0^1(B_1(y))$ with $|D\phi| \leq C_1$. Let $\phi_\varepsilon(x) = \varepsilon^{-n}\phi(\frac{x}{\varepsilon})$ (then $|D\phi_\varepsilon| \leq C_1\varepsilon^{-1}$). From [?], a function g is in $\mathcal{H}^1(\Omega)$ if

$$\sup_{\varepsilon>0} g * \phi_\varepsilon \in L^1(\Omega) \text{ and } \|g\|_{\mathcal{H}^1} = \|g\|_{L^1(\Omega)} + \|\sup_{\varepsilon>0} g * \phi_\varepsilon\|_{L^1(\Omega)}.$$

Furthermore, concerning the A_γ classes, it is well known (e.g. [11, Corollary 2.5]) that if w belongs to the A_γ class for some $\gamma > 1$ then there are positive ε, ν depending on $[w]_\gamma$ such that $w^{1+\varepsilon}$ belongs to the $A_{\gamma-\nu}$ class and $[w^{1+\varepsilon}]_{\gamma-\nu}$ can be controlled by $[w]_\gamma$. Hence, by GN.2) there are positive constants α, β depending on the quantity $[\Phi^{\frac{2}{p+2}}]_{\frac{p}{p+2}+1}$ such that

$$\alpha > 2/(2+p), \beta < p/(p+2) \text{ and } [\Phi^\alpha]_{\beta+1} \leq C([\Phi^{\frac{2}{p+2}}]_{\frac{p}{p+2}+1}). \quad (3.6)$$

A simple use of Hölder's inequality also gives

$$[w^\delta]_\gamma \leq [w]_\gamma^\delta \quad \forall \delta \in (0, 1). \quad (3.7)$$

Proof: Integrating by parts, we have

$$\int_\Omega \Phi^2(u) |DU|^{2p+2} dx = - \int_\Omega U \operatorname{div}(\Phi^2(u) |DU|^{2p} DU) dx. \quad (3.8)$$

We will show that $g = \operatorname{div}(\Phi^2(u) |DU|^{2p} DU)$ belongs to the Hardy space \mathcal{H}^1 and

$$\|g\|_{\mathcal{H}^1} = \int_\Omega \sup_\varepsilon |g * \phi_\varepsilon| dx \leq C([\Phi^{\frac{2}{p+2}}(u)]_{\frac{p}{p+2}+1}) \left[k_1(I_1 + \hat{I}_1) + k_2 I_1^{\frac{1}{2}} I_2^{\frac{1}{2}} \right]. \quad (3.9)$$

Once this is established, (3.8) and the duality of the BMO and Hardy spaces give (3.4).

We write $g = g_1 + g_2$ with $g_i = \operatorname{div} V_i$, setting

$$V_1 = \Phi(u) |DU|^{p+1} \left(\Phi(u) |DU|^{p-1} DU - \int_{B_\varepsilon} \Phi(u) |DU|^{p-1} DU dx \right),$$

and

$$V_2 = \Phi(u) |DU|^{p+1} \int_{B_\varepsilon} \Phi(u) |DU|^{p-1} DU dx.$$

Let us consider g_1 first and define $h = \Phi(u) |DU|^{p-1} DU$. For any $y \in \Omega$ and $B_\varepsilon = B_\varepsilon(y) \subset \Omega$, we use integration by parts, the property of ϕ_ε and then Hölder's inequality for any $s > 1$ to have the following

$$\begin{aligned} |g_1 * \phi_\varepsilon| &= \left| \int_{B_\varepsilon} D\phi\left(\frac{x-y}{\varepsilon}\right) (h - h_{B_\varepsilon}) \Phi(u) |DU|^{p+1} dx \right| \\ &\leq \frac{C_1}{\varepsilon} \left| \int_{B_\varepsilon} |h - h_{B_\varepsilon}| \Phi(u) |DU|^{p+1} dx \right| \\ &\leq \frac{C_1}{\varepsilon} \left(\int_{B_\varepsilon} |h - h_{B_\varepsilon}|^s dx \right)^{\frac{1}{s}} \left(\int_{B_\varepsilon} \Phi^{s'}(u) |DU|^{(p+1)s'} dx \right)^{\frac{1}{s'}}. \end{aligned}$$

By Sobolev-Poincaré's inequality, with $s_* = ns/(n+s)$, we have the following estimate in noting that $|Dh| \leq |D\Phi||DU|^p + p\Phi|DU|^{p-1}|D^2U|$

$$\begin{aligned} \frac{C_1}{\varepsilon} \left(\int_{B_\varepsilon} |h - h_{B_\varepsilon}|^s dx \right)^{\frac{1}{s}} &\leq C \left(\int_{B_\varepsilon} |Dh|^{s_*} dx \right)^{\frac{1}{s_*}} \\ &\leq C \left[\int_{B_\varepsilon} |D\Phi|^{s_*} |DU|^{ps_*} dx + \int_{B_\varepsilon} \Phi^{s_*} |DU|^{(p-1)s_*} |D^2U|^{s_*} dx \right]^{\frac{1}{s_*}} \end{aligned} \quad (3.10)$$

Take $s = 2n/(n-1)$ then $s_* = s' = 2n/(n+1) < 2$. From the definitions of $k_1 = \sup_{u \in \mathbb{R}^m} \frac{|\Phi_u|}{\Phi}$ and $k_2 = \sup_{u \in \mathbb{R}^m} \frac{\Phi}{\Phi_0}$ in GN.1) we have

$$\left(\int_{B_\varepsilon} |D\Phi|^{s_*} |DU|^{ps_*} dx \right)^{\frac{1}{s_*}} \leq k_1 \left(\int_{B_\varepsilon} \Phi |DU|^{ps_*} |Du|^{s_*} dx \right)^{\frac{1}{s_*}},$$

and

$$\left(\int_{B_\varepsilon} \Phi^{s_*} |DU|^{(p-1)s_*} |D^2U|^{s_*} dx \right)^{\frac{1}{s_*}} \leq k_2 \left(\int_{B_\varepsilon} \Phi_0^{s_*} |DU|^{(p-1)s_*} |D^2U|^{s_*} dx \right)^{\frac{1}{s_*}}.$$

Using the above estimates in (3.10), we get

$$\frac{C_1}{\varepsilon} \left(\int_{B_\varepsilon} |h - h_{B_\varepsilon}|^s dx \right)^{\frac{1}{s}} \leq C [k_1 \Psi_1 + k_2 \Psi_2], \quad (3.11)$$

where

$$\Psi_1(y) = (M(\Phi^{s_*} |DU|^{ps_*} |Du|^{s_*})(y))^{\frac{1}{s_*}}, \quad \Psi_2(y) = (M(\Phi_0^{s_*} |DU|^{(p-1)s_*} |D^2U|^{s_*})(y))^{\frac{1}{s_*}}.$$

Putting these estimates together we thus have

$$\sup_{\varepsilon > 0} |g_1 * \phi_\varepsilon| \leq C \Psi_1 [k_1 \Psi_1 + k_2 \Psi_2]. \quad (3.12)$$

Because $2 > 2n/(n+1) = s_*$, we can use Young's inequality and then (3.5) to get

$$\begin{aligned} \left(\int_{\Omega} \Psi_1^2 dx \right)^{\frac{1}{2}} &\leq \|M(|\Phi|^{s_*} |DU|^{(p+1)s_*})\|_{\frac{2}{s_*}}^{\frac{1}{2}} + \|M(|\Phi|^{s_*} |Du|^{(p+1)s_*})\|_{\frac{2}{s_*}}^{\frac{1}{2}} \\ &\leq C (\|\Phi^2 |DU|^{2(p+1)}\|_2 + \|\Phi^2 |Du|^{2(p+1)}\|_2), \end{aligned}$$

and

$$\left(\int_{\Omega} \Psi_2^2 dx \right)^{\frac{1}{2}} = \|M(\Phi_0^{s_*} |DU|^{(p-1)s_*} |D^2u|^{s_*})\|_{\frac{2}{s_*}}^{\frac{1}{2}} \leq C \|\Phi_0^2 |DU|^{p-1} |D^2U|\|_2.$$

Therefore, by Holder's inequality and the above estimates

$$\int_{\Omega} \sup_{\varepsilon} |g_1 * \phi_\varepsilon| dx \leq C \left[k_1 (I_1 + \hat{I}_1) + k_2 I_1^{\frac{1}{2}} I_2^{\frac{1}{2}} \right]. \quad (3.13)$$

We now turn to g_2 and note that $|\operatorname{div} V_2| \leq C(J_1 + J_2)$ for some constant C and

$$J_1 := \sup \frac{|\Phi u|}{\Phi} \Phi |DU|^{p+1} |Du| J_3, \quad J_2 := \Phi |DU|^p |D^2 U| J_3 \text{ with } J_3 := \left| \int_{B_\varepsilon} \Phi |DU|^p dx \right|.$$

We consider J_1 . For any $r > 1/(p+1)$ we denote $r^* = 1 - \frac{1}{r(p+1)}$. We also write $F = \Phi |DU|^{p+1}$ and $\hat{F} = \Phi |Du|^{p+1}$. For any $s > 0$, Hölder's inequality yields

$$\left(\int_{B_\varepsilon} \Phi^{\frac{-s}{(p+1)}} \Phi^{\frac{s}{p+1}} |Du|^s dx \right)^{\frac{1}{s}} \leq \left(\int_{B_\varepsilon} \Phi^{\frac{-s}{r^*(p+1)}} dx \right)^{\frac{r^*}{s}} \left(\int_{B_\varepsilon} \hat{F}^{sr} dx \right)^{\frac{1}{rs(p+1)}}.$$

Similarly, if $r_1 > 1/(p+1)$ we have the following estimate for J_3

$$J_3 = \int_{B_\varepsilon} \Phi^{\frac{1}{(p+1)}} \Phi^{\frac{p}{p+1}} |DU|^p dx \leq \left(\int_{B_\varepsilon} \Phi^{\frac{1}{r_1^*(p+1)}} dx \right)^{r_1^*} \left(\int_{B_\varepsilon} F^{pr_1} dx \right)^{\frac{1}{r_1(p+1)}}.$$

Using Hölder's inequality for the integrand $\Phi |DU|^{p+1} |Du| = F |Du|$ in J_1 and the above estimates, we obtain

$$\sup_\varepsilon |\phi_\varepsilon * J_1| \leq C(\Phi, r, r_1, s) \sup \frac{|\Phi u|}{\Phi} M(F^{s'})^{\frac{1}{s'}} M(\hat{F}^{sr})^{\frac{1}{rs(p+1)}} M(F^{pr_1})^{\frac{1}{r_1(p+1)}},$$

where

$$C(\Phi, r, r_1, s) = C \sup_\varepsilon \left(\int_{B_\varepsilon} \Phi^{\frac{1}{r_1^*(p+1)}} dx \right)^{r_1^*} \left(\int_{B_\varepsilon} \Phi^{\frac{-s}{r^*(p+1)}} dx \right)^{\frac{r^*}{s}}. \quad (3.14)$$

By the definition of weights, it is clear that

$$C(\Phi, r, r_1, s) \leq \left[\Phi^{\frac{1}{r_1^*(p+1)}} \right]_{\frac{r^*}{r_1^*s} + 1}^{r_1^*} = \left[\Phi^{\frac{-s}{r^*(p+1)}} \right]_{\frac{r_1^*s}{r^*} + 1}^{\frac{r^*}{s}}. \quad (3.15)$$

We now choose s, r, r_1 such that $s' = sr = pr_1$ and $sr < 2$. This is the case if $r < 1$, $s = (r+1)/r$ and $r_1 = (r+1)/p$. Let

$$\alpha(r) = \frac{1}{r_1^*(p+1)} = \frac{1}{p+1 - \frac{p}{r+1}} \text{ and } \beta(r) = \frac{r^*}{r_1^*s} = \frac{r(p+1) - 1}{r(p+1) + 1}.$$

We then have $C(\Phi, r, r_1, s) \leq [\Phi^{\alpha(r)}]_{\beta(r)+1}^{r_1^*}$. Clearly, $\alpha(r)$ decreases to $2/(p+2)$ and $\beta(r)$ increases to $p/(p+2)$ as $r \rightarrow 1$. Thus, by the choice α, β as in (3.6), if we choose r close to 1 then $\alpha(r) < \alpha$ and $\beta(r) > \beta$ and thus $[\Phi^{\alpha(r)}]_{\beta(r)+1} \leq C([\Phi^\alpha]_{\beta+1})$, see (3.7). Hence, the above estimates give

$$\sup_\varepsilon |\phi_\varepsilon * J_1| \leq C([\Phi^\alpha]_{\beta+1}) \sup \frac{|\Phi u|}{\Phi} M(F^{sr})^{\frac{2p+1}{rs(p+1)}} M(\hat{F}^{sr})^{\frac{1}{rs(p+1)}}.$$

By Young's inequality we have

$$M(F^{sr})^{\frac{2p+1}{rs(p+1)}} M(\hat{F}^{sr})^{\frac{1}{rs(p+1)}} \leq C(M(F^{sr})^{\frac{2}{rs}} + M(\hat{F}^{sr})^{\frac{2}{rs}}).$$

We then use (3.5) for $q = rs < 2$ to see that

$$\int_{\Omega} \sup_{\varepsilon} |\phi_{\varepsilon} * J_1| dx \leq C([\Phi^{\alpha}]_{\beta+1}) \sup \frac{|\Phi_u|}{\Phi} (I_1 + \hat{I}_1). \quad (3.16)$$

Next, we write $J_2 = \Phi |DU|^{p-1} |D^2U| |DU| J_3$. By Hölder's inequality,

$$\phi_{\varepsilon} * J_2 \leq \left(\int_{B_{\varepsilon}} \Phi |DU|^{p-1} |D^2U|^{s'} dx \right)^{\frac{1}{s'}} \left(\int_{B_{\varepsilon}} \Phi^{\frac{-s}{(p+1)}} \Phi^{\frac{s}{p+1}} |DU|^s dx \right)^{\frac{1}{s}} J_3.$$

Again, Hölder's inequality gives

$$\left(\int_{B_{\varepsilon}} \Phi^{\frac{-s}{(p+1)}} \Phi^{\frac{s}{p+1}} |DU|^s dx \right)^{\frac{1}{s}} \leq \left(\int_{B_{\varepsilon}} \Phi^{\frac{-s}{r^*(p+1)}} dx \right)^{\frac{r^*}{s}} \left(\int_{B_{\varepsilon}} F^{sr} dx \right)^{\frac{1}{rs(p+1)}}.$$

Using the estimate for J_3 , with different r_1 , we have

$$\sup_{\varepsilon} \phi_{\varepsilon} * J_2 \leq C(\Phi, r, r_1, s) M((\Phi |DU|^{p-1} |D^2U|)^{s'})^{\frac{1}{s'}} M(F^{sr})^{\frac{1}{rs(p+1)}} M(F^{pr_1})^{\frac{1}{r_1(p+1)}}.$$

We will choose s, r_1, r such that $pr_1 = sr < 2$ (and $r_1 > 1/(p+1)$). Again, $C(\Phi, r, r_1, s)$ can be estimated by $C([\Phi^{\alpha(r)}]_{\beta(r)+1})$, where

$$\alpha(r) = \frac{1}{p+1 - \frac{p}{rs}} \text{ and } \beta(r) = \frac{r(p+1) - 1}{rs(p+1) - p}.$$

Obviously, we choose $s > 2$ and s is close to 2, $r < 1$ and r is close to 1 such that $sr < 2$, $\alpha(r) < \alpha$ and $\beta(r) > \beta$, see (3.6). As before, with such choice of s, r , we have

$$\sup_{\varepsilon} \phi_{\varepsilon} * J_2 \leq C([\Phi^{\alpha}]_{\beta+1}) M((\Phi |DU|^{p-1} |D^2U|)^{s'})^{\frac{1}{s'}} M(F^{sr})^{\frac{1}{rs}}.$$

We have by (3.5) and similar estimates for (3.10), with s_* is now s' (which is less than 2 because $s > 2$), the following

$$\int_{\Omega} \sup_{\varepsilon} |\phi_{\varepsilon} * J_2| dx \leq C([\Phi^{\alpha}]_{\beta+1}) \sup \frac{\Phi}{\Phi_0} I_1^{\frac{1}{2}} I_2^{\frac{1}{2}}. \quad (3.17)$$

Combining the estimates (3.13), (3.16), (3.17) and GN.1) we obtain the bound

$$\int_{\Omega} \sup_{\varepsilon} |g_2 * \phi_{\varepsilon}| dx \leq C([\Phi^{\alpha}]_{\beta+1}) \left[k_1 I_1 + k_2 I_1^{\frac{1}{2}} I_2^{\frac{1}{2}} \right]. \quad (3.18)$$

We thus prove (3.9) for the \mathcal{H}^1 norm of g . Therefore, by FS theorem

$$I_1 \leq \|U\|_{BMO} \|g\|_{\mathcal{H}^1} \leq C([\Phi^{\alpha}]_{\beta+1}) \|U\|_{BMO} \left[k_1 (I_1 + \hat{I}_1) + k_2 I_1^{\frac{1}{2}} I_2^{\frac{1}{2}} \right].$$

As we noted before, see (3.6), $[\Phi^{\alpha}]_{\beta+1}$ can be controlled by $[\Phi^{\frac{2}{p+2}}]_{\frac{p}{p+2}+1}$. Hence, the above gives (3.9) and the proof is complete. ■

Remark 3.2 By approximation, see [12], the lemma also holds for $u \in W^{1,2}(\Omega)$ and $U \in W^{2,2}(\Omega)$ provided that the quantities I_1, I_2 and \hat{I}_1 defined in (3.2) and (3.20) are finite. Furthermore, if Φ, Φ_0 are constant functions and $u = U$, the above lemma clearly gives the Gagliardo-Nirenberg inequality established in [12].

In order to make use of the continuity in BMO norms in the assumption V) to obtain the regularity results, we will need the following local version of the above lemma.

Lemma 3.3 *Assume as in Lemma 3.1 and let B_s, B_t be two concentric balls in Ω with radii $t > s > 0$. We set*

$$I_1(t) := \int_{B_t} \Phi^2(u) |DU|^{2p+2} dx, \quad \hat{I}_1(t) := \int_{B_t} \Phi^2(u) |Du|^{2p+2} dx, \quad (3.19)$$

and

$$I_2(t) := \int_{B_t} \Phi_0^2(u) |DU|^{2p-2} |D^2U|^2 dx. \quad (3.20)$$

Let ψ be a C^1 function such that $\psi = 1$ in B_s and $\psi = 0$ outside B_t . For any $\varepsilon > 0$ there is a constant $C(\varepsilon)$ such that

$$\begin{aligned} I_1(s) \leq C(\varepsilon) \|U\|_{BMO(B_t)} \sup_{x \in B_t} |D\psi(x)|^2 \int_{B_t} |\Phi|^2(u) |DU|^{2p} dx \\ + C_\Phi \|U\|_{BMO(B_t)} \left[(k_1 + \varepsilon)(I_1(t) + \hat{I}_1(t)) + k_2 I_1^{\frac{1}{2}}(t) I_2^{\frac{1}{2}}(t) \right]. \end{aligned} \quad (3.21)$$

Proof: We revisit the proof of the previous lemma. Integrating by parts, we have

$$\int_{\Omega} \Phi^2(u) \psi^2 |DU|^{2p+2} dx = - \int_{\Omega} U \operatorname{div}(\Phi^2(u) \psi^2 |DU|^{2p} DU) dx.$$

Again, we will show that $g = \operatorname{div}(\Phi^2 \psi^2 |DU|^{2p} DU)$ belongs to the Hardy space \mathcal{H}^1 . We write $g = g_1 + g_2$ with $g_i = \operatorname{div} V_i$, setting

$$V_1 = \Phi(u) \psi |DU|^{p+1} \left(\Phi(u) \psi |DU|^{p-1} DU - \int_{B_\varepsilon} \Phi(u) \psi |DU|^{p-1} DU dx \right),$$

and

$$V_2 = \Phi(u) \psi |DU|^{p+1} \int_{B_\varepsilon} \Phi(u) \psi |DU|^{p-1} DU dx.$$

In estimating V_1 we follow the proof of Lemma 3.1 and replace $\Phi(u), \Phi_0(u)$ respectively by $\Phi(u) \psi(x)$ and $\Phi_0(u) \psi$. There will be the following extra term in estimating Dh in the right hand side of (3.10) and it can be estimated as follows

$$\left(\int_{B_\varepsilon} \Phi^{s^*}(u) |D\psi|^{s^*} |DU|^{ps^*} dx \right)^{\frac{1}{s^*}} \leq \sup_{x \in B_t} |D\psi| \left(\int_{B_\varepsilon} \Phi^{s^*}(u) |DU|^{ps^*} dx \right)^{\frac{1}{s^*}}.$$

We then use the the following in the right hand side of (3.12) (with $\Omega = B_t$)

$$\begin{aligned} \sup |D\psi| \int_{B_t} \Psi_1 M(\Phi^{s^*}(u) |DU|^{ps^*})^{\frac{1}{s^*}} dx \leq \\ \varepsilon \int_{B_t} \Psi_1^2 dx + C(\varepsilon) \sup_{x \in B_t} |D\psi|^2 \int_{B_t} M(\Phi^{s^*}(u) |DU|^{ps^*})^{\frac{2}{s^*}} dx. \end{aligned}$$

The last term can be bounded via (3.5) by

$$C(\varepsilon) \sup_{x \in B_t} |D\psi|^2 \int_{B_t} \Phi^2(u) |DU|^{2p} dx.$$

Using the fact that $|\psi| \leq 1$ and taking Ω to be B_t , the previous proof can go on and (3.13) now becomes

$$\begin{aligned} \int_{B_t} \sup_{\varepsilon} |g_1 * \phi_{\varepsilon}| dx &\leq C \left[(k_1 + \varepsilon) (I_1(t) + \hat{I}_1(t)) + k_2 I_1^{\frac{1}{2}}(t) I_2^{\frac{1}{2}}(t) \right] \\ &\quad + C(\varepsilon) \sup |D\psi|^2 \int_{B_t} \Phi^2(u) |DU|^{2p} dx. \end{aligned} \quad (3.22)$$

Similarly, in considering $g_2 = \operatorname{div} V_2$, we will have an extra term $\Phi(u) |D\psi| |DU|^{p+1} J_3$ in J_1 . We then use the following estimate

$$\sup_{\varepsilon} |\phi_{\varepsilon} * \Phi(u) |D\psi| |DU|^{p+1} J_3 \leq \sup |D\psi| M(\Phi(u) |DU|^{p+1}) M(\Phi(u) |DU|^p),$$

and via Young's inequality

$$\int_{B_t} \sup_{\varepsilon} |\phi_{\varepsilon} * \Phi(u) |D\psi| |DU|^{p+1} J_3 dx \leq \varepsilon I_1(t) + C(\varepsilon) \sup |D\psi|^2 \int_{B_t} \Phi^2(u) |DU|^{2p} dx.$$

Therefore the estimate (3.18) is now (3.22) with g_1 being replaced by g_2 . Combining the estimates for g_1, g_2 and using Young's inequality, we get

$$\begin{aligned} \int_{B_t} \sup_{\varepsilon} |g * \phi_{\varepsilon}| dx &\leq C(\varepsilon) \sup |D\psi|^2 \int_{B_t} |\Phi|^2 |DU|^{2p} dx \\ &\quad + C([\Phi^{\alpha}]_{\beta+1}) \left[(k_1 + \varepsilon) (I_1(t) + \hat{I}_1(t)) + k_2 I_1^{\frac{1}{2}}(t) I_2^{\frac{1}{2}}(t) \right]. \end{aligned} \quad (3.23)$$

The above gives an estimate for the \mathcal{H}^1 norm of g . By FS theorem, we obtain

$$\begin{aligned} \int_{B_t} \Phi^2(u) \psi^2 |DU|^{2p+2} dx &\leq C(\varepsilon) \|u\|_{BMO(B_t)} \sup |D\psi|^2 \int_{B_t} \Phi^2(u) |DU|^{2p} dx \\ &\quad + C([\Phi^{\alpha}]_{\beta+1}) \|u\|_{BMO(B_t)} \left[(k_1 + \varepsilon) (I_1(t) + \hat{I}_1(t)) + k_2 I_1^{\frac{1}{2}}(t) I_2^{\frac{1}{2}}(t) \right]. \end{aligned}$$

Since $\psi = 1$ in B_s , the above yields (3.21) and the proof is complete. ■

Finally, the following lemma will be crucial in obtaining uniform estimates for the approximation sequence $\{u_k\}$.

Lemma 3.4 *Assume as in Lemma 3.3 and let B_{ρ}, B_R be two concentric balls in Ω with radii $R > \rho > 0$. Assume that there is a constant $C(\Phi, \Phi_0)$ depending on $C(\Phi, \Phi_0)$ such that the constant*

$$C(\Phi, \Phi_0) := C_{\Phi} \left(\sup_u \frac{|\Phi_u|}{\Phi} + \sup_u \frac{\Phi}{\Phi_0} + 1 \right) \quad (3.24)$$

is finite. Then there is ε_0 depending on $C(\Phi, \Phi_0)$ such that if $\|U\|_{BMO(B_R)} < \varepsilon_0$ then there is a constant $C_0(\Phi, \Phi_0)$ such that

$$I_1(s) \leq C_0(\Phi, \Phi_0) \|U\|_{BMO(B_R)} \left[\frac{1}{(t-s)^2} \int_{B_t} \Phi_0^2 |DU|^{2p} dx + \hat{I}_1(t) + I_2(t) \right] \quad (3.25)$$

for any s, t such that $0 < s < t < R$.

For the proof of this lemma and later use, let us recall the following elementary iteration result (e.g., see [6, Lemma 6.1, p.192]).

Lemma 3.5 *Let f, g, h be bounded nonnegative functions in the interval $[\rho, R]$ with g, h being increasing. Assume that for $\rho \leq s < t \leq R$ we have*

$$f(s) \leq [(t-s)^{-\alpha}g(t) + h(t)] + \varepsilon f(t)$$

with $C \geq 0$, $\alpha > 0$ and $0 \leq \varepsilon < 1$. Then

$$f(\rho) \leq c(\alpha, \varepsilon)[(R-\rho)^{-\alpha}g(R) + h(R)].$$

The constant $c(\alpha, \varepsilon)$ can be taken to be $(1-\nu)^{-\alpha}(1-\nu^{-\alpha}\nu_0)^{-1}$ for any ν satisfying $\nu^{-\alpha}\nu_0 < 1$.

We are now ready to give the proof of Lemma 3.4.

Proof: For any s, t, ρ such that $0 < \rho < s < t < R$, let ψ be a cutoff function for B_s, B_t with $|D\psi| \leq 1/(t-s)$. Noting that $\|U\|_{BMO(B_t)} \leq \|U\|_{BMO(B_R)}$. By a simple use of Young's inequality to the last product in (3.21) of Lemma 3.3 and our assumption (3.24), we can see easily that if $\varepsilon, \varepsilon_0$ are sufficiently small then for some $\nu_0 \in (0, 1)$

$$C(\Phi, \Phi_0)\|U\|_{BMO(B_R)} \leq \nu_0,$$

and

$$I_1(s) \leq \nu_0 I_1(t) + C(\Phi, \Phi_0)\|U\|_{BMO(B_R)} \left[\frac{1}{(t-s)^2} \int_{B_t} \Phi^2 |DU|^{2p} dx + \hat{I}_1(t) + I_2(t) \right].$$

Let $C_1(\Phi, \Phi_0) = C(\Phi, \Phi_0) \max\{1, \sup_u \frac{\Phi^2}{\Phi_0^2}\}$. The above yields

$$I_1(s) \leq \nu_0 I_1(t) + C_1(\Phi, \Phi_0)\|U\|_{BMO(B_R)} \left[\frac{1}{(t-s)^2} \int_{B_t} \Phi_0^2 |DU|^{2p} dx + \hat{I}_1(t) + I_2(t) \right].$$

It is clear that we can Lemma 3.5 to $f = I_1$ to get

$$I_1(\rho) \leq C(\nu_0)C_1(\Phi, \Phi_0)\|U\|_{BMO(B_R)} \left[\frac{1}{(R-\rho)^2} \int_{B_R} \Phi_0^2 |DU|^{2p} dx + \hat{I}_1(R) + I_2(R) \right].$$

The constant $C(\nu_0)$ can be taken to be $(1-\nu)^{-2}(1-\nu^{-2}\nu_0)^{-1}$ for any ν satisfying $\nu^{-2}\nu_0 < 1$. We can take $C(\nu_0)$ to be a fixed constant for $\nu_0 \in (0, \frac{1}{2})$. Obviously, the above also holds for ρ, R being replaced by s, t and we proved the lemma. ■

4 Proof of the main theorems

We now go back to the iterative scheme (2.6) and prove our main theorems in this section.

The following lemma is the main vehicle of the proof of Theorem 2.3.

Lemma 4.1 *Assume A.1)-A.3), R) and V). Let $p \geq 1$ be a number such that*

$$\frac{2p-2}{2p} < \sup_{u \in \mathbb{R}^m} \frac{\lambda(u)}{\Lambda(u)}. \quad (4.1)$$

If R is sufficiently small then for any two concentric balls $B_\rho \subset B_R$ with center in $\bar{\Omega}$ there is a constant $C(T)$ such that the following holds for all intergers $k \geq 1$

$$\begin{aligned} & \sup_{t \in (0, T)} \int_{B_\rho \cap \Omega} |Du_k|^{2p} dx + \iint_{Q_\rho} \lambda(u_{k-1}) |Du_k|^{2p-2} |D^2 u_k|^2 dz \leq \\ & C_1(T) \iint_{Q_R} \left[\lambda(u_1) |Du_2|^{2p-2} |D^2 u_2|^2 + \frac{|A_u(u_1)|^2}{\lambda(u_1)} |Du_2|^{2p+2} \right] dz \\ & + C_1(T) \frac{1}{(R-\rho)^2} \max_{1 \leq i \leq k} \iint_{Q_R} \lambda((u_{k-1})) |Du_k|^{2p} dz. \end{aligned} \quad (4.2)$$

Here, $Q_R = (B_R \cap \Omega) \times (0, T)$.

Before going to the proof, we recall the following elementary fact in [3, Lemma 2.1].

Lemma 4.2 *Assume the ellipticity condition A). Let α be a number such that there is $\delta_\alpha \in (0, 1)$ such that $\frac{\alpha}{2+\alpha} = \delta_\alpha \frac{\lambda}{\Lambda}$. We then have*

$$AD\zeta D(\zeta|\zeta|^\alpha) \geq \hat{\lambda} \zeta |\zeta|^\alpha |D\zeta|^2, \quad \hat{\lambda} = (1 - \delta_\alpha^2) \lambda. \quad (4.3)$$

Furthermore, since u_{k-1}, u_k are C^2 in x , we can differentiate (2.6) with respect to x to get

$$(Du_k)_t = \operatorname{div}((A(u_{k-1})D^2 u_k + A_u(u_{k-1})Du_{k-1}Du_k) + Df(u_{k-1}, Du_k)) \quad k \geq 1. \quad (4.4)$$

Proof: (Proof of Lemma 4.1) We consider the interior case $B_\rho \subset B_R \subset \Omega$ and leave the boundary case, when the center of B_R is on the boundary $\partial\Omega$, to Remark 4.4 following the proof. For any s, t such that $0 \leq s < t \leq R$ let ψ be a cutoff function for B_s, B_t . That is, $\psi \equiv 1$ in B_s and $\psi \equiv 0$ outside B_t with $|D\psi| \leq 1/(t-s)$. Testing (4.4) with $|Du_k|^{2p-2} Du_k \psi^2$. The assumption (4.1) shows that $\alpha = 2p-2$ satisfies the condition of Lemma 4.2 so that we can find a positive constant $C(p)$ such that

$$\begin{aligned} & \sup_{\tau \in (0, T)} \int_{\Omega} |Du_k|^{2p} \psi^2 dx + C(p) \iint_Q \lambda(u_{k-1}) |Du_k|^{2p-2} |D^2 u_k|^2 \psi^2 dz \leq \\ & - \iint_Q [A_u(u_{k-1}) Du_{k-1} Du_k D(|Du_k|^{2p-2} Du_k \psi^2) + Df(u_{k-1}, Du_k) |Du_k|^{2p-2} Du_k \psi^2] dz. \end{aligned}$$

For simplicity, we will assume in the sequel that $f \equiv 0$. The presence of f will be discussed in Remark 4.3 after the proof. Therefore,

$$\begin{aligned} & \sup_{\tau \in (0, T)} \int_{\Omega} |Du_k|^{2p} \psi^2 dx + C(p) \iint_Q \lambda(u_{k-1}) |Du_k|^{2p-2} |D^2 u_k|^2 \psi^2 dz \leq \\ & C_0(p) \iint_Q |A_u(u_{k-1})| |Du_{k-1}| |Du_k|^{2p-1} |D^2 u_k| \psi^2 + |A_u(u_{k-1})| |Du_{k-1}| |Du_k|^{2p} \psi |D\psi| dz. \end{aligned}$$

Let $\Phi_0^2(u) = \lambda(u)$ and $\Phi^2(u) = \frac{|A_u(u)|^2}{\lambda(u)}$ as in A.2). Applying Young's inequality to the integrand of the first integral on the right of the above and the following

$$\begin{aligned} |A_u(u_{k-1})||Du_{k-1}||Du_k|^{2p}\psi|D\psi| &= \Phi(u_{k-1})|Du_{k-1}||Du_k|^{2p}\psi|\Phi_0||Du_k|^{2p}|D\psi| \\ &\leq \Phi^2(u_{k-1})|Du_{k-1}|^2|Du_k|^{2p}\psi^2 + |D\psi|^2\Phi_0^2(u_{k-1})|Du_k|^{2p}, \end{aligned}$$

we easily deduce

$$\begin{aligned} \sup_{\tau \in (0, T)} \int_{\Omega} |Du_k|^{2p}\psi^2 dx + \iint_Q \Phi_0^2(u_{k-1})|Du_k|^{2p-2}|D^2u_k|^2\psi^2 dz &\leq \\ C_1 \iint_Q \Phi^2(u_{k-1})|Du_{k-1}|^2|Du_k|^{2p}\psi^2 dz + \sup |D\psi|^2 \iint_{Q_t} \Phi_0^2(u_{k-1})|Du_k|^{2p} dz. \end{aligned}$$

Here, we denoted $Q_t = B_t \times (0, T)$. Again, a use of Young's inequality to the first integral on the right yields

$$\begin{aligned} \sup_{\tau \in (0, T)} \int_{\Omega} |Du_k|^{2p}\psi^2 dx + \iint_Q \Phi_0^2(u_{k-1})|Du_k|^{2p-2}|D^2u_k|^2\psi^2 dz &\leq \\ C_2 \iint_Q \Phi^2(u_{k-1})(|Du_{k-1}|^{2p+2} + |Du_k|^{2p+2})\psi^2 dz + C \sup |D\psi|^2 \iint_{Q_t} \Phi_0^2(u_{k-1})|Du_k|^{2p} dz. \end{aligned}$$

By the choice of ψ , we obtain from the above the following

$$\sup_{\tau \in (0, T)} \int_{B_s} |Du_k|^{2p}\psi^2 dx + H(s) \leq C_2(B_0(t) + B_1(t)) + C \frac{1}{(t-s)^2} G(t). \quad (4.5)$$

Here, for any fixed integer $k \geq 1$, we set

$$H(t) = \iint_{Q_t} \Phi_0^2(u_{k-1})|Du_k|^{2p-2}|D^2u_k|^2 dz, \quad B_1(t) = \iint_{Q_t} \Phi^2(u_{k-1})|Du_k|^{2p+2} dz,$$

and

$$B_0(t) = \iint_{Q_t} \Phi^2(u_{k-1})|Du_{k-1}|^{2p+2} dz, \quad G(t) = \iint_{Q_t} \Phi_0^2(u_{k-1})|Du_k|^{2p} dz.$$

We now apply Lemma 3.4 for $u = u_{k-1}$ and $U = u_k$. We will see that our assumptions A.2) and A.3) imply the assumptions GN.1) and GN.2) of Lemma 3.4 for any $p \geq 1$. Indeed, by our assumption (2.2) on Φ_0, Φ in A.2) the constants in of GN.1) are finite. Furthermore, since u_{k-1} is BMO with uniform bounded norm and the assumption A.3), $\Phi^{\frac{2}{3}}(u_{k-1})$ belongs to the $A_{\frac{4}{3}}$ class. As $\frac{2}{3} \geq \frac{2}{p+2}$ and $\frac{4}{3} \leq \frac{p}{p+2} + 1$, $\Phi^{\frac{2}{p+2}}(u_{k-1})$ belongs to the $A_{\frac{p}{p+2}+1}$ class. Thus, the quantity $C(\Phi, \Phi_0)$ defined in (3.24) is finite. Also, our continuity assumption (2.7) on the BMO norm of u_k implies the smallness of $C(\Phi, \Phi_0)\|u_k\|_{BMO(B_R)}$ if R is small. Hence, for any given $\mu_1 > 0$ if $R = R(\mu_1) > 0$ is sufficiently small then we have from (3.21) of Lemma 3.4 the following estimate.

$$\begin{aligned} \int_{B_s} \Phi^2(u_{k-1})|Du_k|^{2p+2} dx &\leq \mu_1 \int_{B_t} \Phi_0^2(u_{k-1})|Du_k|^{2p-2}|D^2u_k|^2 dx + \\ &\mu_1 \left[\int_{B_t} \Phi^2(u_{k-1})|Du_{k-1}|^{2p+2} dx + \frac{1}{(t-s)^2} \int_{B_t} \Phi_0^2(u_{k-1})|Du_k|^{2p} dx \right]. \end{aligned} \quad (4.6)$$

Then (4.6) and (4.5) give a positive constant C_2 , which is redefined and can depend on k_1, k_2 , such that

$$B_1(s) \leq \mu_1[H(t) + B_0(t) + \frac{1}{(t-s)^2}G(t)] \quad \rho < s < t < R, \quad (4.7)$$

$$H(s) \leq C_2B_1(t) + C_2B_0(t) + \frac{1}{(t-s)^2}G(t) \quad \rho < s < t < R. \quad (4.8)$$

Let $t' = s + (t-s)/2$. Using (4.7) with s being t' in the inequality (4.8) with t being t' and the fact that H, B_0, G are increasing, we get

$$H(s) \leq C_2\mu_1H(t) + C_2(\mu_1 + 1)B_0(t) + \frac{4(\mu_1 + 1)}{(t-s)^2}G(t) \quad \rho < s < t < R. \quad (4.9)$$

We can assume that $\mu_2 = C_2\mu_1 < 1$. By Lemma 3.5, (4.9) yields

$$H(\rho) \leq C_3[C_2(\mu_1 + 1)B_0(R) + \frac{4(\mu_1 + 1)}{(R-\rho)^2}G(R)].$$

Here, $C_3 = (1-\nu)^{-2}(1-\nu^{-2}\mu_2)^{-1}$ for any ν satisfying $\nu^{-2}\mu_2 < 1$. Obviously, the above also hold with ρ, R replaced by s, t with $\rho \leq s < t \leq R$. So,

$$H(s) \leq C_3[C_2(\mu_1 + 1)B_0(t) + \frac{4(\mu_1 + 1)}{(t-s)^2}G(t)]. \quad (4.10)$$

We now let $t' = (s+t)/2$ and use (4.7) with t being t' and then (4.10) with s being t' to see that

$$\begin{aligned} B_1(s) &\leq \mu_1[H(t') + B_0(t') + \frac{1}{(t'-s)^2}G(t')] \\ &\leq \mu_1[C_3[C_2(\mu_1 + 1)B_0(t) + \frac{4(\mu_1 + 1)}{(t-t')^2}G(t)] + B_0(t') + \frac{1}{(t'-s)^2}G(t'). \end{aligned}$$

Since B_1, G are increasing functions, the above yields

$$B_1(s) \leq \mu_3B_0(t) + C_4\frac{1}{(t-s)^2}G(t), \quad (4.11)$$

where $\mu_3 = \mu_1(C_3C_2(\mu_1 + 1) + 1)$ and $C_4 = 4(4\mu_1(\mu_1 + 1) + \mu_1)$.

We now consider B_0 . Applying Lemma 3.4 with $u = U = u_{k-1}$, so that $I_1 = \hat{I}_1$, and $\Phi(u) = \Phi_0(u)$, we see that if $\|u_{k-1}\|_{BMO(B_R)}$, or R , is sufficiently small then there is a constant $C_0(\Phi, \Phi_0)$ such that for any s, t satisfying $0 < s < t < R$

$$I_1(s) \leq C_0(\Phi)\|u_{k-1}\|_{BMO(B_R)} \left[\frac{1}{(t-s)^2} \iint_{Q_t} |\Phi(u_{k-1})|^2 |Du_{k-1}|^{2p} dz + I_2(t) \right] \quad (4.12)$$

with

$$\begin{aligned} I_1(t) &= \iint_{Q_t} \Phi^2(u_{k-1}) |Du_{k-1}|^{2p+2} dz = B_0(t), \\ I_2(t) &= \iint_{Q_t} \Phi^2(u_{k-1}) |Du_{k-1}|^{2p-2} |D^2u_{k-1}|^2 dz. \end{aligned}$$

Going in back to the notation $\Phi_0(u) = \lambda^{\frac{1}{2}}(u)$, by (2.8), we can split Q_t into two disjoint sets

$$\begin{aligned} Q_{(1)} &:= \{(x, t) \in Q_t : \Phi(u_{k-1})(x, t) \leq C(T)\Phi(u_{k-2})(x, t)\}, \\ Q_{(2)} &:= \{(x, t) \in Q_t : \Phi_0(u_{k-1})(x, t) \leq C(T)\Phi_0(u_{k-2})(x, t)\}. \end{aligned}$$

Recall that $\Phi(u) \leq k_2\Phi_0(u)$ by A.2). On $Q_{(1)}$, we have $\Phi(u_{k-1}) \leq C(T)\Phi(u_{k-2}) \leq C(T)k_2\Phi_0(u_{k-2})$. Meanwhile, on $Q_{(2)}$, $\Phi(u_{k-1}) \leq k_2\Phi_0(u_{k-1}) \leq C(T)k_2\Phi_0(u_{k-2})$. Thus,

$$\begin{aligned} I_2(t) &= \iint_{Q_t} \Phi^2(u_{k-1})|Du_{k-1}|^{2p-2}|D^2u_{k-1}|^2 dz \\ &\leq C(T)k_2 \iint_{Q_t} \Phi_0^2(u_{k-2})|Du_{k-1}|^{2p-2}|D^2u_{k-1}|^2 dz. \end{aligned}$$

Similarly,

$$\iint_{Q_t} |\Phi(u_{k-1})|^2|Du_{k-1}|^{2p} dz \leq C(T)k_2 \iint_{Q_t} |\Phi_0(u_{k-2})|^2|Du_{k-1}|^{2p} dz. \quad (4.13)$$

Using these estimates in (4.12), we obtain

$$B_0(s) \leq C_1(\Phi, \Phi_0, T)\|u_{k-1}\|_{BMO(B_R)} \left[\frac{1}{(t-s)^2} G_0(t) + H_0(t) \right],$$

where

$$\begin{aligned} G_0(t) &= \iint_{Q_t} |\Phi_0(u_{k-2})|^2|Du_{k-1}|^{2p} dz, \\ H_0(t) &= \iint_{Q_t} \Phi_0^2(u_{k-2})|Du_{k-1}|^{2p-2}|D^2u_{k-1}|^2 dz. \end{aligned}$$

Using the above estimate for B_0 in (4.10) and (4.11) and adding the results, we can easily see that if $\|u_{k-1}\|_{BMO(B_R)}$ is sufficiently small then

$$H(s) + B_1(s) \leq \mu_4 H_0(t) + \frac{C_5}{(t-s)^2} [G(t) + G_0(t)], \quad (4.14)$$

for some C_5 depends on $\Phi, \Phi_0, k_1, k_2, T$ and

$$\mu_4 = C_1(\Phi, \Phi_0, T)\|u_{k-1}\|_{BMO(B_R)} [C_3[C_2(\mu_1 + 1) + \mu_3].$$

We now define

$$\begin{aligned} \mathcal{B}_k(t) &= \iint_{Q_t} [\lambda(u_{k-1})|Du_k|^{2p-2}|D^2u_k|^2 + \Phi^2(u_{k-1})|Du_k|^{2p+2}] dz, \\ \mathcal{G}_k(t) &= \iint_{Q_t} [\lambda(u_{k-1})|Du_k|^{2p} + \lambda(u_{k-2})|Du_{k-1}|^{2p}] dz. \end{aligned}$$

We then have from (4.14) that

$$\mathcal{B}_k(s) \leq \mu_4 \mathcal{B}_{k-1}(t) + \frac{C_5}{(t-s)^2} \mathcal{G}_k(t). \quad (4.15)$$

As before, we can assume that R is sufficiently small such that $\mu_4 < 1$. For any $a \in (0, 1)$ such that $\mu_4 a^{-2} < 1$ we define the sequences $t_0 = \rho$ and $t_{i+1} = t_i + (1-a)a^i(R-\rho)$. Iterate the above $k-2$ times to get

$$\begin{aligned} \mathcal{B}_k(\rho) &\leq \mu_4^k \mathcal{B}_2(t_{k-2}) + \sum_{i=0}^{k-2} \mu_4^i a^{-2i} \frac{C_5}{(1-a)^2 (R-\rho)^2} \mathcal{G}_{k-i}(t_{i+1}) \\ &\leq \mathcal{B}_2(R) + \frac{C_6(a, \mu_4)}{(R-\rho)^2} \max_{2 \leq i \leq k} \mathcal{G}_i(R). \end{aligned}$$

This shows that the quantity

$$\iint_{Q_t} [\lambda(u_{k-1}) |Du_k|^{2p-2} |D^2 u_k|^2 + \Phi^2(u_{k-1}) |Du_k|^{2p+2}] dz, \quad t \geq \rho, \quad k \geq 1$$

can be bounded by

$$C_1(T) \iint_{Q_R} \left[\lambda(u_1) |Du_2|^{2p-2} |D^2 u_1|^2 + \frac{|A_u(u_1)|^2}{\lambda(u_1)} |Du_2|^{2p+2} + \frac{1}{(R-\rho)^2} \max_{1 \leq i \leq k} \lambda((u_{k-1})) |Du_k|^{2p} \right] dz.$$

Using this and (4.13) in (4.5), and the estimate for $B_0(t)$, we obtain (4.2) of the lemma. \blacksquare

Remark 4.3 If $f \neq 0$ then the growth assumption F) gives

$$|f(u_{k-1}, Du_k)| \leq C |Du_k| + C |u_{k-1}|^b + C. \quad (4.16)$$

$$|Df(u_{k-1}, Du_k)| \leq C |D^2 u_k| + C |u_{k-1}|^{b-1} |Du_{k-1}| + C. \quad (4.17)$$

Testing (4.4) with $|Du_k|^{2p-2} Du_k \psi^2$, we will have the extra term $Df(u_{k-1}, Du_k) |Du_k|^{2p-1} \psi^2$ on the right of our estimates in the proof. For any positive $\varepsilon > 0$ we can use Young's inequality to have

$$\begin{aligned} Df(u_{k-1}, Du_k) |Du_k|^{2p-1} &\leq C (|D^2 u_k| + |u_{k-1}|^{b-1} |Du_{k-1}| + C) |Du_k|^{2p-1} \\ &\leq \varepsilon |Du_k|^{2p-2} |D^2 u_k|^2 + \\ &\quad C(p, \varepsilon) [\Phi^{-\frac{kp}{p+1}} |u_{k-1}|^{k_p(b-1)} + \Phi^2 |Du_{k-1}|^{2p+2} + |Du_k|^{2p}], \end{aligned}$$

where $\frac{1}{k_p} = 1 - \frac{2p-1}{2p} - \frac{1}{2p+2}$. Since u_k, u_{k-1} are BMO and Φ is bounded from below, the integral of the first term on the right of the above inequalities is bounded. (4.8) now becomes

$$H(s) \leq \varepsilon H(t) + C_2 B_1(t) + C_2 B_0(t) + \frac{1}{(t-s)^2} G(t) + C(\varepsilon) \quad \rho < s < t < R. \quad (4.18)$$

Choosing ε small, we can see that the iteration arguments in the proof are still in force and the proof can continue.

Remark 4.4 We discuss the case when the centers of B_ρ, B_R are on the boundary $\partial\Omega$. Let us first consider the case u satisfies the Dirichlet condition $u = 0$ on $\partial\Omega$. By flattening the boundary we can assume that $B_R \cap \Omega$ is the set

$$B^+ = \{x : x = (x_1, \dots, x_n) \text{ with } x_n \geq 0 \text{ and } |x| < R\}.$$

For any point $x = (x_1, \dots, x_n)$ we denote by \bar{x} its reflection across the plane $x_n = 0$, i.e., $\bar{x} = (x_1, \dots, -x_n)$. Accordingly, we denote by B^- the reflection of B^+ . For $u = u_k$ we define the odd reflection of u by \bar{u} , i.e. $\bar{u}(x, t) = -u(\bar{x}, t)$ for $x \in B^-$. We then consider the odd extension U in $B = B^+ \cup B^-$ of u

$$U(x, t) = \begin{cases} u(x, t) & \text{if } x \in B^+, \\ \bar{u}(x, t) & \text{if } x \in B^-. \end{cases}$$

It is easy to see that \bar{u} satisfies in B^- a system similar to (4.4) for u in B^+ . As in the proof of the lemma, we test the system for u_k with $|Du_k|^{2p-2} Du_k \psi^2$ and the system for \bar{u} with $|D\bar{u}_k|^{2p-2} D\bar{u}_k \psi^2$ and then sum the results. The integration parts results the extra boundary terms along the flat boundary parts ∂B^+ and ∂B^- . Using the facts that either $D_{x_i} u = D_{x_i} \bar{u} = 0$ for $i \neq n$ or $D_{x_n} u = D_{x_n} \bar{u}$ and the outward normal vectors of B^+ and B^- are opposite we can easily see that those boundary terms are either zero or cancel each others in the summation. Thus, we can obtain (4.5) again with u_{k-1}, u_k being replaced by U_{k-1}, U_k . Since U_k belong to $W^{2,\infty}(B)$ the argument can continue and the lemma holds for U_k and then u_k .

The same argument applies for the Neumann boundary condition if we use the even extension for u_k .

We now give the proof of Theorem 2.3.

Proof: We test the systems (2.6) with u_k and use Young's inequality to have

$$\sup_{t \in [0, T]} \int_{\Omega} |u|^2 dx + \iint_Q \lambda(u_{k-1}) |Du_k|^2 dz \leq C \iint_Q [|u_{k-1}|^2 + |u_{k-1}|^{b+1} + 1] dz.$$

The uniform bound assumption on the BMO norms of u_{k-1} yields that the right hand side is bounded uniformly for all k . Thus, there is a constant C such that

$$\iint_Q \lambda(u_{k-1}) |Du_k|^2 dz \leq C \quad \forall k.$$

Now, for any $0 < \rho < R$ and concentric balls B_ρ, B_R with centers in $\bar{\Omega}$ let us assume that there is some $p \geq 1$ such that there is a constant $C_0(\rho, R, u_0)$ depending on ρ, R and $\sup_{t \in (0, T)} \|u_0(\cdot, t)\|_{C^1(\Omega)}$ on such that ($Q_R = B_R \times (0, T)$)

$$\iint_{Q_\rho} \lambda(u_{k-1}) |Du_k|^{2p} dz \leq C_0(\rho, R, u_0) \quad \forall k. \quad (4.19)$$

It is well known that the C^1 norms of u_1 and u_2 can be bounded by that of u_0 . Now, if p satisfies (4.1) then Lemma 4.1 and (4.19) establish the existence of a constant $C_1(\rho, R)$ such that the following holds for all integers k

$$\sup_{t \in (0, T)} \int_{B_\rho} |Du_k|^{2p} dx + \iint_{Q_\rho} \Phi_0^2(u_{k-1}) |Du_k|^{2p-2} |D^2 u_k|^2 dz \leq C_1(\rho, R, u_0) \quad (4.20)$$

if $0 < \rho < R$ and R is sufficiently small.

Let χ_0 be any number such that $1 < \chi_0 < 1 + \frac{2}{n}$. Denote $V = |Du_k|^p$ and use Hölder's inequality to get

$$\iint_Q \lambda V^{2\chi_0} dz \leq \left(\iint_Q \lambda^r dz \right)^{\frac{1}{r}} \left(\iint_Q V^{2(1+\frac{2}{n})} dz \right)^{\frac{1}{r}}$$

where $\lambda = \lambda(u_{k-1})$ and r is a number such that $r'\chi_0 = 1 + \frac{2}{n}$.

Recall the Sobolev imbedding inequality

$$\|V\|_{L^{\frac{2(n+2)}{n}}(Q)} \leq C \sup_t \|V(\cdot, t)\|_{L^2(\Omega)} + C \left(\iint_Q |DV|^2 dz \right)^{\frac{1}{2}}$$

and the fact that u_k is BMO so that $\lambda(u_{k-1})$ belongs to $L^r(\Omega)$ for any $r > 1$ (see [6]). The above estimates for $Q = Q_\rho$ show that there is a constant $C(\rho)$ such that

$$\iint_{Q_\rho} \lambda V^{2\chi_0} dz \leq C(\rho) \left[\sup_t \|V(\cdot, t)\|_{L^2(B_\rho)} + \left(\iint_{Q_\rho} |DV|^2 dz \right)^{\frac{1}{2}} \right]. \quad (4.21)$$

From the ellipticity condition A) and (4.20) we see that the right hand side is bounded. Hence

$$\iint_Q \lambda(u_{k-1}) |Du_k|^{2p\chi_0} dz \leq C_2(\rho, R, u_0) \quad \forall k.$$

Therefore, (4.19) holds again with p now is $p\chi_0$. We already showed that (4.19) is valid for $p = 1$. Thus, we can repeat the argument k times until $2\chi_0^k > n$ as long as the ratio condition (4.1) of Lemma 4.1 is verified for $p = \chi_0^k$. The assumption R) shows that we can choose χ_0, k such that $1 < \chi_0 < 1 + \frac{2}{n}$, $2\chi_0^k > n$ and the ratio condition (4.1) holds for $p = \chi_0^k$. Therefore, (4.20) holds for $2p = 2\chi_0^k$. We now cover Ω with finitely many balls of radius $R/2$ to obtain

$$\sup_{t \in (0, T)} \int_\Omega |Du_k|^{2p} dx + \iint_Q \Phi_0^2(u_{k-1}) |Du_k|^{2p-2} |D^2u_k|^2 dz \leq C(\Omega, T, u_0). \quad (4.22)$$

For each $t \in (0, T)$, (4.22) shows that the norms $\|u_k(\cdot, t)\|_{W^{1, 2p}(\Omega)}$ for some $2p > n$ are bounded uniformly in t by a constant depending only on the size of Ω, T and u_0 . By Sobolev's imbedding theorem $\{u_k(\cdot, t)\}$ is a bounded sequence in $C^\alpha(\Omega)$ for some $\alpha > 0$. From the system for u_k , (4.22) with $p = 1$ also shows that $\|(u_k)_t\|_{L^2(Q)}$ is uniformly bounded. Together, these facts show that the solutions u_k are uniformly Hölder continuous in (x, t) and that $\{u_k\}$ is bounded in $C^\beta(Q)$ for some $\beta > 0$. We then see that there is a relabeled subsequence $\{u_k\}$ converges in $C^0(Q)$ to some u . Using difference quotient in t we see that $u_t \in L^2(Q)$. The above estimate (4.22) also shows that we can assume $Du_{k+1}(\cdot, t)$ converges weakly to $Du(\cdot, t)$ in $L^2(\Omega)$ for each $t \in (0, T)$. By the continuity of A in its variable u , we see that u weakly solves (1.1).

By the semicontinuity of norms, (4.22) implies

$$\sup_{t \in (0, T)} \int_\Omega |Du|^{2p} dx + \iint_Q \lambda(u) |Du|^{2p-2} |D^2u|^2 dz \leq C(\Omega, u_0). \quad (4.23)$$

Since $2p > n$, the above implies that u is Hölder continuous and its regularity in x . Since u_t is in $L^2(Q)$. It is easy to derive from these facts that u is Hölder in (x, t) . By [?], Du is Hölder in (x, t) and then u is a classical solution. ■

We now turn to the proof of our second theorem.

Proof: (*Proof of Theorem 2.5*) Let u be the classical solution of the system (1.1) in $\Omega \times (0, T_0)$. We can differentiate (1.1) to have

$$(Du)_t = \operatorname{div}((A(u)D^2u + A_u(u)DuDu) + Df(u, Du)). \quad (4.24)$$

For any s, t such that $0 < s < t < R$ let ψ be a cutoff function for two concentric balls B_s, B_t with centers in $\bar{\Omega}$. That is, $\psi \equiv 1$ in B_s and $\psi \equiv 0$ outside B_t with $|D\psi| \leq 1/(t-s)$. As in the proof of the previous theorem, we test (4.24) with $|Du|^{2p-2}Du\psi^2$. Since $u \in C^{1,2}(\bar{\Omega} \times (0, T_0))$, the local Gagliardo-Nirenberg inequality of Lemma 3.3 applies here for $U = u$. Thus, as in the proof of Lemma 4.1, we can use Lemma 3.4 to obtain

$$\begin{aligned} \int_{B_s} \Phi^2 |Du|^{2p+2} dx &\leq \mu_1 \int_{B_t} \Phi_0^2 |Du|^{2p-2} |D^2u|^2 dx + \\ &\mu_1 \left[\int_{B_t} \Phi^2 |Du|^{2p+2} dx + \frac{1}{(t-s)^2} \int_{B_t} \Phi^2 |Du|^{2p} dx \right]. \end{aligned} \quad (4.25)$$

Here, $\Phi_0^2 = \lambda(u)$ and $\Phi^2 = \frac{A^2(u)}{\lambda(u)}$. We now set

$$\begin{aligned} H(t) &= \iint_{Q_t} \Phi_0^2(u) |Du|^{2p-2} |D^2u|^2 dz, \quad B(t) = \iint_{Q_t} \Phi^2(u) |Du|^{2p+2} dz, \\ G(t) &= \iint_{Q_t} \Phi^2(u) |Du|^{2p} dz. \end{aligned}$$

Because A.4) obviously holds for $u_k = u_{k-1} = u$, it is clear that the proof of Lemma 4.1 with $H, H, B_0, B_1, \mathcal{B}_k$, and G being replaced by the new definitions, and $B_0 = B_1 = \mathcal{B}_k = B$, now leads to

$$B(s) \leq \mu_4 B(t) + C_4 \frac{1}{(t-s)^2} G(t), \quad 0 < \rho < s < t < R, \quad (4.26)$$

where $\mu_4 < 1$. For any $a \in (0, 1)$ such that $\mu_4 a^{-2} < 1$ we define the sequences $t_0 = \rho$ and $t_{i+1} = t_i + (1-a)a^i(R-\rho)$. Iterate the above to get

$$B(\rho) \leq \mu_4^k B(t_k) + \sum_{i=0}^{k-1} \mu_4^i a^{-2i} \frac{C_4}{(1-a)^2 (R-\rho)^2} G(t_k).$$

Let k tend to infinity and use the fact that $\mu_4 \in (0, 1)$ and $B(R)$ is finite to get

$$B(\rho) \leq \frac{C_5(a, \mu_4)}{(R-\rho)^2} G(R).$$

We now see that a similar argument in the proof of Theorem 2.3 with u_k being u now gives

$$\sup_{t \in (0, T)} \int_{B_\rho} |Du|^{2p} dx + \iint_{Q_\rho} \Phi_0^2 |Du|^{2p-2} |D^2u|^2 dz \leq C_1(\rho, R) \quad (4.27)$$

if $0 < \rho < R$ and R is sufficiently small and some p such that $2p > n$. Finite covering Ω with balls $B_{R/2}$ yields

$$\sup_{t \in (0, T)} \int_{\Omega} |Du|^{2p} dx + \iint_Q \lambda(u) |Du|^{2p-2} |D^2u|^2 dz \leq C(\Omega, R). \quad (4.28)$$

Hence u is Hölder continuous and its regularity in x . From the system for u and the above, with $p = 1$, we see that u_t is in $L^2(Q)$. It is now standard to show that u is Hölder in (x, t) and Du is Hölder continuous. We now can refer to Amann's results to see that u exists globally. ■

References

- [1] H. Amann. Dynamic theory of quasilinear parabolic equations II. Reaction-diffusion systems, *Differential Integral Equations*, Vol. 3, no. 1 (1990), pp. 13-75.
- [2] H. Amann. Dynamic theory of quasilinear parabolic systems III. Global existence, *Math Z.* 202 (1989), pp. 219-250.
- [3] S. Ahmad and D. Le. Global and Blow Up Solutions to Cross Diffusion Systems. Nonlinear Analysis Series A: TMA. In press.
- [4] A. Friedman. *Partial Differential Equations*, New York, 1969.
- [5] B. Franchi, C. Perez and R. L. Wheeden. Self-Improving Properties of John Nirenberg and Poincaré Inequalities on Spaces of Homogeneous Type. *J. Functional Analysis*, 153, 108–146, 1998.
- [6] E. Giusti. *Direct Methods in the Calculus of Variations*. World Scientific, 2003.
- [7] R. L. Johnson and C. J. Neugebauer. Properties of BMO functions whose reciprocals are also BMO. *Z. Anal. Anwendungen*, 12(1):3-11, 1993.
- [8] D. Le. Regularity of BMO weak solutions to nonlinear parabolic systems via homotopy. *Trans. Amer. Math. Soc.* 365 (2013), no. 5, 2723–2753.
- [9] D. Le. Global existence results for near triangular nonlinear parabolic systems. *Adv. Nonlinear Studies.* 13 (2013), no. 4, 933-944.
- [10] D. Le, L. Nguyen and T. Nguyen. Coexistence in Cross Diffusion systems. *Indiana Univ. J. Math.* Vol.56, No. 4, pp.1749-1791, 2007.
- [11] J. Oróbitg and C. Pérez. A_p weights for nondoubling measures in \mathbb{R}^n and applications. newblock *Transactions of the American mathematical society*, 354 (2002), 2013-2033.
- [12] P. Strzelecki. Gagliardo Nirenberg inequalities with a BMO term. *Bull. London Math. Soc.* Vol. 38, pp. 294-300, 2006.
- [13] N. Shigesada, K. Kawasaki and E. Teramoto. *Spatial segregation of interacting species*. *J. Theor. Biol.*, 79(1979), 83– 99.

- [14] E. M. Stein. *Harmonic Analysis, Real Variable Methods, Orthogonality and Oscillatory Integrals*. Princeton Univ. Press, Princeton, NJ, 1993.