

Weak singularities of 3-D Euler equations and restricted regularity of Navier Stokes equation solutions with time dependent force terms

Jörg Kampen

October 16, 2019

Abstract

A class of singular solutions of the 3-D Euler equation corresponds to a class of singular solutions of the incompressible Navier Stokes equation with time dependent force terms, as the time-dependent force term can be chosen such that it cancels the viscosity term of the incompressible Navier Stokes equation. In this context classical solution branches of the three dimensional incompressible Euler equation are constructed where a vorticity component blows up at some point after finite time for regular velocity component data. Furthermore, we show that there are classical solution branches with $C^k \cap H^k$ -data for $k \geq 2$ which develop weak singularities (kinks) at some point of space time in $C^{k-1} \setminus C^k$ for any $k \geq 2$ after finite time. The time-local solution branches of the Euler equation are obtained by viscosity limits of solutions of Navier-Stokes-type viscosity (ν)-extensions of time-reversed Euler-type equations. Here, Lipschitz continuity of the Leray projection term and the Burgers term and Lipschitz continuity of spatial first order derivatives of the Leray projection term and the Burgers term are used in local time iteration schemes in terms of convolutions with the Gaussian of dispersion ν and first order spatial derivatives of this Gaussian. For the time reversed equations and their Navier Stokes type equation data with singular vorticity at the origin and strong polynomial decay are used. The functional solution increment and derivatives up to some order, i.e. the solution minus the spatially convoluted data and spatial derivatives of this increment up to some order, preserve strong polynomial decay of some order. This polynomial decay of the local time solution is strong enough such that compactness arguments can be constructed in classical Banach spaces. The Lipschitz continuity of the first order derivatives of the Leray projection term and of the Burgers term implies that regularity is gained after any finite time for the time-reversed equation. This result can be sharpened in the sense that the second order derivatives of the Leray projection term of a local solution are Lipschitz even for a viscosity limit. As a consequence, the original Euler equation has a solution branch which develops singular solutions or kinks from regular data after some finite time.

2010 Mathematics Subject Classification. 35Q31, 76N10

1 Idea of construction and statement of weak singularity theorems

We determine data with weakly singular short-time solutions of the three dimensional incompressible Euler equation on the whole Euclidean space. Let v_i , $1 \leq i \leq D$ denote the velocity component functions in dimension $D = 3$, and let

$$\omega = \text{curl}(v) = \left(\frac{\partial v_3}{\partial x_2} - \frac{\partial v_2}{\partial x_3}, \frac{\partial v_1}{\partial x_3} - \frac{\partial v_3}{\partial x_1}, \frac{\partial v_2}{\partial x_1} - \frac{\partial v_1}{\partial x_2} \right) \quad (1)$$

denote the corresponding vorticity functions. The incompressible Euler equation in vorticity form is

$$\frac{\partial \omega}{\partial \tau} + v \cdot \nabla \omega = \frac{1}{2} (\nabla v + \nabla v^T) \omega, \quad (2)$$

and for the corresponding Cauchy problem this equation has to be solved with some initial data $\text{curl}(f)$ at time $\tau = 0$ of some function $f = (f_1, f_2, f_3)^T$ with $f_i \in \mathbb{R}^3 \rightarrow \mathbb{R}$. It is well-known that (for sufficiently regular ω) the velocity is determined by

$$v(\tau, x) = \int_{\mathbb{R}^3} K_3(x-y) \omega(\tau, y) dy, \text{ where } K_3(x)h = \frac{1}{4\pi} \frac{x \times h}{|x|^3}. \quad (3)$$

This Biot-Savart law is also important for our construction of a viscosity limit. This limit is not unique and is a local time solution of the Euler vorticity equation which is Lipschitz and of some order of polynomial decay at spatial infinity. The Biot-Savart law can then be imposed in order to obtain Lipschitz continuity (and some order of polynomial decay at spatial infinity) for first order spatial derivatives of the velocity components, which in turn leads to higher regularity of the velocity solution increment (velocity solution minus initial data) for a viscosity limit. In order to construct local-time and long-time singularities we consider first a related equation which is obtained from (2) by the simple time transformation

$$\tau = -t, \omega_i^-(t, \cdot) = \omega_i(\tau, \cdot), v_i^-(t, \cdot) = v_i(\tau, \cdot), 1 \leq i \leq D. \quad (4)$$

Spatial derivatives are untouched, so multiplying the spatial part of transformed equation by -1 and writing the equation in coordinates we have

$$\frac{\partial \omega_i^-}{\partial t} - \sum_{j=1}^3 v_j^- \frac{\partial \omega_i^-}{\partial x_j} = - \sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^-}{\partial x_j} + \frac{\partial v_j^-}{\partial x_i} \right) \omega_j^-, 1 \leq i \leq D = 3. \quad (5)$$

We refer to the equation in (5) as the 'time-reversed Euler equation'. We shall observe that the 'Navier-Stokes type' extension of the latter equation with an additional viscosity term, i.e., the equation

$$\frac{\partial \omega_i^{\nu,-}}{\partial t} - \nu \Delta \omega_i^{\nu,-} - \sum_{j=1}^3 v_j^- \frac{\partial \omega_i^{\nu,-}}{\partial x_j} = - \sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-}}{\partial x_j} + \frac{\partial v_j^{\nu,-}}{\partial x_i} \right) \omega_j^{\nu,-}, \quad (6)$$

for $1 \leq i \leq D$ (and with a positive constant $\nu > 0$), has a regular solution on a time interval $[0, T]$ for some $T > 0$ for a certain class of data in H^1 which have

a singular vorticity at some point. Regularity results for this specific class of data $\omega_i^{\nu,-}(0, \cdot) = \omega_i^f$, $1 \leq i \leq D$ lead to the conclusion that short or long-time solution families with parameter $\nu > 0$ have a viscosity limit solution branch

$$\omega_i^-(t, \cdot) = \lim_{\nu \downarrow 0} \omega_i^{\nu,-}(t, \cdot), \quad 1 \leq i \leq D \quad (7)$$

of the time-reversed Euler equation with the same initial data, where some regularity is preserved. The initial data ω_i^f , $1 \leq i \leq D$ of the time-reversed Euler equation are then 'final data' of the original incompressible Euler equation, which, hence, has a vorticity singularity after finite time. If the solution branch constructed for the time-reversed Euler equation is short-time, then the solution for the original Euler equation is short-time with a short-time singularity, and if the solution branch constructed for the time-reversed Euler equation is long-time, then the solution for the original Euler equation is long-time with a long-time singularity. We state these results more precisely in the two theorems at the end of this section. In this paper we shall prove the existence of short time singularities for regular data. The proof of long-time singularities follows from the construction of global regular upper bounds, which imply the existence of global regular solution branches. This is not the main interest of this paper, so we propose a sketch of some possible arguments for the existence of global regular solution branches of the Euler equation in a remark below. In the following as usual let C^∞ be the function space of real-valued functions on \mathbb{R}^D with continuous derivatives of any order, and let H^s denote the Sobolev function space of order $s \geq 0$ (for real numbers s). Furthermore, we denote the space of functions (defined on the domain $[0, T) \times \mathbb{R}^D$) with continuous time derivatives up to order m (nonnegative integer) and multivariate spatial derivatives up to order n (nonnegative integer) by $C^{m,n}([0, T) \times \mathbb{R}^D)$ or just by $C^{m,n}$ if the reference to the domain is known from the context. We have

Theorem 1.1. *Let $D = 3$. There exist data $\omega_i^f = \omega_i(0, \cdot) \in H^2 \cap C^2$, $1 \leq i \leq D$ and a vorticity solution ω_i , $1 \leq i \leq D$ of the three dimensional incompressible Euler equation Cauchy problem such that after some finite time $T > 0$ there is a blow-up of the classical solution, i.e.,*

i) for some $T > 0$ there is a solution function $\omega_i : [0, T) \times \mathbb{R}^D \rightarrow \mathbb{R}$, $1 \leq i \leq D$ in $C^1([0, T), H^2 \cap C^2)$ which satisfies the incompressible Euler equation pointwise on the domain $[0, T) \times \mathbb{R}^D$ in a classical sense with data $\omega_i(0, \cdot) = \omega_i^f$, $1 \leq i \leq D$;

ii) for the solution in item i) we have

$$\sup_{\tau \in [0, T)} |\omega_i(\tau, x)| = \infty, \quad (8)$$

i.e., there is no finite upper bound for the left side of (8) on the time interval $[0, T)$.

Even a long time singularity version of Theorem 1.1 holds. We have

Theorem 1.2. *A stronger version of Theorem 1.1 holds with the same text except for the replacement of the quantor 'after some small finite time $T > 0$ ' by 'after any finite time $T > 0$ '.*

Remark 1.3. Theorem 1.2 follows from Theorem 1.1 if the existence of global regular solution branches of the Euler equation can be shown. Strictly speaking following our argument we need to construct global regular solution branches of the time-reversed Euler equation for regular data. However, the following argument does not depend on the signs of the nonlinear terms such that it suffices to construct global regular solution branches of the incompressible Euler equation. In the next remark we note some simple observations which lead to the construction of global regular solution branches.

Remark 1.4. It is not a simple scaling argument which leads to the conclusion that a certain classes of nonlinear equations have global regular solution branches. In order to construct global regular solution branches we have to exploit some special structure of the equations. Nevertheless the structural features needed are of a quite general type. An essential characteristic of the classes of equations we have in mind is that local time solutions can be represented in terms of convolutions with first order spatial derivatives of the Gaussian. In case of the incompressible Navier Stokes equation the incompressibility condition helps in order to obtain local time representations of the value function itself in terms of convolutions with the first order spatial derivatives of the Gaussian (note that this is not possible for typical blow-up equations as $v_{i,\tau} = v_i^2$, $1 \leq i \leq D$). For the construction of global regular solution branches of the incompressible Euler equation we need the construction of a viscosity limit $\nu \downarrow 0$ in addition. This is possible if the convoluted data of the nonlinear terms in the local time representation of the solution of the Navier Stokes equation and their first order derivatives are Lipschitz. This leads to upper bounds in terms of second moments of the Gaussian, where the upper bound of this second moment constant is an essential difference to other equations which do not have local representations in terms of convolutions with spatial first order derivatives of the Gaussian. Let us now sketch an argument starting with the local time solution of the Navier Stokes equation. We introduce the scaling $\tau = \rho\tau_\rho$ which is also useful later when we consider local contraction results and a their independence from the viscosity $\nu > 0$. Let $v_i^\rho(\tau_\rho, \cdot) = v_i(\tau, \cdot)$. We have

$$v_{i,\tau_\rho} = \rho v_{i,\tau}. \quad (9)$$

Let us assume that for some time $\tau_{\rho 0} \geq 0$ we have

$$v_i^\rho(\tau_{\rho 0}, \cdot) \in H^m \cap C^m, \quad |v_i^\rho(\tau_{\rho 0}, \cdot)|_{H^m \cap C^m} \leq C_m \quad \text{for some } m \geq 2. \quad (10)$$

For the construction of global regular solution branches the parameter $\rho > 0$ will depend only on the spatial $|\cdot|_{H^m \cap C^m}$ -norm and on a constant related to the local L^1 -norms of the (first order derivatives of the) Laplacian kernel, i.e., on

$$C_K := 1 + \max_{1 \leq i \leq D} \int_{B_1(0)} |K_{,i}(y)| dy + \int_{B_1(0)} |K(y)| dy \quad (11)$$

where $B_1(0)$ is the ball of radius 1 around the origin in \mathbb{R}^D . Next we note that local contraction arguments lead to the conclusion of existence of local time solutions in a time interval $[\tau_0, \tau_0 + \Delta]$ for some $\Delta > 0$. For $1 \leq i \leq D$, and multiindices β, γ with $|\beta| = k = |\gamma| + 1$, where $\beta_k = \gamma_k + 1$ and $\beta_j = \gamma_j$ for

$j \in \{1, \dots, D\} \setminus \{k\}$ we have for all $1 \leq i \leq D$ the local time representation

$$\begin{aligned}
D_x^\beta v_i^{\rho, \nu}(\tau_\rho, \cdot) &= D_x^\beta v_i^{\rho, \nu}(t_0, \cdot) *_{sp} G_\nu^\rho(\tau_\rho, \cdot) \\
&- \rho \int_0^{\tau_\rho} \int_{\mathbb{R}^D} \sum_{j=1}^D D_x^\gamma (v_j^{\rho, \nu} v_i^{\rho, \nu})(\sigma, y) \left(\frac{-2(\cdot - y)_k}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho dy d\sigma \\
&+ \rho \int_0^{\tau_\rho} \int_{\mathbb{R}^D} \left(\int_{\mathbb{R}^D} (K_{D,i}(y - z)) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\rho, \nu}}{\partial x_j} \frac{\partial v_j^{\rho, \nu}}{\partial x_m} \right) (\sigma, z) dz \right) \times \\
&\times \left(\frac{-2(\cdot - y)_k}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho(\tau_\rho - \sigma, \cdot - y) dy d\sigma,
\end{aligned} \tag{12}$$

where the first derivatives of the spatially scaled Gaussian G_ν^ρ are given by

$$\begin{aligned}
G_{\nu, k}^\rho(\tau_\rho, z; 0, 0) &= \left(\frac{-2z_k}{4\rho\nu\tau_\rho} \right) \frac{1}{\sqrt{4\pi\rho\nu\tau_\rho}^D} \exp\left(-\frac{|z|^2}{4\rho\nu\tau_\rho}\right) \\
&=: \left(\frac{-2z_k}{4\rho\nu\tau_\rho} \right) G_\nu^\rho(\tau_\rho, z; 0, 0) =: \left(\frac{-2z_k}{4\rho\nu\tau_\rho} \right) G_\nu^\rho(\tau_\rho, z),
\end{aligned} \tag{13}$$

where the latter definition is an abbreviation sometimes used in the following. Now it is interesting that the value functions themselves have a local time representation in terms of convolutions with the first order spatial derivatives of the Gaussian. Indeed, due to incompressibility we have

$$\begin{aligned}
v_i^{\rho, \nu}(\tau_\rho, \cdot) &= v_i^{\rho, \nu}(t_0, \cdot) *_{sp} G_\nu^\rho(\tau_\rho, \cdot) \\
&- \rho \int_0^{\tau_\rho} \int_{\mathbb{R}^D} \sum_{j=1}^D (v_j^{\rho, \nu} v_i^{\rho, \nu})(\sigma, y) \left(\frac{-2(\cdot - y)_i}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho dy d\sigma \\
&+ \rho \int_0^{\tau_\rho} \int_{\mathbb{R}^D} \left(\int_{\mathbb{R}^D} (K_D(y - z)) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\rho, \nu}}{\partial x_j} \frac{\partial v_j^{\rho, \nu}}{\partial x_m} \right) (\sigma, z) dz \right) \times \\
&\times \left(\frac{-2(\cdot - y)_i}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho(\tau_\rho - \sigma; \cdot - y) dy d\sigma.
\end{aligned} \tag{14}$$

Here the incompressibility condition is used in order to rewrite the Burgers term. We may abbreviate

$$B_i^0 * G_\nu^\rho := \int_0^{\tau_\rho} \int_{\mathbb{R}^D} \sum_{j=1}^D (v_j^{\rho, \nu} v_i^{\rho, \nu})(\sigma, y) \left(\frac{-2(\cdot - y)_i}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho dy d\sigma, \tag{15}$$

and

$$\begin{aligned}
L^0 * G_\nu^\rho &:= \int_0^{\tau_\rho} \int_{\mathbb{R}^D} \left(\int_{\mathbb{R}^D} (K_D(y - z)) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\rho, \nu}}{\partial x_j} \frac{\partial v_j^{\rho, \nu}}{\partial x_m} \right) (\sigma, z) dz \right) \times \\
&\times \left(\frac{-2(\cdot - y)_i}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho(\tau_\rho - \sigma; \cdot - y) dy d\sigma,
\end{aligned} \tag{16}$$

which implicitly defines the symbols B_i^0 and L^0 , where we note that the latter does not depend on i . From the equation (12) we may define abbreviations B_i and L_i for the convoluted Burgers and Leray data analogously. Spatial derivatives of these functions are denoted in Einstein notation. We observe that the representation in (14) as well as the representation in (12) have the

parameter ρ as a coefficient. Furthermore for local time regular solutions $v_i^{\tau,\nu}(\tau, \cdot) \in H^m \cap C^m$, $1 \leq i \leq D$ with $m \geq 2$ and $\tau \in [\tau_0, \tau_0 + \Delta]$ the Burgers-and Leray functions B_i^0, L^0 , $1 \leq i \leq D$, B_i, L_i , $1 \leq i \leq D$, and $B_{i,j}, L_{i,j}$, $1 \leq i, j \leq D$ are all Lipschitz such that we get interesting upper bounds of the convolutions

$$B_i^0 * G_{\nu,k}^\rho, L^0 * G_{\nu,k}^\rho, B_i * G_{\nu,k}, L_i * G_{\nu,k}, B_{i,j} * G_{\nu,k}^\rho, L_{i,j} * G_{\nu,k}^\rho. \quad (17)$$

Note that the first order spatial derivatives of the Gaussian are antisymmetric in the sense that

$$\left(\frac{-2y_i}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho(\tau_\rho, y) = \left(\frac{2y_i^-}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho(\tau_\rho, y^-), \quad (18)$$

where $y^- = (y_1^-, \dots, y_D^-)$ with $y_i^- = -y_i$ and $y_j^- = y_j$ for $j \in \{1, \dots, D\} \setminus \{i\}$. For any spatially global Lipschitz continuous function F we have

$$\begin{aligned} & \left| \rho F * G_{\nu,i}^\rho = \rho \int F(\tau_\rho - \sigma, x - y) \left(\frac{-2y_i}{4\nu\rho(\sigma)} \right) G_\nu^\rho(\sigma, y) dy d\sigma \right| \\ &= \left| \rho \int \int_{y_i \geq 0} (F(\tau_\rho - \sigma, x - y) - F(\tau_\rho - \sigma, x - y^-)) \left(\frac{-2y_i}{4\nu\rho(\tau_\rho - \sigma)} \right) G_\nu^\rho(\sigma, y) dy d\sigma \right| \\ &\leq \left| \rho L \int \int_{y_i \geq 0} \left(\frac{4y_i^2}{4\nu\rho\sigma} \right) G_\nu^\rho(\sigma, y) dy d\sigma \right| = 4\rho L M_2, \end{aligned} \quad (19)$$

where M_2 is a finite second moment constant of the Gaussian which is small in the following sense. We consider the case $D = 3$. First note that for any small time $\sigma > 0$ the main mass of the spatial integral is on a ball of radius $\sqrt{\rho\nu}$. Now, given $\nu > 0$ using partial integration we observe

$$\begin{aligned} & \left| \rho L \int \int_{\{y_i \geq 0\} \cap B_{\sqrt{\rho\nu}}} \left(\frac{4y_i^2}{4\nu\rho\sigma} \right) G_\nu^\rho(\sigma, y) dy d\sigma \right| \\ &\leq \left| \rho L \int \left(\frac{y_i^3}{3\nu\rho\sigma} \right) \frac{1}{\sqrt{4\pi\rho\nu\sigma}^D} \exp\left(-\frac{|y|^2}{4\rho\nu\sigma}\right) \Big|_0^{\sqrt{\rho\nu}} d\sigma \right| \\ &+ \left| \rho L \int \int_{\{y_i \geq 0\} \cap B_{\sqrt{\rho\nu}}} \left(\frac{y_i^3}{3\nu\rho\sigma} \right) \left(\frac{-2y_i}{4\nu\rho\sigma} \right) \frac{1}{\sqrt{4\pi\rho\nu\sigma}^D} \exp\left(-\frac{|y|^2}{4\rho\nu\sigma}\right) dy d\sigma \right| \quad (20) \\ &\leq \left| \rho L \int \frac{1}{3\sigma} \sqrt{\nu\rho} \frac{1}{\sqrt{4\pi\rho\nu\sigma}^D} \exp\left(-\frac{1}{4\sigma}\right) d\sigma \right| \\ &+ \left| \rho L \int \int_{\{y_i \geq 0\} \cap B_{\sqrt{\rho\nu}}} \left(\frac{y_i^4}{6(\nu\rho\sigma)^2} \right) \frac{1}{\sqrt{4\pi\rho\nu\sigma}^D} \exp\left(-\frac{|y|^2}{4\rho\nu\sigma}\right) dy d\sigma \right|. \end{aligned}$$

Iterating this partial integration procedure we get a geometric series and the upper bound

$$\begin{aligned} & \left| \rho L \int \int_{\{y_i \geq 0\} \cap B_{\sqrt{\rho\nu}}} \left(\frac{4y_i^2}{4\nu\rho\sigma} \right) G_\nu^\rho(\sigma, y) dy d\sigma \right| \\ &\leq \left| \rho L \int \frac{2}{3\sigma} \sqrt{\nu\rho} \frac{1}{\sqrt{4\pi\rho\nu\sigma}^D} \exp\left(-\frac{1}{4\sigma}\right) d\sigma \right|. \end{aligned} \quad (21)$$

Given $\nu > 0$ and as $D = 3$ the choice

$$\text{stepsize } \Delta = \nu \quad (22)$$

leads to a scheme with small upper bounds for local solution increments (related to the nonlinear terms), where

$$M_2 \sim \sqrt{\rho} \quad (23)$$

Note that this additional scaling is obtained by a) the Lipschitz continuity of the nonlinear terms in the probabilistic local time solution representation, and b) by the combination of this Lipschitz continuity with the possibility to rewrite the equation in terms of convolutions with first order derivatives of the Gaussian. Note that spatial transformations $y_i = rx_i$, $1 \leq i \leq D$ with

$$v_{i,j} = v_{i,j}^r \frac{dy_j}{dx_j} = v_{i,j}^r r \quad (24)$$

lead to analogous upper bounds with second moment constants of order

$$M_2^* \sim r, \quad (25)$$

and combinations of time and spatial transformation even to analogous upper bounds with second moment constants of order

$$M_2^* \sim \sqrt{\rho} r. \quad (26)$$

The schemes with step size ν become infinite of course as $\nu \downarrow 0$ such that an additional compactness argument has to be applied in order to use these estimates in the viscosity limit (similarly as this is done locally in this paper below).

For a local solution with data as in (10) you can check that $B_i^0, L^0, B_i, L_i, B_{i,j}, L_{i,j}$ are Lipschitz for $1 \leq i, j \leq D$. These considerations together with application of Young inequalities lead to the conclusion that for all $\tau_\rho \in [\tau_{\rho 0}, \tau_{\rho 0} + \Delta_0]$ we have

$$\max_{1 \leq i \leq D} |B_i^0 * G_{\nu,k}^\rho(\tau_\rho, \cdot)|_{H^2 \cap C^2} + |L^0 * G_{\nu,k}^\rho(\tau_\rho, \cdot)|_{H^2 \cap C^2} \leq \rho c_D C_m^2 C_K M_2 \quad (27)$$

or for all $1 \leq i \leq D$

$$|B_i * G_\nu^\rho(\tau_\rho, \cdot)|_{H^2 \cap C^2} + |L_i * G_{\nu,k}^\rho(\tau_\rho, \cdot)|_{H^2 \cap C^2} \leq \rho c_D C_m^2 C_K M_2, \quad (28)$$

where c_D is a finite constant which depends only on dimension D , and where $m = 2$ may be chosen. This is an upper bound of growth which can be offset by an autocontrol scheme. We may choose any ρ with

$$0 < \rho \leq \frac{1}{4c_D C_m^2 C_K} \quad (29)$$

For the Navier Stokes equation, i.e., for fixed $\nu > 0$ it is possible to analyze the damping term and compare it with upper bounds of the increment of the nonlinear terms over the time interval $[\tau_0, \tau_0 + \Delta]$. We cannot do this in the case of the Euler equation, of course. However the upper bound constructed above indicates that we can set up an autocontrol scheme which works without viscosity damping estimates. Here it is essential to have estimates which are independent of $\nu > 0$, i.e., estimates which work in the case of the Navier Stokes equation uniformly with respect to the viscosity $\nu > 0$ (we then may invoke Lipschitz continuity of the convoluted Leray projection term and Burgers

term, and apply some compactness argument in order to obtain a viscosity limit as in the main text). So let us consider fixed $\nu > 0$ in this remark.

For $\tau_{\rho 0} \geq 0$ and a time interval $[\tau_{\rho 0}, \tau_{\rho 0} + \Delta_0]$ for some $\Delta_0 \in (0, 1)$ we consider a comparison function u_i^{ρ, ν, τ_0} , $1 \leq i \leq D$ by a global time transformation with local time dilatation on a short time interval, i.e., we consider the transformation

$$(1 + \tau_\rho)u_i^{\rho, \nu, \tau_{\rho 0}}(s, y) = v_i^{\rho, \nu}(\tau_\rho, y), \quad (30)$$

where $s = \frac{\tau_\rho - \tau_{\rho 0}}{\sqrt{1 - (\tau_\rho - \tau_{\rho 0})^2}}$, $\tau_\rho - \tau_{\rho 0} \in [0, \Delta_0]$, $1 \leq i \leq D$

where $\Delta_0 \in [0, 1)$. Using the abbreviation $\Delta\tau_\rho = \tau_\rho - \tau_{\rho 0}$, we have

$$v_{i, \tau_\rho}^{\rho, \nu} = u_i^{\rho, \nu, \tau_{\rho 0}} + (1 + \tau_\rho)u_{i, s}^{\rho, \nu, \tau_{\rho 0}} \frac{ds}{d\tau_\rho}, \quad \frac{ds}{d\tau_\rho} = \frac{1}{\sqrt{1 - \Delta\tau_\rho^2}}. \quad (31)$$

We choose a time interval length $\Delta_0 \leq 0.5$ such that local time contraction, and, hence, a local time solution exists on the interval $[\tau_{\rho 0}, \tau_{\rho 0} + \Delta_0]$. The equation for $u_i^{\rho, \nu, \tau_{\rho 0}}$, $1 \leq i \leq D$ becomes

$$\begin{aligned} & u_{i, s}^{\rho, \nu, \tau_{\rho 0}} + \frac{\sqrt{1 - \Delta\tau_\rho^2}}{1 + \tau_\rho} \rho \nu \Delta u_i^{\rho, \nu, \tau_{\rho 0}} + \rho \frac{\sqrt{1 - \Delta\tau_\rho^2}}{1 + \tau_\rho} \sum_{j=1}^D \left(u_j^{\rho, \nu, \tau_{\rho 0}} \frac{\partial u_i^{\rho, \nu, \tau_{\rho 0}}}{\partial x_j} \right) \\ & - \rho \frac{\sqrt{1 - \Delta\tau_\rho^2}}{1 + \tau_\rho} \int_{\mathbb{R}^D} K_{D, i}(\cdot - y) \sum_{j, m=1}^D \left(\frac{\partial u_m^{\rho, \nu, \tau_{\rho 0}}}{\partial x_j} \frac{\partial u_j^{\rho, \nu, \tau_{\rho 0}}}{\partial x_m} \right) (\cdot, y) dy \\ & - \frac{\sqrt{1 - \Delta\tau_\rho^2}}{1 + \tau_\rho} u_i^{\rho, \nu, \tau_{\rho 0}} = 0, \\ & u^{\rho, \nu, \tau_{\rho 0}}(0, \cdot) = v^{\rho, \nu}(\tau_{\rho 0}, \cdot). \end{aligned} \quad (32)$$

This equation is defined on a dilated time interval $[0, \Delta_d]$, where the relation of Δ_d to Δ_0 follows from the time transformation. Now assume that an arbitrary large but fixed time horizon $T_\rho > 0$ is given. Note that this corresponds to an original time horizon $T = \rho T_\rho$ with ρ as in (29). Since ρ depends only on dimension, a spatial norm of the initial data (and the spatial kernel constant C_K) global results with arbitrary horizon T_ρ transfer to global results with respect to original time. Assume that for $m \geq 2$ and for some $0 \leq \tau_{\rho 0} < T$ we have a global upper bound

$$\max_{1 \leq i \leq D} \left| v_i^{\rho, \nu}(t_0, \cdot) \right|_{H^m \cap C^m} \leq C(1 + \tau_0) < \infty. \quad (33)$$

We may assume that $C \geq 3$. Then

$$\max_{1 \leq i \leq D} \left| u_i^{\rho, \nu, \tau_{\rho 0}}(0, \cdot) \right|_{H^m \cap C^m} = \frac{1}{1 + \tau_{\rho 0}} \max_{1 \leq i \leq D} \left| v_i^{\rho, \nu}(\tau_{\rho 0}, \cdot) \right|_{H^m \cap C^m} \leq C, \quad (34)$$

if ρ is chosen as in (29). Hence,

$$\max_{1 \leq i \leq D} \left| u_i^{\rho, \nu, \tau_{\rho 0}}(\Delta_d, \cdot) \right|_{H^m \cap C^m} \leq C. \quad (35)$$

This implies that

$$\begin{aligned} & \max_{1 \leq i \leq D} \left| v_i^{\rho, \nu}(\tau_{\rho 0} + \Delta_0, \cdot) \right|_{H^m \cap C^m} \\ & \leq \max_{1 \leq i \leq D} (1 + \tau_{\rho 0} + \Delta_0) \left| u_i^{\rho, \nu, \tau_{\rho 0}}(\Delta_d, \cdot) \right|_{H^m \cap C^m} \leq C(1 + \tau_{\rho 0} + \Delta_0). \end{aligned} \quad (36)$$

Iterating this procedure for all integers $k \geq 0$ with $k\Delta_0 \leq T_\rho$ we have

$$\begin{aligned} & \max_{1 \leq i \leq D} \left| v_i^{\rho, \nu}(k\Delta_0, \cdot) \right|_{H^m \cap C^m} \leq \max_{1 \leq i \leq D} (1 + k\Delta_0) \left| u_i^{\rho, \nu, \tau_{\rho 0}}(0, \cdot) \right|_{H^m \cap C^m} \\ & \leq C(1 + k\Delta_0). \end{aligned} \quad (37)$$

Then by a local time contraction we can interpolate and have for all $0 \leq \tau_\rho \leq T_\rho$

$$\begin{aligned} & \max_{1 \leq i \leq D} \left| v_i^{\rho, \nu}(\tau_\rho, \cdot) \right|_{H^m \cap C^m} \leq \max_{1 \leq i \leq D} (1 + \tau_\rho) \left| u_i^{\rho, \nu, \tau_{\rho 0}}(0, \cdot) \right|_{H^m \cap C^m} \\ & \leq 2C(1 + \tau_\rho). \end{aligned} \quad (38)$$

Remark 1.5. The argument below shows that singularities can be constructed even for data $\omega_i^f \in C_0^\infty$, $1 \leq i \leq D$, where C_0^∞ the space of smooth functions vanishing at infinity. However, $\omega_i^f \in H^2 \cap C^2$, $1 \leq i \leq D$ is essential, and we consider further regularity results and long-time kinks and singularities elsewhere. Here we are interested mainly in the connection to singular solution or kinks of the Navier Stokes equation with regular time dependent force terms.

The extended statement of Theorem 1.2 is considered elsewhere. Theorem 1.1 can be extended in the sense that there are solution branches with weak singularities of any integer order k in the sense that a solution function is only in $C^{k-1} \setminus C^k$. In order to have a succinct statement we introduce the concept of a 'spatial kink of order k ' of a classical solution.

Definition 1.6. We say that a classical solution branch ω_i , $1 \leq i \leq D$ of an incompressible Euler Cauchy problem with data ω_i^f , $1 \leq i \leq D$ with $\omega_i^f \in C^m$ for $m \geq k$ has spatial a kink of order $k \geq 1$, if there is a space-time point (τ, x) with $\tau > 0$ and $x \in \mathbb{R}^D$ such that

$$\omega_i \in C^{k-1} \setminus C^k \text{ at } (\tau, x) \quad (39)$$

for some integer $k \geq 1$.

Corollary 1.7. *Let $D = 3$. For any $k \geq 2$ and $s \geq 0$ there exist data $\omega_i^f \in H^m \cap C^m$, $1 \leq i \leq D$ with $m \geq k + 2$ and a vorticity solution ω_i , $1 \leq i \leq D$ of the three dimensional incompressible Euler equation Cauchy problem with data ω_i^f , $1 \leq i \leq D$ such that after some finite time the solution has a kink of order k .*

Remark 1.8. We treat the latter result as a Corollary because the proof method is the same, and only the choice of data for the time reversed Euler-type Cauchy problem has to be adapted.

Next we draw consequences for the Navier Stokes equation. A classical solution

$$\omega_i, 1 \leq i \leq D, \omega_i \in C^1([0, T], H^2 \cap C^2) \quad (40)$$

of the incompressible Euler equation (on the interval $[0, T)$ with out the point T and) with a blow-up of vorticity at time T satisfies the vorticity form of the Navier Stokes equation

$$\frac{\partial \omega}{\partial \tau} - \nu \Delta \omega + v \cdot \nabla \omega = \frac{1}{2} (\nabla v + \nabla v^T) \omega + F, \quad (41)$$

with force term $F = (F_1, F_2, F_3)^T$ (on the same time interval $[0, T)$) if

$$\text{for all } t \ F_i(t, \cdot) = -\nu \Delta \omega_i(t, \cdot) \in L^2 \cap C, 1 \leq i \leq 3. \quad (42)$$

The analysis below shows that F_i is also L^2 with respect to time on the time interval $[0, T]$, where $T > 0$ is the time where the vorticity of the Euler equation blows up (this is because we choose data of the time-reversed Euler equation in H^2). Similarly a regular classical solution on the time interval $[0, T)$

$$\omega_i, 1 \leq i \leq D, \omega_i(\tau, \cdot) \in H^{2+k} \cap C^{2+k} \quad (43)$$

of the incompressible Euler equation with a kink of order k at time $T > 0$ such that

$$\omega(T, \cdot) \in C^{k-1} \setminus C^k, \quad (44)$$

satisfies the vorticity form of the Navier Stokes equation (41), where

$$\text{for all } t \in [0, T) \ F_i = -\nu \Delta \omega_i \in H^k \cap C^k, 1 \leq i \leq 3. \quad (45)$$

We conclude

Theorem 1.9. *For the Cauchy problem for the Navier Stokes equation (41) for some time $T > 0$ time dependent force terms with*

$$F_i(\tau, \cdot) \in H^k \cap C^k, k \geq 0, \text{ for all } \tau \in [0, T) \quad (46)$$

and data $\omega_i(0, \cdot) \in H^{2+k} \cap C^{2+k}, 1 \leq i \leq 3, k \geq 0$ can be chosen such that a regular classical solution of the Navier Stokes equation on the time interval $[0, T]$ (and beyond) has a blow-up (case $k = 0$) or a kink of order k at time T .

2 Proof of Theorem 1.1 and Corollary 1.7

For the Gaussian fundamental solution G_ν of the equation

$$p_{,t} - \nu \Delta p = 0 \quad (47)$$

we consider on some time interval $[0, T]$ with time horizon $T > 0$ the iteration scheme $v_i^{\nu, -, k}$, $1 \leq i \leq D$, $k \geq 0$, where for $k \geq 1$

$$\begin{aligned} v_i^{\nu, -, k} &= v_i^f *_{sp} G_\nu + \sum_{j=1}^D \left(v_j^{\nu, -, k-1} \frac{\partial v_i^{\nu, -, k-1}}{\partial x_j} \right) * G_\nu \\ &- \left(\sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu, -, k-1}}{\partial x_j} \frac{\partial v_j^{\nu, -, k-1}}{\partial x_m} \right) (\cdot, y) dy \right) * G_\nu, \end{aligned}$$

and where this scheme is initialized by

$$v_i^{\nu, -, 0} := v_i^f(\cdot) *_{sp} G_\nu. \quad (48)$$

Here the velocity data v_i^f , $1 \leq i \leq D$ are determined by the vorticity data ω_i^f , $1 \leq i \leq D$ via the Biot-Savart law. Note that we start the iteration scheme with smooth data $v_i^f(\cdot) *_{sp} G_\nu$ which appear in the first approximating increment $\delta v_i^{\nu, 1, -} = v_i^{\nu, -, 1} - v_i^f *_{sp} G_\nu$. This makes it possible to estimate first derivative approximations $v_i^f *_{sp} G_{\nu, i}$ for the functional increments. In a former version of this paper we started the iteration scheme with the data, and this has the effect that the smoothing in the increment is felt only after the second iteration. Smoothing coefficients is often useful (sic Leray).

We note that the symbol $*_{sp}$ denotes convolution with respect to the spatial variables, and $*$ denotes convolution with respect to the time variable and the spatial variables. It is an advantage to start the scheme with the spatially convoluted data $v_i^f(\cdot) *_{sp} G_\nu$, since we choose the data v_i^f , $1 \leq i \leq D$ to be Lipschitz (and smooth in the complement of the origin and with decay at spatial infinity). A consequence is that the first derivatives have the representation

$$v_j^{\nu, -, 0} := v_i^f(\cdot) *_{sp} G_{\nu, j}, \quad (49)$$

where we shall use the fact that Lipschitz continuous data convoluted with the first order derivatives of the Gaussian have convenient upper bounds, which are even ν -independent. However, this is not decisive since we have this effect for $v_i^{\nu, -, k}$ for $k \geq 2$ anyway if we start the scheme with $v_i^f(\cdot)$ instead of $v_i^f(\cdot) *_{sp} G_\nu$ at $k = 0$. We choose an index $i_0 \in \{1, \dots, D\}$ and data $v_{i_0}^f$ and complement data v_j^f , $j \in \{1, \dots, D\} \setminus \{i_0\}$ such that corresponding vorticity component $\omega_{i_0}^f$ are singular at one point. Furthermore the complement data v_j^f , $j \in \{1, \dots, D\} \setminus \{i_0\}$ are constructed with the iteration scheme $v_i^{\nu, -, k}$, $1 \leq i \leq D$, $k \geq$ such that for all k

$$\text{for all } t \in (0, T] \quad \sum_{i=1}^D v_{i, i}^{\nu, -, k}(t, \cdot) = 0. \quad (50)$$

The basic idea to realise the additional constraint in (50) is as follows. Local

contraction leads to a local representation

$$\begin{aligned}
v_i^{\nu,-} &= v_i^f *_{sp} G_\nu + \sum_{j=1}^D \left(v_j^{\nu,-} \frac{\partial v_i^{\nu,-}}{\partial x_j} \right) * G_\nu \\
&- \left(\sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu,-}}{\partial x_j} \frac{\partial v_j^{\nu,-}}{\partial x_m} \right) (\cdot, y) dy \right) * G_\nu \quad (51) \\
&=: v_i^f *_{sp} G_\nu + \delta v_i^{\nu,-}.
\end{aligned}$$

We shall choose the data $v_{i_0}^f$ to be Lipschitz continuous such that the symmetry of the first order spatial derivative of the Gaussian (cf. (76) and the related estimate below) such that

$$v_{i_0, i_0}^{\nu,-}(t, \cdot) = v_{i_0}^f *_{sp} G_{\nu, i_0}(t, \cdot) + \delta v_{i_0, i_0}^{\nu,-}(t, \cdot) \quad (52)$$

has an appropriate upper bound for $t \in (0, T]$, where regularity observations concerning the increment $\delta v_{i_0, i_0}^{\nu,-}(t, \cdot)$ come into play. In the latter context we define for $k \geq 1$

$$\delta v_i^{\text{init}, \nu, -, k} = v_i^{\nu, -, k} - v_i^f *_{sp} G_\nu, \quad (53)$$

and

$$\delta v_i^{\nu, -, k} = v_i^{\nu, -, k} - v_i^{\nu, -, k-1}. \quad (54)$$

We show that for some time horizon $T > 0$ the local time solution function $v_i^{\nu,-}$, $1 \leq i \leq D$ of the Navier Stokes type extension of the time-reversed Euler equation has a representation of the form

$$v_i^{\nu,-} = v_i^f *_{sp} G_\nu + \delta v_i^{\text{init}, \nu, -, 2} + \sum_{k=3}^{\infty} \delta v_i^{\nu, -, k}, \quad (55)$$

which is spatially in $H^3 \cap C^3$ for evaluations at all time $t \in (0, T]$.

Remark 2.1. Note the choice $k = 2$ in (55) with respect to the initial increment (cf. (53)). However, as we start with the iteration scheme with the data $v_i^{\nu, -, 0} := v_i^f(\cdot) *_{sp} G_\nu$ this is not necessary and the alternative representation

$$v_i^{\nu,-} = v_i^f *_{sp} G_\nu + \delta v_i^{\text{init}, \nu, -, 1} + \sum_{k=3}^{\infty} \delta v_i^{\nu, -, k}, \quad (56)$$

is also sufficient for this improved scheme. If we start with data $v_i^f(\cdot)$ at $k = 0$ (as we did in a former version of this paper) instead, then the choice $k = 1$ is not sufficient as the first initial increment in (56) is then of lower regularity -at least for the choice of data considered below.

First note that the data v_i^f , $1 \leq i \leq D$ are ν -independent Hölder continuous data such that we have

$$\lim_{\nu \downarrow 0} v_i^f *_{sp} G_\nu = v_i^f, \quad 1 \leq i \leq D. \quad (57)$$

Moreover for Lipschitz continuous data v_i^f we have even a pointwise upper bound for first order derivatives $v_i^f *_{sp} G_{\nu, j}$ which can be used in the iteration scheme.

The convergence in (57) is in H^2 spatially (for the data chosen below). Hence we have Hölder continuous and pointwise convergence in case of dimension $D = 3$. However, our main interest concerns the regularity of the increment

$$\delta v_i^- = \lim_{\nu \downarrow 0} v_i^{\nu, -} - v_i^f. \quad (58)$$

For this purpose we need ν -independent estimates of convolutions

$$g_\nu * G_\nu, \quad g_\nu * G_{\nu, i} \quad (59)$$

with data

$$g_\nu \in S := \left\{ \delta v_i^{\text{init}, \nu, -, 2}, \delta v_i^{\nu, -, k}, k \geq 3 \right\}. \quad (60)$$

Note that $G_{\frac{\nu}{2}}$ is a probability density, so G_ν is essentially one, i.e., up to a constant. The hypothesis of an existing continuous viscosity limit

$$g = \lim_{\nu \downarrow 0} g_\nu \quad (61)$$

for $g_\nu \in S$ implies that the viscosity limit

$$\lim_{\nu \downarrow 0} g_\nu * G_\nu \text{ exists.} \quad (62)$$

Indeed for $g_\nu \in S$ for $z_i = \frac{y_i}{\sqrt{\nu}}$, $1 \leq i \leq D$ we have

$$\begin{aligned} g_\nu * G_\nu &= \int_0^t \int_{\mathbb{R}^D} g_\nu(s, x - y) \frac{1}{\sqrt{4\pi\nu s}^D} \exp\left(-\frac{|y|^2}{4\nu s}\right) dy ds = \\ &\int_0^t \int_{\mathbb{R}^D} g_\nu(s, x - \sqrt{\nu}z) \frac{1}{\sqrt{4\pi s}^D} \exp\left(-\frac{|z|^2}{4s}\right) dz ds \rightarrow \int_0^t g(s, x) ds \text{ as } \nu \downarrow 0. \end{aligned} \quad (63)$$

Similar considerations can be applied for multivariate spatial derivatives of order γ if $D_x^\gamma g_\nu$ is at least continuous and $D_x^\gamma g = \lim_{\nu \downarrow 0} D_x^\gamma g_\nu$ exists. In (48) for $k = 1$ we have $v_i^{\nu, -, k-1} = v_i^{\nu, -, 0} = v_i^f$, $1 \leq i \leq D$. Hence for $k = 1$ the data v_i^f in the convolutions $v_i^f * G_\nu$ and $v_i^f *_{sp} G_\nu$ are independent of ν and there are natural viscosity limits of these terms. For $k = 2$ convolutions of the form $D_x^\gamma (v_i^f *_{sp} G_\nu)$ appear in the recursive scheme (especially with $0 \leq |\gamma| \leq 1$) and as part of the function $v_i^{\nu, -, 1}$. We observe (according to the previous remarks) that these functions $D_x^\gamma (v_i^f *_{sp} G_\nu) = (D_x^\gamma v_i^f) *_{sp} G_\nu$, $0 \leq |\gamma| \leq 1$ have a ν -independent upper bound.

If we apply spatial derivatives to the scheme in (48), then it is natural to consider convolutions with first order spatial derivatives of the Gaussian $G_{\nu, i}$. We prove the existence of ν -independent upper bounds h_2, h_k , $k \geq 3$, where for some $T > 0$

$$\sup_{t \in [0, T]} |\delta v_i^{\text{init}, \nu, -, 2} * G_{\nu, i}(t, \cdot)|_{H^3 \cap C^3} \leq h_2, \quad (64)$$

and where for $k \geq 3$

$$\sup_{t \in [0, T]} |(\delta v_i^{\nu, -, k} * G_{\nu, i})(t, \cdot)|_{H^3 \cap C^3} \leq h_k. \quad (65)$$

Here we shall use spatial Lipschitz continuity of the convoluted data

$$\delta v_i^{\text{init}, \nu, -, 2}, \delta v_i^{\nu, -, k}, \quad 1 \leq i \leq D, \text{ for } k \geq 3.$$

Local time ν -independent contraction in spatial $|\cdot|_{H^3 \cap C^3}$ -norms of the higher order increments $\delta v_i^{\nu, -, k}$, $1 \leq i \leq D$ for $k \geq 3$ leads to a regular limit for the functional series in (55) on the time interval $(0, T]$ for some time horizon $T > 0$. Furthermore we shall observe that strong polynomial decay of order $2(D+1)$ is preserved for the increments, i.e., for $k \geq 3$ there is a $C > 0$ independent of ν such that for all $|x| \geq 1$ and $0 \leq |\gamma| \leq 3$

$$|(D_x^\gamma \delta v_i^{\nu, -, k} * G_\nu)(t, x)| \leq \frac{C}{1 + |x|^{2(D+1)}}, \quad (66)$$

and such that for all $|x| \geq 1$ and $0 \leq |\beta| \leq 2$

$$|(D_x^\beta \delta v_i^{\nu, -, k} * G_{\nu, i})(t, x)| \leq \frac{C}{1 + |x|^{2(D+1)}}. \quad (67)$$

We can then use compactness arguments or ν -independent local time contraction in order to verify the regularity of the limit for the functional series in (55).

In order to obtain ν -independent estimates for convolutions of type $D_x^\beta g_\nu * G_{\nu, i}$ for $g_\nu \in S$ we may use Lipschitz continuity of the convoluted data. For the first increment member $\delta v_i^{\text{init}, \nu, -, 2}$ in (55) this means that for multiindices $0 \leq |\beta| \leq 2$ and $\forall y, y' \in \mathbb{R}^D$ we have

$$|D_x^\beta \delta v_i^{\text{init}, \nu, -, 2}(t, y) - D_x^\beta \delta v_i^{\text{init}, \nu, -, 2}(t, y')| \leq c|y - y'| \quad (68)$$

for some finite constant $c > 0$. Lipschitz continuity turns out to hold for the higher order increments $D_x^\beta \delta v_i^{\nu, -, k}$, $k \geq 3$ for $0 \leq |\beta| \leq 2$ as well. We can have c independent of ν . Here we observe that the convolution with the Gaussian and the first order derivative of the Gaussian degenerates outside a ball $B_{\nu^{0.5-\epsilon}}(x)$ of radius $\nu^{0.5-\epsilon}$ for $0 < \epsilon < 0.5$ as ν becomes small. Indeed this holds for all spatial derivatives of the Gaussian as we have

$$\begin{aligned} & \left| D_x^\gamma G_\nu(t, x; s, y) \right|_{|x-y| \geq \nu^{0.5-\epsilon}} \\ & \leq \left| \frac{H_{\gamma, \nu(t-s)}(x-y)}{\sqrt{4\pi\nu(t-s)}^3} \exp\left(-\frac{|x-y|^2}{4\nu(t-s)}\right) \right|_{|x-y| \geq \nu^{0.5-\epsilon}} \downarrow 0 \text{ as } \nu \downarrow 0, \end{aligned} \quad (69)$$

where $H_{\gamma, \nu(t-s)}(x-y)$ is a Hermite-type polynomial of order $|\gamma|$ parameterized by $1/\nu(t-s)$. If the viscosity limit $\lim_{\nu \downarrow 0} g_\nu * D_x^\gamma G_\nu(t, x)$ exists (is finite for some x), then the 'mass' is concentrated in a ball of radius $B_{\nu^{0.5-\epsilon}}(x)$, i.e., assuming that the limits exist we have

$$\begin{aligned} \lim_{\nu \downarrow 0} g_\nu * D_x^\gamma G_\nu(t, x) &= \lim_{\nu \downarrow 0} \int_0^t \int_{\mathbb{R}^3} g_\nu(s, y) D_x^\gamma G_\nu(t, x; s, y) dy ds \\ &= \lim_{\nu \downarrow 0} \int_0^t \int_{B_{\nu^{0.5-\epsilon}}(x)} g_\nu(s, y) D_x^\gamma G_\nu(t, x; s, y) dy ds. \end{aligned} \quad (70)$$

Locally, we use simple Gaussian estimates of the form

$$\begin{aligned} & |D^\gamma G_\nu(t, x; s, y)| \\ & \leq \frac{|H_{\gamma, \nu(t-s)}(x-y)|}{\sqrt{4\pi}^3} \frac{1}{\nu^\delta (t-s)^\delta} (|x-y|^2)^{\delta - \frac{3}{2}} \left(\frac{|x-y|^2}{\nu(t-s)}\right)^{\frac{3}{2} - \delta} \exp\left(-\frac{|x-y|^2}{4\nu(t-s)}\right) \\ & \leq \frac{C_{|\gamma|}}{\nu^\delta t^\delta |x-y|^{3+|\gamma|-2\delta}} \end{aligned} \quad (71)$$

for $\delta \in (0, 1)$. We remark that for $0 \leq |\gamma| \leq 1$ the constant $C_{|\gamma|}$ in (71) can be chosen to be

$$C_0 = \sup_{z>0, \delta \in (0,1)} z^{\frac{3}{2}-\delta} \exp\left(-\frac{z^2}{4}\right) \quad (72)$$

and

$$C_1 = \sup_{z>0, \delta \in (0,1)} z^{\frac{5}{2}-\delta} \exp\left(-\frac{z^2}{4}\right). \quad (73)$$

Note that C_0, C_1 in (72), (73) are independent of ν . For first order spatial derivatives and regular data h with finite upper bound C_h we observe that for any $\epsilon > 0$

$$\begin{aligned} & \left| \int_{y \in \mathbb{R}^3 \setminus B_{\nu^{0.5-\epsilon}}^3(x)} h(y) \frac{(x-y)_j}{4\nu(t-s)\sqrt{4\pi\nu(t-s)}^3} \exp\left(-\frac{|x-y|^2}{4\nu(t-s)}\right) dy_1 dy_2 dy_3 \right| \\ & \leq \left| \frac{cC_h}{\sqrt{4\pi\nu(t-s)}} \exp\left(-\frac{1}{4\nu^{2\epsilon}(t-s)}\right) \right| \downarrow 0 \text{ as } \nu \downarrow 0, \end{aligned} \quad (74)$$

where we use Lipschitz continuity of the data, and where $c > 0$ is a positive constant. The observation in (74) tells us that any mass of a convolution with $G_{\nu,j}$ evaluated at (t, x) is concentrated in a ball $B_{\sqrt{\nu}}(x)$ of radius $\sqrt{\nu}$ around x as ν becomes small. In order to obtain ν -independent estimates we may use Lipschitz continuity for terms

$$h * G_{\nu,k} \text{ (which are equivalent for } \nu > 0) \quad (75)$$

with data $h = D_x^\beta g_\nu$ for $g_\nu \in S$ and $0 \leq |\beta| \leq 2$. Observe the symmetry

$$G_{\nu,i}(t, x; s, y) = -G_{\nu,i}(t, x^{i,-}; s, y^{i,-}), \quad (76)$$

where $x = (x_1, \dots, x_n)$, $x^{i,-} = (x_1^{i,-}, \dots, x_n^{i,-})$ and $x_j^{i,-} = x_j - 2\delta_{ij}x_j$ with the Kronecker δ_{ij} and where the symbol $y^{i,\cdot}$ is defined analogously. Then Lipschitz continuity (or Hölder continuity with exponent $\delta_1 = 1$) of the shifted convoluted function $h_x(\cdot) = h(x - \cdot)$ and transformation $z = x - y$ (with $r = \sqrt{z_1^2 + z_2^2 + z_3^2}$ for brevity) leads to

$$\begin{aligned} & \left| \int_{B_{\nu^{0.5}}(x)} h(y) \frac{2(x_i - y_i)}{4\nu s \sqrt{4\pi\nu s}^D} \exp\left(-\frac{|x-y|^2}{4\nu s}\right) dy \right| \\ & \leq \left| \int_{B_{\nu^{0.5}}, y_i \geq 0} \left| h_x(z) - h_x(z^{i,-}) \right| \frac{2z_j}{4\nu s \sqrt{\pi\nu s}^D} \exp\left(-\frac{|z|^2}{4\nu s}\right) dz \right| \\ & \leq \int_{B_{\nu^{0.5}}, y_i \geq 0} 4|r|^{\delta_1} \frac{C_1}{\nu^\delta t^\delta |r|^{4-2\delta}} r^2 dr \\ & 4 \frac{\tilde{C}}{\nu^\delta t^\delta} |r|^{\delta_1+2\delta-1} \Big|_0^{r=\nu^{0.5}} = 4 \frac{\tilde{C}}{t^\delta} |\nu|^{(0.5)(\delta_1+2\delta-1)-\delta} = 4 \frac{\tilde{C}}{t^\delta} \quad \forall \nu \geq 0 \text{ and } \delta_1 = 1. \end{aligned} \quad (77)$$

Here, $\delta \in (0.5, 1)$ and the latter equation holds even for $\nu = 0$ if the usual definition $0^0 = 1$ is used. This estimate can be used if the Leray data function, i.e., the Leray projection term applied to the data, is Lipschitz. Lipschitz Leray data function may correspond to a weak singularity of the vorticity, i.e. an oscillatory bounded vorticity which is not continuous. However, we shall observe

that even Hölder continuous data for a velocity component can be chosen and there is still a global solution branch of the reversed Euler equation with such data. This global solution branch is obtained by a viscosity limit of a scheme derived from a Navier Stokes type equation.

For well chosen initial data v_i^f , $1 \leq i \leq 3$ (for the viscosity extension of the time reversed problem) we observe that $\delta v_i^{\text{init},\nu,-,2}$ and $\delta v_i^{\nu,-,2}$ gain regularity independently of ν as k increases from 1 to 2 and observe that for all $t \geq 0$ the function

$$v_i^{\nu,-}(t, \cdot) - v_i^f *_{sp} G_\nu = \delta v_i^{\text{init},\nu,-,2}(t, \cdot) + \sum_{l=3}^{\infty} \delta v_i^{\nu,-,l}(t, \cdot) \quad (78)$$

have ν -independent spatial regular upper bounds with respect to the $H^3 \cap C^3$ -norm. Moreover we shall observe that the functional series in (78) can be differentiated twice with respect to the spatial variable member by member. Here, we may use strong polynomial decay at spatial infinity, which leads to stronger compactness arguments. The property of strong polynomial decay at spatial infinity is considered below (indeed, we shall observe that a spatial polynomial decay at spatial infinity of order $2(D+1)$ is preserved by the increments in (78) and for spatial derivatives of these increments up to order 3). Moreover we shall observe that

$$\lim_{\nu \downarrow 0} \nu \Delta v_i^{\nu,-} \equiv 0, \quad (79)$$

and conclude that $v_i^{\nu,-}$, $1 \leq i \leq D$ is a regular classical solution of the viscosity extension of the time-reversed incompressible Euler equation which has a regular viscosity limit which solves the time reversed incompressible Euler equation.

Remark 2.2. Note that the viscosity limit $\nu \downarrow 0$ for $D_x^\beta v_i^{\nu,-}$, $1 \leq i \leq D$, $0 \leq |\beta| \leq 2$ can be obtained by limits

$$D_x^\beta v_i^{\nu,-} - v_i^f *_{sp} D_x^\beta G_\nu = D_x^\beta \delta v_i^{\text{init},\nu,-,2} + \sum_{l=3}^{\infty} D_x^\beta \delta v_i^{\nu,-,l}, \quad (80)$$

(evaluated at $t > 0$), and where for $0 \leq |\beta| \leq 2$ we can use for $t > 0$

$$\begin{aligned} D_x^\beta \delta v_i^{\text{init},\nu,-,2} &= D_x^\beta v_i^{\nu,-,2} - v_i^f *_{sp} D_x^\beta G_\nu \\ &= \sum_{j=1}^D \left(D_x^\beta \left(v_j^{\nu,-,1} \frac{\partial v_i^{\nu,-,1}}{\partial x_j} \right) \right) * G_\nu \\ &\quad - \left(\sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(D_x^\beta \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu,-,1}}{\partial x_j} \frac{\partial v_j^{\nu,-,1}}{\partial x_m} \right) \right) (\cdot, y) dy \right) * G_\nu, \end{aligned} \quad (81)$$

and variations of this formula obtained by the convolution rule. Furthermore,

for $k \geq 3$ and $t > 0$ we may use

$$\begin{aligned}
D^\beta \delta v_i^{\nu, -, k} &= + \sum_{j=1}^D \left(D_x^\beta \left(v_j^{\nu, -, k-1} \frac{\partial v_i^{\nu, -, k-1}}{\partial x_j} \right) \right) * G_\nu \\
&- \left(\sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(D_x^\beta \left(\frac{\partial v_m^{\nu, -, k-1}}{\partial x_j} \frac{\partial v_j^{\nu, -, k-1}}{\partial x_m} \right) (\cdot, y) dy \right) \right) * G_\nu \\
&- \sum_{j=1}^D \left(D_x^\beta \left(v_j^{\nu, -, k-2} \frac{\partial v_i^{\nu, -, k-2}}{\partial x_j} \right) \right) * G_\nu \\
&+ \left(\sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(D_x^\beta \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu, -, k-2}}{\partial x_j} \frac{\partial v_j^{\nu, -, k-2}}{\partial x_m} \right) (\cdot, y) dy \right) \right) * G_\nu, \tag{82}
\end{aligned}$$

or variations by shifts of one spatial derivative to the Gaussian by the convolution rule. For $0 \leq |\beta| \leq 2$ we can consider the viscosity limit of a functional series of convolutions with the probability density G_ν . There is no degeneracy issue for this classical limit.

We summarize the six steps of the proof.

- i) As usual let $H^{2,1}$ be the Sobolev space where weak derivatives up to second order are in L^1 . We choose data v_i^f , $1 \leq i \leq D$, which determine the vorticity data $\omega_i^f = \omega_i^{\nu, -}(0, \cdot)$, $1 \leq i \leq D$ by the curl operation. Recall that the velocity data can be recovered from the vorticity data by the Biot-Savart law. For a positive number $\beta_0 \in (2, 2 + \alpha_0)$, and $\alpha_0 \in (0, \frac{1}{2})$ we consider velocity component data $v_i^f \in H^{2,1}$, $1 \leq i \leq 3$ where for one index $i_0 \in \{1, 2, 3\}$ we have $v_{i_0}^f(x) = g_{(0)}(r)$ for some univariate function $g_{(0)}$ which we are going to define next. Here the function $g_{(0)}$ is dependent on the radial component of spherical polar coordinates $r = \sqrt{x_1^2 + x_2^2 + x_3^2} \in \mathbb{R}_+$, where \mathbb{R}_+ denotes the set of nonnegative real numbers. This function $g_{(0)} : \mathbb{R}_+ \rightarrow \mathbb{R}$ satisfies

$$g_{(0)}(r) = \phi_1(r) r^{\beta_0} \sin\left(\frac{1}{r^{1+\alpha_0}}\right), \quad g_{(0)}(0) = 0, \tag{83}$$

where $\phi \in C_c^\infty$ is defined by

$$\phi_1(r) = \begin{cases} 1 & \text{if } r \leq 1, \\ \phi_1(r) = \alpha_*(r) & \text{if } 1 \leq r \leq 2, \\ 0 & \text{if } r \geq 2. \end{cases} \tag{84}$$

Furthermore, α_* is a smooth function with bounded derivatives for $1 \leq r \leq 2$. Such functions are well-known in the context of partitions of unity. As usual, C_c^∞ denotes the function space of smooth functions with compact support. In general $\beta_0 - 1$ will be chosen close to $1 + \alpha_0$. Note that the data function $v_{i_0}^f$ is only locally Lipschitz. The regularity gain of the local solution $v_i^{\nu, -}$, $1 \leq i \leq D$ is obtained for the solution increment $\delta v_i^{\nu, -} = v_i^{\nu, -} - v_i^f$.

Remark 2.3. In case $\beta_0 = 2 + \alpha_0$ we have Lipschitz continuous data. The singularities of the vorticities are then bounded and of oscillatory type. The following argument holds also in this case, but the result is weaker.

Remark 2.4. In order to prove the existence of kinks or weak singularities we may consider for $k \geq 2$ and $\beta_0^k \in (k+1, k+1+\alpha_0)$, $\alpha_0 \in (0, \frac{1}{2})$ velocity component data $v_i^f \in H^{2+k}$, $1 \leq i \leq 3$ where for one index $i_0 \in \{1, 2, 3\}$ we have $v_{i_0}^f(x) = g_{(k)}(r)$ and where $g_{(k)} : \mathbb{R}_+ \rightarrow \mathbb{R}$ is defined by

$$g_{(k)}(r) = \phi_1(r)r^{\beta_0^k} \sin\left(\frac{1}{r^{1+\alpha_0}}\right). \quad (85)$$

For $j \in \{1, 2, 3\} \setminus \{i_0\}$ we may choose velocity component data such that

$$\sum_{j=1}^D v_{j,j}^f = 0, \quad \sum_{j=1}^D \omega_{j,j}^f = 0 \quad (86)$$

is satisfied in a distributive sense (and in a classical sense outside the origin). Indeed choosing similar functions v_j^f for $j \neq i_0$, we $\sum_{j=1}^D v_{j,j}^f = 0$ in H^1 sense. With such a choice we may also choose the functions v_j^f , $1 \leq j \leq D$, $j \neq i_0$ such that for all $t > 0$

$$\sum_{k=1}^D v_k^f *_{sp} G_{\nu,k}(t, \cdot) = 0, \quad (87)$$

where the relation holds in the H^1 -sense at $t = 0$. The local time solution scheme below then implies that the local solution of a Navier Stokes type equation (related to the time reversed Euler equation) satisfies incompressibility for $t > 0$ (as does the viscosity limit). Note that we cannot impose a incompressible condition at $t = 0$ in a classical sense, as we have a singular point at time $t = 0$. We shall show that for data v_j^f , $j \neq i_0$ chosen below the local solution $v_i^{\nu,-}$, $1 \leq i \leq D$ and the associated time-reversed Euler equation satisfy the incompressibility equation, i.e. for some time horizon $T > 0$ we have

$$\forall t \in (0, T] : \sum_{i=1}^D v_{i,i}^{\nu,-}(t, \cdot) = 0. \quad (88)$$

Note that for $k = 0$ and $\beta_0 = \beta_0^0$ close to $1 + \alpha_0$ we have $v_i^f \in H^{2,1}$ $\omega_i^f \in H^{1,1}$ for all $1 \leq i \leq D = 3$ as the singularity of $v_{i_0}^f = v_{i_0}^f$ (resp. $\omega_{i_0}^f = \omega_1^f$) at $r = 0$ (polar coordinates) is of order $|\beta_0^0 - 4 - 2\alpha_0| < 2.5$ for β_0^0 close to $2 + \alpha_0$ as $\alpha_0 \in (0, 0.5)$. Similarly for any $l \geq 0$ we have choices $\beta_0^k \in (k+1, k+1+\alpha_0]$, $\alpha_0 \in (0, \frac{1}{2})$ such that

$$g_{(k)}(\cdot) \in C^{l-1} \setminus C^l. \quad (89)$$

We just have to choose $l = \min\{m \geq 0 | \beta_0^k - m(2 + \alpha_0) < 0\}$. For this choice of l we have $g_{(k)}(\cdot) \in H^{l,1}$.

- ii) For given $\nu > 0$, $1 \leq i \leq 3$ and for all $0 \leq |\gamma| \leq 1$, $x \in \mathbb{R}^3 \setminus \{0\}$, $r = \sqrt{x_1^2 + x_2^2 + x_3^2} \in \mathbb{R} \setminus \{0\}$ we have for the first iteration ($k = 1$)

$$|D_x^\gamma \delta v_i^{\text{init}, \nu, -, 1}(t, x)| = |D_x^\gamma \delta v_i^{\nu, -, 1}(t, x)| \leq Cr^{2\beta_0 - 1 - |\gamma|}, \quad (90)$$

and for the second iteration ($k = 2$) and all $\delta \in (0.5, 1)$

$$|D_x^\gamma \delta v_i^{\text{init}, \nu, -, 2}(t, x)| \leq Cr^{2\beta_0 - 1 - |\gamma|}. \quad (91)$$

For the higher order increments and for $k \geq 3$ we also have for all $\delta \in (0.5, 1)$

$$|D_x^\gamma \delta v_i^{\nu, -, k}(t, x)| \leq Cr^{2(\beta_0 - 1) + 1 - |\gamma|}. \quad (92)$$

Indeed these increments gain regularity at each iteration step, but we shall not consider this here.

Inheritance of polynomial decay (cf. item iii)) then implies for $1 \leq i \leq 3$ and for all $k \geq 3$

$$\delta v_i^{\nu, -, k}(t, \cdot) \in H^3 \cap C^3, \quad (93)$$

and

$$\delta v_i^{\text{init}, \nu, -, 2}(t, \cdot) \in H^3 \cap C^3. \quad (94)$$

Note that at this stage we do not consider dependence on ν . The independence of ν requires an additional argument and is considered in item v).

- iii) For a short time horizon $T > 0$ for all $\nu > 0$ polynomial decay of order $2(D + 1)$ at spatial infinity is inherited by the increments

$$D_x^\gamma \delta v_i^{\text{init}, \nu, -, 2}, D_x^\gamma \delta v_i^{\nu, -, k}, \quad k \geq 3, \quad 0 \leq |\gamma| \leq 3.$$

More precisely for $1 \leq i \leq 3$, and $k \geq 3$ there exists a finite constant $c > 0$ such that for all $t \in [0, T]$ and $|x| \geq 1$

$$\forall k \geq 1 \quad \forall 0 \leq |\gamma| \leq 3 \quad \left| D_x^\gamma \delta v_i^{\text{init}, \nu, -, 2}(t, x) \right| \leq \frac{c}{1 + |x|^{2(D+1)}}, \quad (95)$$

and

$$\forall k \geq 1 \quad \forall 0 \leq |\gamma| \leq 3 \quad \left| D_x^\gamma \delta v_i^{\nu, -, k}(t, x) \right| \leq \frac{c}{1 + |x|^{2(D+1)}}. \quad (96)$$

For small $T > 0$ it is observed that the constant $c > 0$ can be chosen independently of the iteration index k . Furthermore, for $0 \leq |\gamma| \leq 4$

$$\left| v_i^f *_{sp} D_x^\gamma G_\nu(t, \cdot) \right| \leq \frac{c}{1 + |x|^{2(D+1)}} \quad t > 0 \text{ and } c \text{ depends on } t > 0, \quad (97)$$

such that a corresponding statement holds for the value functions $v_i^{\nu, -, k}$, $1 \leq i \leq 3$ and $k \geq 2$.

- iv) For the parameter choices of β_0, α_0 with $\beta_0 = 1.5 - \epsilon$ for small $\epsilon > 0$ of item i), for $1 \leq i \leq 3$, $\nu > 0$ and for $k \geq 3$ and some time $T > 0$

$$\left| \delta v_i^{\nu, -, k+1}(t, \cdot) \right|_{H^3 \cap C^3} \leq \frac{1}{2} \left| \delta v_i^{\nu, -, k}(t, \cdot) \right|_{H^3 \cap C^3}. \quad (98)$$

Hence, for all $t \in [0, T]$

$$v_i^{\nu, -}(t, \cdot) - v_i^f *_{sp} G_\nu(t, \cdot) = \delta v_i^{\text{init}, \nu, -, 2} + \sum_{l=3}^{\infty} \delta v_i^{\nu, -, l} \in H^3 \cap C^3. \quad (99)$$

- v) There exists a $T > 0$ independent of $\nu > 0$ and a finite constant $C > 0$ independent of $\nu > 0$ such that

$$\sup_{t \in (0, T]} |v_i^{\nu, -}(t, \cdot) - v_i^f *_{sp} G_\nu(t, \cdot)|_{H^3 \cap C^3} \leq C. \quad (100)$$

Furthermore, for such $T > 0$ independent of $\nu > 0$ such that

$$\text{for all } t \in (0, T]: \sum_{i=1}^D v_{i,i}^{\nu, -}(t, \cdot) = 0. \quad (101)$$

We note that the relation in (101) hold for all $\nu > 0$. The trick here is to use the smooth coefficients in the iteration scheme in order to estimate the functional increments (cf. below ad v)).

- vi) Choose a time horizon $T > 0$ as in the previous step such that contraction holds for the higher order increments $\delta v_i^{\nu, -, k}$ with $k \geq 3$ as in (98). For an upper bound $C > 0$ independent of $\nu > 0$ we have

$$\begin{aligned} & |\delta v_i^{\text{init}, \nu, -, 2}(t, \cdot) + \sum_{l=3}^{\infty} \delta v_i^{\nu, -, l}(t, \cdot)|_{H^3 \cap C^3} \\ &= |v_i^{\nu, -}(t, \cdot) - v_i^f *_{sp} G_\nu(t, \cdot)|_{H^3 \cap C^3} \leq C. \end{aligned} \quad (102)$$

For all $\nu > 0$ and for some $T > 0$ (independent of ν) the function

$$v_i^{\nu, -}(t, \cdot) = v_i^f *_{sp} G_\nu(t, \cdot) + \delta v_i^{\text{init}, \nu, -, 2} + \sum_{l=3}^{\infty} \delta v_i^{\nu, -, l} \quad (103)$$

satisfies the Navier stokes equation on the time interval $[0, T]$ such that

$$\begin{aligned} \nu \Delta v_i^{\nu, -} &= \frac{\partial v_i^{\nu, -}}{\partial t} - \sum_{j=1}^D v_j^{\nu, -} \frac{\partial v_i^{\nu, -}}{\partial x_j} \\ &+ \sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu, -}}{\partial x_j} \frac{\partial v_j^{\nu, -}}{\partial x_m} \right) (t, y) dy, \end{aligned} \quad (104)$$

where the right side of (104) is the time-reversed Euler equation operator applied to $v_i^{\nu, -}$. We choose a sequence $(\nu_p)_{p \geq 1}$ converging to zero and use probability density representations (convolutions with G_ν itself) of increments $D_x^\beta \delta v_i^{\text{init}, \nu_p, -, 2}$ and $D_x^\beta \delta v_i^{\nu, -, l}$, $l \geq 3$ for $0 \leq |\beta| \leq 2$ in

$$v_i^{-, \nu_p} := v_i^f *_{sp} G_{\nu_p} + \delta v_i^{-, \nu_p} \quad (105)$$

in order to show that the viscosity limit v_i^- , $1 \leq i \leq D$ is a classical solution of the time-reversed Euler equation.exists . Here

$$\delta v_i^- := \lim_{\nu_p \downarrow 0} \delta v_i^{\text{init}, \nu_p, -, 2} + \sum_{l=3}^{\infty} \delta v_i^{\nu, -, l} \in C^1((0, T], H^3 \cap C^3). \quad (106)$$

Furthermore the function $v_i^- = v_i^f + \delta v_i^-$, $1 \leq i \leq D$ solves the time-reversed incompressible Euler equation and satisfies the incompressibility condition in the time interval $(0, T]$, i.e.,

$$\text{for all } t \in (0, T] \sum_{i=1}^D v_{i,i}^-(t, \cdot) = 0. \quad (107)$$

We conclude that the original Euler equation develops in opposite time direction from data at time $\tau = 0$ (corresponding to time $s = T$ of the reversed-time Euler equation) a weak singularity at time $T > 0$ (corresponding to data at time $t = 0$ the time-reversed Euler equation).

ad i) first we note that multiindices are denoted by $\alpha = (\alpha_1, \dots, \alpha_D), \beta = (\beta_1, \dots, \beta_D), \dots$ in this paper, and α_0, β_0 just refer to positive real numbers. For the derivative of the data $v_{i_0}^f = v_1^f$ and $k = 0$ we compute for $r \neq 0$ and $r \leq 1$

$$g'(r) = \frac{d}{dr} r^{\beta_0} \sin\left(\frac{1}{r^{1+\alpha_0}}\right) = \beta_0 r^{\beta_0-1} \sin\left(\frac{1}{r^{1+\alpha_0}}\right) - (1+\alpha_0) r^{\beta_0-2-\alpha_0} \cos\left(\frac{1}{r^{1+\alpha_0}}\right). \quad (108)$$

The derivative g' of the function g at $r = 0$ is strongly singular for $\beta_0 \in (2, 2+\alpha_0)$ and $\alpha_0 \in (0, \frac{1}{2})$. Note that it is 'oscillatory' singular bounded for $\beta_0 = \alpha_0 \in (0, \frac{1}{2})$. Note that for data $v_{i_0}^f(x_1, x_2, x_3) = g(r)$ we have (for $r \neq 0$)

$$v_{i_0,j}^f = g'(r) \frac{\partial r}{\partial x_j} = g'(r) \frac{x_j}{r}. \quad (109)$$

In polar coordinates $(r, \theta, \phi) \in [0, \infty) \times [0, \pi] \times [0, 2\pi]$ with

$$x_1 = r \sin(\theta) \cos(\phi), \quad x_2 = r \sin(\theta) \sin(\phi), \quad x_3 = r \cos(\theta), \quad (110)$$

(where for $r \neq 0$ and $x_1 \neq 0$ we have $r = \sqrt{x_1^2 + x_2^2 + x_3^2}$, $\theta = \arccos\left(\frac{x_3}{r}\right)$, $\phi = \arctan\left(\frac{x_2}{x_1}\right)$) we get

$$\begin{aligned} v_{i_0,1}^f &= g'(r) \frac{x_1}{r} = g'(r) \sin(\theta) \cos(\phi), \\ v_{i_0,2}^f &= g'(r) \frac{x_2}{r} = g'(r) \sin(\theta) \sin(\phi), \\ v_{i_0,3}^f &= g'(r) \frac{x_3}{r} = g'(r) \cos(\theta), \end{aligned} \quad (111)$$

such that we have

$$v_{i_0}^f \in H^{1,1} \text{ obviously.} \quad (112)$$

The second derivative of g is

$$\begin{aligned} g''(r) &= \frac{d^2}{dr^2} r^{\beta_0} \sin\left(\frac{1}{r^{1+\alpha_0}}\right) \\ &= \frac{d}{dr} \left(\beta_0 r^{\beta_0-1} \sin\left(\frac{1}{r^{1+\alpha_0}}\right) - (1+\alpha_0) r^{\beta_0-2-\alpha_0} \cos\left(\frac{1}{r^{1+\alpha_0}}\right) \right) \\ &= \beta_0(\beta_0-1) r^{\beta_0-2} \sin\left(\frac{1}{r^{1+\alpha_0}}\right) - (1+\alpha_0) \beta_0 r^{\beta_0-3-\alpha_0} \cos\left(\frac{1}{r^{1+\alpha_0}}\right) \\ &\quad + (1+\alpha_0)(2+\alpha_0-\beta_0) r^{\beta_0-2-\alpha_0} \cos\left(\frac{1}{r^{1+\alpha_0}}\right) \\ &\quad - (1+\alpha_0)^2 r^{\beta_0-4-2\alpha_0} \sin\left(\frac{1}{r^{1+\alpha_0}}\right). \end{aligned} \quad (113)$$

We have $v_{i_0}^f \in H^{2,1}$, since

$$\beta_0 - 4 - 2\alpha_0 > -3. \quad (114)$$

Note that

$$v_{i_0}^f(0, \cdot) \in C^\delta(\mathbb{R}^3), \quad (115)$$

for Hölder constants of order $\delta \in (0, \frac{1}{2})$. Similarly we have

$$\omega_{i_0}^f = \omega_1^f = \omega_1(0, \cdot) \in H^{1,1}. \quad (116)$$

We do not specify the data v_j^f for $j \neq i_0$ at this time. The choice is not arbitrary since we have to ensure the incompressibility condition at times $t \in (0, T]$ for some time $T > 0$. At time $t = 0$ the incompressibility condition breaks down (at least in a classical sense). It is quite remarkable that the Lipschitz continuity of the data $v_{i_0}^f$ makes it possible to choose data v_j^f , $j \neq 0$ which are smooth and bounded (cf. below).

ad ii) For $k = 1$ (48) becomes

$$\begin{aligned} v_i^{\nu, -, 1} &= v_i^f(\cdot) *_{sp} G_\nu + \sum_{j=1}^D \left(\left(v_j^f *_{sp} G_\nu \right) \left(v_i^f *_{sp} G_{\nu, j} \right) \right) * G_\nu \\ &- \sum_{j, m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j, m=1}^D \left(\left(v_m^f *_{sp} G_{\nu, j} \right) \left(v_j^f *_{sp} G_{\nu, m} \right) \right) (t, y) dy * G_\nu \\ &=: v_i^f(\cdot) *_{sp} G_\nu + B_0 * G_\nu - L_0 * G_\nu, \end{aligned} \quad (117)$$

where B_0 and L_0 denote abbreviations of the lowest order approximations of the Burgers term and the Leray projection term. Recall that at this stage we consider a given $\nu > 0$, and do not consider dependence of ν . Nevertheless the estimates are of a form such that an additional argument in item v) leadsto ν -independence of upper bounds. For $t \geq 0$ we have

$$\left| \left(v_j^f *_{sp} G_\nu \right) (t, \cdot) \right| \leq cr^{\beta_0} \quad (118)$$

and

$$\left| \left(v_i^f *_{sp} G_{\nu, j} \right) (t, \cdot) \right| \leq cr^{\beta_0 - 1}, \quad (119)$$

for some finite constant $c > 0$ which is independent of $\nu > 0$.

Consequently, we have

$$\left| \left(v_j^f *_{sp} G_\nu \right) \left(v_i^f *_{sp} G_{\nu, j} \right) (t, \cdot) \right| \leq cr^{2\beta_0 - 1} \quad (120)$$

and as $\frac{\partial}{\partial x_i} K_{D, i}(\cdot - y) \sim \frac{1}{r^2}$ for $D = 3$ we have

$$\begin{aligned} &\left| \int_{\mathbb{R}^D} K_{D, i}(\cdot - y) \sum_{j, m=1}^D \left(\left(v_j^f *_{sp} G_{\nu, m} \right) \left(v_m^f *_{sp} G_{\nu, j} \right) \right) (t, y) dy \right| \\ &\leq cr^{2\beta_0 - 2 + 1} = cr^{2\beta_0 - 1}. \end{aligned} \quad (121)$$

Using Gaussian estimates above for multiindices γ with $0 \leq |\gamma| \leq 1$ for the convoluted Burgers term B^0 we get

$$\left| B^0 * D_x^\gamma G_\nu(t, \cdot) \right| \leq cr^{2\beta_0 - 1 - |\gamma|}, \quad (122)$$

and as $\frac{\partial}{\partial x_i} K_{D, i}(\cdot - y) \sim \frac{1}{r^2}$ for $D = 3$ for the convoluted Leray projection term L^0 we have

$$\left| L^0 * D_x^\gamma G_\nu(t, \cdot) \right| \leq cr^{2\beta_0 - 1 - |\gamma|} \quad (123)$$

Hence the statement in (90) follows. Next we observe that

$$\forall t > 0 \quad v_i^{\nu, -1}(t, \cdot) \in C^2 \cap H^2. \quad (124)$$

First we consider the Leray projection term. For the second order spatial derivatives with indices p, q we have

$$\begin{aligned} L_{i,p,q}^* &:= L_{i,p}^0 * G_{\nu,q} = - \sum_{j,m=1}^D \int_{\mathbb{R}^D} K_{D,i,p}(\cdot - y) \times \\ &\times \sum_{j,m=1}^D \left((v_m^f *_{sp} G_{\nu,j}) \left(v_j^f *_{sp} G_{\nu,m} \right) \right) (t, y) dy * G_{\nu,q} \end{aligned} \quad (125)$$

The singularity order of $K_{D,p,q}$ at $r = 0$ is r^{-n} such that for all x in a finite ball around zero (recall $r = |x| = \sqrt{\sum_{j=1}^D x_j^2}$) we have for all $t > 0$ (fixed)

$$\begin{aligned} &\left| \sum_{j,m=1}^D \int_{\mathbb{R}^D} K_{D,i,p}(x - y) \sum_{j,m=1}^D \left((v_m^f *_{sp} G_{\nu,j}) \left(v_j^f *_{sp} G_{\nu,m} \right) \right) (t, y) dy \right| \\ &\leq cr^{2\beta_0-2} \end{aligned} \quad (126)$$

where c is independent of $\nu > 0$ (and of t). Hence, for all x in a finite ball around the origin we have

$$\left| L_{i,p,q}^0(t, x) \right| \leq cr^{2\beta_0-3}. \quad (127)$$

The latter estimate holds a fortiori for lower order derivatives of L_i . It follows that L_i and spatial derivatives of L_i up to second order are Lipschitz, especially continuous. Here, note that we easily conclude that for all $t > 0$

$$L_{i,p,q}^*(t, \cdot) \in L^2. \quad (128)$$

This holds a fortiori for the lower order spatial derivatives such that we conclude that for all $t > 0$

$$L_i^*(t, \cdot) \in H^2 \cap C^2. \quad (129)$$

Next consider the Burgers term

$$B_i^* = \sum_{j=1}^D \left((v_j^f *_{sp} G_{\nu}) \left(v_i^f *_{sp} G_{\nu,j} \right) \right) * G_{\nu}. \quad (130)$$

We have

$$\begin{aligned} B_{i,p,q}^* &= \sum_{j=1}^D \left((v_j^f *_{sp} G_{\nu,p}) \left(v_i^f *_{sp} G_{\nu,j} \right) \right) * G_{\nu,q} \\ &+ \sum_{j=1}^D \left((v_j^f *_{sp} G_{\nu}) \left(v_i^f *_{sp} G_{\nu,j,p} \right) \right) * G_{\nu,q} \end{aligned} \quad (131)$$

Since we assume $\nu > 0$, we have to show that for all $1 \leq i, j \leq D$ the functions

$$\left| B_{i,p,q}^0(t, x) \right| \leq cr^{2\beta_0-3}, \quad (132)$$

and we can argue similarly as in the case of the Leray projection term in order to conclude that for all $t > 0$

$$B_i^*(t, \cdot) \in H^2 \cap C^2. \quad (133)$$

Moreover, as $\nu > 0$ the nonlinear terms $B_i^*(t, \cdot)$ and $L_i^*(t, \cdot)$ and their derivatives up to second order are Lipschitz continuous (note the difference to the initial data which are only locally Lipschitz continuous). It follows that after the first iteration the nonlinear terms $B_i^1(t, \cdot)$ and $L_i^1(t, \cdot)$ and their derivatives up to second order are Lipschitz continuous. For $k = 2$ (48) becomes

$$\begin{aligned} v_i^{\nu, -, 2} &= v_i^f(\cdot) *_{sp} G_\nu + \sum_{j=1}^D v_j^{\nu, -, 1} \frac{\partial v_i^{\nu, -, 1}}{\partial x_j} * G_\nu \\ &- \sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{l,m=1}^D \left(\frac{\partial v_m^{\nu, -, 1}}{\partial x_j} \frac{\partial v_j^{\nu, -, 1}}{\partial x_m} \right) (t, y) dy * G_\nu \quad (134) \\ &=: v_i^f(\cdot) *_{sp} G_\nu + B_1 * G_\nu - L_1 * G_\nu, \end{aligned}$$

where B_1 and L_1 denote abbreviations of the next order of approximation of the Burgers term and the Leray projection term. As we have $(v_i^f(\cdot) *_{sp} G_\nu)(t, \cdot) \in H^3 \cap C^3$ for $t > 0$. and B_1 and L_1 are $C^2 \cap H^2$, and Lipschitz continuity of B_1 and L_1 and their derivatives up to second order we conclude that

$$\forall t > 0 \quad (B_1 * G_\nu)(t, \cdot) \in H^3 \cap C^3, \quad (L_1 * G_\nu)(t, \cdot) \in H^3 \cap C^3. \quad (135)$$

ad iii) For $k \geq 1$ choose a number m is such that for $t \in [0, T]$

$$\forall 0 \leq |\gamma| \leq m \quad D_x^\gamma \delta v_i^{init, \nu, -, k}(t, \cdot) \text{ is continuous and bounded.} \quad (136)$$

We have observed that for $k = 2$ we can choose $m = 3$ such that the upper bound is independent of $\nu > 0$. For $0 \leq |\gamma| \leq m$ and for $|\beta| + 1 = |\gamma|$, $\beta_j + 1 = \gamma_j$ (if $|\gamma| \geq 1$) we consider a representation of $D_x^\gamma v_i^{\nu, -, k}$, $1 \leq i \leq D$, $k \geq 1$ of the form

$$\begin{aligned} D_x^\gamma v_i^{\nu, -, k} &= v_i^f *_{sp} D_x^\gamma G_\nu - D_x^\beta \left(\sum_{j=1}^D v_j^{\nu, -, k-1} \frac{\partial v_i^{\nu, -, k-1}}{\partial x_j} \right) * G_{\nu, j} \\ &- D_x^\beta \left(\sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{l,m=1}^D \left(\frac{\partial v_m^{\nu, -, k-1}}{\partial x_l} \frac{\partial v_l^{\nu, -, k-1}}{\partial x_m} \right) (t, y) dy \right) * G_{\nu, j}. \end{aligned} \quad (137)$$

Here, recall G_ν is the fundamental solution of the heat equation $p_{,t} - \nu \Delta p = 0$, $*$ denotes the convolution, $*_{sp}$ denotes the spatial convolution, and K_D denotes the fundamental solution of the Laplacian equation for dimension $D \geq 3$. In the following the constant $c > 0$ is generic. Note that for $1 \leq i \leq D$ the initial data v_i^f have polynomial decay of any order at spatial infinity, i.e. we have for $|x| \geq 1$

$$|v_i^f(x)| \leq \frac{c}{1 + |x|^{2(D+1)+D+m}} \quad (138)$$

for some finite constant $c > 0$ and $t \geq 0$. Hence, for multiindices $0 \leq |\gamma| \leq m$ and $t > 0$ we have for some finite constant $c > 0$ and for $|x| \geq 1$

$$|v_i^f *_{sp} D_x^\gamma G_\nu(t, x)| \leq \frac{c}{1 + |x|^{2(D+1)}}. \quad (139)$$

Assuming inductively

$$\forall l \leq k-1 \quad \forall 0 \leq |\gamma| \leq m \quad \left| D_x^\gamma v_i^{\nu, -, l}(\cdot) \right| \leq \frac{c}{1 + |x|^{2(D+1)}} \quad (140)$$

we have or some finite constant $c > 0$ and for for $|x| \geq 1$

$$|D_x^\beta B^{k-1}| := \left| \sum_{j=1}^D D_x^\beta \left(v_j^{\nu,-,k-1} \frac{\partial v_i^{\nu,-,k-1}}{\partial x_j} \right) (t, \cdot) \right| \leq \frac{c}{1 + |x|^{4(D+1)}}, \quad (141)$$

where $|\beta| + 1 + |\gamma|$ is defined as above, and

$$|D_x^\beta L^{k-1}| \leq \frac{c}{1 + |x|^{4D+3}}, \quad (142)$$

where

$$D_x^\beta L^{k-1} \equiv \sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(D_x^\beta \left(\frac{\partial v_m^{\nu,-,k-1}}{\partial x_j} \frac{\partial v_j^{\nu,-,k-1}}{\partial x_m} \right) \right) (t, y) dy. \quad (143)$$

Convolutions with G_ν or $G_{\nu,i}$ weaken this polynomial decay by order D at most such that we (generously) get

$$|D_x^\beta B^{k-1} * G_{\nu,j}| \leq \frac{c}{1 + |x|^{3(D+1)}} \quad (144)$$

and

$$|D_x^\beta L^{k-1} * G_{\nu,j}| \leq \frac{c}{1 + |x|^{2(D+1)}}. \quad (145)$$

Hence using the representation (48) and (139),(144),(145) we get

$$\forall l \leq k \quad \forall 0 \leq |\gamma| \leq m \quad |D_x^\gamma v_i^{\nu,-,l}(\cdot)| \leq \frac{c}{1 + |x|^{2(D+1)}} \quad (146)$$

and by (140) the same holds for the increments $D_x^\gamma \delta v_i^{\nu,-,k}(\cdot)$ a fortiori.

Note that similar conclusions can be made using a vorticity iteration scheme $(\omega_i^{\nu,-,k})_{k \geq 0, 1 \leq i \leq D}$, where for $k = 0$

$$\omega_i^{\nu,-,0} = f_i, \text{ for } 1 \leq i \leq D, \quad (147)$$

and where for $k \geq 1$ the function $\omega_i^{\nu,-,k}$, $1 \leq i \leq D$ is determined as the time-local solution of the Cauchy problem

$$\begin{cases} \frac{\partial \omega_i^{\nu,-,k}}{\partial s} - \nu \Delta \omega_i^{\nu,-,k} - \sum_{j=1}^3 v_j^{\nu,-,k-1} \frac{\partial \omega_i^{\nu,-,k-1}}{\partial x_j} \\ = - \sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-,k-1}}{\partial x_j} + \frac{\partial v_j^{\nu,-,k-1}}{\partial x_i} \right) \omega_j^{\nu,-,k-1}, \\ \omega_i^{\nu,-,k}(0, \cdot) = \omega_i^f, \end{cases} \quad (148)$$

on a domain $[0, T] \times \mathbb{R}^D$ for some time horizon $T > 0$. Recall that in the limit $k \uparrow \infty$ (if existent) we have

$$v^{\nu,-}(t, x) = \int_{\mathbb{R}^3} (K_3(y) \omega^{\nu,-}(t, x - y)) dy, \quad (149)$$

by the Bio-Savart law .

ad iv) For $k \geq 3$ we have $\delta v_i^{\nu,-,k}(t, \cdot) \in H^3 \cap C^3$ corresponding to $\delta \omega_i^{\nu,-,k}(t, \cdot) \in H^2 \cap C^2$. For convenience of the reader we do some explicit very simple computations concerning the recursive relation of increments. For the scheme $\omega_i^{\nu,-,k}(s, \cdot)$, $k \geq 1$ with

$$\begin{aligned} \omega_i^{\nu,-,k}(s, \cdot) &= f_i *_{sp} G_\nu + \left(\sum_{j=1}^3 v_j^{\nu,-,k-1} \frac{\partial \omega_i^{\nu,-,k-1}}{\partial x_j} \right) * G_\nu \\ &- \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-,k-1}}{\partial x_j} + \frac{\partial v_j^{\nu,-,k-1}}{\partial x_i} \right) \omega_j^{\nu,-,k-1} \right) * G_\nu, \end{aligned} \quad (150)$$

we have

$$\begin{aligned} \omega_i^{\nu,-,k}(s, \cdot) - \omega_i^{\nu,-,k-1}(s, \cdot) &= \left(\sum_{j=1}^3 v_j^{\nu,-,k-1} \frac{\partial \omega_i^{\nu,-,k-1}}{\partial x_j} \right) * G_\nu \\ &- \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-,k-1}}{\partial x_j} + \frac{\partial v_j^{\nu,-,k-1}}{\partial x_i} \right) \omega_j^{\nu,-,k-1} \right) * G_\nu \\ &- \left(\sum_{j=1}^3 v_j^{\nu,-,k-2} \frac{\partial \omega_i^{\nu,-,k-1}}{\partial x_j} \right) * G_\nu \\ &+ \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-,k-2}}{\partial x_j} + \frac{\partial v_j^{\nu,-,k-2}}{\partial x_i} \right) \omega_j^{\nu,-,k-1} \right) * G_\nu \\ &+ \left(\sum_{j=1}^3 v_j^{\nu,-,k-2} \frac{\partial \omega_i^{\nu,-,k-1}}{\partial x_j} \right) * G_\nu \\ &- \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-,k-2}}{\partial x_j} + \frac{\partial v_j^{\nu,-,k-2}}{\partial x_i} \right) \omega_j^{\nu,-,k-1} \right) * G_\nu \\ &- \left(\sum_{j=1}^3 v_j^{\nu,-,k-2} \frac{\partial \omega_i^{\nu,-,k-2}}{\partial x_j} \right) * G_\nu \\ &+ \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-,k-2}}{\partial x_j} + \frac{\partial v_j^{\nu,-,k-2}}{\partial x_i} \right) \omega_j^{\nu,-,k-2} \right) * G_\nu \\ &= \left(\sum_{j=1}^3 \left(v_j^{\nu,-,k-1} - v_j^{\nu,-,k-2} \right) \frac{\partial \omega_i^{\nu,-,k-1}}{\partial x_j} \right) * G_\nu \\ &+ \sum_{j=1}^3 v_j^{\nu,-,k-2} \left(\omega_j^{\nu,-,k-1} - \omega_j^{\nu,-,k-2} \right) * G_\nu \\ &- \left(\sum_{j=1}^3 \left(\frac{1}{2} \left(\frac{\partial \delta v_i^{\nu,-,k-1}}{\partial x_j} + \frac{\partial \delta v_j^{\nu,-,k-1}}{\partial x_i} \right) \right) \omega_j^{\nu,-,k-1} \right) * G_\nu \\ &- \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-,k-2}}{\partial x_j} + \frac{\partial v_j^{\nu,-,k-2}}{\partial x_i} \right) \delta \omega_j^{\nu,-,k-1} \right) * G_\nu. \end{aligned} \quad (151)$$

Hence the increments

$$\delta \omega_i^{\nu,-,k} := \omega_i^{\nu,-,k} - \omega_i^{\nu,-,k-1} \quad (152)$$

satisfy the recursion

$$\begin{aligned}
\delta\omega_i^{\nu,-,k} &= \left(\sum_{j=1}^3 \delta v_j^{\nu,-,k-1} \frac{\partial\omega_i^{\nu,-,k-1}}{\partial x_j} \right) * G_\nu \\
&+ \left(\sum_{j=1}^3 v_j^{\nu,-,k-1} \frac{\partial\delta\omega_i^{\nu,-,k-1}}{\partial x_j} \right) * G_\nu \\
&- \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial\delta v_i^{\nu,-,k-1}}{\partial x_j} + \frac{\partial\delta v_j^{\nu,-,k-1}}{\partial x_i} \right) \omega_j^{\nu,-,k-1} \right) * G_\nu \\
&- \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-,k-2}}{\partial x_j} + \frac{\partial v_j^{\nu,-,k-2}}{\partial x_i} \right) \delta\omega_j^{\nu,-,k-1} \right) * G_\nu,
\end{aligned} \tag{153}$$

where

$$\delta v_i^{\nu,-,k-1} := v_i^{\nu,-,k-1} - v_i^{\nu,-,k-2}. \tag{154}$$

Elementary arguments (we considered similar arguments elsewhere) lead to the first part of the next lemma. The second part (ν -independent estimates) is proved in the next item.

Lemma 2.5. *Given $\nu > 0$ there exists $T > 0$ and a constant $c \in (0, 1)$ (dependent on ν) such that for all $k \geq 3$ we have*

$$\max_{1 \leq i \leq D} \sup_{s \in [0, T]} |\delta\omega_i^{\nu,-,k+1}(s, \cdot)|_{H^2 \cap C^2} \leq c \max_{1 \leq i \leq D} \sup_{s \in [0, T]} |\delta\omega_i^{\nu,-,k}(s, \cdot)|_{H^2 \cap C^2}, \tag{155}$$

and for all $k \geq 3$

$$\max_{1 \leq i \leq D} \sup_{s \in [0, T]} |\delta v_i^{\nu,-,k+1}(s, \cdot)|_{H^3 \cap C^3} \leq c \max_{1 \leq i \leq D} \sup_{s \in [0, T]} |\delta v_i^{\nu,-,k}(s, \cdot)|_{H^3 \cap C^3}. \tag{156}$$

These estimates have the ν -independent counterparts with a loss of regularity. There exists $T > 0$ and a constant $c \in (0, 1)$ independent on ν such that for all $k \geq 3$ we have

$$\max_{1 \leq i \leq D} \sup_{s \in [0, T]} |\delta\omega_i^{\nu,-,k+1}(s, \cdot)|_{H^1 \cap C^1} \leq c \max_{1 \leq i \leq D} \sup_{s \in [0, T]} |\delta\omega_i^{\nu,-,k}(s, \cdot)|_{H^1 \cap C^1}, \tag{157}$$

and for all $k \geq 3$

$$\max_{1 \leq i \leq D} \sup_{s \in [0, T]} |\delta v_i^{\nu,-,k+1}(s, \cdot)|_{H^2 \cap C^2} \leq c \max_{1 \leq i \leq D} \sup_{s \in [0, T]} |\delta v_i^{\nu,-,k}(s, \cdot)|_{H^2 \cap C^2}. \tag{158}$$

Moreover, the statement of (155) and (156) hold for each $t_0 > 0$ in a time interval $[t_0, T]$ for some $T > 0$.

ad v) The previous arguments depend on $\nu > 0$, and we have to show how a viscosity limit can be obtained which preserves essential properties such as contraction with lower regularity or contraction with higher regularity after finite time (as indicated in the last lemma). We have to show va) the estimate above are essentially independent of the viscosity $\nu > 0$, and vb) that this viscosity limit preserve incompressibility.

Ad va) The main observation here is the estimate of

$$v_{i_0}^f *_{sp} G_{\nu, j} \tag{159}$$

for given $1 \leq j \leq D$. First note that for $r = |x|$ we have an upper bound

$$|G_{\nu,j}(t, x)| \leq \frac{c}{(\nu t)^\delta r^{D+1-2\delta}}, \quad (160)$$

where integrability is given for $\delta \in (0.5, 1)$ and c is independent of ν . Independence of c from ν follows from

$$\begin{aligned} |G_{\nu,k}(t, z)| &= \left| \left(\frac{-2z_k}{4\rho\nu t} \right) \frac{1}{\sqrt{4\pi\nu t}^D} \exp\left(-\frac{|z|^2}{4\nu t}\right) \right| \\ &\leq \left| \left(\frac{-2z_k}{4\rho\nu t} \right) \exp\left(-\frac{|z|^2}{8\nu t}\right) \frac{\sqrt{2}}{\sqrt{8\pi\nu t}^D} \exp\left(-\frac{|z|^2}{8\nu t}\right) \right| \\ &\leq \left| \left(\frac{2|z|^2}{4\rho\nu t} \right) \exp\left(-\frac{|z|^2}{8\nu t}\right) \frac{1}{|z|} \frac{\sqrt{2}}{\sqrt{8\pi\nu t}^D} \exp\left(-\frac{|z|^2}{8\nu t}\right) \right| \\ &\leq \left| c_0 \frac{1}{|z|} \frac{\sqrt{2}}{\sqrt{8\pi\nu t}^D} \exp\left(-\frac{|z|^2}{8\nu t}\right) \right| \end{aligned} \quad (161)$$

for some finite constant c_0 which is independent of ν . Then the standard point-wise upper bound estimate for G_ν can be employed and multiplied with $\frac{1}{|z|}$ in order to obtain that c in (163) is independent of ν . Next

$$\begin{aligned} |v_{i_0}^f *_{sp} G_{\nu,j}(t, x)| &\leq \int (r_0)^{\beta_0} \frac{\tilde{c}}{(\nu t)^\delta |r-r_0|^{D+1-2\delta}} r_0^{D-1} dr_0 \\ &\leq c^* + \int |r_0|^{\beta_0-1} \frac{\tilde{c}^*}{(\nu t)^\delta |r-r_0|^{D-2\delta}} r_0^{D-1} dr_0 \\ &= c^* + \int |r_0 - r|^{\beta_0-1} \frac{\tilde{c}^*}{(\nu t)^\delta |r_0|^{D-2\delta}} r_0^{D-1} dr_0 \end{aligned} \quad (162)$$

upon partial integration in a sufficiently large ball and with some finite constants c^* and \tilde{c}^* which do not depend on ν . For each $\nu > 0$ we may consider the right side of the integral in a ball of radius $\sqrt{\nu}$. Outside such a ball we have strong decay due to the Gaussian. Hence we get for given $1 \leq j \leq D$ and $r = |x|$

$$|v_{i_0}^f *_{sp} G_{\nu,j}(t, x)| \leq C^* r^{\beta_0-1}, \quad (163)$$

for a C^* independent of ν . A local solution of the equation

$$\frac{\partial \omega_i^{\nu,-}}{\partial \tau} - \nu \Delta \omega_i^{\nu,-} - \sum_{j=1}^3 v_j^{\nu,-} \frac{\partial \omega_i^{\nu,-}}{\partial y_j} = - \sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-}}{\partial x_j} + \frac{\partial v_j^{\nu,-}}{\partial x_i} \right) \omega_j^{\nu,-}. \quad (164)$$

may be represented in the form

$$\begin{aligned} \omega_i^\nu &= + \left(\sum_{j=1}^3 v_j^{\nu,-} \frac{\partial \omega_i^{\nu,-}}{\partial y_j} \right) * G_\nu - \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-}}{\partial x_j} + \frac{\partial v_j^{\nu,-}}{\partial x_i} \right) \omega_j^{\nu,-} \right) * G_\nu \\ &= - \left(\sum_{j=1}^3 v_{j,j}^{\nu,-} \omega_i^\nu \right) * G_\nu + \sum_{j=1}^3 (v_j^{\nu,-} \omega_i^{\nu,-}) * G_{\nu,j} \\ &\quad - \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-}}{\partial x_j} + \frac{\partial v_j^{\nu,-}}{\partial x_i} \right) \omega_j^{\nu,-} \right) * G_\nu \\ &= \sum_{j=1}^3 (v_j^{\nu,-} \omega_i^{\nu,-}) * G_{\nu,j} - \left(\sum_{j=1}^3 \frac{1}{2} \left(\frac{\partial v_i^{\nu,-}}{\partial x_j} + \frac{\partial v_j^{\nu,-}}{\partial x_i} \right) \omega_j^{\nu,-} \right) * G_\nu, \end{aligned} \quad (165)$$

where we use incompressibility in the latter step. Next in our iteration scheme the first order approximation of the short time vorticity increment value is

$$\begin{aligned} \delta\omega_i^{\nu,-,1} &= \sum_{j=1}^3 \left(v_j^f *_{sp} G_\nu \right) \left(\omega_i^f *_{sp} G_\nu \right) * G_{\nu,j} \\ &- \left(\sum_{j=1}^3 \frac{1}{2} \left(v_i^f *_{sp} G_{\nu,j} + v_j^f *_{sp} G_{\nu,i} \right) \omega_j^f *_{sp} G_\nu \right) * G_\nu, \end{aligned} \quad (166)$$

For the first derivative of the increment we have

$$\begin{aligned} \delta\omega_{i,k}^{\nu,-,1} &= \sum_{j=1}^3 \left(\left(v_j^f *_{sp} G_\nu \right) \left(\omega_i^f *_{sp} G_\nu \right) \right)_{,k} * G_{\nu,j} \\ &- \left(\sum_{j=1}^3 \frac{1}{2} \left(v_i^f *_{sp} G_{\nu,j} + v_j^f *_{sp} G_{\nu,i} \right) \omega_j^f *_{sp} G_\nu \right) * G_{\nu,k}, \end{aligned} \quad (167)$$

From the estimate in (159) it follows that $\sum_{j=1}^3 \left(\left(v_j^f *_{sp} G_\nu \right) \left(\omega_i^f *_{sp} G_\nu \right) \right)_{,k}$ and are spatially Lipschitz such that

$$\delta\omega_i^{\nu,-,1}(t, \cdot) \in H^1 \cap C^1 \quad (168)$$

and the contraction results in (157) and (158) follow straightforwardly. For $t_0 > 0$ we can use this regularity at t_0 and may use the same iteration scheme starting in the interval $[t_0, t_0 + \Delta_0]$ in order to obtain the contraction results in (157) and (158) for some T and all times $t \in [t_0, T]$ for given $t_0 > 0$.

Ad vb) for $1 \leq i \leq D$ consider the local solution of the Navier Stokes type equation (related to the time reversed Euler equation)

$$\begin{aligned} v_i^{\nu,-} &= v_i^{\nu,-}(\cdot) *_{sp} G_\nu - \sum_{j=1}^D v_j^{\nu,-} \frac{\partial v_i^{\nu,-}}{\partial x_j} * G_\nu \\ &- \sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu,-}}{\partial x_j} \frac{\partial v_j^{\nu,-}}{\partial x_m} \right) (t, y) dy * G_\nu \end{aligned} \quad (169)$$

on the time interval $[0, T]$. We have to show that the incompressibility condition holds on the restricted time interval $(0, T]$, i.e.,

$$\text{for all } t \in (0, T] \quad \sum_{i=1}^D v_{i,i}^{\nu,-}(t, \cdot) = 0 \quad (170)$$

The local solution in (169) satisfies the equation

$$\begin{aligned} \frac{\partial v_i^{\nu,-}}{\partial t} - \nu \Delta v_i^{\nu,-} - \sum_{j=1}^D v_j^{\nu,-} \frac{\partial v_i^{\nu,-}}{\partial x_j} \\ + \sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu,-}}{\partial x_j} \frac{\partial v_j^{\nu,-}}{\partial x_m} \right) (t, y) dy. \end{aligned} \quad (171)$$

For $t \in (0, T]$ $v_i^{\nu,-}(t, \cdot) \in H^3 \cap C^3$ such that we can apply the divergence operator in a classical sense to the equation in (171). We get

$$\begin{aligned} \frac{\partial \sum_{i=1}^D v_{i,i}^{\nu,-}}{\partial t} - \nu \Delta \sum_{i=1}^D v_{i,i}^{\nu,-} - \sum_{i=1}^D \sum_{j=1}^D v_{j,i}^{\nu,-} \frac{\partial \sum_{i=1}^D v_{i,i}^{\nu,-}}{\partial x_j} \\ - \sum_{i=1}^D \sum_{j=1}^D v_{j,i}^{\nu,-} v_{i,j}^{\nu,-} \\ + \sum_{i=1}^D \sum_{j,m=1}^D \int_{\mathbb{R}^D} (K_{D,i,i}(\cdot - y)) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu,-}}{\partial x_j} \frac{\partial v_j^{\nu,-}}{\partial x_m} \right) (t, y) dy. \end{aligned} \quad (172)$$

The last two terms in (172) encode the Poisson equation $\Delta p^- = \sum_{i=1}^D \sum_{j=1}^D v_{j,i}^{\nu,-} v_{i,j}^{\nu,-}$ (with p^- the analogue of pressure), and we have $\sum_{i=1}^D v_{i,i}^f = 0$ in H^1 sense. Hence, $\sum_{i=1}^D v_{i,i}^{\nu,-}(t, \cdot) = 0$ in classical sense is consistent with (171). Note that we have chosen v_j^f , $1 \leq i \leq D$, $j \neq i_0$ at item i) above such that

$$\sum_{i=1}^D v_i^f *_{sp} G_{\nu,i}(t, \cdot) = 0. \quad (173)$$

We can use the local iteration scheme and check directly that the incompressibility condition holds, for the limit, i.e., $\sum_{i=1}^D v_{i,i}^{\nu,-}(t, \cdot) = 0$.

ad vi) Choose a time horizon $T > 0$ as in the previous step such that contraction holds for the higher order increments $\delta v_i^{\nu,-,k}$ with $k \geq 3$ as in (3). We choose the increment $\delta v_i^{\text{init},\nu,-,2}$ obtained after $k = 2$ iterations of the scheme. We observed that there is an upper bound $C > 0$ independent of ν such that

$$\begin{aligned} & \left| \delta v_i^{\text{init},\nu,-,2} + \sum_{l=3}^{\infty} \delta v_i^{\nu,-,l} \right|_{H^2 \cap C^2} \\ & = \left| v_i^{\nu,-}(t, \cdot) - v_i^f *_{sp} G_{\nu}(t, \cdot) \right|_{H^2 \cap C^2} \leq C \end{aligned} \quad (174)$$

where $C > 0$ is independent of $\nu > 0$. For all $\nu > 0$ the function

$$v_i^{\nu,-}(t, \cdot) = v_i^f *_{sp} G_{\nu}(t, \cdot) + \delta v_i^{\text{init},\nu,-,2} + \sum_{l=3}^{\infty} \delta v_i^{\nu,-,l} \quad (175)$$

satisfies the Navier stokes equation on the time interval $[0, T]$ such that

$$\begin{aligned} \nu \Delta v_i^{\nu,-} &= \frac{\partial v_i^{\nu,-}}{\partial t} - \sum_{j=1}^D v_j^{\nu,-} \frac{\partial v_i^{\nu,-}}{\partial x_j} \\ &+ \sum_{j,m=1}^D \int_{\mathbb{R}^D} \left(\frac{\partial}{\partial x_i} K_D(\cdot - y) \right) \sum_{j,m=1}^D \left(\frac{\partial v_m^{\nu,-}}{\partial x_j} \frac{\partial v_j^{\nu,-}}{\partial x_m} \right) (t, y) dy, \end{aligned} \quad (176)$$

where the right side of (176) is the time-reversed Euler equation operator applied to $v_i^{\nu,-}$. Recall that the time horizon $T > 0$ is structurally independent of ν in the contraction result. Next we may use the strong polynomial decay at spatial infinity in order to apply a strong compactness argument. Here we note that Rellich's theorem holds only with restrictions for unbounded domains. It is convenient to use the strong polynomial decay at spatial infinity in order to transform on a bounded domain. Moreover spatial transformations to bounded domains allow for convergence constructions in C^m -Banach spaces. Here recall that

Proposition 2.6. *For open and bounded $\Omega \subset \mathbb{R}^n$ and consider the function space*

$$C^m(\Omega) := \left\{ f : \Omega \rightarrow \mathbb{R} \mid \partial^\alpha f \text{ exists for } |\alpha| \leq m \right. \quad (177)$$

and $\partial^\alpha f$ has an continuous extension to $\bar{\Omega}$ }

where $\alpha = (\alpha_1, \dots, \alpha_n)$ denotes a multiindex and ∂^α denote partial derivatives with respect to this multiindex. Then the function space $C^m(\bar{\Omega})$ with the norm

$$|f|_m := |f|_{C^m(\bar{\Omega})} := \sum_{|\alpha| \leq m} |\partial^\alpha f| \quad (178)$$

is a Banach space. Here,

$$|f| := \sup_{x \in \Omega} |f(x)|. \quad (179)$$

We choose a sequence $(\nu_p)_{p \geq 1}$ converging to zero and consider the spatial transformation

$$\delta v_i^{c, \text{init}, -, \nu_p, 2}(t, y) = \delta v_i^{\text{init}, -, \nu_p, 2}(t, x) \quad (180)$$

for $y_j = \arctan(x_j)$, $1 \leq j \leq D$ and for all $t \in [0, T]$. Note that for

$$|\delta v_i^{\text{init}, -, \nu_p, 2}(t, x)| \leq \frac{c}{1 + |x|^{2m}} \quad (181)$$

with a finite ν -independent constant c and multiindices γ with $0 \leq |\gamma| \leq 3 + \epsilon$ for some $\epsilon > 0$ we have for all $t \in [0, T]$ and all $x \in \mathbb{R}^D$

$$|D_y^\gamma \delta v_i^{c, \text{init}, -, \nu_p, 2}(t, y)| \leq c_0(1 + |x|^{2m}) |D_x^\gamma \delta v_i^{\text{init}, -, \nu_p, 2}(t, x)| \leq C \quad (182)$$

for some finite constants c_0, C . This implies

$$\delta v_i^{c, \text{init}, -, 0, 2}(t, \cdot) := \lim_{\nu_p \downarrow 0} \delta v_i^{c, \text{init}, -, \nu_p, 2}(t, \cdot) \in H^2 \cap C^2 \quad \text{for all } 0 \leq t \leq T, \quad (183)$$

and as for some finite $C > 0$ independent of ν_p

$$\sup_{\nu_l > 0} |(1 + |x|^{2m}) |D_x^\gamma \delta v_i^{\text{init}, -, \nu_p, 2}(t, x)| \leq C \quad (184)$$

we conclude that for all $t \in [0, T]$

$$\delta v_i^{c, \text{init}, -, 0, 2}(t, \cdot) \in H^2 \cap C^2. \quad (185)$$

The latter statement transfers to $\delta v_i^{\text{init}, 0, -, 2}(t, \cdot)$, $1 \leq i \leq D$. Similarly for the higher order increments $\delta v_i^{\nu, -, k}$, $1 \leq i \leq D$ for $k \geq 3$. Hence the viscosity limit v_i^- , $1 \leq i \leq D$ satisfies

$$v_i^-(t, \cdot) - v_i^f *_{sp} G_\nu(t, \cdot) = \delta v_i^-(t, \cdot) \in H^2 \cap C^2 \quad (186)$$

for all $t \in [0, T]$, where indeed

$$\delta v_i^- := \lim_{\nu_p \downarrow 0} \delta v_i^{\text{init}, \nu_p, -, 2} + \sum_{l=2}^{\infty} \delta v_i^{\nu_l, -, l} \in C^1((0, T], H^2 \cap C^2). \quad (187)$$

Next, we observe that the function v_i^- , $1 \leq i \leq D$ of the described regularity is a classical solution of the time-reversed Euler equation. Indeed, for some finite $C > 0$

$$\lim_{\nu_p \downarrow 0} \nu_p |\Delta \delta v_i^-(t, \cdot)|_{C \cap L^2} \leq \lim_{\nu_p \downarrow 0} \nu_p \sup_{\nu_p} |\delta v_i^{\nu_p, -, 2}(t, \cdot)|_{H^2 \cap C^2} \leq \nu_p C \downarrow 0 \quad \text{as } \nu_p \downarrow 0 \quad (188)$$

for the left side in (176), and hence the right side of (176) is also 0. Note that the set of continuous functions on \mathbb{R}^D which vanish at spatial infinite is closed. In our context of functions with very strong polynomial decay we may even transform to a bounded domain. Hence we may consider norms $|f|_{C^0} =$

$\sup_x |f(x)|$ and similar norms $|f|_{C^m}$ for derivatives up to order $1 \leq m \leq 3$ in our context. Furthermore, we have

$$\lim_{\nu_p \downarrow 0} \nu_p |\Delta(v_i^f *_{sp} G_\nu)(t, \cdot)|_{C^0} = \lim_{\nu_p \downarrow 0} \nu_p^{1-\delta} \left| \sum_j v_{i,j}^f *_{sp} \nu^\delta G_{\nu,j}(t, \cdot) \right|_{C^0} \downarrow 0 \text{ as } \nu_p \downarrow 0, \quad (189)$$

where in the last step we may use the local upper bound

$$\begin{aligned} & \left| \int_{|x-y| \leq 1} (\sum_j v_{i,j}^f)(t, y) \nu^\delta G_{\nu,j}(t, x; 0, y) \right| \leq \int c_0 r^{\beta_0 - (2 + \alpha_0)} \frac{c_1}{t^\delta |x-y|^{4-2\delta}} dy \\ & \leq c_2 r^{\beta_0 + (2\delta - 1) - (2 + \alpha_0)} \text{ for } 0 \leq r = \sqrt{y_1^2 + y_2^2 + y_3^2} \end{aligned} \quad (190)$$

in a ball $B_1(x)$ of radius 1 around x , $\delta \in (0.5, 1)$, and for some finite constants c_0, c_1, c_2 which are independent of ν . For β_0 close to $2 + \alpha_0$ this integral is bounded. Note that this is also achieved for weaker parameter conditions (for α_0, β_0) if δ is chosen to be $1 - \epsilon$ for small $\epsilon > 0$. Note that the integral outside this ball clearly converges to zero as $\nu_p \downarrow 0$. Here, note that we used the factor ν^δ for the latter conclusion in order to ensure that a standard estimate of the Gaussian is independent of ν . Similar arguments show that the velocity viscosity limit v^- has spatial regularity $H^3 \cap C^3$ for positive time corresponding to a spatial vorticity regularity $H^2 \cap C^2$ for positive time $t > 0$. We conclude that the original Euler equation develops in opposite time direction from data at time $t = 0$ (corresponding to time $s = T$ of the reversed-time Euler equation) a weak singularity at time $T > 0$ (corresponding to data at time $s = 0$ the time-reversed Euler equation).

References

- [1] LANDAU, L., LIFSCHITZ, E. *Lehrbuch der Theoretischen Physik VI, Hydrodynamik*, Akademie Verlag, Berlin, J., (1978).
- [2] MAJDA, A., BERTOZZI, L. *Vorticity and Incompressible Flow (Cambridge Texts in Applied Mathematics)* Cambridge University Press, 2001.