

Searching for degenerate Higgs bosons

A profile likelihood ratio method to test for mass-degenerate states in the presence of incomplete data and uncertainties

André David^{a,1}, Jaana Heikkilä^{b,1,2}, Giovanni Petrucciani^{c,1}

¹PH Department, CERN, Switzerland

²Helsinki Institute of Physics, Finland

Received: date / Accepted: date

Abstract Using the likelihood ratio test statistic, we present a method which can be employed to test the hypothesis of a single Higgs boson using the matrix of measured signal strengths. This method can be applied in the presence of incomplete data and takes into account uncertainties on the measurements. The p-value against the hypothesis of a single Higgs boson is defined from the expected distribution of the test statistic, generated using pseudo-experiments. The applicability of the likelihood-based test is demonstrated using numerical examples with uncertainties and missing matrix elements.

Keywords Degenerate states · Matrix rank · Profile likelihood ratio

PACS 12.60.Fr · 14.80.Bn

Mathematics Subject Classification (2010) 15A03 · 62G10

1 Introduction

A new resonance, consistent with the standard model (SM) Higgs boson and with a mass of approximately 125 GeV, has been observed by ATLAS and CMS collaborations at the Large Hadron Collider (LHC) [1–3]. For each production mode and decay mode of the boson, the signal strength can be defined in terms of the observed production cross section and branching ratio relative to the SM expectation as:

$$\mu_{i,j} = \frac{(\sigma_i \cdot \mathcal{B}_j)_{\text{obs}}}{(\sigma_i \cdot \mathcal{B}_j)_{\text{SM}}}, \quad (1)$$

^ae-mail: Andre.David@cern.ch

^be-mail: Jaana.Kristiina.Heikkilae@cern.ch

^ce-mail: Giovanni.Petrucciani@cern.ch

where the indexes i and j stand for the different production modes and decay modes, respectively. The available experimental information can be arranged in a matrix, \mathcal{M} , of signal strengths:

$$\begin{array}{c|ccccc} & \text{H} \rightarrow \gamma\gamma & \text{H} \rightarrow \text{WW} & \text{H} \rightarrow \text{ZZ} & \text{H} \rightarrow \tau\tau & \text{H} \rightarrow \text{bb} \\ \hline \text{ggH} & \mu_{\text{ggH},\gamma\gamma} & \mu_{\text{ggH},\text{WW}} & \mu_{\text{ggH},\text{ZZ}} & \mu_{\text{ggH},\tau\tau} & \mu_{\text{ggH},\text{bb}} \\ \text{VBF} & \mu_{\text{VBF},\gamma\gamma} & \mu_{\text{VBF},\text{WW}} & \mu_{\text{VBF},\text{ZZ}} & \mu_{\text{VBF},\tau\tau} & \mu_{\text{VBF},\text{bb}} \\ \text{VH} & \mu_{\text{VH},\gamma\gamma} & \mu_{\text{VH},\text{WW}} & \mu_{\text{VH},\text{ZZ}} & \mu_{\text{VH},\tau\tau} & \mu_{\text{VH},\text{bb}} \\ \text{ttH} & \mu_{\text{ttH},\gamma\gamma} & \mu_{\text{ttH},\text{WW}} & \mu_{\text{ttH},\text{ZZ}} & \mu_{\text{ttH},\tau\tau} & \mu_{\text{ttH},\text{bb}} \end{array} \quad (2)$$

The rows of \mathcal{M} represent the four main production modes: gluon fusion (ggH), vector boson fusion (VBF), associated production with a Z or W boson (VH), and associated production with a top quark pair (ttH). The columns represent five decay modes¹: $\gamma\gamma$, WW, ZZ, $\tau\tau$ and bb.

It has been suggested that there may be more than one state nearly-degenerate in mass, which could couple differently to the SM particles and produce the observed signal. If more than one state is present, it may be possible to use the rank of \mathcal{M} to test for multiple states without having to directly resolve the resonant peak(s), an approach limited by the experimental mass resolution to few GeV [4–6]. To that end, there are two main challenges that need to be overcome in order to make use of all the experimental information encoded in \mathcal{M} :

1. Some $\mu_{i,j}$ have not yet been measured, such as $\mu_{\text{ttH},\text{ZZ}}$, and are therefore missing, raising the issue of incomplete data.
2. The precision to which different $\mu_{i,j}$ have been measured varies greatly, with the smaller uncertainties usually being related to the processes with the largest production cross-sections.

¹We imply charge conjugation throughout; bb stands for $b\bar{b}$, $\tau\tau$ stands for $\tau^+\tau^-$, etc.

Inspired by the work presented in Ref. [4], we noticed that while their approach tackled the uncertainty in the measurements, it was not able to take into account the missing data aspect. The method presented here addresses the question of whether the experimental observations are compatible with a single Higgs boson using the full information encoded in \mathcal{M} , i.e. coping with missing $\mu_{i,j}$ and taking into account the uncertainties on the $\mu_{i,j}$.

2 The 2×2 case

Consider a 2×2 sub-matrix of \mathcal{M} , namely the one with the ggH and VBF production modes and the $\gamma\gamma$ and WW decay channels:

$$\begin{array}{c|cc} & \text{H} \rightarrow \gamma\gamma & \text{H} \rightarrow \text{WW} \\ \hline \text{ggH} & \mu_{\text{ggH},\gamma\gamma} & \mu_{\text{ggH},\text{WW}} \\ \text{VBF} & \mu_{\text{VBF},\gamma\gamma} & \mu_{\text{VBF},\text{WW}} \end{array} \quad (3)$$

Ignoring the uncertainties on the $\mu_{i,j}$ elements, if there is only one Higgs boson the determinant of this 2×2 matrix is zero and consequently the matrix has rank 1.

However, if there are two particles involved, each signal strength in the matrix can be written as the sum of two terms:

$$\mu_{i,j} = \mu_{i,j}^{(1)} + \mu_{i,j}^{(2)}, \quad (4)$$

where the superscripts 1 and 2 concern each of the two states. When replacing all four matrix elements with sums of form Eq. (4), the determinant becomes, in general, not equal to zero and in that case the matrix rank will be 2.

Instead of the determinant, when studying the rank of a 2×2 matrix, a double ratio can be used without loss of generality:

$$\rho = \frac{\mu_{\text{ggH},\gamma\gamma} \mu_{\text{VBF},\text{WW}}}{\mu_{\text{ggH},\text{WW}} \mu_{\text{VBF},\gamma\gamma}}. \quad (5)$$

This is equivalent to the double ratios introduced in Ref. [7] and the expectation is that $\rho = 1$ if there is only one state. If there are two states, one has

$$\rho = \frac{(\mu_{\text{ggH},\gamma\gamma}^{(1)} + \mu_{\text{ggH},\gamma\gamma}^{(2)}) (\mu_{\text{VBF},\text{WW}}^{(1)} + \mu_{\text{VBF},\text{WW}}^{(2)})}{(\mu_{\text{ggH},\text{WW}}^{(1)} + \mu_{\text{ggH},\text{WW}}^{(2)}) (\mu_{\text{VBF},\gamma\gamma}^{(1)} + \mu_{\text{VBF},\gamma\gamma}^{(2)})}, \quad (6)$$

and ρ can take a range of values, implying the rank of the matrix is 2.

The rank is a discrete quantity that can be exactly computed if the elements of the matrix are also exactly known. If there are independent uncertainties on the matrix elements, the rank cannot be precisely determined. Nevertheless, it is still possible to assess the statistical compatibility of the matrix with one of rank 1. In previous works [4, 7], it was proposed to assess the statistical compatibility via quantities such as ρ which can be assigned an uncertainty obtained by propagating the uncertainties on the matrix elements.

3 The profile likelihood ratio test

In order to evaluate the statistical compatibility of the matrix of observations having rank 1, a model of the matrix elements must be assumed. Such a model defines the correlations between the different $\mu_{i,j}$, which are parameters of interest in the fit to data, but may also include all other parameters that affect the measurements, such as systematic uncertainties. The latter can be treated as nuisance parameters using the profile likelihood method and their values are obtained from a fit to the data [8].

The likelihood is the probability to observe the data assuming that a given model is true. Using the likelihood function $L(\text{data}|\alpha, \beta)$, where α are parameters of interest and β are nuisance parameters, we define the test statistic $q = q(\alpha)$ based on the profile likelihood ratio [8–13]:

$$q(\alpha) = -2 \ln \frac{L(\text{data}|\alpha, \hat{\beta})}{L(\text{data}|\hat{\alpha}, \hat{\beta})}, \quad (7)$$

where $\hat{\beta}$ is the value of β that maximizes the likelihood function for a particular value of α , whereas $\hat{\alpha}$ and $\hat{\beta}$ are the values that maximize the likelihood function overall.

3.1 Application to the 2×2 case

Let's consider again a 2×2 matrix and no nuisance parameters. In this case, the test statistic $q(\rho)$ can be constructed for the double ratio ρ from Eq. (5):

$$q(\rho) = -2 \ln \frac{L(\text{data}|\rho)}{L(\text{data}|\hat{\rho})}. \quad (8)$$

When testing for compatibility with rank 1, the relevant quantity is $q(\rho = 1)$, where the numerator is the likelihood to observe the data assuming a rank 1 matrix, whereas the denominator is the likelihood to observe the data assuming the most general 2×2 matrix.

3.2 Application to \mathcal{M}

The likelihood ratio defined in Eq. (8) can be generalized to any matrix size, which includes \mathcal{M} , a 5×4 matrix. In that case, the denominator is the profile likelihood for the model of the most general 5×4 matrix while the numerator uses the model of a general rank 1 matrix, which we compose as the tensor product of two vectors: $(\mu_{\gamma\gamma}, \mu_{\text{WW}}, \mu_{\text{ZZ}}, \mu_{\tau\tau}, \mu_{\text{bb}})$ and $(1, \lambda_{\text{VBF}}, \lambda_{\text{VH}}, \lambda_{\text{tH}})$, where $\mu_j = \mu_{\text{ggH},j}$ is the signal strength for gluon fusion production and decay mode j , and $\lambda_i = \mu_{i,j} / \mu_{\text{ggH},j}$ are the relative scaling factors between the signal strength in production mode i and that for gluon fusion. Note that, as expected, the rank 1 model entails that the λ_i are the same for all decay modes.

The matrix under each of the alternative hypotheses is:

1. A general rank 1 matrix with eight parameters μ_j , λ_{VBF} , λ_{VH} , and λ_{ttH} :

	H \rightarrow $\gamma\gamma$	H \rightarrow WW	H \rightarrow ZZ	H \rightarrow $\tau\tau$	H \rightarrow bb
ggH	$\mu_{\gamma\gamma}$	μ_{WW}	μ_{ZZ}	$\mu_{\tau\tau}$	μ_{bb}
VBF	$\lambda_{\text{VBF}} \cdot \mu_{\gamma\gamma}$	$\lambda_{\text{VBF}} \cdot \mu_{\text{WW}}$	$\lambda_{\text{VBF}} \cdot \mu_{\text{ZZ}}$	$\lambda_{\text{VBF}} \cdot \mu_{\tau\tau}$	$\lambda_{\text{VBF}} \cdot \mu_{\text{bb}}$
VH	$\lambda_{\text{VH}} \cdot \mu_{\gamma\gamma}$	$\lambda_{\text{VH}} \cdot \mu_{\text{WW}}$	$\lambda_{\text{VH}} \cdot \mu_{\text{ZZ}}$	$\lambda_{\text{VH}} \cdot \mu_{\tau\tau}$	$\lambda_{\text{VH}} \cdot \mu_{\text{bb}}$
ttH	$\lambda_{\text{ttH}} \cdot \mu_{\gamma\gamma}$	$\lambda_{\text{ttH}} \cdot \mu_{\text{WW}}$	$\lambda_{\text{ttH}} \cdot \mu_{\text{ZZ}}$	$\lambda_{\text{ttH}} \cdot \mu_{\tau\tau}$	$\lambda_{\text{ttH}} \cdot \mu_{\text{bb}}$

2. The most general 5×4 matrix with twenty parameters μ_j , λ_{VBF}^j , λ_{VH}^j , and λ_{ttH}^j :

	H \rightarrow $\gamma\gamma$	H \rightarrow WW	H \rightarrow ZZ	H \rightarrow $\tau\tau$	H \rightarrow bb
ggH	$\mu_{\gamma\gamma}$	μ_{WW}	μ_{ZZ}	$\mu_{\tau\tau}$	μ_{bb}
VBF	$\lambda_{\text{VBF}}^{\gamma\gamma} \cdot \mu_{\gamma\gamma}$	$\lambda_{\text{VBF}}^{\text{WW}} \cdot \mu_{\text{WW}}$	$\lambda_{\text{VBF}}^{\text{ZZ}} \cdot \mu_{\text{ZZ}}$	$\lambda_{\text{VBF}}^{\tau\tau} \cdot \mu_{\tau\tau}$	$\lambda_{\text{VBF}}^{\text{bb}} \cdot \mu_{\text{bb}}$
VH	$\lambda_{\text{VH}}^{\gamma\gamma} \cdot \mu_{\gamma\gamma}$	$\lambda_{\text{VH}}^{\text{WW}} \cdot \mu_{\text{WW}}$	$\lambda_{\text{VH}}^{\text{ZZ}} \cdot \mu_{\text{ZZ}}$	$\lambda_{\text{VH}}^{\tau\tau} \cdot \mu_{\tau\tau}$	$\lambda_{\text{VH}}^{\text{bb}} \cdot \mu_{\text{bb}}$
ttH	$\lambda_{\text{ttH}}^{\gamma\gamma} \cdot \mu_{\gamma\gamma}$	$\lambda_{\text{ttH}}^{\text{WW}} \cdot \mu_{\text{WW}}$	$\lambda_{\text{ttH}}^{\text{ZZ}} \cdot \mu_{\text{ZZ}}$	$\lambda_{\text{ttH}}^{\tau\tau} \cdot \mu_{\tau\tau}$	$\lambda_{\text{ttH}}^{\text{bb}} \cdot \mu_{\text{bb}}$

If the matrix has rank 1, the rows in these two models are the same, i.e. $\lambda_{\text{VBF}} \equiv \lambda_{\text{VBF}}^j$, $\lambda_{\text{VH}} \equiv \lambda_{\text{VH}}^j$, and $\lambda_{\text{ttH}} \equiv \lambda_{\text{ttH}}^j$. The parameters of interest are λ_{VBF} (λ_{VBF}^j), λ_{VH} (λ_{VH}^j), and λ_{ttH} (λ_{ttH}^j), while the signal strength parameters are profiled as they have no bearing on the rank of the matrix.

The test statistic q_λ is defined using ratio between the profile likelihood of the two aforementioned models:

$$q_\lambda = -2 \ln \frac{L(\text{data} | \lambda_{\text{VBF}}^j = \hat{\lambda}_{\text{VBF}}^j, \lambda_{\text{VH}}^j = \hat{\lambda}_{\text{VH}}^j, \lambda_{\text{ttH}}^j = \hat{\lambda}_{\text{ttH}}^j)}{L(\text{data} | \hat{\lambda}_{\text{VBF}}^j, \hat{\lambda}_{\text{VH}}^j, \hat{\lambda}_{\text{ttH}}^j)} \quad (9)$$

The numerator is the profile likelihood of data assuming the most general rank 1 matrix and the denominator is the profile likelihood of data assuming the most general 5×4 matrix.

As noted above, if there is only one Higgs boson the λ_i do not depend on the decay mode and the value of the q_λ is not very large. However, if the rank is not equal to 1, the most general 5×4 matrix model will fit the data better than the general rank 1 matrix model and the value of q_λ will be large.

When dealing with more general parameters than the double ratio ρ , the expected distribution of the test statistic is generated using a Monte Carlo simulation. The p-value for the observation is evaluated from the expected test statistic distribution under the hypothesis of the SM Higgs boson.

4 Numerical examples

For ease of comparison, and even if more recent data are available [6, 14–27], the following numerical examples are

carried out using the same signal strength matrix presented in Ref. [4], namely:

	H \rightarrow $\gamma\gamma$	H \rightarrow WW	H \rightarrow ZZ	H \rightarrow $\tau\tau$	H \rightarrow bb
ggH	1.6 ± 0.35	0.8 ± 0.3	1.0 ± 0.3	1.2 ± 0.8	–
VBF	2.1 ± 0.9	-0.2 ± 0.6	–	0.3 ± 0.7	–
VH	1.9 ± 2.6	-0.3 ± 2.1	–	1.0 ± 1.8	0.8 ± 0.6
ttH	–	–	–	–	< 3.8

In the following sub-sections we will consider, in turn, sub-matrices of Eq. (10), building up to the analysis of all the information it provides. While any correlated uncertainties between the matrix elements have not been taken into account, experiments can and should take them into account when using this method.

4.1 (ggH, VBF) \times ($\gamma\gamma$, WW, $\tau\tau$)

As a first example, consider the 2×3 sub-matrix spanned by (ggH, VBF) \times ($\gamma\gamma$, WW, $\tau\tau$) with signal strengths taken from Eq. (10) above. As an approximation, each signal strength is modeled by a Gaussian distribution using the likelihood function for a counting experiment with large event yields. The counting experiment is set up as follows: the expected number of background events B_{exp} is fixed to 1000 and the number of observed events N_{obs} and the expected number of signal events S_{exp} are chosen to reproduce the observed signal strength $\mu_{i,j}$ and its uncertainty. Using this approach, the observed value of the test statistic is $q_\lambda^{\text{obs}} = 4.02$.

Pseudo-experiments are used to estimate the expected distribution of the test statistic under the hypothesis of a single particle. The pseudo-data are generated according to the expectation for the SM Higgs boson hypothesis, where $\mu_j = \lambda_{\text{VBF}} = \lambda_{\text{VBF}}^j = 1$. The expected test statistic distribution is shown in Fig. 1. Small negative values of q_λ are expected from the limited numerical precision when evaluating q_λ , when $q_\lambda \sim 0$. On the other hand, large negative values are due to the failure to fit the pseudo-data with the more general model (in the denominator of q_λ) while the rank 1 model fit converges. These are rare cases, present in this particular scenario that has large input uncertainties. This is in contrast to the scenario considered in the following sub-section, where these large negative values are absent, as expected from the reduced uncertainties.

The p-value for the observed outcome can be derived using the distribution of the test statistic in pseudo-data shown in Fig. 1 by counting how many outcomes have $q_\lambda \geq q_\lambda^{\text{obs}}$:

$$p_{\text{SM}} = P(q_\lambda \geq q_\lambda^{\text{obs}} | \mu_j \equiv 1, \lambda_i^j \equiv 1) \\ = 0.2090.$$

This means that if the matrix has rank 1 we would expect an outcome as extreme, or more extreme, than the observation in $(20.90 \pm 0.03)\%$ of the cases, where the uncertainty




Fig. 1 The distribution of the test statistic, generated using the 2×3 sub-matrix of the Eq. (10), spanned by $(ggH, VBF) \times (\gamma\gamma, WW, \tau\tau)$ and assuming the SM Higgs boson hypothesis. The observed value of is $q_\lambda^{\text{obs}} = 4.02$ and, under the SM Higgs boson hypothesis, the fraction of pseudo-experiments for which $q_\lambda \geq 4.02$ is 0.2090.

quoted is purely statistical and is calculated from the 68.2% Clopper-Pearson interval [28]. The p-value can be translated to into a z score and for that conversion we use the one-tail convention. The z score for the observed p-value above is 0.81 standard deviations (σ) and represents a low significance against the rank 1 hypothesis. This result is in contrast with the findings of Ref. [4], namely 0.4σ . However, both results are compatible with the null hypothesis.

As would be expected, there is no change in the results when extending the matrix to include also the ZZ information from Eq. (10). The reason is that having one single element in any given row or column makes no statement about the rank-related parameters, λ_i .

4.2 Luminosity extrapolation

For a different type of numerical example, let us examine the same 2×3 sub-matrix as above, but scaling the signal strength uncertainties with $1/\sqrt{L} = 1/\sqrt{10}$, a naive way of simulating a 10-fold increase in the amount of data, and also done in Ref. [4]. Because of the decreased uncertainties, the observed value of the test statistic would become $q_\lambda^{\text{high-lumi}} = 38.7$. The expected test statistic distribution can be seen in Fig. 2 and the p-value is found to be:

$$p_{\text{SM,high-lumi}} = P(q_\lambda \geq q_\lambda^{\text{high-lumi}} | \mu_j \equiv 1, \lambda_i^j \equiv 1) \\ = (1.6 \pm 0.3) \times 10^{-5},$$

which would correspond to a $4.15 \pm 0.04\sigma$ significance for rank larger than unity, to be compared with 3.9σ estimated in Ref. [4].

From the comparison of the two test statistic distributions in Figs. 1 and 2, one notes the effect that the reduction




Fig. 2 The distribution of the test statistic, generated using the 2×3 sub-matrix of the Eq. (10), spanned by $(ggH, VBF) \times (\gamma\gamma, WW, \tau\tau)$ and assuming the SM Higgs boson hypothesis. The signal strength uncertainties were scaled with $1/\sqrt{L} = 1/\sqrt{10}$. The observed value of q_λ is $q_\lambda^{\text{high-lumi}} = 38.7$ and, under the SM Higgs boson hypothesis, the fraction of pseudo-experiments for which $q_\lambda \geq 38.7$ is $(1.6 \pm 0.3) \times 10^{-5}$.

of the uncertainties on the signal strengths has in decreasing the number of outcomes with $q_\lambda < 0$, as previously described.

4.3 $(ggH, VBF, VH) \times (\gamma\gamma, WW, \tau\tau)$

Let us now turn to the 3×3 sub-matrix in Eq. (10) that is spanned by $(ggH, VBF, VH) \times (\gamma\gamma, WW, \tau\tau)$.

The observed value of the test statistic is found to be $q_\lambda^{3 \times 3} = 4.28$. The expected test statistic distribution can be seen in Fig. 3 and the p-value is:

$$p_{\text{SM},3 \times 3} = P(q_\lambda \geq q_\lambda^{3 \times 3} | \mu_j \equiv 1, \lambda_i^j \equiv 1) \\ = 0.35,$$

which corresponds to a 0.4σ significance for rank larger than unity.

4.4 $(ggH, VBF, VH) \times (\gamma\gamma, WW, \tau\tau)$ with removed elements

To demonstrate the way in which the proposed technique deals with incomplete data, two different tests with removed elements are presented.

Two elements removed In a first test, the two elements with large uncertainties, namely $\mu_{\text{VBF}, \gamma\gamma}$ and $\mu_{\text{VH}, WW}$, have been removed.




Fig. 3 The distribution of the test statistic, generated using the 3×3 sub-matrix of the Eq. (10), spanned by $(ggH, VBF, VH) \times (\gamma\gamma, WW, \tau\tau)$ and assuming the SM Higgs boson hypothesis. The observed value of q_λ is $q_\lambda^{3 \times 3} = 4.28$ and, under the SM Higgs boson hypothesis, the fraction of pseudo-experiments for which $q_\lambda \geq 4.28$ is 0.35.

The observed value of the test statistic is found to be $q_{\lambda, \text{mis.2}}^{3 \times 3} = 0.28$ and the p-value, defined from the test statistic distribution in Fig. 4, is

$$p_{\text{SM}, 3 \times 3, \text{mis.2}} = P(q_\lambda \geq q_{\lambda, \text{mis.2}}^{3 \times 3} | \mu_j \equiv 1, \lambda_i^j \equiv 1) = 0.81,$$

which corresponds to a -0.86σ significance for rank larger than unity, where the negative sign is a reflection of $q_{\lambda, \text{mis.2}}^{3 \times 3}$ being lower than the median of the SM distribution.




Fig. 4 The distribution of the test statistic, generated using the 3×3 sub-matrix of the Eq. (10), spanned by $(ggH, VBF, VH) \times (\gamma\gamma, WW, \tau\tau)$, with the signal strengths $\mu_{VBF, \gamma\gamma}$ and $\mu_{VH, WW}$ removed, and assuming the SM Higgs boson hypothesis. The observed value of q_λ is $q_{\lambda, \text{mis.2}}^{3 \times 3} = 0.28$ and, under the SM Higgs boson hypothesis, the fraction of pseudo-experiments for which $q_\lambda \geq 0.28$ is 0.81.

Three elements removed In a second test, we remove the $\mu_{VH, \gamma\gamma}$, $\mu_{VBF, WW}$, and $\mu_{ggH, \tau\tau}$ elements from the $(ggH, VBF, VH) \times (\gamma\gamma, WW, \tau\tau)$ matrix.

In this case, a signal strength measurement $\mu_{\tau\tau}$ is absent, implying that the parameterization for the most general 5×4 matrix has more free parameters than there are measurements. This presents a technical difficulty for the numerical minimization used in calculating the profile likelihood. Nevertheless, and without loss of generality, the missing μ_j can be set to unity in the denominator of q_λ , Eq. (9).

The observed value of the test statistic is found to be $q_{\lambda, \text{mis.3}}^{3 \times 3} = 0.31$. The test statistic distribution under the SM Higgs hypothesis can be seen in Fig. 5 and the p-value is

$$p_{\text{SM}, 3 \times 3, \text{mis.3}} = P(q_\lambda \geq q_{\lambda, \text{mis.3}}^{3 \times 3} | \mu_j \equiv 1, \lambda_i^j \equiv 1) = 0.44,$$

which corresponds to a 0.15σ significance for rank larger than unity.




Fig. 5 The distribution of the test statistic, generated using the 3×3 sub-matrix of the Eq. (10), spanned by $(ggH, VBF, VH) \times (\gamma\gamma, WW, \tau\tau)$, after removing three elements, and assuming the SM Higgs boson hypothesis. The observed value of q_λ is $q_{\lambda, \text{mis.3}}^{3 \times 3} = 0.31$ and, under the SM Higgs boson hypothesis, the fraction of pseudo-experiments for which $q_\lambda \geq 0.31$ is 0.44.

Conclusion

We have developed a method that can be used to test for the presence of multiple Higgs bosons. The method can be directly used by the ATLAS and CMS collaborations as it builds upon the profile likelihood techniques already in use for Higgs measurements at the LHC. The main features of method are that it can test for (in)dependence in arbitrarily-sized linear systems in the presence of uncertainties and of missing elements of data.

References

1. ATLAS Collaboration, “Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC”, *Phys. Lett. B* **716** (2012) 1, doi:10.1016/j.physletb.2012.08.020, arXiv:1207.7214.
2. CMS Collaboration, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC”, *Phys. Lett. B* **716** (2012) 30, doi:10.1016/j.physletb.2012.08.021, arXiv:1207.7235.
3. CMS Collaboration, “Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, *J. High Energy Phys.* **06** (2013) 081, doi:10.1007/JHEP06(2013)081, arXiv:1303.4571.
4. Y. Grossman, Z. Surujon, and J. Zupan, “How to test for mass degenerate Higgs resonances”, *J. High Energy Phys.* **03** (2013) 176, doi:10.1007/JHEP03(2013)176, arXiv:1301.0328.
5. ATLAS Collaboration, “Measurement of the Higgs boson mass from the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ channels with the ATLAS detector using 25 fb⁻¹ of pp collision data”, *Phys. Rev. D* **90** (2014) 052004, doi:10.1103/PhysRevD.90.052004, arXiv:1406.3827.
6. CMS Collaboration, “Observation of the diphoton decay of the 125 GeV Higgs boson and measurement of its properties”, *Eur. Phys. J. C* **74** (2014) 3076, doi:10.1140/epjc/s10052-014-3076-z, arXiv:1407.0558.
7. J. F. Gunion, Y. Jiang, and S. Kraml, “Diagnosing Degenerate Higgs Bosons at 125 GeV”, *Phys. Rev. Lett.* **110** (2013) 051801, doi:10.1103/PhysRevLett.110.051801, arXiv:1208.1817.
8. G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011), no. 2, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727.
9. J. Neyman and E. S. Pearson, “On the Use and Interpretation of Certain Test Criteria for Purposes of Statistical Inference: Part I”, *Biometrika* **20A** (1928), no. 1-2, 175–240, doi:10.1093/biomet/20A.1-2.175.
10. J. Neyman and E. S. Pearson, “On the Use and Interpretation of Certain Test Criteria for Purposes of Statistical Inference: Part II”, *Biometrika* **20A** (1928), no. 3-4, 263–294, doi:10.1093/biomet/20A.3-4.263.
11. Particle Data Group, K. A. Olive et al., “Review of Particle Physics”, *Chin. Phys. C* **38** (2014), no. 9, 090001, doi:10.1088/1674-1137/38/9/090001.
12. G. Cowan, “Statistical data analysis”. Oxford Univ. Press, Oxford, 1998.
13. A. Stuart, J. K. Ord, and S. Arnold, “Kendall’s advanced theory of statistics. Vol. 2A, Classical inference and the linear model”. Arnold, London, 6th edition, 2004.
14. CMS Collaboration, “Search for the standard model Higgs boson produced in association with a W or a Z boson and decaying to bottom quarks”, *Phys. Rev. D* **89** (2014) 012003, doi:10.1103/PhysRevD.89.012003, arXiv:1310.3687.
15. CMS Collaboration, “Measurement of Higgs boson production and properties in the WW decay channel with leptonic final states”, *J. High Energy Phys.* **01** (2014) 096, doi:10.1007/JHEP01(2014)096, arXiv:1312.1129.
16. CMS Collaboration, “Measurement of the properties of a Higgs boson in the four-lepton final state”, *Phys. Rev. D* **89** (2014) 092007, doi:10.1103/PhysRevD.89.092007, arXiv:1312.5353.
17. CMS Collaboration, “Evidence for the 125 GeV Higgs boson decaying to a pair of τ leptons”, *J. High Energy Phys.* **05** (2014) 104, doi:10.1007/JHEP05(2014)104, arXiv:1401.5041.
18. CMS Collaboration, “Search for a Higgs boson decaying into a Z and a photon in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, *Phys. Lett. B* **726** (2013) 587, doi:10.1016/j.physletb.2013.09.057, arXiv:1307.5515.
19. ATLAS Collaboration, “Search for Higgs boson decays to a photon and a Z boson in pp collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS detector”, *Phys. Lett. B* **732** (2014) 8, doi:10.1016/j.physletb.2014.03.015, arXiv:1402.3051.
20. ATLAS Collaboration, “Search for Invisible Decays of a Higgs Boson Produced in Association with a Z Boson in ATLAS”, *Phys. Rev. Lett.* **112** (2014) 201802, doi:10.1103/PhysRevLett.112.201802, arXiv:1402.3244.
21. CMS Collaboration, “Search for invisible decays of Higgs bosons in the vector boson fusion and associated ZH production modes”, *Eur. Phys. J. C* **74** (2014) 2980, doi:10.1140/epjc/s10052-014-2980-6, arXiv:1404.1344.
22. CMS Collaboration, “Search for the associated production of the Higgs boson with a top-quark pair”,

-
- J. High Energy Phys.* **09** (2014) 087,
[doi:10.1007/JHEP09\(2014\)087](https://doi.org/10.1007/JHEP09(2014)087),
[arXiv:1408.1682](https://arxiv.org/abs/1408.1682). [Erratum:
[doi:10.1007/JHEP10\(2014\)106](https://doi.org/10.1007/JHEP10(2014)106)].
23. ATLAS Collaboration, “Search for the Standard Model Higgs boson decay to $\mu^+\mu^-$ with the ATLAS detector”, *Phys. Lett. B* **738** (2014) 68,
[doi:10.1016/j.physletb.2014.09.008](https://doi.org/10.1016/j.physletb.2014.09.008),
[arXiv:1406.7663](https://arxiv.org/abs/1406.7663).
24. ATLAS Collaboration, “Measurements of fiducial and differential cross sections for Higgs boson production in the diphoton decay channel at $\sqrt{s} = 8$ TeV with ATLAS”, *J. High Energy Phys.* **09** (2014) 112,
[doi:10.1007/JHEP09\(2014\)112](https://doi.org/10.1007/JHEP09(2014)112),
[arXiv:1407.4222](https://arxiv.org/abs/1407.4222).
25. ATLAS Collaboration, “Fiducial and differential cross sections of Higgs boson production measured in the four-lepton decay channel in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Lett. B* **738** (2014) 234,
[doi:10.1016/j.physletb.2014.09.054](https://doi.org/10.1016/j.physletb.2014.09.054),
[arXiv:1408.3226](https://arxiv.org/abs/1408.3226).
26. ATLAS Collaboration, “Measurements of Higgs boson production and couplings in the four-lepton channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector”, *Phys. Rev. D* **91** (2015) 012006, [doi:10.1103/PhysRevD.91.012006](https://doi.org/10.1103/PhysRevD.91.012006),
[arXiv:1408.5191](https://arxiv.org/abs/1408.5191).
27. ATLAS Collaboration, “Measurement of Higgs boson production in the diphoton decay channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector”, *Phys. Rev. D* **90** (2014) 112015, [doi:10.1103/PhysRevD.90.112015](https://doi.org/10.1103/PhysRevD.90.112015),
[arXiv:1408.7084](https://arxiv.org/abs/1408.7084).
28. C. J. Clopper and E. S. Pearson, “The Use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial”, *Biometrika* **26** (1934), no. 4, 404–413,
[doi:10.1093/biomet/26.4.404](https://doi.org/10.1093/biomet/26.4.404).