

On heteroclinic separators of magnetic fields in electrically conducting fluids

V. Grines¹, T. Medvedev¹, O. Pochinka², and E. Zhuzhoma²

¹Lobachevsky State University of Nizhni Novgorod, 23 Prospekt Gagarina, 603950, Nizhni Novgorod, Russia

²National Research University Higher School of Economics, 25/12 Bolshaya Pecherskaya, 603005, Nizhni Novgorod, Russia

April 21, 2022

Abstract

In this paper we partly solve the problem of existence of separators of a magnetic field in plasma. We single out in plasma a 3-body with a boundary in which the movement of plasma is of special kind which we call an (a-d)-motion. We prove that if the body is the 3-annulus or the “fat” orientable surface with two holes then the magnetic field necessarily has a heteroclinic separator. The statement of the problem and the suggested method for its solution lead to some theoretical problems from Dynamical Systems Theory which are of interest of their own.

Keywords— magnetic field, plasma, separator, fan

1 Introduction

The paper is about the topological structure of magnetic fields in electrically conducting fluids (collision-dominated plasmas such as stellar interiors). Such magnetic fields are common in the Universe and the problem of their origin and evolution is important for the understanding of many astrophysical and geophysical processes. It is probably safe to say that a magnetic field is a normal accompaniment of any cosmic body that is both fluid (wholly or in part) and rotating. The interactions between a magnetic field and a given conducting fluid are adequately described by the equations of magnetohydrodynamics (MHD). The velocities considered are orders of magnitude smaller than the speed of light c . The evolution of a magnetic field \vec{H} in a moving conductor is described by the induction equation

$$\frac{\partial \vec{H}}{\partial t} = \text{rot} [\vec{v} \vec{H}] + \eta \nabla^2 \vec{H}$$

where \vec{v} is the velocity of the conducting fluid and η is the magnetic diffusivity of the medium (for more details see [1, 2, 3]). Literature on MHD often uses the magnetic induction \vec{B} , for which $\vec{B} = \mu\vec{H}$ where μ is the magnetic permeability of the free space.

Hannes Alfvén [4] showed that in a fluid with a large magnetic Reynolds number $\frac{1}{\eta}$ the field lines move as though they are "frozen" into the medium. As a consequence the topological structure of the magnetic field preserves under a short-time steady motion, though it can bifurcate under a long-time and turbulent, in sense, motion of the medium. The concept of frozenness implies the existence of locations where the magnetic field changes in either directions or magnitude or both (so called x -points). According to *Ampère's law* $\mu\vec{j} = \nabla \times \vec{B}$, the density of the electric current \vec{j} is high in the neighborhood of the x -points and this gives rise to special configurations of the current such as current lines and current sheets. These current configurations may include points in which the magnetic field vanishes (null points or neutral points). Null points can occur if the magnetized medium is either a conducting plasma or a neutral gas. In plasmas, null points typically give rise to current sheets [5, 6]. Linear null points in three dimensions typically look like a saddle of the vector field \vec{H} . Moreover, such a saddle is a conservative one with nonzero eigenvalues $\lambda_1, \lambda_2, \lambda_3$ that satisfy the condition $\lambda_1 + \lambda_2 + \lambda_3 = 0$, due to $\nabla \cdot \vec{H} = 0$. Two quite distinct families of field lines pass through a null point. The null *spine* is the isolated field line which approaches or recedes from the null. Its neighboring field lines form two bundles which spread out as they move away from the null and approach a surface which constitutes the null *fan*, see Fig. 1 (a). Thus, from the point of view of Dynamical Systems Theory, a spine and fan are one-dimensional and two-dimensional separatrices respectively and the term "separatrix" means either of them. A magnetic line joining two nulls and representing the intersection of two fans is called a *separator* [5, 6]. A separator is *heteroclinic* if it is a transversal intersection of the fans, see Fig. 1 (b).




Figure 1: The structure of null point (a), and heteroclinic separator (b).

The topological structure of a magnetic field is defined by null points, spines, fans and separators, the union of those forming the so-called *skeleton* of the magnetic field. To study the global magnetic topology we have to consider first the problem of the existence of null points and separators. Experiments and obser-

vations show that the evolution of the structure of the magnetic field is similar to relaxation processes. At first plasma evolves slowly for some considerable time but at some point there occurs a topological restructuring of the magnetic configuration (reconnection) [5, 6]. We consider the problem of the global magnetic topology under steady fluids when the skeleton is invariant (before reconnection).

We suggest to approach this problem in the following way. We consider a body in plasma (i.e. a part of plasma) of special kind and we study movements of plasma such that all the boundary components of the body move inside or outside of the body so that after some time interval all the boundary components are parallel to the initial boundary components (see exact definitions below). The statement of the problem implies that during the movement the topological structure of the magnetic field remains unchanged, therefore it is natural to assume the skeleton of the magnetic field to be invariant with respect to the movement of plasma. Notice that we do not demand all the points of the skeleton to be fixed, we just assume that the points on the skeleton, while moving, remain on the skeleton. We also assume (and this is the only essential restriction) that the null points are not only null hyperbolic points of the magnetic field but they also are null hyperbolic points of the movement of plasma. Taking into consideration that the Kupka-Smale theorem from Dynamical System Theory states that for every typical movement all periodical points (including fixed points) are hyperbolic [7], we consider only the class of typical movements of plasma. The suggested approach lets us apply methods and results of Dynamical Systems Theory because it turns out to be possible to extend the motion of plasma to some 3-manifold in such a way that one gets a classical dynamical system. Though doing so we loose information on the structure of the magnetic field outside the considered body, on the other hand we get an instrument to study its structure in some part of space. Besides, the global structure on the considered manifold sheds light on possible real structures of the magnetic field. The suggested approach to the problem and the method lead to theoretical issues of Dynamical Systems Theory whose solution is of interest in its own.

2 Formulation of the main results

Let M_p^2 be an orientable closed surface of genus $p \geq 0$ smoothly embedded in \mathbb{R}^3 . Two surfaces $M_{p_1,1}^2, M_{p_2,2}^2 \subset \mathbb{R}^3$ are called *parallel* if they bound the set which is homeomorphic to $M_p^2 \times (0; 1)$. As a consequence, $p_1 = p_2 = p$ and $M_{p,1}^2 \cap M_{p,2}^2 = \emptyset$. The closure of $M_p^2 \times [0; 1]$, in short a *fat surface*, is a 3-manifold with the boundary $M_{p,1}^2 \cup M_{p,2}^2$. In particular, $\mathcal{S} = \mathbb{S}^2 \times [0; 1]$ is a *fat sphere* where \mathbb{S}^2 is the 2-sphere. One of its boundary 2-spheres, say $S^2 \times \{0\} = S_{int}$, bounds an open 3-ball in \mathbb{R}^3 which is disjoint from \mathcal{S} . We call S_{int} *interior* while the sphere $S^2 \times \{1\} = S_{ext}$ is called *exterior*. Let B_{int} and B_{ext} be the 3-balls bounded by S_{int} and S_{ext} respectively.

Let G_p be a fat surface bounded by parallel surfaces $M_{p,1}^2$ and $M_{p,2}^2$ of genus $p \geq 1$ and let it contain two 2-spheres S_1 and S_2 such that $S_1 \cap S_2 = \emptyset$ and the

balls B_1 and B_2 , bounded by S_1 and S_2 respectively, do not intersect. Denote by \mathcal{M} the fat surface G_p with two holes, i.e. the set $G_p \setminus \{B_1 \cup B_2\}$.

Let M be either \mathcal{S} or \mathcal{M} which is smoothly embedded into \mathbb{R}^3 . Suppose M be a part of space with plasma of some astrophysical object with a magnetic field \vec{B} . Denote by \vec{B}_0 the restriction of \vec{B} to M and let all the null points of \vec{B} be typical (i.e. hyperbolic). As a consequence we have that M has only finitely many null points.

We assume that

1. the separatrices of the null points intersect transversally (if at all);
2. if M is the fat surface with holes \mathcal{M} the separatrices intersect the components S_1 and S_2 transversally (if intersect at all).

A map $f_0 : M \rightarrow f_0(M) \subset \mathbb{R}^3$ is said to be an (a-d)-*motion* if it satisfies the following conditions:

- (a) f_0 is an orientation preserving diffeomorphism to its image. The non-wandering set of f_0 consists of fixed hyperbolic points which coincide with the null points of the vector field \vec{B}_0 .
- (b) The boundary components of $f_0(M)$ are pairwise disjoint from the boundary components of M .
- (c) If M is the fat sphere \mathcal{S} then the interior boundary sphere maps inside \mathcal{S} while the external boundary sphere maps outside \mathcal{S} , i.e. $f_0(S_{int}) \subset \mathcal{S}$ and $f_0(S_{ext}) \subset \mathbb{R}^3 \setminus (\mathcal{S} \cup B_{int})$ (see Figure 2).

If M is the fat surface with holes \mathcal{M} then one of the boundary spheres, say S_1 , maps inside \mathcal{M} while the other boundary sphere S_2 maps outside \mathcal{M} , one of the boundary surfaces $M_{p,i}^2$, say $M_{p,1}^2$, maps inside \mathcal{M} while the other boundary surface $M_{p,2}^2$ maps outside \mathcal{M} and the restriction $f_0|_{M_{p,i}^2} : M_{p,i}^2 \rightarrow f_0(M_{p,i}^2)$ is homotopy trivial for each $i = 1, 2$. Moreover, every $x \in M_{p,1}^2$ which does not belong to a separatrix leaves \mathcal{M} through S_2 , i.e. $f_0^k(x) \in B_2$ for some $k > 1$ whereas for the reverse motion every $x \in M_{p,2}^2$ which does not belong to a separatrix leaves \mathcal{M} through S_1 , i.e. $f_0^{-k}(x) \in B_1$ for some $k > 1$.

- (d) The fans and the spines are invariant with respect to f_0 and the fixed points of f_0 are of the same type as the null points of the field \vec{B}_0 .

Some explanation on homotopy triviality. Unlike sphere for which there is only one homotopy class of orientation preserving diffeomorphisms, surfaces of genus greater than zero have a countable family of such classes. Since the surfaces $M_{p,i}^2$, $i = 1, 2$ are parallel, the generators of their fundamental groups are naturally isomorphic. The homotopy triviality then means that the restrictions $f_0|_{M_{p,i}^2}$ are homotopy identical.




Figure 2: (a-d)-motion of the fat sphere \mathcal{S} .

Notice that we do not assume that the magnetic lines of the field $\vec{\mathbf{B}}_0$ intersect the boundary components of M transversally. Therefore, the fans and the spines can, in general, intersect the boundary components of M by several connected components. This mathematical model physically means that we consider a motion of plasma in a period during which the singular points and their fans and spines are preserved. It follows from these properties that the separators (if they exist) are invariant with respect to f_0 and their number (including zero) does not change during this time period.

Our main results for an (a-d)-motion of a fat sphere of plasma are the following.

Theorem 1. *Let $f_0 : \mathcal{S} \rightarrow f_0(\mathcal{S}) \subset \mathbb{R}^3$ be an (a-d)-motion of a fat sphere \mathcal{S} of plasma with a magnetic field $\vec{\mathbf{B}}_0$. Let $\vec{\mathbf{B}}_0$ have in \mathcal{S} null points. Then the number of these points is even (and there are at least two of them), their fans intersect and there are finitely many (not zero) heteroclinic separators.*

Corollary 1. *Let the assumptions of Theorem 1 hold and assume that the spines of the magnetic field $\vec{\mathbf{B}}_0$ do not intersect the fans in \mathcal{S} . Then the fan of each null point contains at least one heteroclinic separator in \mathcal{S} .*

For an (a-d)-motion of a fat surface \mathcal{M} of plasma the results are the following.

Theorem 2. *Let $f_0 : \mathcal{M} \rightarrow f_0(\mathcal{M}) \subset \mathbb{R}^3$ be an (a-d)-motion of a fat surface \mathcal{M} which belongs to some domain of plasma with a magnetic field $\vec{\mathbf{B}}_0$. Then the field $\vec{\mathbf{B}}_0$ has in \mathcal{M} at least two null points such that their fans intersect and there are finitely many (more than zero) heteroclinic separators.*

Corollary 2. *Let the assumptions of Theorem 2 hold and assume that the spines of the null points of the vector field $\vec{\mathbf{B}}_0$ do not intersect the fans in \mathcal{M} . The fan of each null point has in \mathcal{M} at least one heteroclinic separator.*

3 Proof of the main results

We now recall some notions and facts about Morse-Smale diffeomorphisms. More detailed explanations can be found for example in [7, 8, 9, 10]. Denote the set of the non-wandering points of a diffeomorphism f by $NW(f)$. For $p \in NW(f)$ denote by $W^s(p)$ (by $W^u(p)$) the stable (respectively unstable) manifold of the point p . A diffeomorphism f is said to be *Morse-Smale* if its non-wandering set $NW(f)$ is hyperbolic, $NW(f)$ consists of finitely many points and for any two distinct points $p, q \in NW(f)$ the invariant manifolds $W^s(p), W^u(q)$ intersect transversally (if intersect at all). A Morse-Smale diffeomorphism is said to be *gradient-like*, if for any two periodic points $p, q \in NW(f)$ from $W^u(p) \cap W^s(q) \neq \emptyset$ it follows that $\dim W^s(p) < \dim W^s(q)$. A point $x \in M$ is said to be *heteroclinic* if x is the point of transversal intersection of invariant manifolds $W^s(p), W^u(q)$ where $p, q \in NW(f)$ and $\dim W^s(p) = \dim W^s(q)$. It is proved that a Morse-Smale diffeomorphism is gradient-like if and only if it has no heteroclinic points [10]. A Morse-Smale diffeomorphism is said to be *polar* if its non-wandering set consists of exactly one source and one sink.

If $W^u(p) \cap W^s(q) \neq \emptyset$ and $\dim W^s(p) < \dim W^s(q)$ then a connected component of $W^u(p) \cap W^s(q)$ is said to be *heteroclinic submanifold*. If the dimension of the manifold equals 3 then every heteroclinic submanifold is either a simple closed curve (which is homeomorphic to the circle) or a non-closed curve without self-intersections (homeomorphic to the open interval). We call such curves *heteroclinic*.

Let p be a periodic point of an orientation preserving Morse-Smale diffeomorphism f . The *Morse index* $u(p)$ of p is the topological dimension of the unstable manifold $W^u(p)$, $u(p) \stackrel{\text{def}}{=} \dim W^u(p)$. The *Kronecker-Poincaré index* of p is $\text{ind}(p, f) \stackrel{\text{def}}{=} (-1)^{u(p)}$.

Denote by S^3 the 3-sphere. The following lemma is of key importance for the proof of Theorem 1.

Lemma 1. *There is an embedding $\mathcal{S} \subset S^3$ and there is an extension of f_0 to a polar Morse-Smale diffeomorphism $f : S^3 \rightarrow S^3$ such that the non-wandering set $NW(f)$ is the union of a source, a sink and the fixed points of f_0 .*

Proof. Glue the balls B_{int}, B_{ext} to the respective boundary components S_{int}, S_{ext} of the fat sphere \mathcal{S} . Then we get the closed manifold $S^3 = \mathcal{S} \cup B_{int} \cup B_{ext}$ diffeomorphic to the 3-sphere and we get the natural embedding $\mathcal{S} \subset S^3$. From the properties of the diffeomorphism $f_0 : \mathcal{S} \rightarrow f_0(\mathcal{S})$ it follows that the 2-sphere S_{int} maps inside the fat sphere \mathcal{S} . Therefore f_0 can be extended to B_{int} in such a way that a hyperbolic source appears in the interior of B_{int} . Analogously f_0 can be extended to the ball B_{ext} so that a hyperbolic sink appears in the interior of B_{ext} . Denote this extension of f_0 by $f : S^3 \rightarrow S^3$. Clearly f_0 can be extended in such a way that f would be a diffeomorphism whose non-wandering set is the union of the non-wandering set of the diffeomorphism f_0 and the two new fixed points.

Thus, the non-wandering set of the diffeomorphism f is finite and it consists

of hyperbolic fixed points. Since the separatrices of the fixed points intersect transversally by the assumption, the diffeomorphism f is Morse-Smale. Moreover, f is polar because it has only two node fixed points. \square

The following lemma is the base for the proof of Theorem 2. Recall that S^1 denotes the circle and M_p^2 denotes the closed orientable surface of genus p . Everywhere below we assume $p \geq 1$.

Lemma 2. *There is an embedding $\mathcal{M} \subset M_p^2 \times S^1$ and there is an extension of f_0 to a polar Morse-Smale diffeomorphism $f : M_p^2 \times S^1 \rightarrow M_p^2 \times S^1$ such that the non-wandering set $NW(f)$ is the union of a source, a sink and the non-wandering points of the diffeomorphism f_0 .*

Proof. Glue balls B_1, B_2 to the respective boundary components S_1, S_2 of the fat surface \mathcal{M} . Then we get the body $\mathcal{M} \cup B_1 \cup B_2$ and the natural embedding $\mathcal{M} \subset \mathcal{M} \cup B_1 \cup B_2$. By the condition (d) the 2-sphere S_1 maps inside \mathcal{M} . Therefore f_0 can be extended to the ball B_1 in such a way that a hyperbolic source appears inside B_1 . Analogously f_0 can be extended to the ball B_2 so that a hyperbolic sink appears inside B_2 . The body $\mathcal{M} \cup B_1 \cup B_2$ is homeomorphic to the direct product $M_p^2 \times [0; 1]$ with the boundary components $T_1 = M_p^2 \times \{0\}$, $T_2 = M_p^2 \times \{1\}$, therefore below we identify $\mathcal{M} \cup B_1 \cup B_2$ and $M_p^2 \times [0; 1]$. It is known that from homotopy triviality of the restrictions $f_0|_{T_1}$, $f_0|_{T_2}$ it follows that each of these restrictions is isotopic to the identity map. Hence, there is $\varepsilon > 0$ and there is an extension of f_0 to the body $M_p^2 \times [-\varepsilon; 1 + \varepsilon]$ such that the restrictions $f_0|_{M_p^2 \times \{-\varepsilon\}}$ and $f_0|_{M_p^2 \times \{1 + \varepsilon\}}$ are the identities on the first coordinate. Without loss of generality one can assume each of $f_0|_{M_p^2 \times \{-\varepsilon\}}$, $f_0|_{M_p^2 \times \{1 + \varepsilon\}}$ to be a shift along the second coordinate $\{\cdot\} \times \{-\varepsilon\} \rightarrow \{\cdot\} \times \{-\varepsilon + \delta\}$ and $\{\cdot\} \times \{1 + \varepsilon\} \rightarrow \{\cdot\} \times \{1 + \varepsilon + \delta\}$ respectively.

Having glued the boundary components $M_p^2 \times \{-\varepsilon\}$, $M_p^2 \times \{1 + \varepsilon\}$ by the identity map we get the manifold $M_p^2 \times S^1$ from $M_p^2 \times [-\varepsilon; 1 + \varepsilon]$. It follows from the previous construction that there is the desired extension of the diffeomorphism f_0 to $M_p^2 \times S^1$. \square

The result below follows from the finiteness of every heteroclinic chain, so called ‘‘beh’’ (see also [11], details of the proof we leave to the reader).

Lemma 3. *Let M^3 be a closed orientable 3-manifold and let $f : M^3 \rightarrow M^3$ be a polar Morse-Smale diffeomorphism with a nonempty set of saddle periodic points of Morse indexes 1 and 2. Let α be the source and ω be the sink of f . Then there are saddle periodic points σ_1, σ_2 of Morse indexes $u(\sigma_1) = 1$ and $u(\sigma_2) = 2$ respectively such that the following inclusions hold*

$$W^u(\sigma_1) \setminus \{\sigma_1\} \subset W^s(\omega), \quad W^s(\sigma_2) \setminus \{\sigma_2\} \subset W^u(\alpha).$$

Lemma 4. *Let the assumptions of Lemma 3 be satisfied. Then the sets*

$$C_\omega \stackrel{\text{def}}{=} \{\omega\} \cup W^u(\sigma_1) \text{ and } C_\alpha \stackrel{\text{def}}{=} \{\alpha\} \cup W^s(\sigma_2)$$

are embeddings of the circles.

Proof. We prove the lemma only for C_ω because for C_α the proof is analogous. There is a C^1 -immersion $\varphi : \mathbb{R} \rightarrow W^u(\sigma_1)$, where $\varphi(0) = \sigma_1$, which is a one-to-one map to its image. We now show that the immersion φ extends to a homeomorphism

$$\varphi : S^1 \cong \mathbb{R} \cup \{\infty\} \rightarrow W^u(\sigma_1) \cup \{\sigma_1\},$$

if we set $\varphi(\pm\infty) = \omega$. Since $W^u(\sigma_1) \setminus \sigma_1 \subset W^s(\omega)$ the desired assertion follows from the fact that the ω -limit set of every point from the set $W^u(\sigma_1) \setminus \sigma_1$ is the point ω , see figure 3. \square




Figure 3: The curves C_α and C_ω .

Let M^3 be a smooth orientable closed 3-manifold. A 2-sphere S^2 in M^3 is said to be *cylindrical* or *cylindrically embedded* into M^3 , if there is a topological embedding $h : \mathbb{S}^2 \times [-1, 1] \rightarrow M^3$ such that $h(\mathbb{S}^2 \times \{0\}) = S^2$. A manifold M^3 is said to be *irreducible*, if every 2-sphere, cylindrically embedded into M^3 , bounds a 3-ball in M^3 . The manifolds S^3 and $M_p^2 \times S^1$ are known to be irreducible. Therefore, to prove the main results we have to answer the question if a polar Morse-Smale diffeomorphism on an irreducible orientable closed 3-manifold can have heteroclinic curves.

It follows from [12] that every Morse-Smale diffeomorphism on any irreducible 3-manifold except the sphere necessarily has heteroclinic curves (compact or noncompact). The following theorem generalizes this result for polar diffeomorphisms.

Theorem 3. *Let $f : M^3 \rightarrow M^3$ be a polar Morse-Smale diffeomorphism on an irreducible closed orientable 3-manifold M^3 and let f have at least one saddle point. Then there is a saddle periodic point such that its invariant 2-manifold contains non-compact heteroclinic curve.*

Proof. Denote the source, the sink and a saddle of the diffeomorphism f by α, ω and σ respectively. Without loss of generality the point $\sigma = \sigma_1$ can be assumed to satisfy the conditions of Lemma 4 (i.e. the manifold W_σ^s can be assumed to be of dimension 2, otherwise one can consider a suitable degree of f). We now show that W_σ^s contains a non-compact heteroclinic curve. Assume the

contrary, that either the manifold W_σ^s contains no heteroclinic curves or all these curves are compact. Then there is a simple smooth closed curve $\gamma \subset W_\sigma^s$ which is disjoint from the heteroclinic curves and which bounds a 2-disk $d_\gamma \subset W_\sigma^s$ such that the point σ lies in its interior. According to [10], $M^3 = \bigcup_{p \in \Omega_f} W_p^u = \bigcup_{p \in \Omega_f} W_p^s$

and therefore $\gamma \subset W_\alpha^u$. By assumption $cl W_\sigma^u = W_\alpha^u \cup \omega$ and $cl W_\sigma^u$ is a topologically embedded circle while W_σ^u is a smooth curve. Now making use of the disk d_γ we are going to prove that in M^3 there is a cylindrically embedded 2-sphere S which contains d_γ and which transversally intersects the circle $cl W_\sigma^u$ at the unique point σ . This would contradict the fact that the intersection $cl W_\sigma^u \cap S$ must consist of even number of points because the sphere S bounds a 3-ball in the irreducible manifold M^3 .

Construction of the sphere S . Consider in W_α^u a smooth 3-ball \tilde{B} containing α . Since $\gamma \subset W_\alpha^u$ there is a natural k for which $f^{-k}(\gamma) \subset \text{int } \tilde{B}$. Set $\tilde{\gamma} = f^{-k}(\gamma)$ and $Q = f^{-k}(d_\gamma)$. Pick a natural k_0 such that the 3-ball $B_0 = f^{-k_0}(\tilde{B})$ is inside the ball \tilde{B} . Set $\gamma_0 = f^{-k_0}(\tilde{\gamma})$ and $\Sigma = f^{-k_0}(Q)$. Without loss of generality the 2-disk Σ can be assumed to be transversal to the 2-spheres $\tilde{S} = \partial \tilde{B}$ and $S_0 = \partial B_0$. By construction the smooth ball B_0 satisfies the following conditions:

(*) $B_0 \subset \tilde{B} \subset W_\alpha^u$, $\alpha \in B_0$, $B_0 \cap Q = \emptyset$, the 2-sphere S_0 is transversal to Σ .

There are two cases: 1) every curve in the set $\Sigma \cap S_0$ separates γ_0 and σ in Σ ; 2) there is a curve in $\Sigma \cap S_0$ which does not separate γ_0 and σ in Σ . We now construct the desired sphere S in each case.

1) If every curve in the set $\Sigma \cap S_0$ separates γ_0 and σ in Σ then by construction the number of such curves is odd. Denote them by $\{c_1, \dots, c_m\}$. Up to reenumeration of c_i one can assume that $\Sigma \setminus (c_1 \cup \dots \cup c_m)$ consists of finitely many connected components $\Delta_1, \Delta_2, \dots, \Delta_{k+1}$ such that Δ_1 is homeomorphic to the 2-disk bounded by c_1 and such that $\sigma \in \Delta_1$; for $2 \leq i \leq k$ the component Δ_i is homeomorphic to the 2-annulus bounded by c_{i-1} and c_i ; the annulus Δ_{k+1} is bounded by the curves c_k and γ_0 . Let $d_1 \subset S_0$ be the 2-disk bounded by the curve c_1 . Then $S = \Delta_1 \cup d_1$ is the desired sphere.

2) Let there be a curve in $\Sigma \cap S_0$ which does not separate γ_0 and σ in Σ . Denote by \mathcal{C}_{B_0} the set of the curves in the intersection $\Sigma \cap S_0$ and denote by $\mathcal{C}_{B_0}^+$ ($\mathcal{C}_{B_0}^-$) the curves from \mathcal{C}_{B_0} which separate (do not separate) γ_0 and σ in Σ . Construct a ball B_1 such that it satisfies (*) and such that the number of the curves in $\mathcal{C}_{B_1}^-$ is less than the number of the curves in the set $\mathcal{C}_{B_0}^-$.

Every curve of the set $\mathcal{C}_{B_0}^-$ bounds on Σ a disk which contains neither γ_0 nor σ . Let $c \in \mathcal{C}_{B_0}^-$ be the innermost curve, i.e. c bounds on Σ the disk d_c that contains neither γ_0 nor σ and besides $\text{int } d_c$ contains no curves of the set $\mathcal{C}_{B_0}^-$. Since $c \cap Q = \emptyset$ and $\sigma \in Q$ we have $d_c \cap Q = \emptyset$ and therefore $d_c \subset \tilde{B}$. On the other hand c bounds on S_0 2-disks δ' and δ'' . Notice that $S' = d_c \cup \delta'$ and $S'' = d_c \cup \delta''$ are tamely embedded 2-spheres lying inside \tilde{B} . Denote by B' and B'' the 3-balls which are contained in \tilde{B} and which are bounded by S' and S''




Figure 4: Illustration to the proof of Theorem 3

respectively (see figure 4). Since $S' \cap Q = \emptyset$, $S'' \cap Q = \emptyset$ and $y_0 \in Q$ we have $B' \cap Q = \emptyset$, $B'' \cap Q = \emptyset$. And since $(B' \cup B'') \supset B_0$ we have $\alpha \in B'$ or $\alpha \in B''$.

To be definite let $\alpha \in B'$. Then we can pick a smooth ball B_1 which is arbitrarily close to B' , which satisfies (*) and such that $\mathcal{C}_{B_1}^- = \mathcal{C}_{B_0}^- \setminus c$. We continue this process and get a ball B which satisfies (*) and for which $\mathcal{C}_B^- = \emptyset$. \square

Corollary 3. *The invariant 2-manifold of every saddle point of a polar gradient-like diffeomorphism on an irreducible manifold contains a non-compact heteroclinic curve.*

Proof. The assertion is immediate from the proof of Theorem 3 and the fact that every one-dimensional separatrix of a gradient-like diffeomorphism is disjoint from the 2-dimensional separatrices. \square

Proof of Theorem 1. Taking into account Lemma 1, to prove that the number of null points is even it is enough to show that a Morse-Smale diffeomorphism $f : M^3 \rightarrow M^3$ necessarily has even number of periodic points on the closed 3-manifold M^3 . Since f preserves orientation of the ambient manifold by condition (a), we can assume it to preserve orientation of all separatrices of the

saddle fixed points (otherwise we consider a suitable iteration of f). Such polar diffeomorphism of the 3-sphere has the Lefschetz number equal to zero. From this and from the Lefschetz formula it follows that the number of the saddle fixed points of Morse index 1 equals the number of the saddle fixed points of Morse index 2. Hence, f has even number of the periodic points. Notice that we have also proved that f necessarily have saddle points of distinct Morse index. Now Theorem 1 follows from Theorem 3. \square

Proof of Theorem 2. First we show that f has at least one saddle point. Suppose the contrary, then f is a diffeomorphism of “source-sink” type. According to [13] the ambient manifold is then homeomorphic to the 3-sphere. This contradiction proves that f has at least one saddle point.

Since f preserves orientation of the ambient manifold by condition (a), we can assume it to preserve orientation of all separatrices of the saddle fixed points (otherwise we consider a suitable iteration of f). According to the condition (d) for $f_0 : M_p^2 \times S^1 \rightarrow f_0(M_p^2 \times S^1)$ the diffeomorphism f is homotopy trivial (i.e. it induces the identity map of the fundamental group of the torus). Therefore, its Lefschetz number equals to zero. The sum of the Morse indexes of the source and of the sink equals to zero, $(-1)^0 + (-1)^3 = 0$. From this and from the Lefschetz formula it follows that existence of one saddle fixed point courses existence of another fixed point with different Morse index. Now Theorem 2 follows from Theorem 3. \square

Corollaries 1, 2 follow from the proofs of Theorems 1, 2 and Corollary 3.

Acknowledgments

This work was supported by the Russian Foundation for Basic Research, Grants 12-01-00672-a and 13-01-12452-ofi-m, and partly by the Russian Science Foundation, Grant 14-11-00446. We especially thank Konstantin Kirsenko for financial support. We also thank the unknown reviewer for his(her) useful suggestions.

References

- [1] H. Alfvén, C. Fälthammar, *Cosmical Electrodynamics: Fundamental Principles*, Oxford, 1963.
- [2] T. G. Cowling, *Magnetohydrodynamics*, Interscience Publ. Inc., New York, 1957.
- [3] L. D. Landau, E. M. Lifshitz, *Course of theoretical physics, Vol. 8. Electrodynamics of Continuous Media*, Butterworth-Heinemann, 1984.
- [4] H. Alfvén, On sunspots and the solar cycle, *Ark. for Astronomi* 29 (1943) 1–17.
- [5] E. R. Priest, *Solar Magneto-Hydrodynamics*, D. Reidel, 1982.

- [6] E. Priest, T. Forbes, Magnetic reconnection: MHD theory and applications, Cambridge University Press: Cambridge, 2000.
- [7] Z. Nitecki, Differentiable dynamics: an introduction to the orbit structure of diffeomorphisms, MIT press Cambridge, 1971.
- [8] V. Z. Grines, O. V. Pochinka, Introduction to Topological Classification of Diffeomorphisms on 2- and 3-Manifolds (Russian), *Regulyarnaya i Chaoticheskaya Dinamika*, 2011.
- [9] D. Anosov, V. Arnold, Encyclopaedia of mathematical sciences: Dynamical systems I, Springer, Berlin, 1988.
- [10] S. Smale, Differentiable dynamical systems, *Bull. of the American Math. Soc.* 73 (6) (1967) 747–817.
- [11] C. Bonatti, V. Grines, Knots as topological invariants for gradient-like diffeomorphisms of the sphere S^3 , *J. of Dyn. and Control Syst.* 6 (4) (2000) 579–602.
- [12] C. Bonatti, V. Grines, V. Medvedev, E. Pécou, Three-manifolds admitting Morse–Smale diffeomorphisms without heteroclinic curves, *Topology and its Appl.* 117 (3) (2002) 335–344.
- [13] V. Z. Grines, E. V. Zhuzhoma, V. S. Medvedev, O. V. Pochinka, Global attractor and repeller of Morse-Smale diffeomorphisms, *Proc. of the Steklov Inst. of Math.* 271 (1) (2010) 103–124.