Bases of the Galois Ring $GR(p^r, m)$ over the Integer Ring \mathbb{Z}_{p^r}

Virgilio Sison*
Institute of Mathematical Sciences and Physics
University of the Philippines Los Baños
College, Laguna 4031, Philippines
Email: vpsison@uplb.edu.ph

August 6, 2018

Abstract

The Galois ring $GR(p^r, m)$ of characteristic p^r and cardinality p^{rm} , where p is a prime and $r, m \geq 1$ are integers, is a Galois extension of the residue class ring \mathbb{Z}_{p^r} by a root ω of a monic basic irreducible polynomial of degree m over \mathbb{Z}_{p^r} . Every element of $GR(p^r, m)$ can be expressed uniquely as a polynomial in ω with coefficients in \mathbb{Z}_{p^r} and degree less than or equal to m-1, thus $GR(p^r, m)$ is a free module of rank m over \mathbb{Z}_{p^r} with basis $\{1, \omega, \omega^2, \ldots, \omega^{m-1}\}$. The ring \mathbb{Z}_{p^r} satisfies the invariant dimension property, hence any other basis of $GR(p^r, m)$, if it exists, will have cardinality m.

This paper was motivated by the code-theoretic problem of finding the homogeneous bound on the p^r -image of a linear block code over $GR(p^r, m)$ with respect to any basis. It would be interesting to consider the dual and normal bases of $GR(p^r, m)$.

By using a Vandermonde matrix over $GR(p^r, m)$ in terms of the generalized Frobenius automorphism, a constructive proof that every basis of $GR(p^r, m)$ has a unique dual basis is given. The notion of normal bases was also generalized from the classic case for Galois fields.

Keywords – Galois rings, trace function, Frobenius automorphism, Vandermonde matrix, dual basis, normal basis

1 Introduction

It was proved in [1] that every basis of the Galois ring GR(4, m) has a dual basis, by treating each linear transformation from GR(4, m) to \mathbb{Z}_4 as being uniquely determined in terms of the generalized trace function on GR(4, m). In this paper this

^{*}This author gratefully acknowledges financial grant from the UPLB Diamond Jubilee-Development Fund Professorial Chair Award.

result is generalized to the Galois ring $GR(p^r, m)$ following the alternate method of MacWilliams and Sloane [8] which constructs the dual basis using simple matrix algebra involving the generalized Frobenius automorphism. The material is organized as follows: Section 2 gives the preliminaries and basic definitions, while Section 3 gives the results.

2 Preliminaries and definitions

An overview of Galois fields and Galois rings, the Frobenius automorphism and the trace function, is presented in this section. For a thorough discussion of these topics, we refer the reader to [7], [8] and [11].

2.1 Galois fields and Galois rings

Let p be a prime number and $r \geq 1$ an integer. Consider the ring \mathbb{Z}_{p^r} of integers modulo p^r . When r = 1 the ring \mathbb{Z}_p with p elements is a field and is usually denoted by \mathbb{F}_p . Let $\mathbb{Z}_{p^r}[x]$ be the ring of polynomials in the indeterminate x with coefficients in \mathbb{Z}_{p^r} .

The Galois field with p^m elements, denoted \mathbb{F}_{p^m} , is a field extension $\mathbb{F}_p[\alpha]$ of \mathbb{F}_p by a root α of an irreducible polynomial $\pi(x)$ of degree m in $\mathbb{F}_p[x]$. Thus every element z of \mathbb{F}_{p^m} can be expressed uniquely as a polynomial in α of the form

$$z = a_0 + a_1 \alpha + a_2 \alpha^2 + \ldots + a_{m-1} \alpha^{m-1}$$
 (1)

with degree at most m-1 with the coefficients a_i coming from \mathbb{F}_p , and hence can also be written as an m-tuple $(a_0, a_1, \ldots, a_{m-1})$ in \mathbb{F}_p^m . Elements of \mathbb{F}_{p^m} may also be described as residue classes of the polynomials in x with coefficients in \mathbb{F}_p reduced modulo $\pi(x)$. When m=1 we again have the field \mathbb{F}_p .

The canonical projection homomorphism $\mu: \mathbb{Z}_{p^r} \to \mathbb{F}_p$ is the mod-p reduction map, and can be extended naturally as a map from $\mathbb{Z}_{p^r}[x]$ onto $\mathbb{F}_p[x]$. This extended map is a ring homomorphism with kernel $(p) = \mathbb{Z}_{p^r}[x]p = \{f(x)p \mid f(x) \in \mathbb{Z}_{p^r}[x]\}.$

Let g(x) be a monic polynomial of degree $m \geq 1$ in $\mathbb{Z}_{p^r}[x]$. If $\mu(g(x))$ is irreducible in $\mathbb{F}_p[x]$, then g(x) is said to be *monic basic irreducible*. If $\mu(g(x))$ is primitive in $\mathbb{F}_p[x]$, then g(x) is said to be *monic basic primitive*. Clearly, monic basic primitive polynomials in $\mathbb{Z}_{p^r}[x]$ are monic basic irreducible.

In the general sense, a Galois ring is a finite commutative local ring with identity $1 \neq 0$ such that the set of zero divisors together with the zero element forms the unique maximal principal ideal (p1) for some prime number p. The residue class ring $\mathbb{Z}_{p^r}[x]/(h(x))$, where h(x) is a monic basic irreducible polynomial of degree m in $\mathbb{Z}_{p^r}[x]$, is a Galois ring with characteristic p^r and cardinality p^{rm} . The elements of $\mathbb{Z}_{p^r}[x]/(h(x))$ are residue classes of the form

$$a_0 + a_1 x + \ldots + a_{m-1} x^{m-1} + (h(x))$$
 (2)

where $a_i \in \mathbb{Z}_{p^r}$. The identity is 1 + (h(x)) and the zero element is (h(x)). The principal ideal (p[1 + (h(x))]) = (p + (h(x))) consists of all the zero divisors and the zero element, and is the only maximal ideal.

If deg h(x) = 1 then $\mathbb{Z}_{p^r}[x]/(h(x))$ is the ring \mathbb{Z}_{p^r} . If r = 1, the canonical homomorphism μ becomes the identity map and $\mathbb{Z}_{p^r}[x]/(h(x)) = \mathbb{F}_p[x]/(h(x)) \cong \mathbb{F}_{p^m}$. Now let $\omega = x + (h(x))$, then $h(\omega) = 0$ and every element z of $\mathbb{Z}_{p^r}[x]/(h(x))$ can be expressed uniquely in the form

$$z = a_0 + a_1 \omega + \ldots + a_{m-1} \omega^{m-1}$$
 (3)

where $a_i \in \mathbb{Z}_{p^r}$. We can thus think of $\mathbb{Z}_{p^r}[x]/(h(x))$ as a Galois extension $\mathbb{Z}_{p^r}[\omega]$ of \mathbb{Z}_{p^r} by ω . The elements take the *additive representation* (3), a generalization of (1) for \mathbb{F}_{p^m} . Since any two Galois rings of the same characteristic and the same cardinality are isomorphic, we simply use the notation $GR(p^r, m)$ for any Galois ring with characteristic p^r and cardinality p^{rm} .

The Galois ring $\mathcal{R} = GR(p^r, m)$ is a finite chain ring of length r, its ideals $p^i\mathcal{R}$ with $p^{(r-i)m}$ elements are linearly ordered by inclusion,

$$\{0\} = p^r \mathcal{R} \subset p^{r-1} \mathcal{R} \subset \ldots \subset p \mathcal{R} \subset \mathcal{R}$$

$$\tag{4}$$

The quotient ring $\mathcal{R}/p\mathcal{R} \cong \mathbb{F}_{p^m}$ is the residue field of \mathcal{R} . There exists a nonzero element ξ of order p^m-1 , which is a root of a unique monic basic primitive polynomial h(x) of degree m over \mathbb{Z}_{p^r} and dividing $x^{p^m-1}-1$ in $\mathbb{Z}_{p^r}[x]$. Consider the set

$$\mathcal{T} = \{0, 1, \xi, \xi^2, \dots, \xi^{p^m - 2}\}$$
 (5)

of Tëichmuller representatives. In this case, every element z of $GR(p^r, m)$ has a unique mutiplicative or p-adic representation as follows

$$z = z_0 + pz_1 + p^2 z_2 + \ldots + p^{r-1} z_{r-1}$$
(6)

where $z_i \in \mathcal{T}$. We have that z is a unit if and only if $z_0 \neq 0$, and z is a zero divisor or 0 if and only if $z_0 = 0$. The units form a multiplicative group of order $(p^m - 1)p^{(r-1)m}$, which is a direct product $\langle \omega \rangle \times \mathcal{E}$, where $\langle \omega \rangle$ is the cyclic group of order $p^m - 1$ that is isomorphic to \mathbb{Z}_{p^m-1} and $\mathcal{E} = \{1 + \pi \mid \pi \in (p)\}$ is a group of order $p^{(r-1)m}$. Let $\mu(\xi) = \alpha$. It can be shown that α is a primitive element in \mathbb{F}_{p^m} , and thus $\mu(\mathcal{T}) = \mathbb{F}_{p^m}$. The p-adic representation in (6) is a generalization of the power representation of an element of \mathbb{F}_{p^m} .

We realize that $GR(p^r, m)$ is a free module of rank m over \mathbb{Z}_{p^r} with the set

$$\mathcal{P}_m(\omega) = \{1, \omega, \omega^2, \dots, \omega^{m-1}\}$$
 (7)

as a free basis. The set $\mathcal{P}_m(\omega)$ is called the *standard* or *polynomial basis* of $GR(p^r, m)$. The ring \mathbb{Z}_{p^r} satisfies the invariant dimension property, hence any other basis of $GR(p^r, m)$, if it exists, will have cardinality m.

2.2 Generalized Frobenius automorphism and trace

The Generalized Frobenius map f on the Galois ring $\mathcal{R} = GR(p^r, m)$ is defined by

$$z^{f} := z_{0}^{p} + pz_{1}^{p} + p^{2}z_{2}^{p} + \ldots + p^{r-1}z_{r-1}^{p}$$
(8)

where z has the p-adic representation given in (6). The map f satisfies the following properties.

- (i) f is a ring automorphism of \mathcal{R} .
- (ii) f fixes every element of \mathbb{Z}_{p^r} .
- (iii) f is of order m and generates the cyclic Galois group of \mathcal{R} over \mathbb{Z}_{p^r} .

When r=1, the automorphism f reduces to the usual Frobenius automorphism σ of \mathbb{F}_{p^m} defined by $\sigma(z)=z^p$.

The generalized trace map T from \mathcal{R} down to \mathbb{Z}_{p^r} is given by

$$T(z) := z + z^f + z^{f^2} + \ldots + z^{f^{m-1}}$$
(9)

and satisfies the following properties.

- (i) T is surjective and $\mathcal{R}/\ker T \cong \mathbb{Z}_{p^r}$.
- (ii) T takes on each value of \mathbb{Z}_{p^r} equally often $p^{r(m-1)}$ times.
- (iii) $T(\alpha + \beta) = T(\alpha) + T(\beta)$ for all $\alpha, \beta \in \mathcal{R}$.
- (iv) $T(\lambda \alpha) = \lambda T(\alpha)$ for all $\lambda \in \mathbb{Z}_{p^r}, \alpha \in \mathcal{R}$.
- (v) $T(\alpha^f) = (T(\alpha))^f = T(\alpha)$ for all $\alpha \in \mathcal{R}$.

Again when r=1 the generalized trace map T reduces to the trace map $t: \mathbb{F}_{p^m} \to \mathbb{F}_p$ defined by

$$t(\beta) = \beta + \beta^p + \beta^{p^2} + \ldots + \beta^{p^{m-1}}.$$
 (10)

2.3 Homogeneous weight on $GR(p^r, m)$

Let R be a finite ring with identity $1 \neq 0$, and \mathbb{T} be the multiplicative group of unit complex numbers. The group \mathbb{T} is a one-dimensional torus. A character of R (considered as an additive abelian group) is a group homomorphism $\chi: R \to \mathbb{T}$. The set of all characters \widehat{R} (called the character module of R) is a right (resp. left) R-module whose group operation is pointwise multiplication of characters and scalar multiplication is given by $\chi^r(x) = \chi(rx)$ (resp. ${}^r\chi(x) = \chi(xr)$). A character χ of R is called a right (resp. left) generating character if the mapping $\phi: R \to \widehat{R}$ given by $\phi(r) = \chi^r$ (resp. $\phi(r) = {}^r\chi$) is an isomorphism of right (resp. left) R-modules. The ring R is called Frobenius if and only if R admits a right or a left generating character, or alternatively, if and only if R as right generating character if and only if it is a left generating character. Further χ is a right generating character if and only if R is a left generating character. Further R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if and only if R is a right generating character if R is a right generating R is a ri

Let \mathbb{R} be the set of real numbers. We define a homogeneous weight on an arbitrary finite ring R with identity in the sense of [4]. Let Rx denote the principal (left) ideal generated by $x \in R$.

Definition 2.1 A weight function $w : R \to \mathbb{R}$ on a finite ring R is called (left) homogeneous if w(0) = 0 and the following is true.

- (i) If Rx = Ry, then w(x) = w(y) for all $x, y \in R$.
- (ii) There exists a real number $\Gamma > 0$ such that

$$\sum_{y \in Rx} w(y) = \Gamma \cdot |Rx|, \text{ for all } x \in R \setminus \{0\}.$$
 (11)

Right homogeneous weights are defined accordingly. If a weight is both left homogeneous and right homogeneous, we call it simply as a homogeneous weight. The constant Γ in (11) is called the average value of w. A homogeneous weight is said to be normalized if its average value is 1. We can normalize the weight w in Definition 2.1 by replacing it with $\widetilde{w} = \Gamma^{-1}w$ [6]. The weight w is extended naturally to R^n , the free module of rank n consisting of n-tuples of elements from R, via $w(z) = \sum_{i=0}^{n-1} w(z_i)$ for $z = (z_0, z_1, \ldots, z_{n-1}) \in R^n$. The homogeneous distance metric $\delta: R^n \times R^n \longrightarrow \mathbb{R}$ is defined by $\delta(x, y) = w(x - y)$, for $x, y \in R^n$.

It was proved in [5] that, if R is Frobenius with generating character χ , then every homogeneous weight w on R can be expressed in terms of χ as follows.

$$w(x) = \Gamma \left[1 - \frac{1}{|R^{\times}|} \sum_{u \in R^{\times}} \chi(xu) \right]$$
 (12)

where R^{\times} is the group of units of R.

For the Galois ring $GR(p^r, m)$ we apply the following homogeneous weight given in [3] for finite chain rings.

$$w_{\text{hom}}(x) = \begin{cases} 0 & \text{if } x = 0\\ p^{m(r-1)} & \text{if } x \in (p^{r-1}) \setminus \{0\}\\ (p^m - 1)p^{m(r-2)} & \text{otherwise} \end{cases}$$
 (13)

where (p^{r-1}) is the principal ideal generated by the element p^{r-1} of $GR(p^r, m)$. Since the Galois ring $GR(p^r, m)$ is a commutative Frobenius ring with identity whose generating character is $\chi(z) = \xi^{b_{m-1}}$, where $\xi = \exp(2\pi i/p^r)$ for $z = \sum_{i=0}^{m-1} b_i \omega^i$, the weight (13) can be derived from (12). The group of units of $GR(p^r, m)$ has cardinality $p^{m(r-1)}(p^m-1)$ and it easy to compute from (11) that its average value is equal to

$$\Gamma = (p^m - 1)p^{m(r-2)} \tag{14}$$

which is its minimum non-zero value. When r=1, we have $\Gamma=(p^m-1)/p^m$ and w_{hom} is just the usual Hamming weight w_{Ham} on \mathbb{F}_{p^m} . When m=1, the average value is $\Gamma=(p-1)p^{r-2}$ for the integer ring \mathbb{Z}_{p^r} .

2.4 Codes over $GR(p^r, m)$

A block code C of length n over an arbitrary finite ring R is a nonempty subset of R^n . The code C is called right (resp. left) R-linear if C is a right (resp. left) R-submodule of R^n . If C is both left R-linear and right R-linear, we simply call C a linear block code over R. A $k \times n$ matrix over R is called a generator matrix of a

linear block code C if the rows span C and no proper subset of the rows generates C.

Let the set $\mathcal{B}_m = \{\beta_0, \beta_1, \dots, \beta_{m-1}\}$ be a basis of the Galois ring \mathcal{R} over \mathbb{Z}_{p^r} , and B be a linear block code of length n over \mathcal{R} . We consider the map $\tau : \mathcal{R} \longrightarrow \mathbb{Z}_{p^r}^m$ defined by

$$\tau(z) = (a_0, a_1, \dots, a_{m-1}) \tag{15}$$

for $z = a_0\beta_0 + a_1\beta_1 + \ldots + a_{m-1}\beta_{m-1} \in \mathcal{R}$, $a_i \in \mathbb{Z}_{p^r}$. This map is a bijection and can be extended coordinate-wise to \mathcal{R}^n . Thus, if $c \in B$ and $c = (c_0, c_1, \ldots, c_{n-1})$, $c_i = \sum_{j=0}^{m-1} a_{ij}\beta_j$, $a_{ij} \in \mathbb{Z}_{p^r}$, then

$$\tau(c) = (a_{00}, \dots, a_{0,m-1}, \dots, a_{n-1,0}, \dots a_{n-1,m-1})$$
(16)

in $\mathbb{Z}_{p^r}^{mn}$. The image $\tau(B)$ of B under τ with respect to \mathcal{B}_m is called the p^r -ary image of B, and is obtained by simply substituting each element of \mathcal{R} by the m-tuple of its coordinates over B. It is easy to prove that $\tau(B)$ is a linear block code of length mn over \mathbb{Z}_{p^r} . For the degenerate case m=1, the block code B is a code over \mathbb{Z}_{p^r} and the map τ is the identity map on B. We equip $\tau(B)$ with a homogeneous distance metric with respect to the weight w_{hom} as given in (13).

The following lemma from [2] will be very useful in the succeeding discussion.

Lemma 2.2 (Constantinescu, Heise and Honold, 1996) For any linear block code $C \subseteq \mathbb{Z}_{p^r}^n$ we have

$$\frac{w_{\text{hom}}(C)}{|C|} = \Gamma \cdot |\{i \mid \pi_i(C) \neq 0\}|$$

where $w_{\text{hom}}(C)$ is the sum of the homogeneous weights of all codewords of C, and π_i is the projection from $\mathbb{Z}_{p^r}^n$ onto the *i*-th coordinate.

3 Major Results

We denote by $w_{hom}(S)$ the sum of the homogeneous weights of the elements of set S, that is,

$$w_{\text{hom}}(S) = \sum_{x \in S} w_{\text{hom}}(x) . \tag{17}$$

Proposition 3.1 For any basis $\mathcal{B}_m = \{\beta_0, \beta_1, \dots, \beta_{m-1}\}\ of\ GR(p^r, m)\ over\ \mathbb{Z}_{p^r}\ we$ have

$$\sum_{x \in GR(p^r, m)} w_{\text{hom}}(\tau(x)) = m(p-1)p^{rm+r-2}.$$
 (18)

Proof: Let $S = \{x \mid x \in GR(p^r, m)\}$. Then $\tau(S)$ is a linear block code over \mathbb{Z}_{p^r} of length m and cardinality p^{rm} . Applying Lemma 2.2 to $\tau(S)$ gives us

$$\frac{w_{\text{hom}}(\tau(S))}{|\tau(S)|} = \Gamma \cdot w_s(\tau(S)) .$$

Therefore we have $w_{\text{hom}}(\tau(S)) = |\tau(S)| \cdot \Gamma \cdot w_s(S)$. The value of Γ is given in (14), and the support size $w_s(\tau(S))$ of $\tau(S)$ is m. Using the notation in (17), the result now follows.

This proposition gives the simple corollary below which was used to prove the bound of Rabizzoni in [9, Theorem 1].

Corollary 3.2 For any basis $\mathcal{B}_m = \{\beta_0, \beta_1, \dots, \beta_{m-1}\}$ of \mathbb{F}_{p^m} over \mathbb{F}_p we have

$$\sum_{x \in \mathbb{F}_{p^m}} w_{\mathtt{Ham}}(\tau(x)) = m(p-1)p^{m-1} \; .$$

Proof: The Galois ring GR(p,m) is the Galois field \mathbb{F}_{p^m} , and the homogeneous weight w_{hom} given in (13) is the Hamming weight w_{Ham} on \mathbb{F}_p with $\Gamma = (p-1)/p$.

The bound of Rabizzoni in [9, Theorem 1] was extended to linear block codes over Galois rings in [10].

Denote by $\operatorname{Mat}_m(\mathcal{R})$ the ring of $m \times m$ matrices over the Galois ring $\mathcal{R} = GR(p^r, m)$. It is known that a matrix A in $\operatorname{Mat}_m(\mathcal{R})$ is nonsingular (or invertible) if and only if $\det A$ is a unit in \mathcal{R} . We will also use the usual notation |A| for the determinant of A. The matrix A is symmetric if and only if $A = A^t$, and is orthogonal if and only if $AA^t = A^tA = I$, where A^t is the transpose of A and I is the identity matrix. We propose the following definition.

Definition 3.3 Two bases $\{\alpha_i\} = \{\alpha_1, \alpha_2, \dots, \alpha_m\}$ and $\{\beta_j\} = \{\beta_1, \beta_2, \dots, \beta_m\}$ of $GR(p^r, m)$ over \mathbb{Z}_{p^r} are said to be dual) if $T(\beta_i \alpha_j) = \delta_{ij}$, where δ_{ij} is the Kronecker delta.

Lemma 3.4 The matrix $\Omega \in Mat_m(\mathcal{R})$ given by

$$\Omega = \begin{pmatrix}
1 & 1 & 1 & \dots & 1 \\
\omega & \omega^f & \omega^{f^2} & \dots & \omega^{f^{m-1}} \\
\omega^2 & (\omega^2)^f & (\omega^2)^{f^2} & \dots & (\omega^2)^{f^{m-1}} \\
\vdots & & & \vdots \\
\omega^{m-1} & (\omega^{m-1})^f & (\omega^{m-1})^{f^2} & \dots & (\omega^{m-1})^{f^{m-1}}
\end{pmatrix}$$

is nonsingular.

Proof: By the definition of the generalized Frobenius automorphism (8), it easy to show that $(\omega^j)^{f^i} = (\omega^{p^i})^j$ for $i, j = 0, 1, \dots, m-1$. Hence,

$$\Omega = \begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ \omega & \omega^{p} & \omega^{p^{2}} & \dots & \omega^{p^{m-1}} \\ \omega^{2} & (\omega^{p})^{2} & (\omega^{p^{2}})^{2} & \dots & (\omega^{p^{m-1}})^{2} \\ \vdots & & & \vdots \\ \omega^{m-1} & (\omega^{p})^{m-1} & (\omega^{p^{2}})^{m-1} & \dots & (\omega^{p^{m-1}})^{m-1} \end{pmatrix}$$

which is a Vandermonde matrix whose determinant is

$$\det \Omega = \prod_{i=1}^{m-1} \prod_{j=i+1}^{m} (\omega^{p^{i-1}} - \omega^{p^{j-1}})$$
(19)

Each factor in this product is a unit of \mathcal{R} so that det Ω is a unit in \mathcal{R} .

Lemma 3.5 Let $\{\beta_j\} = \{\beta_1, \beta_2, \dots, \beta_m\}$ be a basis. The matrix

$$B = \begin{pmatrix} \beta_1 & \beta_1^f & \beta_1^{f^2} & \dots & \beta_1^{f^{m-1}} \\ \beta_2 & \beta_2^f & \beta_2^{f^2} & \dots & \beta_2^{f^{m-1}} \\ \vdots & & & \vdots \\ \beta_m & \beta_m^f & \beta_m^{f^2} & \dots & \beta_m^{f^{m-1}} \end{pmatrix}$$
(20)

is invertible.

Proof: Express the polynomial basis $\mathcal{P}_m(\omega)$ in (7) in terms of the basis $\{\beta_j\}$ as follows.

$$\begin{pmatrix} 1 \\ \omega \\ \omega^{2} \\ \vdots \\ \omega^{m-1} \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ a_{31} & a_{32} & \dots & a_{3m} \\ \vdots & & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mm} \end{pmatrix} \begin{pmatrix} \beta_{1} \\ \beta_{2} \\ \beta_{3} \\ \vdots \\ \beta_{m} \end{pmatrix}$$

where $A = (a_{ij})$ is a nonsingular matrix over \mathbb{Z}_{p^r} . We compute the matrix product AB. The fact that the Frobenius automorphism f fixes each a_{ij} implies that AB is the Vandermonde matrix Ω . Hence by Lemma 3.4, det AB is a unit in \mathcal{R} . Consequently, det B is a unit in \mathcal{R} .

We shall call the matrix B the automorphism matrix of \mathcal{R} relative to the basis $\{\beta_i\}$.

Corollary 3.6 $(\det B)^2$ is a unit in \mathbb{Z}_{p^r} .

Proof: It can be shown that

$$BB^{t} = \begin{pmatrix} T(\beta_{1}^{2}) & T(\beta_{1}\beta_{2}) & \dots & T(\beta_{1}\beta_{m}) \\ T(\beta_{2}\beta_{1}) & T(\beta_{2}^{2}) & \dots & T(\beta_{2}\beta_{m}) \\ \vdots & & & \vdots \\ T(\beta_{m}\lambda_{1}) & T(\beta_{m}\lambda_{2}) & \dots & T(\beta_{m}^{2}) \end{pmatrix}$$

$$(21)$$

which is a matrix over \mathbb{Z}_{p^r} . It follows that $(\det B)^2$ is an element of \mathbb{Z}_{p^r} . By Lemma 3.5, we get the result.

Of course, det B is not necessarily a unit in the base ring \mathbb{Z}_{p^r} , although it is a unit in \mathcal{R} according to Lemma 3.5. Please see Example 3.8.

Theorem 3.7 Every basis has a unique dual basis.

Proof: We show the proof for m = 3 without loss of essential generality. Let $\{\beta_1, \beta_2, \beta_3\}$ be a basis, and consider the automorphism matrix

$$B = \begin{pmatrix} \beta_1 & \beta_1^f & \beta_1^{f^2} \\ \beta_2 & \beta_2^f & \beta_2^{f^2} \\ \beta_3 & \beta_3^f & \beta_3^{f^2} \end{pmatrix}$$

which is nonsingular by Lemma 3.5. Let adj B = (b_{ij}) where $b_{ij} = (-1)^{i+j} |B_{ji}|$. Then

$$\text{adj B} = \left(\begin{array}{ccc} \lambda_1 & \lambda_2 & \lambda_3 \\ \lambda_1^f & \lambda_2^f & \lambda_3^f \\ \lambda_1^{f^2} & \lambda_2^{f^2} & \lambda_3^{f^2} \end{array} \right)$$

where $\lambda_1 = \beta_2^f \beta_3^{f^2} - \beta_2^{f^2} \beta_3^f$, $\lambda_2 = \beta_1^{f^2} \beta_3^f - \beta_1^f \beta_3^{f^2}$, and $\lambda_3 = \beta_1^f \beta_2^{f^2} - \beta_1^{f^2} \beta_2^f$ so that $B^{-1} = |B|^{-1}$ adj B. Note that

$$BB^{-1} = |B|^{-1} \begin{pmatrix} T(\beta_1 \lambda_1) & T(\beta_1 \lambda_2) & T(\beta_1 \lambda_3) \\ T(\beta_2 \lambda_1) & T(\beta_2 \lambda_2) & T(\beta_2 \lambda_3) \\ T(\beta_3 \lambda_1) & T(\beta_3 \lambda_2) & T(\beta_3 \lambda_3) \end{pmatrix}$$

Following the argument of [1] it can be shown by using the generalized trace that $\{\lambda_1/|B|, \lambda_2/|B|, \lambda_3/|B|\}$ is a linearly independent set, and hence is the unique dual of $\{\beta_i\}$.

Example 3.8 The polynomial basis for GR(4,2) is the set $\{1,\omega\}$ where $1+\omega+\omega^2=0$. The automorphism matrix is

$$B = \left(\begin{array}{cc} 1 & 1\\ \omega & 3 + 3\omega \end{array}\right)$$

with determinant $3 + 2\omega$ which is a unit in GR(4,2). Observe that $(3 + 2\omega)^2 = 1$ is a unit in \mathbb{Z}_4 . The inverse

$$B^{-1} = \begin{pmatrix} 3+\omega & 1+2\omega \\ 2+3\omega & 3+2\omega \end{pmatrix}$$

gives $\{3 + \omega, 1 + 2\omega\}$ as the dual of the polynomial basis.

Example 3.9 The polynomial basis for GR(4,3) is the set $\{1,\omega,\omega^2\}$ where ω is the root of the basic primitive polynomial $x^3 + 2x^2 + x - 1$ over \mathbb{Z}_4 . The automorphism matrix is given by

$$B = \left(\begin{array}{ccc} 1 & 1 & 1\\ \omega & \omega^2 & \omega^4\\ \omega^2 & \omega^4 & \omega \end{array}\right)$$

with determinant 3. The inverse is given by

$$B^{-1} = \begin{pmatrix} \omega + 3\omega^3 & w + 3\omega^4 & w^2 + 3\omega^4 \\ \omega^2 + 3\omega^6 & 3\omega + \omega^2 & 3\omega + \omega^4 \\ \omega^4 + 3\omega^5 & 3\omega^2 + \omega^4 & \omega + 3\omega^2 \end{pmatrix}$$

so that $\{3 + 2\omega + 2\omega^2, 2 + 2\omega + \omega^2, 2 + \omega + 2\omega^2\}$ is the dual basis. This corrects the mistake in [1, Example 1].

Example 3.10 The dual of the polynomial basis for $\mathbb{Z}_8[\omega]$, where ω is the root of the basic primitive polynomial $7 + 5x + 6x^2 + x^3$ over \mathbb{Z}_8 , is the set $\{3 + 6\omega + 6\omega^2, 6 + 2\omega + 5\omega^2, 6 + 5\omega + 2\omega^2\}$.

We apply Definition 3.3 for the notion of self-dual basis.

Definition 3.11 The basis $\{\beta_1, \beta_2, \dots, \beta_m\}$ is self-dual if $T(\beta_i \beta_j) = \delta_{ij}$.

Definition 3.12 A normal basis of $GR(p^r, m)$ is a basis of the form $\{\alpha, \alpha^f, \alpha^{f^2}, \dots, \alpha^{f^{m-1}}\}$ where $\alpha \in GR(p^r, m)$ and f is the generalized Frobenius automorphism given in (8). In this case we say that α generates β_m .

We have the following immediate results.

Theorem 3.13 Let $\{\beta_j\}$ be a basis with automorphism matrix B. Then B is orthogonal if and only if $\{\beta_j\}$ is self-dual.

Proof: From (21) we get $BB^t = I \Leftrightarrow T(\beta_i \beta_j) = \delta_{ij}$.

Theorem 3.14 Let $\{\beta_j\}$ be a basis with automorphism matrix B. Then the following statements are equivalent.

- (i) The basis $\{\beta_i\}$ is a normal basis.
- (ii) The automorphism matrix B is a symmetric matrix.
- (iii) The Frobenius automorphism f is the m-cycle $\beta_1 \mapsto \beta_2, \beta_2 \mapsto \beta_3, \dots, \beta_m \mapsto \beta_1$.

Proof: This equivalence is evident from the construction of the automorphism matrix in (20). The basis $\mathcal{B}_m = \{\beta_j\}$ is normal $\Leftrightarrow \beta_1$ generates \mathcal{B}_m , that is, $\beta_2 = \beta_1^f, \beta_3 = \beta_2^f = \beta_1^{f^2}, \beta_3 = \beta_2^f = \beta_1^{f^3}, \dots, \beta_{m-1} = \beta_{m-2}^f = \beta_1^{f^{m-1}}, \beta_m = \beta_{m-1}^f = \beta_1^{f^{m-1}}, \beta_m^f = \beta_1^{f^m} = \beta_1 \Leftrightarrow B$ is symmetric $\Leftrightarrow f$ is the m-cycle $\beta_1 \mapsto \beta_2, \beta_2 \mapsto \beta_3, \dots, \beta_m \mapsto \beta_1$.

Example 3.15 The set $\mathcal{B}_2 = \{\omega, \omega^2 = 3 + 3\omega\}$ is a normal basis for $\mathbb{Z}_4[x]/(x^2 + x + 1)$. The automorphism matrix relative to this basis is given by

$$\left(\begin{array}{cc} \omega & 3+3\omega \\ 3+3\omega & \omega \end{array}\right)$$

which is not orthogonal, hence \mathcal{B}_2 is not self-dual. However B is symmetric.

Example 3.16 The set $\mathcal{B}_3 = \{1+\omega, 1+\omega^2, 3+3\omega+3\omega^2\}$ of $GR(4,3) = \mathbb{Z}_4[x]/(x^3+2x^2+x+3)$ is a self-dual normal basis as the automorphism matrix is both orthogonal and symmetric.

References

- [1] B. Abrahamsson, Architectures for Multiplications in Galois Rings, Ph.D Thesis, Linköpings Universitet, 2004.
- [2] I. Constantinescu, W. Heise, and T. Honold, "Monomial extensions of isometries between codes over \mathbb{Z}_M ," in Proceedings of the 5th International Workshop on Algebraic and Combinatorial Coding Theory (ACCT '96), Unicorn Shumen, pp. 98 104, 1996.

- [3] M. Greferath and S.E. Schmidt, "Gray isometries for finite chain rings and a nonlinear ternary (36, 3¹², 15) code," *IEEE Trans. Inform. Theory*, vol. 45, no. 7, pp. 2522–2524, November 2001.
- [4] M. Greferath and S.E. Schmidt, "Finite-ring combinatorics and MacWilliams Equivalence Theorem," *J. Combin Theory*, Series A, vol. 92, pp. 17–28, 2000.
- [5] T. Honold, "A characterization of finite Frobenius rings", Arch. Math. (Basel), vol. 76, pp. 406–415, 2001.
- [6] Th. Honold and A. Nechaev, "Weighted modules and representations of codes", *Probl. Inform. Transm.*, vol. 35, no. 3, pp. 205–222, 1999.
- [7] B.R. MacDonald, Finite Rings with Identity, Marcel Dekker, 1974.
- [8] F. J. MacWilliams and N. J. A. Sloane, *The Theory of Error-Correcting Codes*. Amsterdam, The Netherlands: North-Holland, 1977.
- [9] P. Rabizzoni, "Relation between the minimum weight of a linear code over $GF(q^m)$ and its q-ary image over GF(q)," Lecture Notes in Computer Science, vol. 388, Berlin, Germany: Springer-Verlag, 1989, pp. 209 212.
- [10] P. Solé and V. Sison, "Bounds on the minimum homogeneous distance of the p^r -ary image of linear block codes over the Galois ring $GR(p^r, m)$," *IEEE Trans. Inform. Theory*, vol. 53, no. 6, pp. 2270–2273, June 2007.
- [11] Z.X. Wan, Lectures on Finite Fields and Galois Rings, World Scientific, 2003.