# A Global Trace Formula for Reductive Lie Algebras and the Harish-Chandra Transform on the Space of Characteristic Polynomials

Shuyang Cheng

#### Abstract

In this paper an integral transform between spaces of nonstandard test functions on the affine space of dimension n is constructed. The integral transform satisfies a summation formula of Poisson type, which is derived from an analogue of the Arthur-Selberg trace formula for the Lie algebra of  $n \times n$  matrices.

### Introduction

The classical summation formula of Poisson

$$\sum_{n \in \mathbb{Z}} f(n) = \sum_{n \in \mathbb{Z}} \hat{f}(n)$$

has been related to the modularity of theta functions and the functional equation of zeta functions since the days of Jacobi and Riemann.

In his seminal paper [1859] Riemann derived the functional equation of the zeta function which now bears his name from the modular identity of Jacobi's theta function, which is a consequence of the Poisson summation formula.

In his thesis [Ta50] Tate recast Riemann's argument over the adeles and derived the functional equation of a large class of L-functions directly from his adelic Poisson summation formula. Tate's method was later generalized to the standard L-functions of higher rank general linear groups and central simple algebras by Godement and Jacquet in [GJ72]. In their argument the group GL(n) is embedded into its Lie algebra gl(n), and the functional equation of standard L-functions of GL(n) is derived from the adelic Poisson summation formula for the vector space gl(n).

Recently in the work of Braverman and Kazhdan [BK00], Ngô [Ng12], Sakellaridis [Sa12] and Lafforgue [La13], the argument of Godement and Jacquet was partially generalized to more general automorphic L-functions:

Let G be a reductive group and let  $\rho$  be a representation of the dual group <sup>L</sup>G satisfying some mild assumptions, then the group G embeds into an algebraic monoid  $M_{\rho}$  which depends on  $\rho$ , and the functional equation of the automorphic *L*-functions  $L(s, \pi, \rho)$  would follow from a conjectural summation formula of Poisson type for the monoid  $M_{\rho}$ . In general  $M_{\rho}$ is nonlinear and possibly singular, and the conjectural Poisson summation formula would involve a nonstandard Fourier transform and spaces of nonstandard test functions. The purpose of this paper is to construct a toy example of such a Poisson summation formula for the space  $\mathcal{A}_n$  of characteristic polynomials of  $n \times n$  matrices:

**Proposition (5.4.7)** There exists a dense subspace  $S_0(\mathcal{A}_n(\mathbb{A}))$  of  $L^2(\mathcal{A}_n(\mathbb{A}))$  of nonstandard Schwartz functions on  $\mathcal{A}_n(\mathbb{A})$  and an endomorphism  $\mathcal{H}_0$  of  $S_0(\mathcal{A}_n(\mathbb{A}))$  such that:

For each point X in  $\mathcal{A}_n(\mathbb{Q})$  there exists a nonzero constant a(X) such that for all  $\varphi$  in  $\mathcal{S}_0(\mathcal{A}_n(\mathbb{A}))$  which is cuspidal at two distinct place of  $\mathbb{Q}$ 

$$\sum_{X \in \mathcal{A}_n(\mathbb{Q})} a(X) \cdot \varphi(X) = \sum_{X \in \mathcal{A}_n(\mathbb{Q})} a(X) \cdot \mathcal{H}_0(\varphi)(X),$$

where the summand for a polynomial X with repeated roots needs to be interpreted appropriately.

The Poisson summation formula for  $\mathcal{A}_n$  is deduced from an analogue of the Arthur-Selberg trace formula for the Lie algebra  $\mathfrak{gl}(n)$ . In this paper such a trace formula is established for a restricted class of test functions subject to some mild local conditions on general reductive Lie algebra  $\mathfrak{g}$ . Some possible problems for future investigation include removing the restrictions on the test function  $\varphi$ , and establishing the Poisson summation formula for the affine space  $\mathcal{A}_{\mathrm{G}}$  defined as the adjoint quotient of  $\mathfrak{g}$  for a general reductive group G. Such a Poisson summation formula will likely require a stable trace formula for  $\mathfrak{g}$ .

### Acknowledgements

I wish to express my gratitude to my advisor Ngô Bao Châu, without whose continual support and encouragement this work would not have been possible. I wish to thank Pierre-Henri Chaudouard for spotting an error in an earlier version of this paper and providing very valuable suggestions. I also wish to thank Bill Casselman, Eddie Herman, Robert Kottwitz and Yiannis Sakellaridis for illuminating conversations. The proof of Corollary (5.4.9) is suggested by Sakellaridis.

## Contents

In	troduction	1				
Ac	cknowledgements	<b>2</b>				
1	Preliminaries         1.1       Notations and definitions	<b>4</b> 4				
2	The non-invariant trace formula2.1A motivating example2.2The non-invariant trace formula of Chaudouard	<b>10</b> 10 12				
3	Refined expansions         3.1       Weighted orbital integrals         3.2       Nilpotent orbits         3.3       General orbits         3.4       The refined trace formula	<ol> <li>18</li> <li>21</li> <li>30</li> <li>45</li> </ol>				
4	<ul> <li>A simple invariant trace formula</li> <li>4.1 Parabolic descent and parabolic induction</li></ul>	<b>46</b> 47 51				
5	The Harish-Chandra transform on the space         of characteristic polynomials         5.1       Preliminaries         5.2       The local and global Schwartz spaces         5.3       The Harish-Chandra transform         5.4       The Poisson summation formula	<b>52</b> 52 56 58 61				
A	A Appendix: Scissors congruence and orbital integrals					
в	Appendix: The example of $GL(2)$	77				
Lis	st of symbols	87				
Re	eferences	92				

### **1** Preliminaries

### **1.1** Notations and definitions

(1.1.1) Let G be a connected reductive group defined over the field of rational numbers  $\mathbb{Q}$ , let  $\mathfrak{g}$  be its Lie algebra equipped with the adjoint action of G from the right. More generally an algebraic group will be denoted by a capital roman letter, its Lie algebra by the corresponding lowercase fraktur letter, except for  $\mathfrak{a}$  which is reserved for a Euclidean vector space.

Denote by  $\mathbb{Q}_v$  the completion of  $\mathbb{Q}$  at a place v. If S is a finite set of places, denote by  $\mathbb{Q}_S$  the direct product of  $\mathbb{Q}_v$  for all v in S. Denote by  $\mathbb{A}$  the ring of adeles of  $\mathbb{Q}$ . Define S-local and global norms by

(1.1.1.1) 
$$\forall x \in \mathbb{Q}_S \quad |x|_S = \prod_{v \in S} |x|_v, \quad \forall x \in \mathbb{A} \quad |x|_{\mathbb{A}} = \lim_S |x|_S.$$

The local norms are normalized in such a way that

$$(1.1.1.2) \qquad \forall x \in \mathbb{Q} - \{0\} \qquad |x|_{\mathbb{A}} = 1.$$

The groups  $G(\mathbb{Q}_v)$ ,  $G(\mathbb{Q}_S)$  and  $G(\mathbb{A})$  of  $\mathbb{Q}_v$ ,  $\mathbb{Q}_S$  and  $\mathbb{A}$ -valued points of G are locally compact with respect to the analytic topology. The group  $G(\mathbb{Q})$  of  $\mathbb{Q}$ -valued points of G is a discrete subgroup of  $G(\mathbb{A})$  with respect to the analytic topology.

(1.1.2) Fix a minimal parabolic subgroup  $P_0$  of G. Fix a Levi subgroup  $M_0$  of  $P_0$  with split component  $A_0$ . A parabolic subgroup P of G is said to be *standard* if P contains  $P_0$ . Denote by  $N_P$  the unipotent radical of P, by  $M_P$  the unique Levi subgroup of P containing  $M_0$ , by  $A_P$  the split component of  $M_P$ . Such a Levi subgroup  $M_P$  is said to be *standard*. Denote by  $\overline{P}$  the parabolic subgroup group opposite to P, by  $\overline{N}_P$  its unipotent radical. To simplify notations the standard Levi, split and unipotent components of  $P_i$  will be denoted by  $M_i$ ,  $A_i$  and  $N_i$  where *i* is a natural number.

Let M and L be Levi subgroups of G such that M is contained in L, denote by  $\mathcal{F}^{L}(M)$ the set of parabolic subgroups of L that contain M, by  $\mathcal{P}^{L}(M)$  the set of parabolic subgroups of L whose Levi component is M, by  $\mathcal{L}^{L}(M)$  the set of Levi subgroups of L that contain M. To simplify notations denote by  $\mathcal{F}(M)$  the set  $\mathcal{F}^{G}(M)$ , by  $\mathcal{F}^{L}$  the set  $\mathcal{F}^{L}(M_{0})$ , by  $\mathcal{F}$  the set  $\mathcal{F}^{G}(M_{0})$ . Similar notations apply to  $\mathcal{P}$  and  $\mathcal{L}$ .

Let P be a parabolic subgroup of G, denote by  $X(M_P)$  the group of rational characters of  $M_P$ 

$$(1.1.2.1) X(M_P) = \operatorname{Hom}_{\operatorname{Grp}/\mathbb{Q}}(M_P, \operatorname{GL}(1, \mathbb{Q})).$$

Let  $\mathfrak{a}_{\mathrm{P}}$  denote the real vector space

(1.1.2.2) 
$$\mathfrak{a}_{\mathrm{P}} = \operatorname{Hom}_{\mathbb{Z}}(X(\mathrm{M}_{\mathrm{P}}), \mathbb{R}),$$

let  $\mathfrak{a}_{\mathrm{P}}^*$  denote the dual space

(1.1.2.3) 
$$\mathfrak{a}_{\mathrm{P}}^* = X(\mathrm{M}_{\mathrm{P}}) \otimes_{\mathbb{Z}} \mathbb{R}.$$

Let

(1.1.2.4) 
$$\Phi_{\mathrm{P}}, \ \Delta_{\mathrm{P}} \subset \mathfrak{a}_{\mathrm{P}}^{*}, \quad \Phi_{\mathrm{P}}^{\vee}, \ \Delta_{\mathrm{P}}^{\vee} \subset \mathfrak{a}_{\mathrm{P}}$$

denote respectively the set of roots, simple roots, coroots, simple coroots of  $A_P$  in  $\mathfrak{g}$  and  $\mathfrak{n}_P$ . The quadruple  $(X(M_0), \Phi_0, X(M_0)^*, \Phi_0^{\vee})$  is called the root datum of G.

Let  $P_1$  and  $P_2$  be parabolic subgroups of G with  $P_1$  contained in  $P_2$ , denote

(1.1.2.5) 
$$N_1^2 = N_1 \cap M_2, \quad \overline{N}_1^2 = \overline{N}_1 \cap M_2$$

Let  $\Delta_1^2$  and  $\Delta_1^{2,\vee}$  be the set of simple roots and coroots of  $A_1$  in  $\mathfrak{n}_1^2$ . There are canonical splittings

(1.1.2.6) 
$$\mathfrak{a}_1 = \mathfrak{a}_1^2 \oplus \mathfrak{a}_2, \qquad \mathfrak{a}_1^* = \mathfrak{a}_1^{2,*} \oplus \mathfrak{a}_2^*.$$

The sets  $\Delta_1^2$  and  $\Delta_1^{2,\vee}$  form bases of  $\mathfrak{a}_1^{2,*}$  and  $\mathfrak{a}_1^2$ . The respective dual bases are called the *coweights* and *weights* and denoted by  $\hat{\Delta}_1^2$  and  $\hat{\Delta}_1^{2,\vee}$ .

Let  $W_0^G$  be the Weyl group of the pair (G, A<sub>0</sub>). Let M<sub>1</sub> and M<sub>2</sub> be two Levi subgroups of G, define the Weyl set  $W(\mathfrak{a}_1, \mathfrak{a}_2)$  to be the set of linear isomorphisms from  $\mathfrak{a}_1$  to  $\mathfrak{a}_2$  obtained by restricting the action of elements of the Weyl group. The group  $W_0^G$  operates on the root datum of G, hence on  $\mathfrak{a}_0$  and  $\mathfrak{a}_0^*$ . Fix Euclidean inner products on  $\mathfrak{a}_0$  and  $\mathfrak{a}_0^*$  which are compatible with each other and the underlying root datum, hence invariant under the Weyl group action. The inner product on  $\mathfrak{a}_0$  induces an inner product on  $\mathfrak{a}_1^2$ .

Let  $\tau_1^2$  denote the characteristic function on  $\mathfrak{a}_0$  of the points that are positive with respect to every element of  $\Delta_1^2$ , let  $\hat{\tau}_1^2$  denote the characteristic function on  $\mathfrak{a}_0$  of the set of points that are positive with respect to every element of  $\hat{\Delta}_1^{2,\vee}$ :

(1.1.2.7) 
$$\tau_1^2 = \mathbb{I}_{\left\{H \in \mathfrak{a}_0: \ \alpha(H) > 0, \ \forall \alpha \in \Delta_1^2\right\}}, \qquad \hat{\tau}_1^2 = \mathbb{I}_{\left\{H \in \mathfrak{a}_0: \ \varpi(H) > 0, \ \forall \varpi \in \hat{\Delta}_1^{2,\vee}\right\}}.$$

To simplify notations denote by  $\tau_1$  the set  $\tau_1^{\rm G}$ , by  $\hat{\tau}_1$  the set  $\hat{\tau}_1^{\rm G}$ .

(1.1.3) Define  $G(\mathbb{A})^1$  to be the subgroup of  $G(\mathbb{A})$  consisting of elements g of  $G(\mathbb{Q})$  such that

(1.1.3.1) 
$$\forall \chi \in X(\mathbf{G}) \quad |\chi(g)|_{\mathbb{A}} = 1.$$

Fix an admissible maximal compact subgroup

(1.1.3.2) 
$$\mathbf{K} = \prod_{v} \mathbf{K}_{v}$$

of  $G(\mathbb{A})$  such that the Iwasawa decomposition

holds. Denote by

$$H_{\mathrm{P}} : \mathrm{G}(\mathbb{A}) \longrightarrow \mathfrak{a}_{\mathrm{P}}$$

the natural projection.

The Tamagawa measure on  $G(\mathbb{A})$  is the measure induced from the choice of a basis of rational 1-forms on G, which is well-defined by the product formula (1.1.1.2). The Euclidean vector space  $\mathfrak{a}_G$  has a translation invariant measure, which without loss of generality assigns the coweight lattice

$$(1.1.3.4) \qquad \qquad \operatorname{Hom}_{\mathbb{Z}}(X(\mathbf{G}),\mathbb{Z}) \subset \mathfrak{a}_{\mathbf{G}}$$

covolume one.

The various measures are compatible under the Iwasawa decomposition in the sense that

(1.1.3.5) 
$$\int_{\mathcal{G}(\mathbb{A})} f(g) \, dg$$
$$= \int_{\mathcal{M}_{\mathcal{P}}(\mathbb{A})^{1}} \int_{\mathfrak{a}_{\mathcal{P}}} \int_{\mathcal{N}_{\mathcal{P}}(\mathbb{A})} \int_{\mathcal{K}} f(mank) \, dk dn da dm$$
$$= \int_{\mathcal{M}_{\mathcal{P}}(\mathbb{A})^{1}} \int_{\mathfrak{a}_{\mathcal{P}}} \int_{\mathcal{N}_{\mathcal{P}}(\mathbb{A})} \int_{\mathcal{K}} f(nmak) e^{-2\rho_{\mathcal{P}}(H_{0}(a))} \, dk dn da dm$$

where dn is the Tamagawa measure on  $N_P(\mathbb{A})$  which could also be characterized by assigning  $N_P(\mathbb{Q})$  covolume one in  $N_P(\mathbb{A})$ , the point  $\rho_P$  in  $\mathfrak{a}_0^*$  is the Weyl vector defined as the half sum of the roots of  $\mathfrak{n}_P$ . The choices of measures on  $G(\mathbb{A})$  and  $\mathfrak{a}_G$  determine a Haar measure on  $G(\mathbb{A})^1$ , hence a measure on the automorphic quotient  $\mathbb{G}(\mathbb{Q})\backslash G(\mathbb{A})^1$ .

Let T' be a point in  $\mathfrak{a}_0$ , let  $\omega$  be a compact subset of  $N_0(\mathbb{A})M_0(\mathbb{A})^1$ . The Siegel set  $\mathfrak{S}(T', \omega)$  is the subset of  $G(\mathbb{A})^1$  defined as

(1.1.3.6) 
$$\left\{ x = pak : \ p \in \omega, \ a \in \exp(\mathfrak{a}_0), \ k \in \mathcal{K}, \ \beta(H_0(a) - T') \ge 0 \ \forall \beta \in \Delta_0 \right\}.$$

A Siegel set is said to be a *Siegel domain* if it contains a fundamental domain for  $G(\mathbb{Q})\setminus G(\mathbb{A})^1$ :

(1.1.3.7) 
$$G(\mathbb{A})^1 = G(\mathbb{Q})\mathfrak{S}(T',\omega).$$

Let T be a sufficiently positive point in  $\mathfrak{a}_0$ , such a T is said to be a truncation parameter. Let  $\mathfrak{S}(T', \omega)$  be a Siegel domain. The truncated Siegel domain  $\mathfrak{S}^T(T', \omega)$  is defined as

(1.1.3.8) 
$$\left\{ x \in \mathfrak{S}(T',\omega) : \ \varpi(H_0(x) - T) \le 0 \ \forall \varpi \in \hat{\Delta}_0 \right\}.$$

**Proposition** (Borel, Harish-Chandra)

There exist a point T' and a compact set  $\omega$  such that  $\mathfrak{S}(T', \omega)$  is a Siegel domain.

*Proof.* See §13 of [Bo69].

The Siegel domain  $\mathfrak{S}(T',\omega)$ , therefore  $G(\mathbb{Q})\backslash G(\mathbb{A})^1$ , has finite volume. The truncated Siegel domain  $\mathfrak{S}^T(T',\omega)$  is compact and exhausts  $\mathfrak{S}(T',\omega)$  as T approaches infinity. Fix a pair  $(T',\omega)$  such that  $\mathfrak{S}(T',\omega)$  is a Siegel domain, denote by  $F^G(x,T)$  the characteristic function on  $G(\mathbb{Q})\backslash G(\mathbb{A})^1$  of  $\mathfrak{S}^T(T',\omega)$ . There are analogues  $F^P(x,T)$  on  $P(\mathbb{Q})\backslash G(\mathbb{A})^1$ .

(1.1.4) The space  $\mathcal{S}(\mathfrak{g}(\mathbb{A}))$  of *Schwartz functions* on  $\mathfrak{g}(\mathbb{A})$  is defined as the tensor product

(1.1.4.1) 
$$\bigotimes_{p}^{\operatorname{res}} C_{c}^{\infty}(\mathfrak{g}(\mathbb{Q}_{p})) \otimes \mathcal{S}(\mathfrak{g}(\mathbb{R}))$$

restricted at all but finitely many finite primes with respect to the unit vector  $\mathbb{I}_{\mathfrak{g}(\mathbb{Z}_p)}$  in  $C_c^{\infty}(\mathfrak{g}(\mathbb{Q}_p))$ , equipped with the final topology with respect to

(1.1.4.2) 
$$\mathcal{S}(\mathfrak{g}(\mathbb{A})) = \varinjlim_{S} \mathcal{S}(\mathfrak{g}(\mathbb{Q}_{S})).$$

Fix a nondegenerate G-invariant rational bilinear form  $\langle \ , \ \rangle$  on  $\mathfrak{g}$ , fix a global additive unitary character  $\psi$  on  $\mathbb{A}$ 

(1.1.4.3) 
$$\psi : \mathbb{A} \to \mathrm{U}(1)$$

such that

(1.1.4.4) 
$$\forall x \in \mathbb{Q} \quad \psi(x) = 1,$$

and  $\psi(\langle , \rangle)$  identifies  $\mathbb{Q}$  and  $\mathbb{A}/\mathbb{Q}$  as Pontryagin duals of each other. The Fourier transform on  $\mathcal{S}(\mathfrak{g}(\mathbb{A}))$  is defined by

(1.1.4.5) 
$$\hat{f}(X) = \int_{\mathfrak{g}(\mathbb{A})} f(Y)\psi(\langle X, Y \rangle) \, \mathrm{d}Y.$$

The global Fourier transform on  $\mathfrak{g}(\mathbb{A})$  factorizes as the tensor product of the local Fourier transforms on  $\mathfrak{g}(\mathbb{Q}_v)$  with respect to compatible choices of  $\psi_v(\langle , \rangle)$ . Denote by  $\vee$  the inverse Fourier transform.

**Proposition** (Poisson summation formula, Tate) For every Schwartz function f on  $\mathfrak{g}(\mathbb{A})$ ,

(1.1.4.6) 
$$\sum_{X \in \mathfrak{g}(\mathbb{Q})} f(X) = \sum_{X \in \mathfrak{g}(\mathbb{Q})} f^{\hat{}}(X),$$

the sums are the absolutely convergent.

*Proof.* See  $\S4.2$  of [Ta50].

(1.1.5) Define an equivalence relation  $\sim$  on  $\mathfrak{g}(\mathbb{Q})$  by

(1.1.5.1) 
$$X \sim Y \quad \text{if} \quad \exists g \in \mathcal{G}(\mathbb{Q}) \ Y_{ss} = X_{ss} \cdot \operatorname{ad}(g).$$

In general  $\sim$  is weaker than conjugacy by  $G(\mathbb{Q})$ . A typical equivalence class will be denoted by  $\mathfrak{o}$ .

Let D be the discriminant function on  $\mathfrak{g}(\mathbb{Q})$ . Let X be an element of  $\mathfrak{g}(\mathbb{Q})$ , define D(X) to be the coefficient of the characteristic polynomial of  $\operatorname{ad}(X)$ , acting on  $\mathfrak{g}(\mathbb{Q})$  as a linear endomorphism, in degree r, the absolute rank of G:

(1.1.5.2) 
$$r = \operatorname{rank}_{\overline{\mathbb{Q}}}(G \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}).$$

Alternatively D could be defined as the product of the roots of  $\mathfrak{g}$  over an algebraically closed field. Denote by  $D^{\mathrm{M}}$  the discriminant function on  $\mathfrak{m}$  for a Levi subgroup M of G.

Let X be a semisimple element of  $\mathfrak{g}(\mathbb{Q})$ . Then X is said to be *regular* if D(X) does not vanish. Denote by  $\mathfrak{g}_{reg.ss}$  the locus of regular semisimple points on  $\mathfrak{g}$ . If an equivalence class  $\mathfrak{o}$  contains a regular semisimple element, the set  $\mathfrak{o}$  is a  $G(\mathbb{Q})$ -orbit consisting of regular semisimple elements. Such an  $\mathfrak{o}$  is said to be *regular*, otherwise  $\mathfrak{o}$  is said to be *singular*. A semisimple element X is said to be  $\mathbb{Q}$ -*elliptic* if it is stablized under the adjoint action by a maximal torus that is anisotropic over  $\mathbb{Q}$  modulo the center of G.

Denote by  $G_X$  the centralizer of X in G, let  $G_X^0$  be the connected component of the identity of  $G_{\Sigma}$ , let  $\pi_0(G_X)$  be the group of connected components of  $G_X$ . There is an exact sequence

$$(1.1.5.3) 1 \longrightarrow \mathcal{G}_X^0 \longrightarrow \mathcal{G}_X \longrightarrow \pi_0(\mathcal{G}_X) \longrightarrow 1 .$$

(1.1.6) Let S be a finite set of places of  $\mathbb{Q}$ , denote by  $G_S$  the base change  $G \otimes_{\mathbb{Q}} \mathbb{Q}_S$  of G. Let v be a place of  $\mathbb{Q}$ , denote by  $G_v$  the base change  $G \otimes_{\mathbb{Q}} \mathbb{Q}_v$  of G. Similar notations apply to the Lie algebra  $\mathfrak{g}$ . The underlying topological groups of  $G(\mathbb{Q}_S)$  and  $G_S(\mathbb{Q}_S)$  are the same, however

(1.1.6.1) 
$$X(\mathbf{G}) = \operatorname{Hom}_{\operatorname{Grp}/\mathbb{Q}}(\mathbf{G}, \operatorname{GL}(1, \mathbb{Q})),$$
$$X(\mathbf{G}_S) = \operatorname{Hom}_{\operatorname{Grp}/\mathbb{Q}_S}(\mathbf{G}_S, \operatorname{GL}(1, \mathbb{Q}_S))$$

are in general different.

All the constructions above generalize to the local and S-local settings with similar caveats:

- Fix a minimal parabolic subgroup  $P_{S,0}$  of  $G_S$  contained in  $P_{0,S}$ , fix a Levi subgroup  $M_{S,0}$  contained in  $M_{0,S}$  with split component  $A_{S,0}$  containing  $A_{0,S}$ . The choice of  $P_{S,0}$  is equivalent to a choice of a minimal parabolic  $P_{v,0}$  of  $G_v$  for each v in S, similarly for  $M_{S,0}$  and  $A_{S,0}$ .
- Let  $M_S$  and  $L_S$  be Levi subgroups of  $G_S$  such that  $M_S$  is contained in  $L_S$ , denote by  $\mathcal{F}^{L_S}(M_S)$ ,  $\mathcal{P}^{L_S}(M_S)$  and  $\mathcal{L}^{L_S}(M_S)$  the analogous sets of parabolic and Levi subgroups. The Levi subgroups  $M_S$  and  $L_S$  determine local Levi subgroups  $M_v$  and  $L_v$  for each v in S. There are bijections

(1.1.6.2) 
$$\mathcal{F}^{\mathcal{L}_{S}}(\mathcal{M}_{S}) = \prod_{v \in S} \mathcal{F}^{\mathcal{L}_{v}}(\mathcal{M}_{v}),$$
$$\mathcal{P}^{\mathcal{L}_{S}}(\mathcal{M}_{S}) = \prod_{v \in S} \mathcal{P}^{\mathcal{L}_{v}}(\mathcal{M}_{v}),$$

$$\mathcal{L}^{\mathcal{L}_S}(\mathcal{M}_S) = \prod_{v \in S} \mathcal{L}^{\mathcal{L}_v}(\mathcal{M}_v).$$

If  $M_S$  and  $L_S$  are the base change of Levi subgroups M and L of G from  $\mathbb{Q}$  to  $\mathbb{Q}_S$ , there are diagonal inclusions

- (1.1.6.3)  $\mathcal{F}^{L}(M) \subset \mathcal{F}^{L_{S}}(M_{S}),$  $\mathcal{P}^{L}(M) \subset \mathcal{P}^{L_{S}}(M_{S}),$  $\mathcal{L}^{L}(M) \subset \mathcal{L}^{L_{S}}(M_{S}).$
- Let  $P_S$  be a parabolic subgroup of  $G_S$ , define real vector spaces
  - (1.1.6.4)  $\mathfrak{a}_{\mathrm{P}_S} = \mathrm{Hom}_{\mathbb{Z}}(X(\mathrm{M}_{\mathrm{P}_S}), \mathbb{R}), \quad \mathfrak{a}_{\mathrm{P}_S}^* = X(\mathrm{M}_{\mathrm{P}_S}) \otimes_{\mathbb{Z}} \mathbb{R}.$

If  $P_S$  is the base change of a parabolic subgroup P of G from  $\mathbb{Q}$  to  $\mathbb{Q}_S$ , there are diagonal inclusions

(1.1.6.5) 
$$\mathfrak{a}_{\mathrm{P}} \subset \mathfrak{a}_{\mathrm{P}_{S}} = \bigoplus_{v \in S} \mathfrak{a}_{\mathrm{P}_{v}}, \qquad \mathfrak{a}_{\mathrm{P}}^{*} \subset \mathfrak{a}_{\mathrm{P}_{S}}^{*} = \bigoplus_{v \in S} \mathfrak{a}_{\mathrm{P}_{v}}^{*}.$$

• A maximal torus  $T_S$  of  $G_S$  is equivalent to the choice of a maximal torus  $T_v$  of  $G_v$  for each v in S. The associated Cartan subalgebra  $\mathfrak{t}_S$ , which is a free  $\mathbb{Q}_S$ -module, is equal as an abelian group to the direct sum of the  $\mathbb{Q}_v$ -vector spaces  $\mathfrak{t}_v$  for all v in S.

A maximal torus  $T_S$  is said to be *elliptic* in  $G_S$  if it is anisotropic modulo  $A_{G_S}$  over  $\mathbb{Q}_S$ . A maximal torus  $T_S$  is elliptic in  $G_S$  if and only if  $T_v$  is elliptic in  $G_v$  for each v in S. If this is the case,  $T_S(\mathbb{Q}_S)$  is compact modulo  $A_{G_S}(\mathbb{Q}_S)$  in the analytic topology.

Denote by  $\mathcal{T}_{ell}(G_S)$  the set of conjugacy classes of elliptic maximal tori of  $G_S$ .

• Let  $W_{S,0}^{G_S}$  be the Weyl group of the pair  $(G_S, A_{S,0})$ , there is a bijection

(1.1.6.6) 
$$W_{S,0}^{G_S} = \prod_{v \in S} W_{v,0}^{G_v},$$

the linear representation of  $W_{S,0}^{G_S}$  on  $\mathfrak{a}_{S,0}$  is the direct sum of the local representations. Denote by  $W(G_S, T_S)$  the Weyl group of the pair  $(G_S(\mathbb{Q}_S), T_S(\mathbb{Q}_S))$ . There is a bijection

(1.1.6.7) 
$$W(\mathbf{G}_S, \mathbf{T}_S) = \prod_{v \in S} W(\mathbf{G}_v, \mathbf{T}_v).$$

• The Schwartz space  $\mathcal{S}(\mathfrak{g}_S(\mathbb{Q}_S))$ , the Fourier transform  $\wedge$ , the discriminant function  $D^{\mathcal{G}_S}$ , and the regular semisimple locus  $\mathfrak{g}_{S, \operatorname{reg.ss}}(\mathbb{Q}_S)$  are unchanged under base change from  $\mathbb{Q}$  to  $\mathbb{Q}_S$ .

**Proposition** (Weyl integration formula) If  $f_S$  is a Schwartz function on  $\mathfrak{g}_S(\mathbb{Q}_S)$ , then

(1.1.6.8) 
$$\int_{\mathfrak{g}_{S}(\mathbb{Q}_{S})} f_{S}(X) \, \mathrm{d}X$$
$$= \sum_{\mathcal{M}_{S}\in\mathcal{L}^{\mathcal{G}_{S}}} |\mathcal{W}_{S,0}^{\mathcal{M}_{S}}| |\mathcal{W}_{S,0}^{\mathcal{G}_{S}}|^{-1} \sum_{\mathcal{T}_{S}\in\mathcal{T}_{\mathrm{ell}}(\mathcal{M}_{S})} |\mathcal{W}(\mathcal{M}_{S},\mathcal{T}_{S})|^{-1} \times$$
$$\times \int_{\mathfrak{t}_{S}(\mathbb{Q}_{S})} |D^{\mathcal{G}_{S}}(X)|_{S} \int_{\mathcal{A}_{\mathcal{M}_{S}}(\mathbb{Q}_{S})\backslash\mathcal{G}_{S}(\mathbb{Q}_{S})} f_{S}(X \cdot \mathrm{ad}(x)) \, \mathrm{d}x \mathrm{d}X.$$

*Proof.* For the *p*-adic case see §7.11 of [Ko05]. For the real case see Lemma 2 on page 35 of [Va77] and the references therein.  $\Box$ 

**Definition** Let X be an element of  $\mathfrak{g}_{S,\operatorname{reg.ss}}(\mathbb{Q}_S)$ . Define the S-local orbital integral  $I_G^G(X, )$  to be the invariant distribution on  $\mathfrak{g}_S(\mathbb{Q}_S)$  such that

(1.1.6.9) 
$$\forall f_S \in \mathcal{S}(\mathfrak{g}_S(\mathbb{Q}_S))$$
$$I_G^G(X, f_S) = |D^G(X)|_S^{1/2} \int_{G^0_{S,X}(\mathbb{Q}_S) \setminus G_S(\mathbb{Q}_S)} f_S(X \cdot \operatorname{ad}(x)) \, \mathrm{d}x$$

where  $G_{S,X}^0$  denotes the connected component of the identity of the stablizer subgroup of X in  $G_S$ .

### 2 The non-invariant trace formula

In this chapter a preliminary version of the trace formula for the reductive Lie algebra  $\mathfrak{g}$  established in Chaudouard [Ch02a] is recalled.

### 2.1 A motivating example

(2.1.1) Definition Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , let  $\mathfrak{o}$  be a ~ equivalence class on  $\mathfrak{g}(\mathbb{Q})$ . Define the *kernel functions* K(x, f) and  $K_{\mathfrak{o}}(x, f)$  by

(2.1.1.1) 
$$\forall x \in \mathcal{G}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{A}) \qquad K(x, f) = \sum_{X \in \mathfrak{g}(\mathbb{Q})} f(X \cdot \operatorname{ad}(x)),$$
$$K_{\mathfrak{o}}(x, f) = \sum_{X \in \mathfrak{o}} f(X \cdot \operatorname{ad}(x)).$$

(2.1.2) **Remark** By the Poisson summation formula (1.1.4.6), the function K(x, f) satisfies the functional equation

(2.1.2.1) 
$$K(x, f) = K(x, f^{\hat{}}).$$

(2.1.3) **Proposition** Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ . If G is anisotropic over  $\mathbb{Q}$ , then

$$(2.1.3.1) \qquad \lim_{S} \sum_{\substack{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim \\ regular}} \operatorname{Vol}(\mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{Q}) \backslash \mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{A})) \cdot I_{\mathrm{G}}^{\mathrm{G}}(X_{\mathfrak{o}}, f_{S}) \\ + \sum_{\substack{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim \\ singular}} \operatorname{Vol}(\mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{Q}) \backslash \mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{A})) \cdot I_{\mathfrak{o}}(f) \\ = \lim_{S} \sum_{\substack{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim \\ regular}} \operatorname{Vol}(\mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{Q}) \backslash \mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{A})) \cdot I_{\mathrm{G}}^{\mathrm{G}}(X_{\mathfrak{o}}, f^{\widehat{}}_{S}) \\ + \sum_{\substack{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim \\ singular}} \operatorname{Vol}(\mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{Q}) \backslash \mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{A})) \cdot I_{\mathfrak{o}}(f^{\widehat{}}), \end{cases}$$

where for each class  $\mathfrak{o}$  choose an element  $X_{\mathfrak{o}}$  of  $\mathfrak{o}$ , for each singular class  $\mathfrak{o}$  define

(2.1.3.2) 
$$I_{\mathfrak{o}}(f) = \int_{\mathrm{G}^{0}_{X_{\mathfrak{o}}}(\mathbb{A})\backslash\mathrm{G}(\mathbb{A})} f(X_{\mathfrak{o}} \cdot \mathrm{ad}(x)) \, \mathrm{d}x.$$

*Proof.* The function K(x, f) is continuous. By assumption G is anisotropic, so  $G(\mathbb{Q})\setminus G(\mathbb{A})$  is compact. Therefore K(x, f) is absolutely integrable, hence

$$(2.1.3.3) \qquad \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})} K(x,f) \, \mathrm{d}x$$

$$= \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} K_{\mathfrak{o}}(x,f) \, \mathrm{d}x$$

$$= \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})} K_{\mathfrak{o}}(x,f) \, \mathrm{d}x$$

$$= \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} \operatorname{Vol}(\mathcal{G}_{X_{\mathfrak{o}}}^{0}(\mathbb{Q})\backslash \mathcal{G}_{X_{\mathfrak{o}}}^{0}(\mathbb{A})) \int_{\mathcal{G}_{X_{\mathfrak{o}}}^{0}(\mathbb{A})\backslash \mathcal{G}(\mathbb{A})} f(X_{\mathfrak{o}} \cdot \operatorname{ad}(x)) \, \mathrm{d}x$$

$$(2.1.3.4) \qquad = \lim_{S} \sum_{\substack{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim\\ \operatorname{regular}}} \operatorname{Vol}(\mathcal{G}_{X_{\mathfrak{o}}}^{0}(\mathbb{Q})\backslash \mathcal{G}_{X_{\mathfrak{o}}}^{0}(\mathbb{A})) \cdot I_{\mathcal{G}}^{\mathcal{G}}(X_{\mathfrak{o}}, f_{S})$$

$$+ \sum_{\substack{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim\\ \operatorname{singular}}} \operatorname{Vol}(\mathcal{G}_{X_{\mathfrak{o}}}^{0}(\mathbb{Q})\backslash \mathcal{G}_{X_{\mathfrak{o}}}^{0}(\mathbb{A})) \cdot I_{\mathfrak{o}}(f^{\wedge}),$$

the equality (2.1.3.4) follows from

(2.1.3.5) 
$$\forall X_{\mathfrak{o}} \in \mathfrak{g}(\mathbb{Q}) \ \exists S \ \forall S' \supset S \qquad |D(X_{\mathfrak{o}})|_{S'} = 1$$

by the product formula (1.1.1.2). The proposition follows from the functional equation (2.1.2.1).  $\hfill \Box$ 

### 2.2 The non-invariant trace formula of Chaudouard

(2.2.1) Definition Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , let  $\mathfrak{o}$  be a ~ equivalence class on  $\mathfrak{g}(\mathbb{Q})$ , let T be a truncation parameter. Define the *truncated kernel function*  $k_{\mathfrak{o}}^{T}(\cdot, f)$  on  $G(\mathbb{Q})\backslash G(\mathbb{A})^{1}$  by

(2.2.1.1) 
$$\forall x \in \mathcal{G}(\mathbb{Q}) \setminus \mathcal{G}(\mathbb{A})^{1}$$
$$k_{\mathfrak{o}}^{T}(x, f) = \sum_{\substack{\mathcal{P} \in \mathcal{F} \\ \text{standard}}} (-1)^{\dim(\mathcal{A}_{\mathcal{P}}/\mathcal{A}_{\mathcal{G}})} \sum_{\delta \in \mathcal{P}(\mathbb{Q}) \setminus \mathcal{G}(\mathbb{Q})} \hat{\tau}_{\mathcal{P}}(H_{0}(\delta x) - T) K_{\mathcal{P},\mathfrak{o}}(\delta x)$$

where

(2.2.1.2) 
$$K_{\mathcal{P},\mathfrak{o}}(x,f) = \sum_{X \in \mathfrak{m}_{\mathcal{P}}(\mathbb{Q}) \cap \mathfrak{o}} \int_{\mathfrak{n}_{\mathcal{P}}(\mathbb{A})} f((X+N) \cdot \operatorname{ad}(x)) \, \mathrm{d}N.$$

Define distributions  $J^T_{\mathfrak{o}}$  and  $J^T$  on  $\mathfrak{g}(\mathbb{A})$  by

(2.2.1.3) 
$$\forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A})) \qquad J^T_{\mathfrak{o}}(f) = \int_{\mathcal{G}(\mathbb{Q}) \setminus \mathcal{G}(\mathbb{A})^1} k^T_{\mathfrak{o}}(x, f) \, \mathrm{d}x,$$
$$J^T(f) = \int_{\mathcal{G}(\mathbb{Q}) \setminus \mathcal{G}(\mathbb{A})^1} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} k^T_{\mathfrak{o}}(x, f) \, \mathrm{d}x$$

(2.2.2) **Proposition** Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , let T be a truncation parameter, then

(2.2.2.1) 
$$\int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^1} \sum_{\mathfrak{o}\in\mathfrak{g}(\mathbb{Q})/\sim} \left| k_\mathfrak{o}^T(x,f) \right| \, \mathrm{d}x < \infty.$$

*Proof.* This is Théorème 3.1 of [Ch02a]. The following is a sketch of the argument, for the details please refer to loc. cit.

The proof depends on the following lemma.

**Lemma** (Combinatorial lemma of Langlands) Let  $P_1$  and  $P_3$  be parabolic subgroups of G such that  $P_1$  is contained in  $P_3$ , then

(2.2.2.2) 
$$\sum_{\substack{P_2 \in \mathcal{F} \\ P_1 \subset P_2 \subset P_3}} (-1)^{\dim(A_2/A_3)} \tau_1^2(H) \hat{\tau}_2^3(H) = \begin{cases} 1 & \text{if } P_1 = P_3, \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* See  $\S6$  of [Ar78].

By (2.2.2.2)

(2.2.2.3) 
$$\int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^1} \sum_{\mathfrak{o}\in\mathfrak{g}(\mathbb{Q})/\sim} \left| k_{\mathfrak{o}}^T(x,f) \right| \, \mathrm{d}x$$

$$\leq \sum_{\substack{P_1, P_4 \in \mathcal{F} \\ \text{standard} \\ P_1 \subset P_4}} \int_{P_1(\mathbb{Q}) \setminus G(\mathbb{A})^1} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} F^{P_1}(x, T) \sigma_1^4(H_0(x) - T) \times \\ \times \left| \sum_{\substack{P_3 \in \mathcal{F} \\ P_1 \subset P_3 \subset P_4}} (-1)^{\dim(A_3/A_G)} K_{P_3, \mathfrak{o}}(x, f) \right| dx$$

where

(2.2.2.4) 
$$\sigma_1^4(H) = \sum_{\substack{P_5 \in \mathcal{F} \\ P_4 \subset P_5}} (-1)^{\dim(A_4/A_5)} \tau_1^5(H) \hat{\tau}_5(H).$$

The second factor of the integrand of the right hand side of (2.2.2.3) satisfies the inequality

$$(2.2.2.5) \qquad \left| \sum_{\substack{P_{3}\in\mathcal{F}\\P_{1}\subset P_{3}\subset P_{4}}} (-1)^{\dim(A_{3}/A_{G})} K_{P_{3},\mathfrak{o}}(x,f) \right|$$

$$\leq \sum_{\substack{P_{2}\in\mathcal{F}\\P_{1}\subset P_{2}\subset P_{4}}} \left| \sum_{\substack{P_{3}\in\mathcal{F}\\P_{2}\subset P_{3}\subset P_{4}}} (-1)^{\dim(A_{3}/A_{G})} \sum_{X\in\mathfrak{m}_{1}^{2}(\mathbb{Q})'\cap\mathfrak{o}} \right|$$

$$\times \sum_{Y\in\mathfrak{m}_{2}^{3}(\mathbb{Q})} \int_{\mathfrak{n}_{3}(\mathbb{A})} f((X+Y+N)\cdot\mathrm{ad}(x)) \,\mathrm{d}N \right|$$

$$(2.2.2.6) \qquad = \left| \sum_{X\in\mathfrak{m}_{1}^{4}(\mathbb{Q})'\cap\mathfrak{o}} \int_{\mathfrak{n}_{4}(\mathbb{A})} f((X+N)\cdot\mathrm{ad}(x)) \,\mathrm{d}N \right| +$$

$$+ \sum_{\substack{P_{2}\in\mathcal{F}\\P_{1}\subset P_{2}\subseteq P_{4}}} \left| \sum_{\substack{P_{3}\in\mathcal{F}\\P_{2}\subset P_{3}\subset P_{4}}} (-1)^{\dim(A_{3}/A_{G})} \sum_{X\in\mathfrak{m}_{1}^{2}(\mathbb{Q})'\cap\mathfrak{o}} \right|$$

$$\times \sum_{\overline{Y\in\mathfrak{m}_{2}^{3}(\mathbb{Q})'}} \int_{\mathfrak{n}_{2}(\mathbb{A})} f((X+N)\cdot\mathrm{ad}(x))\psi(\langle N,\overline{Y}\rangle) \,\mathrm{d}N \right|$$

where  $\mathfrak{m}_{P}^{Q}(\mathbb{Q})'$  denotes the set of points of  $\mathfrak{m}_{P}^{Q}(\mathbb{Q})$  not contained in any proper parabolic subalgebra of  $\mathfrak{q}$ . Similar notation applies to  $\mathfrak{n}$ . The equality (2.2.2.6) follows from the Poisson summation formula applied to  $\mathfrak{n}_{2}^{3}(\mathbb{Q})$  as a lattice in  $\mathfrak{n}_{2}^{3}(\mathbb{A})$ .

By the inclusion-exclusion principle applied to

(2.2.2.7) 
$$\sum_{\substack{P_3 \in \mathcal{F} \\ P_2 \subset P_3 \subset P_4}} (-1)^{\dim(A_3/A_G)}$$

the last expression in (2.2.2.5) reduces to a majorant of the form

(2.2.2.8) 
$$\prod_{\alpha \in \Delta_1^4} \int_0^\infty (1+t_\alpha)^{p_\alpha} e^{-q_\alpha t_\alpha} \, \mathrm{d}t_\alpha$$

for some natural numbers  $p_{\alpha}$  and  $q_{\alpha}$ , which is finite.

(2.2.3) **Definition** Let  $P_2$  be a standard parabolic subgroup of G, let T be a truncation parameter in  $\mathfrak{a}_0$ . Define the geometric gamma' function  $\Gamma'_{P_2}(, T)$  on  $\mathfrak{a}_0$  by

(2.2.3.1) 
$$\forall H \in \mathfrak{a}_0 \quad \Gamma'_{P_2}(H,T) = \sum_{\substack{P_3 \in \mathcal{F} \\ P_2 \subset P_3}} (-1)^{\dim(A_3/A_G)} \tau_2^3(H) \hat{\tau}_3(H-T).$$

(2.2.4) **Remark** For each parabolic subgroup  $P_1$  of G the geometric gamma' functions satisfy the identity

(2.2.4.1) 
$$\hat{\tau}_1(H-T) = \sum_{\substack{P_2 \in \mathcal{F} \\ P_1 \subset P_2}} (-1)^{\dim(A_2/A_G)} \hat{\tau}_1^2(H) \Gamma'_{P_2}(H,T).$$

For a proof see page 13 of [Ar81].

(2.2.5) **Proposition** Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , let  $\mathfrak{o}$  be a ~ equivalence class, let T be a truncation parameter. Then  $J^T_{\mathfrak{o}}(f)$  and  $J^T(f)$  are polynomials in T of degree at most dim $(A_0/A_G)$ .

*Proof.* This is Théorème 4.2 of [Ch02a]. The following is a sketch of the argument for  $J_{\mathfrak{o}}^T(f)$ , the argument for  $J^T(f)$  is the same. For the details please refer to loc. cit.

Fix a point T' in  $\mathfrak{a}_0$  that is sufficiently positive and assume that T dominates T'. By (2.2.4.1)

(2.2.5.1)  $J_{o}^{T}(f)$ 

$$= \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} \left( \sum_{\substack{P_{1}\in\mathcal{F}\\\text{standard}}} (-1)^{\dim(A_{1}/A_{G})} \times \right) \\ \times \sum_{\delta\in P_{1}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{Q})} \hat{\tau}_{1}(H_{0}(\delta x) - T) \cdot K_{P_{1},\mathfrak{o}}(\delta x) dx$$
$$= \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} \left( \sum_{\substack{P_{1},P_{2}\in\mathcal{F}\\\text{standard}\\P_{1}\subset P_{2}}} (-1)^{\dim(A_{1}/A_{2})} \times \right) \\ \times \sum_{\delta\in P_{1}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{Q})} \hat{\tau}_{1}^{2}(H_{P_{1}}(\delta x) - T')\Gamma'_{P_{2}}(H_{P_{2}}(\delta x) - T', T - T') \times$$

$$\times \sum_{X \in \mathfrak{m}_{1}(\mathbb{Q}) \cap \mathfrak{o}} \int_{\mathfrak{n}_{1}(\mathbb{A})} f((X+N) \cdot \operatorname{ad}(\delta x)) \, \mathrm{d}N \right) \, \mathrm{d}x$$

$$= \sum_{\substack{P_{2} \in \mathcal{F} \\ \operatorname{standard}}} \int_{M_{2}(\mathbb{Q}) \setminus M_{2}(\mathbb{A})^{1}} \left( \sum_{\substack{P_{1} \in \mathcal{F} \\ \operatorname{standard}}} (-1)^{\operatorname{dim}(A_{1}/A_{2})} \times \right)$$

$$\times \sum_{\delta \in P_{1}(\mathbb{Q}) \cap M_{2}(\mathbb{Q}) \setminus M_{2}(\mathbb{Q})} \hat{\tau}_{1}^{2} (H_{P_{1}}(\delta x) - T') \sum_{X \in \mathfrak{m}_{1}(\mathbb{Q}) \cap \mathfrak{o}} \int_{\mathfrak{n}_{1}^{2}(\mathbb{A})}$$

$$\times \left( \int_{K} \int_{\mathfrak{n}_{2}(\mathbb{A})} f((X+N) \cdot \operatorname{ad}(\delta xk) + N' \cdot \operatorname{ad}(k)) \, \mathrm{d}N' \mathrm{d}k \right) \, \mathrm{d}N \right) \, \mathrm{d}x \times$$

$$\times \int_{\mathfrak{a}_{2}^{G}} \Gamma'_{P_{2}} (H - T', T - T') \, \mathrm{d}H$$

$$= \sum_{\substack{P_{2} \in \mathcal{F} \\ \operatorname{standard}}} J_{\mathfrak{o}}^{M_{2},T'} (f_{P_{2}}) \int_{\mathfrak{a}_{2}^{G}} \Gamma'_{P_{2}} (H, T - T') \, \mathrm{d}H$$

where  $J^{\mathrm{M}}_{\mathfrak{o}}$  is defined to be the sum of  $J^{\mathrm{M}}_{\mathfrak{o}'}$  over all the  $\mathrm{M}(\mathbb{Q}) \sim$  equivalence classes  $\mathfrak{o}'$  contained in  $\mathfrak{o}$  and  $f_{\mathrm{P}}$  is defined by

(2.2.5.2) 
$$f_{\mathrm{P}}(X) = \int_{\mathrm{K}} \int_{\mathfrak{n}_{\mathrm{P}}(\mathbb{A})} f((X+N) \cdot \mathrm{ad}(k)) \, \mathrm{d}N \mathrm{d}k.$$

Because  $\int \Gamma'_{\rm P}(H, T - T') \, dH$  is a polynomial in T which is homogeneous of degree  $\dim(A_{\rm P}/A_{\rm G})$ , Proposition (2.2.5) follows by induction.

(2.2.6) Proposition Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , let T be a truncation parameter. For every positive  $\epsilon$ 

$$(2.2.6.1) \quad \left| J^T(f) - \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^1} F^{\mathcal{G}}(x,T) \sum_{X \in \mathfrak{g}(\mathbb{Q})} f(X \cdot \operatorname{ad}(x)) \, \mathrm{d}x \right| = O(e^{-\epsilon \|T\|})$$

where  $\| \|$  denotes the Euclidean norm on  $\mathfrak{a}_0$ , as T approaches infinity such that T is uniformly bounded away from the walls of the positive chamber.

*Proof.* This is a corollary of the proof of Lemma 3.2.2.1.

(2.2.7) **Proposition** (Non-invariant trace formula of Chaudouard) Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , then

(2.2.7.1) 
$$\sum_{\mathfrak{o}\in\mathfrak{g}(\mathbb{Q})/\sim} J^T_{\mathfrak{o}}(f) = \sum_{\mathfrak{o}\in\mathfrak{g}(\mathbb{Q})/\sim} J^T_{\mathfrak{o}}(f^{\hat{}})$$

holds as an equality between polynomials in T.

*Proof.* By (2.1.2.1)

(2.2.7.2)  

$$\int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} F^{\mathcal{G}}(x,T) \sum_{X \in \mathfrak{g}(\mathbb{Q})} f(X \cdot \operatorname{ad}(x)) \, \mathrm{d}x$$

$$= \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} F^{\mathcal{G}}(x,T) K(x,f) \, \mathrm{d}x$$

$$= \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} F^{\mathcal{G}}(x,T) K(x,f^{\hat{}}) \, \mathrm{d}x.$$

By (2.2.6.1) the difference between  $J^T(f)$  and  $J^T(f^{\hat{}})$  converges to zero as T approaches infinity. By Proposition (2.2.5)  $J^T(f)$  and  $J^T(f^{\hat{}})$  are both polynomials in T, hence equal to each other.

(2.2.8) Lemma There exists a unique point  $T_0$  in  $\mathfrak{a}_0^G$  such that

(2.2.8.1)  $\forall s \in W_0^G \quad H_0(w_s^{-1}) = T_0 - s^{-1}T_0$ 

where  $w_s$  denotes a representative of s in  $G(\mathbb{Q})$ .

*Proof.* See Lemma 1.1 of [Ar81].

(2.2.9) **Definition** Define distributions J and  $J_{\mathfrak{o}}$  on  $\mathfrak{g}(\mathbb{A})$  by

(2.2.9.1) 
$$\forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A})) \qquad J(f) = J^{T_0}(f), \ J_{\mathfrak{o}}(f) = J_{\mathfrak{o}}^{T_0}(f).$$

(2.2.10) **Remark** The coefficients of the polynomials  $J^T(f)$  and  $J^T_{\mathfrak{o}}(f)$  in positive degrees depend only on the orbital integrals of  $f_P$  along proper Levi subalgebras of  $\mathfrak{g}$ . Therefore the constant terms J(f) and  $J_{\mathfrak{o}}(f)$  contain the essential information. The choice of  $T_0$  implies that the distributions J and  $J_{\mathfrak{o}}$  are independent of the choice of the minimal parabolic subgroup  $P_0$ .

(2.2.11) **Proposition** Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ . If  $\mathfrak{o}$  is the conjugacy class of a regular semisimple element X which is contained in a standard parabolic  $\mathfrak{p}$  and elliptic in its Levi component  $\mathfrak{m}$ , then

$$(2.2.11.1) \quad J_{\mathfrak{o}}(f) = |\pi_0(\mathcal{G}_X)|^{-1} \operatorname{Vol}(\mathcal{M}^0_X(\mathbb{Q}) \setminus \mathcal{M}^0_X(\mathbb{A})^1) \int_{\mathcal{G}^0_X(\mathbb{A}) \setminus \mathcal{G}(\mathbb{A})} f(X \cdot \operatorname{ad}(x)) \ v_{\mathcal{M}}(x) \mathrm{d}x$$

where the weight factor  $v_{\rm M}(x)$  is defined to be the volume of the convex hull of

(2.2.11.2) 
$$\left\{ -H_{\mathbf{P}}(x) : \mathbf{P} \in \mathcal{P}(\mathbf{M}) \right\} \subset \mathfrak{a}_{\mathbf{M}}^{\mathbf{G}}.$$

*Proof.* This is a corollary of Theorem 3.3.11.1. See also (5.4) of [Ch02a].

(2.2.12) **Proposition** Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , let  $\mathfrak{o}$  be a ~ equivalence class, then

(2.2.12.1) 
$$\forall x \in \mathcal{G}(\mathbb{A})^1 \qquad J_{\mathfrak{o}}(f \circ \mathrm{ad}(x)) = \sum_{\mathcal{P} \in \mathcal{F}} |\mathcal{W}_0^{\mathcal{M}_{\mathcal{P}}}| |\mathcal{W}_0^{\mathcal{G}}|^{-1} J_{\mathfrak{o}}^{\mathcal{M}_{\mathcal{P}}}(f_{\mathcal{P},x})$$

where  $J^{\mathrm{M}}_{\mathfrak{o}}$  is defined to be the sum of  $J^{\mathrm{M}}_{\mathfrak{o}'}$  over all the  $\mathrm{M}(\mathbb{Q}) \sim$  equivalence classes  $\mathfrak{o}'$  contained in  $\mathfrak{o}$ , and the function  $f_{\mathrm{P},x}$  on  $\mathfrak{m}_{\mathrm{P}}(\mathbb{A})$  is defined as

(2.2.12.2) 
$$f_{\mathbf{P},x}(X) = \int_{\mathbf{K}} \int_{\mathfrak{n}_{\mathbf{P}}(\mathbb{A})} f((X+N) \cdot \mathrm{ad}(k)) \ v'_{\mathbf{P}}(kx) \mathrm{d}N \mathrm{d}k$$

where the weight factor  $v'_{\rm P}(x)$  is defined by

(2.2.12.3) 
$$v'_{\rm P}(x) = \int_{\mathfrak{a}_{\rm P}^{\rm G}} \Gamma'_{\rm P}(H, -H_{\rm P}(x)) \, \mathrm{d}H$$

Proof. By definition

$$(2.2.12.4) \quad J^{T}_{\mathfrak{o}}(f \circ \operatorname{ad}(x)) = \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} \left( \sum_{\substack{\mathbf{P}_{1} \in \mathcal{F} \\ \operatorname{standard}}} (-1)^{\operatorname{dim}(\mathbf{A}_{1}\mathbf{A}_{G})} \times \sum_{\delta \in \mathcal{P}_{1}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{Q})} \hat{\tau}_{1}(H_{0}(\delta yx) - T) \cdot K_{\mathcal{P}_{1},\mathfrak{o}}(\delta yx) \, \mathrm{d}y \right).$$

By (2.2.4.1)

$$(2.2.12.5) \qquad \hat{\tau}_1(H_{P_1}(\delta yx) - T) = \sum_{\substack{P_2 \in \mathcal{F} \\ P_1 \subset P_2}} (-1)^{\dim(A_2/A_G)} \hat{\tau}_1^2(H_{P_1}(\delta y) - T) \times \\ \times \Gamma'_{P_2}(H_{P_1}(\delta y) - T, -H_{P_1}(kx))$$

where k is a K component of  $\delta y$  under the decomposition of  $G(\mathbb{A})$  as  $P_1(\mathbb{A})K$ , the point  $H_{P_1}(kx)$  is independent of the choice of the element k. Hence the right hand side of (2.2.12.4) is equal to

$$(2.2.12.6) \qquad \sum_{\substack{P_2 \in \mathcal{F} \\ \text{standard}}} \int_{M_2(\mathbb{Q}) \setminus M_2(\mathbb{A})^1} \left( \sum_{\substack{P_1 \in \mathcal{F} \\ \text{standard}}} (-1)^{\dim(A_1/A_2)} \times \right) \\ \times \sum_{\delta \in P_1(\mathbb{Q}) \cap M_2(\mathbb{Q}) \setminus M_2(\mathbb{Q})} \hat{\tau}_1^2 (H_{P_1}(\delta y) - T) \sum_{X \in \mathfrak{m}_1(\mathbb{Q}) \cap \mathfrak{o}} \int_{\mathfrak{n}_1^2(\mathbb{A})} \\ \times \left( \int_K \int_{\mathfrak{n}_2(\mathbb{A})} f((X + N) \cdot \operatorname{ad}(\delta yx) + N' \cdot \operatorname{ad}(kx)) \times \right) \right)$$

$$\times \left( \int_{\mathfrak{a}_2^{\mathbf{G}}} \Gamma'_{\mathbf{P}_2}(H - T, -H_{\mathbf{P}_2}(kx)) \, \mathrm{d}H \right) \, \mathrm{d}N' \mathrm{d}k \ \right) \, \mathrm{d}N \right) \, \mathrm{d}y.$$

Substituting  $T_0$  for T, the expression (2.2.12.6) becomes

(2.2.12.7) 
$$\sum_{\substack{\mathbf{P}_{2} \in \mathcal{F} \\ \text{standard}}} J_{\mathfrak{o}}^{\mathbf{M}_{2}}(f_{\mathbf{P}_{2},x}) = \sum_{\mathbf{P}_{2} \in \mathcal{F}} |\mathbf{W}_{0}^{\mathbf{M}_{2}}| |\mathbf{W}_{0}^{\mathbf{G}}|^{-1} J_{\mathfrak{o}}^{\mathbf{M}_{2}}(f_{\mathbf{P}_{2},x})$$

since the distribution  $J_{\mathfrak{o}}^{\mathrm{M}}$  is independent of the choice of the minimal parabolic subgroup  $\mathbf{P}_{0}$ .

### **3** Refined expansions

In this chapter the distribution  $J_{\mathfrak{o}}$  is decomposed as a linear combination of weighted orbital integrals following the methods of Arthur [Ar85] [Ar86].

#### 3.1 Weighted orbital integrals

(3.1.1) **Definition** Let M be a standard Levi subgroup of G. A collection of complexvalued functions

(3.1.1.1) 
$$\left\{ c_{\mathbf{P}} \in C^{\infty}(i\mathfrak{a}_{\mathbf{M}}^{*}) : \mathbf{P} \in \mathcal{P}(\mathbf{M}) \right\}$$

is said to be a (G, M)-family if for each pair of adjacent parabolic subgroups P and P' in  $\mathcal{P}(M)$ , the functions  $c_P$  and  $c_{P'}$  agree on the hyperplane spanned by the common wall of the positive chambers of  $i\mathfrak{a}_M^*$  defined by P and P'.

Let  $(c_{\rm P})$  be a (G, M)-family, define the function  $c_{\rm M}$  on the complement of the coroot hyperplanes in  $i\mathfrak{a}_{\rm M}^*$  by

(3.1.1.2) 
$$\forall \lambda \in i\mathfrak{a}_{\mathrm{M}}^{*} \quad c_{\mathrm{M}}(\lambda) = \sum_{\mathrm{P} \in \mathcal{P}(\mathrm{M})} c_{\mathrm{P}}(\lambda) \theta_{\mathrm{P}}(\lambda)^{-1}$$

where

(3.1.1.3) 
$$\theta_{\mathrm{P}}(\lambda) = \operatorname{Vol}(\mathfrak{a}_{\mathrm{M}}^{\mathrm{G}}/\mathbb{Z}(\Delta_{\mathrm{P}}^{\vee})^{-1}) \prod_{\alpha \in \Delta_{\mathrm{P}}} \langle \alpha^{\check{}}, \lambda \rangle.$$

The function  $c_{\rm M}$  extends smoothly over  $i\mathfrak{a}_{\rm M}^*$ . Denote by  $c_{\rm M}$  its value at the origin of  $i\mathfrak{a}_{\rm M}^*$ .

(3.1.2) Definition Let Q be a parabolic subgroup in  $\mathcal{F}(M)$  with Levi component L, let  $(c_{\rm P})$  be a (G, M)-family. Let

$$(3.1.2.1) i_{\mathbf{Q}}^{\mathbf{G}} : \mathcal{P}^{\mathbf{L}}(\mathbf{M}) \to \mathcal{P}^{\mathbf{G}}(\mathbf{M})$$

be the map that sends a parabolic subgroup P in  $\mathcal{P}^{L}(M)$  to the unique parabolic subgroup in  $\mathcal{P}^{G}(M)$  that is contained in Q whose intersection with L is P. Denote by  $(c_{P}^{Q})$  the (L, M)family

(3.1.2.2) 
$$\left\{c_{i_{\mathbf{Q}}^{\mathbf{G}}(\mathbf{P})}: \mathbf{P} \in \mathcal{P}^{\mathbf{L}}(\mathbf{M})\right\}.$$

Let

$$(3.1.2.3) j_{\mathrm{M}}^{\mathrm{L}} : i\mathfrak{a}_{\mathrm{L}}^{*} \to i\mathfrak{a}_{\mathrm{M}}^{*}$$

be the natural inclusion map. Denote by the  $(c_{\rm P})$  the (G, L)-family

(3.1.2.4) 
$$\left\{ j_{\mathbf{M}}^{\mathbf{L},*}(c_{\mathbf{P}'}): \mathbf{P} \in \mathcal{P}^{\mathbf{G}}(\mathbf{L}) \right\}$$

where P' is a parabolic subgroup in  $\mathcal{P}^{G}(M)$  contained in P, and the function  $j_{M}^{L,*}(c_{P'})$  is independent of the choice of P' in  $\mathcal{P}^{G}(M)$ .

Let  $(c_{\rm P})$  and  $(d_{\rm P})$  be two (G, M)-families, denote by  $((cd)_{\rm P})$  the product of  $(c_{\rm P})$  and  $(d_{\rm P})$ , which is a (G, M)-family. For each parabolic subgroup Q in  $\mathcal{F}(M)$  there exists a function  $c'_{\rm Q}$ on  $i\mathfrak{a}^*_{\rm M}$  such that

(3.1.2.5) 
$$\forall \lambda \in i\mathfrak{a}_{\mathcal{M}}^{*} \quad (cd)_{\mathcal{M}}(\lambda) = \sum_{\mathcal{Q} \in \mathcal{F}(\mathcal{M})} c'_{\mathcal{Q}}(\lambda) d^{\mathcal{Q}}_{\mathcal{M}}(\lambda).$$

Denote by  $c'_{\rm Q}$  the value  $c'_{\rm Q}(0)$ .

(3.1.3) Definition Let M be a standard Levi subgroup of G. A collection of points

(3.1.3.1) 
$$\mathcal{Y}_{\mathrm{M}} = \left\{ Y_{\mathrm{P}} \in \mathfrak{a}_{\mathrm{M}} : \mathrm{P} \in \mathcal{P}(\mathrm{M}) \right\}$$

is said to be a (G, M)-orthogonal set if for each pair of adjacent parabolic subgroups P and P' in  $\mathcal{P}(M)$ , the vector

$$(3.1.3.2) Y_{\rm P} - Y_{\rm P'} \in \mathfrak{a}_{\rm M}$$

is orthogonal to the hyperplane spanned by the common wall of the positive chambers defined by P and P'.

Let  $\alpha_{P'}^{P}$  be the unique coroot that is positive for P and negative for P'. A (G, M)-orthogonal set  $\mathcal{Y}_M$  is *positive* if

(3.1.3.3) 
$$\exists t > 0 \qquad Y_{\rm P} - Y_{\rm P'} = t\alpha_{\rm P'}^{\rm P}.$$

(3.1.4) **Remark** Let  $\mathcal{Y}_M$  be a positive (G, M)-orthogonal set, then the collection of functions

(3.1.4.1) 
$$\left\{ v_{\mathrm{P}}(\mathcal{Y}_{\mathrm{M}})(\lambda) = e^{\langle \lambda, Y_{\mathrm{P}} \rangle} : \mathrm{P} \in \mathcal{P}(\mathrm{M}) \right\}$$

forms a (G, M)-family. The associated constant  $v_{\rm M}(\mathcal{Y}_{\rm M})$  as in (3.1.1.2) is equal to the volume of the convex hull of  $\mathcal{Y}_{\rm M}$  in  $\mathfrak{a}_{\rm M}$ .

(3.1.5) **Definition** Let M be a standard Levi subgroup of G, let x be an element of  $G(\mathbb{Q}_S)$ . The collection of points

(3.1.5.1) 
$$\mathcal{Y}_{\mathcal{M}}(x) = \left\{ -H_{\mathcal{P}}(x) : \mathcal{P} \in \mathcal{P}(\mathcal{M}) \right\}$$

forms a positive (G, M)-orthogonal set. Define the weight factor  $v_{\rm M}(x)$  to be the associated constant

(3.1.5.2) 
$$v_{\rm M}(x) = v_{\rm M}(\mathcal{Y}_{\rm M}(x)).$$

Let X be a point in  $\mathfrak{m}(\mathbb{Q}_S)$  such that  $\mathrm{G}^0_X(\mathbb{Q}_S)$  is contained in  $\mathrm{M}(\mathbb{Q}_S)$ , define the weighted orbital integral  $J^{\mathrm{G}}_{\mathrm{M}}(X, \cdot)$  to be the distribution on  $\mathfrak{g}(\mathbb{Q}_S)$  such that

(3.1.5.3) 
$$\forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$$
$$J_{\mathcal{M}}^{\mathcal{G}}(X, f) = |D^{\mathcal{G}}(X)|_{S}^{1/2} \int_{\mathcal{G}_{X}^{0}(\mathbb{Q}_S) \setminus \mathcal{G}(\mathbb{Q}_S)} f(X \cdot \mathrm{ad}(x)) v_{\mathcal{M}}(x) \mathrm{d}x.$$

For a general element X in  $\mathfrak{m}(\mathbb{Q}_S)$  define the *weighted orbital integral*  $J^{\mathrm{G}}_{\mathrm{M}}(X, \cdot)$  to be the distribution on  $\mathfrak{g}(\mathbb{Q}_S)$  such that

(3.1.5.4) 
$$\forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$$
$$J_{\mathrm{M}}^{\mathrm{G}}(X, f) = \lim_{A \to 0} \sum_{\mathrm{L} \in \mathcal{L}(\mathrm{M})} r_{\mathrm{M}}^{\mathrm{L}}(\exp(X_{\mathrm{nil}}), \exp(A)) J_{\mathrm{L}}^{\mathrm{G}}(X + A, f)$$

where A is a sequence of  $\mathbb{Q}_S$ -points of Lie(A<sub>M</sub>) such that

$$(3.1.5.5) G^0_{X+A}(\mathbb{Q}_S) \subset M(\mathbb{Q}_S)$$

and  $r_{\rm P}^{\rm L}(x, a)$  is an auxiliary (L, M)-family constructed by Arthur in §5 of [Ar88a], see also §2.4 of [HW13] and §III.2 of [Wa95].

(3.1.6) Lemma Let M be a standard Levi subgroup of G. Let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ , let x be an element of  $G(\mathbb{Q}_S)$ . Let f be a Schwartz function on  $\mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$ . Then

(3.1.6.1) 
$$J_{\mathcal{M}}^{\mathcal{G}}(X, f \circ \operatorname{ad}(x)) = \sum_{\mathcal{P} \in \mathcal{F}(\mathcal{M})} J_{\mathcal{M}}^{\mathcal{M}_{\mathcal{P}}}(X, f_{\mathcal{P},x})$$

where  $f_{\mathbf{P},x}$  is the Schwartz function on  $\mathfrak{m}_{\mathbf{P}}(\mathbb{Q}_S)$  defined as

(3.1.6.2) 
$$f_{\mathbf{P},x}(X) = \int_{\mathbf{K}} \int_{\mathfrak{n}_{\mathbf{P}}(\mathbb{A})} f((X+N) \cdot \mathrm{ad}(k)) v'_{\mathbf{P}}(kx) \mathrm{d}N \mathrm{d}k$$

where for each y in  $G(\mathbb{Q}_S)$  and for each parabolic subgroup Q in  $\mathcal{F}(M)$  the constant  $v'_Q(y)$  is the constant  $v'_Q$  intervening in 3.1.2.5 associated to the (G, M)-orthogonal set  $\mathcal{Y}_M(x)$  defined in 3.1.5.1.

Proof. See III.3.(f) of [Wa95].

### 3.2 Nilpotent orbits

(3.2.1) **Definition** Let  $\mathfrak{g}_{nil}$  be the ~ equivalence class of the origin in  $\mathfrak{g}(\mathbb{Q})$ , hence the nilpotent locus of  $\mathfrak{g}$ . Let  $J_{nil}$  and  $J_{nil}^T$  denote respectively  $J_{\mathfrak{g}_{nil}}$  and  $J_{\mathfrak{g}_{nil}}^T$ . Let  $J_{nil}^G$  denote  $J_{nil}$  for convenience in inductive arguments involving Levi subgroups.

(3.2.2) Lemma There exists a continuous seminorm  $\| \|$  on  $S(\mathfrak{g}(\mathbb{A}))$  such that for every truncation parameter T in  $\mathfrak{a}_0$ 

$$(3.2.2.1) \qquad \forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A})) \\ \left| J_{\mathrm{nil}}^{T}(f) - \int_{\mathrm{G}(\mathbb{Q})\backslash\mathrm{G}(\mathbb{A})^{1}} F^{\mathrm{G}}(x,T) \sum_{X \in \mathfrak{g}_{\mathrm{nil}}(\mathbb{Q})} f(X \cdot \mathrm{ad}(x)) \, \mathrm{d}x \right| \leq \|f\| e^{-\frac{\mathrm{d}(T)}{2}}$$

where d(T) denotes the distance from T to the root hyperplanes.

*Proof.* The argument is valid for a general class  $\boldsymbol{o}$  and the sum of all the classes  $\boldsymbol{o}$ , hence contains 2.2.6.1 as a special case. The necessary estimates are established by Chaudouard in the proof of Proposition 4.4 of [Ch02a]. The following is a sketch of the argument, for the details please refer to loc. cit. See also Theorem 3.1 of [Ar85].

Following the first part of the proof of (2.2.2.1), by the combinatorial lemma of Langlands, expand  $J_{\text{nil}}^T(f)$  as a sum indexed by triples of nested standard parabolic subgroups whose leading term corresponding to (G, G, G) is

(3.2.2.2) 
$$\int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^1} F^{\mathcal{G}}(x,T) \sum_{X \in \mathfrak{g}_{\mathrm{nil}}(\mathbb{Q})} f(X \cdot \mathrm{ad}(x)) \, \mathrm{d}x.$$

Hence the left hand side of the inequality in (3.2.2.1) is bounded by

$$(3.2.2.3) \sum_{\substack{\mathrm{P}_1,\mathrm{P}_2,\mathrm{P}_3\in\mathcal{F}\\\mathrm{standard}\\\mathrm{P}_1\subset\mathrm{P}_2\subseteq\mathrm{P}_3}} \int_{\mathrm{P}_1(\mathbb{Q})\backslash\mathrm{G}(\mathbb{A})^1} F^{\mathrm{P}_1}(x,T) \sigma_1^3(H_0(x)-T) \sum_{X\in\mathfrak{m}_1^2(\mathbb{Q})'\cap\mathfrak{o}} \left|\sum_{\overline{Y}\in\tilde{\mathfrak{n}}_2^3(\mathbb{Q})'} \Phi_X(x,\overline{Y})\right| \,\mathrm{d}x$$

where  $\Phi$  denotes the partial Fourier transform of f

(3.2.2.4) 
$$\Phi_X(x,\overline{Y}) = \int_{\mathfrak{n}_2(\mathbb{A})} f((X+N) \cdot \operatorname{ad}(x)) \cdot \psi(\langle N,\overline{Y} \rangle) \, \mathrm{d}N.$$

Changing variables each summand of (3.2.2.3) is bounded by

$$(3.2.2.5) \qquad \sup_{y \in \Gamma} \left( \int_{\mathcal{A}_{1}^{G}(\mathbb{R})_{0}} \int_{\mathcal{A}_{0,T'}^{1,T}(\mathbb{R})} \delta_{0}^{2}(a_{1}a)^{-1} \sigma_{1}^{3}(H_{\mathrm{P}_{1}}(a_{1}) - T) \times \right) \\ \times \sum_{X \in \mathfrak{m}_{1}^{2}(\mathbb{Q})' \cap \mathfrak{o}} \sum_{\overline{Y} \in \overline{\mathfrak{n}}_{2}^{3}(\mathbb{Q})'} |\Phi_{X \cdot \mathrm{ad}(a_{1}a)}(y, \overline{Y} \cdot \mathrm{ad}(a_{1}a))| \, \mathrm{d}a\mathrm{d}a_{1} \right)$$

where  $\Gamma$  is a fixed compact subset of  $M_0(\mathbb{A})^1$  which is independent of the truncation parameter T, the subset  $A_{0,T'}^{1,T}(\mathbb{R})$  of  $A_0^1(\mathbb{R})$  is the T', T-truncated part as in (1.1.3.8), and  $\delta_0^2$  is the modulus function of  $P_2$ .

Let *n* be a natural number, let  $\mathcal{D}$  be an invariant differential operator on  $\mathfrak{g}(\mathbb{R})$  of degree *n*, denote by  $\Phi^{\mathcal{D}}$  the partial Fourier transform of  $\mathcal{D}f$  where  $\mathcal{D}$  operates on *f* via its archimedean component  $f_{\infty}$ , then

$$(3.2.2.6) \left| \Phi_{X \cdot \mathrm{ad}(a_1 a)}(y, \overline{Y} \cdot \mathrm{ad}(a_1 a)) \right| = C(y) \cdot \left| \overline{Y} \cdot \mathrm{ad}(a_1 a) \right|^{-n} \cdot \left| \Phi_{X \cdot \mathrm{ad}(a_1 a)}^{\mathcal{D}}(y, \overline{Y} \cdot \mathrm{ad}(a_1 a)) \right|$$

for some constant C(y) that depends continuously on y. For a Schwartz function f on  $\mathfrak{g}(\mathbb{A})$  define N(f) to be the smallest natural number N such that f is supported on  $\frac{1}{N}\widehat{\mathbb{Z}} \times \mathbb{R}$  where  $\widehat{\mathbb{Z}}$  denotes the profinite completion of  $\mathbb{Z}$ . Let the natural number n be large enough so that

(3.2.2.7) 
$$\sum_{\overline{Y}\in\overline{\mathfrak{n}}_{2}^{3}(\frac{1}{N(f)}\mathbb{Z})'} \left|\overline{Y}\cdot\operatorname{ad}(a_{1}a)\right|^{-n} \leq C\prod_{\alpha\in\Delta_{0}^{3}} e^{-k_{\alpha}\alpha(H_{0}(a_{1}a))}$$

for some constant C and natural numbers  $k_{\alpha}$ . Define the seminorm  $\| \|'$  by

$$(3.2.2.8) ||f||' = \sup_{X \in \mathfrak{g}(\mathbb{A})} |\mathcal{D}f(X)|.$$

A Schwartz function on the  $\mathbb{A}$ -valued points of a rational vector space is bounded by a product of Schwartz functions on each coordinate, hence for Z in  $\mathfrak{p}_2(\mathbb{A})$ 

(3.2.2.9) 
$$|\mathcal{D}f(Z \cdot \mathrm{ad}(y))| \leq \left(\prod_{\mu \in \Phi_0 - \Phi_2} \phi_{\mu}(Z_{\mu})\right) \phi_{\mathfrak{n}_2}(Z_{\mathfrak{n}_2})$$

where  $Z_{\mu}$  and  $Z_{\mathfrak{n}_2}$  are the components of Z on the weight space of  $\mu$  and  $\mathfrak{n}_2$ , and  $\phi_{\bullet}$  are positive Schwartz functions. If  $a_0$  is an element of  $A_0(\mathbb{R})$ , denote by  $\Psi(a_0)$  for the sum

(3.2.2.10) 
$$\Psi(a_0) = \sum_{X \in \mathfrak{m}_1^2(\frac{1}{N(f)}\mathbb{Z})' \cap \mathfrak{o}} \left(\prod_{\mu \in \Phi_0 - \Phi_2} \phi_\mu(\mu(a_0)^{-1}X_\mu)\right).$$

Then (3.2.2.5) is bounded by

(3.2.2.11) 
$$\sup_{y \in \Gamma} C(y) \int_{A_{1}^{G}(\mathbb{R})} \int_{A_{0,T'}^{1,T}(\mathbb{R})} \left( \delta_{0}^{2}(a_{1}a)^{-1} \sigma_{1}^{2}(H_{P_{1}}(a_{1}) - T) \times \Psi(a_{1}a) \cdot C \prod_{\alpha \in \Delta_{0}^{2}} e^{-k_{\alpha}\alpha(H_{0}(a_{1}a))} \right) dada_{1},$$

which only depends on the Schwartz function f via its componentwise bounds  $\phi_{\bullet}$  and the lattice  $\overline{\mathfrak{n}}_{2}^{3}(\frac{1}{N(f)}\mathbb{Z})$ , hence is proportional to the seminorm  $\| \|$  defined by

(3.2.2.12) 
$$\forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A})) \qquad \|f\| = \sup_{y \in \Gamma} C(y) \cdot C \cdot \|f\|' \cdot N(f)^n.$$

The constant  $\sup_{y \in \Gamma} C(y) \cdot C$  is independent of both the Schwartz function f and the truncation parameter T.

Substituting the definition of the seminorm  $\| \|$ , the majorant (3.2.2.11) reduces to

(3.2.2.13) 
$$\|f\| \cdot \operatorname{Vol}(\mathcal{A}_{0,T'}^{1,T}(\mathbb{R})) \prod_{\alpha \in \Delta_1^3} \left( e^{-\alpha(T)} \int_0^\infty p(t_\alpha) e^{-t_\alpha} \, \mathrm{d}t_\alpha \right)$$

where p(t) is a polynomial. Because

(3.2.2.14) 
$$\prod_{\alpha \in \Delta_1^3} e^{-\alpha(T)} \int_0^\infty p(t) e^{-t} \, \mathrm{d}t \leq e^{-\mathrm{d}(T)},$$

and  $\operatorname{Vol}(A^{1,T}_{0,T'}(\mathbb{R}))$  is of polynomial growth in T, the majorant (3.2.2.13) reduces to

$$(3.2.2.15) ||f||e^{-\frac{\mathrm{d}(T)}{2}}$$

	_		

(3.2.3) **Definition** Let v be a place of  $\mathbb{Q}$ , let  $\beta_v$  be a bump function on  $\mathbb{Q}_v$ , let  $\nu$  be a G-orbit contained in  $\mathfrak{g}_{nil}$ , let  $\{p_1, \ldots, p_l\}$  be a collection of polynomials rational with rational coefficients cutting out the Zariski closure  $\overline{\nu}$ . Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , let  $\epsilon$  be a positive real number. Define the *truncated function*  $f_{\nu,\nu}^{\epsilon}$  on  $\mathfrak{g}(\mathbb{A})$  as in [Ar85] by

$$(3.2.3.1) \qquad \forall X \in \mathfrak{g}(\mathbb{A}) \qquad f_{\nu,\nu}^{\epsilon}(X) = f(X)\beta_{\nu}(\epsilon^{-1}|p_1(X)|_{\nu})\dots\beta_{\nu}(\epsilon^{-1}|p_l(X)|_{\nu}).$$

(3.2.4) Lemma Let the place v, the bump function  $\beta_v$ , the orbit  $\nu$  and the polynomials  $\{p_1, \ldots, p_l\}$  be as in (3.2.3), then there exists a natural number m and another seminorm  $\|\|_1$  for which the inequality (3.2.2.1) holds such that

(3.2.4.1) 
$$\forall \epsilon \text{ such that } 0 < \epsilon < 1 \ \forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A})) \qquad \|f_{\nu,\nu}^{\epsilon}\| \le \epsilon^{-ml} \|f\|_{1}.$$

*Proof.* It is enough to consider the case when there is a single polynomial p. There are two cases:

• If v is the archimedean place then

(3.2.4.2) 
$$N(f_{\nu,\nu}^{\epsilon}) = N(f),$$

so by properties of the derivative

(3.2.4.3)  $||f_{\nu,\nu}^{\epsilon}||' \leq \epsilon^{-m} C ||f||'.$ 

Define  $||f||'_1$  to be C||f||.

• If v is finite then

(3.2.4.4) 
$$\|f_{\nu,\nu}^{\epsilon}\|' = \|f\|',$$

 $\mathbf{SO}$ 

$$(3.2.4.5) N(f_{\nu,\nu}^{\epsilon}) \leq \epsilon^{-m} N(f)$$

since N(f) only depends on the support of  $f_{\text{finite}}$  and the assignment

(3.2.4.6) 
$$f \mapsto f_{\nu,v}^{\epsilon} = f \prod \beta_v(\epsilon^{-1}|p|_v)$$

shrinks the support of f by a factor of  $\epsilon^m$ , up to a multiplicative constant. Define ||f||' to be ||f||.

Hence (3.2.4.1) follows since

$$(3.2.4.7) ||f|| = ||f||' N(f).$$

(3.2.5) Lemma Let the place v, the bump function  $\beta_v$ , the orbit  $\nu$  and the polynomials  $\{p_1, \ldots, p_l\}$  be as in (3.2.3), then there exists a positive real number r such that

$$(3.2.5.1) \quad \forall \epsilon > 0 \ \forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A})) \\ \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^1} F^{\mathcal{G}}(x,T) \sum_{X \in \mathfrak{g}(\mathbb{Q}) - \overline{\nu}(\mathbb{Q})} \left| f^{\epsilon}_{\nu,\nu}(X \cdot \mathrm{ad}(x)) \right| \, \mathrm{d}x \le \|f\|\epsilon^r (1+|T|)^{d_0}$$

where  $d_0$  is dim(A<sub>0</sub>/A<sub>G</sub>), the split rank of G.

*Proof.* This is Lemma 4.1 of [Ar85].

(3.2.6) **Proposition** Let T be a truncation parameter. For each nilpotent orbit  $\nu$  there exists a distribution  $J_{\nu}^{T}$  on  $\mathfrak{g}(\mathbb{A})$  such that for each Schwartz function f on  $\mathfrak{g}(\mathbb{A})$ , the expression  $J_{\nu}^{T}(f)$  is a polynomial in T of degree at most  $d_{0}$ , and

(3.2.6.1) 
$$J_{\text{nil}}^T(f) = \sum_{\substack{\nu \subset \mathfrak{g}_{\text{nil}} \\ orbit}} J_{\nu}^T(f).$$

There exists a continuous seminorm  $\| \|$  on  $\mathcal{S}(\mathfrak{g}(\mathbb{A}))$  and a positive real number  $\epsilon$  such that

(3.2.6.2) 
$$\left| \begin{aligned} \forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A})) \\ \left| J_{\nu}^{T}(f) - \int_{\mathrm{G}(\mathbb{Q})\backslash\mathrm{G}(\mathbb{A})^{1}} F^{\mathrm{G}}(x,T) \sum_{X \in \nu(\mathbb{Q})} f(X \cdot \mathrm{ad}(x)) \, \mathrm{d}x \right| \leq \|f\| e^{-\epsilon \mathrm{d}(T)}. \end{aligned} \right|$$

*Proof.* Define the polynomials recursively by the formula

(3.2.6.3) 
$$J_{\overline{\nu}}^T(f) = \lim_{\epsilon \to 0} J_{\mathrm{nil}}^T(f_{\nu,\nu}^\epsilon)$$

where  $J_{\overline{\nu}}^T$  denotes the sum of  $J_{\nu'}^T$  for those orbits  $\nu'$  contained in  $\overline{\nu}$ .

The limit of  $J_{\text{nil}}^T(f_{\nu,v}^{\epsilon})$  as  $\epsilon$  approaches 0 exists because

(3.2.6.4) 
$$\sum_{X \in \overline{\nu}(\mathbb{Q})} f(X \cdot \operatorname{ad}(x)) = \sum_{X \in \overline{\nu}(\mathbb{Q})} f_{\nu,v}^{\epsilon}(X \cdot \operatorname{ad}(x))$$

implies that

$$(3.2.6.5) \qquad \left| J_{\mathrm{nil}}^{T}(f_{\nu,\nu}^{\epsilon}) - \int_{\mathrm{G}(\mathbb{Q})\backslash\mathrm{G}(\mathbb{A})^{1}} F^{\mathrm{G}}(x,T) \sum_{X\in\overline{\nu}(\mathbb{Q})} f(X\cdot\mathrm{ad}(x)) \,\mathrm{d}x \right| \\ \leq \left| J_{\mathrm{nil}}^{T}(f_{\nu,\nu}^{\epsilon}) - \int_{\mathrm{G}(\mathbb{Q})\backslash\mathrm{G}(\mathbb{A})^{1}} F^{\mathrm{G}}(x,T) \sum_{X\in\mathfrak{g}_{\mathrm{nil}}(\mathbb{Q})} f_{\nu,\nu}^{\epsilon}(X\cdot\mathrm{ad}(x)) \,\mathrm{d}x \right| + \\ + \int_{\mathrm{G}(\mathbb{Q})\backslash\mathrm{G}(\mathbb{A})^{1}} F^{\mathrm{G}}(x,T) \sum_{X\in\mathfrak{g}_{\mathrm{nil}}(\mathbb{Q})-\overline{\nu}(\mathbb{Q})} \left| f_{\nu,\nu}^{\epsilon}(X\cdot\mathrm{ad}(x)) \right| \,\mathrm{d}x.$$

By (3.2.2.1) the first summand of the right hand side of (3.2.6.5) is bounded by

(3.2.6.6) 
$$\|f_{\nu,\nu}^{\epsilon}\|e^{-\frac{\mathrm{d}(T)}{2}} \leq \epsilon^{-lm} \cdot \|f\|_1 \cdot e^{-\frac{\mathrm{d}(T)}{2}}$$

which follows from (3.2.4.1).

The second summand of the right hand side of (3.2.6.5) is bounded by summing over the complement of  $\overline{\nu}(\mathbb{Q})$  in  $\mathfrak{g}(\mathbb{Q})$  instead of in  $\mathfrak{g}_{nil}(\mathbb{Q})$ , hence by (3.2.5.1) is bounded by

(3.2.6.7) 
$$||f||_1 \cdot \epsilon^r \cdot (1+|T|)^{d_0}.$$

Therefore the right hand side of (3.2.6.5) is bounded by

(3.2.6.8) 
$$||f||_1 \left( \epsilon^{-lm} e^{-\frac{\mathrm{d}(T)}{2}} + \epsilon^r (1+|T|)^{d_0} \right).$$

It suffices to take  $\delta^n$  as  $\epsilon$  with  $\delta$  bounded strictly between 0 and 1 and n a sequence natural numbers approaching infinity. Then the majorant (3.2.6.8) satisfies the inequality

$$(3.2.6.9) ||f||_1 \left( e^{|\log(\delta)|lmn - \frac{d(T)}{2}} + \delta^{rn} (1 + |T|)^{d_0} \right) \leq ||f|| \cdot \delta^{rn} \cdot (1 + |T|)^{d_0}$$

provided d(T) is bounded below by  $C|\log(\delta)|n$  for some constant C and || || is another continuous seminorm with the same properties.

Therefore for every natural number n and every T in the open subcone

(3.2.6.10) 
$$\left\{ T : d(T) > C |\log(\delta)|(n+1) \right\}$$

of the positive chamber, the following inequality holds

$$(3.2.6.11) |J_{\rm nil}^T(f_{\nu,\nu}^{\delta^n}) - J_{\rm nil}^T(f_{\nu,\nu}^{\delta^{n+1}})| \leq 2||f||(1+|T|)^{d_0}\delta^{rn}.$$

The left hand side is a polynomial in T of degree at most  $d_0$ , hence the polynomial extrapolation lemma in Lemma 5.2 of [Ar82] applies:

There exists a constant A such that for all T,

$$(3.2.6.12) \quad |J_{\rm nil}^T(f_{\nu,\nu}^{\delta^n}) - J_{\rm nil}^T(f_{\nu,\nu}^{\delta^{n+1}})| \leq A \cdot ||f|| \cdot (1+|T|)^{d_0} \cdot (|\log(\delta)|(n+1))^{d_0} \cdot \delta^{rn}.$$

Since

(3.2.6.13) 
$$\sum_{n=0}^{\infty} \left( |\log(\delta)|(n+1) \right)^{d_0} \delta^{rn} < \infty,$$

by telescoping the series

(3.2.6.14) 
$$\sum_{n=0}^{\infty} J_{\text{nil}}^{T}(f_{\nu,\nu}^{\delta^{n}}) - J_{\text{nil}}^{T}(f_{\nu,\nu}^{\delta^{n+1}})$$

the sequence  $J_{\text{nil}}^T(f_{\nu,v}^{\delta^n})$  has a limit as *n* approaches infinity. The limit is a polynomial in *T* of degree at most  $d_0$ , denoted by  $J_{\overline{\nu}}^T(f)$ .

By construction

$$(3.2.6.15) \qquad \left| J_{\overline{\nu}}^{T}(f) - \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} F^{\mathcal{G}}(x,T) \sum_{X \in \overline{\nu}(\mathbb{Q})} f(X \cdot \operatorname{ad}(x)) \, \mathrm{d}x \right| \\ \leq \left| J_{\operatorname{nil}}^{T}(f_{\nu,\nu}^{\delta^{n}}) - J_{\overline{\nu}}^{T}(f) \right| + \left| J_{\operatorname{nil}}^{T}(f_{\nu,\nu}^{\delta^{n}}) - \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} F^{\mathcal{G}}(x,T) \sum_{X \in \overline{\nu}(\mathbb{Q})} f(X \cdot \operatorname{ad}(x)) \, \mathrm{d}x \right| \\ (3.2.6.16) \leq \sum_{n=0}^{\infty} A \|f\| (1+|T|)^{d_{0}} \big( |\log(\delta)|(n+1) \big)^{d_{0}} \delta^{rn} + \|f\| \delta^{rn} (1+|T|)^{d_{0}}$$

where (3.2.6.16) holds for a fixed  $\delta$ , a sufficiently positive T and n the largest natural number such that

$$(3.2.6.17) d(T) \geq C |\log(\delta)| n.$$

Therefore it is possible choose a new seminorm  $\| \|$  and a new constant  $\epsilon > 0$  such that

$$(3.2.6.18) \quad \left| J_{\overline{\nu}}^T(f) - \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^1} F^{\mathcal{G}}(x,T) \sum_{X \in \overline{\nu}(\mathbb{Q})} f(X \cdot \operatorname{ad}(x)) \, \mathrm{d}x \right| \leq ||f|| e^{-\epsilon \operatorname{d}(T)}.$$

The corresponding statements for  $\nu$  instead of  $\overline{\nu}$  follow recursively by setting

(3.2.6.19) 
$$J_{\overline{\nu}}^T(f) = \sum_{\substack{\nu' \subset \overline{\nu} \\ \text{orbit}}} J_{\nu'}^T(f).$$

(3.2.7) Proposition Let S be a finite set of places of  $\mathbb{Q}$  containing the archimedean place, let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$  such that  $f_p$  is the characteristic function of the standard lattice  $\mathfrak{g}(\mathbb{Z}_p)$  whenever p is not contained in S. Let M be a standard Levi subgroup of G. Denote by  $\mathfrak{m}_{nil}(\mathbb{Q})_{M,S}$  the set of  $M(\mathbb{Q}_S)$ -conjugacy classes in  $\mathfrak{m}_{nil}(\mathbb{Q})$ . For each nilpotent conjugacy class  $\nu$  in  $\mathfrak{m}_{nil}(\mathbb{Q})_{M,S}$  here exists a constant  $a^M(S, \nu)$  such that

(3.2.7.1) 
$$J_{\rm nil}^{\rm G}(f) = \sum_{{\rm M}\in\mathcal{L}} |{\rm W}_0^{\rm M}| |{\rm W}_0^{\rm G}|^{-1} \sum_{\nu\in\mathfrak{m}_{\rm nil}(\mathbb{Q})_{{\rm M},S}} a^{\rm M}(S,\nu) J_{\rm M}^{\rm G}(\nu,f_S).$$

**Remark** By Lemma 7.1 of [Ar85], the sets  $\mathfrak{m}_{nil}(\mathbb{Q})_{M,S}$  and  $\mathfrak{m}_{nil}(\mathbb{Q}_S)/M(\mathbb{Q}_S)^1$  are in natural bijection, hence the expression  $J_M^G(\nu, f_S)$  makes sense.

*Proof.* The argument is based on the following two lemmas:

**Lemma** For every x in  $G(\mathbb{A})^1$ 

(3.2.7.2) 
$$J_{\rm nil}^{\rm G}(f \circ {\rm ad}(x)) = \sum_{{\rm Q} \in \mathcal{F}} |{\rm W}_0^{\rm M_{\rm Q}}| |{\rm W}_0^{\rm G}|^{-1} J_{\rm nil}^{\rm M_{\rm Q}}(f_{{\rm Q},x}),$$

where the function  $f_{Q,x}$  on  $\mathfrak{m}_Q(\mathbb{A})$  is defined as in (2.2.12.2).

*Proof.* This is a special case of (2.2.12.1).

**Lemma** Let M be a standard Levi subgroup. For every point x in  $G(\mathbb{Q}_S)^1$  and  $\nu$  in  $\mathfrak{m}_{nil}(\mathbb{Q}_S)/M(\mathbb{Q}_S)^1$ 

(3.2.7.3) 
$$J_{M}^{G}(\nu, f_{S} \circ ad(x)) = \sum_{Q \in \mathcal{F}(M)} J_{M}^{M_{Q}}(\nu, f_{S,Q,x}).$$

*Proof.* This is a special case of (3.1.6.1).

Argue by induction. Assume that the constants  $a^{L}(S,\nu)$  and the identities

(3.2.7.4) 
$$J_{\rm nil}^{\rm L}(f) = \sum_{{\rm M}\in\mathcal{L}^{\rm L}} |{\rm W}_0^{\rm M}| |{\rm W}_0^{\rm L}|^{-1} \sum_{\nu\in\mathfrak{m}_{\rm nil}(\mathbb{Q})_{{\rm M},S}} a^{\rm M}(S,\nu) J_{\rm M}^{\rm L}(\nu,f_S)$$

are known for every proper Levi subgroup L of G.

Define the distribution  $T^{\mathrm{G}}$  on  $\mathfrak{g}(\mathbb{Q}_S)$  by

$$(3.2.7.5) \qquad \forall f_S \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$$
$$T^{\mathrm{G}}(f_S) = J^{\mathrm{G}}_{\mathrm{nil}} \Big( f_S \otimes \bigotimes_{p \notin S} \mathbb{I}_{\mathfrak{g}(\mathbb{Z}_p)} \Big) - \sum_{\substack{\mathrm{M} \in \mathcal{L} \\ \mathrm{M} \neq \mathrm{G}}} |\mathrm{W}^{\mathrm{M}}_0| |\mathrm{W}^{\mathrm{G}}_0|^{-1} \times$$
$$\times \sum_{\nu \in \mathfrak{m}_{\mathrm{nil}}(\mathbb{Q})_{\mathrm{M},S}} a^{\mathrm{M}}(S,\nu) J^{\mathrm{G}}_{\mathrm{M}}(\nu, f_S).$$

The distribution  $T^{\mathrm{G}}$  is supported on  $\mathfrak{g}_{\mathrm{nil}}(\mathbb{Q}_S)$ . By (3.2.7.2) and (3.2.7.3)

$$(3.2.7.6) \qquad T^{\mathcal{G}}(f_{S} \circ \operatorname{ad}(x)) - T^{\mathcal{G}}(f_{S})$$

$$= \left(\sum_{Q \in \mathcal{F}} |W_{0}^{\mathcal{M}_{Q}}| |W_{0}^{\mathcal{G}}|^{-1} J_{\operatorname{nil}}^{\mathcal{M}_{Q}} \left( (f_{S} \otimes \bigotimes_{p \notin S} \mathbb{I}_{\mathfrak{g}(\mathbb{Z}_{p})})_{Q,x} \right) \right)$$

$$- \sum_{\substack{M \in \mathcal{L} \\ M \neq G}} \sum_{Q \in \mathcal{F}(M)} |W_{0}^{\mathcal{M}}| |W_{0}^{\mathcal{G}}|^{-1} \sum_{\nu \in \mathfrak{m}_{\operatorname{nil}}(\mathbb{Q})_{\mathcal{M},S}} a^{\mathcal{M}}(S,\nu) J_{\mathcal{M}}^{\mathcal{M}_{Q}}(\nu, f_{S,Q,x}) \right)$$

$$- \left( J_{\operatorname{nil}}^{\mathcal{G}} \left( f_{S} \otimes \bigotimes_{p \notin S} \mathbb{I}_{\mathfrak{g}(\mathbb{Z}_{p})} \right) - \sum_{\substack{M \in \mathcal{L} \\ M \neq G}} |W_{0}^{\mathcal{M}}| |W_{0}^{\mathcal{G}}|^{-1} \times \sum_{\nu \in \mathfrak{m}_{\operatorname{nil}}(\mathbb{Q})_{\mathcal{M},S}} a^{\mathcal{M}}(S,\nu) J_{\mathcal{M}}^{\mathcal{G}}(\nu, f_{S}) \right)$$

$$= \sum_{\substack{Q \in \mathcal{F} \\ Q \neq G}} |W_{0}^{\mathcal{M}_{Q}}| |W_{0}^{\mathcal{G}}|^{-1} \left( J_{\operatorname{nil}}^{\mathcal{M}_{Q}} \left( (f_{S} \otimes \bigotimes_{p \notin S} \mathbb{I}_{\mathfrak{g}(\mathbb{Z}_{p})})_{Q,x} \right) - \sum_{\substack{M \in \mathcal{L}} M^{\mathcal{M}_{Q}}} |W_{0}^{\mathcal{M}}| |W_{0}^{\mathcal{M}}|^{-1} \sum_{\nu \in \mathfrak{m}_{\operatorname{nil}}(\mathbb{Q})_{\mathcal{M},S}} a^{\mathcal{M}}(S,\nu) J_{\mathcal{M}}^{\mathcal{M}_{Q}}(\nu, f_{S,Q,x}) \right)$$

which vanishes by the inductive hypothesis that  $T^{L}$  is invariant applied to the Levi subgroup  $M_Q$  with Q a proper parabolic subgroup of G. Therefore  $T^G$  is invariant under the action of  $\mathcal{G}(\mathbb{Q}_S)^1$ .

Construct the constants  $a^{G}(S,\nu)$  subject to the new identity

(3.2.7.7) 
$$T^{\mathcal{G}}(f_S) = \sum_{\nu \in \mathfrak{g}_{\mathrm{nil}}(\mathbb{Q})_{\mathcal{G},S}} a^{\mathcal{G}}(S,\nu) J^{\mathcal{G}}_{\mathcal{G}}(\nu,f_S).$$

Stratify the nilpotent locus  $\mathfrak{g}_{nil}(\mathbb{Q}_S)$  equivariantly by codimension. More precisely define for each natural number d the open set

(3.2.7.8) 
$$\mathfrak{g}_{\operatorname{nil},d}(\mathbb{Q}_S) = \bigcup_{\substack{\nu \in \mathfrak{g}_{\operatorname{nil}}(\mathbb{Q})_{\mathrm{G},S} \\ \operatorname{codim}(\nu) \leq d}} \nu(\mathbb{Q}_S).$$

Denote by  $T_d^G$  the distribution obtained by restricting  $T^G$  to  $\mathfrak{g}_{\operatorname{nil},d}(\mathbb{Q}_S)$ . The open set  $\mathfrak{g}_{\operatorname{nil},0}(\mathbb{Q}_S)$  is the regular nilpotent orbit. Since  $T_0^G$  is invariant it is equal to a multiple of  $J_{\rm G}^{\rm G}(\nu_{\rm reg})$ , where  $\nu_{\rm reg}$  denotes the regular nilpotent orbit. Define  $a^{\rm G}(S, \nu_{\rm reg})$  to be the constant of proportionality.

The other constants  $a^{G}(S,\nu)$  are constructed by induction on the codimension d which ranges among  $0, 1, 2, \ldots, \dim(\mathfrak{g}_{nil})$ . Let  $T^{G,d}$  be the distribution on the complement of  $\mathfrak{g}_{\operatorname{nil},d-1}(\mathbb{Q}_S)$  in  $\mathfrak{g}_{\operatorname{nil}}(\mathbb{Q}_S)$  defined by

(3.2.7.9) 
$$T^{\mathrm{G},d}(f_S) = T^{\mathrm{G}}(f_S) - \sum_{\substack{\nu \in \mathfrak{g}_{\mathrm{nil}}(\mathbb{Q})_{\mathrm{G},S} \\ \mathrm{codim}(\nu) < d}} a^{\mathrm{G}}(S,\nu) J^{\mathrm{G}}_{\mathrm{G}}(\nu,f_S).$$

Denote by  $T_d^{\mathrm{G},d}$  its restriction to the complement of  $\mathfrak{g}_{\mathrm{nil},d-1}(\mathbb{Q}_S)$  in  $\mathfrak{g}_{\mathrm{nil},d}(\mathbb{Q}_S)$ .

Since the complement of  $\mathfrak{g}_{\operatorname{nil},d-1}(\mathbb{Q}_S)$  in  $\mathfrak{g}_{\operatorname{nil},d}(\mathbb{Q}_S)$  has an open partition by the nilpotent orbits of codimension d, and the distribution  $T_d^{\mathrm{G},d}$  is invariant, there exist constants  $a^{\mathrm{G}}(S,\nu)$ , one for each  $\nu$  of codimension d, such that

(3.2.7.10) 
$$T_d^{\mathrm{G},d}(f_S) = \sum_{\substack{\nu \in \mathfrak{g}_{\mathrm{nil}}(\mathbb{Q})_{\mathrm{G},S} \\ \mathrm{codim}(\nu) = d}} a^{\mathrm{G}}(S,\nu) J_{\mathrm{G}}^{\mathrm{G}}(\nu,f_S).$$

The constants  $a^{\rm G}(S,\nu)$  are required to satisfy (3.2.7.7), which is equivalent to

$$(3.2.7.11) \quad T^{\mathrm{G}}(f_S) = T_0^{\mathrm{G},0}(f_S) + T_1^{\mathrm{G},1}(f_S) + T_2^{\mathrm{G},2}(f_S) + \dots + T_{\mathrm{dim}(\mathfrak{g}_{\mathrm{nil}})}^{\mathrm{G},\mathrm{dim}(\mathfrak{g}_{\mathrm{nil}})}(f_S).$$

Let  $v \in S$  be a place of  $\mathbb{Q}_S$ , let  $\nu$  be a nilpotent orbit, define the function  $f_{S,\nu,v}^{\epsilon}$  by the same formula (3.2.3.1) as for  $f_{\nu,v}^{\epsilon}$ . Let  $\nu_d$  denote the complement of  $\mathfrak{g}_{\operatorname{nil},d-1}$  in  $\mathfrak{g}_{\operatorname{nil},d}$ , the union of the nilpotent orbits  $\nu$  of codimension d. Then the expression  $f_{S,\nu_d,v}^{\epsilon}$  makes sense, and

(3.2.7.12) 
$$T_{d}^{G,d}(f_{S}) = \lim_{\epsilon \to 0} T^{G}(f_{S,\nu_{d},v}^{\epsilon}) - \lim_{\epsilon \to 0} T^{G}(f_{S,\nu_{d+1},v}^{\epsilon}).$$

Therefore

$$(3.2.7.13) \ T_{0}^{G,0}(f_{S}) + T_{1}^{G,1}(f_{S}) + T_{2}^{G,2}(f_{S}) + \dots + T_{\dim(\mathfrak{g}_{\operatorname{nil}})}^{G,\dim(\mathfrak{g}_{\operatorname{nil}})}(f_{S})$$

$$= \lim_{\epsilon \to 0} \left( \underbrace{T^{G}(f_{S,\nu_{0},v}^{\epsilon}) - T^{G}(f_{S,\nu_{1},v}^{\epsilon}) + T^{G}(f_{S,\nu_{1},v}^{\epsilon}) - T^{G}(f_{S,\nu_{2},v}^{\epsilon}) + T^{G}(f_{S,\nu_{2},v}^{\epsilon}) - \dots \right)$$

$$\dots - T^{G}(f_{S,\nu_{\dim(\mathfrak{g}_{\operatorname{nil}})},v}) + T^{G}(f_{S,\nu_{\dim(\mathfrak{g}_{\operatorname{nil}})},v}^{\epsilon}) - T^{G}(\underbrace{f_{S,\nu_{\dim(\mathfrak{g}_{\operatorname{nil}})+1},v}^{\epsilon}}_{\operatorname{this is void}}) \right)$$

$$= T^{G}(f_{S}).$$

(3.2.8) **Remark** If  $\nu$  is the orbit consisting of the origin, the coefficient  $a^{M}(S,\nu)$  is independent of S and equal to the corresponding Tamagawa number

(3.2.8.1) 
$$a^{\mathcal{M}}(S,0) = \operatorname{Vol}(\mathcal{M}(\mathbb{Q}) \setminus \mathcal{M}(\mathbb{A})^{1}).$$

#### 3.3 General orbits

(3.3.1) Definition Let  $\mathfrak{o}$  be a ~ equivalence class on  $\mathfrak{g}(\mathbb{Q})$ . Let  $P_1$  be a parabolic subgroup of G, let  $M_1$  be the Levi component of  $P_1$  which is standard. Fix a semisimple element  $\Sigma$  in  $\mathfrak{o}$  such that  $\Sigma$  is contained in the Levi subalgebra  $\mathfrak{m}_1$ , but not in any proper parabolic subalgebra of  $\mathfrak{p}_1$ . The group  $P_{1,\Sigma}^0$  is a minimal parabolic subgroup of  $G_{\Sigma}^0$  with minimal Levi component  $M_{1,\Sigma}^0$ . Denote by  $\mathcal{F}^{\Sigma}$  the set of parabolic subgroups of  $G_{\Sigma}^0$  containing  $M_{1,\Sigma}^0$ . A parabolic subgroup Q in  $\mathcal{F}^{\Sigma}$  is said to be standard if Q contains  $P_{1,\Sigma}^0$ . Fix a maximal compact subgroup  $K_{\Sigma}$  of  $G_{\Sigma}^0(\mathbb{A})$  that is admissible with respect to  $M_{1,\Sigma}^0$  such that for each parabolic subgroup Q in  $\mathcal{F}^{\Sigma}$  there is the associated function

$$(3.3.1.1) H_{\mathbf{Q}}: \quad \mathbf{G}^{0}_{\Sigma}(\mathbb{A}) \to \mathfrak{a}_{\mathbf{Q}}$$

There is a unique point  $T_{\Sigma,1}$  in  $\mathfrak{a}_1$  modulo  $\mathfrak{a}_{G_{\Sigma}^0}$  defined in the same manner as  $T_0$  in  $\mathfrak{a}_0^G$  that satisfies an identity analogous to (2.2.8.1). Let L be a Levi subgroup of  $G_{\Sigma}^0$  containing  $M_{1,\Sigma}^0$ , denote by  $W_1^L$  the Weyl group of L with respect to the split torus  $A_1$ .

Let  $\pi_{\Sigma}$  be the surjection from  $\mathcal{F}(M_1)$  onto  $\mathcal{F}^{\Sigma}$  defined by

(3.3.1.2) 
$$\forall \mathbf{P} \in \mathcal{F}(\mathbf{M}_1) \qquad \pi_{\Sigma}(\mathbf{P}) = \mathbf{P}_{\Sigma}^0.$$

Let Q be a parabolic subgroup in  $\mathcal{F}^{\Sigma}$ . Let  $\mathcal{F}_Q(M_1)$  be the inverse image  $\pi_{\Sigma}^{-1}(Q)$ . Define subsets  $\mathring{\mathcal{F}}_Q(M_1)$  and  $\overline{\mathcal{F}}_Q(M_1)$  of  $\mathcal{F}(M_1)$  by

(3.3.1.3) 
$$\hat{\mathcal{F}}_{Q}(M_{1}) = \left\{ P \in \mathcal{F}(M_{1}) : \pi_{\Sigma}(P) = Q, \ \mathfrak{a}_{P} = \mathfrak{a}_{Q} \right\}$$
$$\bar{\mathcal{F}}_{Q}(M_{1}) = \left\{ P \in \mathcal{F}(M_{1}) : \pi_{\Sigma}(P) \supset Q \right\}.$$

There are inclusions

(3.3.1.4) 
$$\forall Q \in \mathcal{F}^{\Sigma} \quad \mathring{\mathcal{F}}_{Q}(M_{1}) \subset \mathcal{F}_{Q}(M_{1}) \subset \overline{\mathcal{F}}_{Q}(M_{1}) \subset \mathcal{F}(M_{1}).$$

(3.3.2) Definition Let  $P_1$ ,  $M_1$ ,  $\mathfrak{o}$ ,  $\Sigma$  be as in (3.3.1). Let Q be a parabolic subgroup in  $\mathcal{F}^{\Sigma}$ . Let  $\mathcal{Y}$  be a collection of points

(3.3.2.1) 
$$\mathcal{Y} = \left\{ Y_{\mathrm{P}} \in \mathfrak{a}_{0} : \mathrm{P} \in \mathring{\mathcal{F}}_{\mathrm{Q}}(\mathrm{M}_{1}) \right\}$$

satisfying the compatibility conditions defining a (G, M<sub>1</sub>)-orthogonal set in (3.1.3), namely that for each pair of adjacent parabolic subgroups P and P' in  $\mathring{\mathcal{F}}_Q(M_1)$  the difference between  $Y_P$  and  $Y_{P'}$  is orthogonal to the common wall of the positive chambers defined by P and P'.

The collection  $\mathcal{Y}$  has a unique extension from  $\mathring{\mathcal{F}}_Q(M_1)$  to  $\overline{\mathcal{F}}_Q(M_1)$ . Let Q' be a parabolic subgroup in  $\mathcal{F}^{\Sigma}$  containing Q, denote by  $\mathcal{Y}_{Q'}$  the collection of points

(3.3.2.2) 
$$\mathcal{Y}_{Q'} = \left\{ Y_{P} : P \in \mathcal{F}_{Q'}(M_1) \right\}.$$

Let  $Q_3$  be a parabolic subgroup in  $\mathcal{F}^{\Sigma}$ . Define the gamma' function  $\Gamma'_{Q_3}(, \mathcal{Y}_{Q_3})$  on  $\mathfrak{a}_1$  by (3.3.2.3)  $\forall H \in \mathfrak{a}_1$  $\Gamma'_{Q_3}(H, \mathcal{Y}_{Q_3}) = \sum_{\substack{Q_4 \in \mathcal{F}^{\Sigma} \\ Q_4 \supset Q_4}} \tau_3^4(H) \left(\sum_{P \in \mathcal{F}_{Q_4}(M_1)} (-1)^{\dim(A_P/A_G)} \hat{\tau}_P(H - Y_P)\right).$ 

(3.3.3) **Remark** The function  $\Gamma'_{Q_3}(, \mathcal{Y}_{Q_3})$  factorizes through the projection from  $\mathfrak{a}_1$  onto  $\mathfrak{a}_3^G$  and depends continuously on  $\mathcal{Y}$ . For each parabolic subgroup  $Q_2$  in  $\mathcal{F}^{\Sigma}$ 

$$(3.3.3.1) \sum_{\mathbf{P}\in\mathcal{F}_{\mathbf{Q}_{2}}(\mathbf{M}_{1})} (-1)^{\dim(\mathbf{A}_{\mathbf{P}}/\mathbf{A}_{\mathbf{G}})} \hat{\tau}_{\mathbf{P}}(H-Y_{\mathbf{P}}) = \sum_{\substack{\mathbf{Q}_{3}\in\mathcal{F}^{\Sigma}\\\mathbf{Q}_{3}\supset\mathbf{Q}_{2}}} (-1)^{\dim(\mathbf{A}_{2}/\mathbf{A}_{3})} \hat{\tau}_{2}^{3}(H) \Gamma_{\mathbf{Q}_{3}}'(H,\mathcal{Y}_{\mathbf{Q}_{3}}).$$

See  $\S4$  of [Ar86].

(3.3.4) Lemma Let Q be a parabolic subgroup in  $\mathcal{F}^{\Sigma}$ , let  $\mathcal{Y}$  be a collection of points as in (3.3.2.2). The function  $\Gamma'_{Q}(, \mathcal{Y}_{Q})$  is compactly supported as a function on  $\mathfrak{a}_{Q}^{G}$ .

Let  $(c_P)$  be the  $(G, M_1)$ -family associated with a  $(G, M_1)$ -orthogonal set that contains  $\mathcal{Y}_Q$ as in (3.1.4.1). Let  $c'_P$  be the functions intervening in (3.1.2.5). Let  $\Gamma'_Q(, \mathcal{Y}_Q)$  denote the Fourier transform of  $\Gamma'_Q(, \mathcal{Y}_Q)$ . Then

(3.3.4.1) 
$$\forall \lambda \in i\mathfrak{a}_{\mathbf{Q}}^{\mathbf{G},*} \quad \Gamma'_{\mathbf{Q}}(\lambda,\mathcal{Y}_{\mathbf{Q}}) = \sum_{\mathbf{P} \in \mathring{\mathcal{F}}_{\mathbf{Q}}(\mathbf{M}_{1})} c'_{\mathbf{P}}(\lambda).$$

*Proof.* See Lemma 4.1 of [Ar86].

(3.3.5) **Remark** The main ingredient of the proof is the following generalization of (2.2.2.2):

Let Q be a parabolic subgroup in  $\mathcal{F}^{\Sigma}$ , let P be a parabolic subgroup in  $\overline{\mathcal{F}}_Q(M_1)$ , then

$$(3.3.5.2)\sum_{\substack{\mathbf{P}'\in\bar{\mathcal{F}}_{\mathbf{Q}}(\mathbf{M}_{1})\\\mathbf{P}'\subset\mathbf{P}}} (-1)^{\dim(\mathbf{A}_{\mathbf{P}'}/\mathbf{A}_{\mathbf{P}})} \tau_{\mathbf{Q}}^{\pi_{\Sigma}(\mathbf{P}')}(H)\hat{\tau}_{\mathbf{P}'}^{\mathbf{P}}(H) = \begin{cases} 1 & \text{if } \mathbf{P}\in\mathcal{F}_{\mathbf{Q}}(\mathbf{M}_{1}) \text{ and } H\in\mathfrak{a}_{\mathbf{P}}, \\ 0 & \text{otherwise.} \end{cases}$$

This is Lemma 4.2 of [Ar86].

(3.3.6) Lemma (Global semisimple descent) Let  $P_1$ ,  $M_1$ ,  $\mathfrak{o}$ ,  $\Sigma$  be as in (3.3.1), let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$ , then

$$(3.3.6.1) \quad J_{\mathfrak{o}}(f) = |\pi_{0}(\mathcal{G}_{\Sigma})|^{-1} \int_{\mathcal{G}_{\Sigma}^{0}(\mathbb{A})\backslash \mathcal{G}(\mathbb{A})} \left( \sum_{\mathcal{Q}\in\mathcal{F}^{\Sigma}} |\mathcal{W}_{1}^{\mathcal{M}_{Q}}| |\mathcal{W}_{1}^{\mathcal{G}_{\Sigma}^{0}}|^{-1} J_{\mathrm{nil}}^{\mathcal{M}_{Q}}(\Phi_{\mathcal{Q},x}^{T_{0}-T_{\Sigma,1}}) \right) \, \mathrm{d}x$$

where for a truncation parameter T in  $\mathfrak{a}_0$  the function  $\Phi_{Q,x}^T$  on  $\mathfrak{m}_Q(\mathbb{A})$  is defined by (3.3.6.2)  $\forall X \in \mathfrak{m}_Q(\mathbb{A})$ 

$$\Phi_{\mathbf{Q},x}^{T}(X) = \int_{\mathbf{K}_{\Sigma}} \int_{\mathfrak{n}_{\mathbf{Q}}(\mathbb{A})} f\Big( \big(\Sigma + (X+N) \cdot \mathrm{ad}(k)\big) \cdot \mathrm{ad}(x) \Big) \ v_{\mathbf{Q}}'(kx,T) \mathrm{d}N \mathrm{d}k$$

where the weight factor  $v_{\rm Q}^\prime$  is defined as

(3.3.6.3) 
$$v'_{\mathbf{Q}}(kx,T) = \int_{\mathfrak{a}_{\mathbf{Q}}^{\mathbf{G}}} \Gamma'_{\mathbf{Q}}(H,\mathcal{Y}_{\mathbf{Q}}^{T}(k,x)) \, \mathrm{d}H$$

where  $\mathcal{Y}^{T}(k, x)$  is the collection of points defined by

(3.3.6.4) 
$$\forall \mathbf{P} \in \mathcal{F}_{\mathbf{Q}}(\mathbf{M}_1) \qquad Y_{\mathbf{P}}^T(k, x) = -H_{\mathbf{P}}(kx) - T_{\Sigma} + T$$

where  $T_{\Sigma}$  is a truncation parameter in  $\mathfrak{a}_1$  such that  $T_{\Sigma} - T_{\Sigma,1}$  is the projection of  $T - T_0$ .

Proof. This argument follows the proof of Lemma 6.2 of [Ar86].

Let T be a truncation parameter in  $\mathfrak{a}_0$ . Define a second truncated kernel function  $j_{\mathfrak{o}}^T(, f)$ on  $\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^1$  by

(3.3.6.5) 
$$\forall x \in \mathcal{G}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{A})^{1}$$
$$j_{\mathfrak{o}}^{T}(x, f) = \sum_{\substack{\mathcal{P} \in \mathcal{F} \\ \text{standard}}} (-1)^{\dim(\mathcal{A}_{\mathcal{P}}/\mathcal{A}_{\mathcal{G}})} \sum_{\delta \in \mathcal{P}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{Q})} \hat{\tau}_{\mathcal{P}}(H_{0}(\delta x) - T) J_{\mathcal{P},\mathfrak{o}}(\delta x)$$

where

$$(3.3.6.6) \quad J_{\mathrm{P},\mathfrak{o}}(x,f) = \sum_{X \in \mathfrak{m}_{\mathrm{P}}(\mathbb{Q}) \cap \mathfrak{o}} \sum_{\eta \in \mathrm{N}_{\mathrm{P},X_{\mathrm{ss}}}(\mathbb{Q}) \setminus \mathrm{N}_{\mathrm{P}}(\mathbb{Q})} \int_{\mathfrak{n}_{\mathrm{P},X_{\mathrm{ss}}}(\mathbb{A})} f((X+N) \cdot \mathrm{ad}(\eta x)) \, \mathrm{d}N.$$

**Lemma** The function  $j_{\mathfrak{o}}^{T}(\cdot, f)$  is integrable on  $G(\mathbb{Q}) \setminus G(\mathbb{A})^{1}$ , and

(3.3.6.7) 
$$J^T_{\mathfrak{o}}(f) = \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^1} j^T_{\mathfrak{o}}(x,f) \, \mathrm{d}x.$$

*Proof.* Following the proof of (2.2.2.1), by the combinatorial lemma of Langlands (2.2.2.2), it suffices to prove that

(3.3.6.8) 
$$\sum_{\substack{P_2,P_5\in\mathcal{F}\\\text{standard}\\P_2\subset P_5}} \int_{P_2(\mathbb{Q})\backslash G(\mathbb{A})^1} F^{P_2}(x,T) \sigma_2^5(H_0(x)-T) \times \left| \sum_{\substack{P_4\in\mathcal{F}\\P_2\subset P_4\subset P_5}} (-1)^{\dim(A_4/A_G)} J_{P_4,\mathfrak{o}}(x,f) \right| dx$$

is finite. Decompose the sum defining  $J_{\mathrm{P}_4,\mathfrak{o}}(x,f)$  over the set of parabolic subgroups contained in  $\mathrm{P}_4$ ,

$$(3.3.6.9)$$
  $J_{P_4,o}(x, f)$ 

$$\begin{aligned} &= \sum_{\substack{P_{3} \in \mathcal{F} \\ P_{2} \subset P_{3} \subset P_{4}}} \sum_{X \in \mathfrak{m}_{2}^{2}(\mathbb{Q})' \cap \mathfrak{o}} \sum_{Y \in \mathfrak{n}_{3}^{4}(\mathbb{Q})} \\ &\times \sum_{\eta \in \mathcal{N}_{4,(X+Y)_{\mathrm{ss}}}(\mathbb{Q}) \setminus \mathcal{N}_{4}(\mathbb{Q})} \int_{\mathfrak{n}_{4,(X+Y)_{\mathrm{ss}}}(\mathbb{A})} f((X+Y+N) \cdot \mathrm{ad}(\eta x)) \, \mathrm{d}N \\ (3.3.6.10) &= \sum_{\substack{P_{3} \in \mathcal{F} \\ P_{2} \subset P_{3} \subset P_{4}}} \sum_{X \in \mathfrak{m}_{2}^{2}(\mathbb{Q})' \cap \mathfrak{o}} \sum_{Z \in \mathfrak{n}_{3,X_{\mathrm{ss}}}^{4}(\mathbb{Q})} \sum_{\delta \in \mathcal{N}_{4,X_{\mathrm{ss}}}^{4}(\mathbb{Q}) \setminus \mathcal{N}_{4}^{4}(\mathbb{Q})} \\ &\times \sum_{\eta \in \mathcal{N}_{4,(X+Z)_{\mathrm{ss}} \cdot \mathrm{ad}(\delta)}(\mathbb{Q}) \setminus \mathcal{N}_{4}(\mathbb{Q})} \\ &\times \int_{\mathfrak{n}_{4,(X+Z)_{\mathrm{ss}} \cdot \mathrm{ad}(\delta)}(\mathbb{Q}) \setminus \mathcal{N}_{4}(\mathbb{Q})} \\ &= \sum_{\substack{P_{3} \in \mathcal{F} \\ P_{2} \subset P_{3} \subset P_{4}}} \sum_{X \in \mathfrak{m}_{2}^{2}(\mathbb{Q})' \cap \mathfrak{o}} \sum_{Z \in \mathfrak{n}_{3,X_{\mathrm{ss}}}^{4}(\mathbb{Q})} \sum_{\eta \in \mathcal{N}_{3,X_{\mathrm{ss}}}(\mathbb{Q}) \setminus \mathcal{N}_{3}(\mathbb{Q})} \\ &\times \int_{\mathfrak{n}_{4,(X+Z)_{\mathrm{ss}}}(\mathbb{A})} f((X+Z+N) \cdot \mathrm{ad}(\eta x)) \, \mathrm{d}N \end{aligned}$$

$$(3.3.6.11) = \sum_{\substack{P_{3} \in \mathcal{F} \\ P_{2} \subset P_{3} \subset P_{4}}} \sum_{X \in \mathfrak{m}_{2}^{2}(\mathbb{Q})' \cap \mathfrak{o}} \sum_{\eta \in \mathcal{N}_{3,X_{\mathrm{ss}}}(\mathbb{Q}) \setminus \mathcal{N}_{3}(\mathbb{Q})} \\ &\times \sum_{Z \in \mathfrak{n}_{3,X_{\mathrm{ss}}}(\mathbb{Q})} \int_{\mathfrak{n}_{3,X_{\mathrm{ss}}}(\mathbb{A})} f((X+N) \cdot \mathrm{ad}(\eta x)) \cdot \psi(\langle N, \overline{Z} \rangle) \, \mathrm{d}N \end{aligned}$$

where (3.3.6.10) follows from Corollary 2.4 of [Ch02a] which states that

$$(3.3.6.12) \qquad \sum_{Z \in \mathfrak{n}_{X_{ss}}(\mathbb{Q})} \sum_{\delta \in \mathcal{N}_{X_{ss}}(\mathbb{Q}) \setminus \mathcal{N}(\mathbb{Q})} f((X+Z) \cdot \mathrm{ad}(\delta)) = \sum_{Y \in \mathfrak{n}(\mathbb{Q})} f(X+Y)$$

and (3.3.6.11) follows from the Poisson summation formula. The right hand side of (3.3.6.11) is independent of  $P_4$ , hence the alternating sum in  $P_4$  cancels by the inclusion-exclusion principle, therefore the same estimates in the proof of (2.2.2.1) implies the integrability of  $j_{\mathfrak{o}}^T(, f)$ .

For the integral representation of  $J^T_{\mathfrak{o}}(f)$ , consider the (P<sub>2</sub>, P<sub>5</sub>) summand of the integral of  $j^T_{\mathfrak{o}}(f)$ , f) over  $G(\mathbb{Q})\backslash G(\mathbb{A})^1$ :

(3.3.6.13) 
$$\int_{P_{2}(\mathbb{Q})\backslash G(\mathbb{A})^{1}} F^{P_{2}}(x,T) \sigma_{2}^{5}(H_{0}(x)-T) \times \\ \times \sum_{\substack{P_{4}\in\mathcal{F}\\P_{2}\subset P_{4}\subset P_{5}}} (-1)^{\dim(A_{4}/A_{G})} J_{P_{4},\mathfrak{o}}(x,f) \, \mathrm{d}x$$

$$\begin{aligned} &= \int_{M_{2}(\mathbb{Q})N_{2}(\mathbb{A})\backslash G(\mathbb{A})^{1}} F^{P_{2}}(x,T)\sigma_{2}^{5}(H_{0}(x)-T) \times \\ &\times \sum_{\substack{P_{4}\in\mathcal{F}\\P_{2}\subset P_{4}\subset P_{5}}} (-1)^{\dim(A_{4}/A_{G})} \sum_{X\in\mathfrak{m}_{4}(\mathbb{Q})\cap\mathfrak{o}} \left(\int_{N_{2}(\mathbb{Q})\backslash N_{2}(\mathbb{A})} \sum_{\eta\in N_{4,X_{88}}(\mathbb{Q})\backslash N_{4}(\mathbb{Q})} \\ &\times \left(\int_{\mathfrak{n}_{4,X_{88}}(\mathbb{A})} f((X+N)\cdot\mathrm{ad}(\eta n_{2}x)) \,\mathrm{d}N\right) \mathrm{d}n_{2}\right) \mathrm{d}x \end{aligned}$$

$$(3.3.6.14) = \int_{M_{2}(\mathbb{Q})N_{2}(\mathbb{A})\backslash G(\mathbb{A})^{1}} F^{P_{2}}(x,T)\sigma_{2}^{5}(H_{0}(x)-T) \times \\ &\times \sum_{\substack{P_{4}\in\mathcal{F}\\P_{2}\subset P_{4}\subset P_{5}}} (-1)^{\dim(A_{4}/A_{G})} \sum_{X\in\mathfrak{m}_{4}(\mathbb{Q})\cap\mathfrak{o}} \left(\int_{N_{2}(\mathbb{Q})\backslash N_{2}(\mathbb{A})} \int_{N_{4,X_{88}}(\mathbb{Q})\backslash N_{4}(\mathbb{A})} \\ &\times \left(\int_{\mathfrak{n}_{4,X_{88}}(\mathbb{A})} f((X+N)\cdot\mathrm{ad}(nn_{2}x)) \,\mathrm{d}N\right) \mathrm{d}ndn_{2}\right) \mathrm{d}x \end{aligned}$$

$$(3.3.6.15) = \int_{M_{2}(\mathbb{Q})N_{2}(\mathbb{A})\backslash G(\mathbb{A})^{1}} F^{P_{2}}(x,T)\sigma_{2}^{5}(H_{0}(x)-T) \sum_{\substack{P_{4}\in\mathcal{F}\\P_{2}\subset P_{4}\subset P_{5}}} (-1)^{\dim(A_{4}/A_{G})} \times \\ &\times \left(\sum_{X\in\mathfrak{m}_{4}(\mathbb{Q})\cap\mathfrak{o}} \int_{N_{2}(\mathbb{Q})\backslash N_{2}(\mathbb{A})} \int_{\mathfrak{n}_{4}(\mathbb{A})} f((X+N)\cdot\mathrm{ad}(n_{2}x)) \,\mathrm{d}N\mathrm{d}n_{2}\right) \mathrm{d}x. \end{aligned}$$

The equality (3.3.6.14) holds since  $N_4(\mathbb{Q})\setminus N_4(\mathbb{A})$  has volume 1, and the equality (3.3.6.15) follows from Corollary 2.5 of [Ch02a] which is the integral analogue of (3.3.6.12). Reversing the combinatorial manipulations to the right hand side of (3.3.6.15) skipping the step (3.3.6.14),

$$(3.3.6.16) \qquad \int_{\mathcal{M}_{2}(\mathbb{Q})\mathcal{N}_{2}(\mathbb{A})\backslash \mathcal{G}(\mathbb{A})^{1}} F^{\mathcal{P}_{2}}(x,T) \sigma_{2}^{5}(H_{0}(x)-T) \sum_{\substack{\mathcal{P}_{4}\in\mathcal{F}\\\mathcal{P}_{2}\subset\mathcal{P}_{4}\subset\mathcal{P}_{5}}} (-1)^{\dim(\mathcal{A}_{4}/\mathcal{A}_{G})} \times \\ \times \left(\sum_{X\in\mathfrak{m}_{4}(\mathbb{Q})\cap\mathfrak{o}} \int_{\mathcal{N}_{2}(\mathbb{Q})\backslash\mathcal{N}_{2}(\mathbb{A})} \int_{\mathfrak{n}_{4}(\mathbb{A})} f((X+N)\cdot\operatorname{ad}(n_{2}x)) \, \mathrm{d}N\mathrm{d}n_{2}\right) \mathrm{d}x \\ = \int_{\mathcal{P}_{2}(\mathbb{Q})\backslash\mathcal{G}(\mathbb{A})^{1}} F^{\mathcal{P}_{2}}(x,T) \sigma_{2}^{5}(H_{0}(x)-T) \times \\ \times \sum_{\substack{\mathcal{P}_{4}\in\mathcal{F}\\\mathcal{P}_{2}\subset\mathcal{P}_{4}\subset\mathcal{P}_{5}}} (-1)^{\dim(\mathcal{A}_{4}/\mathcal{A}_{G})} K_{\mathcal{P}_{4},\mathfrak{o}}(x,f) \, \mathrm{d}x,$$

which is the  $(\mathbf{P}_2, \mathbf{P}_5)$  summand of the integral of  $k_{\mathbf{o}}^T(\cdot, f)$  over  $\mathbf{G}(\mathbb{Q}) \setminus \mathbf{G}(\mathbb{A})^1$ , hence  $J_{\mathbf{o}}^T(f)$ .  $\Box$ 

By (3.3.6.7) it is enough to consider the function  $J_{\mathrm{P},\mathfrak{o}}(\ ,f)$  instead of  $K_{\mathrm{P},\mathfrak{o}}(\ ,f)$  for a parabolic subgroup P of G containing P<sub>1</sub> for the proof of (3.3.6.1). Every element X in  $\mathfrak{m}_{\mathrm{P}}(\mathbb{Q}) \cap \mathfrak{o}$  is conjugate under the adjoint action of  $\mathrm{G}(\mathbb{Q})$  to the sum of  $\Sigma$  and  $N_{\Sigma}$  for some  $N_{\Sigma}$  in  $\mathfrak{g}_{\Sigma,\mathrm{nil}}(\mathbb{Q})$ . More precisely

$$(3.3.6.17) \qquad \exists \mathbf{P}_1' \in \mathcal{F}, \ \mathbf{P}_1' \text{ standard}, \ \mathbf{P}_1' \subset \mathbf{P} \quad \exists s \in \mathbf{W}(\mathfrak{a}_{\mathbf{P}_1}, \mathfrak{a}_{\mathbf{P}_1'}) \quad \exists \mu \in \mathbf{M}_{\mathbf{P}}^0(\mathbb{Q}) \\ \exists N_{\Sigma} \in \mathfrak{m}_{\mathbf{P}}(\mathbb{Q}) \cdot \mathrm{ad}(w_s) \cap \mathfrak{g}_{\Sigma, \mathrm{nil}}(\mathbb{Q}) \quad X = (\Sigma + N_{\Sigma}) \cdot \mathrm{ad}(w_s^{-1}\mu)$$

where

- the element  $w_s$  in  $G(\mathbb{Q})$  is a representative of s;
- the double coset

$$(3.3.6.18) \qquad [s] \in W_0^{M_P} \backslash W(\mathfrak{a}_{P_1}, \mathfrak{a}_{P'_1}) / W_1^{G_{\Sigma}^0}$$

is uniquely determined;

• for a fixed choice of the element s, the coset

$$(3.3.6.19) \qquad \qquad [\mu] \in \mathcal{M}_{\mathcal{P}}(\mathbb{Q}) \cap w_s \mathcal{G}_{\Sigma}(\mathbb{Q}) w_s^{-1} \backslash \mathcal{M}_{\mathcal{P}}(\mathbb{Q})$$

is uniquely determined;

• for a fixed choice of the representative  $w_s$  and the element  $\mu$ , the element  $N_{\Sigma}$  is uniquely determined.

Let  $W(\mathfrak{a}_1; M_P^+, G_{\Sigma}^{0,+})$  be the subset of the Weyl group of G defined by

$$(3.3.6.20) \qquad W(\mathfrak{a}_{1}; \mathrm{M}_{\mathrm{P}}^{+}, \mathrm{G}_{\Sigma}^{0,+}) \\ = \bigcup_{\substack{\mathrm{P}_{1}' \in \mathcal{F} \\ \text{standard} \\ \mathrm{P}_{1}' \subset \mathrm{P}}} \left\{ s \in \mathrm{W}(\mathfrak{a}_{\mathrm{P}_{1}}, \mathfrak{a}_{\mathrm{P}_{1}'}) : \ \forall \alpha \in \Delta_{\mathrm{P}_{1}'}^{\mathrm{P}} \ s^{-1}\alpha > 0, \ \forall \beta \in \Delta_{1}^{\mathrm{G}_{\Sigma}^{0}} \ s\beta > 0 \right\}.$$

The map

$$(3.3.6.21) \qquad \left( \mathbf{W}(\mathfrak{a}_{1}; \mathbf{M}_{\mathbf{P}}^{+}, \mathbf{G}_{\Sigma}^{0,+}) \right) \times \left( \mathbf{M}_{\mathbf{P}}(\mathbb{Q}) \cap w_{s} \mathbf{G}_{\Sigma}^{0}(\mathbb{Q}) w_{s}^{-1} \setminus \mathbf{M}_{\mathbf{P}}(\mathbb{Q}) \right) \times \\ \times \left( \mathfrak{m}_{\mathbf{P}}(\mathbb{Q}) \cdot \mathrm{ad}(w_{s}) \cap \mathfrak{g}_{\Sigma,\mathrm{nil}}(\mathbb{Q}) \right) \\ \longrightarrow \qquad \mathfrak{m}_{\mathbf{P}}(\mathbb{Q}) \cap \mathfrak{o}$$

defined by

$$(3.3.6.22) \qquad (s,\mu,N_{\Sigma}) \mapsto (\Sigma+N_{\Sigma}) \cdot \operatorname{ad}(w_s^{-1}\mu)$$

is surjective and the group  $\pi_0(G_{\Sigma})$  operates simply transitively on each fiber. Hence

$$(3.3.6.23) \qquad \forall x \in \mathcal{G}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{A})^1$$

$$J_{\mathrm{P},\mathfrak{o}}(x,f)$$

$$= \sum_{s \in W(\mathfrak{a}_{1}; M_{P}^{+}, G_{\Sigma}^{0, +})} \sum_{\mu \in M_{P}(\mathbb{Q}) \cap w_{s} G_{\Sigma}^{0}(\mathbb{Q}) w_{s}^{-1} \setminus M_{P}(\mathbb{Q})} \times \sum_{N_{\Sigma} \in \mathfrak{m}_{P}(\mathbb{Q}) \cdot \mathrm{ad}(w_{s}) \cap \mathfrak{g}_{\Sigma, \mathrm{nil}}(\mathbb{Q})} \times |\pi_{0}(G_{\Sigma})|^{-1} \sum_{\eta \in N_{P, \Sigma \cdot \mathrm{ad}(w_{s}^{-1}\mu)}(\mathbb{Q}) \setminus N_{P}(\mathbb{Q})} \int_{\mathfrak{n}_{P, \Sigma \cdot \mathrm{ad}(w_{s}^{-1}\mu)}(\mathbb{A})} \times f\left(\left((\Sigma + N_{\Sigma}) \cdot \mathrm{ad}(w_{s}^{-1}\mu) + N\right) \cdot \mathrm{ad}(\eta x)\right)\right) dN$$

$$(3.3.6.24) = |\pi_{0}(G_{\Sigma})|^{-1} \sum_{s \in W(\mathfrak{a}_{1}; M_{P}^{+}, G_{\Sigma}^{0, +})} \sum_{N_{\Sigma} \in \mathfrak{m}_{P}(\mathbb{Q}) \cdot \mathrm{ad}(w_{s}) \cap \mathfrak{g}_{\Sigma, \mathrm{nil}}(\mathbb{Q})} \times \sum_{\pi \in P(\mathbb{Q}) \cap w_{s} G_{\Sigma}^{0}(\mathbb{Q}) w_{s}^{-1} \setminus P(\mathbb{Q})} \int_{\mathfrak{n}_{P, \Sigma}(\mathbb{A}) \cdot \mathrm{ad}(w_{s})} \sum_{\times f\left((\Sigma + N_{\Sigma} + N) \cdot \mathrm{ad}(w_{s}^{-1}\pi x)\right)} dN$$

where (3.3.6.24) follows from the change of variables

$$(3.3.6.25) \qquad \qquad \mathbf{M}_{\mathbf{P}}(\mathbb{Q}) \cap w_{s} \mathbf{G}_{\Sigma}^{0}(\mathbb{Q}) w_{s}^{-1} \Big\backslash \mathbf{M}_{\mathbf{P}}(\mathbb{Q}) \times \mathbf{N}_{\mathbf{P}, \Sigma \cdot \mathrm{ad}(w_{s}^{-1}\mu)}(\mathbb{Q}) \setminus \mathbf{N}_{\mathbf{P}}(\mathbb{Q}) \\ \xrightarrow{\sim} \qquad \mathbf{P}(\mathbb{Q}) \cap w_{s} \mathbf{G}_{\Sigma}^{0}(\mathbb{Q}) w_{s}^{-1} \Big\backslash \mathbf{P}(\mathbb{Q}).$$

Substitute (3.3.6.23) into the formula (3.3.6.7),

$$(3.3.6.26) \qquad J_{\mathfrak{o}}^{T}(f)$$

$$= |\pi_{0}(\mathcal{G}_{\Sigma})|^{-1} \int_{\mathcal{G}(\mathbb{Q})\backslash \mathcal{G}(\mathbb{A})^{1}} \left( \sum_{\substack{\mathcal{P}\in\mathcal{F}\\\text{standard}}} (-1)^{\dim(\mathcal{A}_{\mathcal{P}}/\mathcal{A}_{\mathcal{G}})} \sum_{\delta\in\mathcal{P}(\mathbb{Q})\backslash\mathcal{G}(\mathbb{Q})} \right)$$

$$\times \sum_{s\in\mathcal{W}(\mathfrak{a}_{1};\mathcal{M}_{\mathcal{P}}^{+},\mathcal{G}_{\Sigma}^{0,+})} \sum_{N_{\Sigma}\in\mathfrak{m}_{\mathcal{P}}(\mathbb{Q})\cdot\mathrm{ad}(w_{s})\cap\mathfrak{g}_{\Sigma,\mathrm{nil}}(\mathbb{Q})} \sum_{\pi\in\mathcal{P}(\mathbb{Q})\cap w_{s}\mathcal{G}_{\Sigma}^{0}(\mathbb{Q})w_{s}^{-1}\backslash\mathcal{P}(\mathbb{Q})}$$

$$\times \int_{\mathfrak{n}_{\mathcal{P},\Sigma}(\mathbb{A})\cdot\mathrm{ad}(w_{s})} f\left((\Sigma+N_{\Sigma}+N)\cdot\mathrm{ad}(w_{s}^{-1}\pi\delta x)\right) dN \times$$

$$\times \hat{\tau}_{\mathcal{P}}(H_{\mathcal{P}}(\delta x)-T) dx$$
$$\begin{aligned} (3.3.6.27) &= |\pi_{0}(\mathbf{G}_{\Sigma})|^{-1} \int_{\mathbf{G}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{A})^{1}} \left( \sum_{\substack{\mathbf{P}\in\mathcal{F}\\\text{standard}}} (-1)^{\dim(\mathbf{A}_{\mathbf{P}}/\mathbf{A}_{\mathbf{G}})} \sum_{s\in\mathbf{W}(\mathfrak{a}_{1};\mathbf{M}_{\mathbf{P}}^{+},\mathbf{G}_{\Sigma}^{0,+})} \right. \\ &\times \sum_{\xi\in\mathbf{Q}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{Q})} \sum_{N_{\Sigma}\in\mathfrak{m}_{\mathbf{Q},\mathrm{nil}}(\mathbb{Q})} \int_{\mathfrak{n}_{\mathbf{Q}}(\mathbb{A})} f((\Sigma+N_{\Sigma}+N)\cdot\mathrm{ad}(\xi x)) \, \mathrm{d}N \times \\ &\times \hat{\tau}_{\mathbf{P}}(H_{\mathbf{p}}(w_{s}\xi x)-T) \right) \, \mathrm{d}x \end{aligned} \\ &= |\pi_{0}(\mathbf{G}_{\Sigma})|^{-1} \int_{\mathbf{G}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{A})^{1}} \left( \sum_{\substack{\mathbf{Q}\in\mathcal{F}^{\Sigma}\\\mathrm{standard}}} \sum_{\xi\in\mathbf{Q}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{Q})} \\ &\times \sum_{N_{\Sigma}\in\mathfrak{m}_{\mathbf{Q},\mathrm{nil}}(\mathbb{Q})} \int_{\mathfrak{n}_{\mathbf{Q}}(\mathbb{A})} f((\Sigma+N_{\Sigma}+N)\cdot\mathrm{ad}(\xi x)) \, \mathrm{d}N \times \\ &\times \sum_{\substack{N_{\Sigma}\in\mathfrak{m}_{\mathbf{Q},\mathrm{nil}}(\mathbb{Q})\\\mathrm{standard}}} (-1)^{\dim(\mathbf{A}_{\mathbf{P}}/\mathbf{A}_{\mathbf{G}})} \hat{\tau}_{\mathbf{P}}(H_{\mathbf{p}}(w_{s}\xi x)-T) \right) \, \mathrm{d}x. \end{aligned}$$

where in (3.3.6.27)  $\xi$  denotes the product  $w_s^{-1}\pi\delta$  and Q denotes the standard parabolic subgroup of  ${\rm G}^0_\Sigma$  defined by

$$(3.3.6.28) \qquad \qquad \mathbf{Q} = w_s^{-1} \mathbf{P} w_s \cap \mathbf{G}_{\Sigma}^0$$

with Levi decomposition

$$(3.3.6.29) \qquad \qquad \mathfrak{m}_{\mathbf{Q}} = \mathfrak{m}_{\mathbf{P}} \cdot \mathrm{ad}(w_s) \cap \mathfrak{g}_{\Sigma}, \qquad \mathfrak{n}_{\mathbf{Q}} = \mathfrak{n}_{\mathbf{P}} \cdot \mathrm{ad}(w_s) \cap \mathfrak{g}_{\Sigma}.$$

The assignment

$$(3.3.6.30) \qquad (\mathbf{P},s) \quad \mapsto \quad \mathbf{P}' = w_s^{-1} \mathbf{P} w_s$$

defines a bijection

$$(3.3.6.31) \qquad \left\{ \mathbf{P} \in \mathcal{F}, \ s \in \mathbf{W}(\mathfrak{a}_1; \mathbf{M}_{\mathbf{P}}^+, \mathbf{G}_{\Sigma}^{0,+}) : \ \mathbf{P} \text{ standard}, \ w_s^{-1} \mathbf{P} w_s \cap \mathbf{G}_{\Sigma}^0 = \mathbf{Q} \right\}$$
$$\xrightarrow{\sim} \mathcal{F}_{\mathbf{Q}}(\mathbf{M}_1),$$

hence

 $(3.3.6.32) J_{\mathfrak{o}}^T(f)$ 

$$= |\pi_0(G_{\Sigma})|^{-1} \int_{G(\mathbb{Q})\backslash G(\mathbb{A})^1} \left(\sum_{\substack{Q\in\mathcal{F}^{\Sigma}\\ standard}} \sum_{\xi\in Q(\mathbb{Q})\backslash G(\mathbb{Q})}\right)$$

$$\times \sum_{N_{\Sigma} \in \mathfrak{m}_{Q,nil}(\mathbb{Q})} \int_{\mathfrak{n}_{Q}(\mathbb{A})} f((\Sigma + N_{\Sigma} + N) \cdot \operatorname{ad}(\xi x)) \, \mathrm{d}N \times$$
$$\times \sum_{\mathrm{P}' \in \mathcal{F}_{Q}(\mathrm{M}_{1})} (-1)^{\dim(\mathrm{A}_{\mathrm{P}'}/\mathrm{A}_{\mathrm{G}})} \hat{\tau}_{\mathrm{P}'}(H_{\mathrm{P}'}(\xi x) - s^{-1}(T - T_{0}) - T_{0}) \right) \, \mathrm{d}x$$

where s denotes the second component of the inverse image of P' under the map defined in (3.3.6.30).

Let  $T_{\Sigma}$  be a truncation parameter for the triple  $(G_{\Sigma}(\mathbb{A})^0, M_{1,\Sigma}(\mathbb{A}), K_{\Sigma})$  in  $\mathfrak{a}_1$  such that  $T_{\Sigma} - T_{\Sigma,1}$  is the projection of  $T - T_0$ . By (3.3.3.1)

$$(3.3.6.33) \qquad \sum_{\mathbf{P}\in\mathcal{F}_{\mathbf{Q}}(\mathbf{M}_{1})} (-1)^{\dim(\mathbf{A}_{\mathbf{P}}/\mathbf{A}_{\mathbf{G}})} \hat{\tau}_{\mathbf{P}} \left( H_{\mathbf{P}}(\delta xy) - s^{-1}(T - T_{0}) - T_{0} \right)$$
$$= \sum_{\mathbf{P}\in\mathcal{F}_{\mathbf{Q}}(\mathbf{M}_{1})} (-1)^{\dim(\mathbf{A}_{\mathbf{P}}/\mathbf{A}_{\mathbf{G}})} \hat{\tau}_{\mathbf{P}} \left( (H_{\mathbf{Q}}(\delta x) - T_{\Sigma}) - Y_{\mathbf{P}}^{T}(\delta x, y) \right)$$
$$= \sum_{\substack{\mathbf{Q}'\in\mathcal{F}^{\Sigma}\\\mathbf{Q}'\supset\mathbf{Q}}} (-1)^{\dim(\mathbf{A}_{\mathbf{Q}}/\mathbf{A}_{\mathbf{Q}'})} \hat{\tau}_{\mathbf{Q}}^{\mathbf{Q}'} \left( H_{\mathbf{Q}}(\delta x) - T_{\Sigma} \right) \times$$
$$\times \Gamma'_{\mathbf{Q}'} \left( H_{\mathbf{Q}'}(\delta x) - T_{\Sigma}, \mathcal{Y}_{\mathbf{Q}'}^{T}(\delta x, y) \right)$$

where the family  $\mathcal{Y}_{\mathbf{Q}'}^T(\delta x, y)$  is defined by

(3.3.6.34) 
$$\forall \mathbf{P} \in \mathcal{F}_{\mathbf{Q}}(\mathbf{M}_{1}) \quad Y_{\mathbf{P}}^{T}(\delta x, y) = -H_{\mathbf{P}}(ky) + s^{-1}(T - T_{0}) - T_{\Sigma} + T_{0},$$

where k is the  $K_{\Sigma}$  component of  $\delta x$  under the Iwasawa decomposition with respect to  $P_{\Sigma}$ and s is the second component of the inverse image of P under the map defined in (3.3.6.30). On the right hand side of (3.3.6.32) make the change of variables

$$(3.3.6.35) \qquad Q(\mathbb{Q}) \setminus G(\mathbb{Q}) \times G(\mathbb{Q}) \setminus G(\mathbb{A})^1 \\ \xrightarrow{\sim} \qquad Q(\mathbb{Q}) \setminus G_{\Sigma}^0(\mathbb{Q}) \times G_{\Sigma}^0(\mathbb{Q}) \setminus G_{\Sigma}^0(\mathbb{A}) \cap G(\mathbb{A})^1 \times G_{\Sigma}^0(\mathbb{A}) \setminus G(\mathbb{A}).$$

Then

$$(3.3.6.36) J_{\mathfrak{o}}^T(f)$$

$$= |\pi_{0}(\mathbf{G}_{\Sigma})|^{-1} \int_{\mathbf{G}_{\Sigma}^{0}(\mathbb{A})\backslash\mathbf{G}(\mathbb{A})} \int_{\mathbf{G}_{\Sigma}^{0}(\mathbb{Q})\backslash\mathbf{G}_{\Sigma}^{0}(\mathbb{A})\cap\mathbf{G}(\mathbb{A})^{1}} \left(\sum_{\substack{\mathbf{Q}\in\mathcal{F}^{\Sigma}\\ \text{standard}}} \sum_{\substack{\mathbf{Q}'\in\mathcal{F}^{\Sigma}\\ \mathbf{Q}'\supset\mathbf{Q}}} \times \sum_{\delta\in\mathbf{Q}(\mathbb{Q})\backslash\mathbf{G}_{\Sigma}^{0}(\mathbb{Q})} \sum_{N_{\Sigma}\in\mathfrak{m}_{\mathbf{Q},\text{nil}}(\mathbb{Q})} (-1)^{\dim(\mathbf{A}_{\mathbf{Q}}/\mathbf{A}_{\mathbf{Q}'})} \times \int_{\mathfrak{n}_{\mathbf{Q}}(\mathbb{A})} f((\Sigma+N_{\Sigma}+N)\cdot\operatorname{ad}(\delta xy)) \, \mathrm{d}N \times \right)$$

where on the right hand side of (3.3.6.37)  $\Phi_{\mathbf{Q}',a,k,y}^T$  denotes the function on  $\mathfrak{m}_{\mathbf{Q}'}(\mathbb{A})$  defined by

$$(3.3.6.38) \qquad \forall X \in \mathfrak{m}_{Q'}(\mathbb{A})$$

$$\Phi_{Q',a,k,y}^{T}(X) = \int_{\mathfrak{m}_{Q'}(\mathbb{A})} f((\Sigma + X + N) \cdot \operatorname{ad}(ky)) \, \mathrm{d}N \times \\ \times \Gamma_{Q'}'(H_{Q'}(a) - T_{\Sigma}, \mathcal{Y}_{Q'}^{T}(k, y)),$$

and the equality (3.3.6.37) follows from the changes of variables

$$\begin{array}{ll} (3.3.6.39) & Q(\mathbb{Q}) \backslash G^0_{\Sigma}(\mathbb{Q}) \times G^0_{\Sigma}(\mathbb{Q}) \backslash G^0_{\Sigma}(\mathbb{A}) \cap G(\mathbb{A})^1 \\ & \xrightarrow{\sim} & Q(\mathbb{Q}) \cap M_{Q'}(\mathbb{Q}) \backslash M_{Q'}(\mathbb{Q}) \times Q'(\mathbb{Q}) \backslash G^0_{\Sigma}(\mathbb{A}) \cap G(\mathbb{A})^1 \end{array}$$

and

$$(3.3.6.40) \qquad \qquad \mathfrak{n}_{\mathbf{Q}}(\mathbb{A}) \xrightarrow{\sim} \mathfrak{n}_{\mathbf{Q}}(\mathbb{A}) \cap \mathfrak{m}_{\mathbf{Q}'}(\mathbb{A}) \times \mathfrak{n}_{\mathbf{Q}'}(\mathbb{A})$$

and the Iwasawa decomposition for  $G^0_{\Sigma}(\mathbb{A}) \cap G(\mathbb{A})^1$ . Evaluate T at  $T_0$  and  $T_{\Sigma}$  at  $T_{\Sigma,1}$ , and apply (3.3.4.1), then

$$(3.3.6.41) J_{\mathfrak{o}}(f) = |\pi_{0}(\mathcal{G}_{\Sigma})|^{-1} \int_{\mathcal{G}_{\Sigma}^{0}(\mathbb{A})\backslash \mathcal{G}(\mathbb{A})} \sum_{\substack{\mathbf{Q}'\in\mathcal{F}^{\Sigma}\\\text{standard}}} J_{\mathrm{nil}}^{\mathcal{M}_{\mathbf{Q}'}}(\Phi_{\mathbf{Q}',y}^{T_{0}-T_{\Sigma,1}}) \, \mathrm{d}y.$$

By the defining property (2.2.8.1) of the points  $T_0$  and  $T_{\Sigma,1}$ ,

(3.3.6.42) 
$$J_{\text{nil}}^{M_{Q'}}(\Phi_{Q',y}^{T_0-T_{\Sigma,1}}) = J_{\text{nil}}^{M_{Q''}}(\Phi_{Q'',y}^{T_0-T_{\Sigma,1}})$$

whenever Q' and Q'' are parabolic subgroups in  $\mathcal{F}^{\Sigma}$  conjugate under the Weyl group of  $G_{\Sigma}^{0}$ , hence

$$(3.3.6.43) \qquad J_{\mathfrak{o}}(f) = |\pi_{0}(\mathbf{G}_{\Sigma})|^{-1} \int_{\mathbf{G}_{\Sigma}^{0}(\mathbb{A})\backslash \mathbf{G}(\mathbb{A})} \left( \sum_{\mathbf{Q}\in\mathcal{F}^{\mathbf{G}_{\Sigma}}} |\mathbf{W}_{1}^{\mathbf{M}_{\mathbf{Q}}}| |\mathbf{W}_{1}^{\mathbf{G}_{\Sigma}^{0}}|^{-1} J_{\mathrm{nil}}^{\mathbf{M}_{\mathbf{Q}}}(\Phi_{\mathbf{Q},x}^{T_{0}-T_{\Sigma,1}}) \right) \, \mathrm{d}x.$$

(3.3.7) **Definition** Fix a finite set of places S. Let M be a standard Levi subgroup of G, let  $\Xi$  be a semisimple element of  $\mathfrak{m}(\mathbb{Q})$ , let  $\nu$  be a nilpotent element of  $\mathfrak{m}_{\Xi}(\mathbb{Q}_S)$  defined upto the adjoint action of  $M_{\Xi}(\mathbb{Q}_S)$ . The element

$$(3.3.7.1) X = \Xi + \nu \in \mathfrak{m}(\mathbb{Q}_S)$$

is well-defined modulo  $\mathcal{M}(\mathbb{Q}_S)$ . Denote by  $D^{\mathcal{G}}$  the discriminant function on  $\mathfrak{g}$ .

Let T be a point in  $\mathfrak{a}_M$ , let x be an element of  $G(\mathbb{Q}_S)$ . Let  $(v_P(x,T))$  be the (G,M)-family defined by

(3.3.7.2) 
$$\forall \mathbf{P} \in \mathcal{F}(\mathbf{M}) \ \forall \lambda \in i\mathfrak{a}_{\mathbf{M}}^* \qquad v_{\mathbf{P}}(x,T)(\lambda) = v_{\mathbf{P}}(x)(\lambda) \times e^{\langle \lambda,T \rangle}$$

Let Q be a parabolic subgroup of  $G^0_{\Xi}$  containing  $M^0_{\Xi}$ . Define the weight factor  $v'_Q$  by

(3.3.7.3) 
$$v'_{Q}(x,T) = \sum_{P \in \mathring{\mathcal{F}}_{Q}(M)} v'_{P}(x,T)$$

where the set  $\mathring{\mathcal{F}}_{Q}(M)$  is defined with respect to the semisimple element  $\Xi$ . Let  $T_{\Xi,M}$  be the point in  $\mathfrak{a}_{M_{\Xi}^{0}}$  modulo  $\mathfrak{a}_{G_{\Xi}^{0}}$  defined in the same manner as  $T_{0}$  in  $\mathfrak{a}_{0}^{G}$  that satisfies an identity analogous to (2.2.8.1).

(3.3.8) Lemma Let L be a Levi subgroup in  $\mathcal{L}(M)$ , let T be a point in  $\mathfrak{a}_M$ , then

$$(3.3.8.1) \qquad \forall x \in \mathcal{G}(\mathbb{Q}_S) \ \forall y \in \mathcal{G}^0_{\Xi}(\mathbb{Q}_S) \qquad v_{\mathcal{L}}(yx) = \sum_{\mathcal{Q} \in \mathcal{F}^{\Xi}(\mathcal{M}^0_{\Xi})} v^{\mathcal{Q}}_{\mathcal{L}^0_{\Xi}}(y) v'_{\mathcal{Q}}(kx,T)$$

where  $v_{\rm L}(x)$  is the weight factor defined in (3.1.5.2) and k is the K<sub>\mathbb{E}</sub>-component of y under the Iwasawa decomposition of  $G^0_{\mathbb{E}}(\mathbb{Q}_S)$  with respect to the parabolic subgroup  $Q^0_{\mathbb{E}}$ .

*Proof.* This is Corollary 8.4 of [Ar88a].

(3.3.9) Lemma (Local semisimple descent)

Let  $f_S$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$ , let X and  $\nu$  be as in (3.3.7.1), then

$$(3.3.9.1) J_{\mathrm{M}}^{\mathrm{G}}(X, f_{S}) = |D^{\mathrm{G}}(\Xi)|_{S}^{1/2} \int_{\mathrm{G}_{\Xi}^{0}(\mathbb{Q}_{S})\backslash\mathrm{G}(\mathbb{Q}_{S})} \left(\sum_{\mathrm{Q}\in\mathcal{F}^{\Xi}(\mathrm{M}_{\Xi}^{0})} J_{\mathrm{M}_{\Xi}^{0}}^{\mathrm{M}_{\mathrm{Q}}}(\nu, \Phi_{S,\mathrm{Q},x}^{T_{0}-T_{\Xi,\mathrm{M}}})\right) \,\mathrm{d}x$$

where  $\Phi_{S,Q,x}^T$  is the function on  $\mathfrak{m}_Q(\mathbb{Q}_S)$  defined by

$$(3.3.9.2) \qquad \forall Y \in \mathfrak{m}_{\mathbb{Q}}(\mathbb{Q}_{S})$$

$$\Phi_{S,\mathbb{Q},x}^{T}(Y)$$

$$= \int_{K_{\Xi}} \int_{\mathfrak{n}_{\mathbb{Q}}(\mathbb{Q}_{S})} f_{S}\Big(\Big(\Xi + (Y+N) \cdot \mathrm{ad}(k)\Big) \cdot \mathrm{ad}(x)\Big) \ v_{\mathbb{Q}}'(kx,T) \mathrm{d}N \mathrm{d}k$$
where the weight factor  $v_{\mathbb{Q}}'(x,T)$  is defined as in (3.3.7.3)

where the weight factor  $v'_{\rm Q}(x,T)$  is defined as in (3.3.7.3).

*Proof.* This argument follows the proof of Corollary 8.7 of [Ar88a], see also the discussion in §2.6 of [HW13].

Since the point  $T_0 - T_{\Xi,M}$  lies in  $\mathfrak{a}_M$ , the identity (3.3.8.1) is valid for the weight factor  $v'_Q(x, T_0 - T_{\Xi,M})$ , hence

$$(3.3.9.3) \qquad J_{M}^{G}(X, f_{S})$$

$$= \lim_{A \to 0} \left( |D^{G}(X+A)|_{S}^{1/2} \int_{G_{Xss}^{0}(\mathbb{Q}_{S}) \setminus G(\mathbb{Q}_{S})} \int_{G_{X+A}^{0}(\mathbb{Q}_{S}) \setminus G_{Xss}^{0}(\mathbb{Q}_{S})} \times f_{S}((X+A) \cdot \operatorname{ad}(yx)) \times \right)$$

$$\times \left( \sum_{L \in \mathcal{L}(M)} r_{M}^{L} (\exp(X_{\operatorname{nil}}), \exp(A)) v_{L}(yx) \right) dy dx \right)$$

$$= \lim_{A \to 0} \left( |D^{G}(X+A)|_{S}^{1/2} \int_{G_{\Xi}^{0}(\mathbb{Q}_{S}) \setminus G(\mathbb{Q}_{S})} \int_{M_{X}^{0}(\mathbb{Q}_{S}) \setminus G_{\Xi}^{0}(\mathbb{Q}_{S})} \times f_{S}((X+A) \cdot \operatorname{ad}(yx)) \times \right)$$

$$\times \left( \sum_{Q \in \mathcal{I}^{\Xi}(M_{\Xi}^{0})} \sum_{L \in \mathcal{L}^{M_{Q}}(M_{\Xi}^{0})} r_{M_{\Xi}^{0}}^{L} (\exp(X_{\operatorname{nil}}), \exp(A)) \times \right)$$

$$\times v_{L}^{Q}(y) v_{Q}'(kx, T_{0} - T_{\Xi,M}) dy dx \right)$$

$$(3.3.9.4) \qquad = \lim_{A \to 0} \left( |D^{G}(X+A)|_{S}^{1/2} |D^{G_{\Xi}}(X+A)|_{S}^{-1/2} \int_{G_{\Xi}^{0}(\mathbb{Q}_{S}) \setminus G(\mathbb{Q}_{S})} \right)$$

$$\times \left( \sum_{\mathbf{Q} \in \mathcal{F}^{\Xi}(\mathbf{M}_{\Xi}^{0})} \sum_{\mathbf{L} \in \mathcal{L}^{\mathbf{M}_{\mathbf{Q}}}(\mathbf{M}_{\Xi})} r_{\mathbf{M}_{\Xi}^{0}}^{\mathbf{L}} \left( \exp(X_{\mathrm{nil}}), \exp(A) \right) \times J_{\mathbf{L}}^{\mathbf{M}_{\mathbf{Q}}} (X_{\mathrm{nil}} + A, \Phi_{S,\mathbf{Q},y}^{T_{0} - T_{\Xi,\mathbf{M}}}) \right) \mathrm{d}y \right)$$
$$= |D^{\mathbf{G}}(\Xi)|_{S}^{1/2} \int_{\mathbf{G}_{\Xi}^{0}(\mathbb{Q}_{S}) \setminus \mathbf{G}(\mathbb{Q}_{S})} \left( \sum_{\mathbf{Q} \in \mathcal{F}^{\Xi}(\mathbf{M}_{\Xi}^{0})} J_{\mathbf{M}_{\Xi}}^{\mathbf{M}_{\mathbf{Q}}} (X_{\mathrm{nil}}, \Phi_{S,\mathbf{Q},y}^{T_{0} - T_{\Xi,\mathbf{M}}}) \right) \mathrm{d}y$$

where the equality (3.3.9.4) follows from the change of variables

$$(3.3.9.5) \qquad \qquad \mathbf{M}^{0}_{X}(\mathbb{Q}_{S}) \backslash \mathbf{G}^{0}_{\Xi}(\mathbb{Q}_{S}) \xrightarrow{\sim} \mathbf{M}_{X}(\mathbb{Q}_{S}) \backslash \mathbf{M}_{\mathbf{Q}^{0}_{\Xi}}(\mathbb{Q}_{S}) \times \mathbf{N}_{\mathbf{Q}}(\mathbb{Q}_{S}) \times \mathbf{K}_{\Xi}$$

for each parabolic subgroup Q in  $\mathcal{F}^{\Xi}(M^0_{\Xi})$ .

(3.3.10) Definition Let  $\equiv$  be the equivalence relation on  $\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o}$  defined by

$$\begin{array}{ll} (3.3.10.1) & \forall X \in \mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o} \quad \forall Y \in \mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o} \quad X \equiv Y \quad \text{if} \\ & \exists \delta \in \mathcal{M}(\mathbb{Q}) \quad \exists \eta \in \mathcal{M}^0_{X_{\text{ss}}}(\mathbb{Q}_S) \quad X_{\text{ss}} = Y_{\text{ss}} \cdot \operatorname{ad}(\delta) \quad X_{\text{nil}} = Y_{\text{nil}} \cdot \operatorname{ad}(\delta\eta). \end{array}$$

Let  $(\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o})_{M,S}$  denote the collection of  $\equiv$  equivalence classes in  $\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o}$ . By abuse of notation denote the  $\equiv$  equivalence class of an element X in  $\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o}$  by the same symbol X.

(3.3.11) **Proposition** Let  $\mathfrak{o}$  be  $a \sim$  equivalence class on  $\mathfrak{g}(\mathbb{Q})$ . Let  $S_{\mathfrak{o}}$  be a sufficiently large finite set of places of  $\mathbb{Q}$ . Then for each standard Levi subgroup M of G and for each  $\equiv$  equivalence class X in  $\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o}$  there exists a constant  $a^{\mathrm{M}}(S_{\mathfrak{o}}, X)$  such that

 $(3.3.11.1) \qquad \forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A}))$ 

$$J^{\mathcal{G}}_{\mathfrak{o}}(f) = \sum_{\mathcal{M} \in \mathcal{L}} |\mathcal{W}^{\mathcal{M}}_{0}| |\mathcal{W}^{\mathcal{G}}_{0}|^{-1} \sum_{X \in (\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o})_{\mathcal{M}, S_{\mathfrak{o}}}} a^{\mathcal{M}}(S_{\mathfrak{o}}, X) J^{\mathcal{G}}_{\mathcal{M}}(X, f_{S_{\mathfrak{o}}}).$$

*Proof.* Retain the notations of (3.3.6.1) and (3.3.9.1). Let  $\Xi$  be equal to  $\Sigma$ , hence

(3.3.11.2) 
$$X = \Sigma + \nu \in \mathfrak{m}(\mathbb{Q}_S)/\mathrm{ad}(\mathrm{M}(\mathbb{Q}_S)),$$

and

$$(3.3.11.3) T_0 - T_{\Xi,M} = T_0 - T_{\Sigma,1}.$$

Without loss of generality assume that the finite set  $S_{\mathfrak{o}}$  is large enough such that

$$\begin{array}{ll} (3.3.11.4) & \forall p \notin S_{\mathfrak{o}} & \bullet \Sigma \in \mathfrak{g}(\mathbb{Z}_p); \\ & \bullet |D(\Sigma)|_p = 1; \end{array}$$

•
$$\mathfrak{g}(\mathbb{Z}_p) \cdot \mathrm{ad}(\mathrm{K}_p) = \mathfrak{g}(\mathbb{Z}_p);$$
  
• $\mathrm{K}_p \cap \mathrm{G}^0_{\Sigma}(\mathbb{Q}_p) = \mathrm{K}_{\Sigma,p}$  is hyperspecial in  $\mathrm{G}^0_{\Sigma}(\mathbb{Q}_p);$   
• $\forall x_p \in \mathrm{G}(\mathbb{Q}_p) \ (\Sigma + \mathfrak{g}_{\Sigma,\mathrm{nil}}(\mathbb{Q}_p)) \cdot \mathrm{ad}(x_p) \cap \mathfrak{g}(\mathbb{Z}_p) \neq \emptyset$   
 $\Rightarrow x_p \in \mathrm{G}^0_{\Sigma}(\mathbb{Q}_p)\mathrm{K}_p.$ 

The last condition in (3.3.11.4) is satisfied for large enough  $S_{\mathfrak{o}}$  by Lemma 6.1 of [Ar86]. Since

(3.3.11.5) 
$$\left( H \mapsto \Gamma'_{\mathcal{Q}} \left( H - T_{\Sigma,1}, \mathcal{Y}^{T_0}_{\mathcal{Q}}(k, x) \right) \right)^{\wedge} = v'_{\mathcal{Q}}(kx, T_0 - T_{\Sigma,1})(\lambda)$$

the conditions (3.3.11.4) imply that

$$(3.3.11.6) \qquad \Phi_{\mathbf{Q},x}^{T_0-T_{\Sigma,1}} = \Phi_{S_{\mathfrak{o}},\mathbf{Q},x_{S_{\mathfrak{o}}}}^{T_0-T_{\Sigma,1}} \otimes \bigotimes_{p \notin S_{\mathfrak{o}}} \mathbb{I}_{\mathbf{G}_{\Sigma}^0(\mathbb{Q}_p)\mathbf{K}_p}(x) \cdot \mathbb{I}_{\mathfrak{m}_{\mathbf{Q}}(\mathbb{Z}_p)}.$$

By(3.3.6.1)) and (3.2.7.1),

$$(3.3.11.7) \qquad J_{\mathfrak{o}}^{\mathrm{G}}(f) = |\pi_{0}(\mathrm{G}_{\Sigma})|^{-1} \int_{\mathrm{G}_{\Sigma}^{0}(\mathbb{Q}_{S\mathfrak{o}})\backslash\mathrm{G}(\mathbb{Q}_{S\mathfrak{o}})} \left(\sum_{\mathrm{Q}\in\mathcal{F}^{\Sigma}} \sum_{\mathrm{L}\in\mathcal{L}^{\mathrm{M}_{\mathrm{Q}}}(\mathrm{M}_{1,\Sigma}^{0})} |\mathrm{W}_{1}^{\mathrm{L}}||\mathrm{W}_{1}^{\mathrm{G}_{\Sigma}^{0}}|^{-1} \times \sum_{\nu\in(\mathfrak{l}_{\mathrm{nil}}(\mathbb{Q}))_{\mathrm{L},S\mathfrak{o}}} a^{\mathrm{L}}(S_{\mathfrak{o}},\nu) J_{\mathrm{L}}^{\mathrm{M}_{\mathrm{Q}}}(\nu,\Phi_{S\mathfrak{o},\mathrm{Q},x}^{T_{0}-T_{\Sigma,1}})\right) \mathrm{d}x$$

$$(3.3.11.8) = |\pi_{0}(\mathrm{G}_{\Sigma})|^{-1} \sum_{\mathrm{M}\in\mathring{\mathcal{L}}_{\Sigma}(\mathrm{M}_{1})} |\mathrm{W}_{1}^{\mathrm{M}_{\Sigma}^{0}}||\mathrm{W}_{1}^{\mathrm{G}_{\Sigma}^{0}}|^{-1} \times \sum_{\nu\in(\mathfrak{m}_{\Sigma,\mathrm{nil}}(\mathbb{Q}))_{\mathrm{M}_{\Sigma}^{0},S\mathfrak{o}}} a^{\mathrm{M}_{\Sigma}^{0}}(S_{\mathfrak{o}},\nu) J_{\mathrm{M}}^{\mathrm{G}}(\Sigma+\nu,f_{S\mathfrak{o}})$$

where on the right hand side of (3.3.11.8) the symbol  $\mathring{\mathcal{L}}_{\Sigma}(M_1)$  denotes the set

(3.3.11.9) 
$$\mathring{\mathcal{L}}_{\Sigma}(M_1) = \left\{ M \in \mathcal{L}(M_1) : A_M = A_{M_{\Sigma}^0} \right\}.$$

The equality (3.3.11.8) follows from (3.3.9.1) and the bijection

(3.3.11.10) 
$$\pi_{\Sigma}: \quad \mathring{\mathcal{L}}_{\Sigma}(\mathbf{M}_{1}) \xrightarrow{\sim} \mathcal{L}^{\mathbf{G}_{\Sigma}^{0}}(\mathbf{M}_{1,\Sigma}^{0})$$

defined by

(3.3.11.11) 
$$\forall \mathbf{M} \in \mathring{\mathcal{L}}_{\Sigma}(\mathbf{M}_{1}) \qquad \pi_{\Sigma}(\mathbf{M}) = \mathbf{M}_{\Sigma}^{0} = \mathbf{L}$$

and the equation

$$(3.3.11.12) |D(\Sigma)|_{S_{\mathfrak{o}}} = 1$$

which follows from (3.3.11.4).

Define the constant  $a^{\mathrm{M}}(S_{\mathfrak{o}}, X)$  by

$$(3.3.11.13) a^{\mathcal{M}}(S_{\mathfrak{o}}, X) = |\pi_{0}(\mathcal{M}_{\Sigma})|^{-1} \sum_{\substack{\nu \in (\mathfrak{m}_{\Sigma, \operatorname{nil}})_{\mathcal{M}_{\Sigma}^{0}, S_{\mathfrak{o}}} \\ \Sigma + \nu \equiv X \operatorname{mod}(\mathcal{M}, S_{\mathfrak{o}})}} a^{\mathcal{M}_{\Sigma}^{0}}(S_{\mathfrak{o}}, \nu)$$

if  $X_{ss}$  is  $\mathbb{Q}$ -elliptic in  $\mathfrak{m}(\mathbb{Q})$  and zero otherwise. For a fixed  $\Sigma$  and let M vary in  $\mathcal{L}(M_1)$ , then  $\Sigma$  is  $\mathbb{Q}$ -elliptic in  $\mathfrak{m}(\mathbb{Q})$  if and only if  $A_M$  and  $A_{M_{\Sigma}}$  are equal, hence by (3.3.11.7)

$$(3.3.11.14) J_{\mathfrak{o}}^{\mathrm{G}}(f) = \sum_{\mathrm{M}\in\mathcal{L}(\mathrm{M}_{1})} |\mathrm{W}_{1}^{\mathrm{M}_{\Sigma}^{0}}| |\mathrm{W}_{1}^{\mathrm{G}_{\Sigma}^{0}}|^{-1} |\pi_{0}(\mathrm{M}_{\Sigma})| |\pi_{0}(\mathrm{G}_{\Sigma})|^{-1} \times \\ \times \sum_{\substack{X\in(\mathfrak{m}(\mathbb{Q})\cap\mathfrak{o})_{\mathrm{M},S_{\mathfrak{o}}}\\X_{\mathrm{ss}}=\Sigma}} a^{\mathrm{M}}(S_{\mathfrak{o}}, X) J_{\mathrm{M}}^{\mathrm{G}}(X, f_{S_{\mathfrak{o}}}).$$

The set

(3.3.11.15) 
$$\begin{cases} M \in \mathcal{L}, \sigma \text{ is a semisimple } M(\mathbb{Q})-\\ (M, \sigma): & \text{-orbit in } \mathfrak{m}(\mathbb{Q}), \Sigma \text{ is conjugate}\\ & \text{to a point in } \sigma \text{ under } G(\mathbb{Q}). \end{cases}$$

admits a natural action by  $W_0^G$  which preserves  $a^M(S_{\mathfrak{o}}, X)$  and  $J_M^G(X, f_{S_{\mathfrak{o}}})$ , where each pair  $(M, \sigma)$  has stablizer  $W_0^M$ . Denote by  $W_1^{G_{\Sigma}}$  the quotient of the normalizer of  $A_1$  in  $G_{\Sigma}$  by  $M_{1,\Sigma}$ . There is an exact sequence

$$(3.3.11.16) 1 \longrightarrow W_1^{G_{\Sigma}^0} \longrightarrow W_1^{G_{\Sigma}} \longrightarrow \pi_0(G_{\Sigma}) \longrightarrow 1.$$

Similar notations apply to  $M_{\Sigma}$ . The subset of (3.3.11.15)

(3.3.11.17) 
$$\left\{ (\mathbf{M}, \sigma) : \mathbf{M} \in \mathcal{L}(\mathbf{M}_1) \text{ and } \sigma = \Sigma \cdot \mathrm{ad}(\mathbf{M}(\mathbb{Q})) \right\}$$

admits a natural action of  $W_1^{G_{\Sigma}}$ , where each pair  $(M, \sigma)$  has stablizer  $W_1^{M_{\Sigma}}$ . The quotients

(3.3.11.18) 
$$\begin{cases} M \in \mathcal{L}, \sigma \text{ is a semisimple } M(\mathbb{Q})\text{-} \\ (M, \sigma) : & \text{-orbit in } \mathfrak{m}(\mathbb{Q}), \Sigma \text{ is conjugate} \\ & \text{to a point in } \sigma \text{ under } G(\mathbb{Q}). \end{cases} \\ \end{cases} / W_0^G$$

and

(3.3.11.19) 
$$\left\{ \begin{array}{l} (\mathbf{M},\sigma): \ \mathbf{M} \in \mathcal{L}(\mathbf{M}_1) \text{ and } \sigma = \Sigma \cdot \mathrm{ad}(\mathbf{M}(\mathbb{Q})) \end{array} \right\} \middle/ \mathbf{W}_1^{\mathbf{G}_{\Sigma}}$$

are in bijection, therefore

(3.3.11.20) 
$$J^{\rm G}_{\mathfrak{o}}(f)$$

$$= \sum_{\mathbf{M}\in\mathcal{L}(\mathbf{M}_{1})} |\mathbf{W}_{1}^{\mathbf{M}_{\Sigma}}| |\mathbf{W}_{1}^{\mathbf{G}_{\Sigma}}|^{-1} \sum_{\substack{X\in(\mathfrak{m}(\mathbb{Q})\cap\mathfrak{o})_{\mathbf{M},S_{\mathfrak{o}}}\\X_{ss}=\Sigma}} a^{\mathbf{M}}(S_{\mathfrak{o}}, X) J_{\mathbf{M}}^{\mathbf{G}}(X, f_{S_{\mathfrak{o}}})}$$

$$= \sum_{\mathbf{M}\in\mathcal{L}} \frac{|\mathbf{W}_{0}^{\mathbf{M}}|}{|\mathbf{W}_{0}^{\mathbf{G}}|} \left(\frac{|\mathbf{W}_{1}^{\mathbf{M}_{\Sigma}}|}{|\mathbf{W}_{1}^{\mathbf{G}_{\Sigma}}|}\right)^{-1} |\mathbf{W}_{1}^{\mathbf{M}_{\Sigma}}| |\mathbf{W}_{1}^{\mathbf{G}_{\Sigma}}|^{-1} \times$$

$$\times \sum_{\substack{X\in(\mathfrak{m}(\mathbb{Q})\cap\mathfrak{o})_{\mathbf{M},S_{\mathfrak{o}}}} a^{\mathbf{M}}(S_{\mathfrak{o}}, X) J_{\mathbf{M}}^{\mathbf{G}}(X, f_{S_{\mathfrak{o}}})$$

$$= \sum_{\mathbf{M}\in\mathcal{L}} |\mathbf{W}_{0}^{\mathbf{M}}| |\mathbf{W}_{0}^{\mathbf{G}}|^{-1} \sum_{\substack{X\in(\mathfrak{m}(\mathbb{Q})\cap\mathfrak{o})_{\mathbf{M},S_{\mathfrak{o}}}} a^{\mathbf{M}}(S_{\mathfrak{o}}, X) J_{\mathbf{M}}^{\mathbf{G}}(X, f_{S_{\mathfrak{o}}}).$$

(3.3.12) **Remark** For a semisimple element X in  $\mathfrak{o}$ ,

(3.3.12.1)  $a^{\mathcal{M}}(S_{\mathfrak{o}}, X) = |\pi_0(\mathcal{M}_X)|^{-1} \mathrm{Vol}\big(\mathcal{M}^0_X(\mathbb{Q}) \setminus \mathcal{M}^0_X(\mathbb{A})^1\big)$ 

if X is  $\mathbb{Q}$ -elliptic in  $\mathfrak{m}(\mathbb{Q})$ , and vanishes otherwise. If X is in addition assumed to be regular, the identity (3.3.11.1) reduces to (2.2.11.1).

#### 3.4 The refined trace formula

(3.4.1) **Proposition** (Refined trace formula)

For each sufficiently large finite set S of places of  $\mathbb{Q}$ , for each ~ equivalence class  $\mathfrak{o}$  in  $\mathfrak{g}(\mathbb{Q})$ , for each  $\equiv$  equivalence class X in  $(\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o})_{M,S}$ , there exists a constant  $a^M(S,X)$  such that

$$(3.4.1.1) \qquad \forall f \in \mathcal{S}(\mathfrak{g}(\mathbb{A}))$$

$$\begin{split} &\lim_{S} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} \sum_{\mathbf{M} \in \mathcal{L}} |\mathbf{W}_{0}^{\mathbf{M}}| |\mathbf{W}_{0}^{\mathbf{G}}|^{-1} \sum_{X \in (\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o})_{\mathbf{M},S}} a^{\mathbf{M}}(S,X) J_{\mathbf{M}}^{\mathbf{G}}(X,f_{S}) \\ &= \lim_{S} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} \sum_{\mathbf{M} \in \mathcal{L}} |\mathbf{W}_{0}^{\mathbf{M}}| |\mathbf{W}_{0}^{\mathbf{G}}|^{-1} \sum_{X \in (\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o})_{\mathbf{M},S}} a^{\mathbf{M}}(S,X) J_{\mathbf{M}}^{\mathbf{G}}(X,f_{S}^{*}). \end{split}$$

*Proof.* By (2.2.7.1) it is enough to show that the left hand side of (3.4.1.1) is equal to

(3.4.1.2) 
$$J(f) = \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} J_{\mathfrak{o}}(f)$$

For each individual class  $\boldsymbol{o}$  by (3.3.11.1)

(3.4.1.3) 
$$J_{\mathfrak{o}}(f) = \lim_{S} \sum_{M} |W_{0}^{M}| |W_{0}^{G}|^{-1} \sum_{X} a^{M}(S, X) J_{M}^{G}(X, f_{S})$$

where the limit stablizes as S grows large enough.

**Lemma** Let  $\Gamma$  be a compact subset of  $\mathfrak{g}(\mathbb{A})$ . Then there are only finitely many classes  $\mathfrak{o}$  such that  $\mathfrak{o} \cdot \mathrm{ad}(\mathrm{G}(\mathbb{A}))$  intersects  $\Gamma$ .

*Proof.* The lemma follows from Corollary A.2 of [Ar86] which states that there exists a compact set  $G_{\Gamma}$  contained in  $G(\mathbb{A})^1$  such that

(3.4.1.4) 
$$\forall x \in \mathcal{G}(\mathbb{A})^1 - \mathcal{G}(\mathbb{Q})\mathcal{G}_{\Gamma} \qquad \mathfrak{g}(\mathbb{Q})' \cdot \mathrm{ad}(x) \cap \Gamma = \emptyset$$

where  $\mathfrak{g}(\mathbb{Q})'$  denotes the set of points of  $\mathfrak{g}(\mathbb{Q})$  not contained in any proper parabolic subalgebra. This is established by a reduction theory argument.

To prove the lemma first consider a class  $\mathfrak{o}$  contained in  $\mathfrak{g}(\mathbb{Q})'$ . There exists a point X in  $\mathfrak{o}$  such that  $X \cdot \mathrm{ad}(\mathrm{G}_{\Gamma})$  is contained in  $\Gamma$ , hence there are only finitely many such X, hence finitely many such  $\mathfrak{o}$ .

Next for an arbitrary class  $\mathfrak{o}$  there is some Levi subalgebra  $\mathfrak{m}$  such that  $\mathfrak{o}$  intersects  $\mathfrak{m}(\mathbb{Q})'$ . Then by the Iwasawa decomposition

$$\begin{array}{ll} (3.4.1.5) & \quad \mathfrak{o} \cdot \mathrm{ad}(\mathrm{G}(\mathbb{A})^1) \\ &= & \quad \mathfrak{o} \cdot \mathrm{ad}(\mathrm{M}(\mathbb{A})^1) \cdot \mathrm{ad}(\mathrm{N}(\mathbb{Q}) \backslash \mathrm{N}(\mathbb{A})) \cdot \mathrm{ad}(\mathrm{K}) \\ &= & \quad (\mathfrak{o} \cap \mathfrak{m}(\mathbb{Q})') \cdot \mathrm{ad}(\mathrm{M}_{\Gamma \cap \mathfrak{m}(\mathbb{A})}) \cdot \mathrm{ad}(\mathrm{N}(\mathbb{Q}) \backslash \mathrm{N}(\mathbb{A})) \cdot \mathrm{ad}(\mathrm{K}), \end{array}$$

the lemma follows by induction.

The Schwartz function f is compactly supported if its component at infinity  $f_{\infty}$  is compactly supported. For such an f the limit on the left hand side of (3.4.1.1) stablizes for every S such that

$$(3.4.1.6) S \supset \bigcup_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} S_{\mathfrak{o}}$$

where the union is supported on a finite set of  $\mathfrak{o}$ . Hence for a compactly supported f the left hand side of (3.4.1.1) is equal to J(f).

However the function spaces  $C_c^{\infty}(\mathfrak{g}(\mathbb{R}))$  and hence  $C_c^{\infty}(\mathfrak{g}(\mathbb{Q}_S))$  are dense in  $\mathcal{S}(\mathfrak{g}(\mathbb{R}))$  and  $\mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$  respectively, therefore  $C_c^{\infty}(\mathfrak{g}(\mathbb{A}))$  is dense in  $\mathcal{S}(\mathfrak{g}(\mathbb{A}))$  which by (1.1.4.2) is defined as

$$(3.4.1.7) \qquad \qquad \mathcal{S}(\mathfrak{g}(\mathbb{A})) = \varinjlim \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$$

equipped with the final topology.

Since J(f) extends continuously to all Schwartz functions, the limit on the left hand side of (3.4.1.1) exists, and the interchangeability of the operations of taking the limit as Sapproaches infinity and taking the sum over all classes  $\mathfrak{o}$  extends from  $C_c^{\infty}(\mathfrak{g}(\mathbb{A}))$  to  $\mathcal{S}(\mathfrak{g}(\mathbb{A}))$ . This establishes the identity (3.4.1.1) for all Schwartz functions f on  $\mathfrak{g}(\mathbb{A})$ .

## 4 A simple invariant trace formula

In this chapter following the arguments in §7 of [Ar88c] the refined trace formula (3.4.1.1) is reduced to an identity between invariant distributions on  $\mathfrak{g}(\mathbb{A})$  for a suitably restricted class of test functions.

#### 4.1 Parabolic descent and parabolic induction

(4.1.1) **Definition** Let P be a parabolic subgroup in  $\mathcal{F}$  with Levi component M and unipotent radical N. Let  $f_S$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$ . Define  $f_{S,P}$  to be the Schwartz function on  $\mathfrak{m}(\mathbb{Q}_S)$  by *parabolic descent* along P

(4.1.1.1) 
$$\forall X \in \mathfrak{m}(\mathbb{Q}_S)$$
$$f_{S,P}(X) = \int_{\mathcal{K}_S} \int_{\mathfrak{n}(\mathbb{Q}_S)} f_S((X+N) \cdot \mathrm{ad}(k)) \, \mathrm{d}N \mathrm{d}k.$$

For each parabolic subgroup  $P_S$  in  $\mathcal{F}^{G_S}$  define the Schwartz function  $f_{S,P_S}$  on  $\mathfrak{m}_S(\mathbb{Q}_S)$  analogously.

(4.1.2) **Remark** Parabolic descent preserves orbital integrals, i.e.

(4.1.2.1) 
$$\forall X \in \mathfrak{m}_{\mathrm{reg},\mathrm{ss}}(\mathbb{Q}_S) \qquad I_{\mathrm{G}}^{\mathrm{G}}(X, f_S) = I_{\mathrm{M}}^{\mathrm{M}}(X, f_{S,\mathrm{P}}).$$

See §13.12 of [Ko05]. Hence the orbital integrals of  $f_{S,P}$  on  $\mathfrak{m}_{reg,ss}(\mathbb{Q}_S)$  is independent of the choice of the parabolic subgroup P in  $\mathcal{P}(M)$ . In this case denote  $f_{S,P}$  by the alternative notation  $f_{S,M}$  such that

(4.1.2.2) 
$$\forall X \in \mathfrak{m}_{\mathrm{reg},\mathrm{ss}}(\mathbb{Q}_S) \qquad I^{\mathrm{M}}_{\mathrm{M}}(X, f_{S,\mathrm{M}}) = I^{\mathrm{M}}_{\mathrm{M}}(X, f_{S,\mathrm{P}})$$

is well-defined.

(4.1.3) **Remark** With compatible choices of Fourier transforms on  $\mathfrak{g}(\mathbb{Q}_S)$  and  $\mathfrak{m}(\mathbb{Q}_S)$ , parabolic descent intertwines the Fourier transforms, i.e.

$$(4.1.3.1) (f_S)_P = (f_{S,P})^{\hat{}}.$$

See §13.13 of [Ko05].

(4.1.4) **Definition** Let P be a parabolic subgroup in  $\mathcal{F}$  with Levi component M and unipotent radical N. Let X be an element in  $\mathfrak{m}(\mathbb{Q}_S)$ . Define  $\mathrm{Ind}_{M,P}^G(X)$  to be the  $\mathrm{ad}(G(\mathbb{Q}_S))$ invariant subset of  $\mathfrak{g}(\mathbb{Q}_S)$  by *parabolic induction* along P

(4.1.4.1) 
$$\operatorname{Ind}_{M,P}^{G}(X) = \left( \left( X \cdot \operatorname{ad}(M) + \mathfrak{n} \right) \cdot \operatorname{ad}(G) \right)_{\operatorname{reg}}(\mathbb{Q}_{S})$$

where the subscript reg denotes the regular locus, the set of points with minimal dimensional isotropy group in G.

(4.1.5) **Remark** With a fixed choice of M, parabolic induction along P is independent of the choice of the parabolic subgroup P in  $\mathcal{P}(M)$ , hence adopt the alternative notation  $\operatorname{Ind}_{M}^{G}(X)$ . For a proof see Satz 2.6 of [Bo81]. If the inducing Levi subgroup M is understood tacitly, denote  $\operatorname{Ind}_{M}^{G}(X)$  by the alternative notation  $X^{G}$ .

(4.1.6) **Remark** If X is regular semisimple then  $\operatorname{Ind}_{M}^{G}(X)$  is equal to the  $\operatorname{ad}(G(\mathbb{Q}_{S}))$ -orbit of X. In general  $\operatorname{Ind}_{M}^{G}(X)$  is a finite union of  $\operatorname{ad}(G(\mathbb{Q}_{S}))$ -orbits which are geometrically conjugate to each other, but  $\operatorname{Ind}_{M}^{G}(X)$  is still called the *induced orbit* of X. See page 255 of [Ar88a] and §2.1 of [Bo81]. If L is a Levi subgroup in  $\mathcal{L}(M)$ , define the invariant weighted orbital integral along the induced orbit  $X^{L}$  by

(4.1.6.1) 
$$\forall f_S \in \mathfrak{g}(\mathbb{Q}_S) \qquad J_{\mathrm{L}}^{\mathrm{G}}(X^{\mathrm{L}}, f_S) = \sum_{\substack{\Omega \subset X^{\mathrm{L}} \\ \mathrm{ad}(\mathrm{L}(\mathbb{Q}_S)) \text{-orbit}}} J_{\mathrm{L}}^{\mathrm{G}}(\Omega, f_S).$$

(4.1.7) **Remark** Parabolic induction is transitive in nested chains of Levi subgroups, i.e.

(4.1.7.1)  $\forall \mathbf{L} \in \mathcal{L}^{\mathbf{G}}(\mathbf{M}) \qquad \mathrm{Ind}_{\mathbf{M}}^{\mathbf{G}}(X) = \mathrm{Ind}_{\mathbf{L}}^{\mathbf{G}}(\mathrm{Ind}_{\mathbf{M}}^{\mathbf{L}}(X)).$ 

See  $\S2.3$  of [Bo81].

(4.1.8) **Remark** Parabolic induction is compatible with Jordan decomposition, i.e.

(4.1.8.2) 
$$\operatorname{Ind}_{M}^{G}(X) = \left(X_{ss} + \operatorname{Ind}_{M_{X_{ss}}}^{G_{X_{ss}}^{0}}(X_{nil})\right) \cdot \operatorname{ad}(G(\mathbb{Q}_{S})).$$

See §2.4 of [Bo81]. Also see Lemma 2 of [Ho13].

(4.1.9) Lemma (Descent and splitting of (G, M)-families) Let M be a Levi subgroup in  $\mathcal{L}$ . There exist a function  $d_{M}^{G}(,)$ 

 $(4.1.9.1) d_{\mathbf{M}}^{\mathbf{G}}: \quad \mathcal{L}(\mathbf{M}) \times \mathcal{L}(\mathbf{M}) \longrightarrow \mathbb{R},$ 

and a partially defined map s(, )

$$(4.1.9.2) \qquad s: \quad \mathcal{L}(\mathbf{M}) \times \mathcal{L}(\mathbf{M}) \longrightarrow \quad \mathcal{F}(\mathbf{M}) \times \mathcal{F}(\mathbf{M})$$

whose domain contains the pairs  $(L_1, L_2)$  for which  $d_M^G(L_1, L_2)$  is nonzero, such that

• if  $(L_1, L_2)$  is contained in the domain of s, then

$$(4.1.9.3) \qquad \qquad s(\mathbf{L}_1, \mathbf{L}_2) \in \mathcal{P}(\mathbf{L}_1) \times \mathcal{P}(\mathbf{L}_2);$$

• if  $(c_P)$  is a (G, M)-family and L is a Levi subgroup in  $\mathcal{L}(M)$ , then

(4.1.9.4) 
$$c_{\rm L} = \sum_{{\rm L}' \in \mathcal{L}({\rm M})} d_{\rm M}^{\rm G}({\rm L},{\rm L}') c_{\rm M}^{\rm Q}$$

where Q' denotes the second component of s(L, L');

• if  $(c_{\rm P})$  and  $(d_{\rm P})$  are (G, M)-families, then

(4.1.9.5) 
$$(cd)_{\mathrm{M}} = \sum_{\mathrm{L}_{1},\mathrm{L}_{2}\in\mathcal{L}(\mathrm{M})} d_{\mathrm{M}}^{\mathrm{G}}(\mathrm{L}_{1},\mathrm{L}_{2}) c_{\mathrm{M}}^{\mathrm{Q}_{1}} d_{\mathrm{M}}^{\mathrm{Q}_{2}}$$

where

$$(4.1.9.6) (Q_1, Q_2) = s(L_1, L_2).$$

Analogous results hold for the groups  $G_v$  and  $G_s$ .

*Proof.* See  $\S7$  of [Ar88b].

(4.1.10) **Remark** The constant  $d_M^G(L_1, L_2)$  is defined to be the volume in  $\mathfrak{a}_M^G$  of the image of a fundamental parallelotope in the direct sum of  $\mathfrak{a}_M^{L_1}$  and  $\mathfrak{a}_M^{L_2}$  under the natural map

$$(4.1.10.1) \qquad \qquad \mathfrak{a}_{\mathrm{M}}^{\mathrm{L}_{1}} \oplus \mathfrak{a}_{\mathrm{M}}^{\mathrm{L}_{2}} \longrightarrow \mathfrak{a}_{\mathrm{M}}^{\mathrm{G}}$$

if (4.1.10.1) is an isomorphism, and zero otherwise. In the former case  $d_M^G(L_1, L_2)$  is equal to the volume in  $\mathfrak{a}_{L_2}^G$  of the image of a fundamental parallelotope in  $\mathfrak{a}_M^{L_1}$  under the natural isomorphism

$$(4.1.10.2) \qquad \qquad \mathfrak{a}_{\mathrm{M}}^{\mathrm{L}_{1}} \longrightarrow \mathfrak{a}_{\mathrm{L}_{2}}^{\mathrm{G}}.$$

(4.1.11) **Remark** The map s depends on the choice of a vector  $\xi$  in general position in  $\mathfrak{a}_{M}^{G}$ . Let  $L_1$  and  $L_2$  be Levi subgroups in  $\mathcal{L}(M)$  such that (4.1.10.1) is an isomorphism, then

(4.1.11.1) 
$$\exists \xi_1 \in \mathfrak{a}_{L_1}^G \ \exists \xi_2 \in \mathfrak{a}_{L_2}^G \qquad \xi = \frac{\xi_1}{2} - \frac{\xi_2}{2}.$$

Define  $s(L_1, L_2)$  to be  $(Q_1, Q_2)$  where  $Q_i$  is the parabolic subgroup in  $\mathcal{P}(L_i)$  whose corresponding positive chamber in  $\mathfrak{a}_{L_i}^G$  contains the vector  $\xi_i$  where the index i is 1 or 2.

(4.1.12) Lemma (Descent and splitting of weighted orbital integrals) Let M be a Levi subgroup in  $\mathcal{L}$ , let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ . Let  $f_S$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$ . Let  $\xi$  be a vector in general position in  $\mathfrak{a}_M^G$ , let

$$(4.1.12.1) \qquad \qquad d_{\mathrm{M}}^{\mathrm{G}} : \mathcal{L}^{\mathrm{G}}(\mathrm{M}) \times \mathcal{L}^{\mathrm{G}}(\mathrm{M}) \longrightarrow \mathbb{R}$$
$$s : \mathcal{L}^{\mathrm{G}}(\mathrm{M}) \times \mathcal{L}^{\mathrm{G}}(\mathrm{M}) \longrightarrow \mathcal{F}^{\mathrm{G}}(\mathrm{M}) \times \mathcal{F}^{\mathrm{G}}(\mathrm{M})$$

be defined as in Remark (4.1.10) and Remark (4.1.11) with respect to  $\xi$ .

• Let L be a Levi subgroup in  $\mathcal{L}^{G}(M)$ , then

(4.1.12.2) 
$$J_{\rm L}^{\rm G}(X^{\rm L}, f_S) = \sum_{{\rm L}' \in \mathcal{L}({\rm M})} d_{\rm M}^{\rm G}({\rm L}, {\rm L}') J_{\rm M}^{\rm L'}(X, f_{S, {\rm Q}'})$$

where Q' denotes the second component of s(L, L').

• Let S be the set  $\{v_1, v_2\}$ . Let  $f_S$  be of the form  $f_{v_1} \otimes f_{v_2}$  where  $f_{v_i}$  is a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_{v_i})$  where the index i is 1 or 2. Then

(4.1.12.3) 
$$J_{\mathrm{M}}^{\mathrm{G}}(X, f_{S}) = \sum_{\mathrm{L}_{1}, \mathrm{L}_{2} \in \mathcal{L}^{(\mathrm{M})}} d_{\mathrm{M}}^{\mathrm{G}}(\mathrm{L}_{1}, \mathrm{L}_{2}) J_{\mathrm{M}}^{\mathrm{L}_{1}}(X, f_{v_{1}, \mathrm{Q}_{1}}) J_{\mathrm{M}}^{\mathrm{L}_{2}}(X, f_{v_{2}, \mathrm{Q}_{2}})$$

where

$$(4.1.12.4) (Q_1, Q_2) = s(L_1, L_2).$$

Local identities analogous to (4.1.12.2) also hold for  $G_v$ .

*Proof.* For a regular semisimple X this follows from (4.1.9.4) and (4.1.9.5). For a general X this follows from the limit formula (3.1.5.4).

(4.1.13) Corollary (Descent and splitting of orbital integrals) Let X be an element of  $\mathfrak{g}(\mathbb{Q}_S)$ . Let  $f_S$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$ .

• Let L be a Levi subgroup in  $\mathcal{L}$  such that X is contained in  $\mathfrak{l}(\mathbb{Q}_S)$ , then

(4.1.13.1) 
$$I_{\rm G}^{\rm G}(X^{\rm G}, f_S) = I_{\rm L}^{\rm L}(X, f_{S,{\rm L}}).$$

• Let S be the set  $\{v_1, v_2\}$ . Let  $f_S$  be of the form  $f_{v_1} \otimes f_{v_2}$  where  $f_{v_i}$  is a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_{v_i})$  where the index i is 1 or 2. Then

(4.1.13.2) 
$$I_{\rm G}^{\rm G}(X, f_{v_1} \otimes f_{v_2}) = I_{\rm G}^{\rm G}(X, f_{v_1}) I_{\rm G}^{\rm G}(X, f_{v_2}).$$

Local identities analogous to (4.1.13.1) also hold for  $G_v$ .

*Proof.* The identity (4.1.13.1) follows from (4.1.12.2) where the only nonzero summand on the right hand side corresponds to the Levi subgroup L in  $\mathcal{L}(L)$ .

The identity (4.1.13.2) follows from (4.1.12.3) where the only summand on the right hand side corresponds to the pair (G, G) in  $\mathcal{L}(G) \times \mathcal{L}(G)$ .

(4.1.14) **Remark** If X is a regular semisimple element of  $\mathfrak{m}(\mathbb{Q})$  then the local analogue of (4.1.12.2) becomes

(4.1.14.1) 
$$J_{\mathcal{L}_{v}}^{\mathcal{G}_{v}}(X,f_{v}) = \sum_{\mathcal{L}_{v}'\in\mathcal{L}^{\mathcal{G}_{v}}(\mathcal{M}_{v})} d_{\mathcal{M}_{v}}^{\mathcal{G}_{v}}(\mathcal{L}_{v},\mathcal{L}_{v}') J_{\mathcal{M}_{v}}^{\mathcal{L}_{v}'}(X,f_{v,\mathcal{Q}_{v}'}).$$

Repeatedly applying the identities (4.1.12.3) and (4.1.14.1) reduces a regular semisimple weighted orbital integral  $J_{\mathrm{M}}^{\mathrm{G}}(X, )$  that appears generically in the refined trace formula (3.4.1.1) to a linear combination of products of local weighted orbital integrals  $J_{\mathrm{M}'_{v}}^{\mathrm{G}_{v}}(X, )$  where  $\mathrm{M}'_{v}$  is contained in  $\mathrm{M}_{v}$  and X is elliptic in  $\mathfrak{m}'_{v}(\mathbb{Q}_{v})$ .

#### 4.2 A simple form of the trace formula

(4.2.1) **Definition** Let v be a place of  $\mathbb{Q}$ , let  $f_v$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_v)$ . The function  $f_v$  is said to be *cuspidal* if it is a finite linear combination of Schwartz functions  $f_{v,i}$  on  $\mathfrak{g}(\mathbb{Q}_v)$  such that

(4.2.1.1)  $\forall \text{index } i \quad \forall \mathbf{P}_v \in \mathcal{F}^{\mathbf{G}_v} - \{\mathbf{G}_v\} \qquad f_{v,i,\mathbf{P}_v} = 0.$ 

Denote the space of cuspidal Schwartz functions on  $\mathfrak{g}(\mathbb{Q}_v)$  by  $\mathcal{S}_{\text{cusp}}(\mathfrak{g}(\mathbb{Q}_v))$ .

(4.2.2) **Remark** A Schwartz function  $f_v$  on  $\mathfrak{g}(\mathbb{Q}_v)$  is cuspidal if

$$(4.2.2.1) I_{\rm G}^{\rm G}(X, f_v) = 0$$

whenever X is a regular semisimple element of  $\mathfrak{g}(\mathbb{Q}_v)$  which is not  $\mathbb{Q}_v$ -elliptic.

(4.2.3) **Remark** The space  $S_{\text{cusp}}(\mathfrak{g}(\mathbb{Q}_v))$  is stable under the Fourier transform on  $\mathfrak{g}(\mathbb{Q}_v)$  by (4.1.3.1).

#### (4.2.4) **Proposition** (Simple invariant trace formula)

For each sufficiently large finite set S of places of  $\mathbb{Q}$ , for each  $\sim$  equivalence class  $\mathfrak{o}$  in  $\mathfrak{g}(\mathbb{Q})$ , for each  $\equiv$  equivalence class X in  $(\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o})_{M,S}$ , there exists a constant  $a^M(S,X)$  such that for each Schwartz function f on  $\mathfrak{g}(\mathbb{A})$  which is cuspidal at two distinct places of  $\mathbb{Q}$ 

(4.2.4.1)  
$$\lim_{S} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} \sum_{X \in (\mathfrak{g}(\mathbb{Q}) \cap \mathfrak{o})_{\mathrm{G},S}} a^{\mathrm{G}}(S,X) I^{\mathrm{G}}_{\mathrm{G}}(X,f_{S})$$
$$= \lim_{S} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} \sum_{X \in (\mathfrak{g}(\mathbb{Q}) \cap \mathfrak{o})_{\mathrm{G},S}} a^{\mathrm{G}}(S,X) I^{\mathrm{G}}_{\mathrm{G}}(X,f_{S}).$$

*Proof.* This argument follows the proof of Theorem 7.1(b) of [Ar88c].

Let  $v_1$  and  $v_2$  denote two places at which f is cuspidal. Without loss of generality the set S contains both  $v_1$  and  $v_2$ , and there exist two subsets  $S_1$  and  $S_2$  which partition S such that  $S_i$  contains  $v_i$  where the index i is 1 or 2.

With a choice of  $\xi$  as in Remark (4.1.11), it follows from (4.1.12.3) that

(4.2.4.2) 
$$J_{\mathrm{M}}^{\mathrm{G}}(X, f_{S}) = \sum_{\mathrm{L}_{1}, \mathrm{L}_{2} \in \mathcal{L}^{(\mathrm{M})}} d_{\mathrm{M}}^{\mathrm{G}}(\mathrm{L}_{1}, \mathrm{L}_{2}) J_{\mathrm{M}}^{\mathrm{L}_{1}}(X, f_{S_{1}, \mathrm{Q}_{1}}) J_{\mathrm{M}}^{\mathrm{L}_{2}}(X, f_{S_{2}, \mathrm{Q}_{2}})$$

where

$$(4.2.4.3) (Q_1, Q_2) = s(L_1, L_2).$$

Since f is cuspidal at  $v_1$  and  $v_2$ , each summand on the right hand side of (4.2.4.2) vanishes unless the parabolic subgroups  $Q_1$  and  $Q_2$  are both equal to G. Hence the Levi subgroups  $M_1$  and  $M_2$  are both equal to G, which implies that

$$(4.2.4.4) d_{\rm M}^{\rm G}({\rm L}_1,{\rm L}_2) = d_{\rm M}^{\rm G}({\rm G},{\rm G})$$

vanishes unless the Levi subgroup M is equal to G by Remark (4.1.10).

Then (4.2.4.1) follows from the refined trace formula (3.4.1.1) and Remark (4.2.3).

(4.2.5) **Remark** Each distribution that appears in the summands of the simple invariant trace formula (4.2.4.1) factorizes into local distributions that are invariant under the adjoint action of G on  $\mathfrak{g}$ .

# 5 The Harish-Chandra transform on the space of characteristic polynomials

In this chapter an integral transform on the space of characteristic polynomials satisfying a summation formula of Poisson type is constructed.

### 5.1 Preliminaries

(5.1.1) In this chapter G denotes the general linear group  $GL(n, \mathbb{Q})$  for some natural number n, with the standard choice of the minimal Levi subgroup  $M_0$  and the Borel subgroup B to be the subgroup consisting of the diagonal matrices and the upper triangular matrices respectively.

(5.1.2) Let  $\mathcal{A}_{G}$  denote the affine space of characteristic polynomials of  $n \times n$  matrices over  $\mathbb{Q}$ , i.e.

(5.1.2.1) 
$$\mathcal{A}_{\mathrm{G}} = \mathrm{gl}(n, \mathbb{Q}) /\!\!/ \mathrm{GL}(n, \mathbb{Q})$$

where  $/\!\!/$  denotes the affine quotient and  $\operatorname{GL}(n, \mathbb{Q})$  acts on  $\operatorname{gl}(n, \mathbb{Q})$  from the right by conjugation. The discriminant function D on  $\operatorname{gl}(n, \mathbb{Q})$  descends to a polynomial on  $\mathcal{A}_{\mathrm{G}}$ , denote by  $\mathcal{A}_{\mathrm{G,reg}}$  the open subset where D does not vanish.

Let M be a standard Levi subgroup of G. Denote by  $\mathcal{A}_{M}$  the affine quotient of  $\mathfrak{m}$  by the adjoint action of M, then  $\mathcal{A}_{M}$  is an affine space and there exists a partition  $(n_1, n_2, \ldots, n_r)$  of n such that

(5.1.2.2) 
$$\mathcal{A}_{\mathrm{M}} = \mathcal{A}_{\mathrm{G}_1} \times \mathcal{A}_{\mathrm{G}_2} \times \cdots \times \mathcal{A}_{\mathrm{G}_r}.$$

The embedding of  $\mathfrak{m}$  into  $\mathfrak{g}$  induces a map

(5.1.2.3) 
$$\pi_{\mathrm{M}}: \mathcal{A}_{\mathrm{M}} \longrightarrow \mathcal{A}_{\mathrm{G}}$$

which is finite of degree  $|W_0^M|^{-1}|W_0^G|$  and étale over  $\mathcal{A}_{G,reg}$ .

(5.1.3) Let v be a place of  $\mathbb{Q}$ . Equip  $\mathcal{A}_{G}(\mathbb{Q}_{v})$  with the measure

(5.1.3.1) 
$$|D(X_v)|_v^{-1/2} \mathrm{d}X_v$$

where  $dX_v$  denotes the standard translation invariant measure on the affine space  $\mathcal{A}_G(\mathbb{Q}_v)$ . This is equal to the pushforward of the translation invariant measure on  $\mathfrak{m}_0(\mathbb{Q}_v)$  along the Chevalley morphism

(5.1.3.2) 
$$\pi_{M_0}: \quad \mathfrak{m}_0 = \mathcal{A}_{M_0} \quad \longrightarrow \quad \mathfrak{m}_0 /\!\!/ W_0^G = \mathcal{A}_G.$$

The complement of  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$  in  $\mathcal{A}_G(\mathbb{Q}_v)$  is a null set.

For each standard Levi subgroup M of G equip  $\mathcal{A}_{M}(\mathbb{Q}_{v})$  with the product measure

(5.1.3.3) 
$$\prod_{i=1}^{r} |D^{\mathbf{G}_{i}}(X_{i,v})|_{v}^{-1/2} \mathrm{d}X_{i,v}$$

where  $G_1, G_2, \ldots, G_r$  are related to M as in (5.1.2.2).

Denote by  $\mathcal{A}_{M,reg}(\mathbb{Q}_v)_{ell}$  the subset of  $\mathcal{A}_{M,reg}(\mathbb{Q}_v)$  consisting of the images of the  $\mathbb{Q}_v$ -elliptic elements in  $\mathfrak{m}(\mathbb{Q}_v)$ . Then

(5.1.3.4) 
$$\mathcal{A}_{\mathrm{G,reg}}(\mathbb{Q}_v) = \prod_{\mathrm{M}\in\mathcal{L}} \pi_{\mathrm{M}}(\mathcal{A}_{\mathrm{M,reg}}(\mathbb{Q}_v)_{\mathrm{ell}}).$$

This decomposition is compatible with the measures on  $\mathcal{A}_{G}(\mathbb{Q}_{v})$  and  $\mathcal{A}_{M}(\mathbb{Q}_{v})$ .

The local measures induce S-local and global measures on  $\mathcal{A}_{G}(\mathbb{Q}_{S})$  and  $\mathcal{A}_{G}(\mathbb{A})$ .

(5.1.4) **Definition** Let S be a finite set of places of  $\mathbb{Q}$ , let M be a standard Levi subgroup of G, let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ . Define the *invariant weighted orbital integral*  $I_{\mathrm{M}}^{\mathrm{G}}(X, )$ to be the distribution on  $\mathfrak{g}(\mathbb{Q}_S)$  such that

(5.1.4.1) 
$$\forall f_S \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$$

$$\begin{split} I_{\mathrm{M}}^{\mathrm{G}}(X, f_{S}) &= J_{\mathrm{M}}^{\mathrm{G}}(X, f_{S}) - \sum_{\substack{\mathrm{L} \in \mathcal{L}(\mathrm{M}) \\ \mathrm{L} \neq \mathrm{G}}} \left( \sum_{\substack{\mathrm{L}'_{S} \in \mathcal{L}^{\mathrm{L}_{S}}}} |\mathrm{W}_{S,0}^{\mathrm{L}'_{S}}| |\mathrm{W}_{S,0}^{\mathrm{L}_{S}}|^{-1} \times \right. \\ & \times \sum_{\mathrm{T}_{S} \in \mathcal{T}_{\mathrm{ell}}(\mathrm{L}'_{S})} |\mathrm{W}(\mathrm{L}'_{S}, \mathrm{T}_{S})|^{-1} \times \\ & \times \int_{\mathfrak{t}_{S}(\mathbb{Q}_{S})} J_{\mathrm{L}}^{\mathrm{G}}(Y, f_{S}^{\,\check{}}) \Big( I_{\mathrm{M}}^{\mathrm{L}}(X, \,\,)^{\,\check{}}(Y) \Big) \, |D^{\mathrm{L}}(Y)|_{S}^{1/2} \mathrm{d}Y \end{split}$$

where  $I_{\mathrm{M}}^{\mathrm{M}}(X, \cdot)$  denotes the standard orbital integral on  $\mathfrak{m}(\mathbb{Q}_{S})$  defined by (1.1.6.9).

(5.1.5) **Remark** The invariant weighted orbital integral  $I_{\mathrm{M}}^{\mathrm{G}}(X, \cdot)$  is well-defined.

- The distribution  $I_{\mathrm{M}}^{\mathrm{G}}(X, \cdot)$  does not depend on the choice of the Fourier transforms on  $\mathfrak{g}(\mathbb{Q}_S)$  and its Levi subalgebras  $\mathfrak{l}(\mathbb{Q}_S)$  as long as these are compatible. See [Wa95] Lemme VI.5 for the *p*-adic case, the same argument also works for the real and *S*-local cases.
- The integral in (5.1.4.1) converges since the distribution  $I_{\mathrm{M}}^{\mathrm{L}}(X, )$  is represented by a smooth function supported on  $\mathfrak{g}_{\mathrm{reg.ss}}(\mathbb{Q}_S)$  such that the function

(5.1.5.1) 
$$Y \mapsto (I_{\mathrm{M}}^{\mathrm{L}}(X, )^{\hat{}}(Y)) |D^{\mathrm{L}}(Y)|_{S}^{1/2}$$

is locally

(5.1.5.2) 
$$O\left(\max\left\{1, -\log(|D^{G}(Y)|_{S})^{N}\right\}\right)$$

on  $\mathfrak{t}(\mathbb{Q}_S)$  for some natural number N and tempered at infinity, and the function

$$(5.1.5.3) Y \mapsto J_{\mathrm{L}}^{\mathrm{G}}(Y, f_{S})$$

is locally

(5.1.5.4) 
$$O\bigg(\max\{1, -\log(|D^{G}(Y)|_{S})^{M}\}\bigg)$$

on  $\mathfrak{t}(\mathbb{Q}_S)$  for some natural number M and rapidly decreasing at infinity. For the p-adic case see Lemme VI.3(iv) and Corollaire III.6 of [Wa95]. For the real case see Proposition 9 on page 108 of [Va77] and Corollary 7.4 of [Ar76].

(5.1.6) **Proposition** Let M be a standard Levi subgroup of G, let X be an element of  $\mathfrak{m}(\mathbb{Q})$ . Then  $I^{\mathrm{G}}_{\mathrm{M}}(X, )$  is an invariant distribution on  $\mathfrak{g}(\mathbb{Q}_{S})$ , i.e.

(5.1.6.1) 
$$\forall x \in \mathcal{G}(\mathbb{Q}_S) \ \forall f_S \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S)) \\ I^{\mathcal{G}}_{\mathcal{M}}(X, f_S \circ \mathrm{ad}(x)) = I^{\mathcal{G}}_{\mathcal{M}}(X, f_S)$$

*Proof.* For the *p*-adic case see Proposition VI.1 of [Wa95], the same argument also works for the *S*-local case.  $\Box$ 

(5.1.7) **Remark** To quote Waldspurger from §Introduction of [Wa95]:

- " On dispose de deux ensembles de distributions invariantes sur G:
  - (1) les intégrales orbitales associées aux éléments semi-simples de G;
  - (2) les caractères de représentations tempérées irréductibles de G.

 $(\ldots)$ 

Remplaçons G par son algèbre de Lie  $\mathfrak{g}$  et considérons l'espace de distributions invariantes par l'action adjointe de G. L'ensemble (1) a un analogue évident: les intégrales orbitales associées aux éléments semi-simples de  $\mathfrak{g}$ . Le seul but de cet article est de fournir un support un peu consistant à l'idée, d'ailleurs banale, que l'analogue de (2) est l'ensemble des transformées de Fourier des intégrales orbitales précédents (invariantes). "

The duality between the orbital integrals and the Fourier transforms of the invariant weighted orbital integrals is embodied in the local trace formulae on the Lie algebra  $\mathfrak{g}$ .

(5.1.8) **Proposition** (Local invariant trace formula of Waldspurger) Let v be a place of  $\mathbb{Q}$ , let  $f_v$  and  $g_v$  be Schwartz functions on  $\mathfrak{g}_v(\mathbb{Q}_v)$ , then

(5.1.8.1) 
$$\sum_{M_{v} \in \mathcal{L}^{G_{v}}} |W_{v,0}^{M_{v}}| |W_{v,0}^{G_{v}}|^{-1} \sum_{T_{v} \in \mathcal{T}_{ell}(M_{v})} |W(M_{v}, T_{v})|^{-1} \times \int_{\mathfrak{t}_{v}(\mathbb{Q}_{v})} (-1)^{\dim(A_{M_{v}}/A_{G_{v}})} I_{M_{v}}^{G_{v}}(X_{v}, f_{v}^{*}) I_{G_{v}}^{G_{v}}(X_{v}, g_{v}) dX_{v}$$
$$= \sum_{M_{v} \in \mathcal{L}^{G_{v}}} |W_{v,0}^{M_{v}}| |W_{v,0}^{G_{v}}|^{-1} \sum_{T_{v} \in \mathcal{T}_{ell}(M_{v})} |W(M_{v}, T_{v})|^{-1} \times \int_{\mathfrak{t}_{v}(\mathbb{Q}_{v})} (-1)^{\dim(A_{M_{v}}/A_{G_{v}})} I_{M_{v}}^{G_{v}}(X_{v}, g_{v}^{*}) I_{G_{v}}^{G_{v}}(X_{v}, f_{v}) dX_{v}.$$

*Proof.* For the *p*-adic case see Théorème VII.1 of [Wa95]. The argument only uses combinatorial properties of  $(G_v, M_v)$ -families and standard results from harmonic analysis on a  $\mathbb{Q}_v$ -vector space, hence works equally well in the real case.

(5.1.9) **Definition** Let S be a finite set of places of  $\mathbb{Q}$ , let M be a Levi subgroup in  $\mathcal{L}$ , let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ . Define the vector-valued orbital integral  $\mathcal{I}_{\mathrm{M}}^{\mathrm{G}}(X, \cdot)$  to be the vector-valued distribution on  $\mathfrak{g}(\mathbb{Q}_S)$  taking values in  $\bigoplus_{\mathbb{Q}\in\mathcal{F}(\mathrm{M})}\mathbb{C}$  such that

 $(5.1.9.1) \qquad \forall f_S \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$ 

$$\mathcal{I}_{\mathcal{M}}^{\mathcal{G}}(X, f_{S}) = \left( (-1)^{\dim(\mathcal{A}_{\mathcal{M}}/\mathcal{A}_{\mathcal{M}_{\mathcal{Q}}})} I_{\mathcal{M}}^{\mathcal{M}_{\mathcal{Q}}}(X, f_{S,\mathcal{Q}}) \right)_{\mathcal{Q}\in\mathcal{F}(\mathcal{M}).}$$

(5.1.10) Lemma (Induction and splitting)

Let S be a finite set of places of  $\mathbb{Q}$ , let M be a Levi subgroup in  $\mathcal{L}$ , let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ , let  $f_S$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$ .

- If L is a Levi subgroup in  $\mathcal{L}(M)$ , then  $\mathcal{I}_{L}^{G}(X, f_{S})$  is completely determined by  $\mathcal{I}_{M}^{G}(X, f_{S})$ .
- If S is the set  $\{v_1, v_2, \ldots, v_r\}$  for some natural number r and

(5.1.10.1) 
$$\forall i = 1, 2, \dots, r \; \exists f_{v_i} \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_{v_i}))$$
$$f_S = f_{v_1} \otimes f_{v_2} \otimes \cdots \otimes f_{v_r},$$

then  $\mathcal{I}_{\mathrm{M}}^{\mathrm{G}}(X, f_{S})$  is completely determined by  $\mathcal{I}_{\mathrm{M}}^{\mathrm{G}}(X_{v_{i}}, f_{v_{i}})$  where *i* ranges among  $1, 2, \ldots, r$ .

*Proof.* This follows from analogues of the descent and splitting identities (4.1.12.2) and (4.1.12.3) for the invariant weighted orbital integrals  $I_{\rm M}^{\rm G}(X, \cdot)$ , which could be deduced by the same arguments as in §7 of [Ar88b].

#### (5.1.11) Lemma (Local trace formula)

Let v be a place of  $\mathbb{Q}$ , let  $f_v$  and  $g_v$  be Schwartz functions on  $\mathfrak{g}(\mathbb{Q}_v)$ , then

$$(5.1.11.1) \qquad \sum_{\mathbf{M}\in\mathcal{L}} |\mathbf{W}_{0}^{\mathbf{M}}| |\mathbf{W}_{0}^{\mathbf{G}}|^{-1} \sum_{\mathbf{T}_{v}\in\mathcal{T}_{\mathrm{ell}}(\mathbf{M})} |\mathbf{W}(\mathbf{M},\mathbf{T}_{v})|^{-1} \times \\ \times \int_{\mathbf{t}_{v}(\mathbb{Q}_{v})} \mathcal{I}_{\mathbf{M}}^{\mathbf{G}}(X_{v},f_{v}^{*}) I_{\mathbf{G}}^{\mathbf{G}}(X_{v},g_{v}) \, \mathrm{d}X_{v} \\ = \sum_{\mathbf{M}\in\mathcal{L}} |\mathbf{W}_{0}^{\mathbf{M}}| |\mathbf{W}_{0}^{\mathbf{G}}|^{-1} \sum_{\mathbf{T}_{v}\in\mathcal{T}_{\mathrm{ell}}(\mathbf{M})} |\mathbf{W}(\mathbf{M},\mathbf{T}_{v})|^{-1} \times \\ \times \int_{\mathbf{t}_{v}(\mathbb{Q}_{v})} \mathcal{I}_{\mathbf{M}}^{\mathbf{G}}(X_{v},g_{v}^{*}) I_{\mathbf{G}}^{\mathbf{G}}(X_{v},f_{v}) \, \mathrm{d}X_{v}.$$

*Proof.* This follows from (5.1.8.1).

#### (5.1.12) Lemma (Global trace formula)

Let f be a Schwartz function on  $\mathfrak{g}(\mathbb{A})$  which is cuspidal at two distinct places of  $\mathbb{Q}$ , then

(5.1.12.1) 
$$\lim_{S} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} \sum_{X \in (\mathfrak{g}(\mathbb{Q}) \cap \mathfrak{o})_{\mathrm{G},S}} a^{\mathrm{G}}(S,X) I_{\mathrm{G}}^{\mathrm{G}}(X,f_{S})$$
$$= \lim_{S} \sum_{\mathfrak{o} \in \mathfrak{g}(\mathbb{Q})/\sim} \sum_{X \in (\mathfrak{g}(\mathbb{Q}) \cap \mathfrak{o})_{\mathrm{G},S}} a^{\mathrm{G}}(S,X) \mathcal{I}_{\mathrm{G}}^{\mathrm{G}}(X,f_{S})$$

where the limit is taken over finite sets S of places of  $\mathbb{Q}$ , and the coefficients  $a^{\mathrm{M}}(S, X)$  are defined as in (3.3.11.13).

*Proof.* This follows from (4.2.4.1).

#### 5.2 The local and global Schwartz spaces

(5.2.1) Definition Let v be a place of  $\mathbb{Q}$ . Let  $X_v$  be an element of  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$ . Define the maximal orbital integral  $\mathcal{I}_{max}^G(X_v, \cdot)$  to be the vector-valued distribution on  $\mathfrak{g}(\mathbb{Q}_v)$  such that

(5.2.1.1) 
$$\forall f_v \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_v)) \qquad \mathcal{I}^{\mathrm{G}}_{\max}(X_v, f_v) = \mathcal{I}^{\mathrm{G}}_{\mathrm{M}[\widetilde{X}_v - \mathrm{ell}]}(\widetilde{X}_v, f_v)$$

where  $\widetilde{X}_v$  is a regular semisimple element in  $\mathfrak{g}(\mathbb{Q}_v)$  lifting  $X_v$ , and  $M[\widetilde{X}_v\text{-ell}]$  is a standard Levi subgroup of G such that  $\mathfrak{m}[\widetilde{X}_v\text{-ell}](\mathbb{Q}_v)$  contains  $\widetilde{X}_v$  as a  $\mathbb{Q}_v$ -elliptic element.

(5.2.2) **Definition** Let v be a place of  $\mathbb{Q}$ . Define the *local Schwartz space*  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$  to be the space of complex-valued functions on  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$  such that

(5.2.2.1) 
$$\varphi_v \in \mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$$
$$\Leftrightarrow \exists f_v \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_v)) \; \forall X_v \in \mathcal{A}_{G, reg}(\mathbb{Q}_v)$$

$$\varphi_v(X_v) = I_{\mathcal{G}}^{\mathcal{G}}(X_v, f_v).$$

Define the *local Schwartz space*  $S_1(\mathcal{A}_G(\mathbb{Q}_v))$  to be the space of vector-valued functions on  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$  such that

(5.2.2.2)  

$$\varphi_{v} \in \mathcal{S}_{1}(\mathcal{A}_{G}(\mathbb{Q}_{v}))$$

$$\Leftrightarrow \exists f_{v} \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_{v})) \; \forall X_{v} \in \mathcal{A}_{G, reg}(\mathbb{Q}_{v})$$

$$\varphi_{v}(X_{v}) = \mathcal{I}_{max}^{G}(X_{v}, f_{v}).$$

(5.2.3) **Definition** Let v be a place of  $\mathbb{Q}$ . If v is p-adic, denote by  $\Lambda_0$  the standard lattice

(5.2.3.1) 
$$\Lambda_0 = \operatorname{gl}(n, \mathbb{Z})(\mathbb{Z}_p) \subset \operatorname{gl}(n, \mathbb{Q})(\mathbb{Q}_p)$$

and denote by  $\mathbb{I}_{\Lambda_0}$  its characteristic function. If v is archimedean, denote by  $\mathcal{E}$  the Gaussian function on  $\mathfrak{g}(\mathbb{R})$  which is self-dual with respect to the unitary Fourier transform, i.e.

$$\forall M \in \mathfrak{g}(\mathbb{R}) \qquad \mathcal{E}(M) = e^{-\pi \operatorname{Tr}(M^{\mathrm{T}}M)}$$

where  $M^{\mathrm{T}}$  denotes the transpose of M and Tr denotes the trace of an  $n \times n$  matrix.

Define the basic function  $\phi_{0,v}$  to be the element of  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$  such that

(5.2.3.2) 
$$\forall X_v \in \mathcal{A}_{G,reg}(\mathbb{Q}_v)$$

$$\phi_{0,v}(X_v) = \begin{cases} I_G^G(X_v, \mathbb{I}_{\Lambda_0}) & \text{if } v \text{ is } p\text{-adic,} \\ I_G^G(X_v, \mathcal{E}) & \text{if } v \text{ is archimedean.} \end{cases}$$

Define the basic function  $\phi_{1,v}$  to be the element of  $\mathcal{S}_1(\mathcal{A}_G(\mathbb{Q}_v))$  such that

(5.2.3.3) 
$$\forall X_v \in \mathcal{A}_{G,reg}(\mathbb{Q}_v)$$

$$\phi_{1,v}(X_v) = \begin{cases} \mathcal{I}_{\max}^G(X_v, \mathbb{I}_{\Lambda_0}) & \text{if } v \text{ is } p\text{-adic,} \\ \\ \mathcal{I}_{\max}^G(X_v, \mathcal{E}) & \text{if } v \text{ is archimedean.} \end{cases}$$

(5.2.4) Lemma If v is a p-adic place, then a complex-valued function  $\varphi_v$  on  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$  belongs to  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$  if and only if the following conditions are satisfied:

- $\varphi_v$  is a locally constant function on  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$ ;
- after extending by zero to  $\mathcal{A}_{G}(\mathbb{Q}_{v}), \varphi_{v}$  is compactly supported;
- for each singular point  $Z_v$  in the complement of  $\mathcal{A}_{G,reg}(\mathbb{Q}_S)$  in  $\mathcal{A}_G(\mathbb{Q}_v)$ , there exists an open neighborhood  $U_{\varphi_v, Z_v}$  of  $Z_v$  in  $\mathcal{A}_G(\mathbb{Q}_v)$  such that

(5.2.4.1) 
$$\varphi_{v}()|_{U_{\varphi_{v},Z_{v}}} \in \operatorname{span} \begin{cases} \Gamma_{G}^{G}(,\nu)|_{U_{\varphi_{v},Z_{v}}} : & \nu \subset \mathfrak{g}_{\widetilde{Z}_{v}}(\mathbb{Q}_{v}) \\ \text{nilpotent orbit} \end{cases}$$

where  $\varphi|_U$  denotes the restriction of the function  $\varphi$  to the open subset U, the point  $\widetilde{Z}_v$  is a regular semisimple element in  $\mathfrak{g}(\mathbb{Q}_v)$  lifting  $Z_v$ , and  $\Gamma_{\mathrm{G}}^{\mathrm{G}}(,\nu)$  denotes the Shalika germ at the nilpotent orbit  $\nu$ , see §17 of [Ko05].

*Proof.* Away from the singular locus the function  $\varphi_v$  is a linear combination of the characteristic functions of small compact open sets, and these all have lifts in  $\mathcal{S}(\mathfrak{g}(\mathbb{Q}_v))$ .

Near the singular locus the theory of Shalika germs implies that the asymptotic behavior of  $\varphi_v$  near the singular point  $Z_v$  is necessary. It remains to show that all the Shalika germs appear in  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$ , in other words there is no nontrivial linear relation among the possible asymptotes of  $I_G^G(X_v, f_v)$  as  $f_v$  ranges over  $\mathcal{S}(\mathfrak{g}(\mathbb{Q}_v))$ . The Shalika germ associated to the nilpotent orbit  $\nu$  is homogeneous of degree equal to the codimension of  $\nu$  in the nilpotent locus, hence the only possible linear relations are among the Shalika germs associated to the nilpotent orbits of the same dimension. But such nilpotent orbits are separated, so there are Schwarts functions  $f_v$  on  $\mathfrak{g}(\mathbb{Q}_v)$  that vanish on all but one of the nilpotent orbits of a given dimension, hence these too are linearly independent.

(5.2.5) Remark Let v be a p-adic place. It is conjectured by Jacquet for a general reductive Lie algebra and proven by Waldspurger for gl(n) that a Schwartz function  $f_v$  on  $\mathfrak{g}(\mathbb{Q}_v)$  has the property that

(5.2.5.1) 
$$\exists \lambda \in \mathbb{C} \ \forall X_v \in \mathcal{A}_{G,reg}(\mathbb{Q}_v)$$
$$I_G^G(X_v, f_v) = \lambda \cdot \phi_{0,v}(X_v)$$

if and only if  $I_{G}^{G}(X_{v}, f_{v})$  and  $I_{G}^{G}(X_{v}, f_{v})$  are both supported on the subset of  $\mathcal{A}_{G}(\mathbb{Q}_{v})$  consisting of the characteristic polynomials with coefficients in  $\mathbb{Z}_{v}$ .

(5.2.6) Definition Define the global Schwartz spaces  $S_0(\mathcal{A}_G(\mathbb{A}))$  and  $S_1(\mathcal{A}_G(\mathbb{A}))$  to be the tensor products of local Schwartz spaces as defined in (5.2.2.1) and (5.2.2.2)

(5.2.6.1) 
$$\mathcal{S}_{0}(\mathcal{A}_{G}(\mathbb{A})) = \bigotimes_{v}^{\text{res}} \mathcal{S}_{0}(\mathcal{A}_{G}(\mathbb{Q}_{v}))$$
$$\mathcal{S}_{1}(\mathcal{A}_{G}(\mathbb{A})) = \bigotimes_{v}^{\text{res}} \mathcal{S}_{1}(\mathcal{A}_{G}(\mathbb{Q}_{v}))$$

restricted with respect to the basic functions  $\phi_{0,v}$  and  $\phi_{1,v}$  as defined in (5.2.3.2) and (5.2.3.3).

#### 5.3 The Harish-Chandra transform

(5.3.1) **Definition** Let v be a place of  $\mathbb{Q}$ . For an element  $\varphi_v$  in  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$ , choose a Schwartz function  $f_v$  on  $\mathfrak{g}(\mathbb{Q}_v)$  such that

(5.3.1.1) 
$$\forall X_v \in \mathcal{A}_{G,reg}(\mathbb{Q}_v) \qquad \varphi_v(X_v) = I_G^G(X_v, f_v).$$

Let  $f_v$  be the Fourier transform of  $f_v$ , denote by  $\mathcal{H}_v(\varphi_v)$  the element in  $\mathcal{S}_1(\mathcal{A}_G(\mathbb{Q}_v))$  such that

(5.3.1.2) 
$$\forall X_v \in \mathcal{A}_{G,reg}(\mathbb{Q}_v) \qquad \mathcal{H}_v(\varphi_v)(X_v) = \mathcal{I}_{\max}^G(X_v, f_v).$$

Define the local Harish-Chandra transform to be the linear operator

$$(5.3.1.3) \qquad \qquad \mathcal{H}_v: \quad \mathcal{S}_0(\mathcal{A}_{\mathrm{G}}(\mathbb{Q}_v)) \quad \longrightarrow \quad \mathcal{S}_1(\mathcal{A}_{\mathrm{G}}(\mathbb{Q}_v))$$

defined by

(5.3.1.4) 
$$\mathcal{H}_v: \quad \varphi_v \quad \mapsto \quad \mathcal{H}_v(\varphi_v).$$

(5.3.2) Definition Let v be a place of  $\mathbb{Q}$ . If  $X_v$  is an element of  $\mathcal{A}_{G,reg}(\mathbb{Q})$ , then the orbital integral  $\mathcal{I}_{max}^G(X_v, \)$  is a vector-valued tempered distribution on  $\mathfrak{g}(\mathbb{Q}_v)$ . Denote by  $\mathcal{I}_{max}^G(X_v, \)^{\hat{}}$  its componentwise Fourier transform, which is represented by a conjugation invariant function on  $\mathfrak{g}(\mathbb{Q}_v)$ , hence descends to a function on  $\mathcal{A}_G(\mathbb{Q}_v)$  denoted by

(5.3.2.1) 
$$Y_v \mapsto \left(\mathcal{I}^{\mathrm{G}}_{\mathrm{max}}(X_v, )^{\hat{}}\right)(Y_v).$$

Define the local Harish-Chandra kernel function  $\mathcal{K}_v(, )$  to be the vector-valued bivariate function on  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$  such that

(5.3.2.2) 
$$\forall X_v, Y_v \in \mathcal{A}_{G, reg}(\mathbb{Q}_v)$$
$$\mathcal{K}_v(X_v, Y_v) = |D(Y_v)|_v^{1/2} \Big( \mathcal{I}_{\max}^G(X_v, )^{\hat{}} \Big) (Y_v).$$

(5.3.3) **Remark** In the case when v is a p-adic place, each component of the vectorvalued kernel function  $\mathcal{K}_v$  is denoted by  $\hat{i}_{\mathrm{M}}^{\mathrm{G}}(X_v, Y_v)$  for some Levi subgroup M by Waldspurger in [Wa95].

(5.3.4) Lemma Let v be a place of  $\mathbb{Q}$ . The local Harish-Chanrda transform  $\mathcal{H}_v$  is an integral operator with integral kernel  $\mathcal{K}_v$ , i.e.

(5.3.4.1)  $\forall \varphi_v \in \mathcal{S}_0(\mathcal{A}_{\mathrm{G}}(\mathbb{Q}_v)) \ \forall X_v \in \mathcal{A}_{\mathrm{G,reg}}(\mathbb{Q}_v)$ 

$$\mathcal{H}_{v}(\varphi_{v})(X_{v}) = \int_{\mathcal{A}_{\mathrm{G}}(\mathbb{Q}_{v})} \mathcal{K}_{v}(X_{v}, Y_{v})\varphi_{v}(Y_{v}) |D(Y_{v})|_{v}^{1/2} \mathrm{d}Y_{v}$$

In particular the operator  $\mathcal{H}_v$  is well-defined.

*Proof.* Every  $X_v$  in  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$  has a  $\delta$ -sequence in  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$ . More precisely there exists a sequence of functions  $\delta_{X_v,1}, \delta_{X_v,2}, \delta_{X_v,3}, \ldots$  in  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$  such that

(5.3.4.2) 
$$\lim_{i \to \infty} \delta_{X_v,i} = \delta_{X_v}$$

as distributions on  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$ , where  $\delta_{X_v}$  denotes the Dirac distribution at  $X_v$ . For each natural number *i* choose a Schwartz function  $\widetilde{\delta}_{X_v,i}$  on  $\mathfrak{g}(\mathbb{Q}_v)$  such that

(5.3.4.3) 
$$\forall Y_v \in \mathcal{A}_{G,reg}(\mathbb{Q}_v) \qquad \delta_{X_v,i}(Y_v) = I_G^G(Y_v, \widetilde{\delta}_{X_v,i}).$$

Let  $f_v$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_v)$  such that

(5.3.4.4) 
$$\forall Y_v \in \mathcal{A}_{G,reg}(\mathbb{Q}_v) \qquad \varphi_v(Y_v) = I_G^G(Y_v, f_v).$$

By the local trace formula (5.1.11.1)

(5.3.4.5) 
$$\int_{\mathcal{A}_{\mathrm{G}}(\mathbb{Q}_{v})} \mathcal{I}_{\mathrm{max}}^{\mathrm{G}}(Y_{v}, f_{v}^{\hat{}}) I_{\mathrm{G}}^{\mathrm{G}}(Y_{v}, \widetilde{\delta}_{X_{v},i}) |D(Y_{v})|_{v}^{-1/2} \mathrm{d}Y_{v}$$
$$= \int_{\mathcal{A}_{\mathrm{G}}(\mathbb{Q}_{v})} \mathcal{I}_{\mathrm{max}}^{\mathrm{G}}(Y_{v}, \widetilde{\delta}_{X_{v},i}^{\hat{}}) I_{\mathrm{G}}^{\mathrm{G}}(Y_{v}, f_{v}) |D(Y_{v})|_{v}^{-1/2} \mathrm{d}Y_{v},$$

hence

(5.3.4.6) 
$$\mathcal{H}_v(\varphi_v)(X_v)$$

$$= \int_{\mathcal{A}_{G}(\mathbb{Q}_{v})} \mathcal{H}_{v}(\varphi_{v})(Y_{v})\delta_{X_{v}}(Y_{v}) |D(Y_{v})|_{v}^{-1/2} dY_{v}$$

$$= \lim_{i \to \infty} \int_{\mathcal{A}_{G}(\mathbb{Q}_{v})} \mathcal{H}_{v}(\varphi_{v})(Y_{v})\delta_{X_{v},i}(Y_{v}) |D(Y_{v})|_{v}^{-1/2} dY_{v}$$

$$= \lim_{i \to \infty} \int_{\mathcal{A}_{G}(\mathbb{Q}_{v})} \mathcal{I}_{\max}^{G}(Y_{v}, \widetilde{\delta}_{X_{v},i}) \varphi_{v}(Y_{v}) |D(Y_{v})|_{v}^{-1/2} dY_{v}.$$

Hence  $\mathcal{H}_v$  is an integral operator with integral kernel

(5.3.4.7) 
$$\lim_{i \to \infty} \mathcal{I}_{\max}^{\mathrm{G}}(Y_v, \widetilde{\delta}_{X_v, i}) = |D(Y_v)|_v^{1/2} \left( \mathcal{I}_{\max}^{\mathrm{G}}(X_v, )^{\widehat{}} \right) (Y_v)$$

which is independent of the choice of the sequence  $(\widetilde{\delta}_{X_v,i})_{i=1}^{\infty}$  and equal to the local Harish-Chandra kernel  $\mathcal{K}_v(X_v, Y_v)$ .

(5.3.5) **Remark** The local Harish-Chandra transform preserves the basic functions, i.e. (5.3.5.1)  $\mathcal{H}_v: \phi_{0,v} \mapsto \phi_{1,v}.$ 

(5.3.6) Definition For a finite set S of places of  $\mathbb{Q}$ , define the S-local Harish-Chandra transform  $\mathcal{H}_S$  to be the tensor product

(5.3.6.1) 
$$\mathcal{H}_S = \bigotimes_{v \in S} \mathcal{H}_v : \bigotimes_{v \in S} \mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v)) \longrightarrow \bigotimes_{v \in S} \mathcal{S}_1(\mathcal{A}_G(\mathbb{Q}_v)).$$

By (5.3.5.1) the limit of  $\mathcal{H}_S$  as S approaches infinity defines a linear operator

(5.3.6.2) 
$$\lim_{S} \mathcal{H}_{S} : \bigotimes_{v}^{\operatorname{res}} \mathcal{S}_{0}(\mathcal{A}_{G}(\mathbb{Q}_{v})) \longrightarrow \bigotimes_{v}^{\operatorname{res}} \mathcal{S}_{1}(\mathcal{A}_{G}(\mathbb{Q}_{v})),$$

which is defined to be the global Harish-Chandra transform

$$(5.3.6.3) \qquad \qquad \mathcal{H}: \quad \mathcal{S}_0(\mathcal{A}_G(\mathbb{A})) \quad \longrightarrow \quad \mathcal{S}_1(\mathcal{A}_G(\mathbb{A})).$$

#### 5.4 The Poisson summation formula

(5.4.1) **Definition** Let Z be a singular point in the complement of  $\mathcal{A}_{G,reg}(\mathbb{Q})$  in  $\mathcal{A}_G(\mathbb{Q})$ . Denote by  $\mathfrak{o}_Z$  the  $\sim$  equivalence class in  $\mathfrak{g}(\mathbb{Q})$  which is the fiber of Z. Let S be a finite set of places of  $\mathbb{Q}$ , let  $\varphi_S$  be an element of  $\bigotimes_{v \in S} \mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$ , let  $f_S$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$  such that

(5.4.1.1) 
$$\forall X \in \mathcal{A}_{G,reg}(\mathbb{Q}_S) \qquad \varphi_S(X) = I_G^G(X, f_S)$$

By (3.1.5.4), if Z' is a  $\equiv$  equivalence class in  $(\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o}_Z)_{M,S}$ , then  $I_G^G(Z', f_S)$  is determined by  $\varphi_S(X)$  where X ranges over the elements of  $\mathcal{A}_{G,reg}(\mathbb{Q}_S)$  close to Z'. Denote

(5.4.1.2) 
$$\varphi_S(Z') = I_G^G(Z', f_S).$$

(5.4.2) **Definition** Let X be an element of  $\mathcal{A}_{G,reg}(\mathbb{Q})$ , let  $\widetilde{X}$  be a regular semisimple element of  $\mathfrak{g}(\mathbb{Q})$  lifting X, then by (3.3.12.1) the constant  $a^G(S, \widetilde{X})$  as defined in (3.3.11.13) is independent of the finite set of places S and determined by X. Denote

(5.4.2.1) 
$$a(X) = a^{G}(S, X).$$

(5.4.3) **Definition** Let v be a place of  $\mathbb{Q}$ , let  $\varphi_v$  be an element in  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$ , then  $\varphi_v$  is said to be *cuspidal* if

$$(5.4.3.1) \qquad \qquad \varphi_v(X_v) = 0$$

whenever  $X_v$  is the image of a point  $\widetilde{X}_v$  in  $\mathfrak{g}(\mathbb{Q}_v)$  which is regular semisimple and not  $\mathbb{Q}_v$ -elliptic.

#### (5.4.4) **Proposition** (Poisson summation formula)

Let  $\varphi$  be an element of  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{A}))$  which is cuspidal at two distinct places of  $\mathbb{Q}$ , then

$$(5.4.4.1) \qquad \sum_{X \in \mathcal{A}_{G,reg}(\mathbb{Q})} a(X)\varphi(X) + \\ + \sum_{Z \in \mathcal{A}_G(\mathbb{Q}) - \mathcal{A}_{G,reg}(\mathbb{Q})} \lim_{S} \left( \sum_{Z' \in (\mathfrak{g}(\mathbb{Q}) \cap \mathfrak{o}_Z)_{G,S}} a^G(S, Z')\varphi_S(Z') \right) \\ = \sum_{X \in \mathcal{A}_{G,reg}(\mathbb{Q})} a(X)\mathcal{H}(\varphi)(X) + \\ + \sum_{Z \in \mathcal{A}_G(\mathbb{Q}) - \mathcal{A}_{G,reg}(\mathbb{Q})} \lim_{S} \left( \sum_{Z' \in (\mathfrak{g}(\mathbb{Q}) \cap \mathfrak{o}_Z)_{G,S}} a^G(S, Z')\mathcal{H}_S(\varphi_S)(Z') \right).$$

*Proof.* This follows from the global trace formula (5.1.12.1) and Remark (4.2.2).

(5.4.5) **Remark** The Poisson summation formula (5.4.4.1) has the general form

(5.4.5.1) 
$$\sum_{X \in \mathcal{A}_{G, reg}(\mathbb{Q})} a(X)\varphi(X) + \sum_{Z \in \mathcal{A}_{G}(\mathbb{Q}) - \mathcal{A}_{G, reg}(\mathbb{Q})} (\dots)$$
$$= \sum_{X \in \mathcal{A}_{G, reg}(\mathbb{Q})} a(X)\mathcal{H}(\varphi)(X) + \sum_{Z \in \mathcal{A}_{G}(\mathbb{Q}) - \mathcal{A}_{G, reg}(\mathbb{Q})} (\dots).$$

(5.4.6) Corollary Let v and w be two places of  $\mathbb{Q}$ . Let  $\varphi$  be an element of  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{A}))$ which is cuspidal at two distinct places of  $\mathbb{Q}$  such that  $\varphi_v$  vanishes on a neighborhood of the complement of  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$  in  $\mathcal{A}_G(\mathbb{Q}_v)$  and  $\mathcal{H}_w(\varphi_w)$  vanishes on a neighborhood of the complement of  $\mathcal{A}_{G,reg}(\mathbb{Q}_w)$  in  $\mathcal{A}_G(\mathbb{Q}_w)$ . Then

(5.4.6.1) 
$$\sum_{X \in \mathcal{A}_{G,reg}(\mathbb{Q})} a(X)\varphi(X) = \sum_{X \in \mathcal{A}_{G,reg}(\mathbb{Q})} a(X)\mathcal{H}(\varphi)(X).$$

*Proof.* This follows from (5.4.5.1).

#### (5.4.7) Corollary (Poisson summation formula for $\mathcal{H}_0$ )

There exists an endomorphism  $\mathcal{H}_0$  of  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{A}))$ , defined similarly as  $\mathcal{H}$  as a tensor product of local endomorphisms  $\mathcal{H}_{0,v}$ , such that for each Schwartz function  $\varphi$  on  $\mathcal{A}_G(\mathbb{A})$  which is cuspidal at two distinct places of  $\mathbb{Q}$ 

$$(5.4.7.1) \qquad \sum_{X \in \mathcal{A}_{G, reg}(\mathbb{Q})} a(X)\varphi(X) + \\ + \sum_{Z \in \mathcal{A}_{G}(\mathbb{Q}) - \mathcal{A}_{G, reg}(\mathbb{Q})} \lim_{S} \left( \sum_{Z' \in (\mathfrak{g}(\mathbb{Q}) \cap \mathfrak{o}_{Z})_{G, S}} a^{G}(S, Z')\varphi_{S}(Z') \right) \\ = \sum_{X \in \mathcal{A}_{G, reg}(\mathbb{Q})} a(X)\mathcal{H}_{0}(\varphi)(X) + \\ + \sum_{Z \in \mathcal{A}_{G}(\mathbb{Q}) - \mathcal{A}_{G, reg}(\mathbb{Q})} \lim_{S} \left( \sum_{Z' \in (\mathfrak{g}(\mathbb{Q}) \cap \mathfrak{o}_{Z})_{G, S}} a^{G}(S, Z')\mathcal{H}_{0, S}(\varphi_{S})(Z') \right)$$

Proof. The local endomorphisms  $\mathcal{H}_{0,v}$  are defined by sending the orbital integral  $I_{\mathrm{G}}^{\mathrm{G}}(X, f_v)$  to the orbital integral of the Fourier transform  $I_{\mathrm{G}}^{\mathrm{G}}(X, f_v)$  for a Schwartz function  $f_v$  on  $\mathfrak{g}(\mathbb{Q}_v)$ . If v is a p-adic place, the endomorphism  $\mathcal{H}_{0,v}$  is well-defined due to the density of regular semisimple orbital integrals. See §27 of [Ko05]. If v is archimedean, then  $\mathcal{H}_{0,v}$  is well-defined due to the second last identity on page 104 of [Va77]. The local basic function  $\phi_{0,v}$  is preserved by  $\mathcal{H}_{0,v}$ , hence the global endomorphism  $\mathcal{H}_0$  is also well-defined.

The Poisson summation formula (5.4.7.1)follows from (5.4.4.1) since the cuspidality condition on  $\varphi$  implies that only those X for which the orbital integrals  $I_{\mathrm{G}}^{\mathrm{G}}(X, f_{v})$  and  $\mathcal{I}_{\mathrm{max}}^{\mathrm{G}}(X, f_{v})$ are equal, where  $f_{v}$  denotes a local Schwartz function on  $\mathfrak{g}(\mathbb{Q}_{v})$  whose orbital integral is equal to  $\varphi_{v}$ , contribute to the identity. (5.4.8) **Remark** The endomorphism  $\mathcal{H}_0$  is the Harish-Chandra transform originally considered by Jacquet. If v is a p-adic place, then it is conjectured by Jacquet for a general reductive Lie algebra and proven by Waldspurger for gl(n) that a Schwartz function  $\varphi_v$  on  $\mathcal{A}_G(\mathbb{Q}_v)$  is proportional to  $\phi_{0,v}$  if and only if  $\varphi_v$  and  $\mathcal{H}_{0,v}(\varphi_v)$  are both supported on the subset of  $\mathcal{A}_G(\mathbb{Q}_v)$  consisting of the characteristic polynomials with coefficients in  $\mathbb{Z}_v$ .

(5.4.9) Corollary Let v be a place of  $\mathbb{Q}$ . The local Harish-Chandra transform  $\mathcal{H}_v$  is a bijection from  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$  onto  $\mathcal{S}_1(\mathcal{A}_G(\mathbb{Q}_v))$ .

*Proof.* The following argument is suggested by Sakellaridis.

The local Harish-Chandra transform  $\mathcal{H}_v$  is surjective by definition. For injectivity assume for contradiction that there exists an element  $\varphi_v$  in  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_v))$  and a point  $X_v$  in  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$ such that

(5.4.9.1) 
$$\forall Y_v \in \mathcal{A}_{G,reg}(\mathbb{Q}_v) \qquad \mathcal{H}_v(\varphi_v)(Y_v) = 0$$

and

(5.4.9.2) 
$$\varphi_v(X_v) \neq 0.$$

Since  $\varphi_v$  is smooth on  $\mathcal{A}_{G,reg}(\mathbb{Q}_v)$ , without loss of generality  $X_v$  lies in the dense subset  $\mathcal{A}_{G,reg}(\mathbb{Q})$ . By parabolic descent along a suitable parabolic subgroup, the argument is reduced to the special case that  $X_v$  is the image of a  $\mathbb{Q}_v$ -elliptic element of  $\mathfrak{g}(\mathbb{Q}_v)$ .

Let  $\varphi$  be en element of  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{A}))$  which is cuspidal at two other places and whose local component at v is equal to  $\varphi_v$ , then by the Poisson summation formula (5.4.4.1)

(5.4.9.3) 
$$\sum_{X \in \mathcal{A}_{G,reg}(\mathbb{Q})} a(X)\varphi(X) + \sum_{Z \in \mathcal{A}_G(\mathbb{Q}) - \mathcal{A}_{G,reg}(\mathbb{Q})} \begin{pmatrix} \text{contribution from} \\ \text{the singular locus} \end{pmatrix} = 0$$

since  $\mathcal{H}(\varphi)$  vanishes identically at the place v. There are two cases:

• If v is p-adic, then there exists an integer N such that  $X_v$  lies in the image of  $\mathfrak{g}_{\text{reg.ss}}(N^{-1}\mathbb{Z})$  in  $\mathcal{A}_{G,\text{reg}}(\mathbb{Q})$  under the natural projection.

At a finite place w distinct from v, denote by  $\Lambda_{N,w}$  the lattice

(5.4.9.4) 
$$\Lambda_{N,w} = \mathfrak{g}(N^{-1}\mathbb{Z}_w) \subset \mathfrak{g}(\mathbb{Q}_w).$$

Let  $\varphi_w$  be the element of  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_w))$  such that

(5.4.9.5) 
$$\forall Y_w \in \mathcal{A}_{G,reg}(\mathbb{Q}_w) \qquad \varphi_w(Y_w) = I_G^G(Y_w, \mathbb{I}_{\Lambda_{N,w}})$$

where  $\mathbb{I}_{\Lambda_{N,w}}$  denotes the characteristic function of  $\Lambda_{N,w}$ .

At infinity, the set of rational points where  $\bigotimes_{w<\infty} \varphi_w$  is nonzero is contained in the discrete subset

(5.4.9.6) 
$$\pi_{\infty}(\mathfrak{g}_{\mathrm{reg.ss}}(N^{-1}\mathbb{Z})) \subset \mathcal{A}_{\mathrm{G,reg}}(\mathbb{R})$$

where  $\pi_{\infty}$  denotes the natural projection from  $\mathfrak{g}(\mathbb{R})$  to  $\mathcal{A}_{G}(\mathbb{R})$ . Choose  $\varphi_{\infty}$  to be a bump function supported away from the complement of  $\mathcal{A}_{G,reg}(\mathbb{R})$  in  $\mathcal{A}_{G}(\mathbb{R})$  such that  $X_{v}$  is the only point contained in

(5.4.9.7) 
$$\operatorname{supp}(\varphi_{\infty}) \cap \pi_{\infty}(\mathfrak{g}_{\operatorname{reg.ss}}(N^{-1}\mathbb{Z})).$$

Choose  $\varphi$  to be  $\bigotimes_w \varphi_w$ , then the left hand side of (5.4.9.3) is equal to

(5.4.9.8) 
$$a(X_v)\varphi(X_v) = \operatorname{Vol}\left(\operatorname{T}_{X_v}(\mathbb{Q})\backslash \operatorname{T}_{X_v}(\mathbb{A})^1\right)\lim_{S}\left(\prod_{w\in S}\varphi_w(X_v)\right)$$

for some torus  $T_{X_v}$ , which is nonzero.

• If v is archimedean, then choose a regular semisimple element  $\widetilde{X}_{\infty}$  of  $\mathfrak{g}(\mathbb{Q})$  lifting  $X_{\infty}$ and choose a Schwartz function  $f_{\infty}$  on  $\mathfrak{g}(\mathbb{A})$  such that

(5.4.9.9) 
$$\forall Y_{\infty} \in \mathcal{A}_{G,reg}(\mathbb{R}) \qquad \varphi_{\infty}(Y_{\infty}) = I_{G}^{G}(Y_{\infty}, f_{\infty}).$$

If N is an integer, denote by  $\Lambda_{N,\infty}$  the lattice

(5.4.9.10) 
$$\Lambda_{N,\infty} = \mathfrak{g}(N\mathbb{Z}) \subset \mathfrak{g}(\mathbb{R}).$$

Since  $f_{\infty}$  is a Schwartz function on  $\mathfrak{g}(\mathbb{R})$ , the quantity  $\varphi_{\infty}(Y_{\infty})$  is rapidly decreasing as  $Y_{\infty}$  approaches infinity in  $\mathcal{A}_{G,reg}(\mathbb{R})$  in such a way that  $|D(Y_{\infty})|$  is uniformly bounded below, hence for every positive real numbers  $\epsilon$  and r there exists a natural number  $N_{\epsilon}$  such that

(5.4.9.11) 
$$\forall N \ge N_{\epsilon} \qquad \sum_{\substack{Y \in \pi_{\infty}(\tilde{X}_{\infty} + \Lambda_{N,\infty}) \\ Y \neq X_{\infty}, \ |D(Y)| \ge 1}} |\varphi_{\infty}(Y)| < \epsilon N^{-r}$$

where  $\pi_{\infty}$  denotes the natural projection from  $\mathfrak{g}(\mathbb{R})$  to  $\mathcal{A}_{G}(\mathbb{R})$ . Choose

(5.4.9.12) 
$$\epsilon = \frac{1}{2(n!)^{n \cdot n!}} |a(X_{\infty})\varphi_{\infty}(X_{\infty})|$$

where n is the rank of G and choose r to be  $2\dim(\mathfrak{g})$ . Let N be a natural number greater than  $N_{\epsilon}$  such that

(5.4.9.13) 
$$\forall Y \in \pi_{\infty}(\widetilde{X}_{\infty} + \Lambda_{N,\infty}) \quad D(Y) \neq 0.$$

At a finite place w, denote by  $\Lambda_{N,w}$  the lattice

(5.4.9.14) 
$$\Lambda_{N,w} = \mathfrak{g}(N\mathbb{Z}_w) \subset \mathfrak{g}(\mathbb{Q}_w)$$

Let  $\varphi_w$  be the element of  $\mathcal{S}_0(\mathcal{A}_G(\mathbb{Q}_w))$  such that

(5.4.9.15) 
$$\forall Y_w \in \mathcal{A}_{G,reg}(\mathbb{Q}_w) \qquad \varphi_w(Y_w) = I_G^G(Y_w, \mathbb{I}_{\widetilde{X}_\infty + \Lambda_{N,w}})$$

where  $\mathbb{I}_{\widetilde{X}_{\infty}+\Lambda_{N,w}}$  denotes the characteristic function of the translation of  $\Lambda_{N,w}$  by  $\widetilde{X}_{\infty}$ . Choose  $\varphi$  to be  $\bigotimes_{w} \varphi_{w}$ , then the left hand side of (5.4.9.3) is equal to

$$(5.4.9.16) \qquad a(X_{\infty}) \prod_{w < \infty} \varphi_w(X_{\infty}) \cdot \varphi_{\infty}(X_{\infty}) + + \sum_{\substack{Y \in \pi_{\infty}(\tilde{X}_{\infty} + \Lambda_{N,\infty}) \\ Y \neq X_{\infty}, \ |D(Y)| \ge 1}} a(Y) \prod_{w < \infty} \varphi_w(Y) \cdot \varphi_{\infty}(Y) = C \cdot \left( a(X_{\infty})\varphi_{\infty}(X_{\infty}) + \right) + \sum_{\substack{Y \in \pi_{\infty}(\tilde{X}_{\infty} + \Lambda_{N,\infty}) \\ Y \neq X_{\infty}, \ |D(Y)| \ge 1}} \left( a(Y) \frac{\prod_{w < \infty} \varphi_w(Y)}{\prod_{w < \infty} \varphi_w(X_{\infty})} \right) \varphi_{\infty}(Y) \right)$$

where C is a nonzero constant. Let Y be an element of  $\mathcal{A}_{G,reg}(\mathbb{Q})$ . By definition

(5.4.9.17) 
$$a(Y) = \pm |\pi_0(\mathcal{M}_Y)|^{-1} \operatorname{Vol}(\mathcal{T}_Y(\mathbb{Q}) \setminus \mathcal{T}_Y(\mathbb{A})^1)$$

where M is a Levi subgroup in  $\mathcal{L}$  and  $T_Y$  is a maximal torus in G. By the Main theorem in §5 of [On63],

(5.4.9.18) 
$$\operatorname{Vol}(\operatorname{T}_{Y}(\mathbb{Q})\backslash\operatorname{T}_{Y}(\mathbb{A})^{1}) \leq |H_{\operatorname{Gal}}^{1}(\mathbb{Q},\widehat{\operatorname{T}}_{Y})|$$

where  $H_{\text{Gal}}^1$  denotes the first Galois cohomology group and  $\widehat{T}_Y$  denotes the Galois module of algebraic characters of  $T_Y$ . Let F be the splitting field of  $T_Y$ , denote by  $\Gamma$ the Galois group of F over  $\mathbb{Q}$ . Then by the inflation-restriction exact sequence

$$(5.4.9.19) \qquad 0 \longrightarrow H^{1}(\Gamma, \widehat{T}_{Y}) \longrightarrow H^{1}_{\text{Gal}}(\mathbb{Q}, \widehat{T}_{Y}) \longrightarrow H^{1}_{\text{Gal}}(F, \widehat{T}_{Y})^{\Gamma}$$

$$\|$$

$$Hom(\text{Gal}(\overline{F}/F), \mathbb{Z}^{n})^{\Gamma}$$

and the compactness of  $\operatorname{Gal}(\overline{F}/F)$  which implies that the only continuous homomorphism from  $\operatorname{Gal}(\overline{F}/F)$  to  $\mathbb{Z}^n$  is the trivial homomorphism,

$$(5.4.9.20) |H^1(\Gamma, \widehat{T}_Y)| = |H^1_{\text{Gal}}(\mathbb{Q}, \widehat{T}_Y)|.$$

The group  $H^1(\Gamma, \widehat{\mathrm{T}}_Y)$  is annihilated by the order of  $\Gamma$  which is bounded by the factorial of n since the  $\widehat{\mathrm{T}}_Y$  splits over the splitting field of the characteristic polynomial represented by Y which is of degree n. By the bar resolution, the group  $H^1(\Gamma, \widehat{\mathrm{T}}_Y)$ is a subquotient of  $\bigoplus_{g\in\Gamma} \mathbb{Z}^n$  which is generated by at most  $n \cdot n!$  elements, hence by (5.4.9.17), (5.4.9.18) and (5.4.9.20)

(5.4.9.21) 
$$\forall Y \in \mathcal{A}_{G,reg}(\mathbb{Q}) \quad |a(Y)| \le (n!)^{n \cdot n!}.$$

As N approaches infinity, either  $\prod_{w < \infty} \varphi_w(Y)$  vanishes or

(5.4.9.22) 
$$N^{-\dim(\mathfrak{g})} \leq \prod_{w < \infty} \varphi_w(Y) \leq 1,$$

and  $\prod_{w<\infty} \varphi_w(X_\infty)$  is nonzero. Hence without loss of generality N is large enough so that

(5.4.9.23) 
$$\forall Y \in \mathcal{A}_{G,reg}(\mathbb{Q}) \qquad \frac{\prod_{w < \infty} \varphi_w(Y)}{\prod_{w < \infty} \varphi_w(X_\infty)} \le N^r.$$

By (5.4.9.21) and (5.4.9.23),

$$(5.4.9.24) \qquad \sum_{\substack{Y \in \pi_{\infty}(\tilde{X}_{\infty} + \Lambda_{N,\infty}) \\ Y \neq X_{\infty}, \ |D(Y)| \ge 1}} \left| a(Y) \frac{\prod_{w < \infty} \varphi_w(Y)}{\prod_{w < \infty} \varphi_w(X_{\infty})} \varphi_{\infty}(Y) \right| \\ \leq N^r (n!)^{n \cdot n!} \sum_{\substack{Y \in \pi_{\infty}(\tilde{X}_{\infty} + \Lambda_{N,\infty}) \\ Y \neq X_{\infty}, \ |D(Y)| \ge 1}} \left| \varphi_{\infty}(Y) \right| \\ (5.4.9.25) \qquad < N^r (n!)^{n \cdot n!} \epsilon N^{-r}$$

$$(5.4.9.26) = (n!)^{n \cdot n!} \frac{1}{2(n!)^{n \cdot n!}} |a(X_{\infty})\varphi_{\infty}(X_{\infty})|$$
$$= \frac{1}{2} |a(X_{\infty})\varphi_{\infty}(X_{\infty})|$$

where the inequality (5.4.9.25) follows from (5.4.9.11) and the equality (5.4.9.26) follows from (5.4.9.12). Hence the right hand side of (5.4.9.16) is nonzero.

In either case the left hand side of (5.4.9.3) is nonzero, which is a contradiction.

(5.4.10) Remark Injectivity of  $\mathcal{H}_v$  is analogous to the classical result of Harish-Chandra on density of regular semisimple orbital integrals which states that for a *p*-adic reductive Lie algebra  $\mathfrak{g}$ , if f is a Schwartz function on  $\mathfrak{g}$  such that all the regular semisimple orbital integrals of f vanish, then  $\mathcal{D}(f)$  vanishes for every invariant distribution  $\mathcal{D}$  on  $\mathfrak{g}$ . See §27 of [Ko05].

# A Appendix: Scissors congruence and orbital integrals

In this appendix weighted orbital integrals on  $\mathfrak{g}$  are interpreted as scissors congruence classes of polyhedra. Similar results also hold for invariant weighted orbital integrals.

(A.1) Definition Let  $\mathbb{E}^n$  denote the *n*-dimensional Euclidean space. A polytope in  $\mathbb{E}^n$  that is closed with nonempty interior is said to be *proper*. Let *P* and *Q* be proper convex polytopes in  $\mathbb{E}^n$ . Then *P* and *Q* are said to be *translational scissors congruent* if there exist convex polytopes  $P_1, P_2, \ldots, P_l$  and  $Q_1, Q_2, \ldots, Q_l$  in  $\mathbb{E}^n$  such that

(A.1.1) 
$$P = \bigcup_{i=1}^{l} P_i \qquad Q = \bigcup_{i=1}^{l} Q_i$$

and  $P_i$  is a translation of  $Q_i$  for each index *i* among 1, 2, ..., l. Define the scissors group of  $\mathbb{E}^n$ , denoted by  $\mathbb{S}(\mathbb{E}^n)$ , to be the quotient of the free abelian group generated by the proper convex polytopes in  $\mathbb{E}^n$  modulo translational scissors congruence.

(A.2) Definition A flag  $\Phi$  of linear subspaces in  $\mathbb{E}^n$  is said to be strict of length l if

(A.2.1) 
$$\Phi = V_0 \supset V_1 \supset \cdots \supset V_l$$

where  $V_i$  has codimension *i*. Let  $\Phi$  be a strict flag of length *l*. A rigging **r** of  $\Phi$  is a collection

(A.2.2) 
$$\mathbf{r} = \left\{ r_1, r_2, \dots, r_l \right\}$$

where  $r_i$  is a real linear functional on  $V_{i-1}$  with kernel  $V_i$  for each i among  $1, 2, \ldots, l$ . Two riggings  $\mathbf{r}$  and  $\mathbf{r}'$  are *equivalent* if  $r_i$  and  $r'_i$  are positive multiples of each other for each iamong  $1, 2, \ldots, l$ . Denote by  $\mathcal{R}ig(\Phi)$  the collection of equivalence classes of riggings of  $\Phi$ . A rigged flag  $\Phi^{\mathbf{r}}$  is defined to be a strict flag  $\Phi$  together with a choice of an element  $\mathbf{r}$  in  $\mathcal{R}ig(\Phi)$ .

An orientation of  $\mathbb{E}^n$  is an ordered basis of  $\mathbb{E}^n$  defined up to a linear transformation with positive determinant. The product of an orientation of  $\mathbb{E}^n$  with a translation invariant measure on  $\mathbb{E}^n$  is equal to a volume form on  $\mathbb{E}^n$ . Fix an ordered basis  $\mathcal{B}$ 

(A.2.3) 
$$\mathcal{B} = \left(b_1, b_2, \dots, b_n\right)$$

of  $\mathbb{E}^n$  such that

(A.2.4) 
$$\mathcal{B}' = \left(b_{l+1}, b_{l+2}, \dots, b_n\right)$$

is an ordered basis of  $V_l$ . Let **r** be a rigging of  $\Phi$ , choose vectors

(A.2.5) 
$$c_1 \in V_0, \quad c_2 \in V_1, \quad \dots, \quad c_l \in V_{l-1}$$

such that

(A.2.6) 
$$\forall i = 1, 2, \dots, l \quad r_i(c_i) > 0.$$

Denote by  $\mathcal{B}^{\mathbf{r}}$  the ordered basis

(A.2.7) 
$$\mathcal{B}^{\mathbf{r}} = \left(c_1, c_2, \dots, c_l, b_{l+1}, b_{l+2}, \dots, b_n\right)$$

of  $\mathbb{E}^n$ . Define the sign of **r** by

(A.2.8) 
$$\operatorname{sign}(\mathbf{r}) = \begin{cases} 1 & \text{if } \mathcal{B} \text{ and } \mathcal{B}^{\mathbf{r}} \text{ define} \\ & \text{the same orientation of } \mathbb{E}^{n}, \\ -1 & \text{if } \mathcal{B} \text{ and } \mathcal{B}^{\mathbf{r}} \text{ define} \\ & \text{opposite orientations of } \mathbb{E}^{n}. \end{cases}$$

Let  $\Phi^{\mathbf{r}}$  be a rigged flag of length l in  $\mathbb{E}^n$ . Let P be a proper convex polytope in  $\mathbb{E}^n$ . Then the  $\Phi^{\mathbf{r}}$ -boundary  $\partial_{\Phi^{\mathbf{r}}}$  of P is defined by

(A.2.9) 
$$\partial_{\Phi^{\mathbf{r}}} P = r_l^{\min} \left( r_{l-1}^{\min} \left( \dots \left( r_2^{\min} \left( r_1^{\min}(P) \right) \dots \right) \right) \right)$$

where for each subset S of  $V_{i-1}$ 

(A.2.10) 
$$r_{i}^{\min}(S) = \begin{cases} v_{i} + r_{i}^{-1} \left( \min_{s \in S} r_{i}(s) \right) & \text{if } v_{i} \text{ is a vector in } V_{i-1} \\ & \text{such that} \\ & v_{i} + r_{i}^{-1} \left( \min_{s \in S} r_{i}(s) \right) \\ & \text{is a subset of } V_{i} \text{ with} \\ & \text{nonempty interior,} \\ & \emptyset & \text{if no such } v_{i} \text{ exists,} \end{cases}$$

which is a subset of  $V_i$  defined up to translation. Fix a translation invariant measure on the Euclidean space  $V_l$ . This determines the volume of the convex polytope  $\partial_{\Phi^r} P$ . Then define the Hadwiger invariant Had<sub> $\Phi$ </sub> of P with respect to  $\Phi$  by

(A.2.11) 
$$\operatorname{Had}_{\Phi}(P) = \sum_{\mathbf{r} \in \mathcal{R}ig(\Phi)} \operatorname{sign}(\mathbf{r}) \operatorname{Vol}(\partial_{\Phi^{\mathbf{r}}} P).$$

(A.3) **Remark** Each Hadwiger invariant defines a real-valued additive function on  $\mathbb{S}(\mathbb{E}^n)$ .

(A.4) Lemma Let P and Q be proper convex polytopes in  $\mathbb{E}^n$ . Then P are Q are translational scissors congruent if and only if for every l among  $0, 1, 2, \ldots, n$ , for each strict flag  $\Phi$  of length l,

(A.4.1) 
$$\operatorname{Had}_{\Phi}(P) = \operatorname{Had}_{\Phi}(Q).$$

*Proof.* See Corollary 2 in §4 of [Mo93a].

(A.5) Lemma Let  $(H_{\Phi})$  be a collection of real numbers indexed by the set of all strict flags  $\Phi$  in  $\mathbb{E}^n$  which vanishes for all but finitely many  $\Phi$ . Then there exists an element [P]in the scissors group  $\mathbb{S}(\mathbb{E}^n)$  such that

(A.5.1) 
$$\forall \text{ strict flag } \Phi \text{ in } \mathbb{E}^n \quad H_{\Phi} = \operatorname{Had}_{\Phi}([P])$$

if and only if for every l among 0, 1, 2, ..., n, for each strict flag

(A.5.2) 
$$\Phi = V_0 \supset V_1 \supset \cdots \supset V_l$$

of length l, for every i among  $1, 2, \ldots, l-1$ ,

(A.5.3) 
$$\sum_{\Phi' \in \mathcal{F}_{s}(\Phi,i)} H_{\Phi'} = 0$$

where  $\mathcal{F}_{s}(\Phi, i)$  is the set defined by

(A.5.4) 
$$\mathcal{F}_{s}(\Phi, i) = \left\{ \Phi' \text{ strict flag in } \mathbb{E}^{n} : \Phi' = V_{0} \supset \dots \supset V_{i-1} \supset U_{i} \supset V_{i+1} \supset \dots \supset V_{l} \right\},$$

and

(A.5.5) 
$$\sum_{\Phi' \in \mathcal{F}_{s}(\Phi,l)} H_{\Phi'} \wedge \mathbf{d}U_{l} = 0$$

where

(A.5.6) 
$$\Phi' = V_0 \supset V_1 \supset \cdots \supset V_{l-1} \supset U_l$$

and  $\mathbf{d}U_l$  is the element of  $\bigwedge^{n-l} \mathbb{E}^{n,*}$  defined as the product of the fixed orientation and translation invariant measure on  $U_l$ .

*Proof.* See Corollary 3 in §4 of [Mo93a].

(A.6) **Remark** The scissors group  $\mathbb{S}(\mathbb{E}^n)$  has the structure of a real vector space.

(A.7) Definition Let M be a Levi subgroup in  $\mathcal{L}$ . Let  $\mathcal{Y}_M$  and  $\mathcal{Z}_M$  be positive (G, M)orthogonal sets. Then  $\mathcal{Y}_M$  and  $\mathcal{Z}_M$  are said to be scissors congruent as (G, M)-orthogonal sets if the convex hull of  $\mathcal{Y}_M$  in  $\mathfrak{a}_M^G$  is translational scissors congruent to the convex hull of  $\mathcal{Z}_M$  in  $\mathfrak{a}_M^G$ . Define the scissors group of  $\mathfrak{a}_M^G$ , denoted by  $\mathbb{S}(\mathfrak{a}_M^G)$ , to be the quotient of the free abelian group generated by the positive (G, M)-orthogonal sets modulo translational scissors congruence in  $\mathfrak{a}_M^G$ .

(A.8) Remark The Hadwiger invariants of  $\mathcal{Y}_M$  are supported on the strict flags  $\Phi$  in  $\mathfrak{a}_M^G$  of the form

(A.8.1) 
$$\Phi = \mathfrak{a}_{M}^{L^{0}} \supset \mathfrak{a}_{M}^{L^{1}} \supset \cdots \supset \mathfrak{a}_{M}^{L^{l}}$$

where

$$(A.8.2) L^0 \supset L^1 \supset \cdots \supset L^l$$

is a nested chain of Levi subgroups in  $\mathcal{L}(M)$ . Denote the collection of such  $\Phi$  by  $\mathcal{F}_{s}(\mathfrak{a}_{M}^{G})$ . A rigging **r** of  $\Phi$  is equivalent to a nested chain of parabolic subgroups

(A.8.3) 
$$\mathbf{r} = \mathbf{Q}^0 \supset \mathbf{Q}^1 \supset \cdots \supset \mathbf{Q}^l$$

such that  $Q^i$  is a parabolic subgroup in  $\mathcal{P}(L^i)$  for each *i* among  $0, 1, 2, \ldots, l$ .

Fix a vector  $\xi$  in general position in  $\mathfrak{a}_{M}^{G}$ . Then  $\xi$  defines a total order on  $\Delta_{L}^{G}$  for each Levi subgroup L in  $\mathcal{L}(M)$ , which induces a consistent choice of signs for all rigged flags  $\Phi^{\mathbf{r}}$  with  $\Phi$  in  $\mathcal{F}_{s}(\mathfrak{a}_{M}^{G})$ . More precisely choose an element *s* of the Weyl group that stablizes M such that the parabolic subgroup  $sQ^{l}$  is standard. The nested chain

(A.8.4) 
$$sQ^0 \supset sQ^1 \supset \cdots \supset sQ^l$$

determines a sequence of positive roots  $\alpha^i$  where *i* ranges among  $1, 2, \ldots, l$ . Let  $\sigma$  be the permutation on *l* letters such that  $\alpha^{\sigma(i)}$  is strictly increasing with respect to the total order determined by  $\xi$ . Then the sign of the rigging **r** is equal to

(A.8.5) 
$$\operatorname{sign}(\mathbf{r}) = \operatorname{sign}(\operatorname{Det}(s))\operatorname{sign}(\sigma).$$

Each subspace  $\mathfrak{a}_{M}^{L}$  of  $\mathfrak{a}_{M}^{G}$  is equipped with the translation invariant measure determined by the coweight lattice. This choice of orientations and measures determines the numerical values of the Hadwiger invariants  $\operatorname{Had}_{\Phi}(\mathcal{Y}_{M})$ .

(A.9) **Definition** Let M be a Levi subgroup in  $\mathcal{L}$ , let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ . Define the scissors-congruence-valued orbital integral, or orbital integrohedron  $\mathbb{J}_{\mathrm{M}}^{\mathrm{G}}(X, \cdot)$  to be the vector-valued distribution on  $\mathfrak{g}(\mathbb{Q}_S)$  taking values in  $\bigoplus_{\Phi \in \mathcal{F}_{\mathrm{s}}(\mathfrak{a}_{\mathrm{M}}^{\mathrm{G}})} \mathbb{C}$  such that

(A.9.1)  $\forall f_S \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$ 

$$\mathbb{J}_{\mathcal{M}}^{\mathcal{G}}(X, f_{S}) = \left(\sum_{\mathbf{r} \in \mathcal{R}ig(\Phi)} \operatorname{sign}(\mathbf{r}) J_{\mathcal{M}}^{\mathcal{L}^{l}}(X, f_{S, \mathcal{Q}^{l}})\right)_{\Phi \in \mathcal{F}_{\mathrm{s}}(\mathfrak{a}_{\mathcal{M}}^{\mathcal{G}})}$$

where  $\Phi$ ,  $\mathbf{r}$ ,  $\mathbf{L}^{l}$  and  $\mathbf{Q}^{l}$  are related as in (A.8.1), (A.8.2) and (A.8.3).

For each Levi subgroup  $M_S$  in  $\mathcal{L}^{G_S}$  define  $\mathbb{J}_{M_S}^{G_S}(X, \cdot)$  by the analogous formula.

(A.10) Lemma Let M be a Levi subgroup in  $\mathcal{L}$ , let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ , let  $f_S$  be a real-valued Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$ . Then  $\mathbb{J}^{\mathrm{G}}_{\mathrm{M}}(X, f_S)$  defines a unique element of  $\mathbb{S}(\mathfrak{a}^{\mathrm{G}}_{\mathrm{M}})$ .

*Proof.* The Schwartz function  $f_S$  is real-valued, so  $\mathbb{J}_{M}^{G}(X, f_S)$  is a collection of real numbers indexed by the strict flags  $\Phi$  in  $\mathcal{F}_{s}(\mathfrak{a}_{M}^{G})$ . It suffices to verify (A.5.3) and (A.5.5).

The left hand side of (A.5.3) is equal to

(A.10.1) 
$$\sum_{\Phi' \in \mathcal{F}_{s}(\Phi,i)} \sum_{\mathbf{r}' \in \mathcal{R}ig(\Phi')} \operatorname{sign}(\mathbf{r}') J_{\mathrm{M}}^{\mathrm{L}^{l'}}(X, f_{S, \mathrm{Q}^{l'}})$$

$$= \sum_{\substack{\mathbf{Q}^{0'} \supset \mathbf{Q}^{1'} \supset \cdots \supset \mathbf{Q}^{l'} \\ \forall j=0,1,2,\dots,l \ Q^{j'} \in \mathcal{P}(\mathbf{L}^{j'}) \\ \forall j=0,1,2,\dots,l \ j \neq i \Rightarrow \mathbf{L}^{j'} = \mathbf{L}^{j}}} \operatorname{sign}(\mathbf{r}') J_{\mathbf{M}}^{\mathbf{L}^{l'}}(X, f_{S, \mathbf{Q}^{l'}})$$

where  $\Phi$ ,  $L^{j}$  and  $\Phi'$ ,  $\mathbf{r}'$ ,  $L^{j'}$ ,  $Q^{j'}$  are related as in (A.8.1), (A.8.2) and (A.8.3). The summands of the right hand side of (A.10.1) with a fixed minimal term  $Q^{l'}$  are in (1,1) correspondence with the sequences of roots ( $\alpha^{j}$ ) that are positive with respect to  $Q^{l'}$  such that

(A.10.2) 
$$\alpha^1, \alpha^2, \dots, \alpha^{i-1}, \alpha^{i+2}, \alpha^{i+3}, \dots, \alpha^l$$

are determined by  $\Phi$ . Hence the summand

(A.10.3) 
$$J_{\rm M}^{{\rm L}^{\prime\prime}}(X, f_{S,{\rm Q}^{\prime\prime}})$$

appears twice on the right hand side of (A.10.1) with opposite signs, so (A.10.1) vanishes. Hence  $\mathbb{J}_{M}^{G}(X, f_{S})$  satisfies (A.5.3).

The left hand side of (A.5.5) is equal to

(A.10.4) 
$$\sum_{\Phi' \in \mathcal{F}_{s}(\Phi,l)} \sum_{\mathbf{r}' \in \mathcal{R}ig(\Phi')} \operatorname{sign}(\mathbf{r}') J_{\mathrm{M}}^{\mathrm{L}''}(X, f_{S,\mathrm{Q}^{l'}}) \wedge \mathbf{d}\mathfrak{a}_{\mathrm{M}}^{\mathrm{L}^{l'}}$$
$$= \sum_{\substack{\mathrm{Q}^{0} \supset \mathrm{Q}^{1} \supset \cdots \supset \mathrm{Q}^{l-1} \supset \mathrm{Q}^{l'}\\\mathrm{Q}^{l'} \in \mathcal{P}(\mathrm{L}^{l'})}} \operatorname{sign}(\mathbf{r}') J_{\mathrm{M}}^{\mathrm{L}^{l'}}(X, f_{S,\mathrm{Q}^{l'}}) \wedge \mathbf{d}\mathfrak{a}_{\mathrm{M}}^{\mathrm{L}^{l'}}$$

where  $\Phi$ ,  $Q^j$  and  $\Phi'$ ,  $\mathbf{r}'$ ,  $L^{l'}$ ,  $Q^{l'}$  are related as in (A.8.1), (A.8.2) and (A.8.3). It suffices to show that (A.10.4) vanishes as a differential form on  $\mathfrak{a}_{M}^{L^{l-1}}$ . Let L be a Levi subgroup in  $\mathcal{L}^{L^{l-1}}(M)$  such that

(A.10.5) 
$$\dim(A_M/A_L) = 1,$$

denote by  $d\mathfrak{a}_L^{L^{l-1}}$  the differential form on  $\mathfrak{a}_M^{L^{l-1}}$  defined by pulling back the volume form on  $\mathfrak{a}_L^{L^{l-1}}$  along the natural projection

(A.10.6) 
$$\mathfrak{a}_{\mathrm{M}}^{\mathrm{L}^{l-1}} \longrightarrow \mathfrak{a}_{\mathrm{L}}^{\mathrm{L}^{l-1}}.$$

Then the orthogonal projection of (A.10.4) onto the one dimensional subspace

(A.10.7) 
$$\operatorname{span}(\mathbf{d}\mathfrak{a}_{\mathrm{L}}^{L^{l-1}}) \subset \bigwedge^{\dim(\mathrm{A}_{\mathrm{M}}/\mathrm{A}_{\mathrm{G}})-l} \mathfrak{a}_{\mathrm{M}}^{L^{l-1},*}$$

is equal to

(A.10.8) 
$$\sum_{\substack{\mathbf{Q}^0 \supset \mathbf{Q}^1 \supset \cdots \supset \mathbf{Q}^{l-1} \supset \mathbf{Q}^{l\prime} \\ \mathbf{Q}^{l\prime} \in \mathcal{P}(\mathbf{L}^{l\prime})}} \operatorname{sign}(\mathbf{r}') d_{\mathbf{M}}^{\mathbf{L}^{l-1}}(\mathbf{L}, \mathbf{L}^{l\prime}) J_{\mathbf{M}}^{\mathbf{L}^{l\prime}}(X, f_{S, \mathbf{Q}^{l\prime}}) \wedge \mathbf{d}\mathfrak{a}_{\mathbf{L}}^{\mathbf{L}^{l-1}}$$

by the definition of the constant  $d_{\rm M}^{{\rm L}^{l-1}}({\rm L},{\rm L}^{l\prime})$  in Remark (4.1.10). The summation in (A.10.8) is taken over the set

(A.10.9) 
$$\left\{ (\mathbf{L}^{l\prime}, \mathbf{Q}^{l\prime} \cap \mathbf{L}^{l-1}) : \begin{array}{l} \mathbf{L}^{l\prime} \in \mathcal{L}^{\mathbf{L}^{l-1}}(\mathbf{M}), \ \dim(\mathfrak{a}_{\mathbf{L}^{l\prime}}^{\mathbf{G}}) = l, \\ \mathbf{Q}^{l\prime} \in \mathcal{P}(\mathbf{L}^{l\prime}), \ \mathbf{Q}^{l\prime} \subset \mathbf{Q}^{l-1} \end{array} \right\}$$

$$\subset \mathcal{L}^{L^{l-1}}(M) \times \mathcal{F}^{L^{l-1}}(M)$$

Let  $\xi^{l-1}$  be the projection of the vector  $\xi$  in  $\mathfrak{a}_{M}^{G}$  used to define the orientation onto  $\mathfrak{a}_{M}^{L^{l-1}}$ . Then  $\xi^{l-1}$  determines a partial map

(A.10.10) 
$$s^{l-1}: \mathcal{L}^{L^{l-1}}(M) \times \mathcal{L}^{L^{l-1}}(M) \longrightarrow \mathcal{F}^{L^{l-1}}(M) \times \mathcal{F}^{L^{l-1}}(M)$$

as in Remark (4.1.11) which is positive in the sense that

$$(A.10.11) \qquad \qquad \operatorname{sign}(\mathbf{r}) = 1$$

if the rigging **r** corresponds to the element  $(L^{l'}, Q^{l,+})$  in the set (A.10.9) where  $Q^{l,+}$  denotes the second component of  $s^{l-1}(L, L^{l'})$ . Let  $Q^{l,-}$  denote the opposite parabolic of  $Q^{l,+}$ , then (A.10.9) is equal to the disjoint union

(A.10.12) 
$$\begin{cases} (\mathbf{L}^{l\prime}, \mathbf{Q}^{l,+}) : \ \mathbf{L}^{l\prime} \in \mathcal{L}^{\mathbf{L}^{l-1}}(\mathbf{M}) \end{cases} \coprod \\ \prod \left\{ (\mathbf{L}^{l\prime}, \mathbf{Q}^{l,-}) : \ \mathbf{L}^{l\prime} \in \mathcal{L}^{\mathbf{L}^{l-1}}(\mathbf{M}) \right\}, \end{cases}$$

hence (A.10.8) is equal to

(A.10.13)  

$$\begin{pmatrix} \sum_{\mathrm{L}^{l'} \in \mathcal{L}^{\mathrm{L}^{l-1}}(\mathrm{M})} d_{\mathrm{M}}^{\mathrm{L}^{l-1}}(\mathrm{L}, \mathrm{L}^{l'}) J_{\mathrm{M}}^{\mathrm{L}^{l'}}(X, f_{S, \mathrm{Q}^{l, +}}) \\ - \sum_{\mathrm{L}^{l'} \in \mathcal{L}^{\mathrm{L}^{l-1}}(\mathrm{M})} d_{\mathrm{M}}^{\mathrm{L}^{l-1}}(\mathrm{L}, \mathrm{L}^{l'}) J_{\mathrm{M}}^{\mathrm{L}^{l'}}(X, f_{S, \mathrm{Q}^{l, -}}) \end{pmatrix} \wedge \mathbf{d} \mathfrak{a}_{\mathrm{L}}^{\mathrm{L}^{l-1}}$$
(A.10.14)  

$$= \left( J_{\mathrm{L}}^{\mathrm{L}^{l-1}}(X^{\mathrm{L}}, f_{S, \mathrm{Q}^{l-1}}) - J_{\mathrm{L}}^{\mathrm{L}^{l-1}}(X^{\mathrm{L}}, f_{S, \mathrm{Q}^{l-1}}) \right) \wedge \mathbf{d} \mathfrak{a}_{\mathrm{L}}^{\mathrm{L}^{l-1}}$$

$$= 0$$

where the equality (A.10.14) follows from the S-local version of (4.1.12.2) and the fact that  $\mathbf{Q}^{l,-}$  is the second component of  $s^{l-1,-}(\mathbf{L},\mathbf{L}^{l'})$  where  $s^{l-1,-}(,)$  is the partial map determined as in Remark (4.1.11) by the vector  $-\xi^{l-1}$ . Hence (A.10.8) vanishes for every L, so (A.10.4) vanishes as a differential form on  $\mathfrak{a}_{\mathbf{M}}^{\mathbf{G}}$ . Hence  $\mathbb{J}_{\mathbf{M}}^{\mathbf{G}}(X, f_S)$  satisfies (A.5.5).

(A.11) Remark In §23 of [Ko05] Kottwitz defined weight factors and weighted orbital integrals taking values in the complexified K-group of the toric variety of the fan of the root hyperplanes in  $\mathfrak{a}_{M}^{G,*}$ . In §4 of [Mo93b] Morelli proved that this K-group is the additive group of translational scissors congruent classes of positive (G, M)-orthogonal sets whose vertices are contained in the coweight lattice in  $\mathfrak{a}_{M}^{G}$ .
(A.12) Definition Let M be a Levi subgroup in  $\mathcal{L}$ . Define the *total scissors ring* of  $\mathfrak{a}_{M}^{G}$ , denoted by  $\mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^{G})$ , to be the direct sum

(A.12.1) 
$$\mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^{G}) = \bigoplus_{L \in \mathcal{L}^{G}(M)} \mathbb{S}(\mathfrak{a}_{L}^{G}).$$

Define a bilinear product  $\boxtimes$  on  $\mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^G)$  by

(A.12.2) 
$$\forall \mathcal{L}_{1}, \mathcal{L}_{2} \in \mathcal{L}^{\mathcal{G}}(\mathcal{M}) \; \forall [\mathcal{Y}_{\mathcal{L}_{i}}] \in \mathbb{S}(\mathfrak{a}_{\mathcal{L}_{i}}^{\mathcal{G}}) \text{ where } i = 1, 2$$

$$[\mathcal{Y}_{\mathcal{L}_{1}}] \boxtimes [\mathcal{Y}_{\mathcal{L}_{2}}]$$

$$= \begin{cases} j^{*} \left( [\mathcal{Y}_{\mathcal{L}_{1}} \times \mathcal{Y}_{\mathcal{L}_{2}}] \right) & \text{if the natural map} \\ j : \mathfrak{a}_{\mathcal{L}_{1} \cap \mathcal{L}_{2}}^{\mathcal{G}} \longrightarrow \mathfrak{a}_{\mathcal{L}_{1}}^{\mathcal{G}} \oplus \mathfrak{a}_{\mathcal{L}_{2}}^{\mathcal{G}} \\ & \text{is an isomorphism,} \end{cases}$$

$$0 \quad \text{otherwise,}$$

where the  $(G \times G, L_1 \times L_2)$ -family  $\mathcal{Y}_{L_1} \times \mathcal{Y}_{L_2}$  is well-defined upto translational scissors congruence in  $\mathfrak{a}_{L_1}^G \oplus \mathfrak{a}_{L_2}^G$ , and  $j^*$  is the homomorphism

(A.12.3) 
$$j^*: \ \mathbb{S}(\mathfrak{a}_{L_1}^G \oplus \mathfrak{a}_{L_2}^G) \longrightarrow \ \mathbb{S}(\mathfrak{a}_{L_1 \cap L_2}^G)$$

induced by j.

(A.13) **Remark** The total scissors ring  $\mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^{G})$  is graded by

(A.13.1) 
$$\forall n \in \mathbb{N} \quad \left(\mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^{G})\right)_{n} = \bigoplus_{\substack{\mathrm{L} \in \mathcal{L}^{G}(M) \\ \dim(\mathfrak{a}_{\mathrm{L}}^{G}) = n}} \mathbb{S}(\mathfrak{a}_{\mathrm{L}}^{\mathrm{G}}).$$

The grading (A.13.1) has a refinement into the  $\mathcal{L}^{G}(M)$ -grading defined by

(A.13.2) 
$$\forall L \in \mathcal{L}^{G}(M) \quad \left(\mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^{G})\right)_{L} = \mathbb{S}(\mathfrak{a}_{L}^{G})$$

where the monoid structure on  $\mathcal{L}^{G}(M)$  is defined by intersection.

Each Levi subgroup L in  $\mathcal{L}^{G}(M)$  defines a homogeneous ideal

(A.13.3) 
$$\bigoplus_{\substack{L' \in \mathcal{L}^{G}(M) \\ L' \not\supset L}} \mathbb{S}(\mathfrak{a}_{L'}^{G}) \subset \mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^{G}),$$

and taking the quotient of  $\mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^G)$  modulo (A.13.3) defines the homomorphism

(A.13.4) 
$$q_{\mathrm{M}}^{\mathrm{L}} := \mathbb{S}(\mathfrak{a}_{\mathcal{L}(\mathrm{M})}^{\mathrm{G}}) \longrightarrow \mathbb{S}(\mathfrak{a}_{\mathcal{L}(\mathrm{L})}^{\mathrm{G}}).$$

The map  $q_{\mathrm{M}}^{\mathrm{G}}$  is the quotient modulo the augmentation ideal onto the graded component  $\left(\mathbb{S}(\mathfrak{a}_{\mathcal{L}(\mathrm{M})}^{\mathrm{G}})\right)_{\mathrm{G}}$ , which is a copy of  $\mathbb{R}$ .

(A.14) **Definition** Let M be a Levi subgroup in  $\mathcal{L}$ , let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ . Define the total orbital integrohedron  $\mathcal{J}_{\mathrm{M}}^{\mathrm{G}}(X, )$  to be the  $\mathbb{S}(\mathfrak{a}_{\mathcal{L}(\mathrm{M})}^{\mathrm{G}}) \otimes \mathbb{C}$ -valued distribution on  $\mathfrak{g}(\mathbb{Q}_S)$ such that

(A.14.1)  $\forall f_S \in \mathcal{S}(\mathfrak{g}(\mathbb{Q}_S))$ 

$$\mathcal{J}_{\mathbf{M}}^{\mathbf{G}}(X, f_S) = \left( (-1)^{\dim(\mathbf{A}_{\mathbf{L}}/\mathbf{A}_{\mathbf{G}})} |\mathbf{W}_0^{\mathbf{L}}| |\mathbf{W}_0^{\mathbf{G}}|^{-1} \mathbb{J}_{\mathbf{L}}^{\mathbf{G}}(X^{\mathbf{L}}, f_S) \right)_{\mathbf{L} \in \mathcal{L}^{\mathbf{G}}(\mathbf{M})}$$

For each Levi subgroup  $M_S$  in  $\mathcal{L}^{G_S}$  define  $\mathcal{J}_{M_S}^{G_S}(X, \cdot)$  by the analogous formula.

(A.15) Lemma (Induction and splitting of orbital integrohedra) Let M be a Levi subgroup in  $\mathcal{L}$ , let X be an element of  $\mathfrak{m}(\mathbb{Q}_S)$ , let  $f_S$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$ .

• Let L be a Levi subgroup in  $\mathcal{L}(M)$ , then

(A.15.1) 
$$\mathcal{J}_{\mathrm{L}}^{\mathrm{G}}(X^{\mathrm{L}}, f_{S}) = q_{\mathrm{M}}^{\mathrm{L}} \Big( \mathcal{J}_{\mathrm{M}}^{\mathrm{G}}(X, f_{S}) \Big).$$

• Let S be the set  $\{v_1, v_2\}$ , let  $f_S$  be of the form  $f_{v_1} \otimes f_{v_2}$  where  $f_{v_i}$  is a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_{v_i})$  where the index i is 1 or 2, then

(A.15.2) 
$$\mathcal{J}_{\mathrm{M}}^{\mathrm{G}}(X, f_{v_{1}} \otimes f_{v_{2}}) = \mathcal{J}_{\mathrm{M}}^{\mathrm{G}}(X, f_{v_{1}}) \boxtimes \mathcal{J}_{\mathrm{M}}^{\mathrm{G}}(X, f_{v_{2}}).$$

Local identities analogous to (A.15.1) also hold for  $G_v$ .

*Proof.* The induction identity (A.15.1) follows from the definition (A.14.1).

The splitting identity (A.15.2) follows by comparing the Hadwiger invariants of the two sides of (A.15.2). Retain the notations of (A.9.1). Let L be a Levi subgroup in  $\mathcal{L}(M)$ , let  $\Phi$  be a strict flag in  $\mathcal{F}_{s}(\mathfrak{a}_{L}^{G})$ . It suffices to show that the  $\Phi$ -components of the Hadwiger invariants of the L-components of the two sides of (A.15.2) are equal.

The L-component of the right hand side of (A.15.2) is

(A.15.3) 
$$\sum_{\substack{L_1,L_2 \in \mathcal{L}(L) \\ \mathfrak{a}_L^G \to \mathfrak{a}_{L_1}^G \oplus \mathfrak{a}_{L_2}^G \\ \text{isomorphism}}} (-1)^{\dim(A_L/A_G)} |W_0^{L_1}| |W_0^G|^{-1} |W_0^G|^{-1} \times \mathbb{J}_{L_1}^G \oplus \mathfrak{a}_{L_2}^G \\ \times \mathbb{J}_{L_1}^G (X^{L_1}, f_{v_1}) \boxtimes \mathbb{J}_{L_2}^G (X^{L_2}, f_{v_2}),$$

whose Hadwiger invariant has  $\Phi$ -component equal to

(A.15.4) 
$$\sum_{L_1,L_2 \in \mathcal{L}(L)} (-1)^{\dim(A_L/A_G)} |W_0^{L_1}| |W_0^G|^{-1} |W_0^{L_2}| |W_0^G|^{-1} \times d_L^{L^l}(L_1^l, L_2^l) \left(\sum_{\mathbf{r}_1 \in \mathcal{R}ig(\Phi_1)} \operatorname{sign}(\mathbf{r}_1) J_{L_1}^{L_1^l}(X^{L_1}, f_{v_1, Q_1^l})\right) \times d_L^{L^l}(L_1^l, L_2^l) \left(\sum_{\mathbf{r}_1 \in \mathcal{R}ig(\Phi_1)} \operatorname{sign}(\mathbf{r}_1) J_{L_1}^{L_1^l}(X^{L_1}, f_{v_1, Q_1^l})\right) \times d_L^{L^l}(L_1^l, L_2^l) \left(\sum_{\mathbf{r}_1 \in \mathcal{R}ig(\Phi_1)} \operatorname{sign}(\mathbf{r}_1) J_{L_1}^{L_1^l}(X^{L_1}, f_{v_1, Q_1^l})\right) \right) \times d_L^{L^l}(L_1^l, L_2^l) \left(\sum_{\mathbf{r}_1 \in \mathcal{R}ig(\Phi_1)} \operatorname{sign}(\mathbf{r}_1) J_{L_1}^{L_1^l}(X^{L_1}, f_{v_1, Q_1^l})\right) \right) \times d_L^{L^l}(L_1^l, L_2^l) \left(\sum_{\mathbf{r}_1 \in \mathcal{R}ig(\Phi_1)} \operatorname{sign}(\mathbf{r}_1) J_{L_1}^{L_1^l}(X^{L_1}, f_{v_1, Q_1^l})\right) \right) \times d_L^{L^l}(L_1^l, L_2^l) \left(\sum_{\mathbf{r}_1 \in \mathcal{R}ig(\Phi_1)} \operatorname{sign}(\mathbf{r}_1) J_{L_1}^{L_1^l}(X^{L_1}, f_{v_1, Q_1^l})\right) \times d_L^{L^l}(L_1^l, L_2^l) \left(\sum_{\mathbf{r}_1 \in \mathcal{R}ig(\Phi_1)} \operatorname{sign}(\mathbf{r}_1) J_{L_1}^{L^l}(X^{L_1}, f_{v_1, Q_1^l})\right) \right)$$

$$\times \left(\sum_{\mathbf{r}_2 \in \mathcal{R}ig(\Phi_2)} \operatorname{sign}(\mathbf{r}_2) J_{\mathrm{L}_2}^{\mathrm{L}_2^l}(X^{\mathrm{L}_2}, f_{v_2, \mathrm{Q}_2^l})\right)$$

where  $\Phi_i$  denotes the strict flag in  $\mathfrak{a}_{L_i}^G$  induced by  $\Phi$ , and  $\Phi_i$ ,  $\mathbf{r}_i$ ,  $L_i^l$  and  $Q_i^l$  are related as in (A.8.1), (A.8.2) and (A.8.3) where the index *i* is 1 or 2. Combine  $\mathbf{r}_1$  and  $\mathbf{r}_2$  into a rigging  $\mathbf{r}$  of  $\Phi$ , the expression (A.15.4) becomes

$$(A.15.5) \qquad \sum_{L_{1},L_{2}\in\mathcal{L}(L)} (-1)^{\dim(A_{L}/A_{G})} |W_{0}^{L_{1}}||W_{0}^{G}|^{-1} |W_{0}^{L_{2}}||W_{0}^{G}|^{-1} \times \\ \times \sum_{Q_{1}^{l}\in\mathcal{P}(L_{1}^{l})} \sum_{Q_{2}^{l}\in\mathcal{P}(L_{2}^{l})} \operatorname{sign}(\mathbf{r}) d_{L}^{L^{l}}(L_{1}^{l}, L_{2}^{l}) \times \\ \times J_{L_{1}}^{L_{1}^{l}}(X^{L_{1}}, f_{v_{1},Q_{1}^{l}}) J_{L_{2}}^{L_{2}^{l}}(X^{L_{2}}, f_{v_{2},Q_{2}^{l}}) \\ = \sum_{L_{1},L_{2}\in\mathcal{L}(L)} (-1)^{\dim(A_{L}/A_{G})} |W_{0}^{L_{1}}||W_{0}^{G}|^{-1} |W_{0}^{L_{2}}||W_{0}^{G}|^{-1} \times \\ \times \sum_{Q_{1}^{l}\in\mathcal{P}(L_{1}^{l})} \sum_{Q_{2}^{l}\in\mathcal{P}(L_{2}^{l})} \operatorname{sign}(\mathbf{r}) d_{L}^{L^{l}}(L_{1}^{l}, L_{2}^{l}) \times \\ \times \sum_{L_{1}^{\circ}\in\mathcal{L}^{L_{1}^{l}}(L)} d_{L}^{L_{1}^{l}}(L_{1}, L_{1}^{\circ}) J_{L}^{L_{2}^{\circ}}(X^{L}, f_{v_{1},Q_{1}^{l,\circ}}) \times \\ \times \sum_{L_{1}^{\circ}\in\mathcal{L}^{L_{1}^{l}}(L)} d_{L}^{L_{2}^{l}}(L_{2}, L_{2}^{\circ}) J_{L}^{L_{2}^{\circ}}(X^{L}, f_{v_{2},Q_{2}^{l,\circ}}) \\ \end{array}$$

by (4.1.12.2), where  $\mathbf{Q}_i^{l,\circ}$  is the second component of  $s_{\mathbf{Q}_i^l}(\mathbf{L}_i, \mathbf{L}_i^\circ)$  for the choice of a collection of partial maps  $(s_{\mathbf{Q}_i^l})_{\mathbf{Q}_i^l \in \mathcal{P}(\mathbf{L}_1')}$  such that

(A.15.6) 
$$\forall \mathbf{Q}_i^l \in \mathcal{P}(\mathbf{L}_i^l) \qquad \mathbf{Q}_i^{l,\circ} \subset \mathbf{Q}_i^l$$

where the index *i* is 1 or 2. The choice of such a collection is equivalent to the choice of a collection of vectors  $(\xi_{\mathbf{Q}_i^l})_{\mathbf{Q}_i^l \in \mathcal{P}(\mathbf{L}_1')}$ 

(A.15.7) 
$$\forall \mathbf{Q}_i^l \in \mathcal{P}(\mathbf{L}_i^l) \qquad \xi_{\mathbf{Q}_i^l} \in \mathfrak{a}_{\mathbf{L}}^{\mathbf{L}_i^l}$$

in general position and positive with respect to  $\mathbf{Q}_i^l.$  By definition

(A.15.8) 
$$d_{\mathrm{L}}^{\mathrm{L}^{l}}(\mathrm{L}_{1}^{l},\mathrm{L}_{2}^{l})d_{\mathrm{L}}^{\mathrm{L}_{1}^{l}}(\mathrm{L}_{1},\mathrm{L}_{1}^{\circ})d_{\mathrm{L}}^{\mathrm{L}_{2}^{l}}(\mathrm{L}_{2},\mathrm{L}_{2}^{\circ}) = d_{\mathrm{L}}^{\mathrm{L}^{l}}(\mathrm{L}_{1}^{\circ},\mathrm{L}_{2}^{\circ}),$$

hence the right hand side of (A.15.5) is equal to

(A.15.9) 
$$(-1)^{\dim(A_{L}/A_{G})} \sum_{\mathbf{r} \in \mathcal{R}ig(\Phi)} \operatorname{sign}(\mathbf{r}) \times$$

$$\begin{split} & \times \sum_{\mathbf{L}_{1},\mathbf{L}_{2}\in\mathcal{L}(\mathbf{L})} \sum_{\substack{\mathbf{L}_{1}^{\circ}\in\mathcal{L}^{\mathbf{L}_{1}^{l}}(\mathbf{L})\\\mathbf{L}_{2}^{\circ}\in\mathcal{L}^{\mathbf{L}_{2}^{l}}(\mathbf{L})}} & |\mathbf{W}_{0}^{\mathbf{L}_{1}}||\mathbf{W}_{0}^{\mathbf{G}}|^{-1}|\mathbf{W}_{0}^{\mathbf{L}_{2}}||\mathbf{W}_{0}^{\mathbf{G}}|^{-1} \times \\ & \times d_{\mathbf{L}}^{\mathbf{L}^{l}}(\mathbf{L}_{1}^{\circ},\mathbf{L}_{2}^{\circ})J_{\mathbf{L}}^{\mathbf{L}_{2}^{\circ}}(X^{\mathbf{L}},f_{v_{1},\mathbf{Q}_{1}^{l,\circ}})J_{\mathbf{L}}^{\mathbf{L}_{2}^{\circ}}(X^{\mathbf{L}},f_{v_{2},\mathbf{Q}_{2}^{l,\circ}}) \end{split}$$

where  $(Q_1^{l,\circ},Q_2^{l,\circ})$  is the image of  $(L_1^\circ,L_2^\circ)$  under the partial map  $s_{Q_1^l,Q_2^l}$  determined by the vector

(A.15.10) 
$$\frac{\xi_{\mathbf{Q}_1^l}}{2} - \frac{\xi_{\mathbf{Q}_2^l}}{2} \in \mathfrak{a}_{\mathbf{L}}^{\mathbf{L}^l} = \mathfrak{a}_{\mathbf{L}}^{\mathbf{L}_1^l} \oplus \mathfrak{a}_{\mathbf{L}}^{\mathbf{L}_2^l}.$$

For each fixed L and fixed  $\Phi$  with minimal term L<sup>*l*</sup>, the map

(A.15.11) 
$$\begin{cases} (L_1, L_2, L_1^\circ, L_2^\circ) : & L_1 \in \mathcal{L}(L), \ L_2 \in \mathcal{L}(L), \\ L_1^\circ \in \mathcal{L}^{L_1^l}(L), \ L_2^\circ \in \mathcal{L}^{L_2^l}(L) \end{cases} \\ \longrightarrow & \left\{ (L_1^\circ, L_2^\circ) : \ L_1^\circ \in \mathcal{L}^{L^l}(L), \ L_2^\circ \in \mathcal{L}^{L^l}(L) \right\} \end{cases}$$

defined by

(A.15.12) 
$$(L_1, L_2, L_1^{\circ}, L_2^{\circ}) \mapsto (L_1^{\circ}, L_2^{\circ})$$

is surjective. Fix Levi subgroups  $L_1$  and  $L_2$  in  $\mathcal{L}(L)$  and parabolic subgroups  $P_1$  in  $\mathcal{P}(L_1)$ and  $P_2$  in  $\mathcal{P}(L_2)$ , and let  $P_{1,2}$  be the parabolic subgroup in  $\mathcal{P}(L)$  whose unipotent radical  $N_{P_{1,2}}$  contains  $N_{P_1}$  and  $N_{P_2}$ , then the map (A.15.12) restricts to a bijection between Levi subgroups  $L_1^{\circ}$  and  $L_2^{\circ}$  that are standard with respect to  $P_1$  and  $P_2$  or  $P_{1,2}$ . Hence each fiber of the map (A.15.12) containing the pair ( $L_1, L_2$ ) has cardinality

(A.15.13) 
$$\frac{|W_0^G|/|W_0^L|}{\left(|W_0^G|/|W_0^{L_1}|\right)\left(|W_0^G|/|W_0^{L_2}|\right)},$$

hence (A.15.9) is equal to

$$(A.15.14) \qquad (-1)^{\dim(A_{L}/A_{G})}|W_{0}^{L}||W_{0}^{G}|^{-1}\sum_{\mathbf{r}\in\mathcal{R}ig(\Phi)}\operatorname{sign}(\mathbf{r}) \times \\ \times \sum_{L_{1}^{\circ},L_{2}^{\circ}\in\mathcal{L}^{L^{l}}(L)} d_{L}^{L^{l}}(L_{1}^{\circ},L_{2}^{\circ})J_{L}^{L_{1}^{\circ}}(X^{L},f_{v_{1},Q_{1}^{l,\circ}})J_{L}^{L_{2}^{\circ}}(X^{L},f_{v_{2},Q_{2}^{l,\circ}}) \\ = (-1)^{\dim(A_{L}/A_{G})}|W_{0}^{L}||W_{0}^{G}|^{-1}\sum_{\mathbf{r}\in\mathcal{R}ig(\Phi)}\operatorname{sign}(\mathbf{r})J_{L}^{L^{l}}(X^{L},f_{v_{1},Q^{l}}\otimes f_{v_{2},Q^{l}})$$

by (4.1.12.3) where  $\Phi$ ,  $\mathbf{r}$ ,  $\mathbf{L}^{l}$  and  $\mathbf{Q}^{l}$  are related as in (A.8.1), (A.8.2) and (A.8.3), which is equal to the  $\Phi$ -component of the Hadwiger invariant of the L-component of the left hand side of (A.15.2).

(A.16) **Remark** The orbital integrohedra  $\mathbb{J}_{M}^{G}(X, \cdot)$  and  $\mathcal{J}_{M}^{G}(X, \cdot)$  could be alternatively defined as distributions on  $\mathfrak{g}(\mathbb{Q}_{S})$  obtained by integrating against certain polyhedron-valued weight factors with a formula analogous to (3.1.5.3) if X is regular semisimple.

## **B** Appendix: The example of GL(2)

In this appendix some explicit computations for the group GL(2) are carried out. Similar local computations could be found in [Ev98].

(B.1) Definition In this appendix v denotes an odd rational prime p, and || denotes the p-adic absolute value. Let G denote the group  $GL(2, \mathbb{Q}_p)$ , let M denote the minimal Levi subgroup of G consisting of the diagonal matrices, then M is the Levi component of the Borel subgroup B consisting of the upper triangular matrices. The groups G and M are the only standard Levi subgroups of G.

Let  $\mathfrak{g}'$  denote the Lie algebra  $\mathfrak{sl}(2, \mathbb{Q}_p)$ , equipped with the adjoint action of G from the right. Let  $\mathfrak{m}'$  denote the intersection of  $\mathfrak{m}$  and  $\mathfrak{g}'$ . Denote by  $\mathcal{A}$  the affine quotient  $\mathfrak{g}'/\!/ \mathbb{G}$  which is identified with the affine line  $\mathbb{L}$  via the negative determinant map

$$(B.1.1) \qquad -\det: \quad \mathfrak{g}' \longrightarrow \quad \mathbb{L} = \mathcal{A}.$$

(B.2) **Remark** Let  $\mathfrak{z}$  denote the center of  $\mathfrak{g}$  consisting of the diagonal matrices on which G operates trivially. Since

$$(B.2.1) g = g' \oplus \mathfrak{z}$$

as representations of G, for computing the local basic functions and the local Harish-Chandra transforms it suffices to consider  $\mathfrak{g}'$  instead of  $\mathfrak{g}$ .

(B.3) Definition The discriminant function D on  $\mathcal{A}$  is equal to 4X where X denotes the coordinate function on  $\mathcal{A}$ . Let  $\eta$  be an element of  $\mathcal{A}_{reg}(\mathbb{Q}_p)$  which is identified with the subset of units  $\mathbb{Q}_p^{\times}$  of  $\mathbb{Q}_p$ . Then  $\eta$  is said to be

- split if  $\eta$  has a square root in  $\mathbb{Q}_p$ ;
- unramified elliptic if  $\eta$  does not have a square root in  $\mathbb{Q}_p$  but the *p*-adic valuation of  $\eta$  is even;
- ramified elliptic if the p-adic valuation of  $\eta$  is odd.

The image of  $\mathfrak{m}'_{\text{reg.ss}}(\mathbb{Q}_p)$  in  $\mathcal{A}_{\text{reg}}(\mathbb{Q}_p)$  is equal to the subset of split elements.

(B.4) **Definition** The algebraic differential form

(B.4.1) 
$$d_{\eta}X = \frac{dx \wedge dy}{y} = \frac{dy \wedge dz}{2x} = \frac{dz \wedge dx}{z}$$

on the orbit

(B.4.2) 
$$\mathfrak{g}'_{\eta} = \left\{ \left( \begin{array}{cc} x & y \\ z & -x \end{array} \right) \in \mathfrak{g}' : x^2 + yz = \eta \right\}$$

where  $\eta$  lies in  $\mathcal{A}_{reg}$  is invariant under the action of G.

**(B.5)** Lemma Let f be a Schwartz function on  $\mathfrak{g}'(\mathbb{Q}_p)$ , then

(B.5.1) 
$$\forall \eta \in \mathcal{A}_{\mathrm{reg}}(\mathbb{Q}_p) \qquad I_{\mathrm{G}}^{\mathrm{G}}(\eta, f) = \frac{1}{1 + p^{-1}} \int_{\mathfrak{g}'_{\eta}(\mathbb{Q}_p)} f(X) \, |\mathrm{d}_{\eta}X|.$$

*Proof.* There exists a multiplicative constant  $\lambda$  such that

(B.5.2) 
$$\forall f \in \mathcal{S}(\mathfrak{g}'(\mathbb{Q}_p)) \ \forall \eta \in \mathcal{A}_{\mathrm{reg}}(\mathbb{Q}_p)$$

$$I_{\mathcal{G}}^{\mathcal{G}}(\eta, f) = \lambda \cdot \int_{\mathfrak{g}_{\eta}'(\mathbb{Q}_p)} f(X) |\mathrm{d}_{\eta}X|,$$

hence the constant  $\lambda$  is determined by

(B.5.3) 
$$I_{\mathcal{G}}^{\mathcal{G}}(1,\mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)}) = \lambda \cdot \int_{\mathfrak{g}'_1(\mathbb{Q}_p)} \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)}(X) |\mathrm{d}_1 X|.$$

By parabolic descent along the Borel subgroup B, the left hand side of (B.5.3) is equal to

(B.5.4) 
$$I_{\mathrm{M}}^{\mathrm{M}}(1,(\mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)})_{\mathrm{B}}) = I_{\mathrm{M}}^{\mathrm{M}}(1,\mathbb{I}_{\mathfrak{m}'(\mathbb{Z}_p)})$$

which is equal to one. Hence by (B.5.3) the reciprocal of  $\lambda$  is equal to

$$(B.5.5) \qquad \int_{\mathfrak{g}_{1}^{\prime}(\mathbb{Q}_{p})} \mathbb{I}_{\mathfrak{g}^{\prime}(\mathbb{Z}_{p})} \left( \left( \begin{array}{c} x & y \\ z & -x \end{array} \right) \right) \left| \frac{\mathrm{d}x \wedge \mathrm{d}y}{y} \right|$$
$$= \int_{\mathbb{Z}_{p}} \left( \int_{\left\{ y \in \mathbb{Z}_{p} \colon z = \frac{1-x^{2}}{y} \in \mathbb{Z}_{p} \right\}} \frac{\mathrm{d}y}{|y|} \right) \mathrm{d}x$$
$$= \int_{|1+x|<1} \left( \int_{|1-x^{2}| \leq |y| \leq 1} \frac{\mathrm{d}y}{|y|} \right) \mathrm{d}x +$$
$$+ \int_{|1-x|<1} \left( \int_{|1-x^{2}| \leq |y| \leq 1} \frac{\mathrm{d}y}{|y|} \right) \mathrm{d}x +$$
$$+ \int_{|1-x|=1} \left( \int_{|1-x^{2}| \leq |y| \leq 1} \frac{\mathrm{d}y}{|y|} \right) \mathrm{d}x$$
$$= \sum_{i=1}^{\infty} \left( \int_{|1+x|=p^{-i}} \mathrm{d}x \right) \left( \int_{p^{-i} \leq |y| \leq 1} \frac{\mathrm{d}y}{|y|} \right) +$$

$$\begin{split} &+ \sum_{j=1}^{\infty} \left( \int_{|1-x|=p^{-j}} \mathrm{d}x \right) \left( \int_{p^{-j} \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right) + \\ &+ \left( \int_{|1+x|=1}^{|1+x|=1} \mathrm{d}x \right) \left( \int_{|y|=1} \frac{\mathrm{d}y}{|y|} \right) \\ &= \sum_{i=1}^{\infty} \left( \frac{1}{p^{i}} - \frac{1}{p^{i+1}} \right) \left( \frac{(i+1)(p-1)}{p} \right) + \\ &+ \sum_{j=1}^{\infty} \left( \frac{1}{p^{j}} - \frac{1}{p^{j+1}} \right) \left( \frac{(j+1)(p-1)}{p} \right) + \\ &+ \left( \frac{p-2}{p} \right) \left( \frac{p-1}{p} \right) \\ &= \frac{p-1}{p} \left( 2 \cdot \left( \frac{1}{p} + \frac{1}{p-1} \right) + \frac{p-2}{p} \right) \\ &= 1 + p^{-1}. \end{split}$$

(B.6)	Lemma	Let $\eta$ be	$an \ element$	of	$\mathcal{A}_{\mathrm{reg}}(\mathbb{Q}_p),$	then
-------	-------	---------------	----------------	----	---	------

$$(B.6.1) \qquad \phi_{0,v}(\eta) = \begin{cases} 1 & \text{if } \eta \text{ is split} \\ and \text{ contained in } \mathbb{Z}_p - \{0\}, \\ 1 - \frac{2|\eta|^{1/2}}{1+p} & \text{if } \eta \text{ is unramified elliptic} \\ and \text{ contained in } \mathbb{Z}_p - \{0\}, \\ 1 - p^{-1/2}|\eta|^{1/2} & \text{if } \eta \text{ is ramified elliptic} \\ and \text{ contained in } \mathbb{Z}_p - \{0\}, \\ 0 & \text{if } \eta \text{ does not lie in } \mathbb{Z}_p - \{0\}. \end{cases}$$

Proof. If  $\eta$  does not lie in  $\mathbb{Z}_p - \{0\}$ , then the orbit  $\mathfrak{g}'_{\eta}(\mathbb{Q}_p)$  is disjoint from the support of the characteristic function  $\mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)}$ , hence  $\phi_{0,v}(\eta)$  vanishes. If  $\eta$  is split and contained in  $\mathbb{Z} - \{0\}$ , then by (B.5.4) the value  $\phi_{0,v}(\eta)$  is equal to one.

If  $\eta$  is unramified elliptic and contained in  $\mathbb{Z}_p - \{0\}$ , then by (B.5.1)

(B.6.2) 
$$\phi_{0,v}(\eta) = \frac{1}{1+p^{-1}} \int_{\mathfrak{g}_{\eta}'(\mathbb{Q}_p)} \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)}(X) |\mathrm{d}_{\eta}X|$$
$$= \frac{1}{1+p^{-1}} \int_{\mathfrak{g}_{\eta}'(\mathbb{Q}_p)} \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)}\left(\left(\begin{array}{cc} x & y \\ z & -x \end{array}\right)\right) \left|\frac{\mathrm{d}x \wedge \mathrm{d}y}{y}\right|$$

(B.6.3) 
$$= \frac{1}{1+p^{-1}} \int_{|x| \le 1} \left( \int_{|\eta-x^2| \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right) \,\mathrm{d}x$$

Decompose the integral on the right hand side of (B.6.3) over two disjoint regions:

• If  $|x^2|$  is strictly less than  $|\eta|$ , then  $|\eta - x^2|$  is equal to  $|\eta|$ , hence

(B.6.4) 
$$\int_{|x^2| < |\eta|} \left( \int_{|\eta - x^2| \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right) \mathrm{d}x$$
$$= \left( \int_{|x^2| < |\eta|} \mathrm{d}x \right) \left( \int_{|\eta| \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right)$$
$$= \left( p^{-\left(\frac{\operatorname{val}(\eta)}{2} + 1\right)} \right) \left( \frac{(\operatorname{val}(\eta) + 1)(p - 1)}{p} \right)$$

where  $val(\eta)$  denotes the *p*-adic valuation of  $\eta$ .

• If  $|x^2|$  is greater than or equal to  $|\eta|$ , then  $|\eta - x^2|$  is equal to  $|x^2|$ , hence

$$(B.6.5) \qquad \int_{|\eta| \le |x^2| \le 1} \left( \int_{|\eta-x^2| \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right) \mathrm{d}x$$

$$= \int_{|\eta| \le |x^2| \le 1} \left( \int_{|x^2| \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right) \mathrm{d}x$$

$$= \int_{|\eta| \le |x^2| \le 1} (2\mathrm{val}(x) + 1) \left(\frac{p-1}{p}\right) \mathrm{d}x$$

$$= \frac{p-1}{p} \left( 1 - p^{-\left(\frac{\mathrm{val}(\eta)}{2} + 1\right)} + 2 \int_{|\eta| \le |x^2| \le 1} \mathrm{val}(x) \mathrm{d}x \right)$$

$$= \frac{p-1}{p} \left( 1 - p^{-\left(\frac{\mathrm{val}(\eta)}{2} + 1\right)} + 2 \sum_{i=0}^{\frac{\mathrm{val}(\eta)}{2}} i \left(\frac{1}{p^i} - \frac{1}{p^{i+1}}\right) \right)$$

$$= \frac{p-1}{p} \left( 1 - p^{-\left(\frac{\mathrm{val}(\eta)}{2} + 1\right)} + \frac{1 - p^{-\left(\frac{\mathrm{val}(\eta)}{2} + 1\right)}}{p-1} \right)$$

Hence the right hand side of (B.6.3) is equal to

(B.6.6) 
$$\frac{1}{1+p^{-1}} \cdot \frac{p-1}{p} \left( \left( \operatorname{val}(\eta) + 1 \right) p^{-\left( \frac{\operatorname{val}(\eta)}{2} + 1 \right)} + \left( \operatorname{val}(\eta) + 2 \right) p^{-\left( \frac{\operatorname{val}(\eta)}{2} + 1 \right)} + \left( 1 - p^{-\left( \frac{\operatorname{val}(\eta)}{2} + 1 \right)} \right) \left( 1 + \frac{2}{p-1} \right) \right)$$

$$= \frac{p-1}{p+1} \left( -p^{-\left(\frac{\operatorname{val}(\eta)}{2}+1\right)} - \left(\frac{p+1}{p-1}\right) p^{-\left(\frac{\operatorname{val}(\eta)}{2}+1\right)} + \frac{p+1}{p-1} \right)$$
$$= 1 - \frac{2}{p+1} p^{-\frac{\operatorname{val}(\eta)}{2}}.$$

If  $\eta$  is ramified elliptic and contained in  $\mathbb{Z}_p - \{0\}$ , then by (B.5.1)

(B.6.7) 
$$\phi_{0,v}(\eta) = \frac{1}{1+p^{-1}} \int_{|x| \le 1} \left( \int_{|\eta-x^2| \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right) \,\mathrm{d}x$$

Decompose the integral on the right hand side of (B.6.7) over two disjoint regions:

• If  $|x^2|$  is less than  $|\eta|$ , then  $|\eta - x^2|$  is equal to  $|\eta|$ , hence

(B.6.8) 
$$\int_{|x^2| < |\eta|} \left( \int_{|\eta - x^2| \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right) \mathrm{d}x$$
$$= \left( \int_{|x^2| < |\eta|} \mathrm{d}x \right) \left( \int_{|\eta| \le |y| \le 1} \frac{\mathrm{d}y}{|y|} \right)$$
$$= \left( p^{-\left(\frac{\operatorname{val}(\eta) + 1}{2}\right)} \right) \left( \frac{(\operatorname{val}(\eta) + 1)(p - 1)}{p} \right).$$

• If  $|x^2|$  is greater than  $|\eta|$ , then  $|\eta - x^2|$  is equal to  $|x^2|$ , hence

$$(B.6.9) \qquad \int_{|\eta| \le |x^2| \le 1} \left( \int_{|\eta-x^2| \le |y| \le 1} \frac{dy}{|y|} \right) dx \\ = \int_{|\eta| \le |x^2| \le 1} \left( \int_{|x^2| \le |y| \le 1} \frac{dy}{|y|} \right) dx \\ = \int_{|\eta| \le |x^2| \le 1} (2\text{val}(x) + 1) \left(\frac{p-1}{p}\right) dx \\ = \frac{p-1}{p} \left( 1 - p^{-\left(\frac{\text{val}(\eta)+1}{2}\right)} + 2 \int_{|\eta| \le |x^2| \le 1} \text{val}(x) dx \right) \\ = \frac{p-1}{p} \left( 1 - p^{-\left(\frac{\text{val}(\eta)+1}{2}\right)} + 2 \sum_{i=0}^{\frac{\text{val}(\eta)-1}{2}} i \left(\frac{1}{p^i} - \frac{1}{p^{i+1}}\right) \right) \\ = \frac{p-1}{p} \left( 1 - p^{-\left(\frac{\text{val}(\eta)+1}{2}\right)} + \frac{1 - p^{-\left(\frac{\text{val}(\eta)+1}{2}\right)}}{p-1} \right) \right).$$

Hence the right hand side of (B.6.7) is equal to

(B.6.10) 
$$\frac{1}{1+p^{-1}} \cdot \frac{p-1}{p} \left( \left( \operatorname{val}(\eta)+1 \right) p^{-\left(\frac{\operatorname{val}(\eta)+1}{2}\right)} + \left( 1-p^{-\left(\frac{\operatorname{val}(\eta)+1}{2}\right)} \right) \left( 1+\frac{2}{p-1} \right) \right) \right)$$
$$= \frac{p-1}{p+1} \left( 1-p^{-\left(\frac{\operatorname{val}(\eta)+1}{2}\right)} \right) \left( \frac{p+1}{p-1} \right)$$
$$= 1-p^{-1/2} p^{-\frac{\operatorname{val}(\eta)}{2}}.$$

(B.7) **Definition** The weighted orbital integral  $J_{\mathrm{M}}^{\mathrm{G}}(\tilde{\eta}, \cdot)$  where  $\tilde{\eta}$  is an element of  $\mathfrak{g}'_{\mathrm{reg.ss}}(\mathbb{Q}_p)$  lifting  $\eta$  is defined is  $\eta$  is split. For such an  $\eta$  choose a square root  $\sqrt{\eta}$  of  $\eta$  in  $\mathbb{Q}_p$  and fix the lift  $\tilde{\eta}$  to be

(B.7.1) 
$$\widetilde{\eta} = \begin{pmatrix} \sqrt{\eta} \\ -\sqrt{\eta} \end{pmatrix}.$$

(B.8) Remark With respect to the choice of B as the Borel subgroup and  $G(\mathbb{Z}_p)$  as the maximal compact subgroup, the weight factor v() appearing in the definition of  $J^G_M(\tilde{\eta}, )$  is the function on  $G(\mathbb{Q}_p)$  such that v() is invariant under left translation by  $M(\mathbb{Q}_p)$  and right translation by  $G(\mathbb{Z}_p)$  and

(B.8.1) 
$$\forall u \in \mathbb{Q}_p$$
  
 $v\left(\begin{pmatrix} 1 & u \\ & 1 \end{pmatrix}\right) = \begin{cases} 0 & \text{if } u \text{ is contained in } \mathbb{Z}_p, \\ -\text{val}(u) & \text{otherwise.} \end{cases}$ 

See §12.1 of [Ko05].

**(B.9)** Lemma Let  $\eta$  be a split element of  $\mathcal{A}_{reg}(\mathbb{Q}_p)$ , then

(B.9.1) 
$$J_{\mathrm{M}}^{\mathrm{G}}(\tilde{\eta}, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)}) = \begin{cases} \frac{\mathrm{val}(\eta)}{2} - \frac{1}{p-1} + \frac{|\eta|^{1/2}}{p-1} & \text{if } \eta \text{ lies in } \mathbb{Z}_p - \{0\}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. By definition

(B.9.2)  $J_{\mathcal{M}}^{\mathcal{G}}(\widetilde{\eta}, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)})$ 

$$= |D(\eta)|^{1/2} \int_{\mathcal{M}(\mathbb{Q}_p) \setminus \mathcal{G}(\mathbb{Q}_p)} \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)}(\widetilde{\eta} \cdot \operatorname{ad}(g)) v(g) dg$$

$$(B.9.3) = |\eta|^{1/2} \int_{\mathbb{Q}_p} \int_{\mathcal{G}(\mathbb{Z}_p)} \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)} \left( \left( \begin{array}{c} \sqrt{\eta} \\ -\sqrt{\eta} \end{array} \right) \cdot \operatorname{ad} \left( \left( \begin{array}{c} 1 & u \\ 1 \end{array} \right) \right) \operatorname{ad}(k) \right) \times \\ \times v \left( \left( \begin{array}{c} 1 & u \\ 1 \end{array} \right) \right) dk du$$

$$= |\eta|^{1/2} \int_{\mathbb{Q}_p} \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)} \left( \left( \begin{array}{c} \sqrt{\eta} & 2u\sqrt{\eta} \\ -\sqrt{\eta} \end{array} \right) \right) v \left( \left( \begin{array}{c} 1 & u \\ 1 \end{array} \right) \right) du$$

$$= |\eta|^{1/2} \int_{1 < |u| \le |2\sqrt{\eta}|^{-1}} -\operatorname{val}(u) du$$

$$= |\eta|^{1/2} \sum_{i=1}^{\frac{\operatorname{val}(\eta)}{2}} i \left( \frac{1}{p^{-i}} - \frac{1}{p^{-i+1}} \right)$$

$$= |\eta|^{1/2} \left( \frac{\operatorname{val}(\eta)}{2} p^{\frac{\operatorname{val}(\eta)}{2}} - \frac{p^{\frac{\operatorname{val}(\eta)}{2}} - 1}{p - 1} \right)$$

where the equality (B.9.3) follows from the Iwasawa decomposition.

**(B.10)** Lemma Let  $\eta$  be a split element of  $\mathcal{A}_{reg}(\mathbb{Q}_p)$ , then

(B.10.1) 
$$I_{\mathrm{M}}^{\mathrm{G}}(\tilde{\eta}, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)}) = \begin{cases} \frac{\mathrm{val}(\eta)}{2} + \frac{|\eta|^{1/2}}{p-1} - \frac{2p+1}{p^2-1} & \text{if } \eta \text{ lies in } \mathbb{Z}_p - \{0\}, \\ \frac{|\eta|^{-1}}{1-p^{-2}} - \frac{|\eta|^{-1/2}}{1-p^{-1}} & \text{otherwise.} \end{cases}$$

*Proof.* By definition

(B.10.2) 
$$I_{\mathrm{M}}^{\mathrm{G}}(\widetilde{\eta}, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)})$$

$$= J_{\mathrm{M}}^{\mathrm{G}}(\widetilde{\eta}, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_{p})}) - \int_{\mathfrak{m}'(\mathbb{Q}_{p})} J_{\mathrm{M}}^{\mathrm{G}}(X, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_{p})})\psi(\mathrm{Tr}(\widetilde{\eta}^{\mathrm{T}}X)) \, \mathrm{d}X$$
$$= J_{\mathrm{M}}^{\mathrm{G}}(\widetilde{\eta}, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_{p})}) - \int_{\mathbb{Q}_{p}} J_{\mathrm{M}}^{\mathrm{G}}\left(\begin{pmatrix} x \\ & -x \end{pmatrix}, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_{p})}\right)\psi(2\sqrt{\eta}x) \, \mathrm{d}x$$

where  $\psi$  denotes the additive character of  $\mathbb{Q}_p$  involved in the definition of the Fourier transform on a *p*-adic vector space, the superscript T denotes the transpose of a 2 × 2 matrix and Tr denotes the trace of a 2 × 2 matrix.

The Fourier transform of val( ) $\mathbb{I}_{\mathbb{Z}_p}($  ) on  $\mathbb{Q}_p$  is the function such that

(B.10.3) 
$$\forall x \in \mathbb{Q}_p \qquad \left( \operatorname{val}(\ )\mathbb{I}_{\mathbb{Z}_p}(\ ) \right)^{\hat{}}(x)$$

$$= \left(\sum_{i=1}^{\infty} \mathbb{I}_{p^{i}\mathbb{Z}_{p}}(\cdot)\right)^{\hat{}}(x)$$

$$= \sum_{i=1}^{\infty} \frac{1}{p^{i}} \mathbb{I}_{p^{-i}\mathbb{Z}_{p}}(x)$$

$$= \sum_{i=\max\{1,-\operatorname{val}(x)\}}^{\infty} \frac{1}{p^{i}}$$

$$= \frac{p^{-\max\{1,-\operatorname{val}(x)\}}}{1-p^{-1}}$$

$$= \min\left\{\frac{p^{-1}}{1-p^{-1}},\frac{|x|^{-1}}{1-p^{-1}}\right\}$$

The Fourier transform of  $| | \mathbb{I}_{\mathbb{Z}_p}( )$  on  $\mathbb{Q}_p$  is the function such that

(B.10.4) 
$$\forall x \in \mathbb{Q}_{p} \qquad \left( \mid \mid \mathbb{I}_{\mathbb{Z}_{p}}( \mid) \right)^{\hat{}}(x)$$

$$= \left( \mathbb{I}_{\mathbb{Z}_{p}}( \mid) + \sum_{i=1}^{\infty} \left( \frac{1}{p^{i}} - \frac{1}{p^{i-1}} \right) \mathbb{I}_{p^{i}\mathbb{Z}_{p}}( \mid) \right)^{\hat{}}(x)$$

$$= \mathbb{I}_{\mathbb{Z}_{p}}(x) + \sum_{i=1}^{\infty} \left( \frac{1}{p^{2i}} - \frac{1}{p^{2i-1}} \right) \mathbb{I}_{p^{-i}\mathbb{Z}_{p}}(x)$$

$$= \mathbb{I}_{\mathbb{Z}_{p}}(x) + \sum_{i=\max\{1, -\operatorname{val}(x)\}}^{\infty} \left( \frac{1}{(-p)^{2i}} + \frac{1}{(-p)^{2i-1}} \right)$$

$$= \mathbb{I}_{\mathbb{Z}_{p}}(x) + \frac{(-p)^{-\left(2\max\{1, -\operatorname{val}(x)\} - 1\right)}}{1 + p^{-1}}$$

$$= \mathbb{I}_{\mathbb{Z}_{p}}(x) - \min\left\{ \frac{p^{-1}}{1 + p^{-1}}, \frac{p|x|^{-2}}{1 + p^{-1}} \right\}.$$

Hence by (B.9.1) and (B.10.2) the invariant weighted orbital integral  $I_{\mathcal{M}}^{\mathcal{G}}(\tilde{\eta}, \mathbb{I}_{\mathfrak{g}'(\mathbb{Z}_p)})$  is equal to

(B.10.5) 
$$\left( \frac{\operatorname{val}(\eta)}{2} - \frac{1}{p-1} + \frac{|\eta|^{1/2}}{p-1} \right) \mathbb{I}_{\mathbb{Z}_p - \{0\}}(\eta)$$
$$- \left( \left( \operatorname{val}(\ ) - \frac{1}{p-1} + \frac{|\ |}{p-1} \right) \mathbb{I}_{\mathbb{Z}_p}(\ ) \right)^{\wedge} (2\sqrt{\eta})$$
$$= \left( \frac{\operatorname{val}(\eta)}{2} - \frac{1}{p-1} + \frac{|\eta|^{1/2}}{p-1} \right) \mathbb{I}_{\mathbb{Z}_p - \{0\}}(\eta) + \frac{1}{p-1} \mathbb{I}_{\mathbb{Z}_p}(\eta)$$

$$(B.10.6) = \begin{cases} \frac{\operatorname{val}(\eta)}{2} + \frac{|\eta|^{1/2}}{p-1} \\ -\frac{p^{-1}}{1-p^{-1}} - \frac{1}{p-1} \left(1 - \frac{p^{-1}}{1+p^{-1}}\right) & \text{if } \eta \text{ lies in } \mathbb{Z}_p - \{0\}, \\ -\frac{|2\sqrt{\eta}|^{-1}}{1-p^{-1}} - \frac{1}{p-1} \left(-\frac{p|2\sqrt{\eta}|^{-2}}{1+p^{-1}}\right) & \text{otherwise}, \end{cases}$$
$$= \begin{cases} \frac{\operatorname{val}(\eta)}{2} + \frac{|\eta|^{1/2}}{p-1} - \frac{1}{p-1} \left(\frac{2+p^{-1}}{1+p^{-1}}\right) & \text{if } \eta \text{ lies in } \mathbb{Z}_p - \{0\}, \\ -\frac{|\eta|^{-1/2}}{1-p^{-1}} + \frac{|\eta|^{-1}}{p-1} \left(\frac{2+p^{-1}}{1+p^{-1}}\right) & \text{if } \eta \text{ lies in } \mathbb{Z}_p - \{0\}, \end{cases}$$

where the equality (B.10.6) follows from (B.10.3) and (B.10.4).

(B.11) **Remark** If  $\eta$  is a split element of  $\mathcal{A}_{reg}(\mathbb{Q}_p)$ , then  $\phi_{1,v}(\eta)$  has three components indexed by the parabolic subgroups B,  $\overline{B}$  and G, and

(B.11.1) 
$$\phi_{1,v}(\eta)_{\mathbf{B}} = \phi_{1,v}(\eta)_{\overline{\mathbf{B}}} = 1 \qquad \phi_{1,v}(\eta)_{\mathbf{G}} = -I_{\mathbf{M}}^{\mathbf{G}}(\widetilde{\eta}, \mathbb{I}_{\mathbb{Z}_p}).$$

If  $\eta$  is an elliptic element of  $\mathcal{A}_{reg}(\mathbb{Q}_p)$ , then  $\phi_{1,v}(\eta)$  has one component corresponding to the parabolic subgroup G which is equal to  $\phi_{0,v}(\eta)$ .

(B.12) Remark: Towards an invariant trace formula The simple invariant trace formula (4.2.4.1) holds only for test functions which are cuspidal at two distinct places of  $\mathbb{Q}$ . An invariant trace formula, once extended to full generality for all Schwartz functions on  $\mathfrak{g}(\mathbb{A})$ , will likely involve more intricate distributions similar to the invariant weighted orbital integrals  $I_{\mathrm{M}}^{\mathrm{G}}(X)$ , which are absent from (4.2.4.1) due to the cuspidality condition. One such identity could be established when  $\mathfrak{g}$  is gl(2) following the arguments of Langlands in [La80].

Fix a sufficiently large finite set S of places of  $\mathbb{Q}$ , let  $f_S$  be a Schwartz function on  $\mathfrak{g}(\mathbb{Q}_S)$ . It follows from the non-invariant trace formula (3.4.1.1) that

(B.12.1) 
$$\sum_{\mathfrak{o}\in\mathfrak{g}/\sim} \sum_{X\in\mathfrak{o}_{\mathrm{G},S}} a^{\mathrm{G}}(S,X) I^{\mathrm{G}}_{\mathrm{G}}(X,f_{S}) + \\ + \frac{1}{2} \sum_{\mathfrak{o}\in\mathfrak{g}/\sim} \sum_{X\in(\mathfrak{m}\cap\mathfrak{o})_{\mathrm{M},S}} a^{\mathrm{M}}(S,X) J^{\mathrm{G}}_{\mathrm{M}}(X,f_{S}) \\ = \sum_{\mathfrak{o}\in\mathfrak{g}/\sim} \sum_{X\in\mathfrak{o}_{\mathrm{G},S}} a^{\mathrm{G}}(S,X) I^{\mathrm{G}}_{\mathrm{G}}(X,f_{S}) +$$

$$+\frac{1}{2}\sum_{\mathfrak{o}\in\mathfrak{g}/\sim}\sum_{X\in(\mathfrak{m}\cap\mathfrak{o})_{\mathrm{M},S}}a^{\mathrm{M}}(S,X)J^{\mathrm{G}}_{\mathrm{M}}(X,f^{\hat{}}_{S}).$$

The weighted orbital integral  $J_{\mathrm{M}}^{\mathrm{G}}(X, f_S)$ , considered as a function in the variable X on  $\mathfrak{m}(\mathbb{Q}_S)$ , has at worst logarithmic singularity along the discriminant locus. There is a natural way to subtract from  $J_{\mathrm{M}}^{\mathrm{G}}(X, f_S)$  an invariant function in X such that the remainder extends to a continuous function on  $\mathfrak{m}(\mathbb{Q}_S)$  which is denoted by  $J_{\mathrm{M}}^{\mathrm{G},\mathrm{reg}}(X, f_S)$ . In general  $J_{\mathrm{M}}^{\mathrm{G},\mathrm{reg}}(X, f_S)$  is not smooth, as could be seen for instance from (B.9.1). However  $J_{\mathrm{M}}^{\mathrm{G},\mathrm{reg}}(X, f_S)$ , when restricted to a function on the torus  $\mathrm{M}(\mathbb{Q}_S)$ , has integrable Mellin transform, which follows from the estimates in §9 of [La80]. By a suitable adelic extension of the arguments of Ferrar in [Fe37], it could be shown that the Poisson summation formula applies to the function  $J_{\mathrm{M}}^{\mathrm{G},\mathrm{reg}}(X, f_S)$  and implies that

(B.12.2) 
$$\sum_{X \in \mathfrak{m}(\mathbb{Q})} J_{\mathrm{M}}^{\mathrm{G,reg}}(X, f_S) = \sum_{X \in \mathfrak{m}(\mathbb{Q})} J_{\mathrm{M}}^{\mathrm{G,reg}}(X, f_S)^{\mathsf{G}}$$

where the Fourier transform is taken over  $\mathfrak{m}(\mathbb{Q}_S)$ .

By definition the coefficients  $a^{M}(S, X)$  are independent of the point X, hence a suitable linear combination of (B.12.1) and (B.12.2) gives the identity

$$(B.12.3) \qquad \sum_{\mathfrak{o}\in\mathfrak{g}/\sim} \sum_{X\in\mathfrak{o}_{\mathrm{G},S}} a^{\mathrm{G}}(S,X)I_{\mathrm{G}}^{\mathrm{G}}(X,f_{S}) + \\ +\frac{1}{2}\sum_{\mathfrak{o}\in\mathfrak{g}/\sim} \sum_{X\in(\mathfrak{m}\cap\mathfrak{o})_{\mathrm{M},S}} a^{\mathrm{M}}(S,X) \left(J_{\mathrm{M}}^{\mathrm{G}}(X,f_{S}) - J_{\mathrm{M}}^{\mathrm{G},\mathrm{reg}}(X,f_{S})^{\hat{}}\right) \\ = \sum_{\mathfrak{o}\in\mathfrak{g}/\sim} \sum_{X\in\mathfrak{o}_{\mathrm{G},S}} a^{\mathrm{G}}(S,X)I_{\mathrm{G}}^{\mathrm{G}}(X,f_{S}) + \\ +\frac{1}{2}\sum_{\mathfrak{o}\in\mathfrak{g}/\sim} \sum_{X\in(\mathfrak{m}\cap\mathfrak{o})_{\mathrm{M},S}} a^{\mathrm{M}}(S,X) \left(J_{\mathrm{M}}^{\mathrm{G}}(X,f_{S}) - J_{\mathrm{M}}^{\mathrm{G},\mathrm{reg}}(X,f_{S})^{\hat{}}\right).$$

Since

(B.12.4) 
$$I_{\mathrm{G}}^{\mathrm{G}}(X, f_{S}) \quad \text{and} \quad J_{\mathrm{M}}^{\mathrm{G}}(X, f_{S}) - J_{\mathrm{M}}^{\mathrm{G}, \mathrm{reg}}(X, f_{S})^{\widehat{}}$$

are invariant under the adjoint action, the identity (B.12.3) is an invariant trace formula for gl(2) which is valid for all Schwartz functions on  $\mathfrak{g}(\mathbb{Q}_S)$ .

In order to extend this construction to a general reductive Lie algebra  $\mathfrak{g}$ , the singularities of higher rank weighted orbital integrals need to be separated, and the general invariant trace formula for each proper Levi subalgebra  $\mathfrak{m}$  needs to be extended to test functions beyond the Schwartz space  $\mathcal{S}(\mathfrak{m}(\mathbb{Q}_S))$ .

(B.13) Remark Arthur's work [Ar88b] and [Ar88c] on the invariant trace formula for reductive groups G follows an approach which is dual to the approach taken in the previous remark. Instead of working with the geometric side of the trace formula, Arthur devised the

cancellation of singularities on the spectral side of the trace formula, utilizing the underlying complex structure and in particular the powerful techniques of residue calculus. However such an alternative is unavailable in the Lie algebra case since both sides of the trace formula are geometric in nature.

(B.14) **Remark** Distributions similar to  $J_{\mathrm{M}}^{\mathrm{G,reg}}(X, \cdot)$  has been constructed by Chaudouard in [Ch02b]. He has shown that his version of the weighted orbital integral  $J_{\mathrm{M}}^{\mathrm{G,b}}(X, \cdot)$  remains bounded as X traverses  $\mathfrak{m}(\mathbb{Q}_S)$  while also enjoying many other nice properties expected of the usual weighted orbital integrals.

(B.15) **Remark** There has been some important recent work on extending the Arthur-Selberg trace formula beyond the usual space of Schwartz functions. See [FL11] and [Ma11].

## List of symbols

$G, \mathfrak{g}$	a connected reductive group defined over $\mathbb{Q}$ and its Lie algebra	page 4
v, S	a place of $\mathbb{Q}$ and a finite set of places of $\mathbb{Q}$	page 4
$\mathbb{Q}_v, \mathbb{Q}_S, \mathbb{A}$	the completions of $\mathbb{Q}$ at $v$ and $S$ and the ring of adeles of $\mathbb{Q}$	page 4
v,  S,  A	the norms on $\mathbb{Q}_v$ , $\mathbb{Q}_S$ and $\mathbb{A}$	page 4
$\mathrm{G}(\mathbb{Q}_v), \mathrm{G}(\mathbb{Q}_S),$	the topological groups of $\mathbb{Q}_v$ , $\mathbb{Q}_S$ , $\mathbb{A}$ and $\mathbb{Q}$ -valued points	page 4
$\mathrm{G}(\mathbb{A}),\mathrm{G}(\mathbb{Q})$	of G	
$P_0, M_0, A_0$	a minimal parabolic subgroup, a minimal Levi subgroup	page 4
	and a maximal split torus	
$\mathrm{P},\mathrm{M}_{\mathrm{P}},\mathrm{N}_{\mathrm{P}},\mathrm{A}_{\mathrm{P}}$	a parabolic subgroup, its Levi component, its unipotent	page 4
	radical and its split component	4
$\mathbf{P}_i, \mathbf{M}_i, \mathbf{N}_i, \mathbf{A}_i$	a parabolic subgroup, its Levi component, its unipotent	page 4
<u></u>	radical and its split component	
$P, N_P$	the parabolic subgroup opposite to P and its unipotent radical	page 4
$\mathcal{F},\mathcal{P},\mathcal{L}$	finite sets of parabolic subgroups and Levi subgroups	page 4
$X(M_P), X(M_P)^*$	the group of rational characters of $M_P$ and its dual	page 4
$\mathfrak{a}_{\mathrm{P}}, \mathfrak{a}_{\mathrm{P}}^{*}$	a Euclidean space and its dual space	page 4
$\Phi_{\rm P}, \Delta_{\rm P}, \Phi_{\rm P}^{\vee}, \Delta_{\rm P}^{\vee}$	the sets of roots, simple roots, coroots and simple coroots of P	page 5
$\Lambda^2_1  \Lambda^{2,\vee}_1  \hat{\Lambda}^2_1  \hat{\Lambda}^{2,\vee}_1$	sets of simple roots and coroots, coweights and weights	page 5
$\mathbf{a}_{1}^{2}, \mathbf{a}_{2}^{2,*}$	subquotients of $\mathbf{a}$ and $\mathbf{a}^*$	page 5
$W^{G} W(\mathbf{q}_1, \mathbf{q}_2)$	the Weyl group of $G$ and the Weyl set from $\mathbf{a}_1$ to $\mathbf{a}_2$	page 5
$\tau_0^2, \tau_1^2, \tau_2^2$	characteristic functions of positive cones in $a$	page 5
$G(\mathbb{A})^1$	the subgroup of $G(\mathbb{A})$ of elements of norm one	nage 5
K K	global and local admissible maximal compact subgroups	page 5
$H_{\rm D}$	a height function on $G(\mathbb{A})$	page 5
1112	a norgine remotion on $O(M)$	page J

$ ho_{ m P}$	the Weyl vector in $\mathfrak{a}^*$	page 6
T', T	a fixed negative point and a truncation parameter in ${\mathfrak a}$	page 6
ω	a compact subset of $N_0(\mathbb{A})M_0(\mathbb{A})^1$	page 6
$\mathfrak{S}(T',\omega), \mathfrak{S}^T(T',\omega)$	a Siegel domain and a truncated Siegel domain	page 6
$F^{\mathrm{G}}(x,T), F^{\mathrm{P}}(x,T)$	characteristic functions of a truncated Siegel domain	page 6
$\mathcal{S}, C_c^{\infty}$	spaces of Schwartz functions and compactly supported	page $7$
	smooth functions	
$\mathbb{I}_{\mathfrak{g}(\mathbb{Z}_p)}$	the characteristic function of the points in $\mathfrak{g}(\mathbb{Q}_p)$ with p-	page $7$
	adically integral coordinates	
$\langle \ , \ \rangle, \psi, \psi_v$	a nondegenerate invariant bilinear form on $\mathfrak{g}$ , a additive	page $7$
	character on $\mathbb{A}$ and its $\mathbb{Q}_v$ -component	
$\wedge,\vee$	the Fourier transform and the inverse Fourier transform	page $7$
$\sim, \mathfrak{o}$	an equivalence relation on $\mathfrak{g}(\mathbb{Q})$ and an equivalence class	page 7
$X_{\rm ss}, X_{\rm nil}$	the semisimple and nilpotent components of $X$ under the	page 7
	Jordan decomposition	
$D, D^{\mathrm{M}}$	the discriminant functions on $\mathfrak g$ and $\mathfrak m$	page 8
$\mathfrak{g}_{\mathrm{reg.ss}}$	the locus of regular semisimple elements of $\mathfrak{g}$	page 8
$G_X, G_X^0, \pi_0(G_X)$	the centralizer of $X$ in G, its identity component and its	page 8
	group of connected components	
$G_S, G_v$	the base changes of G to $\mathbb{Q}_S$ and $\mathbb{Q}_v$	page 8
$P_{S,0}, M_{S,0}, A_{S,0},$	a pair of minimal parabolic subgroups, minimal Levi sub-	page 8
$P_{v,0}, M_{v,0}, A_{v,0}$	groups and maximal split tori of $G_S$ and $G_v$	
$T_S, T_v, \mathfrak{t}_S, \mathfrak{t}_v$	maximal tori of $G_S$ and $G_v$ and the associated Cartan sub-	page 9
	algebras	
$\mathcal{T}_{\mathrm{ell}}(\mathrm{G}_S)$	the set of conjugacy classes of elliptic maximal tori of $G_S$	page 9
$W(G_S, T_S)$	the Weyl group of the pair $(G_S(\mathbb{Q}_S), T_S(\mathbb{Q}_S))$	page 9
$I_{\mathrm{G}}^{\mathrm{G}}(X, )$	the standard normalized orbital integral	page 10
$\widetilde{K}(x,f), K_{\mathfrak{o}}(x,f)$	the kernel functions of Chaudouard	page 10
$I_{\mathfrak{o}}(f)$	a variant of the standard orbital integral in the anisotropic	page 11
	case	
$K_{\mathrm{P},\mathfrak{o}}(x,f), k_{\mathfrak{o}}^{T}(,f)$	a variant of the kernel function and the truncated kernel	page 12
, ( · · · <b>·</b> ( · · · <i>·</i> )	function of Chaudouard	
$J_{\mathfrak{o}}^T, J^T$	distributions on $\mathfrak{g}(\mathbb{A})$	page 12
$\sigma_{1}^{4}()$	a combinatorial function on $\mathfrak{a}$	page 13
$\mathfrak{m}^{\mathrm{Q}}_{\mathrm{P}},\mathfrak{m}^{\mathrm{Q}}_{\mathrm{P}}(\mathbb{Q})',\mathfrak{n}^{3}_{2}$	the quotient of $\mathfrak{m}_{\mathcal{O}}$ by $\mathfrak{m}_{\mathcal{P}}$ , a subset of $\mathfrak{m}_{\mathcal{P}}^{\mathcal{Q}}(\mathbb{Q})$ and the quo-	page 13
1 / 1 ( 0) / 2	tient of $\mathfrak{n}_2$ by $\mathfrak{n}_3$	10
$\Gamma'_{\rm Po}(,T)$	the geometric gamma' function	page 14
$J_{2}^{M}$	a sum of distributions on $\mathfrak{m}(\mathbb{A})$	page 15
f <sub>P</sub>	the parabolic descent of $f$ along the parabolic subgroup P	page 15
	the Euclidean norm on $\mathfrak{a}$	page 15
$T_0$	a distinguished point in $\mathfrak{a}$	page 16
w <sub>s</sub>	a representative of the element s of the Weyl group in $G(\mathbb{O})$	page 16
$J, J_{\mathfrak{o}}$	distributions on $\mathfrak{g}(\mathbb{A})$	page 16
$v_{\rm M}(x)$	the weight factor	page 16
\ /	$\sim$	1 U -

$f_{\mathrm{P},x}, v'_{\mathrm{P}}(X)$	a noninvariant version of the parabolic descent of $f$ along P and a variant of the weight factor $v_{\rm M}(x)$	page 17
$\theta_{\mathrm{P}}$	a polynomial on $i\mathfrak{a}_{M}^{*}$	page 18
$(c_{\rm P}), c_{\rm M}$	a (G, M)-family and its associated constant	page 18
$(c_{\mathrm{P}}^{\mathrm{Q}}), (c_{\mathrm{P}})$	the (L, M)-family and the (G, L)-family associated to the $(G, M)$ -family ( $c_P$ )	page 18
$c'_{ m Q}$	the function or constant associated to the $(G, M)$ -family $(c_P)$	page 19
$\mathcal{Y}_{\mathrm{M}}, Y_{\mathrm{P}}$	a (G, M)-orthogonal set and an element of $\mathcal{Y}_{M}$	page 19
$(v_{\mathrm{P}}(\mathcal{Y}_{\mathrm{M}}))$	the (G, M)-family associated to the (G, M)-orthogonal set $\mathcal{Y}_{M}$	page 19
$\mathcal{Y}_{\mathrm{M}}(x), v_{\mathrm{M}}(x)$	a positive (G, M)-orthogonal and its associated constant which is the weight factor	page 20
$J_{\mathrm{M}}^{\mathrm{G}}(X, \cdot)$	the weighted orbital integral	page 20
$r_{\rm P}^{\rm L}(x,a)$	an auxiliary (L, M)-family	page 20
$f_{\mathrm{P},x}, v'_{\mathrm{O}}(y)$	an S-local noninvariant parabolic descent of $f$ along P and	page 20
51, <sup>20</sup> / (2(0))	its associated weight factor	10
$\mathfrak{g}_{\mathrm{nil}}, J_{\mathrm{nil}}, J_{\mathrm{nil}}^T$	the nilpotent locus and the distributions supported on $\mathfrak{g}_{nil}$	page 21
$\  \ , \mathbf{d}(T)$	a continuous seminorm on $\mathcal{S}(\mathfrak{g}(\mathbb{A}))$ and the distance from	page 21
	T to the root hyperplanes	10
$\Phi_X(x,\overline{Y})$	a partial Fourier transform of $f$	page 21
$\Gamma, \mathbf{A}_{0,T'}^{1,T}(\mathbb{R}), \delta_0^2$	a compact subset of $M_0(\mathbb{A})^1$ , a compact subset of $A_0^1(\mathbb{R})$	page 22
) 0,1'( )) '0	and the modulus function of $P_2$	1.00
$\mathcal{D}, \Phi^{\mathcal{D}}$	an invariant differential operator on $\mathfrak{q}(\mathbb{R})$ and its action on	page 22
_ , _	$\Phi$ via the Fourier transform	P - 0
N(f)	a natural number determined by a Schwartz function $f$	page 22
$\widehat{\mathbb{Z}}$	the profinite completion of $\mathbb{Z}$	nage 22
/	a seminorm on $\mathcal{S}(\mathfrak{q}(\mathbb{A}))$	nage 22
$Z_{11}$ $Z_{22}$	the components of Z on the weight space of the weight $\mu$	nage 22
$\mu, \mu$	and on $\mathbf{n}$	page 22
$\phi_{\bullet}, \Psi(a_0)$	positive Schwartz functions and the upper bounded deter- mined by the collection of $\phi$	page 22
в	a hump function on $\mathbb{O}$	nage 23
$p_v$ $y \{n_1, \dots, n_r\}$	a pull protection of $Q_v$	page 23
$\nu, (p_1, \ldots, p_l)$	on <i>u</i>	page 20
$f\epsilon$	a function obtained by truncating $f$ around $\nu$	nage 23
$J_{\nu,\nu}$	a seminorm on $S(\mathfrak{q}(\mathbb{A}))$	page 20 page 23
$T^{IIII}$	a distribution supported on $\overline{u}$	page 20 page 24
$J_{\nu}$	a distribution supported on $\nu$ a seminorm on $S(\mathfrak{q}(\mathbb{A}))$	page 24 page 24
II II II II IT	a distribution supported on $\overline{u}$	page 24
$\mathbf{m} = \mathbf{m} (\mathbf{m}) \mathbf{n} \mathbf{c}$	the set of $M(\mathbb{O}_{\alpha})$ -conjugacy classes in $\mathfrak{m}_{\alpha}(\mathbb{O})$	nage 20
$a^{M}(S \nu)$	a global coefficient that depends on $S$	page 27
$T^{G}$	a distribution supported on $\sigma_{\rm es}(\Omega_{\rm e})$	page 21
<b>п</b> н н(Фа)	a distribution supported on $\mathfrak{g}_{ni}(\mathbb{Q}_S)$	page 21
$\mathfrak{v}_{n1l,d}(\mathfrak{V}S)$	an open subset of $\mathfrak{g}_{nil}(\mathscr{C}S)$	page 20

$T_d^{\mathrm{G}}, T^{\mathrm{G},d}, T_d^{\mathrm{G},d}$	distributions constructed from $T^{\mathrm{G}}$ and $\mathfrak{g}_{\mathrm{nil},d}(\mathbb{Q}_S)$	page 28
$\Sigma$	a semisimple element in $\boldsymbol{o}$	page 30
$P_{1,\Sigma}^0, M_{1,\Sigma}^0, K_{\Sigma}$	a minimal parabolic subgroup and a minimal Levi subgroup	page 30
, ,	of $\mathcal{G}^0_{\Sigma}$ and a maximal compact subgroup of $\mathcal{G}^0_{\Sigma}(\mathbb{A})$	
$T_{\Sigma,1}$	a distinguished point in $\mathfrak{a}_1^{\mathrm{G}_{\Sigma}^{\mathrm{o}}}$	page 30
$\mathcal{F}^{\Sigma}, \mathring{\mathcal{F}}_{\mathrm{Q}}, ar{\mathcal{F}}_{\mathrm{Q}}$	finite sets of parabolic subgroups	page 30
$\pi_{\Sigma}$	a surjection from $\mathcal{F}$ to $\mathcal{F}^{\Sigma}$	page 30
${\mathcal Y}, {\mathcal Y}_{\mathrm{Q}}$	variants of a (G, M)-orthogonal set	page 30
$\Gamma'_{\mathrm{Q}}(\mathcal{Y}_{\mathrm{Q}}), \Gamma'_{\mathrm{Q}}(\mathcal{Y}_{\mathrm{Q}})$	a gamma' function and its Fourier transform	page 31
$\Phi^{\tilde{T}}_{\Omega,x}, v'_{\Omega}, \mathcal{Y}^{\tilde{T}}(k,x)$	a variant of the parabolic descent of $f$ along Q and a weight	page 31
-07	factor and a (G, M-orthogonal set involved in its definition	
$T_{\Sigma}$	a truncation parameter in $\mathfrak{a}_1$	page 32
$j_{\mathfrak{o}}^{T}(,f), J_{\mathrm{P},\mathfrak{o}}(x,f)$	variants of a truncated kernel function and its summand	page 32
Ξ	a semisimple element of $\mathfrak{m}(\mathbb{Q})$	page 40
$(v_{\mathrm{P}}(x,T)), v'_{\mathrm{Q}}$	a (G, M)-family and a weight factor	page 40
$T_{\Xi,\mathrm{M}}$	a distinguished point in $\mathfrak{a}_{M^{\underline{O}}}^{G^{\underline{O}}_{\underline{\Xi}}}$	page 40
$\Phi^T_{S,\Omega,x}$	an S-local variant of the parabolic descent of $f_S$ along Q	page 41
≡	an equivalence relation on $\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o}$	page 42
$(\mathfrak{m}(\mathbb{Q})\cap\mathfrak{o})_{\mathrm{M},S}$	the collection of $\equiv$ equivalence classes in $\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o}$	page 42
$S_{\mathfrak{o}}$	a sufficiently large finite set of places of $\mathbb{Q}$	page 42
$a^{\mathrm{M}}(S_{\mathfrak{o}}, X)$	a global coefficient that depends on $S_{\mathfrak{o}}$	page 42
$\mathring{\mathcal{L}}_{\Sigma}(M_1)$	a finite set of Levi subgroups	page 43
$a^{\overline{\mathrm{M}}}(S, X)$	a global coefficient that depends on $S$	page 45
$\Gamma, \mathcal{G}_{\Gamma}, \mathfrak{g}(\mathbb{Q})'$	a compact subset of $\mathfrak{g}(\mathbb{A})$ , a compact subset of $G(\mathbb{A})^1$ and	page 46
	a subset of $\mathfrak{g}(\mathbb{Q})$	
$f_{S,\mathrm{P}}$	the parabolic descent of $f_S$ along P	page 47
$\operatorname{Ind}_{\operatorname{M,P}}^{\operatorname{G}}(X),$	the orbit in $\mathfrak{g}(\mathbb{Q}_S)$ parabolically induced from X along P	page 47
$\operatorname{Ind}_{\operatorname{M}}^{\operatorname{G}}(X), X^{\operatorname{G}}$		
$d_{ m M}^{ m G}(\ ,\ ),\xi,s(\ ,\ )$	a bivariate function on $\mathcal{L}(M)$ , a vector in general position	page 48
	in $\mathfrak{a}$ and the partial map determined by $\xi$	
$\mathrm{G},\mathrm{M}_0,\mathbb{B}$	the general linear group $\operatorname{GL}(n, \mathbb{Q})$ and the standard choice	page $52$
	of the minimal Levi subgroup and the Borel subgroup	
$\mathcal{A}_{\mathrm{G}}, D, \mathcal{A}_{\mathrm{G,reg}}, \pi_{\mathrm{M}}$	the space of characteristic polynomials, the discriminant	page 52
, 0	function, the nonvanishing locus of $D$ and the natural map	
	from $\mathcal{A}_{\mathrm{M}}$ to $\mathcal{A}_{\mathrm{G}}$	
$ D(X_{v}) _{v}^{-1/2} dX_{v}$	a measure on $\mathcal{A}_{G}(\mathbb{Q}_{v})$	page 52
$\mathcal{A}_{\mathrm{M} \mathrm{reg}}(\mathbb{O}_v)_{\mathrm{ell}}$	the subset of $\mathcal{A}_{\mathrm{M}\mathrm{reg}}(\mathbb{Q}_{v})$ consisting of the $\mathbb{Q}_{v}$ -elliptic ele-	page 53
	ments	10
$I_{\mathrm{M}}^{\mathrm{G}}(X, ), I_{\mathrm{M}}^{\mathrm{G}}(X, )^{}$	the invariant weighted orbital integral and its Fourier trans-	page 53
IVI ( / / IVI ( / /	form	
$\mathcal{I}^{\mathrm{G}}_{\mathrm{M}}(X, \ )$	the vector-valued orbital integral	page 55
$\mathcal{I}_{\max}^{\widetilde{\mathrm{G}}}(X_v, )$	the maximal orbital integral	page 56
	~	

$\widetilde{X}_v,  \mathcal{M}[\widetilde{X}_v\text{-ell}]$	a lift of $X_v$ in $\mathfrak{g}(\mathbb{Q}_v)$ and a standard Levi subgroup whose	page 56
	Lie algebra contains $X_v$	
$egin{array}{llllllllllllllllllllllllllllllllllll$	the local Schwartz spaces	page 56
$\Lambda_0, \mathbb{I}_{\Lambda_0}, \mathcal{E}, M^{\mathrm{T}}, \mathrm{Tr}$	the standard lattice in $\mathfrak{g}(\mathbb{Q}_p)$ , its characteristic function, the standard Gaussian on $\mathfrak{g}(\mathbb{R})$ , the transpose of the matrix $M$ and the matrix trace	page 57
$\phi_{0,v},\phi_{1,v}$	the local basic functions	page 57
$Z_v, U_{\varphi_v, Z_v}, \Gamma_{\rm G}^{\rm G}(\ , \nu)$	a singular point, an open neighborhood of $Z_v$ and the Sha- like germ at a nilpotent orbit $u$	page 57
$\mathcal{S}(\Lambda(\mathbb{A}))$	the global Schwartz spaces	nora 59
$\mathcal{S}_0(\mathcal{A}_{\mathrm{G}}(\mathbb{A})),\ \mathcal{S}_1(\mathcal{A}_{\mathrm{G}}(\mathbb{A}))$	the global Schwartz spaces	page 58
$\mathcal{H}_v$	the local Harish-Chandra transform	page 59
$\mathcal{I}_{\max}^{\mathrm{G}}(X_v, )$	the function representing the Fourier transform of the max- imal orbital integral	page 59
$\mathcal{K}_{v}(.)$	the local Harish-Chandra kernel	page 59
$\hat{i}_{\mathrm{M}}^{\mathrm{G}}(.)$	Waldspurger's notation for the local Harish-Chandra kernel	page 59
$\delta_{\mathbf{X}}(\delta_{\mathbf{Y}};) \widetilde{\delta}_{\mathbf{Y}};$	the Dirac distribution at $X_{\nu}$ a $\delta$ -sequence converging to	nage 59
$o_{A_v}, (o_{A_v,i}), o_{A_v,i}$	$\delta_{\mathbf{v}}$ and a lift of $\delta_{\mathbf{v}}$ i on $\mathfrak{q}(\mathbb{Q}_{v})$	page oo
$\mathcal{H}_{S}$ , $\mathcal{H}$	the S-local and global Harish-Chandra transforms	page 60
$Z, \mathfrak{o}_{Z}, Z', \varphi_{S}(Z')$	a singular point in $\mathcal{A}_{\mathcal{C}}(\mathbb{O})$ , its fiber in $\mathfrak{q}(\mathbb{O})$ , a $\equiv$ equivalence	page 61
-,•2,-,+5(-)	class in $(\mathfrak{m}(\mathbb{Q}) \cap \mathfrak{o}_Z)_{M,S}$ and the value determined by the	Page 01
$(\mathbf{V})$	nearby regular values $\varphi_S(X)$	
$a(\Lambda)$	the global coefficient for a regular $A$	page 61
$\Lambda_{N,w},\mathbb{I}_{\Lambda_{N,w}}$	a lattice in $\mathfrak{g}(\mathbb{Q}_w)$ for a finite place $w$ and its characteristic function	page 63
$\pi_{\infty}$	the natural projection from $\mathfrak{g}(\mathbb{R})$ to $\mathcal{A}_{\mathrm{G}}(\mathbb{R})$	page 64
$T_{X_v}$	the centralizer of $X_v$ which is a torus	page 64
$\Lambda_{N,\infty}, \Lambda_{N,w},$	a lattice in $\mathfrak{g}(\mathbb{R}, a \text{ lattice in } \mathfrak{g}(\mathbb{Q}_w) \text{ for a finite place } w, \text{ and}$	page 64
$\mathbb{I}_{\widetilde{X}_{\infty}+\Lambda_{N,w}}$	the characteristic function of the translation of $\Lambda_{N,w}$ by $\widetilde{X}_{\infty}$	
$\mathbf{T}_Y, \widehat{\mathbf{T}}_Y, F, \Gamma$	the centralizer of Y which is a maximal torus, its group of algebraic characters, the splitting field of $T_Y$ and its Galois group	page 65
$\mathcal{D}$	an invariant distribution on a <i>p</i> -adic reductive Lie algebra	page 66
$\mathbb{E}^n, \mathbb{S}(\mathbb{E}^n)$	the <i>n</i> -dimensional Euclidean space and its scissors group	page 67
$\Phi, V_i, \mathbf{r}, \mathcal{R}iq(\Phi)$	a strict flag in $\mathbb{E}^n$ , the <i>i</i> th subspace of $\Phi$ , a rigging of $\Phi$ and	page 67
	the collection of equivalence classes of riggings of $\Phi$	10
$\mathcal{B}, \mathcal{B}^{\mathbf{r}}, \operatorname{sign}(\mathbf{r})$	two ordered bases of $\mathbb{E}^n$ and the sign of $\mathbf{r}$	page 67
$\Phi^{\mathbf{r}}, \partial_{\Phi^{\mathbf{r}}}(P), r_i^{\min}()$	a rigged flag, the $\Phi^{\mathbf{r}}$ -boundary of the polytope P and an	page 68
, - ( ) / ( ( )	auxiliary function	
$\operatorname{Had}_{\Phi}(P)$	the Hadwiger invariant of the polytope $P$	page 68
$(H_{\Phi}), \mathcal{F}_{s}(\Phi, i)$	a collection of real numbers and a collection of strict flags	page 68
	in $\mathbb{E}^n$	10 -

$\mathbb{S}(\mathfrak{a}_M^G), \mathcal{F}_S(\mathfrak{a}_M^G)$	the scissors group of positive (G, M)-orthogonal sets and a collection of strict flags in $\mathfrak{a}_M^G$	page 69
$\mathbb{J}_{\mathrm{M}}^{\mathrm{G}}(X, \ )$	the scissors-congruence-valued orbital integral	page 70
$\mathbf{d}\mathfrak{a}_{\mathrm{L}}^{\mathrm{L}^{l-1}}$	a differential form on $\mathfrak{a}_{M}^{L^{l-1}}$	page 71
$\mathbb{S}(\hat{\mathfrak{a}}_{\mathcal{C}(M)}^{\mathrm{G}})$	the total scissors ring of $\mathfrak{a}_{\mathrm{M}}^{\mathrm{G}}$	page 73
$\boxtimes, j, j^*$	a bilinear product on the total scissors ring and two auxil- iary maps	page 73
$ \left( \mathbb{S}(\mathfrak{a}_{\mathcal{L}(\mathbf{M})}^{\mathbf{G}}) \right)_n, $	graded components of the total scissors ring with respect to two different gradings	page 73
$\left( \mathbb{S}(\mathfrak{a}_{\mathcal{L}(\mathrm{M})}^{\mathbb{C}}) \right)_{\mathrm{L}}$		70
$q_{\mathrm{M}}^{\mathrm{L}}$	a homomorphism from $\mathbb{S}(\mathfrak{a}_{\mathcal{L}(M)}^{G})$ to $\mathbb{S}(\mathfrak{a}_{\mathcal{L}(L)}^{G})$	page 73
$\mathcal{J}_{\mathrm{M}}^{\mathrm{G}}(X, )$	the total orbital integrohedron	page 74
v, p,	an odd rational prime and the $p$ -adic absolute value	page 77
G, M, B	the group $GL(2)$ and the standard choice of the minimal	page $77$
	Levi subgroup and the Borel subgroup	
$\mathfrak{g},\mathfrak{g}',\mathfrak{m}',\mathfrak{z}$	the Lie algebras $gl(2), sl(2), \mathfrak{m} \cap \mathfrak{g}'$ and the center of $\mathfrak{g}$	page 77
$\mathcal{A}, \mathbb{L}, D, \mathcal{A}_{\mathrm{reg}}, \mathbb{Q}_p^{\times}$	the affine quotient of $\mathfrak{g}'$ by G, the affine line, the discrim-	page 77
U F	inant function on $\mathcal{A}$ , the nonvanishing locus of $D$ and the	
	group of units of $\mathbb{Q}_n$	
$\eta, \mathfrak{g}'_n, \mathrm{d}_n X$	a point of $\mathcal{A}_{reg}$ , the fiber of $\eta$ in $\mathfrak{g}'$ and a differential form	page 77
	on $\mathfrak{g}'_n$	- 0
$\operatorname{val}(\eta)$	the <i>p</i> -adic valuation of $\eta$	page 80
$\widetilde{\eta}$	a lift of a split $\eta$ in $\mathfrak{g}'_{reg,ss}(\mathbb{Q}_p)$	page 82
$J_{\mathrm{M}}^{\mathrm{G,reg}}(X, )$	a regularized version of the weighted orbital integral	page 86
IVI ( / /	$J_{\rm M}^{\rm G}(\tilde{X}, \cdot)$ for the Lie algebra gl(2)	
$J_{\mathrm{M}}^{\mathrm{G},b}(X, \ )$	a variant of the weighted orbital integral $J_{M}^{G}(X, \cdot)$ con-	page 87
111	structed by Chaudouard	

## References

- [1859] B. Riemann, Üeber die Anzahl der Primzahlen unter einer gegebenen Grösse, Monatsberichte der Berliner Akademie, 1859, 671-680.
- [Ar76] J. Arthur, The characters of discrete series as orbital integrals, Inv. Math. 32 (1976), 205-261.
- [Ar78] J. Arthur, A trace formula for reductive groups I: terms associated to classes in  $G(\mathbb{Q})$ , Duke Math. J. 245 (1978), 911-952.
- [Ar81] J. Arthur, The trace formula in invariant form, Ann. of Math. 114 (1981), 1-74.
- [Ar82] J. Arthur, On a family of distributions obtained from Eisenstein series I: Application of the Paley-Wiener theorem, Amer. J. Math. 104 (1982), 1243-1288.
- [Ar85] J. Arthur, A measure on the unipotent variety, Canad. J. Math. 37 (1985), 1237-1274.

- [Ar86] J. Arthur, On a family of distributions obtained from orbits, Canad. J. Math. 38 (1986), 179-214.
- [Ar88a] J. Arthur, The local behaviour of weighted orbital integrals, Duke Math. J. 56 (1988), 223-293.
- [Ar88b] J. Arthur, The invariant trace formula I. Local theory, J. Amer. Math. Soc. 1 (1988), 323-383.
- [Ar88c] J. Arthur, The invariant trace formula II. Global theory, J. Amer. Math. Soc. 1 (1988), 501-554.
- [Bo81] W. Borho, Uber Schichten halbeinfacher Lie-Algebren, Inv. Math. 65 (1981), 283-371.
- [BK00] A. Braverman and D. Kazhdan,  $\gamma$ -functions of representations and lifting (with an appendix by V. Vologodsky), Geom. Funct. Anal. 2000, Special Volume, Part I, 237-278.
- [Bo69] A. Borel, *Introduction aux groupes arithmétiques*, Actualitées Sci. Ind. No. 1341, Hermann, Paris, 1969.
- [Ch02a] P.-H. Chaudouard, La formule des traces pour les algèbres de Lie, Math. Ann. 322 (2002), 347-382.
- [Ch02b] P.-H. Chaudouard, Intégrales orbitales pondérées sur les algèbres de Lie : le cas p-adique Canad. J. Math. 54 (2002), 263-302.
- [Ev98] U. Everling, An example of Fourier transforms of orbital integrals and their endoscopic transfer, New York J. Math. 4 (1998), 17-29.
- [Fe37] W. L. Ferrar, Summation formulae and their relation to Dirichlet's series II, Compos. Math. 4 (1937), 394-405.
- [FL11] T. Finis and E. Lapid, On the Arthur-Selberg trace formula for GL(2), Groups Geom. Dyn. 5 (2011), No. 2, Special issue on the occasion of Fritz Grunewald's 60th birthday, 367-391.
- [GJ72] R. Godement and H. Jacquet, Zeta functions of simple algebras, Lecture Notes in Math., Vol. 260, Springer, New York, 1972.
- [Ho13] W. Hoffmann, Induced conjugacy classes, prehomogeneous varieties, and canonical parabolic subgroups, Preprint, 2013, arXiv:1206.3068.
- [HW13] W. Hoffmann and S. Wakatsuki, On the geometric side of the Arthur trace formula for the symplectic group of rank 2, Preprint, 2013, arXiv:1310.0541.
- [Ko05] R. Kottwitz, Harmonic analysis on reductive p-adic groups and Lie algebras, in Harmonic analysis, the trace formula, and Shimura varieties, Clay Math. Proc., Vol. 4, Amer. Math. Soc., Providence, RI, 2005, 391-522.

- [La13] L. Lafforgue, Noyaux dutransfert automorphe deLanglands formules dePoisson 2013,etnonlinéaires, Preprint, http://www.ihes.fr/~lafforgue/publications.html.
- [La80] R. Langlands, Base Change for GL(2), Annals of Math. Studies 96, Princeton U. P. 1980.
- formula for GL(2)[Ma11] J. Matz, Arthur's traceand GL(3)and nonsupported functions, Dissertation, Düsseldorf, 2011,compactly testhttp://www.math.uni-bonn.de/people/matz/.
- [Mo93a] R. Morelli, Translation scissors congruence, Adv. Math. 100 (1993), 1-27.
- [Mo93b] R. Morelli, The K theory of a toric variety, Adv. Math. 100 (1993), 154-182.
- [Ng12] B. C. Ngô, On a certain sum of automorphic L-functions, Preprint, 2012, http://www.math.uchicago.edu/~ngo/.
- [On63] T. Ono, On the Tamagawa number of algebraic tori, Ann. of Math. 78 (1963), 47-73.
- [Sa12] Y. Sakellaridis, Spherical varieties and integral representations of L-functions, Algebra Number Theory, Vol. 6 (2012), No. 4, 611-667.
- [Ta50] J. Tate, Fourier analysis in number fields, and Hecke's zeta-functions, in Algebraic Number Theory, Academic Press, New York, 1967, 305-347.
- [Va77] V. S. Varadarajan, Harmonic analysis on real reductive groups, Part one: Invariant analysis on a real reductive Lie algebra, in Harmonic analysis on real reductive groups, Lecture Notes in Math., Vol. 576, Springer, New York, 1977, 1-174.
- [Wa95] J.-L. Waldspurger, Une formule des traces locale pour les algèbres de Lie p-adiques,
   J. Reine Angew. Math. 465 (1995), 41-99.