

Inflation in scale-invariant theories of gravity

Massimiliano Rinaldi^{*}, Guido Cognola[†], Luciano Vanzo[‡], and Sergio Zerbini[§]

Dipartimento di Fisica, Università di Trento

via Sommarive 14, 38123 Trento, Italia

and

TIFPA (INFN)

via Sommarive 14, 38123 Trento, Italia

(Dated: November 5, 2018)

Abstract

Thanks to the Planck Collaboration, we now know the scalar spectral index of primordial fluctuations with unprecedented precision. In addition, the joint analysis of the data from Planck, BICEP2, and KEK has further constrained the value of the tensor-to-scalar ratio r so that chaotic inflationary scenarios seem to be disfavoured. Inspired by these results, we look for a model that yields a value of r that is larger than the one predicted by the Starobinsky model but is still within the new constraints. We show that purely quadratic, renormalizable, and scale-invariant gravity, implemented by loop-corrections, satisfies these requirements.

^{*} e-mail: massimiliano.rinaldi@unitn.it

[†] e-mail: cognola@science.unitn.it

[‡] e-mail: vanzo@science.unitn.it

[§] e-mail: zerbini@science.unitn.it

INTRODUCTION

The claim of the BICEP2 collaboration [1] that the tensor-to-scalar ratio has the surprisingly large value $r \simeq 0.2$, which points to a robust production of gravitational waves during inflation, has polarized the attention of the physics community. The situation has become less clear when some serious criticisms to the BICEP2 analysis appeared in the literature (see for example [2]). Very recently, the joint analysis of data from Planck, KEK, and BICEP2 has finally settled the issue, fixing a new upper value for the tensor-to-scalar ratio at $r < 0.9$ [3, 4]. In conclusion, although this bound is much lower than initially claimed in [1], we can say that it is still significant enough to allow for alternative theories to the Starobinsky Model (SM).

As showed in [5], cosmic inflation can be studied in models of gravity where the Lagrangian $\sqrt{g}R$ is replaced by a suitable function $\sqrt{g}f(R)$ (see the general analysis in [6]). Along these lines, in [7] we showed that the functional form of the inflationary $f(R)$ can be determined if the scalar spectral index n_s and the tensor-to-scalar ratio r are known with sufficient accuracy. In particular, we have shown that the slow-roll conditions imply that $f(R) \sim R^{2-\delta} \sim R^2(1 - \delta \log R + \dots)$, where δ is a small and weakly time-dependent parameter, in line with the results discussed in [8]. Since it is known that R^2 gravity leads to a scale-invariant spectrum of scalar perturbations, we concluded that small deviations from $f(R) = R^2$ are crucial. Such deviations can appear in various form. For instance, one can add other gravitational quadratic terms and the matter fields of the standard model in such a way that the Planck scale is dynamically generated, as in [9] and in earlier work reviewed in [10]. Alternatively, one can add quantum one-loop corrections to R^2 and resort again to the mechanism advocated by Adler in [10]. However, in our approach, the R^2 term is not interpreted as a loop correction to something else, but as the classical (inflationary) Lagrangian. This choice is also based on theoretical motivations. For example, the addition of the linear Einstein term to R^2 leads to a tachyonic degree of freedom at tree-level and in flat space [11]. This problem persists in the presence of terms like $R_{\mu\nu}^2$ which, in addition, lead to multi-field inflation in the Einstein frame description of the theory. The same occurs for terms such as $R\Box^n R$ [12]. Finally, R^2 has a de Sitter vacuum solution with arbitrary cosmological constant.

In principle, the fundamental Lagrangian should also contain the non-minimally coupled

Higgs field. On the other hand, one can assume that, at the onset of inflation, the standard model matter is in the symmetric vacuum, so that the (vacuum) expectation value of the Higgs field is vanishing. Indeed, in de Sitter space the Higgs potential looks like

$$V = -\frac{m^2}{2}\mathcal{H}^\dagger\mathcal{H} + \frac{\lambda}{4}(\mathcal{H}^\dagger\mathcal{H})^2 + \xi R_{\text{ds}}\mathcal{H}^\dagger\mathcal{H}, \quad (1)$$

where $m \sim 125$ Gev is the standard model Higgs mass and R_{ds} is the de Sitter curvature. As long as $\xi R_{\text{ds}} \geq m^2/2$, the symmetry is unbroken and $\langle\mathcal{H}\rangle = 0$. Then, the Einstein linear term in R vanishes, leaving the responsibility for inflation to the higher-order terms.

Motivated by these considerations, we study an effective Lagrangians of the form R^2 + loop corrections. We first show that a finite number of loop corrections leads to spectral indices incompatibles with data. Therefore, we suggest a phenomenological form which mimics a possible resummed Lagrangian for which the values of r and n_s are consistent with the Planck data. In particular, r is significantly larger than the one of the Starobinsky model (SM) but still within the experimental bounds.

Our approach is based on the effective Lagrangian formalism and is driven by phenomenological considerations. As discussed below, the starting point of our work is the calculation of one-loop corrections to quadratic gravity by expanding around a (Euclidean) de Sitter space, as we believe that this is more appropriate for an effective theory of gravity that describes inflation. Instead, the traditional approach to renormalizable theories of gravity, based on the pioneering work of Stelle [11], treats the quantization of fluctuations around flat space (see e.g [13]). Whether or not these methods are equivalent is an open question.

THE SCALE INVARIANT R^2 MODEL AND ITS DEFORMATIONS

To begin with, let us recall the most general scale-invariant Jordan frame action containing the square of the Ricci scalar, the Weyl invariant, and the Higgs doublet \mathcal{H} non-minimally coupled to gravity:

$$S_J = \int d^4x \sqrt{g} \left[bR^2 + aW + \xi R\mathcal{H}^2 - (\partial\mathcal{H})^2 - \frac{\lambda}{4!}\mathcal{H}^4 + \dots \right]. \quad (2)$$

The dots stand for the other quadratic invariants of the metric and scale invariant operators of the standard model. Here, the parameters b , ξ , and λ are all dimensionless. This action has been thoroughly investigated in the past, see e.g. [14]. Recently, it was reconsidered

within a much more large physical context in [9], where it was shown that it leads to an inflationary model consistent with observations, provided one adds a new scalar field degree of freedom and takes in account the running of b , ξ , and λ . In addition, this model is particularly attractive as it is believed to be renormalizable and asymptotically free [11, 15–17], although ghosts are in general present (for a review, see [18]).

In this paper, we take a different look at this action and we show that inflation may be realized by just the first term of eq. (2), implemented by suitable loop corrections. As explained in the introduction, in the symmetric state, the operators related to the Higgs field vanish until the curvature drops below a certain critical value. In the following, we adopt the point of view that, if by “quantization” we understand the process of functional integration over a set of fundamental, non-composite fields, then the effective action has a loop-expansion in the Nambu sense: take a parameter g , replace the action $I \rightarrow g^{-1}I$ and expand the functional integral in powers of g . Then, one sees that a L loop-connected graph has a coefficient g^{L-1} . Hence, the one-loop term is given by a suitable ratio of functional determinants. The important point is that we integrate the Euclidean action over the four-sphere, which corresponds to de Sitter space, rather than over flat Euclidean space. In fact, the theory R^2 has only a one-parameter family of non-trivial homogeneous and isotropic solutions, among which we find de Sitter space and the radiation-dominated Universe. Together with the latter, the other solutions do not seem to have any Euclidean counterpart and presumably lead to an ill-defined functional integral (see [19] for other remarkable properties of R^2).

In Ref. [7], and on the grounds of the results presented in [20], we showed that the one-loop corrections to R^2 leads to the effective Lagrangian

$$f_{\text{eff}}(R) = R^2 \left[1 - \gamma \ln \left(\frac{R^2}{\mu^2} \right) \right], \quad (3)$$

where γ is a small positive parameter and μ is a constant that fixes the scale of the corrections (see also [21], and, for the asymptotic safety approach, [22]). A similar expression was obtained in [23] by a conformal transformation from an Einstein frame action containing a quadratic potential and a cosmological constant. However, as we will show shortly, this Lagrangian inevitably leads to a blue-tilted scalar spectrum and must be discarded. Motivated by this no-go result, we softly break the scale invariance with a deformation of the classical R^2 Lagrangian that mimics the resummation of higher-loop logarithmic corrections

(in absence of gravity, see, for example [24]), namely

$$f_{\text{eff}}(R) = \frac{R^2}{\left[1 + \gamma \ln\left(\frac{R^2}{\mu^2}\right)\right]}. \quad (4)$$

The scale μ , which has the dimension of a squared mass, is necessary to make the logarithm dimensionless. Of course there is now a troublesome pole at $R_{\text{div}} = \mu \exp(-1/2\gamma)$, but with the reasonable assumption that $\gamma \ll 1$ this is in the region of such a small curvature that the effects induced by the Higgs and other standard model fields cannot be ignored and a different physics should sets in. In any case, we will show below that inflation occurs entirely at $R > R_{\text{div}}$, independently of the value of γ . So, even if we are not really able to fully justify this choice of the Lagrangian (some arguments in favor of this choice will be given in the next Section) we feel authorized in using it during the inflationary phase. However, we may anticipate that the surprising feature of (4) is that it yields an inflationary phase such that the spectral index, its running, and the tensor-to-scalar ratio depend exclusively on the number of e-folds. The only constraint that relates γ and μ comes from amplitude of the scalar power spectrum.

INFLATION IN $f(R)$ THEORIES

In order to obtain the inflationary observables, we introduce a simple and transparent formalism that is valid for all $f(R)$ theories. Let us consider the generic action in Jordan frame (for reviews on $f(R)$ gravity see e.g. [25–27])

$$S_J = \int d^4x \sqrt{|g|} f(R). \quad (5)$$

Our goal is to express the usual inflationary observables in both Einstein and Jordan frame in a simple and universal form. The only vacuum equation of motion for a homogeneous and isotropic Universe with metric $ds^2 = -dt^2 + a^2 d\vec{x}^2$ is

$$3XH^2 = \frac{1}{2}(XR - f) - 3H\dot{X}, \quad (6)$$

where the dot represents a derivative with respect to the (Jordan frame) cosmic time t , $H = a^{-1}\dot{a}$ is the Hubble function, $R \equiv 6(2H^2 + \dot{H})$, and $X \equiv df(R)/dR$. The conformal transformation $\tilde{g}_{\mu\nu} = Xg_{\mu\nu}$ brings the action (5) into the canonical form in Einstein frame

$$S_E = \int d^4x \sqrt{|\tilde{g}|} \left[\frac{M^2}{2} \tilde{R} - \frac{1}{2} (\tilde{\partial}\tilde{\phi})^2 - V(\tilde{\phi}) \right], \quad (7)$$

where

$$V(\tilde{\phi}) = \frac{M^2}{2} \left(\frac{XR - f(R)}{X^2} \right), \quad (8)$$

and X and $\tilde{\phi}$ are related by

$$\tilde{\phi} = \sqrt{\frac{3}{2}} M \ln(X). \quad (9)$$

Usually, the parameter M is identified with the Planck mass m_p under the hypothesis that the action (7) describes also the low-energy limit of the theory. However, since we will deal with the scale-invariant Lagrangian (4), this identification is not strictly speaking justified. Nevertheless, for now we keep a conservative point of view by setting $M = m_p$ and we will comment below on alternative choices. If $X(R)$ is positive definite and invertible, we can always write a derivative with respect to $\tilde{\phi}$ in terms of a derivative with respect to R . In particular, we can express the slow-roll parameters as (the prime indicates a functional derivative with respect to R)

$$\begin{aligned} \epsilon &= \frac{M^2}{2} \left(\frac{d \ln(V)}{d \tilde{\phi}} \right)^2 = \frac{(XR - 2f)^2}{3(XR - f)^2}, \\ \eta &= \frac{M^2}{V} \frac{d^2 V}{d \tilde{\phi}^2} = \frac{2(XR - 4f)}{3(XR - f)} + \frac{2X^2}{3(XR - f)X'}, \\ \xi^2 &= \frac{M^2}{V^2} \frac{dV}{d \tilde{\phi}} \frac{d^3 V}{d \tilde{\phi}^3} = \frac{4(XR - 2f)(X^3 X'' + X'^3 X R - 8X'^3 f + 3X^2 X'^2)}{9X'^3 (XR - f)^2}, \end{aligned} \quad (10)$$

from which we construct the spectral index, its running, and the tensor-to-scalar ratio defined as

$$n_s = 1 - 6\epsilon + 2\eta, \quad r = 16\epsilon, \quad \frac{dn_s}{d \ln k} = 16\epsilon\eta - 24\eta^2 - 2\xi^2. \quad (11)$$

With the help of the definition (9), we can also define the number of e-fold (in Einstein frame) as a function of $\tilde{\phi}$ or R according to

$$\tilde{N}(\tilde{\phi}) = \frac{1}{M^2} \int V \left(\frac{dV}{d \tilde{\phi}} \right)^{-1} d \tilde{\phi} = \frac{3}{2} \int \frac{V}{V'} \frac{X'^2}{X^2} dR. \quad (12)$$

We stress that these formulae are valid for any $f(R)$ theory and hold whenever $X(R)$ is positive definite and invertible.

With the help of these formulae we may try to justify Eq. (4). Suppose that the full Lagrangian is a deformation of the SM form induced by n loop corrections, so it can be

written as

$$f(R) = \xi \mathcal{H}R + a_2 R^2 g(z), \quad z = \ln \left(\frac{R^2}{\mu^2} \right), \quad (13)$$

for some constant a_2 and ξ . Here we stress again that the role of inflaton is played by the gravitational scalaron, and the \mathcal{H} may play a role only at the end of inflation. We can reasonably assume that the function $g(z)$ can be written, at least in the regime of interest, as $g(z) = 1 + \gamma_1 z + \gamma_2 z^2 + \dots + \gamma_n z^n$ with $\gamma_1 \gg \gamma_2 \gg \dots \gg \gamma_n$. By using the first two equations of (10) and the expressions of n_s and r we find that (for small r and $\gamma_1 \ll 1$) $r \simeq 3(1 - 2\gamma_1)(n_s - 1)^2$. This shows that the corrections to the predictions of the SM (i.e. $r \simeq 3(n_s - 1)^2$) are relatively small but in principle measurable, as already noticed in several papers (see e.g. [8]).

Let us now consider eq. (13) with $\xi = 0$. In this case, we find that $n_s \simeq 1 + r/8$, which is patently in contrast with observations. However, if we assume that all the logarithmic corrections give a contribution of the form (4), then $r \simeq 8(1 - n_s)/3$, which lies within the experimental bounds and represents a class of models disjoint from the SM class. For $n_s \simeq 0.968$ [3], this relation yields $r = 0.085$.

As discussed above, eq. (4) is valid only in the range $R > R_{\text{div}}$, where it has a global minimum at $R_{\text{min}} = \mu \exp[(\gamma - 1)/2\gamma]$ so both $f'(R)$ and $f''(R)$ are positive for $R > R_{\text{min}}$, fulfilling the conditions of [6]. The numerical analysis of the equations of motion reveals that R_{div} is an attractor therefore the system, during inflation, evolves from a value $R \gg \mu$ towards R_{div} . However, before reaching this point, we expect some reheating mechanism (for example by the symmetry breaking explained above) after which the expression of (4) is no longer valid.

INFLATIONARY OBSERVABLES

To check the consistency of our model, we must verify that inflation last long enough. With the help of eqs. (10), we find that

$$\epsilon = \frac{4\gamma^2}{3(1 + \gamma z - 2\gamma)^2}, \quad \eta = -\frac{16\gamma^3}{3(1 + \gamma z - 2\gamma)[(1 + \gamma z)^2 - 3\gamma(1 + \gamma z) + 4\gamma^2]}. \quad (14)$$

By combining these expressions with the first two eqs. of (11) we find that

$$n_s - 1 = -\frac{r(\pm 9r + 28\sqrt{3r} \pm 96)}{8(\pm 3r + 4\sqrt{3r} \pm 32)} \simeq -\frac{3r}{8} \mp \mathcal{O}(r^{3/2}), \quad (15)$$

where the second equality holds for small r . This confirms, to leading order, the relation $r \simeq 8(1 - n_s)/3$ found above. It is remarkable that this relation does not depend on γ and μ . To fix a point in the (n_s, r) plane, we must compute the number of e-folds before the end of inflation, which typically occurs when $\epsilon(z, \gamma) = 1$. This equation has two solutions, of which only one is at a value of $R > R_{\min}$, corresponding to

$$z_{\text{end}} = -\frac{1}{\gamma} + 1 + \frac{\sqrt{3}}{3}. \quad (16)$$

We note in passing that this result implies that inflation occurs entirely for $R > R_{\min} > R_{\text{div}}$, namely not only in the range of validity of our theory but also in the stability range, where X and dX/dR are positive definite [6]. By integrating eq. (12) we find that

$$\tilde{N}(z) = \frac{3z^2}{16} - \frac{3z}{2} + \frac{3z}{8\gamma} + \frac{3}{4} \ln \left[\frac{(1 + \gamma z)^4}{(1 + \gamma z - \gamma)} \right]. \quad (17)$$

At a given number \tilde{N}^* of e-folds before the end of inflation, the corresponding value of z_{ex} is then implicitly determined by

$$\tilde{N}(z_{\text{ex}}) - \tilde{N}(z_{\text{end}}) = \tilde{N}^*. \quad (18)$$

The spectral index, its running, and the tensor-to-scalar ratio are finally obtained numerically by inserting z_{ex} in the expressions (14) and (11). One surprising characteristic is that the results do not depend on γ but only on \tilde{N}^* . In table I, we report the numerical values of n_s , r , and $dn_s/d \ln k$ for a range of \tilde{N}^*

\tilde{N}^*	n_s	r	$dn_s/d \ln k$
40	0.9661	0.084	-0.0008
45	0.9697	0.075	-0.0006
50	0.9727	0.068	-0.0005

TABLE I: Values of n_s , its running, and r corresponding to three values of the number of e-folds before the end of inflation.

We note that, in order to fit the experimental value $n_s = 0.968 \pm 0.006$ [3] we need to take a number of e-fold which is lower than the standard interval $50 < \tilde{N}^* < 60$. However, it is known that for non-polynomial (in Einstein frame) models of inflation such a range can be different, depending on the details of the reheating mechanism [29]. We also note that the

tensor-to-scalar ratio is about ten times larger than the one predicted by the SM but still lies within the experimental Planck bound $r \lesssim 0.09$ [35]. The running of the spectral index is negative, small and fully compatible with the Planck result $dn_s/d \ln k = -0.003 \pm 0.007$ [3]. Although n_s , $dn_s/d \ln k$, and r are independent of γ and μ , the amplitude A_s of the power spectrum of the curvature perturbations is not. In our model (with the assumption that M is the same as the Planck mass) we find the expression

$$A_s = \frac{V}{24\pi^2 M^4 \epsilon} = \frac{\mu(1 + \gamma z)^2(1 + \gamma z - 2\gamma)^3}{512 M^2 \pi^2 \gamma^2 (1 + \gamma z - \gamma)^2}, \quad (19)$$

which must be evaluated at the horizon exit $z = z_{\text{ex}}$. By assuming the typical value $A_s \simeq 2 \times 10^{-9}$, we find that $\sqrt{\mu}/M \simeq 5 \times 10^{-5}/\sqrt{\gamma}$. The parameter γ is assumed to be a small number, and only when $\gamma \sim 10^{-9}$ the mass scale $\sqrt{\mu}$ approaches the value of the Planck mass M . With the help of eqs. (9) and (18), we can write $\tilde{\phi}$ at a generic \tilde{N}^* as

$$\frac{\tilde{\phi}^*}{M} = F(\tilde{N}^*) - \frac{\sqrt{6}}{2} \left(\ln \gamma + \frac{1}{2\gamma} \right), \quad (20)$$

where the first term is a complicate algebraic function of \tilde{N}^* only. If, for example, we require that the value of $\tilde{\phi}$ at horizon exit is of the order of $5M$, as in the SM [5], we find that, for $\tilde{N}^* = 40$, $\gamma \simeq 0.087$ in line with the requirement that $\gamma \ll 1$. In Jordan frame, this value corresponds to $R_{\text{ex}} \simeq 3 \times 10^{-8} M^2$ and to a Hubble parameter that can be estimated to be of the order of $H_{\text{ex}} = \sqrt{R_{\text{ex}}/12} \simeq \sqrt{2\mu} = 5 \times 10^{-6} M$, similarly to the SM. This shows that our model can be compared to the SM in terms of energy scales and spectral index. In the recent paper [30], similar results concerning n_s and r have been obtained without including the Hilbert-Einstein term to the action, but taking into account quantum corrections to the potential, coming from to the RG-equations related to scalar electrodynamics.

CONCLUSIONS

In this letter we entertained the idea that the inflationary Universe can be entirely described by a purely quadratic gravitational effective theory, provided loop corrections are taken in account. We find that the spectral index of scalar perturbations matches the Planck data, while the scalar-to-tensor ratio is about ten times larger than the one of the Starobinsky model. Remarkably, these predictions are independent of the parameters of the theory. It is worth noticing that our model is not, in principle, affected by a transplanckian problem.

It is known, from the Lyth bound [34], that in single-field inflation, in order to have a non-negligible value of r , one needs a planckian excursion of the inflaton field, according to $\Delta\phi/m_p \sim \int_0^{\tilde{N}^*} d\tilde{N} \sqrt{\tilde{r}}$. If future data analysis will confirm that r is of the order of $1/10$, it will be difficult to claim that quantum gravitational effects should not be taken in account in these models. In our case, the situation is different as the scale M that appears in (9) does not necessarily match the Planck mass. Thus, in general, the expression $\Delta\phi/m_p$ should contain the ratio M/m_p that can relax the Lyth bound. This is consistent with the fact that the classical part of our model is scale-invariant and physical scales appear only after inflation, when standard model particles emerge.

-
- [1] P. A. R. Ade *et al.* [BICEP2 Collaboration], Phys. Rev. Lett. **112** (2014) 241101.
 - [2] R. Flauger, J. C. Hill and D. N. Spergel, JCAP **1408** (2014) 039.
 - [3] P. A. R. Ade *et al.* [Planck Collaboration], “Planck 2015 results. XIII. Cosmological parameters,” arXiv:1502.01589 [astro-ph.CO].
 - [4] P. A. R. Ade *et al.* [BICEP2 and Planck Collaborations], “A Joint Analysis of BICEP2/Keck Array and Planck Data,” [arXiv:1502.00612 [astro-ph.CO]].
 - [5] A. A. Starobinsky, Phys. Lett. B **91** (1980) 99.
 - [6] S. A. Appleby, R. A. Battye and A. A. Starobinsky, JCAP **1006**, (2010) 005.
 - [7] M. Rinaldi, G. Cognola, L. Vanzo and S. Zerbini, JCAP **1408** (2014) 015.
 - [8] I. Ben-Dayan, S. Jing, M. Torabian, A. Westphal and L. Zarate, JCAP **1409** (2014) 005; A. Codello, J. Joergensen, F. Sannino and O. Svendsen, “Marginally Deformed Starobinsky Gravity,” arXiv:1404.3558 [hep-ph]; J. Joergensen, F. Sannino and O. Svendsen, Phys. Rev. D **90** (2014) 043509.
 - [9] A. Salvio and A. Strumia, JHEP **1406** (2014) 080
 - [10] S. L. Adler, Rev. Mod. Phys. **54**, 729 (1982); [Erratum-ibid. **55**, 837 (1983)].
 - [11] K. S. Stelle, Phys. Rev. D **16** (1977) 953.
 - [12] D. Wands, Class. Quant. Grav. **11**, 269 (1994).
 - [13] F. Brisce, A. Marcian, L. Modesto and E. N. Saridakis, Phys. Rev. D **87** (2013) 8, 083507; T. Biswas, E. Gerwick, T. Koivisto and A. Mazumdar, Phys. Rev. Lett. **108** (2012) 031101; L. Modesto, Phys. Rev. D **86** (2012) 044005.

- [14] I. L. Buchbinder, Sov. Phys. J. **29** (1986) 220.
- [15] E. Tomboulis, Phys. Lett. B **70** (1977) 361; E. Tomboulis, Phys. Lett. B **97** (1980) 77.
- [16] E. S. Fradkin and A. A. Tseytlin, Nucl. Phys. B **201** (1982) 469; E. S. Fradkin and A. A. Tseytlin, Phys. Lett. B **104** (1981) 377.
- [17] I. G. Avramidi and A. O. Barvinsky, Phys. Lett. B **159** (1985) 269.
- [18] I. L. Buchbinder, S. D. Odintsov, and I. L. Shapiro, “Effective Action in Quantum Gravity”, IOP, Bristol (1992).
- [19] H. A. Buchdahl, J. Phys. A - Math. Gen. **11** 871 (1978); H. A. Buchdahl, Int. Journ. Theor. Phys. **17**, 149 (1978).
- [20] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov and S. Zerbini, JCAP **0502** (2005) 010; G. Cognola, E. Elizalde and S. Zerbini, Phys. Rev. D **87** (2013) 4, 044027; K. Bamba, G. Cognola, S. D. Odintsov and S. Zerbini, Phys. Rev. D **90** (2014) 023525.
- [21] L. H. Ford, Phys. Rev. D **35** (1987) 2339.
- [22] A. Bonanno, Phys. Rev. D **85** (2012) 081503.
- [23] S. V. Ketov and N. Watanabe, Phys. Lett. B **741** (2015) 242.
- [24] F. T. Brandt, F. A. Chishtie and D. G. C. McKeon, “The Effective Potential in Non-Conformal Gauge Theories,” arXiv:1409.2820 [hep-th].
- [25] S. Nojiri and S. D. Odintsov, Phys. Rept. **505** (2011) 59.
- [26] T. P. Sotiriou and V. Faraoni, Rev. Mod. Phys. **82** (2010) 451.
- [27] A. De Felice and S. Tsujikawa, Living Rev. Rel. **13** (2010) 3.
- [28] C. Kounnas, D. Lüst and N. Toumbas, arXiv:1409.7076 [hep-th].
- [29] A. R. Liddle and S. M. Leach, Phys. Rev. D **68** (2003) 103503.
- [30] E. Elizalde, S. D. Odintsov, E. O. Pozdeeva and S. Y. Vernov, Phys. Rev. D **90**, 084001 (2014).
- [31] M. B. Mijic, M. S. Morris and W. M. Suen, Phys. Rev. D **34** (1986) 2934.
- [32] L. H. Ford, Phys. Rev. D **35** (1987) 2955.
- [33] G. N. Felder, L. Kofman and A. D. Linde, Phys. Rev. D **60** (1999) 103505; G. N. Felder, L. Kofman and A. D. Linde, Phys. Rev. D **59** (1999) 123523.
- [34] D. H. Lyth, Phys. Rev. Lett. **78** (1997) 1861.
- [35] In the SM one has $r = 192/(4\tilde{N} + 3)^2$, which amounts to $r = 0.007$ for $\tilde{N} = 40$.