

# CRITICAL AND SUBCRITICAL FRACTIONAL PROBLEMS WITH VANISHING POTENTIALS

JOÃO MARCOS DO Ó, OLÍMPIO H. MIYAGAKI, AND MARCO SQUASSINA

ABSTRACT. We investigate a class of nonlinear nonautonomous scalar field equations with fractional diffusion, critical power nonlinearity and a subcritical term. The involved potentials are allowed for vanishing behavior at infinity. The problem is set on the whole space and compactness issues have to be tackled.

## 1. INTRODUCTION AND MAIN RESULTS

We consider existence of solutions for the following class of equations

$$(1.1) \quad (-\Delta)^{\frac{s}{2}}u + V(x)u = K(x)f(u) + \lambda|u|^{2_s^*-2}u \quad \text{in } \mathbb{R}^N.$$

Here  $\lambda \geq 0$  is a parameter,  $s \in (0, 2)$ ,  $2_s^* = 2N/(N - s)$ ,  $N > s$ ,  $(-\Delta)^{\frac{s}{2}}$  is fractional laplacian  $V, K$  are positive functions and  $f$  is a continuous function with quasiscritical growth. Recently, a great attention has been focused on the study nonlinear problems involving fractional laplacian, in view of real-world applications. For instance, this type of operators arise in thin obstacle problems, optimization, finance, phase transitions, stratified materials, anomalous diffusion, crystal dislocation, soft thin films, semipermeable membranes, flame propagation, conservation laws, ultra-relativistic limits of quantum mechanics, quasi-geostrophic flows, multiple scattering, minimal surfaces, materials science and water waves, see [22]. The fractional laplacian  $(-\Delta)^{\frac{s}{2}}$  with  $s \in (0, 2)$  of a function  $\phi : \mathbb{R}^N \rightarrow \mathbb{R}$  is defined by

$$\mathcal{F}((-\Delta)^{\frac{s}{2}}\phi)(\xi) = |\xi|^s \mathcal{F}(\phi)(\xi), \quad \text{for } s \in (0, 2),$$

where  $\mathcal{F}$  is the Fourier transform. We are going explore problem (1.1) with *zero mass* potential, that is when  $V(x) \rightarrow 0$ , as  $|x| \rightarrow \infty$ . This class was studied by several researchers in the local case  $s = 2$ , e.g. in [1, 2, 7, 8, 10, 11, 24, 27] and reference therein, where the main feature is to impose restrictions on  $V, K$  to get some compact embedding into a weighted  $L^p$  space. Recently Alves and Souto, in [3], in addition to improving all the former restrictions on the potentials, handled subcritical nonlinearities  $f$  which do not satisfy the so-called Ambrosetti-Rabinowitz condition, namely,

$$(AR) \quad \text{there exists } \vartheta \in (2, 2_s^*) \text{ with } 0 < \vartheta F(s) \leq s f(s) \text{ for all } s > 0, \quad F(s) = \int_0^s f(t) dt.$$

Conditions weaker than (AR) were used, first time, in [20, 25, 27, 28, 34]. In all the above cited papers, the nonlinearity  $f$  had subcritical growth, that is, in addition to  $\lambda = 0$ , the growth of  $f$  is comparable with  $s^p$  with  $p < (N + 2)/(N - 2)$ , for  $N \geq 3$ . In the case  $s \in (0, 2)$ , nonlocal case, we say that  $f$  has a subcritical growth, if the growth of  $f$  in  $s$  is comparable with  $s^p$  for  $p < (N + s)/(N - s)$ , with  $N > s$ . In this situation, we would like to mention two works, one by Chang and Wang [19], where the authors recovered the Berestycki and Lions [11] results by improving Strauss compactness result [35], and a paper by Secchi [30] where the existence of ground state solutions is established. Motivated

---

2000 *Mathematics Subject Classification.* 35P15, 35P30, 35R11.

*Key words and phrases.* Fractional laplacian, nonautonomous equations, vanishing potentials, variational methods.

Research supported in part by INCTmat/MCT/Brazil, J.M. do Ó was partially supported by CNPq, M. Squassina was supported by MIUR project Variational and Topological Methods in the Study of Nonlinear Phenomena, O.H. Miyagaki was partially supported by CNPq/Brazil and CAPES/Brazil (Proc 2531/14-3).

The paper was completed while the second author was visiting the Department of Mathematics of Rutgers University, whose hospitality he gratefully acknowledges. He would also like to express his gratitude to Prof. Haim Brezis.

by the papers above, we are going to study the nonlocal case, with nonlinearities involving a critical growth and a subcritical perturbation  $f$ . Elliptic problems with critical growth, after the pioneering works by Brezis and Nirenberg [14] have had many progresses in several directions. We would like to mention [6, 39] and the references therein, in local case. For nonlocal case, in bounded domain, we cite [9, 15, 23, 26, 31, 38] and references therein, while in whole space was studied recently in [33] for non vanishing potential. Recently, Caffarelli and Silvestre [16] developed a local interpretation of the fractional Laplacian given in  $\mathbb{R}^N$  by considering a Neumann type operator in the extended domain  $\mathbb{R}_+^{N+1}$  defined by  $\{(x, t) \in \mathbb{R}^{N+1} : t > 0\}$ . A similar extension, for nonlocal problems on bounded domain with the zero Dirichlet boundary condition, was established, for instance, by Cabré and Tan in [15], Tan [37], Capella, Dàvila, Dupaigne and Sire [17], Brändle, Colorado, de Pablo and Sánchez [13]. It is worth noticing that, in a bounded domain, the Fourier definition of the fractional laplacian and its local Caffarelli-Silvestre interpretation do not agree, see the discussion developed [32] for more details. For  $u \in H^{s/2}(\mathbb{R}^N)$ , the solution  $w \in X^s(\mathbb{R}_+^{N+1})$  of

$$(1.2) \quad \begin{cases} -\operatorname{div}(y^{1-s}\nabla w) = 0 & \text{in } \mathbb{R}_+^{N+1} \\ w = u & \text{on } \mathbb{R}^N \times \{0\} \end{cases}$$

is called  $s$ -harmonic extension  $w = E_s(u)$  of  $u$  and it is proved in [16] (see also [13]) that

$$\lim_{y \rightarrow 0^+} y^{1-s} \frac{\partial w}{\partial y}(x, y) = -\frac{1}{k_s} (-\Delta)^{\frac{s}{2}} u(x),$$

where

$$k_s = 2^{1-s} \Gamma(1 - \frac{s}{2}) / \Gamma(\frac{s}{2}).$$

Here the spaces  $X^s(\mathbb{R}_+^{N+1})$  and  $H^{s/2}(\mathbb{R}^N)$  are defined as the completion of  $C_0^\infty(\overline{\mathbb{R}_+^{N+1}})$  and  $C_0^\infty(\mathbb{R}^N)$ , under the norms (which actually do coincide, see [13, Lemma A.2])

$$\begin{aligned} \|w\|_{X^s} &:= \left( \int_{\mathbb{R}_+^{N+1}} k_s y^{1-s} |\nabla w|^2 dx dy \right)^{1/2}, \\ \|u\|_{H^{\frac{s}{2}}} &:= \left( \int_{\mathbb{R}^N} |2\pi\xi|^s |\mathbb{F}(u(\xi))|^2 d\xi \right)^{1/2} = \left( \int_{\mathbb{R}^N} |(-\Delta)^{\frac{s}{2}} u|^2 dx \right)^{1/2}. \end{aligned}$$

Our problem (1.1) will be studied in the half-space, namely,

$$(1.3) \quad \begin{cases} -\operatorname{div}(y^{1-s}\nabla w) = 0 & \text{in } \mathbb{R}_+^{N+1} \\ -k_s \frac{\partial w}{\partial \nu} = -V(x)u + K(x)f(u) + \lambda|u|^{2_s^*-2}u & \text{on } \mathbb{R}^N \times \{0\}, \end{cases}$$

where

$$\frac{\partial w}{\partial \nu} = \lim_{y \rightarrow 0^+} y^{1-s} \frac{\partial w}{\partial y}(x, y).$$

We are looking for a positive solution in the Hilbert space  $E$  defined by

$$E = \left\{ w \in X^s(\mathbb{R}_+^{N+1}) : \int_{\mathbb{R}^N} V(x)w(x, 0)^2 dx < \infty \right\}$$

endowed with norm

$$\|w\| := \left( \int_{\mathbb{R}_+^{N+1}} k_s y^{1-s} |\nabla w|^2 dx dy + \int_{\mathbb{R}^N} V(x)w(x, 0)^2 dx \right)^{1/2}.$$

Consider the Euler-Lagrange functional associated to (1.3) given by

$$(1.4) \quad J_\lambda(w) := \frac{1}{2} \|w\|^2 - \int_{\mathbb{R}^N} K(x)F(w(x, 0)) dx - \frac{\lambda}{2_s^*} \int_{\mathbb{R}^N} w^+(x, 0)^{2_s^*} dx$$

which is  $C^1$  with Gâteaux derivative

$$(1.5) \quad J'_\lambda(w)v = \int_{\mathbb{R}_+^{N+1}} k_s y^{1-s} \nabla w \cdot \nabla v dx dy + \int_{\mathbb{R}^N} V(x)w(x,0)v(x,0)dx \\ - \int_{\mathbb{R}^N} K(x)f(w(x,0))v(x,0)dx - \lambda \int_{\mathbb{R}^N} w^+(x,0)^{2_s^*-1}v(x,0)dx, \quad \text{for all } w, v \in E.$$

We now formulate assumptions for  $V, K, f$  in problem (1.1).

• ASSUMPTIONS ON  $V$  AND  $K$ .

(I) (sign of  $V$  and  $K$ ):  $V, K$  are continuous,  $V, K > 0$  on  $\mathbb{R}^N$  and  $K \in L^\infty(\mathbb{R}^N)$ ;

(II) (decay of  $K$ ): If  $\{A_n\}$  is a sequence of Borel sets of  $\mathbb{R}^N$  with  $|A_n| \leq R$  for some  $R > 0$ ,

$$(1.6) \quad \lim_{r \rightarrow \infty} \int_{A_n \cap B_r^c(0)} K(x)dx = 0, \quad \text{uniformly with respect to } n \in \mathbb{N};$$

(III) (interrelation between  $V$  and  $K$ ): either

$$(1.7) \quad K/V \in L^\infty(\mathbb{R}^N)$$

or there exists  $p \in (2, 2_s^*)$  such that

$$(1.8) \quad \lim_{|x| \rightarrow \infty} \frac{K(x)}{V(x)^\gamma} = 0, \quad \gamma = \frac{ps - N(p-2)}{2s} \in (0, 1).$$

• ASSUMPTIONS ON  $f$ .

(f1) (behaviour at zero):  $f : \mathbb{R} \rightarrow \mathbb{R}^+$  is continuous with  $f = 0$  on  $\mathbb{R}^-$ . If (1.7) holds, then

$$\limsup_{s \rightarrow 0^+} \frac{f(s)}{s} = 0.$$

If condition (1.8) holds, we assume

$$\limsup_{s \rightarrow 0^+} \frac{f(s)}{s^{p-1}} < +\infty.$$

(f2) (quasi-critical growth): If (1.7) holds, then

$$\limsup_{s \rightarrow +\infty} \frac{f(s)}{s^{2_s^*-1}} = 0.$$

If condition (1.8) holds, we assume

$$\limsup_{s \rightarrow +\infty} \frac{F(s)}{s^p} < +\infty.$$

(f3) (super-quadraticity):  $\frac{f(s)}{s}$  is non-decreasing in  $\mathbb{R}^+$ , and

$$\limsup_{s \rightarrow +\infty} \frac{F(s)}{s^2} = +\infty.$$

(f3)' (super-quadraticity):  $\frac{f(s)}{s}$  is non-decreasing in  $\mathbb{R}^+$  and there exist  $C_0 > 0$  and  $q \in (2, 2_s^*)$  with

$$F(s) \geq C_0 s^q, \quad \text{for all } s \in \mathbb{R}^+$$

The following are the main results of the paper.

**Theorem 1.1.** Assume (I)-(III), (f1)-(f3) and  $\lambda = 0$ . Then (1.1) admits a positive solution  $u \in E$ .

**Theorem 1.2.** Assume (I)-(III), (f1)-(f2)-(f3)', (AR),  $\lambda = 1$  and that one of the following hold

- (1)  $N > 2s$ ,
- (2)  $N = 2s$ ,
- (3)  $s < N < 2s$  and  $q > \frac{N}{N-s}$ ,
- (4)  $s < N < 2s$  and  $q < \frac{N}{N-s}$ , with  $C_0$  large enough.

Then (1.1) admits a positive solution  $u \in E$ .

Throughout the paper, unless explicitly stated, the symbol  $C$  will always denote a generic positive constant, which may vary from line to line.

## 2. PRELIMINARY RESULTS

Consider the weighted Banach space:

$$L_K^p = \left\{ u : \mathbb{R}^N \rightarrow \mathbb{R} \text{ measurable and } \int_{\mathbb{R}^N} K(x)|u|^p dx < \infty \right\}, \quad \|\cdot\|_{L_K^p} = \left( \int_{\mathbb{R}^N} K(x)|u|^p dx \right)^{1/p}.$$

The first result, on compact injections for  $E$ , follows by adapting the arguments in [3].

**Proposition 2.1** (Compactness). *The following facts hold:*

- (1)  $E$  is compactly embedded into  $L_K^q$  for all  $q \in (2, 2_s^*)$ , provided that (1.7) holds;
- (2)  $E$  is compactly embedded into  $L_K^p$  provided that (1.8) holds;
- (3) If  $w_n \rightharpoonup w$  in  $E$ , then up to a subsequence,

$$\lim_n \int_{\mathbb{R}^N} K(x)F(w_n(x, 0))dx = \int_{\mathbb{R}^N} K(x)F(w(x, 0))dx;$$

- (4) If  $w_n \rightharpoonup w$  in  $E$ , then up to a subsequence,

$$\lim_n \int_{\mathbb{R}^N} K(x)w_n(x, 0)f(w_n(x, 0))dx = \int_{\mathbb{R}^N} K(x)w(x, 0)f(w(x, 0))dx;$$

- (5) If  $w_n \rightharpoonup w$  in  $E$ , then, up to a subsequence, for any  $v \in E$ ,

$$\lim_n \int_{\mathbb{R}^N} w_n^+(x, 0)^{2_s^*-1}v(x, 0)dx = \int_{\mathbb{R}^N} w^+(x, 0)^{2_s^*-1}v(x, 0)dx.$$

- (6) If  $w_n \rightharpoonup w$  in  $E$ , then up to a subsequence, for any  $v \in E$ ,

$$\lim_n \int_{\mathbb{R}^N} K(x)f(w_n(x, 0))v(x, 0)dx = \int_{\mathbb{R}^N} K(x)f(w(x, 0))v(x, 0)dx.$$

*Proof.* Assume that condition (1.7) holds, let  $q \in (2, 2_s^*)$  and let us prove assertion (1). Let  $\varepsilon > 0$ . Then, there exist  $0 < s_0(\varepsilon) < s_1(\varepsilon)$ , a positive constant  $C(\varepsilon)$  and  $C_0$  depending only on  $V$  and  $K$ , such that

$$(2.1) \quad K(x)|s|^q \leq \varepsilon C_0(V(x)|s|^2 + |s|^{2_s^*}) + C(\varepsilon)K(x)\chi_{[s_0(\varepsilon), s_1(\varepsilon)]}(|s|)|s|^{2_s^*}, \quad \text{for all } s \in \mathbb{R}.$$

Therefore we obtain, for every  $w \in E$  and  $r > 0$ ,

$$(2.2) \quad \int_{B_r^{+c}(0) \cap \{y=0\}} K(x)|w(x, 0)|^q dx \leq \varepsilon Q(w) + C(\varepsilon)s_1(\varepsilon)^{2_s^*} \int_{A_\varepsilon \cap (B_r^{+c}(0) \cap \{y=0\})} K(x)dx,$$

where we have set

$$(2.3) \quad Q(w) := C_0 \int_{\mathbb{R}^N} (V(x)|w(x, 0)|^2 + |w(x, 0)|^{2_s^*})dx, \quad A_\varepsilon := \{x \in \mathbb{R}^N : s_0(\varepsilon) \leq |w(x, 0)| \leq s_1(\varepsilon)\}.$$

If  $(w_n) \subset E$  is such that  $w_n \rightharpoonup w$  weakly in  $E$  for some  $w \in E$ , there exists  $M > 0$  with

$$(2.4) \quad \begin{aligned} \int_{\mathbb{R}_+^{N+1}} k_s |\nabla w_n|^2 dx dy + \int_{\mathbb{R}^N} V(x)|w_n(x, 0)|^2 dx &\leq M, \quad \text{for all } n \in \mathbb{N}, \\ \int_{\mathbb{R}^N} |w_n(x, 0)|^{2_s^*} dx &\leq M, \quad \text{for all } n \in \mathbb{N}, \end{aligned}$$

so that  $Q(w_n)$  is bounded in  $\mathbb{R}$ . On the other hand, if  $A_\varepsilon^n = \{s_0(\varepsilon) \leq |w_n(x, 0)| \leq s_1(\varepsilon)\}$ , we get

$$s_0(\varepsilon)^{2_s^*} |A_\varepsilon^n| \leq \int_{A_\varepsilon^n} |w_n(x, 0)|^{2_s^*} dx \leq \int_{\mathbb{R}^N} |w_n(x, 0)|^{2_s^*} dx \leq M, \quad \text{for all } n \in \mathbb{N}.$$

which implies that  $\sup_{n \in \mathbb{N}} |A_\varepsilon^n| < +\infty$ . Then, in light of (1.6), there exists  $r(\varepsilon) > 0$  such that

$$(2.5) \quad \int_{A_\varepsilon^n \cap (B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\})} K(x) dx < \frac{\varepsilon}{C(\varepsilon)s_1(\varepsilon)^{2_s^*}}, \quad \text{for all } n \in \mathbb{N}.$$

Whence, invoking (2.2), we get

$$(2.6) \quad \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x) |w_n(x, 0)|^q dx \leq (2C_0M + 1)\varepsilon.$$

By the fractional compact embedding [13], we have

$$(2.7) \quad \lim_{n \rightarrow \infty} \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x) |w_n(x, 0)|^q dx = \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x) |w(x, 0)|^q dx.$$

Combining (2.6)-(2.7), yields

$$\lim_n \int_{\mathbb{R}^N \cap \{y=0\}} K(x) |w_n(x, 0)|^q dx = \int_{\mathbb{R}^N \cap \{y=0\}} K(x) |w(x, 0)|^q dx,$$

which concludes the proof of (1).

Assume now that condition (1.8) holds and let us prove assertion (2). By a direct calculation, for any  $x \in \mathbb{R}^N$  and  $s \geq 0$ , if  $\gamma \in (0, 1)$  is the constant introduced in (1.8), we get

$$V(x)s^{2-p} + s^{2_s^*-p} \geq \omega(p, s)V(x)^\gamma, \quad \omega(p, s) = \left(\frac{2_s^* - 2}{2_s^* - p}\right) \left(\frac{p-2}{2_s^* - p}\right)^{\frac{2-p}{2_s^*-2}}.$$

Let  $\varepsilon > 0$ . Combining this inequality with (1.8), there exists  $r(\varepsilon) > 0$  such that

$$(2.8) \quad K(x)|s|^p \leq \varepsilon(V(x)|s|^2 + |s|^{2_s^*}), \quad \text{for all } s \in \mathbb{R} \text{ and } |x| \geq r(\varepsilon).$$

Then, for all  $w \in E$ , we conclude

$$\int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x) |w(x, 0)|^p dx \leq \varepsilon \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} (V(x)|w(x, 0)|^2 + |w(x, 0)|^{2_s^*}) dx.$$

If  $(w_n) \subset E$  and  $w_n \rightharpoonup w$  weakly in  $E$ , there exists  $M > 0$  such that (2.4) hold. Whence, for a suitable radius  $r(\varepsilon) > 0$  there holds

$$(2.9) \quad \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x) |w_n(x, 0)|^p dx \leq 2\varepsilon M, \quad \text{for all } n \in \mathbb{N}.$$

Since  $p \in (2, 2_s^*)$ , by the fractional compact embedding we have

$$(2.10) \quad \lim_n \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x) |w_n(x, 0)|^p dx = \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x) |w(x, 0)|^p dx.$$

Combining (2.9) and (2.10) we get

$$\lim_n \int_{\mathbb{R}^N \cap \{y=0\}} K(x) |w_n(x, 0)|^p dx = \int_{\mathbb{R}^N \cap \{y=0\}} K(x) |w(x, 0)|^p dx,$$

which concludes the proof of assertion (2).

Let us now turn to the proof of (3) and (4) under assumption (1.7). From  $(f_1)$ - $(f_3)$ , fixed  $q \in (2, 2_s^*)$  and given  $\varepsilon > 0$ , there exist  $0 < s_0(\varepsilon) < s_1(\varepsilon)$ ,  $C(\varepsilon) > 0$  and  $C_0$  depending only upon  $V$  and  $K$ , with

$$(2.11) \quad |K(x)F(s)| \leq \varepsilon C_0(V(x)|s|^2 + |s|^{2_s^*}) + C(\varepsilon)K(x)\chi_{[s_0(\varepsilon), s_1(\varepsilon)]}(|s|)|s|^q, \quad \text{for all } s \in \mathbb{R},$$

$$(2.12) \quad |K(x)f(s)s| \leq \varepsilon C_0(V(x)|s|^2 + |s|^{2_s^*}) + C(\varepsilon)K(x)\chi_{[s_0(\varepsilon), s_1(\varepsilon)]}(|s|)|s|^q, \quad \text{for all } s \in \mathbb{R}.$$

Notice that, by (1.6), arguing as for the proof of (1), there exists  $r(\varepsilon) > 0$  such that

$$(2.13) \quad \int_{A_\varepsilon^n \cap B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x) dx \leq \frac{\varepsilon}{C(\varepsilon)s_1(\varepsilon)^q}, \quad \text{for all } n \in \mathbb{N}.$$

Let  $\{w_n\} \in E$  be bounded. Combining the above inequality with (2.4) and (2.11)-(2.12), we have

$$(2.14) \quad \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x)F(w_n(x,0))dx \leq (2C_0M + 1)\varepsilon, \quad \text{for all } n \in \mathbb{N},$$

$$(2.15) \quad \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x)f(w_n(x,0))w_n(x,0)dx \leq (2C_0M + 1)\varepsilon, \quad \text{for all } n \in \mathbb{N}.$$

Since  $(w_n(x,0))$  is bounded in  $L^{2^*_s}(\mathbb{R}^N)$ , by Strauss lemma [11, Theorem A.I p.338], we infer

$$\begin{aligned} \lim_n \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x)F(w_n(x,0))dx &= \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x)F(w(x,0))dx, \\ \lim_n \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x)f(w_n(x,0))w_n(x,0)dx &= \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x)f(w(x,0))w(x,0)dx. \end{aligned}$$

Combining these limits with (2.14)-(2.15) we conclude the proof.

Let us now turn to the proof of (3) and (4) under assumption (1.8). Let  $\varepsilon > 0$ . We learned that there exists  $r(\varepsilon) > 0$  such that (2.8) holds, yielding

$$\begin{aligned} K(x)|F(s)| &\leq \varepsilon(V(x)|F(s)||s|^{2-p} + |F(s)||s|^{2^*_s-p}), \quad \text{for all } s \in \mathbb{R} \text{ and } |x| \geq r(\varepsilon), \\ K(x)f(s)s &\leq \varepsilon(V(x)f(s)s|s|^{2-p} + f(s)s|s|^{2^*_s-p}), \quad \text{for all } s \in \mathbb{R}^+ \text{ and } |x| \geq r(\varepsilon). \end{aligned}$$

From (f<sub>1</sub>)-(f<sub>2</sub>), there exist  $0 < s_0(\varepsilon) < s_1(\varepsilon)$  satisfying

$$\begin{aligned} K(x)|F(s)| &\leq \varepsilon(V(x)|s|^2 + |s|^{2^*_s}), \quad \text{for all } s \in I_\varepsilon \text{ and } |x| \geq r(\varepsilon), \\ K(x)f(s)s &\leq \varepsilon(V(x)|s|^2 + |s|^{2^*_s}), \quad \text{for all } s \in I_\varepsilon \cap \mathbb{R}^+ \text{ and } |x| \geq r(\varepsilon), \end{aligned}$$

where  $I_\varepsilon = \{|s| < s_0(\varepsilon) \text{ or } |s| > s_1(\varepsilon)\}$ . Then, we have

$$(2.16) \quad \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x)F(w_n(x,0))dx \leq \varepsilon Q(w_n) + C(\varepsilon) \int_{A_\varepsilon^n \cap (B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\})} K(x)dx,$$

$$(2.17) \quad \int_{B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\}} K(x)f(w_n(x,0))w_n(x,0)dx \leq \varepsilon Q(w_n) + C(\varepsilon) \int_{A_\varepsilon^n \cap (B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\})} K(x)dx,$$

where

$$C(\varepsilon) = \max \left\{ \max_{[s_0(\varepsilon), s_1(\varepsilon)]} |F(s)|, \max_{[s_0(\varepsilon), s_1(\varepsilon)]} |f(s)s| \right\}.$$

Arguing as for the proof of (1), we have

$$\begin{aligned} \left| \int_{(B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\})} K(x)F(w_n(x,0))dx \right| &\leq (2M + 1)\varepsilon, \quad \text{for all } n \in \mathbb{N}, \\ \left| \int_{(B_{r(\varepsilon)}^{+c}(0) \cap \{y=0\})} K(x)f(w_n(x,0))w_n(x,0)dx \right| &\leq (2M + 1)\varepsilon, \quad \text{for all } n \in \mathbb{N}. \end{aligned}$$

Invoking again Strauss lemma, by the above inequalities, conclusions (3) and (4) follows. To prove (5), it is enough to observe that  $w_n^+(x,0)^{2^*_s-1} \rightharpoonup w^+(x,0)^{2^*_s-1}$  weakly in  $(L^{2^*_s})'$ . Finally, let us prove (6). If (1.7) holds, then the sequence  $(\sqrt{K(x)}f(w_n(x,0))\chi_{\{|w_n(x,0)| \leq 1\}})$  is bounded in  $L^2(\mathbb{R}^N)$  being

$$|\sqrt{K(x)}f(w_n(x,0))\chi_{\{|w_n(x,0)| \leq 1\}}|^2 \leq CV(x)|w_n(x,0)|^2.$$

This, by pointwise convergence, yields for every  $\varphi \in L^2(\mathbb{R}^N)$

$$\lim_k \int_{\mathbb{R}^N} \sqrt{K(x)}f(w_n(x,0))\chi_{\{|w_n(x,0)| \leq 1\}}\varphi(x)dx = \int_{\mathbb{R}^N} \sqrt{K(x)}f(w(x,0))\chi_{\{|w(x,0)| \leq 1\}}\varphi(x)dx.$$

Given  $v \in E$ , since  $\sqrt{K(x)} \leq C\sqrt{V(x)}$ , it follows  $\sqrt{K(x)}v(x,0) \in L^2(\mathbb{R}^N)$ , yielding

$$(2.18) \quad \lim_k \int_{\mathbb{R}^N} K(x)f(w_n(x,0))\chi_{\{|w_n(x,0)| \leq 1\}}v(x,0)dx = \int_{\mathbb{R}^N} K(x)f(w(x,0))\chi_{\{|w(x,0)| \leq 1\}}v(x,0)dx.$$

In a similar fashion, the sequence  $(K(x)f(w_n(x,0))\chi_{\{|w_n(x,0)|\geq 1\}})$  is bounded in  $L^{\frac{2^*}{2^*-1}}(\mathbb{R}^N)$  being

$$|K(x)f(w_n(x,0))\chi_{\{|w_n(x,0)|\geq 1\}}|^{\frac{2^*}{2^*-1}} \leq |w_n(x,0)|^{2^*}.$$

This, by pointwise convergence, and since  $v \in E$ , yields

$$(2.19) \quad \lim_k \int_{\mathbb{R}^N} K(x)f(w_n(x,0))\chi_{\{|w_n(x,0)|\geq 1\}}v(x,0)dx = \int_{\mathbb{R}^N} K(x)f(w(x,0))\chi_{\{|w(x,0)|\geq 1\}}v(x,0)dx.$$

Combining (2.18)-(2.19) yields the assertion. In a similar fashion one can treat the case when (1.8) holds since, by means of (2),  $K^{1/p}v(x,0) \in L^p(\mathbb{R}^N)$  for all  $v \in E$  and, up to a subsequence,

$$|K(x)^{\frac{p-1}{p}}f(w_n(x,0))\chi_{\{|w_n(x,0)|\leq 1\}}|^{p'} \leq K(x)|w_n(x,0)|^p \leq z(x) \in L^1(\mathbb{R}^N).$$

This concludes the proof.  $\square$

From  $(f_1)$ - $(f_2)$  one can prove that  $J_\lambda$  satisfies the Mountain-Pass geometry.

**Lemma 2.2** (Geometry). *The functional  $J_\lambda$  satisfies*

- (1) *There exists  $\beta, \rho > 0$  such that  $J_\lambda(u) \geq \beta$  if  $\|u\| = \rho$ ;*
- (2) *There exists  $e \in E \setminus \{0\}$  with  $\|e\| > \rho$  such that  $J_\lambda(e) \leq 0$ ;*

*Proof.* (2) is obvious. Concerning (1), observe that in light of condition (1.8) on  $V$  and  $K$ , the space  $E$  is continuously embedded into  $L_K^p(\mathbb{R})$  where  $p \in (2, 2^*)$  is the precisely the value which appears in condition (1.8). This can be readily obtained by arguing as in the proof of [12, part (i) of Theorem 4] (see formula (8) therein obtained by Hölder inequality) and by using the fractional Sobolev inequality. This is possible since in any of the two assumptions between  $V$  and  $K$ , we have that

$$\frac{K}{V^\gamma} \in L^\infty(\mathbb{R}), \quad \gamma = \frac{ps - N(p-2)}{2s}.$$

This is the fractional counterpart of the assumption on  $\mathcal{W}$  in [12]. Once this embedding is available, recall that we can write the inequality, for  $\varepsilon_0$  to be fixed small

$$K(x)F(s) \leq \varepsilon_0 V(x)s^2 + Cs^{2^*} + CK(x)s^p, \quad x \in \mathbb{R}, s \in \mathbb{R}^+,$$

and the Mountain-Pass geometry can be proved.  $\square$

Therefore, there exists a sequence  $\{w_n\} \subset E$ , so called *Cerami sequence* [18], such that

$$(2.20) \quad J_\lambda(w_n) \rightarrow c, \quad (1 + \|w_n\|)\|J'_\lambda(w_n)\| \rightarrow 0,$$

where  $c$  is given by

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J_\lambda(\gamma(t)),$$

with

$$\Gamma = \{\gamma \in C([0,1], E) : \gamma(0) = 0 \text{ and } J_\lambda(\gamma(1)) \leq 0\}.$$

Next we turn to the boundedness of  $(w_n)$  in  $E$ .

**Lemma 2.3** (Boundedness). *Let  $\lambda \in \{0, 1\}$ . Then the Cerami sequence  $(w_n) \subset E$  is bounded.*

*Proof.* First of all, we observe that  $w_n^- \in E$  and, by the definition of  $J_\lambda$ ,

$$\begin{aligned} J'_\lambda(w_n)(-w_n^-) &= - \int_{\mathbb{R}_+^{N+1}} k_s y^{1-s} \nabla w_n \cdot \nabla w_n^- dx dy - \int_{\mathbb{R}^N} V(x)w_n(x,0)w_n^-(x,0) \\ &= \int_{\mathbb{R}_+^{N+1}} k_s y^{1-s} |\nabla w_n^-|^2 dx dy + \int_{\mathbb{R}^N} V(x)w_n^-(x,0)^2 = \|w_n^-\|^2. \end{aligned}$$

Since  $(1 + \|w_n\|)J'_\lambda(w_n)(-w_n^-) = o_n(1)$  as  $n \rightarrow \infty$ , it follows that  $J'_\lambda(w_n)(-w_n^-) = o_n(1)$  as  $n \rightarrow \infty$ , which in turn implies that  $\|w_n^-\| = o_n(1)$ , as  $n \rightarrow \infty$ .

**Case  $\lambda = 0$ .** Denote  $J_0 = J$ . Let  $t_n \in [0, 1]$  be such that

$$J(t_n w_n) = \max_{t \in [0, 1]} J(t w_n).$$

We *claim* that  $J(t_n w_n)$  is bounded from above. Without loss of generality, we may assume that  $t_n \in (0, 1)$  for all  $n$ . Then, we have  $J'(t_n w_n)(w_n) = 0$  and

$$(2.21) \quad \begin{aligned} 2J(t_n w_n) &= 2J(t_n w_n) - J'(t_n w_n)(t_n w_n) \\ &= \int_{\mathbb{R}^N} K(x) \mathcal{H}(t_n w_n(x, 0)) dx = \int_{\mathbb{R}^N} K(x) \mathcal{H}(t_n w_n^+(x, 0)) dx, \end{aligned}$$

where  $\mathcal{H}(s) = sf(s) - 2F(s)$  is nondecreasing and  $\mathcal{H} = 0$  on  $\mathbb{R}^-$ . Thus, since  $t_n \in (0, 1)$  and  $w_n^+ \geq 0$ , from formula (2.21) we obtain that

$$\begin{aligned} 2J(t_n w_n) &\leq \int_{\mathbb{R}^N} K(x) \mathcal{H}(w_n^+(x, 0)) dx = \int_{\mathbb{R}^N} K(x) \mathcal{H}(w_n(x, 0)) dx \\ &= 2J(w_n) - J'(w_n)(w_n) = 2J(w_n) + o_n(1), \end{aligned}$$

which proves the claim. Now, we prove that  $(w_n) \subset E$  is bounded. Assume by contradiction that, up to subsequence,  $\|w_n\| \rightarrow +\infty$  as  $n \rightarrow \infty$ . Set  $z_n := w_n / \|w_n\|$  and suppose that  $z_n \rightharpoonup z$ , as  $n \rightarrow \infty$ , in  $E$ . We now *claim* that  $z(x, 0) = 0$  almost everywhere in  $\mathbb{R}^N$ . In fact,

$$o_n(1) + \frac{1}{2} = \int_{\mathbb{R}^N} \frac{K(x)F(w_n(x, 0))}{\|w_n\|^2} dx = \int_{\mathbb{R}^N} \frac{K(x)F(w_n(x, 0))}{|w_n(x, 0)|^2} z_n^2(x, 0) dx.$$

By  $(f_3)$ , given  $\tau > 0$  there exists  $\xi_\tau > 0$  such that  $F(s) \geq \tau s^2$  for all  $|s| \geq \xi_\tau$ . Thus,

$$o_n(1) + \frac{1}{2} \geq \int_{\{|w_n(x, 0)| \geq \xi_\tau\}} \frac{K(x)F(w_n(x, 0))}{|w_n(x, 0)|^2} z_n^2(x, 0) dx \geq \tau \int_{\mathbb{R}^N} K(x) z_n^2(x, 0) \chi_{\{|z_n(x, 0)| \geq \frac{\xi_\tau}{\|w_n\|}\}} dx.$$

Thus, by Fatou lemma, since  $z_n^2(x, 0) \chi_{\{|z_n(x, 0)| \geq \frac{\xi_\tau}{\|w_n\|}\}} \rightarrow z(x, 0)$  a.e., for any  $\tau > 0$ , we conclude

$$\frac{1}{2} \geq \tau \int_{\mathbb{R}^N} K(x) z^2(x, 0) dx.$$

Since  $K > 0$ , it follows  $z(x, 0) = 0$ , by the arbitrariness of  $\tau > 0$  and the claim follows. Now, let  $B > 0$ . Of course  $B\|w_n\|^{-1} \in [0, 1]$  eventually for  $n \geq n_B$ , for some  $n_B \in \mathbb{N}$ . Thus,

$$J(t_n w_n) \geq J(Bz_n) = \frac{B^2}{2} - \int_{\mathbb{R}^N} K(x)F(Bz_n(x, 0)) dx,$$

since  $t_n$  is a maximum point. By Proposition 2.1, it follows

$$\int_{\mathbb{R}^N} K(x)F(Bz_n(x, 0)) dx \rightarrow \int_{\mathbb{R}^N} K(x)F(Bz(x, 0)) = 0,$$

and we have  $J(t_n w_n) + o_n(1) \geq B^2/2$ , which yields  $\sup\{J(t_n w_n) : n \in \mathbb{N}\} \geq B^2/2$ , a contradiction if

$$B = 2\sqrt{\sup\{J(t_n w_n) : n \in \mathbb{N}\}} \in (0, \infty).$$

This concludes the proof.

**Case  $\lambda = 1$ .** Denote  $J_\lambda = J$ . The boundedness of the  $\{w_n\}$  in  $E$  follows easily from (AR), since

$$o_n(1) + c \geq J(w_n) - \frac{1}{\vartheta} J'(w_n)(w_n) \geq \left(\frac{1}{2} - \frac{1}{\vartheta}\right) \|w_n\|^2.$$

This concludes the proof. □

The following Sobolev inequality can be found in [13],

$$(2.22) \quad \int_{\mathbb{R}_+^{N+1}} y^{1-s} |\nabla w|^2 dx dy \geq \mathbb{S}(s, N) \left( \int_{\mathbb{R}_+^N} |w(x, 0)|^{2^*} dx \right)^{\frac{2}{2^*}}, \quad \text{for all } w \in X^s(\mathbb{R}_+^{N+1}),$$



where

$$\mathbb{S}(s, N) = \frac{\Gamma(\frac{s}{2})\Gamma(\frac{1}{2}(N-s))(\Gamma(N))^{\frac{s}{N}}}{2\pi^{\frac{s}{N}}\Gamma(\frac{1}{2}(2-s))\Gamma(\frac{1}{2}(N+s))(\Gamma(\frac{1}{2}N))^{\frac{s}{N}}}.$$

This constant is achieved on the family of functions [13, 21, 31]  $w_\varepsilon = E_s(u_\varepsilon)$  (by [36] for  $s = 2$ ), where

$$u_\varepsilon(x) = \frac{\varepsilon^{\frac{N-s}{2}}}{(|x|^2 + \varepsilon^2)^{\frac{N-s}{2}}}, \quad \varepsilon > 0.$$

Furthermore, take  $\phi(x, y) = \phi_0(|(x, y)|)$ , where  $\phi_0 \in C^\infty(0, \infty)$  is a non increasing cut-off such that

$$\phi_0(s) = 1 \quad \text{if } s \in [0, 1/2], \quad \phi_0(s) = 0, \quad \text{if } s \geq 1.$$

Let  $\phi w_\varepsilon$  which belongs to  $X^s(\mathbb{R}_+^{N+1})$ . By [9, Lemma 3.8] (which is formulated on a bounded domain  $\Omega$ , but which holds with the very same proof when taking  $\Omega = \mathbb{R}^N$ ), we have

**Lemma 2.4** (concentration). *The family  $\{\phi w_\varepsilon\}$ , and its trace on  $\{y = 0\}$ , namely,  $\phi u_\varepsilon$ , satisfy*

$$(2.23) \quad \|\phi w_\varepsilon\|_{X^s}^2 = \|w_\varepsilon\|_{X^s}^2 + \mathcal{O}(\varepsilon^{N-s}),$$

$$(2.24) \quad \|\phi u_\varepsilon\|_{L^2}^2 = \begin{cases} \mathcal{O}(\varepsilon^s) & \text{if } N > 2s, \\ \mathcal{O}(\varepsilon^s \log(1/\varepsilon)) & \text{if } N = 2s, \\ \mathcal{O}(\varepsilon^{N-s}) & \text{if } N < 2s, \end{cases}$$

for  $\varepsilon > 0$  small enough. Define  $\eta_\varepsilon = \phi w_\varepsilon / \|\phi u_\varepsilon\|_{L^2}$ , then

$$(2.25) \quad \|\eta_\varepsilon\|_{X^s}^2 = k_s \mathbb{S}(s, N) + \mathcal{O}(\varepsilon^{N-s}),$$

$$(2.26) \quad \|\eta_\varepsilon(x, 0)\|_{L^2}^2 = \begin{cases} \mathcal{O}(\varepsilon^s) & \text{if } N > 2s, \\ \mathcal{O}(\varepsilon^s \log(1/\varepsilon)) & \text{if } N = 2s, \\ \mathcal{O}(\varepsilon^{N-s}) & \text{if } N < 2s. \end{cases}$$

and

$$(2.27) \quad \|\eta_\varepsilon(x, 0)\|_{L^q}^q = \begin{cases} \mathcal{O}(\varepsilon^{\frac{2N-(N-s)q}{2}}) & \text{if } q > \frac{N}{N-s} \text{ (or } N \geq 2s), \\ \mathcal{O}(\varepsilon^{\frac{(N-s)q}{2}}) & \text{if } q < \frac{N}{N-s}. \end{cases}$$

Here with the notation  $a_\varepsilon = \mathcal{O}(b_\varepsilon)$  we mean that  $a_\varepsilon/b_\varepsilon$  is uniformly bounded with respect to  $\varepsilon$ .

**Remark 2.5.** We remark that, actually, except (2.23) and (2.25), the other estimates follow exactly as in local case (see [14]), because in these cases, we know the explicit expression for  $u_\varepsilon$ . While, for  $w_\varepsilon$ , except for  $s = 1$  (see [38]) and  $s = 2$  (local case), the explicit expressions are not available. But, in [9], the authors were clever to overcome this difficulty, by exploring some properties of the Poisson kernel. The  $s$ -harmonic extension of the  $u_\varepsilon$  has the following explicit expression

$$w_\varepsilon(x, y) = P_y^s * u_\varepsilon(x) = C_{N,s} y^s \int_{\mathbb{R}^N} \frac{u_\varepsilon(\xi)}{(|x - \xi|^2 + y^2)^{\frac{N+s}{2}}} d\xi, \quad \text{for some } C_{N,s} > 0.$$

Noticing that as  $u_\varepsilon$  and  $P_y^s$  are self-similar functions, namely

$$u_\varepsilon(x) = \varepsilon^{\frac{s-N}{2}} u_1(x/\varepsilon), \quad P_y^s(x) = \frac{1}{y^N} P_1^s(x/|y|) = \frac{y^s}{(|x|^2 + y^2)^{\frac{N+s}{2}}},$$

then

$$w_\varepsilon(x, y) = \varepsilon^{\frac{s-N}{2}} w_1(x/\varepsilon, y/\varepsilon).$$

Exploiting this fact, they estimate as follows

$$\int_{\mathbb{R}_+^{N+1}} y^{1-s} w_\varepsilon \phi \nabla \phi \cdot \nabla w_\varepsilon dx dy \leq \mathcal{O}(\varepsilon^{N-s}), \quad \int_{\mathbb{R}_+^{N+1}} y^{1-s} |w_\varepsilon \nabla \phi|^2 dx dy \leq \mathcal{O}(\varepsilon^{N-s}).$$

Combining these inequalities, (2.23) holds. The inequality (2.25) comes as a consequence. Concerning (2.27), we justify it in the case  $q < N/(N-s)$ , the opposite case being similar. We have

$$\begin{aligned}
\|\phi u_\varepsilon\|_{L^q}^q &= \int_{\mathbb{R}^N} |\phi|^q |u_\varepsilon|^q dx \geq \int_{B(0, \frac{1}{2})} |u_\varepsilon|^q dx \\
&= \varepsilon^{\frac{(N-s)q}{2}} \int_{B(0, \frac{1}{2})} \frac{1}{(\varepsilon^2 + |x|^2)^{\frac{(N-s)q}{2}}} dx \\
&= \varepsilon^{\frac{(N-s)q}{2}} \int_{B(0, \frac{1}{2\varepsilon})} \frac{1}{\varepsilon^{(N-s)q} (1 + |y|^2)^{\frac{(N-s)q}{2}}} \varepsilon^N dy \\
&= \varepsilon^{N - \frac{(N-s)q}{2}} \int_{B(0, \frac{1}{2\varepsilon})} \frac{1}{(1 + |y|^2)^{\frac{(N-s)q}{2}}} dy \\
&= \varepsilon^{N - \frac{(N-s)q}{2}} C \int_0^{1/(2\varepsilon)} \frac{1}{(1 + \varrho^2)^{\frac{(N-s)q}{2}}} \varrho^{N-1} d\varrho \\
&= \varepsilon^{N - \frac{(N-s)q}{2}} C \left( \int_0^1 \frac{1}{(1 + \varrho^2)^{\frac{(N-s)q}{2}}} \varrho^{N-1} d\varrho + \int_1^{1/(2\varepsilon)} \frac{1}{(1 + \varrho^2)^{\frac{(N-s)q}{2}}} \varrho^{N-1} d\varrho \right) \\
&\geq \varepsilon^{N - \frac{(N-s)q}{2}} \left( C + C \int_1^{1/(2\varepsilon)} \frac{1}{\varrho^{(N-s)q - N + 1}} d\varrho \right) \\
&\geq \varepsilon^{N - \frac{(N-s)q}{2}} \left( C + C\varepsilon^{(N-s)q - N} \right) \geq C\varepsilon^{\frac{(N-s)q}{2}},
\end{aligned}$$

for  $\varepsilon > 0$  small enough. Analogously, we get

$$\begin{aligned}
\|\phi u_\varepsilon\|_{L^q}^q &= \int_{\mathbb{R}^N} |\phi|^q |u_\varepsilon|^q dx \leq \int_{B(0, 1)} |u_\varepsilon|^q dx \\
&= \varepsilon^{\frac{(N-s)q}{2}} \int_{B(0, 1)} \frac{1}{(\varepsilon^2 + |x|^2)^{\frac{(N-s)q}{2}}} dx \\
&= \varepsilon^{\frac{(N-s)q}{2}} \int_{B(0, \frac{1}{\varepsilon})} \frac{1}{\varepsilon^{(N-s)q} (1 + |y|^2)^{\frac{(N-s)q}{2}}} \varepsilon^N dy \\
&= \varepsilon^{N - \frac{(N-s)q}{2}} \int_{B(0, \frac{1}{\varepsilon})} \frac{1}{(1 + |y|^2)^{\frac{(N-s)q}{2}}} dy \\
&= \varepsilon^{N - \frac{(N-s)q}{2}} C \int_0^{1/\varepsilon} \frac{1}{(1 + \varrho^2)^{\frac{(N-s)q}{2}}} \varrho^{N-1} d\varrho \\
&= \varepsilon^{N - \frac{(N-s)q}{2}} C \left( \int_0^1 \frac{1}{(1 + \varrho^2)^{\frac{(N-s)q}{2}}} \varrho^{N-1} d\varrho + \int_1^{1/\varepsilon} \frac{1}{(1 + \varrho^2)^{\frac{(N-s)q}{2}}} \varrho^{N-1} d\varrho \right) \\
&\leq \varepsilon^{N - \frac{(N-s)q}{2}} \left( C + C \int_1^{1/\varepsilon} \frac{1}{\varrho^{(N-s)q - N + 1}} d\varrho \right) \\
&\leq \varepsilon^{N - \frac{(N-s)q}{2}} \left( C + C\varepsilon^{(N-s)q - N} \right) \leq C\varepsilon^{\frac{(N-s)q}{2}},
\end{aligned}$$

for  $\varepsilon > 0$  small enough. Since  $\|\phi u_\varepsilon\|_{L^{2_s^*}}$  converges to a positive constant, the assertion follows.

The following result will be crucial for the proof of our main result

**Lemma 2.6** (MP energy bound). *Let  $\lambda = 1$  and let (f1)-(f2)-(f3)' hold. Then  $c < \frac{s}{2N}(k_s \mathbb{S}(s, N))^{N/s}$ .*

*Proof.* By definition of  $c$ , it is sufficient to prove that there exists  $\varepsilon > 0$  small enough that

$$\sup_{t \geq 0} J(t\eta_\varepsilon) < \frac{s}{2N}(k_s \mathbb{S}(s, N))^{N/s}, \quad J = J_1.$$

By definition of  $J$ , we have

$$J(t\eta_\varepsilon) = \frac{t^2}{2}\|\eta_\varepsilon\|^2 - \int_{\mathbb{R}^N} K(x)F(t\eta_\varepsilon(x,0))dx - \frac{t^{2_s^*}}{2_s^*}.$$

By the assumptions of  $f$ , there exist  $q \in (2, 2_s^*)$  and  $C_0 > 0$  with  $F(s) \geq C_0 s^q$  for any  $s \in \mathbb{R}^+$ . Then

$$J(t\eta_\varepsilon) \leq \psi(t), \quad \psi(t) = \frac{t^2}{2}\|\eta_\varepsilon\|^2 - C_0 t^q \int_{\mathbb{R}^N} |\eta_\varepsilon(x,0)|^q dx - \frac{t^{2_s^*}}{2_s^*}.$$

Since  $\psi(t) \rightarrow -\infty$  as  $t \rightarrow +\infty$ , we have  $\sup\{\psi(t) : t \geq 0\} = \psi(t_\varepsilon)$  for some  $t_\varepsilon > 0$ , so that

$$\|\eta_\varepsilon\|^2 - C_0 q t_\varepsilon^{q-2} \int_{\mathbb{R}^N} |\eta_\varepsilon(x,0)|^q dx = t_\varepsilon^{2_s^*-2},$$

which yields  $\sigma_0 \leq t_\varepsilon \leq \|\eta_\varepsilon\|^{\frac{2}{2_s^*-2}} \leq K_0$  for some  $\sigma_0, K_0 > 0$  independent of  $\varepsilon$ , in view of Lemma 2.4 and the above equality. Since the map

$$[0, \|\eta_\varepsilon\|^{\frac{2}{2_s^*-2}}] \ni t \mapsto \frac{t^2}{2}\|\eta_\varepsilon\|^2 - \frac{t^{2_s^*}}{2_s^*},$$

increases, we get for some universal constant  $C > 0$ ,

$$\begin{aligned} \sup_{\mathbb{R}^+} \psi &\leq \frac{s}{2N} \left( \|\eta_\varepsilon\|_{X^s}^2 + \int_{\mathbb{R}^N} V(x)\eta_\varepsilon(x,0)^2 dx \right)^{N/s} - C_0 C \|\eta_\varepsilon(x,0)\|_{L^q}^q \\ &\leq \frac{s}{2N} \left( k_s \mathbb{S}(s, N) + \mathcal{O}(\varepsilon^{N-s}) + \int_{\mathbb{R}^N} V(x)\eta_\varepsilon(x,0)^2 dx \right)^{N/s} - C_0 C \|\eta_\varepsilon(x,0)\|_{L^q}^q. \end{aligned}$$

Now, by the elementary inequality  $(a+b)^\alpha \leq a^\alpha + \alpha(a+b)^{\alpha-1}b$ ,  $\alpha \geq 1$  and  $a, b > 0$ , we get by (2.25)

$$\begin{aligned} \sup_{\mathbb{R}^+} \psi &\leq \frac{s}{2N} (k_s \mathbb{S}(s, N))^{N/s} + \mathcal{O}(\varepsilon^{N-s}) + C \int_{\mathbb{R}^N} V(x)\eta_\varepsilon(x,0)^2 dx - C_0 C \|\eta_\varepsilon(x,0)\|_{L^q}^q \\ &\leq \frac{s}{2N} (k_s \mathbb{S}(s, N))^{N/s} + \mathcal{O}(\varepsilon^{N-s}) + C \|\eta_\varepsilon(x,0)\|_{L^2}^2 - C_0 C \|\eta_\varepsilon(x,0)\|_{L^q}^q \end{aligned}$$

• In the case  $N > 2s$ , by means of (2.26) and (2.27), we get

$$\sup_{\mathbb{R}^+} \psi \leq \frac{s}{2N} (k_s \mathbb{S}(s, N))^{N/s} + \mathcal{O}(\varepsilon^{N-s}) + \mathcal{O}(\varepsilon^s) - C_0 \mathcal{O}(\varepsilon^{\frac{2N-(N-s)q}{2}}).$$

Since  $\frac{2N-(N-s)q}{2} < s < N-s$ , we get the conclusion for  $\varepsilon$  sufficiently small.

• If  $N = 2s$  and  $2 < q < 2_s^* = 4$ , by (2.26) and (2.27), we get

$$\sup_{\mathbb{R}^+} \psi \leq \frac{s}{2N} (k_s \mathbb{S}(s, N))^{N/s} + \mathcal{O}(\varepsilon^s(1 + \log(\varepsilon^{-1}))) - C_0 \mathcal{O}(\varepsilon^{\frac{2N-sq}{2}}).$$

Since it holds

$$\lim_{\varepsilon \rightarrow 0} \frac{\varepsilon^{\frac{2N-sq}{2}}}{\varepsilon^s \log(\varepsilon^{-1})} = +\infty,$$

again can we get the conclusion, for  $\varepsilon$  sufficiently small.

• If  $s < N < 2s$  and  $\frac{N}{N-s} < q < 2_s^*$ , by (2.26) and (2.27), we get

$$\sup_{\mathbb{R}^+} \psi \leq \frac{s}{2N} (k_s \mathbb{S}(s, N))^{N/s} + \mathcal{O}(\varepsilon^{N-s}) - C_0 \mathcal{O}(\varepsilon^{\frac{2N-(N-s)q}{2}}).$$

Since  $\frac{2N-(N-s)q}{2} < N-s$  means  $q > \frac{2s}{N-s} (> 2)$ , we get the conclusion for  $\varepsilon$  sufficiently small.

• If  $s < N < 2s$  and  $2 < q < \frac{N}{N-s}$ , by (2.26) and (2.27), we get

$$\sup_{\mathbb{R}^+} \psi \leq \frac{s}{2N} (k_s \mathbb{S}(s, N))^{N/s} + \mathcal{O}(\varepsilon^{N-s}) - C_0 \mathcal{O}(\varepsilon^{\frac{(N-s)q}{2}}),$$

and for  $C_0 = \varepsilon^{-\vartheta}$  with  $\vartheta > \frac{(N-s)(q-2)}{2}$ , we get the conclusion. This concludes the proof.  $\square$

## 3. PROOF OF THEOREM 1.1 COMPLETED

In light of Lemma 2.2, there exists a *Cerami* sequence  $\{w_n\} \subset E$  for  $J = J_0$ . From Lemma 2.3 it follows that  $w_n^- \rightarrow 0$  in  $E$  as  $n \rightarrow \infty$  and that  $\{w_n\}$  is bounded and has a nonnegative weak limit  $w \in E$ . By Proposition 2.1, it follows that  $w$  is a weak nonnegative solution, to which a weak solution  $u \in H^{s/2}(\mathbb{R}^N)$  to (1.1) corresponds. We have  $u > 0$  if  $u \neq 0$ . In fact, if  $u(x_0) = 0$  for some  $x_0 \in \mathbb{R}^N$ , then  $(-\Delta)^{s/2}u(x_0) = 0$  and by the representation formula [22]

$$(-\Delta)^{s/2}u(x) = -\frac{c(N, s/2)}{2} \int_{\mathbb{R}^N} \frac{u(x+y) + u(x-y) - 2u(x)}{|x-y|^{N+s}} dy,$$

one obtains that, at  $x_0$ , that

$$\int_{\mathbb{R}^N} \frac{u(x_0+y) + u(x_0-y)}{|x_0-y|^{N+s}} dy = 0,$$

yielding  $u = 0$ , a contradiction. Let us prove that, indeed,  $u \neq 0$ . We prove that  $w = E_s(u) \neq 0$ . In fact,  $(w_n)$  converges to  $w$  strongly in  $E$ . Indeed, up to a subsequence,  $w_n \rightarrow w$  in  $E$  as  $n \rightarrow \infty$ , and since  $J'(w_n)(w_n) = o_n(1)$ , we have, again by virtue of Proposition 2.1,

$$\begin{aligned} \lim_{n \rightarrow \infty} \|w_n\|^2 &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x) f(w_n(x, 0)) w_n(x, 0) dx \\ &= \int_{\mathbb{R}^N} K(x) f(w(x, 0)) w(x, 0) dx = \|w\|^2, \end{aligned}$$

that is,  $w_n \rightarrow w$  in  $E$ . Hence  $J(w) = c$  and  $J'(w) = 0$ , this implies that  $w \neq 0$ .  $\square$

## 4. PROOF OF THEOREM 1.2 COMPLETED

In light of Lemma 2.2, there exists a *Cerami* sequence  $\{w_n\} \subset E$  for  $J = J_1$ . From Lemma 2.3 it follows that  $w_n^- \rightarrow 0$  in  $E$  as  $n \rightarrow \infty$  and that  $\{w_n\}$  is bounded and has a nonnegative weak limit  $w \in E$ . By Proposition 2.1, it follows that  $w$  is a weak nonnegative solution, to which a weak solution  $u \in H^{s/2}(\mathbb{R}^N)$  to (1.1) corresponds. We have  $u > 0$  if  $u \neq 0$ , arguing as in Section 3. Let us prove that, indeed,  $u \neq 0$ . We prove that  $w = E_s(u) \neq 0$ . By virtue of (2.20), we have

$$\begin{aligned} \frac{1}{2} \|w_n\|^2 - \int_{\mathbb{R}^N} K(x) F(w(x, 0)) dx - \frac{1}{2_s^*} \int_{\mathbb{R}^N} w_n^+(x, 0)^{2_s^*} dx &= c + o_n(1), \\ \|w_n\|^2 - \int_{\mathbb{R}^N} K(x) f(w(x, 0)) w(x, 0) dx - \int_{\mathbb{R}^N} w_n^+(x, 0)^{2_s^*} dx &= o_n(1). \end{aligned}$$

Suppose, by contradiction, that  $w \equiv 0$ . Then, we entail

$$\left( \frac{1}{2} - \frac{1}{2_s^*} \right) \|w_n\|^2 = c + o_n(1),$$

which combined with  $\|w_n\|^2 = \|w_n(x, 0)\|_{2_s^*}^{2_s^*} + o_n(1)$  as  $n \rightarrow \infty$  and the Sobolev inequality

$$\|w_n\|^2 \geq \int_{\mathbb{R}_+^{N+1}} k_s y^{1-s} |\nabla w|^2 dx dy \geq k_s \mathbb{S}(s, N) \|w_n(x, 0)\|_{2_s^*}^2$$

implying

$$c = \lim_n J(w_n) = \left( \frac{1}{2} - \frac{1}{2_s^*} \right) \lim_n \|w_n\|^2 \geq \frac{s}{2N} (k_s \mathbb{S}(s, N))^{N/s}.$$

This contradicts Lemma 2.6. Hence  $w \neq 0$  and the proof is complete.  $\square$

## REFERENCES

- [1] C.O. Alves, M. Montenegro, M.A.S. Souto, *Existence of solution for three classes of elliptic problems in  $\mathbb{R}^N$  with zero mass*, J. Differential Equations **252** (2012), 5735–5750. [1](#)
- [2] C.O. Alves, M.A.S. Souto, *Existence of solutions for a class of elliptic equations in  $\mathbb{R}^N$  with vanishing potentials*, J. Differential Equations **252** (2012), 5555–5568. [1](#)
- [3] C.O. Alves, M.A.S. Souto, *Existence of solutions for a class of nonlinear Schrödinger equations with potential vanishing at infinity*, J. Differential Equations **254** (2013), 1977–1991. [1](#), [4](#)
- [4] A. Ambrosetti, V. Felli, A. Malchiodi, *Ground states of nonlinear Schrödinger equations with potentials vanishing at infinity*, J. Eur. Math. Soc. (JEMS) **7** (2005) 117–144.
- [5] A. Ambrosetti, P. H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Funct. Anal. **14** (1973), 349–381.
- [6] A. Ambrosetti, J. Garcia Azorero, I. Peral, *Elliptic variational problems in  $\mathbb{R}^N$  with critical growth*, J. Differential Equations **168** (2000), 10–32. [2](#)
- [7] A. Ambrosetti, Z.-Q. Wang, *Nonlinear Schrödinger equations with vanishing and decaying potentials*, Differential Integral Equations **18** (2005), 1321–1332. [1](#)
- [8] M. Badiale, L. Pisani, S. Rolando, *Sum of weighted Lebesgue spaces and nonlinear elliptic equations*, NoDEA Nonlinear Differential Equations Appl. **18** (2011), 369–405. [1](#)
- [9] B. Barrios, E. Colorado, A. de Pablo, U. Sánchez, *On some critical problems for the fractional Laplacian operator*, J. Differential Equations **252** (2012), 613–6162. [2](#), [9](#)
- [10] V. Benci, C.R. Grisanti, A.M. Micheletti, *Existence of solutions for the nonlinear Schrödinger equation with  $V(\infty) = 0$* , Progr. Nonlinear Differential Equations Appl. **66** (2005), 53–65. [1](#)
- [11] H. Berestycki, P.L. Lions, *Nonlinear scalar field equations. I. Existence of a ground state*, Arch. Ration. Mech. Anal. **82** (1983), 313–346. [1](#), [6](#)
- [12] D. Bonheure, J. Van Schaftingen, *Ground states for nonlinear Schrödinger equation with potential vanishing at infinity*, Ann. Mat. Pura Appl. **189** (2010), 273–301. [7](#)
- [13] C. Brandle, E. Colorado, A. de Pablo, U. Sánchez, *A concave-convex elliptic problem involving the fractional laplacian*, Proc. Roy. Soc. Edinburgh Sect A **143** (2013), 39–71. [2](#), [5](#), [8](#), [9](#)
- [14] H. Brezis, L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), 437–477. [2](#), [9](#)
- [15] X. Cabré, J.G. Tan, *Positive solutions of nonlinear problems involving the square root of the Laplacian*, Adv. Math. **224** (2010), 2052–2093. [2](#)
- [16] L. Caffarelli, L. Silvestre, *An extension problems related to the fractional Laplacian*, Comm. PDE **32** (2007), 1245–1260. [2](#)
- [17] A. Capella, J. Dávila, L. Dupaigne, Y. Sire, *Regularity of radial extremal solutions for some non local semilinear equations*, Comm. PDE **36** (2011), 1353–1384. [2](#)
- [18] G. Cerami, *Un criterio de esistenza per i punti critici su varietà ilimitate*, Istit. Lombardo Accad. Sc. Lett. Rend. **112** (1978), 332–336. [7](#)
- [19] X. Chang, Z.-Q. Wang, *Ground state of scalar field equations involving a fractional laplacian with general nonlinearity*, Nonlinearity **26** (2013), 479–494. [1](#)
- [20] D.G. Costa, C.A. Magalhaes, *Variational elliptic problems which are nonquadratic at infinity*, Nonlinear Anal. **23** (1994), 1401–1412. [1](#)
- [21] A. Cotsiolis, N.K. Tavoularis, *Best constants for Sobolev inequalities for higher order fractional derivatives*, J. Math. Anal. Appl. **295** (2004), 225–236. [9](#)
- [22] E. Di Nezza, G. Palatucci, E. Valdinoci, *Hitchhiker’s guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136** (2012), 521–573. [1](#), [12](#)
- [23] M.M. Fall, T. Weth, *Nonexistence results for a class of fractional elliptic boundary value problems*, J. Funct. Anal. **263** (2012), 2205–2227. [2](#)
- [24] M. Ghimenti, A.M. Micheletti, *Existence of minimal nodal solutions for the nonlinear Schrödinger equation with  $V(\infty) = 0$* , Adv. Diff. Equations **11** (2006), 1375–1396. [1](#)
- [25] L. Jeanjean, *On the existence of bounded PalaisSmale sequences and application to a LandesmanLazer type problem set on  $\mathbb{R}^N$* , Proc. Roy. Soc. Edinburgh Sect. A **129** (1999), 787–809. [1](#)
- [26] T. Jin, Y. Li, J. Xiong, *On a fractional Nirenberg problem, part I: blow up analysis and compactness of solutions*, J. European Math. Soc. **16** (2014), 1111–1171. [2](#)
- [27] Y. Liu, Z.-Q. Wang, J. Zhang, *Ground states of nonlinear Schrödinger equations with potentials*, Ann. Inst. H. Poincaré Anal.Non Linéaire **23** (2006), 829–837. [1](#)
- [28] Z. Liu, Z.-Q. Wang, *On the Ambrosetti-Rabinowitz superlinear condition*, Adv. Nonlinear Stud. **4** (2004), 563–574. [1](#)
- [29] O.H. Miyagaki, *On a class of semilinear elliptic problems in  $\mathbb{R}^N$  with critical growth*, Nonlinear Analysis **29** (1997), 773–781.

- [30] S. Secchi, *Ground state solutions for nonlinear fractional Schrödinger equations in  $\mathbb{R}^N$* , J. Math. Phys. **54** (2013), 031501-17 pages. [1](#)
- [31] R. Servadei, E. Valdinoci, *A Brezis-Nirenberg result for non-local critical equations in low dimension*, Comm. Pure Appl. Anal. **12** (2013), 2445–2464. [2](#), [9](#)
- [32] R. Servadei, E. Valdinoci, *On the spectrum of two different fractional operators*, Proc. Roy. Soc. Edinburgh Sect. A **144** (2014), 831–855. [2](#)
- [33] X. Shang, J. Zhang, Y. Yang, *On fractional Schrödinger equation in  $\mathbb{R}^N$  with critical growth*, J. Math. Phys. **54** (2013), 121502-19 pages. [2](#)
- [34] M. Schechter, W. Zou, *Superlinear problems*, Pacific J. Math. **214** (2004), 145–160. [1](#)
- [35] W. Strauss, *Existence of solitary waves in higher dimensions*, Comm. Math. Phys. **55** (1977), 149–162. [1](#)
- [36] G. Talenti, *Best constants in Sobolev inequality*, Ann. Mat. Pura Appl. **110** (1976), 353–372. [9](#)
- [37] J. Tan, *Positive solutions for non local elliptic problems*, Disc. Cont. Dyn. Syst. **33** (2013), 837–859. [2](#)
- [38] J. Tan, *The Brezis-Nirenberg type problem involving the square root of the laplacian*, Calc. Var. Partial Differential Equations **42** (2011), 21–41. [2](#), [9](#)
- [39] M. Willem, *Minimax theorems*, Birkhäuser, Boston, 1996. [2](#)

(J.M. do Ó)

DEPARTMENT OF MATHEMATICS  
FEDERAL UNIVERSITY OF PARAÍBA  
58051-900, JOÃO PESSOA-PB, BRAZIL  
*E-mail address:* [jmbo@pq.cnpq.br](mailto:jmbo@pq.cnpq.br)

(O. Miyagaki)

DEPARTMENT OF MATHEMATICS  
FEDERAL UNIVERSITY OF JUIZ DE FORA  
36036-330 JUIZ DE FORA, MINAS GERAIS, BRAZIL  
*E-mail address:* [ohmiyagaki@gmail.com](mailto:ohmiyagaki@gmail.com)

(M. Squassina)

DIPARTIMENTO DI INFORMATICA  
UNIVERSITÀ DEGLI STUDI DI VERONA  
CÁ VIGNAL 2, STRADA LE GRAZIE 15, I-37134 VERONA, ITALY  
*E-mail address:* [marco.squassina@univr.it](mailto:marco.squassina@univr.it)