

Graphs with no induced five-vertex path or antipath

Maria Chudnovsky* Louis Esperet[†] Laetitia Lemoine[‡]
Peter Maceli[§] Frédéric Maffray[¶] Irena Penev^{||}

August 16, 2018

Abstract

We prove that a graph G contains no induced 5-vertex path and no induced complement of a 5-vertex path if and only if G is obtained from 5-cycles and split graphs by repeatedly applying the following operations: substitution, split unification, and split unification in the complement, where split unification is a new class-preserving operation introduced here.

*Princeton University, Princeton, NJ 08544, USA E-mail: mchudnov@math.princeton.edu. Most of this work was conducted while the author was at Columbia University. Partially supported by NSF grants DMS-1001091 and IIS-1117631.

[†]CNRS, Laboratoire G-SCOP, University of Grenoble, France. E-mail: louis.esperet@g-scop.grenoble-inp.fr

[‡]Laboratoire G-SCOP, University of Grenoble, France. E-mail: lae.lemoine@gmail.com

[§]Wesleyan University, Middletown CT 06459, USA. Most of this work was conducted while the author was at Columbia University. E-mail: pmaceli@wesleyan.edu.

[¶]CNRS, Laboratoire G-SCOP, University of Grenoble, France. E-mail: frederic.maffray@g-scop.grenoble-inp.fr.

^{||}Department of Applied Mathematics and Computer Science, Technical University of Denmark, Lyngby, Denmark. Email: ipen@dtu.dk. A part of this work was conducted while the author was at Université de Lyon, LIP, ENS de Lyon, Lyon, France. Partially supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR), and by the ERC Advanced Grant GRACOL, project number 320812.

Authors Esperet, Maffray, Lemoine, and Penev were partially supported by ANR project Stint under reference ANR-13-BS02-0007.

1 Introduction

All graphs in this paper are finite and simple. For fixed $n \geq 1$, let P_n denote the path on n vertices, and for $n \geq 3$, let C_n denote the cycle on n vertices. The graph C_5 is also called a *pentagon*. The complement of a graph G is denoted by \overline{G} . Given graphs G and F , we say that G is *F-free* if no induced subgraph of G is isomorphic to F . Given a family \mathcal{F} of graphs, we say that a graph G is *\mathcal{F} -free* provided that G is F -free for all $F \in \mathcal{F}$.

A graph G is *perfect* if for every induced subgraph H of G the chromatic number of H is equal to the maximum clique size in H . Chudnovsky, Robertson, Seymour, and Thomas [4] solved the long-standing and famous problem known as the Strong Perfect Graph Conjecture by proving that a graph G is perfect if and only if neither G nor \overline{G} contains an induced odd cycle of length at least five.

There are various instances of the collection \mathcal{F} such that \mathcal{F} -free graphs are highly structured in a way that can be described precisely; this fact is interesting in itself, and sometimes, it is also useful for solving various optimization problems on \mathcal{F} -free graphs. The goal of this paper is to understand the structure of $\{P_5, \overline{P_5}\}$ -free graphs. The motivation for this is manifold:

- The class of $\{P_5, \overline{P_5}\}$ -free graphs contains all *cographs* and all *split graphs*. Cographs are also known as P_4 -free graphs; their structure is very well understood (see for example [1, 6]). Split graphs are graphs whose vertex-set can be partitioned into a clique and a stable set, and it is known [8, 9] that they are exactly the $\{C_4, \overline{C_4}, C_5\}$ -free graphs.
- The class of $\{P_5, \overline{P_5}\}$ -free graphs has already been the object of much research. Fouquet [10] proved that the study of this class can be reduced in a certain way (which we recall in more detail below) to the study of $\{P_5, \overline{P_5}, C_5\}$ -free graphs. Moreover, it follows from the results of Chvátal, Hoàng, Mahadev, and de Werra [5], Giakoumakis and Rusu [11], and Hoàng and Lazzarato [13] that several optimization problems can be solved in polynomial time in the class of $\{P_5, \overline{P_5}\}$ -free graphs. However, none of these results gives (or attempts to give) a description of the structure of such graphs.
- The class of $\{P_5, \overline{P_5}, C_5\}$ -free graphs is a subclass of the class of perfect graphs, and it is interesting to have a structure theorem for this subclass since so far, no structure theorem has been proved for the class of all perfect graphs.

Before presenting our results, we need to introduce some notation and definitions. For a graph G , we denote by $V(G)$ its vertex-set and by $E(G)$ its edge-set. Given a set $S \subseteq V(G)$, let $N(S)$ be the set of vertices in $V(G) \setminus S$ that have a neighbor in S . Let $G[S]$ denote the subgraph of G induced by S , and let $G \setminus S$ denote the induced subgraph $G[V(G) \setminus S]$. We say that a vertex v in $V(G) \setminus S$ is *complete* to S if v is adjacent to every vertex of S , and that v is *anticomplete* to S if v has no neighbor in S . A vertex of $V(G) \setminus S$ that is neither complete nor anticomplete to S is *mixed* on S . Given two disjoint sets $S, T \subseteq V(G)$, we say that S is *complete* to T when every vertex of S is complete to T , and we say that S is *anticomplete* to T when every vertex of S is anticomplete to T .

An *anticomponent* of a set $S \subseteq V(G)$ is any subset of S that induces a compo-

ment of the graph $\overline{G}[S]$. A graph G is *anticonnected* if \overline{G} is connected.

A *homogeneous set* is a non-empty set $S \subseteq V(G)$ such that every vertex of $V(G) \setminus S$ is either complete or anticomplete to S . A homogeneous set S is *proper* when $|S| \geq 2$ and $S \neq V(G)$. Let G be a graph that admits a proper homogeneous set S , and let s be any vertex in S . We can decompose G into the two graphs $G[S]$ and $G \setminus (S \setminus s)$. Since S is a homogeneous set, we see that up to isomorphism, the latter graph is the same whatever the choice of s . Moreover, both $G[S]$ and $G \setminus (S \setminus s)$ are induced subgraphs of G . The reverse operation, known as *substitution*, can be defined as follows. Let G and H be two vertex-disjoint graphs and let x be a vertex in G . Make a graph G' with vertex-set $V(G \setminus x) \cup V(H)$, taking the union of the two graphs $G \setminus x$ and H and adding all edges between $V(H)$ and the neighborhood of x in G . Clearly, in G' , the set $V(H)$ is a homogeneous set, $H = G'[V(H)]$, and G is isomorphic to an induced subgraph of G' . Moreover $V(H)$ is a proper homogeneous set if both G and H have at least two vertices. Thus, a graph G is obtained by substitution from smaller graphs if and only if G contains a proper homogeneous set. A graph is *prime* if it has no proper homogeneous set.

The following result about the structure of $\{P_5, \overline{P_5}\}$ -free graphs was proved by Fouquet in [10].

Theorem 1.1 ([10]) *Every $\{P_5, \overline{P_5}\}$ -free graph G satisfies one of the following properties:*

- G contains a proper homogeneous set;
- G is isomorphic to C_5 ;
- G is C_5 -free.

Theorem 1.1 immediately implies that every $\{P_5, \overline{P_5}, C_5\}$ -free graph can be obtained by substitution starting from $\{P_5, \overline{P_5}, C_5\}$ -free graphs and pentagons. Furthermore, it is easy to check that every graph obtained by substitution starting from $\{P_5, \overline{P_5}, C_5\}$ -free graphs and pentagons is $\{P_5, \overline{P_5}\}$ -free. We remark that the Strong Perfect Graph Theorem [4] implies that a $\{P_5, \overline{P_5}\}$ -free graph is perfect if and only if it is C_5 -free. Thus, every $\{P_5, \overline{P_5}\}$ -free graph can be obtained by substitution starting from $\{P_5, \overline{P_5}\}$ -free perfect graphs and pentagons. In view of this, the bulk of this paper focuses on prime $\{P_5, \overline{P_5}, C_5\}$ -free graphs (equivalently: prime $\{P_5, \overline{P_5}\}$ -free perfect graphs).

Our first result, Theorem 2.3, states that every prime $\{P_5, \overline{P_5}, C_5\}$ -free graph that is not split admits a particular kind of partition. Our second result, Theorem 3.1, states that every prime $\{P_5, \overline{P_5}, C_5\}$ -free graph that is not split admits a new kind of decomposition, which we call a “split divide” (see section 3). Next, we reverse the split graph divide decomposition and turn it into a composition that preserves the property of being $\{P_5, \overline{P_5}, C_5\}$ -free. We call this composition “split unification” (see section 4). Finally, combining our results with Theorem 1.1, we prove that every $\{P_5, \overline{P_5}\}$ -free graph is obtained by repeatedly applying substitution, split graph unification, and split graph unification in the complement starting from split graphs and pentagons, and furthermore, we prove that every graph obtained in this way is $\{P_5, \overline{P_5}\}$ -free (see Theorems 5.1 and 5.2).

This paper results from the merging of the two (unpublished) manuscripts [3] and [7] on the same subject; it combines the proofs and results from these two manuscripts so as to present them in the most succinct way.

2 Prime $\{P_5, \overline{P_5}, C_5\}$ -free graphs

Recall that a graph is *split* if its vertex-set can be partitioned into a stable set and a clique. Földes and Hammer [8, 9] gave the following characterization of split graphs (a short proof is given in [12, p. 151]).

Theorem 2.1 ([8, 9]) *A graph is split if and only if it is $\{C_4, \overline{C_4}, C_5\}$ -free.*

Lemma 2.2 *In a $\{P_5, \overline{P_5}, C_5\}$ -free graph G , let A and B be non-empty and disjoint subsets of $V(G)$, and let t be a vertex in $V(G) \setminus (A \cup B)$ such that:*

- t is anticomplete to A and complete to B ,
- every vertex in B has a neighbor in A , and
- A is connected.

Then some vertex of A is complete to B .

Proof. Pick a vertex a in A with the maximum number of neighbors in B . Suppose that a has a non-neighbor y in B . We know that y has a neighbor a' in A . Since A is connected, there is a path $P = a_0 \cdots a_k$ in $G[A]$ with $k \geq 1$, $a_0 = a'$ and $a_k = a$. Choose a' such that k is minimal. So P is chordless and y has no neighbor in $P \setminus \{a_0\}$. Then $k = 1$, for otherwise t, y, a_0, a_1, a_2 induce a P_5 . By the choice of a , since y is adjacent to a' and not to a , there is a vertex z in B adjacent to a and not to a' . Then a, z, t, y, a' induce a C_5 or $\overline{P_5}$ (depending on the pair y, z), a contradiction. Thus a is complete to B . \square

We say that a set, or a graph, is *big* if it contains at least two vertices.

Theorem 2.3 *Let G be a prime $\{P_5, \overline{P_5}, C_5\}$ -free graph that contains a $\overline{C_4}$. Then there are pairwise disjoint subsets $X_0, X_1, \dots, X_m, Y_0, Y_1, \dots, Y_m$, with $m \geq 2$, whose union is equal to $V(G)$, such that the following properties hold, where $X = X_0 \cup X_1 \cup \dots \cup X_m$ and $Y = Y_0 \cup Y_1 \cup \dots \cup Y_m$:*

- (i) *For each $i \in \{1, \dots, m\}$, X_i is connected, $|X_i| \geq 2$, X_0 is a (possibly empty) stable set, and X_0, X_1, \dots, X_m are pairwise anticomplete to each other.*
- (ii) *For each $i \in \{1, \dots, m\}$, $Y_i \neq \emptyset$, every vertex of Y_i is mixed on X_i and complete to $X \setminus (X_i \cup X_0)$, and Y_0 is complete to $X \setminus X_0$.*
- (iii) *Y_0, Y_1, \dots, Y_m are pairwise complete to each other. (So each anticomponent of Y is included in some Y_i with $i \in \{0, \dots, m\}$.)*
- (iv) *No vertex of $X \setminus X_0$ is mixed on any anticomponent of Y .*
- (v) *For each $i \in \{1, \dots, m\}$, X_i contains a vertex that is complete to Y .*

- (vi) Every vertex of X_0 is mixed on at most one anticomponent of Y .
- (vii) For every big anticomponent Z of Y , the set X_Z of vertices of X_0 that are mixed on Z is not empty. Moreover, if Z and Z' are any two distinct big anticomponents of Y , then $X_Z \cap X_{Z'} = \emptyset$.
- (viii) Each big anticomponent Z of Y contains a vertex that is anticomplete to X_Z .
- (ix) If Y is not a clique, there is a big anticomponent Z of Y such that X_Z is anticomplete to all big anticomponents of $Y \setminus Z$.

Proof. Since G contains a $\overline{C_4}$, there is a subset X of $V(G)$ such that $G[X]$ has at least two big components. We choose X maximal with this property. Let X_1, \dots, X_m ($m \geq 2$) be the vertex-sets of the big components of $G[X]$, and let $X_0 = X \setminus (X_1 \cup \dots \cup X_m)$. So (i) holds. Let $Y = V(G) \setminus X$. We claim that:

$$\text{For every } y \in Y \text{ and } i \in \{1, \dots, m\}, y \text{ has a neighbor in } X_i. \quad (1)$$

Proof. If y has no neighbour in X_i , then $X \cup \{y\}$ induces a subgraph of G with at least two big components (one of which is X_i), which contradicts the maximality of X . Thus (1) holds.

$$\text{For every vertex } y \in Y, \text{ there is at most one integer } i \text{ in } \{1, \dots, m\} \text{ such that } y \text{ has a non-neighbor in } X_i. \quad (2)$$

Proof. Suppose that y has a non-neighbor in two distinct components X_i and X_j (with $1 \leq i, j \leq m$) of X . For each $h \in \{i, j\}$, y has a neighbor in X_h by (1), and since X_h is connected, there are adjacent vertices $u_h, v_h \in X_h$ such that y is adjacent to u_h and not to v_h . Then v_i, u_i, y, u_j, v_j induce a P_5 , a contradiction. Thus (2) holds.

An immediate consequence of Claims (1) and (2) is the following.

$$\begin{aligned} &\text{For every vertex } y \in Y, \text{ either } y \text{ is complete to } X \setminus X_0, \\ &\text{or there is a unique integer } i \in \{1, \dots, m\} \text{ such that } y \text{ is} \\ &\text{complete to } X \setminus (X_i \cup X_0) \text{ and } y \text{ is mixed on } X_i. \end{aligned} \quad (3)$$

For each $i \in \{1, \dots, m\}$, let $Y_i = \{y \in Y \mid y \text{ is mixed on } X_i\}$, and let $Y_0 = Y \setminus (Y_1 \cup \dots \cup Y_m)$. By (3), the sets Y_0, Y_1, \dots, Y_m are pairwise disjoint and their union is Y . For each $i \in \{1, \dots, m\}$, since G is prime, X_i is not a homogeneous set, so there exists a vertex in $V(G) \setminus X_i$ that is mixed on X_i ; by (i), any such vertex is in Y , and so $Y_i \neq \emptyset$. Thus (ii) holds.

Now we prove (iii). Suppose that Y_i is not complete to Y_j for some distinct $i, j \in \{0, \dots, m\}$. Let $y \in Y_i$ and $z \in Y_j$ be non-adjacent. Up to symmetry we may assume that $i \neq 0$, say $i = 1$. Since X_1 is connected, there are adjacent vertices u_1 and v_1 in X_1 such that y is adjacent to u_1 and not to v_1 . By (ii), z is complete to $\{u_1, v_1\}$. Furthermore, by (ii), Y_1 is complete to X_2 , and every vertex in Y_j has a neighbor in X_2 ; thus, there exists a vertex $x_2 \in X_2$ such that x_2 is adjacent to both y and z . By (i), x_2 is non-adjacent to u_1 and v_1 . But now z, x_2, y, u_1, v_1 induce a $\overline{P_5}$, a contradiction. So the first sentence of (iii) holds. The second sentence is an immediate consequence of the first. Thus (iii) holds.

Now we prove (iv). Suppose on the contrary, and up to symmetry, that a vertex x in X_1 is mixed on some anticomponent Z of Y . Since Z is anticonnected, there are non-adjacent vertices $y, z \in Z$ such that x is adjacent to y and not to z . By (ii), z has a neighbor u in X_1 , so $z \in Y_1$. Since X_1 is connected, there is a path $u_0 \cdots u_k$ in $G[X_1]$ with $u_0 = u$, $u_k = x$ and $k \geq 1$. Choose u such that k is minimal. By (ii), y has a neighbor x_2 in X_2 , and since $z \in Y_1$, z is adjacent to x_2 . If $k = 1$, then x, y, z, u, x_2 induce a C_5 or \overline{P}_5 (depending on the pair y, u). So $k \geq 2$. The minimality of k implies that z is not adjacent to u_1 or u_2 , and u is not adjacent to u_2 . Then x_2, z, u, u_1, u_2 induce a P_5 , a contradiction.

Now we prove (v). We observe that by (i) and (ii), any vertex t from a big component of $X \setminus X_i$ is complete to Y_i and anticomplete to X_i , and so we can apply Lemma 2.2 to X_i, Y_i , and t . It follows that some vertex a of X_i is complete to Y_i . By (ii), X_i is complete to $Y \setminus Y_i$. Thus a is complete to Y .

Now we prove (vi). Suppose that a vertex x in X_0 is mixed on two anticomponents Z_1 and Z_2 of Y . For each $j \in \{1, 2\}$, since Z_j is anticonnected, there are non-adjacent vertices y_j and z_j in Z_j such that x is adjacent to y_j and not to z_j . Then y_1, z_1, x, z_2, y_2 induce a \overline{P}_5 , a contradiction.

Now we prove (vii). If Z is any big anticomponent of Y , then, since G is prime, Z is not a homogeneous set, and so there exists a vertex of $V(G) \setminus Z$ that is mixed on Z . The definition of Z and (iv) imply that any such vertex is in X_0 . So $X_Z \neq \emptyset$. The second sentence of (vii) follows directly from (vi).

Now we prove (viii). Let Z be a big anticomponent of Y . By (iii), Z is included in one of Y_0, Y_1, \dots, Y_m . By (ii) and (iv), some vertex t of $X \setminus X_0$ is complete to Z , and by (i) t is anticomplete to X_Z . Hence we can apply Lemma 2.2 to Z, X_Z and t in the complementary graph \overline{G} , and we obtain that some vertex in Z is complete (in \overline{G}) to X_Z .

Finally we prove (ix). Suppose that Y is not a clique, and choose a big anticomponent Z of Y that minimizes the number of big anticomponents of Y that are not anticomplete to X_Z . If this number is 1, then Z satisfies the desired property. So suppose that this number is at least 2, that is, there is a vertex $x \in X_Z$ and a big anticomponent Z' of $Y \setminus Z$ that contains a neighbor of x . There are non-adjacent vertices $y, z \in Z$ such that x is adjacent to y and not to z . By (vi), x is complete to Z' . Consider any $t \in X_{Z'}$; there are non-adjacent vertices $y', z' \in Z'$ such that t is adjacent to y' and not to z' . If t has any neighbor in Z , then, by (vi), t is complete to Z , and then z, x, t, z', y' induce a \overline{P}_5 , a contradiction. Since this holds for any $t \in X_{Z'}$, we obtain that $X_{Z'}$ is anticomplete to Z . Now the choice of Z implies that there is a third big anticomponent Z'' of Y (a big anticomponent of $Y \setminus (Z \cup Z')$) such that some vertex u of $X_{Z'}$ has a neighbor y'' in Z'' and X_Z is anticomplete to Z'' . There are non-adjacent vertices $a, b \in Z'$ such that u is adjacent to a and not to b . Then a, b, u, x, y'' induce a \overline{P}_5 , a contradiction. This completes the proof. \square

3 The split divide

A *split divide* of a graph G is a partition (A, B, C, L, T) of $V(G)$ such that:

- $|A| \geq 2$, A is complete to B and anticomplete to $C \cup T$, and some vertex of A is complete to L ;
- L is a non-empty clique, every vertex of L is mixed on A , and L is complete to $B \cup C$;
- $|C| \geq 2$, some vertex of C is complete to B , and no vertex of C is mixed on any anticomponent of B ;
- T is a (possibly empty) stable set and is anticomplete to C .




Figure 1: A split divide. Adjacency between sets is as follows: gray means complete, no edge means anticomplete, and a dashed edge means arbitrary adjacency. Gray border around a set means that the set is a clique, and white border means that the set is stable.

Note that the sets B and T may be empty. The split divide, illustrated in Figure 1, can be thought of as a relaxation of the homogeneous set decomposition: a set $X \subseteq V(G)$ is a homogeneous set in G if no vertex in $V(G) \setminus X$ is mixed on X ; in the case of the split divide, the set A is not homogeneous, but all the vertices that are mixed on A lie in the clique L , and adjacency between L and the rest of the graph is heavily restricted.

Theorem 3.1 *Let G be a prime $\{P_5, \overline{P_5}, C_5\}$ -free graph. Then either G is a split graph or G or \overline{G} admits a split divide.*

Proof. By Theorem 2.1 and up to complementation, we may assume that G contains a $\overline{C_4}$. Consequently G admits the structure described in Theorem 2.3, and we use it with the same notation. All items (i) to (ix) refer to Theorem 2.3.

Suppose that Y is a clique. Let $A = X_1$, $L = Y_1$, $B = Y \setminus Y_1$, $C = X_2 \cup \dots \cup X_m$ and $T = X_0$. Then (A, B, C, L, T) is a split divide of G ; this follows immediately from the definition of the partition $X_0, X_1, \dots, X_m, Y_0, Y_1, \dots, Y_m$, the fact that Y is a clique, and items (i)–(v).

Now suppose that Y is not a clique. We will show that \overline{G} admits a split divide. By (ix), we can choose a big anticomponent Z of Y such that X_Z is anticomplete to all big anticomponents of $Y \setminus Z$. By (vii), $X_Z \neq \emptyset$. By (iii), and up to relabeling, we may assume that $Z \subseteq Y_0 \cup Y_1$. Hence Z is complete to $X_2 \cup \dots \cup X_m$, and every vertex of $X_1 \cup (X_0 \setminus X_Z)$ is either complete or anticomplete to Z . Let K be the union of all anticomponents of Y of size 1. So K is a clique and is complete to $Y \setminus K$. Let:

$$A = Z;$$

$$\begin{aligned}
L &= X_Z; \\
B &= \{x \in X_1 \cup (X_0 \setminus X_Z) \mid x \text{ is anticomplete to } Z\}; \\
C' &= \{x \in X_1 \cup (X_0 \setminus X_Z) \mid x \text{ is complete to } Z\}; \\
T &= \{k \in K \mid k \text{ has a neighbor in } X_Z\}; \\
C &= X_2 \cup \dots \cup X_m \cup (Y \setminus (Z \cup T)) \cup C'.
\end{aligned}$$

We claim that:

$$L \text{ is anticomplete to } B \cup C. \quad (1)$$

Indeed, $X_Z (= L)$ is anticomplete to $X_1 \cup \dots \cup X_m$ because $X_Z \subseteq X_0$, and it is anticomplete to $X_0 \setminus X_Z$ because X_0 is a stable set. Moreover, X_Z is anticomplete to every (big) anticomponent of $(Y \setminus K) \setminus Z$, by the choice of Z , and it is anticomplete to $K \setminus T$ by the definition of T . Thus (1) holds.

$$\text{No vertex of } C \text{ is mixed on any component of } B. \quad (2)$$

For suppose that there is a vertex $c \in C$ and adjacent vertices $u, v \in B$ such that c is adjacent to u and not to v . Since X_0 is a stable set and is anticomplete to X_1 , we have $u, v \in \{x \in X_1 \mid x \text{ is anticomplete to } Z\}$. Since c is adjacent to u , we have $c \in (Y \setminus (Z \cup T)) \cup \{x \in X_1 \mid x \text{ is complete to } Z\}$. Pick any $x \in X_Z$ and any vertex $z \in Z$ adjacent to x . By (1), x is not adjacent to c . Then x, z, c, u, v induce a P_5 , a contradiction. Thus (2) holds.

$$T \text{ is complete to } C. \quad (3)$$

For suppose that there are non-adjacent vertices $t \in T$ and $c \in C$. Since K is complete to $Y \setminus K$ and $T \subseteq K$, we have that $c \notin Y \setminus (Z \cup T)$. Thus, $c \in X_2 \cup \dots \cup X_m \cup C'$. By (ii), Y_0 and Y_1 are complete to $X_2 \cup \dots \cup X_m$; since $Z \subseteq Y_0 \cup Y_1$, it follows that Z is complete to $X_2 \cup \dots \cup X_m$. Thus, $X_2 \cup \dots \cup X_m \cup C'$ is complete to Z , and so c is complete to Z . Further, since $X_2 \cup \dots \cup X_m \cup C' \subseteq X \setminus X_Z$ and X_Z is anticomplete to $X \setminus X_Z$ (because $X_Z \subseteq X_0$), we know that c is anticomplete to X_Z . By the definition of T , t has a neighbor x in X_Z . There are non-adjacent vertices $y, z \in Z$ such that x is adjacent to y and not to z . Since t and c are complete to Z , we see that t, c, y, z, x induce a \overline{P}_5 , a contradiction. Thus (3) holds.

Now we observe that:

- $|A| \geq 2$ because Z is big; A is anticomplete to B by the definition of B ; A is complete to $C \cup T$ by (ii); and some vertex of A is anticomplete to L by (viii).
- L is a non-empty stable set by (i) and (vii); every vertex of L is mixed on A by the definition of L ; and L is anticomplete to $B \cup C$ as shown in (1).
- $|C| \geq 2$ because $X_2 \subseteq C$; some vertex of C is anticomplete to B (every vertex of X_2 has this property); and no vertex of C is mixed on any component of B as proved in (2).
- T is a clique and is complete to C as proved in (3).

These observations mean that (A, B, C, L, T) is a split divide in \overline{G} . This completes the proof. \square

Let G be a graph that admits a split divide (A, B, C, L, T) as above, let a_0 be a vertex of A that is complete to L , and let c_0 be a vertex of C that is complete to B . Let $G_1 = G[A \cup B \cup \{c_0\} \cup L \cup T]$ and $G_2 = G[\{a_0\} \cup B \cup C \cup L \cup T]$. Then we consider that G is decomposed into the two graphs G_1 and G_2 . Note that G_1 and G_2 are induced subgraphs of G and each of them has strictly fewer vertices than G since $|A| \geq 2$ and $|C| \geq 2$.

4 Split unification

We can define a composition operation that “reverses” the split divide decomposition. Let A, B, C, L, T be pairwise disjoint sets, and assume that A and C are non-empty. Let a^*, c^* be distinct vertices such that $a^*, c^* \notin A \cup B \cup C \cup L \cup T$. Let G_1 be a graph with vertex-set $A \cup B \cup L \cup T \cup \{c^*\}$ and adjacency as follows:

- L is a (possibly empty) clique;
- T is a (possibly empty) stable set;
- A is complete to B and anticomplete to T ;
- Some vertex a_0 of A is complete to L ;
- c^* is complete to $B \cup L$ and anticomplete to $A \cup T$.

Let G_2 be a graph with vertex-set $B \cup C \cup L \cup T \cup \{a^*\}$ and adjacency as follows:

- $G_2[B \cup L \cup T] = G_1[B \cup L \cup T]$;
- T is anticomplete to C ;
- L is complete to $B \cup C$;
- a^* is complete to $B \cup L$ and anticomplete to $C \cup T$;
- Some vertex c_0 of C is complete to B , and no vertex of C is mixed on any anticomponent of B .




Figure 2: A composable pair.

Under these circumstances, we say that (G_1, G_2) is a *composable pair* (see Figure 2). The *split unification* of a composable pair (G_1, G_2) is the graph G with vertex-set $A \cup B \cup C \cup L \cup T$ such that:

- $G[A \cup B \cup L \cup T] = G_1 \setminus c^*$;
- $G[B \cup C \cup L \cup T] = G_2 \setminus a^*$;
- A is anticomplete to C in G .

Thus to obtain G from G_1 and G_2 , we “glue” G_1 and G_2 along their common induced subgraph $G_1[B \cup L \cup T] = G_2[B \cup L \cup T]$, where $L \cup T$ induces a split graph (hence the name of the operation).

We say that a graph G is *obtained by split unification* provided that there exists a composable pair (G_1, G_2) such that G is the split unification of (G_1, G_2) . We say that G is *obtained by split unification in the complement* provided that \overline{G} is obtained by split unification. We now prove that every graph that admits a split divide is obtained by split unification from smaller graphs.

Theorem 4.1 *If a graph G admits a split divide, then it is obtained from a composable pair of smaller graphs (each of them isomorphic to an induced subgraph of G) by split unification.*

Proof. Let G be a graph that admits a split divide. Let (A, B, C, L, T) be a split divide of G , let a_0 be a vertex of A that is complete to L , and let c_0 be a vertex of C that is complete to B . Let $G_1 = G[A \cup B \cup L \cup T \cup \{c_0\}]$. Since $|C| \geq 2$, we have $|V(G_1)| < |V(G)|$. Let $G_2 = G[B \cup C \cup L \cup T \cup \{a_0\}]$. Since $|A| \geq 2$, we have $|V(G_2)| < |V(G)|$. Now (G_1, G_2) is a composable pair, and G is obtained from it by split unification. \square

The split unification can be thought of as generalized substitution. Indeed, we obtain the graph G from G_1 and G_2 by first substituting $G_1[A]$ for a^* in G_2 , and then reconstructing the adjacency between A and L in G using the adjacency between A and L in G_1 . We include B, T and c^* in G_1 in order to ensure that split unification preserves the property of being $\{P_5, \overline{P}_5, C_5\}$ -free. In fact, we prove now something stronger than this: split unification preserves the (individual) properties of being P_5 -free, \overline{P}_5 -free, and C_5 -free.

Theorem 4.2 *Let (G_1, G_2) be a composable pair and let G be the split unification of (G_1, G_2) . Then, for each $H \in \{P_5, \overline{P}_5, C_5\}$, G is H -free if and only if both G_1 and G_2 are H -free.*

Proof. We use the same notation as in the definition of the split unification above. First suppose that G is H -free. Observe that G_1 is isomorphic to the induced subgraph $G[A \cup B \cup L \cup T \cup \{c_0\}]$, and G_2 is isomorphic to the induced subgraph $G[B \cup C \cup L \cup T \cup \{a_0\}]$. Hence G_1 and G_2 are H -free. Now suppose that G_1 and G_2 are H -free and that G contains an induced copy of H . Let W be a five-vertex subset of $V(G)$ such that $G[W] \simeq H$. We claim that W must contain two non-adjacent vertices b and c with $b \in W \cap B$ and $c \in W \cap C$. For suppose the contrary. Then $W \cap C$ is complete to $W \cap (L \cup B)$ and anticomplete

to $W \cap (A \cup T)$. If $|W \cap C| \geq 2$, then either $|W \cap C| \leq 4$, so $W \cap C$ is a proper homogeneous set in $G[W]$ (a contradiction since H is prime), or $W \subseteq C$, so W is isomorphic to an induced subgraph of G_2 (a contradiction since G_2 is H -free). So $|W \cap C| \leq 1$, and then W is isomorphic to an induced subgraph of G_1 (where c^* plays the role of the vertex in $W \cap C$ if there is such a vertex), a contradiction since G_1 is H -free. Therefore the claim holds. By a similar argument, W must contain two non-adjacent vertices a and ℓ with $a \in W \cap A$ and $\ell \in W \cap L$. Let w be the fifth vertex in W , so that $W = \{a, b, c, \ell, w\}$. By the definition of the split unification, a, b, ℓ, c induce a P_4 with edges $ab, b\ell, \ell c$. Consequently we must have one of the following two cases:

(i) W induces a P_5 or C_5 . So w is anticomplete to $\{b, \ell\}$ and has a neighbor in $\{a, c\}$. Since w is anticomplete to $\{b, \ell\}$, it cannot be in A, B, L or C , so it is in T . But then w should be anticomplete to $\{a, c\}$.

(ii) W induces a \overline{P}_5 . So w is adjacent to a and c and has exactly one neighbor in $\{b, \ell\}$. Since w is adjacent to a , it is not in $C \cup T$, and since it is adjacent to c , it is not in A . Moreover, since w is adjacent to exactly one of b and ℓ , it is not in L . So $w \in B$, and so it is adjacent to ℓ and, consequently, not to b . Hence b and w lie in the same anticomponent of B , and c is adjacent to exactly one of them, a contradiction (to the last axiom in the definition of a split unification). \square

5 The main theorem

In this section, we use Theorem 1.1 and the results of the preceding sections to prove Theorem 5.1, the main theorem of this paper.

Theorem 5.1 *A graph G is $\{P_5, \overline{P}_5\}$ -free if and only if at least one of the following holds:*

- G is a split graph;
- G is a pentagon;
- G is obtained by substitution from smaller $\{P_5, \overline{P}_5\}$ -free graphs;
- G or \overline{G} is obtained by split unification from smaller $\{P_5, \overline{P}_5\}$ -free graphs.

Proof. We first prove the “if” part. If G is a split graph or a pentagon, then it is clear that G is $\{P_5, \overline{P}_5\}$ -free. Since both P_5 and \overline{P}_5 are prime, we know that the class of $\{P_5, \overline{P}_5\}$ -free graphs is closed under substitution, and consequently, any graph obtained by substitution from smaller $\{P_5, \overline{P}_5\}$ -free graphs is $\{P_5, \overline{P}_5\}$ -free. Finally, if G or \overline{G} is obtained by split unification from smaller $\{P_5, \overline{P}_5\}$ -free graphs, then the fact that G is $\{P_5, \overline{P}_5\}$ -free follows from Theorem 4.2 and from the fact that the complement of a $\{P_5, \overline{P}_5\}$ -free graph is again $\{P_5, \overline{P}_5\}$ -free.

For the “only if” part, suppose that G is a $\{P_5, \overline{P}_5\}$ -free graph. We may assume that G is prime, for otherwise, G is obtained by substitution from smaller $\{P_5, \overline{P}_5\}$ -free graphs, and we are done. If some induced subgraph of G is isomorphic to the pentagon, then by Theorem 1.1, G is a pentagon, and again we are done. Thus we may assume that G is $\{P_5, \overline{P}_5, C_5\}$ -free. By Theorem 3.1,

we know that either G is a split graph, or one of G and \overline{G} admits a split divide. In the former case, we are done. In the latter case, Theorem 4.1 implies that G or \overline{G} is the split unification of a composable pair of smaller $\{P_5, \overline{P_5}, C_5\}$ -free graphs, and again we are done. \square

As an immediate corollary of Theorem 5.1, we have the following.

Theorem 5.2 *A graph is $\{P_5, \overline{P_5}\}$ -free if and only if it is obtained from pentagons and split graphs by repeated substitutions, split unifications, and split unifications in the complement.*

Finally, a proof analogous to the proof of Theorem 5.1 (but without the use of Theorem 1.1) yields the following result for $\{P_5, \overline{P_5}, C_5\}$ -free graphs.

Theorem 5.3 *A graph G is $\{P_5, \overline{P_5}, C_5\}$ -free if and only if at least one of the following holds:*

- G is a split graph;
- G is obtained by substitution from smaller $\{P_5, \overline{P_5}, C_5\}$ -free graphs;
- G or \overline{G} is obtained by split unification from smaller $\{P_5, \overline{P_5}, C_5\}$ -free graphs.

Acknowledgment

We would like to thank Ryan Hayward, James Nastos, Paul Seymour, and Yori Zwols for many useful discussions.

References

- [1] A. Brandstädt, V.B. Le, J.P. Spinrad. *Graph Classes: A Survey*. SIAM Monographs on Discrete Mathematics and Applications, 1999.
- [2] M. Chudnovsky, P. Maceli. Simplicial vertices in graphs with no induced four-edge path or four-edge antipath, and the H6-conjecture. *Journal of Graph Theory* 76 (2014) 249–261.
- [3] M. Chudnovsky, P. Maceli, I. Penev. Excluding four-edge paths and their complements. Manuscript, 2013. ArXiv: <http://arxiv.org/abs/1302.0405>.
- [4] M. Chudnovsky, N. Robertson, P. Seymour, R. Thomas. The strong perfect graph theorem. *Annals of Mathematics* 164 (2006) 51–229.
- [5] V. Chvátal, C.T. Hoàng, N.V.R. Mahadev, D. de Werra. Four classes of perfectly orderable graphs. *Journal of Graph Theory* 11 (1987) 481–495.
- [6] D.G. Corneil, Y. Perl, L.K. Stewart. A linear recognition algorithm for cographs. *SIAM Journal on Computing* 14 (1985) 926–934.
- [7] L. Esperet, L. Lemoine, F. Maffray. On $(P_5, \overline{P_5})$ -free graphs. Manuscript, 2013.
- [8] S. Földes, P.L. Hammer. Split graphs. *Congressus Numerantium XIX*, Utilitas Math. (1977) 311–315.
- [9] S. Földes, P.L. Hammer. Split graphs having Dilworth number two. *Canadian Journal of Mathematics* 29 (1977) 666–672.
- [10] J.-L. Fouquet. A decomposition for a class of $(P_5, \overline{P_5})$ -free graphs. *Discrete Mathematics* 121 (1993) 75–83.
- [11] V. Giakoumakis, I. Rusu. Weighted parameters in $(P_5, \overline{P_5})$ -free graphs. *Discrete Applied Mathematics* 80 (1997) 255–261.
- [12] M.C. Golumbic. *Algorithmic Graph Theory and Perfect Graphs*. 2nd edition, Annals of Discrete Mathematics 57, Elsevier, 2004.
- [13] C.T. Hoàng, D.A. Lazzarato. Polynomial-time algorithms for minimum weighted colorings of $(P_5, \overline{P_5})$ -free graphs and similar graph classes. *Discrete Applied Mathematics* 186 (2015) 106–111.