

Cosmological perturbations for an inflaton field coupled to radiation

Luca Visinelli

Department of Physics and Astronomy, University of Utah, 115 South 1400 East #201,
Salt Lake City, Utah 84112-0830, USA

E-mail: Luca.Visinelli@utah.edu

Abstract. Within the framework of the interacting fluid formalism, we provide the numerical solution to the Boltzmann equation describing the evolution of an inflaton field coupled to radiation. We study the behavior of the system during the slow-roll regime, in the case in which an additional stochastic source term is included in the set of equations, and we recover the expression for the cosmological perturbations previously obtained in the Warm inflation scenarios.

Keywords: Inflation, Warm inflation

Contents

1	Introduction	1
2	Equations for the background fields	2
3	Evolution of perturbations during inflation	5
4	Slow-roll regime with a stochastic source	9
5	Conclusions	12

1 Introduction

The mechanism of inflation [1–6] addresses the problems of the flatness, homogeneity, and the lack of relic monopoles raised by the standard Big-Bang cosmology [7–9], and provides an explanation to the inhomogeneities observed in the Cosmic Microwave Background Radiation (CMBR) [10–14]. It is usually postulated that during the inflationary period, the expansion of the Universe is driven by the potential energy of the so-called “inflaton” field. When the kinetic energy of the inflaton is no longer negligible with respect to its potential energy, the inflaton decays into lighter particles, which might be both hypothetical and Standard Model particles, and the Universe reheats and transits to a radiation-dominated epoch [15–20]. An alternative model postulates that the decay of the inflaton field occurs during the whole inflationary period, as in Warm Inflation (WI) scenarios ([21–37], see also Refs. [38–42]). Other models include multi-inflaton fields [43–49], natural inflation [50–54], and chain inflation [55, 56]; alternative mechanisms include the curvaton [57–60].

The only constraint on the reheating temperature of the Universe $T_{\text{RH}} \gtrsim 4$ MeV, valid in both the standard and WI models, comes from the requirement that a sufficiently large population of thermal neutrinos has to be produced [61–64], since a modification of the neutrino number density would spoil the observed abundance of light elements, the observed CMBR and matter power spectra [65, 66], and the large-scale structure.

As previously mentioned, inhomogeneities might have generated during inflation [67–69], either as quantum fluctuations of the inflaton (standard inflation scenarios [10–14]), or as thermal fluctuations (WI scenarios [21, 22]). These cosmological fluctuations later evolve into perturbations of the matter and radiation power spectra that are observed in the CMBR. The evolution of the fields participating inflation is described by a set of coupled Boltzmann equations, whose first-order perturbation describes the evolution of inhomogeneities. To some extent, these fields can be treated as fluids, with a pressure related to the energy density by an equation of state. The formalism for interacting fluids in an expanding Friedmann-Robertson-Walker (FRW) metric has been extensively discussed in Refs. [68–71].

In this work, we present the equations describing the coupling of the inflaton to a relativistic field using the interacting fluids approach developed in Refs. [70, 71]. The equations for the bulk and the cosmological fluctuations are solved numerically and tested against the analytical approximation around the reheating period. The paper is organized as follows. In Sec. 2, we provide a numerical solution to the basic equations for the background fields, while in Sec. 3 we study the time evolution of the perturbations in the inflaton and radiation fields.

Sec. 4 is devoted to the modifications induced in the equation by an additional stochastic term, which is usually considered in WI scenarios. Results are summarized in Sec. 5.

2 Equations for the background fields

We consider a model in which the energy density of the inflaton field ρ_ϕ dominates the energy density of the Universe, decaying into relativistic matter with energy density ρ_r at a rate Γ . We assume a FRW metric with cosmic time t and scale factor $a = a(t)$,

$$ds^2 = -dt^2 + a^2(t) \delta_{ij} dx^i dx^j. \quad (2.1)$$

If radiation thermalizes on a time scale much shorter than $1/\Gamma$, the potential $U(\phi, T)$ [72, 73], that expresses the interaction between the inflaton and radiation, splits as $U(\phi, T) = U(\phi) + \rho_r$ [22, 31]. Denoting the zero-th order (background) term in the perturbation with a bar over a quantity, we write the background expressions for the energy density and pressure fields as

$$\begin{aligned} \bar{\rho}_\phi &= \frac{1}{2} \dot{\bar{\phi}}^2 + U(\bar{\phi}), & \bar{p}_\phi &= \frac{1}{2} \dot{\bar{\phi}}^2 - U(\bar{\phi}), \\ \bar{\rho}_r &= \frac{\pi^2}{30} g_*(T) T^4, & \bar{p}_r &= c_r^2 \bar{\rho}_r, \end{aligned} \quad (2.2)$$

where $c_r^2 = 1/3$ describes the equation of state for relativistic matter. Under these conditions, Friedmann equations read [9]

$$\begin{aligned} H^2 &= \frac{8\pi}{3 M_{\text{Pl}}^2} \bar{\rho} = \frac{8\pi}{3 M_{\text{Pl}}^2} \left(\frac{1}{2} \dot{\bar{\phi}}^2 + U + \bar{\rho}_r \right), \\ \dot{H} &= -\frac{4\pi}{M_{\text{Pl}}^2} (\bar{p} + \bar{\rho}) = -\frac{4\pi}{M_{\text{Pl}}^2} \left(\dot{\bar{\phi}}^2 + \frac{4}{3} \bar{\rho}_r \right), \end{aligned} \quad (2.3)$$

where $M_{\text{Pl}} = 1.221 \times 10^{19}$ GeV is Planck's mass, $H = \dot{a}/a$ is the Hubble rate, and where a dot indicates a derivation with respect to the cosmic time t .

For each species α , the continuity equation is [68]

$$\dot{\bar{\rho}}_\alpha + 3H (\bar{p}_\alpha + \bar{\rho}_\alpha) = \bar{Q}_\alpha, \quad (2.4)$$

where \bar{Q}_α is a source term describing the conversion between the species α and the other species accounted in the theory. To assure the conservation of total energy density in a co-moving volume, the sum of all source terms must satisfy

$$\sum_\alpha \bar{Q}_\alpha = 0. \quad (2.5)$$

The conversion of the inflaton energy density into radiation energy density is thus described by the set of Boltzmann equations

$$\begin{aligned} \dot{\bar{\rho}}_\phi + 3H (\bar{p}_\phi + \bar{\rho}_\phi) &= -\bar{Q}, \\ \dot{\bar{\rho}}_r + 3H (\bar{p}_r + \bar{\rho}_r) &= \bar{Q}. \end{aligned} \quad (2.6)$$

At early times $t \ll 1/\Gamma$, the source term in the Boltzmann equation can be neglected, and the inflaton energy density is described by

$$\frac{d\bar{\rho}_\phi}{da} + \frac{3}{a} (\bar{p}_\phi + \bar{\rho}_\phi) = 0. \quad (2.7)$$

For a pressure-less fluid $\bar{p}_\phi = 0$, Eq. (2.7) predicts $\bar{\rho}_\phi \propto a^{-3}$ [74]. In this paper instead, we use the expression $\bar{p}_\phi + \bar{\rho}_\phi = 2\epsilon \bar{\rho}_\phi/3$, where ϵ is the first slow-roll parameter defined in Eq. (4.3) below. With this expression, the energy density is constant during ϕ domination as $\bar{\rho}_\phi \propto a^{-2\epsilon}$.

Inserting the expressions for $\bar{\rho}_\phi$ and \bar{p}_ϕ given in Eq. (2.2), together with the source term

$$\bar{Q} = \Gamma (\bar{p}_\phi + \bar{\rho}_\phi), \quad (2.8)$$

into Eq. (2.6), we obtain the evolution of the inflaton and radiation fields in a FRW metric,

$$\begin{aligned} \ddot{\phi} + (3H + \Gamma) \dot{\phi} + U_\phi &= 0, \\ \dot{\rho}_r + 4H \bar{\rho}_r &= \Gamma \dot{\phi}^2, \end{aligned} \quad (2.9)$$

where $U_\phi = \partial U / \partial \phi$. The second line in Eq. (2.9) corresponds to the Boltzmann equation for the massive scalar field ϕ decaying into a massless field ξ via an interaction proportional to $\phi \xi^2$ [20]. Here, we do not account for a microscopical model for the radiation field, which is treated as a fluid. The set in Eq. (2.9) has been often presented various time in the literature. Examples include warm inflation [25–27, 29, 34, 54], the decay of the inflaton [75, 76] or the curvaton field [77, 78], or the early domination of a massive scalar field [74, 79]. We solve the set of Eq. (2.9) numerically, assuming that the inflaton potential be [80]

$$U(\phi) = \frac{\lambda}{4} (\phi^2 - \phi_0^2)^2, \quad (2.10)$$

where λ is a dimensionless parameter describing the strength of the potential, and ϕ_0 is the value of the inflaton field at which the potential reaches its minimum value. Initial conditions for $\dot{\phi}$ and ρ_r are obtained by first solving the set of Eq. (2.9) in the regime $\ddot{\phi} \ll H \dot{\phi}$ and $\dot{\rho}_r \ll H \bar{\rho}_r$, starting from an initial configuration of the inflaton $\phi = \phi_i$. In Fig. 1, we show the energy densities $\bar{\rho}_\phi$ (red solid line) and $\bar{\rho}_r$ (blue solid line) as a function of time t , for the values of the parameters given in Table 1. Also shown is the Hubble rate H , multiplied by the constant $\phi_0^2 \Gamma$ in order to obtain the units of the energy density.

Quantity

$$\begin{aligned} \lambda &= 10^{-8} \\ \Gamma &= 10^{13} \text{ GeV} \\ \phi_0 &= 10^{17} \text{ GeV} \\ \phi_i &= 0.01 \phi_0 \end{aligned}$$

Table 1. Values for the parameters used in Fig. 1.

Defining the reheating time t_{RH} as the time at which

$$\bar{\rho}_\phi(t_{\text{RH}}) = \bar{\rho}_r(t_{\text{RH}}), \quad (2.11)$$

the total energy density is dominated by the inflaton energy density for $t < t_{\text{RH}}$, with a constant value $\bar{\rho}_\phi \approx U(\phi_i)$. When $t \approx t_{\text{RH}}$, the inflaton field starts oscillating around its




Figure 1. The energy density for the inflaton field $\bar{\rho}_\phi$ (red line), the radiation field $\bar{\rho}_r$ (blue line), and the Hubble rate $\phi_0^2 \Gamma H$, as a function of time t .

equilibrium value ϕ_0 , so we write the expansion $\phi = \phi_0 + \varphi$, where φ is a small perturbation. In this approximation, the inflaton potential is $U(\phi_0 + \varphi) \approx \lambda \phi_0^2 \varphi^2$, and Eq. (2.9) reads

$$\begin{aligned} \ddot{\varphi} + \left[\sqrt{24\pi G} \left(\lambda \phi_0^2 \varphi^2 + \frac{\dot{\varphi}^2}{2} + \bar{\rho}_r \right)^{1/2} + \Gamma \right] \dot{\varphi} + 2\lambda \phi_0^2 \varphi &= 0, \\ \dot{\bar{\rho}}_r + 4 \sqrt{\frac{8\pi G}{3}} \left(\lambda \phi_0^2 \varphi^2 + \frac{\dot{\varphi}^2}{2} + \bar{\rho}_r \right)^{1/2} \bar{\rho}_r - \Gamma \dot{\varphi}^2 &= 0. \end{aligned} \quad (2.12)$$

Keeping only the leading terms in φ and $\dot{\varphi}$, we obtain

$$\begin{aligned} \ddot{\varphi} + \Gamma \dot{\varphi} + 2\lambda \phi_0^2 \varphi &= 0, \\ \dot{\bar{\rho}}_r + 4 \sqrt{\frac{8\pi G}{3}} \bar{\rho}_r^{3/2} &= 0, \end{aligned} \quad (2.13)$$

with solution

$$\begin{aligned} \varphi(t) &\propto e^{-\Gamma t/2} \cos\left(\frac{t}{2} \sqrt{\Gamma^2 - 8\lambda \phi_0^2}\right), \\ \bar{\rho}_r(t) &\propto t^{-2}. \end{aligned} \quad (2.14)$$

The inflaton field approaches its equilibrium value ϕ_0 with an exponential decay; for the sake of illustration, in Fig. 2 we show a detail of the value of the inflaton field around $t = t_{\text{RH}}$. The dependence of the energy density of the relativistic field $\bar{\rho}_r(t) \propto t^{-2}$ is a standard result, which can also be inferred from the decoupled conservation equation,

$$\dot{\bar{\rho}}_r + 4H \bar{\rho}_r \approx 0, \quad (2.15)$$

and the fact that $a(t) \propto t^{1/2}$ during the radiation epoch. From this result, it follows that the Hubble rate (the black dashed line in Fig. 1) is $H(t) = 1/2t$ when $t > t_{\text{RH}}$.




Figure 2. The behavior of the inflaton field $\phi(t)$ around the time t_{RH} at which the inflaton converts into relativistic degrees of freedom, as a function of time t .

3 Evolution of perturbations during inflation

We consider the perturbations of the FRW background metric in Eq. (2.1), described by the metric in the longitudinal gauge [67, 69]

$$ds^2 = - (1 + 2\Psi) dt^2 + a^2(t) \delta_{ij} (1 - 2\Psi) dx^i dx^j, \quad (3.1)$$

where Ψ parametrizes the perturbations in the metric. Eq. (3.1) is valid whenever perturbations of the total energy-momentum tensor do not give rise to anisotropic stress.

Following Eqs. (2.15) and (2.17) in Ref. [70] with our notation and in the case in which the shear is zero, the expression for the perturbations in the energy density and pressure fields of a generic species α are

$$\delta\dot{\rho}_\alpha + 3H (\delta\rho_\alpha + \delta p_\alpha) - 3(\bar{\rho}_\alpha + \bar{p}_\alpha) \dot{\Psi} + a^{-2} \nabla^2 (\bar{\rho}_\alpha + \bar{p}_\alpha) \delta u_\alpha - \bar{Q}_\alpha \Psi - \delta Q_\alpha = 0, \quad (3.2)$$

and

$$\delta\dot{u}_\alpha + \left[\frac{\bar{Q}_\alpha}{\bar{\rho}_\alpha + \bar{p}_\alpha} (1 + c_\alpha^2) - 3c_\alpha^2 H \right] \delta u_\alpha + \Psi + \frac{1}{\bar{\rho}_\alpha + \bar{p}_\alpha} [\delta p_\alpha - \bar{Q}_\alpha \delta u] = 0. \quad (3.3)$$

Here, a dot indicates a derivation with respect to t , $\delta\rho_\alpha$ and δp_α are respectively the first-order perturbations in the energy density and pressure fields of the α -fluid, $c_\alpha^2 = \dot{p}_\alpha/\dot{\rho}_\alpha$, δu is the total covariant velocity perturbation, and ∇^2 is the co-moving spatial Laplacian. Defining the Fourier transform of a generic quantity $F(\mathbf{x}, t)$ as

$$F_q \equiv F(\mathbf{q}, t) = \frac{1}{(2\pi)^3} \int d^3x e^{-i\mathbf{q}\cdot\mathbf{x}} F(\mathbf{x}, t), \quad (3.4)$$

we take the Fourier transform of the equations for density and pressure perturbations, and set $\nabla^2 \rightarrow -q^2$. In momentum space, Eq. (3.2) for the inflaton and the radiation fields reads

$$\begin{aligned} \delta\dot{\rho}_{\phi q} + 3H(\delta\rho_{\phi q} + \delta p_{\phi q}) + (\bar{\rho}_\phi + \bar{p}_\phi) \left[\Gamma \Psi_q - 3\dot{\Psi}_q - \frac{q^2}{a^2} \delta u_{\phi q} \right] &= -\delta Q, \\ \delta\dot{\rho}_{r q} + 3H(\delta\rho_{r q} + \delta p_{r q}) - \bar{\rho}_r \left[\frac{4}{3} \frac{q^2}{a^2} \delta u_{r q} + 4\dot{\Psi}_q \right] - \Gamma (\bar{p}_\phi + \bar{\rho}_\phi) \Psi_q &= \delta Q. \end{aligned} \quad (3.5)$$

The first-order perturbation of the source terms in Eq. (2.8) is

$$\delta Q = \Gamma (\delta p_\phi + \delta \rho_\phi), \quad (3.6)$$

while the term $\bar{Q} \Psi_q$ containing the perturbations in the gravitational field has already been included in Eq. (3.5) which, once the term δQ in Eq. (3.6) has been substituted, reads

$$\begin{aligned} \delta\dot{\rho}_{\phi q} + (3H + \Gamma) (\delta\rho_{\phi q} + \delta p_{\phi q}) + (\bar{\rho}_\phi + \bar{p}_\phi) \left[\Gamma \Psi_q - 3\dot{\Psi}_q - \frac{q^2}{a^2} \delta u_{\phi q} \right] &= 0, \\ \delta\dot{\rho}_{r q} + 3H (\delta\rho_{r q} + \delta p_{r q}) - \bar{\rho}_r \left[\frac{4}{3} \frac{q^2}{a^2} \delta u_{r q} + 4\dot{\Psi}_q \right] - \Gamma (\bar{p}_\phi + \bar{\rho}_\phi) \Psi_q &= \Gamma (\delta\rho_{\phi q} + \delta p_{\phi q}). \end{aligned} \quad (3.7)$$

For velocity perturbations, we rewrite Eq. (3.3) for the inflaton and radiation fields as

$$\begin{aligned} \delta\dot{u}_{\phi q} - (3H + \Gamma) c_\phi^2 \delta u_{\phi q} + \Psi_q + \Gamma (\delta u_q - \delta u_{\phi q}) + \frac{\delta p_{\phi q}}{\bar{\rho}_\phi + \bar{p}_\phi} &= 0, \\ \frac{4}{3} \bar{\rho}_r (\delta\dot{u}_{r q} - H \delta u_{r q} + \Psi_q) + \Gamma (\bar{\rho}_\phi + \bar{p}_\phi) \left(\frac{4}{3} \delta u_{r q} - \delta u_q \right) + \delta p_{r q} &= 0. \end{aligned} \quad (3.8)$$

Perturbations in the gravitational field, described by the field Ψ appearing in the metric in Eq. (3.1), are described by [70]

$$\dot{\Psi}_q + H \Psi_q + 4\pi G \left[\frac{4}{3} \bar{\rho}_r \delta u_{r q} + (\bar{\rho}_\phi + \bar{p}_\phi) \delta u_{\phi q} \right] = 0. \quad (3.9)$$

Following Weinberg [9], the perturbations in the energy density, pressure, and velocity of the inflation field are related to the perturbations in the scalar field $\delta\phi$ by

$$\begin{aligned} \delta\rho_{\phi q} &= \dot{\phi} \delta\dot{\phi}_q + U_\phi \delta\phi_q - \dot{\phi}^2 \Psi_q, \\ \delta p_{\phi q} &= \dot{\phi} \delta\dot{\phi}_q - U_\phi \delta\phi_q - \dot{\phi}^2 \Psi_q, \\ \delta u_{\phi q} &= -\frac{\delta\phi_q}{\dot{\phi}}. \end{aligned} \quad (3.10)$$

In addition, we assume that perturbations in the radiation fluid are adiabatic, setting $\delta p_{r q} = c_r^2 \delta\rho_{r q} = \delta\rho_{r q}/3$. Combining the set in Eq. (3.10) with the expressions for the energy density and pressure fields in Eqs. (3.7), (3.8) and (3.9), we obtain

$$\delta\ddot{\phi}_q + (3H + \Gamma) \delta\dot{\phi}_q + \left(U_{\phi\phi} + \frac{q^2}{a^2} \right) \delta\phi_q = 4\dot{\phi} \dot{\Psi}_q - \left(2U_\phi + \Gamma \dot{\phi} \right) \Psi_q, \quad (3.11)$$

$$\delta\dot{\rho}_{r q} + 4H \delta\rho_{r q} - \frac{4}{3} \frac{q^2}{a^2} \bar{\rho}_r \delta u_{r q} = 4\bar{\rho}_r \dot{\Psi}_q + \Gamma \dot{\phi} \left(2\delta\dot{\phi}_q - \dot{\phi} \Psi_q \right), \quad (3.12)$$

$$\frac{4}{3} \bar{\rho}_r (\delta \dot{u}_{rq} - H \delta u_{rq} + \Psi_q) + \Gamma \dot{\phi} \left(\frac{4}{3} \dot{\phi} \delta u_{rq} + \delta \phi_q \right) + \frac{1}{3} \delta \rho_{rq} = 0, \quad (3.13)$$

$$\dot{\Psi}_q + H \Psi_q + 4\pi G \left(\frac{4}{3} \bar{\rho}_r \delta u_{rq} - \dot{\phi} \delta \phi_q \right) = 0. \quad (3.14)$$

In the literature, velocity perturbations in the radiation fluid have often been expressed in terms of the scalar potential velocity v_r , related to the covariant velocity perturbation used here by $\delta u_{rq} = a v_r/q$. When substituting for v_r , Eq. (3.13) reads

$$\dot{v}_r + \frac{\Gamma \dot{\phi}^2}{\bar{\rho}_r} v_r + \frac{q}{a} \left(\Psi_q + \frac{3\Gamma \dot{\phi}}{4\bar{\rho}_r} \delta \phi_q + \frac{\delta \rho_{rq}}{4\bar{\rho}_r} \right) = 0. \quad (3.15)$$

In their Eq. (A20), Moss and Xiong [30] quote the same expression as our Eq. (3.15), once set their $\alpha = \Psi_q$ and their $J = -\Gamma \dot{\phi} \delta \phi_q$ as in their Eq. (A13). Oliveira and Joras [25] quote the same as our Eq. (3.15) in the third line of their Eq. (A2); the second line of their Eq. (A2) agrees with our Eq. (3.11) once set $\Gamma' = 0$, while their first line in Eq. (A2) shows a term with an extra factor of three with respect to our Eq. (3.12). Our Eq. (3.11) shows all terms on the RHS of the equation with the opposite sign with respect to Eq. (61) quoted in Hall *et al.* [27].

To obtain the initial conditions for $\delta \phi_q$ and Ψ_q , we perform a WKB expansion using the fact that, at early times, the term $q/a(t)$ is much larger than H , and radiation can be neglected. Introducing the conformal time $d\tau = dt/a(t)$, we write

$$\delta \phi_q = f \exp(-iq\tau), \quad \text{and} \quad \Psi_q = g \exp(-iq\tau), \quad (3.16)$$

where $f(t)$ and $g(t)$ are slowly-varying functions. Substituting these expressions in Eqs. (3.11) and (3.14), and keeping the leading terms in q/a only, we obtain

$$f \sim a^{-3/2}, \quad \text{and} \quad g = \frac{4\pi i G \dot{\phi} a}{q} f. \quad (3.17)$$

Initial conditions for $\delta \rho_{rq}$ and δu_{rq} are obtained by solving Eqs. (3.11) and (3.14) for these variables, neglecting their time dependence and given the initial conditions for $\delta \phi_q$ and Ψ_q . We solve the set of Eqs. (3.11) (3.12) (3.13) (3.14) numerically using these initial conditions and the numerical solution to the background Eq. (2.9). The scale factor $a(t)$ that appears in the set of equations is obtained as the numerical solution of the differential equation $\dot{a} = H a$, where the Hubble rate H is defined by the Friedmann Eq. (2.3). In Fig. 3, we show results for the numerical resolution of this set of equations with the values for the parameters as in Table 1, for the values $q = 0.01$ (a), $q = 0.1$ (b), $q = 1$ (c).

After the reheating stage, the radiation perturbations dominate over the inflaton perturbations. The first-order perturbations in the radiation field oscillate with a frequency that increases with q . To explain this oscillatory behavior, we set $\Gamma = 0$, and we introduce the variable $\delta_{rq} = \delta \rho_{rq}/\bar{\rho}_r$, so that Eqs. (3.12), (3.14), and (3.15), when $t > t_{\text{RH}}$ are simplified as

$$\begin{aligned} \dot{\delta}_{rq} - \frac{4q}{3a} v_{rq} - 4\dot{\Psi}_q &= 0, \\ \dot{v}_r + \frac{q}{a} \left(\Psi_q + \frac{1}{4} \delta_{rq} \right) &= 0, \\ \dot{\Psi}_q + H \Psi_q + 2H^2 \frac{a}{q} v_{rq} &= 0. \end{aligned} \quad (3.18)$$




Figure 3. Perturbation in the energy density of the (red solid line) and radiation (blue solid line) and gravitational perturbations (black dashed line) in order of $\Gamma^2\phi_0^2$ obtained from the numerical resolution of the set of Eqs. (3.11), (3.12), (3.13), (3.14), with the parameters in Table 1 and for $q = 0.01$ (top figure), $q = 0.1$ (middle figure), and $q = 1$ (bottom figure).

Neglecting the perturbations in the gravitational potential Ψ_q , we find that both $\delta_{r,q}$ and v follow the same differential equation,

$$\ddot{x} + H \dot{x} + \frac{q^2}{3a^2} x = 0, \quad (3.19)$$

where $x(t)$ expresses either δ_{rq} or v . For a radiation-dominated cosmology, setting

$$H = \frac{1}{2t}, \quad a = a_{\text{RH}} \sqrt{\frac{t}{t_{\text{RH}}}}, \quad Q^2 = \frac{q^2}{3a_{\text{RH}}^2}, \quad (3.20)$$

this differential equation has solution

$$x(t) = x_1 \sin\left(2Q\sqrt{\frac{t}{t_{\text{RH}}}}\right) + x_2 \cos\left(2Q\sqrt{\frac{t}{t_{\text{RH}}}}\right), \quad (3.21)$$

where x_1 and x_2 are arbitrary constants.

In Fig. 4, we compare the solution to the expression in Eq. (3.20) in a radiation-dominated Universe (black dashed line) for the case of the density perturbations (top) and velocity perturbations (bottom), with the full numerical solution obtained from Eqs. (3.12) (blue solid line) and (3.13) (green solid line), setting $q = 0.1$. Once the matching initial conditions have been chosen, the analytical solution captures the oscillatory behavior and the trend obtained with the numerical resolution.

4 Slow-roll regime with a stochastic source

Inflation occurs if the potential $U(\phi)$ is sufficiently flat and much larger than all other forms of energy, so that the Hubble expansion rate H is constant. During this period, which is known as the slow-roll regime of the inflaton field, higher derivatives in Eq. (2.9) can be neglected,

$$\ddot{\phi} \ll H \dot{\phi}, \quad \text{and} \quad \dot{\rho}_r \ll H \rho_r. \quad (4.1)$$

In this regime, the Friedmann Eq. (2.3) and Eq. (2.9) for the motion of the inflaton field read

$$\begin{aligned} H^2 &\simeq \frac{8\pi G}{3} U, \\ \dot{\phi} &\simeq -\frac{U_\phi}{3H + \Gamma}, \\ \rho_r &\simeq \frac{\Gamma \dot{\phi}^2}{4H}, \end{aligned} \quad (4.2)$$

where we use the symbol “ \simeq ” for an equality that holds only in the slow-roll regime. Defining the slow-roll parameters,

$$\epsilon = \frac{1}{16\pi G} \left(\frac{U_\phi}{U}\right)^2, \quad \eta = \frac{1}{8\pi G} \frac{U_{\phi\phi}}{U}, \quad \beta = \frac{1}{8\pi G} \left(\frac{\Gamma_\phi U_\phi}{\Gamma U}\right), \quad (4.3)$$

the condition in Eq. (4.1) is met if the slow-roll parameters satisfy [24, 27, 30, 31, 54]

$$\epsilon \ll 1 + \frac{\Gamma}{3H}, \quad |\eta| \ll 1 + \frac{\Gamma}{3H}, \quad |\beta| \ll 1 + \frac{\Gamma}{3H}. \quad (4.4)$$

We now discuss the evolution of perturbations during the slow-roll regime. In the following, we take into account the effects from both quantum and thermal fluctuations, the latter being the predominant source for perturbations in WI scenarios. For this, we introduce a stochastic noise $\xi(\mathbf{x}, t)$ in the source term for $\delta\rho_\phi$, according to the Schwinger-Keldysh




Figure 4. Top: Density perturbations in the radiation fluid $\delta\rho_{rq}$ obtained with the numerical resolution of Eq. (3.12) (blue solid line) and with the analytical solution of Eq. (3.19) with $x(t) = \delta_{rq}$ (black dashed line). Bottom: Velocity perturbations in the radiation fluid δu_{rq} obtained with the numerical resolution of Eq. (3.13) (green solid line) and with the analytical solution of Eq. (3.19) with $x(t) = v$ (black dashed line).

approach to describe the quantum mechanical evolution of a system in a non-equilibrium state [81–83]. This term has been discussed in Ref. [84] in the context of the Boltzmann equation, and in Ref. [21] in the setting of WI. Instead of δQ_ϕ in Eq. (3.6), we thus consider the source term

$$\delta Q_\phi = -\Gamma (\delta p_\phi + \delta \rho_\phi) - \dot{\phi} \Gamma_{\text{eff}} \xi_q, \quad (4.5)$$

where ξ_q is the Fourier transform of the stochastic noise, which follows the statistical average

$$\langle \xi_q \xi_{q'} \rangle = a(t)^{-3} \delta^{(3)}(\mathbf{q} + \mathbf{q}') \delta(t - t'), \quad (4.6)$$

and we have defined the parameter [31]

$$\Gamma_{\text{eff}} = \sqrt{2 [\Gamma + H] T}, \quad (4.7)$$

where the temperature of the radiation bath is defined through $\bar{\rho}_r$ in Eq. (2.2). With this source perturbations, the equation for $\delta\phi$ in Eq. (3.11) modifies as

$$\delta\ddot{\phi}_q + (3H + \Gamma) \delta\dot{\phi}_q + \left(U_{\phi\phi} + \frac{q^2}{a^2} \right) \delta\phi_q - 4\dot{\phi} \dot{\Psi}_q + \left(2U_\phi + \Gamma \dot{\phi} \right) \Psi_q = -\Gamma_{\text{eff}} \xi_q. \quad (4.8)$$

We assume that, during the slow-roll regime, the gravitational field perturbation varies slowly as [85]

$$\dot{\Psi} \ll H \Psi, \quad (4.9)$$

so that, using Eq. (4.2), perturbations for the inflaton field in Eq. (4.8) are given by

$$\delta\ddot{\phi}_q + (3H + \Gamma) \delta\dot{\phi}_q + \left[U_{\phi\phi} + \frac{q^2}{a^2} - \frac{2H(3H + \Gamma)(6H + \Gamma)}{\Gamma} \frac{\bar{\rho}_r}{U} \right] \delta\phi_q \simeq -\Gamma_{\text{eff}} \xi_q. \quad (4.10)$$

We redefine the latter term in the square brackets as

$$- \frac{2H(3H + \Gamma)(6H + \Gamma)}{\Gamma} \frac{\bar{\rho}_r}{U} = \kappa \frac{U_\phi^2}{2U}, \quad (4.11)$$

where we have parametrized the strength of Γ with respect to H , as $\kappa = 1$ when $H \gg \Gamma$ and $\kappa = 1/2$ when $H \ll \Gamma$. With this substitution, the first line of Eq. (4.10) gives the dissipation equation presented in Ref. [31], except for the extra term containing $U_{\phi\phi}$ and U_ϕ^2 ,

$$\delta\ddot{\phi}_q + (3H + \Gamma) \delta\dot{\phi}_q + \left(U_{\phi\phi} + \kappa \frac{U_\phi^2}{2U} + \frac{q^2}{a^2} \right) \delta\phi_q \simeq -\Gamma_{\text{eff}} \xi_q, \quad (4.12)$$

In order to solve for ϕ_q , we switch to the variable

$$z = \frac{q}{H a(t)}, \quad (4.13)$$

and, using the slow-roll Eqs. (4.3) in the form $U_\phi^2 = 3H^2 U \epsilon/2$ and $U_{\phi\phi} = 3H^2 \eta$, we define

$$\nu = \frac{3}{2}(1 + R), \quad \text{and} \quad \mu = \sqrt{\nu^2 - 3(\kappa\epsilon + \eta)}. \quad (4.14)$$

Eq. (4.12) is rewritten as

$$\frac{d^2 \delta\phi}{dz^2} + \frac{1 - 2\nu}{z} \frac{d\delta\phi}{dz} + \left[1 + \frac{3(\kappa\epsilon + \eta)}{z^2} \right] \delta\phi = \frac{[2(\Gamma + H)T]^{1/2}}{H^2 z^2} \xi_q, \quad (4.15)$$

which is a second order stochastic differential equation, with solution for the field and its first derivative given respectively by

$$\begin{aligned}\delta\phi_q(z) &= z^\nu (\delta\phi_1 J_\mu(z) + \delta\phi_2 Y_\mu(z)) - \frac{\pi}{2} \frac{[2(\Gamma + H)T]^{1/2}}{H^2} \int_z^{+\infty} G_\mu^{(1)}(z, z') \left(\frac{z}{z'}\right)^\nu \xi_q(z') \frac{dz'}{z'}, \\ \delta\phi'_q(z) &= \delta\phi'_1 + \delta\phi'_2 - \frac{\pi}{2} \frac{[2(\Gamma + H)T]^{1/2}}{H^2} \int_z^{+\infty} G_\mu^{(2)}(z, z') \left(\frac{z}{z'}\right)^\nu \xi_q(z') \frac{dz'}{z'}.\end{aligned}\tag{4.16}$$

Here, $J_\mu(z)$ and $Y_\mu(z)$ are the Bessel functions of respectively the first and second kind of order μ , $\delta\phi_1$ and $\delta\phi_2$ are arbitrary constants, and we have introduced the functions

$$\begin{aligned}G_\mu^{(1)}(z, z') &= J_\mu(z) Y_\mu(z') - J_\mu(z') Y_\mu(z), \\ G_\mu^{(2)}(z, z') &= J_{\mu-1}(z) Y_\mu(z') - J_\mu(z') Y_{\mu-1}(z).\end{aligned}\tag{4.17}$$

This solution generalizes the result presented in Ref. [31] to the case $\mu \neq \nu$, and it is valid whenever the slow-roll parameters can be considered constant.

5 Conclusions

We shall summarize the main points of this paper. In Sec. 2, we have revised the Boltzmann equations for an inflaton field coupled to radiation, and we have solved the system of equation numerically in the case of a quartic inflaton potential $U(\phi)$, see Eq. (2.10). In Sec. 3, we have applied the interacting fluids formalism, based on the framework given in Refs. [68, 69] and described in Refs. [70, 71], to derive the differential equations governing the cosmological fluctuations arising during the inflationary period in Eqs. (3.11), (3.12), (3.13), (3.14). These equations have been solved numerically for a quartic inflaton potential $U(\phi)$. We give an analytical expression for $\delta\phi$ around the reheating period in Eq. (3.19), which approximates the frequency of oscillation obtained with the numerical solution.

In Sec. 4, we added a stochastic source to the expression describing $\delta\phi_q$, which is the approach usually considered in warm inflation scenarios. In the slow-roll regime, the equation for the inflaton perturbations can be solved exactly, including the stochastic term. The solution extends the result obtained in Ref. [31] to the case in which the slow-roll parameters are not neglected in the differential equation.

Acknowledgments

The author would like to thank Paolo Gondolo and Xuefang Sui for useful discussions on cosmological perturbations.

References

- [1] D. Kazanas, *Dynamics of the universe and spontaneous symmetry breaking*, *Astrophys. J.* **241**, L59 (1980).
- [2] A. A. Starobinsky, *A new type of isotropic cosmological models without singularity*, *Phys. Lett. B* **91**, 99 (1980).
- [3] A. H. Guth, *Inflationary universe: A possible solution to the horizon and flatness problems*, *Phys. Rev. D* **23**, 347 (1981).

- [4] K. Sato, *Cosmological baryon-number domain structure and the first order phase transition of a vacuum*, Phys. Lett. B **99**, 66 (1981).
- [5] A. Albrecht and P. J. Steinhardt, *Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking*, Phys. Rev. Lett. **48**, 1220 (1982).
- [6] A. D. Linde, *A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems*, Phys. Lett. B **108**, 389 (1982); *Chaotic Inflation*, Phys. Lett. B **129**, 177 (1983).
- [7] A. D. Linde, *Particle Physics and Inflationary Cosmology*, Harwood Academic, Switzerland (1990).
- [8] E. W. Kolb and M. S. Turner, *The Early Universe*, Addison-Wesley (1990).
- [9] S. Weinberg, *Cosmology*, Oxford University Press, USA (2008).
- [10] V. F. Mukhanov and G. V. Chibisov, *Quantum fluctuations and a nonsingular Universe*, Pisma Zh. Eksp. Teor. Fiz. **33**, 549 (1981) [JETP Lett. **33**, 532 (1981)].
- [11] A. H. Guth and S. Y. Pi, *Fluctuations in the New Inflationary Universe*, Phys. Rev. Lett. **49**, 1110 (1982).
- [12] S. W. Hawking, *The development of irregularities in a single bubble inflationary universe*, Phys. Lett. B **115**, 295 (1982).
- [13] A. A. Starobinsky, Phys. Lett. B **117**, 175 (1982).
- [14] J. M. Bardeen, P. J. Steinhardt and M. S. Turner, Phys. Rev. D **28**, 679 (1983).
- [15] A. Linde, *Hybrid Inflation*, Phys. Rev. D **49**, 748 (1994) [[astro-ph/9307.002](#)].
- [16] L. Kofman, A. Linde, and A. Starobinsky, *Reheating after Inflation*, Phys. Rev. Lett. **73**, 3195 (1994) [[hep-th/9405.187](#)].
- [17] L. Kofman, A. Linde, and A. Starobinsky, *Towards the Theory of Reheating After Inflation*, Phys. Rev. D **56**, 3258 (1997) [[hep-ph/9704.452](#)].
- [18] D. Boyanovsky, H. J. de Vega, R. Holman, and J. F. J. Salgado, *Analytic and Numerical Study of Preheating Dynamics*, Phys. Rev. D **54**, 7570 (1996) [[hep-ph/9608.205](#)].
- [19] P. B. Greene and L. Kofman, *Preheating of Fermions*, Phys. Lett. B **448**, 6 (1999) [[hep-ph/9807.339](#)].
- [20] J. Braden, L. Kofman, and N. Barnaby, *Reheating the Universe After Multi-Field Inflation*, JCAP **7**, 16 (2010) [[hep-th/1005.2196](#)].
- [21] A. Berera and L. Z. Fang, *Thermally Induced Density Perturbations in the Inflation Era*, Phys. Rev. Lett. **74**, 1912 (1995) [[astro-ph/9501.024](#)].
- [22] A. Berera, *Warm Inflation*, Phys. Rev. Lett. **75**, 3218 (1995) [[astro-ph/9509.049](#)].
- [23] A. Berera, *Warm Inflation in the Adiabatic Regime- a Model, an Existence Proof for Inflationary Dynamics in Quantum Field Theory*, Nucl. Phys. B **595**, 666 (2000) [[hep-ph/9904.409](#)].
- [24] A. N. Taylor and A. Berera, *Perturbation Spectra in the Warm Inflationary Scenario*, Phys. Rev. D **62**, 083517 (2000) [[astro-ph/0006.077](#)].
- [25] H. P. De Oliveira, S. E. Joras, *On Perturbations in Warm Inflation*, Phys. Rev. D **64**, 063513 (2001) [[gr-qc/0103.089](#)].
- [26] H. P. De Oliveira, *Density perturbations in warm inflation and COBE normalization*, Phys. Lett. B **526**, 1 (2002) [[gr-qc/0202.045](#)].
- [27] L. M. H. Hall, I. G. Moss, A. Berera, *Scalar perturbation spectra from warm inflation*, Phys. Rev. D **69** 083525 (2004) [[astro-ph/0305.015](#)].

- [28] M. Bastero-Gil and A. Berera, *Determining the regimes of cold and warm inflation in the supersymmetric hybrid model*, Phys. Rev. D **71**, 063515 (2005) [[hep-ph/0411.144](#)].
- [29] S. del Campo, R. Herrera, D. Pavon, *Cosmological perturbations in warm inflationary models with viscous pressure*, Phys. Rev. D **75**, 083518 (2007) [[astro-ph/0703.604](#)].
- [30] I. G. Moss and C. Xiong, *Non-gaussianity in fluctuations from warm inflation*, JCAP **0704**, 007 (2007) [[astro-ph/0701.302](#)].
- [31] I. G. Moss and C. Xiong, *On the consistency of warm inflation*, JCAP **0811**, 023 (2008) [[astro-ph/0808.0261](#)].
- [32] A. Berera, I. G. Moss, and R. O. Ramos, *Warm Inflation and its Microphysical Basis*, Rept. Prog. Phys. **72**, 026901 (2009) [[hep-ph/0808.1855](#)].
- [33] M. Bastero-Gil and A. Berera, *Warm inflation model building*, Int. J. Mod. Phys. A **24**, 2207 (2009) [[hep-ph/0902.0521](#)].
- [34] C. Graham and I. G. Moss, *Density fluctuations from warm inflation*, JCAP **0907**, 013 (2009) [[astro-ph/0905.3500](#)].
- [35] I. G. Moss and T. Yeomans, *Non-gaussianity in the strong regime of warm inflation*, [[astro-ph/1102.2833](#)] (2011).
- [36] S. Bartrum, A. Berera, and J. G. Rosa, *Warming up for Planck*, JCAP **1306**, 025 (2013) [[astro-ph/1303.3508](#)].
- [37] M. Bastero-Gil, A. Berera, I. G. Moss, and R. O. Ramos, *Cosmological fluctuations of a random field and radiation fluid*, [[astro-ph/1401.1149](#)] (2014).
- [38] A. Hosoya and M.-aki Sakagami, *Time development of Higgs field at finite temperature*, Phys. Rev. D **29**, 2228 (1984).
- [39] I. G. Moss, *Primordial inflation with spontaneous symmetry breaking*, Phys. Lett. B **154**, 120 (1985).
- [40] S. R. Lonsdale and I. G. Moss, *A superstring cosmological model*, Phys. Lett. B **189**, 12 (1987).
- [41] J. Yokoyama and K. Maeda, *On the Dynamics of the Power Law Inflation Due to an Exponential Potential*, Phys. Lett. B **207**, 31 (1988).
- [42] A. R. Liddle, *Power Law Inflation With Exponential Potentials*, Phys. Lett. B **220**, 502 (1989).
- [43] J. Garcia-Bellido, A. D. Linde, and D. Wands, *Density Perturbations and Black Hole Formation in Hybrid Inflation*, Phys. Rev. D **54**, 6040 (1996) [[astro-ph/9605.094](#)].
- [44] L. Randall, M. Soljagic, and A. H. Guth, *Supernatural Inflation: Inflation from Supersymmetry with No (Very) Small Parameters*, Nucl. Phys. B **472**, 377 (1996) [[hep-ph/9512.439](#)].
- [45] S. Tsujikawa, D. Parkinson, and B. A. Bassett, *Correlation-consistency cartography of the double inflation landscape*, Phys. Rev. D **67**, 083516 (2003) [[astro-ph/0210.322](#)].
- [46] D. Parkinson, S. Tsujikawa, B. A. Bassett, L. Amendola, *Testing for double inflation with WMAP*, Phys. Rev. D **71**, 063524 (2005) [[astro-ph/0409.071](#)].
- [47] K. Kadota and E. D. Stewart, *Successful Modular Cosmology*, JHEP **0307**, 013 (2003) [[hep-ph/0304.127](#)].
- [48] D. Wands, *Multiple field inflation*, Lect. Notes Phys. **738**, 275 (2008) [[astro-ph/0702.187](#)].
- [49] B. A. Bassett, S. Tsujikawa, and D. Wands, *Inflation Dynamics and Reheating*, Rev. Mod. Phys. **78**, 537 (2006) [[astro-ph/0507.632](#)].
- [50] K. Freese, J. A. Frieman, A. V. Olinto, *Natural inflation with pseudo Nambu-Goldstone bosons*, Phys. Rev. Lett. **65**, 3233 (1990).
- [51] F. Adams, J. R. Bond, K. Freese, J. Frieman, A. Olinto, *Natural inflation: Particle physics*

- models, power-law spectra for large-scale structure, and constraints from the Cosmic Background Explorer*, Phys. Rev. D **47**, 426 (1993) [[hep-ph/9207245](#)].
- [52] C. Savage K. Freese and W. H. Kinney, *Natural Inflation: status after WMAP 3-year data*, Phys. Rev. D **74**, 123511 (2006) [[hep-ph/0609.144](#)].
- [53] S. Mohanty, A. Nautiyal, *Natural inflation at the GUT scale*, Phys. Rev. D **78**, 123515 (2008) [[hep-ph/0807.0317](#)].
- [54] L. Visinelli, *Natural Warm Inflation*, JCAP **1109**, 013 (2011) [[astro-ph/1107.3523](#)].
- [55] K. Freese and D. Spolyar, *Chain Inflation in the Landscape: “Bubble Bubble Toil and Trouble”*, JCAP **0507**, 007 (2005) [[hep-ph/0412.145](#)].
- [56] K. Freese, J. T. Liu, and D. Spolyar, *Inflating with the QCD Axion*, Phys. Rev. D **72**, 123521 (2005) [[hep-ph/0502.177](#)].
- [57] K. Enqvist and M. S. Sloth, *Adiabatic CMB perturbations in pre-big bang string cosmology*, Nucl. Phys. B **626**, 395 (2002) [[hep-ph/0109.214](#)].
- [58] D. H. Lyth and D. Wands, *Generating the curvature perturbation without an inflaton*, Phys. Lett. B **524**, 5 (2002) [[hep-ph/0110.002](#)].
- [59] T. Moroi and T. Takahashi, *Effects of Cosmological Moduli Fields on Cosmic Microwave Background*, Phys. Lett. B **522**, 215 (2001) [Erratum-ibid. B **539**, 303 (2002)] [[hep-ph/0110.096](#)].
- [60] T. Moroi and T. Takahashi, *Cosmic Density Perturbations from Late-Decaying Scalar Condensations*, Phys. Rev. D **66**, 063501 (2002) [[hep-ph/0206026](#)].
- [61] M. Kawasaki, K. Kohri, and N. Sugiyama, *Cosmological Constraints on Late-time Entropy Production*, Phys. Rev. Lett. **82**, 4168 (1999) [[astro-ph/9811.437](#)].
- [62] M. Kawasaki, K. Kohri, and N. Sugiyama, *MeV-scale Reheating Temperature and Thermalization of Neutrino Background*, Phys. Rev. D **62**, 023506 (2000) [[astro-ph/0002.127](#)].
- [63] S. Hannestad, *What is the lowest possible reheating temperature?*, Phys. Rev. D **70**, 043506 (2004) [[astro-ph/0403.291](#)].
- [64] K. Ichikawa, M. Kawasaki, and F. Takahashi, *The oscillation effects on thermalization of the neutrinos in the universe with low reheating temperature*, Phys. Rev. D **72**, 043522 (2005) [[astro-ph/0505395](#)].
- [65] K. Ichikawa, M. Kawasaki, and F. Takahashi, *Constraint on the Effective Number of Neutrino Species from the WMAP and SDSS LRG Power Spectra*, JCAP **5**, 7 (2007), [[astro-ph/0611784](#)].
- [66] F. de Bernardis, L. Pagano, and A. Melchiorri, *Astroparticle Physics* **30**, 192 (2008).
- [67] J. Bardeen, *Gauge-invariant cosmological perturbations*, Phys. Rev. D **22**, 1882 (1980).
- [68] H. Kodama and M. Sasaki, *Cosmological Perturbation Theory*, Prog. Theor. Phys. Suppl. **78** (1984).
- [69] V. F. Mukhanov, H. A. Feldman, and R. H. Brandenberger, *Theory of cosmological perturbations*, Phys. Rep. **215**, 203 (1992).
- [70] K. A. Malik and D. Wands, *Adiabatic and entropy perturbations with interacting fluids and fields*, JCAP **02**, 007 (2005) [[astro-ph/0411.703](#)].
- [71] K. A. Malik, D. Wands, and C. Ungarelli, *Large-scale curvature and entropy perturbations for multiple interacting fluids*, Phys. Rev. D **67**, 063516 (2003) [[astro-ph/0211.602](#)].
- [72] M. Bellini, *Fresh inflation with increasing cosmological parameter*, Phys. Rev. D **67**, 027303 (2003) [[gr-qc/0211.044](#)].
- [73] T. Matsuda, *Aspects of warm-flat directions*, Int. J. Mod. Phys. A **25**, 4221 (2010)

- [hep-ph/0908.3059].
- [74] A. L. Erickcek, K. Sigurdson, *Reheating Effects in the Matter Power Spectrum and Implications for Substructure*, Phys. Rev. D **84**, 083503 (2011) [[astro-ph/1106.0536](#)].
- [75] R. Allahverdi and M. Drees, *Production of massive stable particles in inflaton decay*, Phys. Rev. Lett. **89**, 091302 (2002) [[hep-ph/0203.118](#)].
- [76] R. Allahverdi and M. Drees, *Thermalization after inflation and production of massive stable particles*, Phys. Rev. D **66**, 063513 (2002) [[hep-ph/0205.246](#)].
- [77] S. Gupta, K. A. Malik, and D. Wands, *Curvature and isocurvature perturbations in a three-fluid model of curvaton decay*, Phys. Rev. D **69**, 063513 (2004) [[astro-ph/0311.562](#)].
- [78] S. Matarrese and A. Riotto, *Large-scale curvature perturbations with spatial and time variation of the inflaton decay rate*, JCAP **0308**, 007 (2003) [[astro-ph/0306.416](#)].
- [79] D. J. H. Chung, E. W. Kolb, A. Riotto, *Production of massive particles during reheating*, Phys. Rev. D **60**, 063504 (1999) [[hep-ph/9809.453](#)].
- [80] D. H. Lyth and A. Riotto, Phys. Rept. **314**, 1 (1999) [[hep-ph/9807.278](#)].
- [81] J. Schwinger, *Brownian motion of a quantum oscillator*, J. Math. Phys. **2**, 407 (1961).
- [82] L. V. Keldysh, *Diagram Technique for Non-equilibrium Processes*, J. Exptl. Theoret. Phys. **47**, 1515 (1964) [<http://www.jetp.ac.ru>].
- [83] J. Rammer and H. Smith, *Quantum field-theoretical methods in transport theory of metals*, Rev. Mod. Phys. **58**, 323 (1986).
- [84] E. Calzetta and B. L. Hu, *Stochastic dynamics of correlations in quantum field theory: From Schwinger-Dyson to Boltzmann-Langevin equation*, Phys. Rev. D **61**, 025012 (2000) [[hep-ph/9903.291](#)].
- [85] D. Polarski and A. A. Starobinsky, Nucl. Phys. B **385**, 623 (1992).
- [86] S. Weinberg, *Adiabatic Modes in Cosmology*, Phys. Rev. D **67**, 123504 (2003) [[astro-ph/0302.326](#)].