

## RENORMALIZATION AND ASYMPTOTIC STATES IN LORENTZ-VIOLATING QFT

M. CAMBIASO

*Universidad Andres Bello, Departamento de Ciencias Físicas,  
Facultad de Ciencias Exactas, Av. República 220, Santiago, Chile.*

*\*E-mail: mcambiaso@unab.cl*

R. LEHNERT

*Indiana University Center for Spacetime Symmetries,  
Bloomington, IN 47405, USA.*

*E-mail: ralehner@indiana.edu*

R. POTTING

*CENTRA, Departamento de Física, Universidade do Algarve,  
8005-139 Faro, Portugal.*

*E-mail: rpotting@ualg.pt*

Radiative corrections in quantum field theories with small departures from Lorentz symmetry alter structural aspects of the theory, in particular the definition of asymptotic single-particle states. Specifically, the mass-shell condition, the standard renormalization procedure as well as the Lehmann-Symanzik-Zimmermann reduction formalism are affected.

In the presence of Lorentz breakdown, previous analyses involving the properties of freely propagating particles have been performed under the tacit assumption that the physics of free particles is determined by the quadratic pieces of the corresponding Lagrangian. However, by analyzing radiative corrections for a sample Lorentz-violating (LV) Lagrangian contained in the SME, we claim that the previous line of reasoning fails.

We consider the bare gauge-invariant Lagrangian density for single-flavor QED within the minimal SME in the presence of the  $c^{\mu\nu}$  and  $\tilde{k}^{\mu\nu}$  only. Both are symmetric and traceless and the latter is defined as a  $\tilde{k}^{\mu\nu} = (k_F)^{\mu\alpha\nu}{}_{\alpha}$ . One-loop multiplicative renormalizability of the model is assumed.<sup>1</sup> For convenience we write the gauge-fixed bare Lagrangian together with appropriate IR regularization terms.<sup>2</sup> To understand the external fermion states we need to consider radiative corrections

to the fermion two-point function, wherefrom the wave-function renormalization “constant” is read-off as the residue of the one-particle pole. As a zeroth-order system on which perturbation theory is set up, we choose the full renormalized quadratic Lagrangian together with the LV part. The remaining non-quadratic contributions to the Lagrangian being taken as perturbations. Thus, the proper two-point function adopts the form  $\Gamma^{(2)}(p) = \Gamma^\mu p_\mu - m - \Sigma(p^\mu)$ , where  $\Gamma^\mu = \gamma^\mu + c^{\mu\nu} \gamma_\nu$  and

$$\Sigma(p^\mu) = \Sigma_{\text{LI}}(\not{p}) + \Sigma_{\text{LV}}(p^2, c_\gamma^p, \tilde{k}_\gamma^p) + \delta\Sigma(p^\mu, c^{\mu\nu}, \tilde{k}^{\mu\nu}), \quad (1)$$

with  $c_\gamma^p \equiv c^{\mu\nu} p_\mu \gamma_\nu$  and similarly for  $\tilde{k}_\gamma^p$ . The first term on the RHS of Eq. (1) denotes the usual Lorentz-symmetric contributions. The rest come from LV terms. The last, however, contains terms that are not present in the original Lagrangian. To linear order in LV and given that electromagnetic interactions preserve C, P, and T, we can write:

$$\Sigma_{\text{LV}}(p^2, c_\gamma^p, \tilde{k}_\gamma^p) = f_2^c(p^2) c_\gamma^p + f_2^{\tilde{k}}(p^2) \tilde{k}_\gamma^p, \quad (2)$$

$$\delta\Sigma(p^\mu, c_p^p, \tilde{k}_p^p) = f_3^c(p^2) \frac{c_p^p}{m} + f_4^c(p^2) \frac{\not{p} c_p^p}{m^2} + f_3^{\tilde{k}}(p^2) \frac{\tilde{k}_p^p}{m} + f_4^{\tilde{k}}(p^2) \frac{\not{p} \tilde{k}_p^p}{m^2}. \quad (3)$$

The functions  $f_i^c(p^2)$  and  $f_i^{\tilde{k}}(p^2)$  are calculable to any order in the fine-structure constant  $\alpha$ . Following Ref. 3 we extract the propagator pole:<sup>a</sup>

$$\Gamma^{(2)}(p) = \mathcal{Z}_R^{-1} \bar{P}(p) + \bar{P}(p) \Sigma_2(\not{p}, c_\gamma^p, \tilde{k}_\gamma^p, (c, k)_p^p) \bar{P}(p), \quad (4)$$

which yields the desired property:  $\Gamma^{(2)}(p)^{-1} = \mathcal{Z}_R \bar{P}(p)^{-1} + \text{finite}$ .

Doing standard perturbative calculations to first order in LV coefficients and to first order in  $\alpha$ , the one-loop corrected wave-function renormalization and the fermion propagator pole, respectively read:

$$\mathcal{Z}_R^{-1} = (\mathcal{Z}_R)_{\text{LI}}^{-1} - \frac{2\alpha}{3\pi m^2} [2c_p^p + \tilde{k}_p^p], \quad (5)$$

$$\bar{P}(p) = \not{p} + (c_{\text{phys}})_\gamma^p - m_{\text{phys}} + \frac{\alpha}{3\pi m} (2(c_{\text{phys}})_p^p - (\tilde{k}_{\text{phys}})_p^p), \quad (6)$$

where:

$$m_{\text{phys}} = m + \frac{\alpha m}{\pi} \left[ 1 - \frac{3}{4} \gamma_E - \frac{3}{4} \ln \left( \frac{m^2}{4\pi\mu^2} \right) \right] \quad (7)$$

is the usual loop-corrected mass in the minimal-subtraction scheme, and

$$(c_{\text{phys}})^{\mu\nu} = c^{\mu\nu} - \frac{\alpha}{3\pi} \left[ \frac{29}{12} - \gamma_E - \ln \left( \frac{m^2}{4\pi\mu^2} \right) \right] (2c^{\mu\nu} - \tilde{k}^{\mu\nu}). \quad (8)$$

<sup>a</sup>The dependence of  $\Gamma^{(2)}(p)$  on  $c_\gamma^p$  and  $\tilde{k}_\gamma^p$  introduce further subtleties. Details of the one-loop calculation together with the necessary Feynman rules can be found in Ref. 4.

Equation (8) expresses the radiatively corrected, physical value of the Lorentz-violating parameter  $c^{\mu\nu}$  in terms of its tree-level value. Note that both  $c^{\mu\nu}$  and the mass scale  $\mu$  are unphysical, renormalization-scheme-dependent quantities, unlike  $(c_{\text{phys}})^{\mu\nu}$ , which is in principle measurable. This is in fact granted by the cancellation between the terms containing  $\mu$  in Eq. (8) with those coming from the running of  $c^{\mu\nu}$ .<sup>1</sup> UV divergent quantities have been regularized with dimensional regularization. Equations (6) and (8) show that LV radiative corrections depend only on  $(2c_p^\mu - \tilde{k}_p^\mu)$  and  $c_{\text{phys}}^{\mu\nu}$  is indeed infrared finite. Furthermore, it can be shown that the  $\mathcal{O}(\alpha)$  LV radiative corrections to the dispersion relation are proportional to  $(2c_p^\mu - \tilde{k}_p^\mu)$  too. This property is a requirement coming from considerations of field redefinitions which imply that, in this context, physically observable radiative effects should depend on  $2c^{\mu\nu} - \tilde{k}^{\mu\nu}$  only.<sup>5</sup> Turning to the description of asymptotic states, Eq. (6) implies a modified equation of motion for in- and out-spinors. Correspondingly an adapted LSZ reduction formula for the fundamental scattering amplitude  $\langle f|i\rangle$ , controlled by Eqs. (5) and (6), is obtained.

Our results allow us to conclude that asymptotic single-particle states of fermions in Lorentz-violating QFT receive concrete modifications due to radiative corrections. Specifically, the corresponding Dirac equation turns out to be modified by Lorentz-violating operators not present in the Lagrangian, a novel feature of LV QFT as it does not occur in the Lorentz-symmetric case. Also, the fermion wave-function renormalization is no longer a constant but rather depends on the LV coefficients under consideration and on the external momentum as well. As shown in Ref. 4 a non-trivial cancellation of IR divergences is also achieved in the fermion's dispersion relation and wave-function renormalization function. Furthermore, IR divergences cancel in the elastic Coulomb scattering cross section when the contribution of soft-photon emission from the corresponding external legs is taken into account.

### Acknowledgments

This work has been supported in part by: the Portuguese Fundação para a Ciência e a Tecnologia, the Mexican RedFAE, UNAB Grant DI-27-11/R, FONDECYT No. 11121633 and by the IUCSS.

### References

1. V.A. Kostelecký, C.D. Lane, and A.G.M. Pickering, Phys. Rev. D **65**, 056006 (2002).

2. M. Cambiaso, R. Lehnert, and R. Potting, Phys. Rev. D **85**, 085023 (2012).
3. R. Potting, Phys. Rev. D **85**, 045033 (2012).
4. M. Cambiaso, R. Lehnert and R. Potting, Phys. Rev. D **90**, 065003 (2014).
5. See, e.g., D. Colladay and P. McDonald, J. Math. Phys. **43**, 3554 (2002); R. Lehnert, Phys. Rev. D **74**, 125001 (2006); Rev. Mex. Fis. **56**, 469 (2010); B. Altschul, J. Phys. A **39**, 13757 (2006).