

SHARP DECAY ESTIMATES FOR AN ANISOTROPIC LINEAR SEMIGROUP AND APPLICATIONS TO THE SQG AND INVISCID BOUSSINESQ SYSTEMS

TAREK M. ELGINDI AND KLAUS WIDMAYER

ABSTRACT. At the core of this article is an improved, sharp dispersive estimate for the anisotropic linear semigroup $e^{R_1 t}$ arising in both the study of the dispersive SQG equation and the inviscid Boussinesq system. We combine the decay estimate with a blow-up criterion to show how dispersion leads to long-time existence of solutions to the dispersive SQG equation, improving the results obtained using hyperbolic methods. In the setting of the inviscid Boussinesq system it turns out that linearization around a specific stationary solution leads to the same linear semigroup, so that we can make use of analogous techniques to obtain stability of the stationary solution for an increased timespan.

CONTENTS

1. Introduction	1
1.1. The SQG Equation	2
1.2. The Inviscid Boussinesq System	3
1.3. Plan of the Article	4
2. Dispersion	5
2.1. Dispersive Estimate	5
2.2. Sharpness of the Dispersive Estimate	10
3. Increased Time of Existence of Solutions to the SQG Equation	11
4. A Stability Result for the Inviscid Boussinesq System	13
4.1. Linearized Equations	14
4.2. Nonlinear Stability	14
References	16

1. INTRODUCTION

The key ingredient of this article is an improved, sharp dispersive estimate for the semigroup $e^{R_1 t}$ of the linear equation

$$\partial_t \theta - R_1 \theta = 0, \quad \theta : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}.$$

2000 *Mathematics Subject Classification.* 76B03, 76B15, 35B35, 35Q31, 35Q35.

Key words and phrases. dispersive surface quasi-geostrophic equation, inviscid Boussinesq system, stability.

Here, R_1 is the Riesz transform with Fourier multiplier with symbol $-i\frac{\xi_1}{|\xi|}$. The presence of the Riesz transform makes this an an-isotropic equation and introduces a degeneracy that precludes the application of the standard stationary phase lemma.

Equipped with such a dispersive estimate we can perturbatively treat nonlinear versions of the above equation. More specifically, here we focus on two problems of independent interest, both with quadratic nonlinearities: The dispersive surface quasi-geostrophic equation (SQG) and the inviscid Boussinesq system, discussed in the following.

1.1. The SQG Equation. We are concerned with the time of existence of solutions $\theta : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}$ to the dispersive surface quasi-geostrophic equation (SQG)

$$\partial_t \theta + u \cdot \nabla \theta = R_1 \theta,$$

where $u = \nabla^\perp (-\Delta)^{-1/2} \theta$. This equation has been suggested as a model in geophysical fluid dynamics and may prove useful in the study of wave-turbulence interactions (see [22]). As the name suggests, this fluid equation exhibits dispersion.

The inviscid and viscous SQG equations have been studied by many authors, particularly because they are expected to model the main features of $3d$ incompressible flow (see [9]). We mention only a few results from the literature: Local existence and uniqueness for H^{2+} solutions and global existence of L^2 solutions were established in [21]. Blow-up criteria of Beale-Kato-Majda type were obtained in [9]. In recent years, the critically dissipative SQG equation was proven to be globally well-posed in H^s spaces for $s \geq 1$ by several authors using a variety of methods ([17], [4], [10], [18]). Aside from local well-posedness and breakdown criteria, not much is known about the well-posedness of the inviscid SQG equation.

On the other hand, the addition of the Riesz transform linearity $R_1 \theta$ on the right hand side of the inviscid SQG equation $\partial_t \theta + u \cdot \nabla \theta = 0$ allows for this problem to be studied from a perturbative point of view. This is a standard approach and has been very fruitful in the setting of dispersive partial differential equations, especially when the decay properties of the linear equation are very strong. Notice, however, that in the case of the dispersive SQG equation the Riesz transform breaks the isotropy of the equation. Apart from other difficulties this entails that – in contrast to many other dispersive equations – getting the optimal decay rate of the linear semigroup associated to the problem is not a simple matter.

The available existence theory for this equation is very basic. From the hyperbolic systems point of view the dispersive and the inviscid SQG equations have the same energy estimates, thus solutions (for either) exist a priori only for a timespan inversely proportional to the size of the initial data. A finer understanding of the dispersive properties allows us to improve this timespan. In particular, here it is crucial to obtain a sharp decay result, which has not been done before for such equations.

In comparison, other works on this type of equations have studied them in the presence of a viscous or dissipative term. This typically resolves the existence question (for small data or a large dispersion constant) and in some cases allows for the use of the dispersion to derive equations in limiting cases (see for example the case of the β -plane equation in [8, Chapter 5], or the work of Babin et al. [2] on the

Navier-Stokes equations with Coriolis forcing). In this context the exact dispersive rate is not of key importance.

In a related work [5], Cannone, Miao and Xue proved the existence of global strong solutions to the dispersive SQG equation with supercritical dissipation. Specifically, the authors of [5] study

$$\partial_t \theta + u \cdot \nabla \theta = -\nu(-\Delta)^\alpha \theta + KR_1 \theta, \quad 0 < \alpha < 1,$$

and prove that for given initial data θ_0 there exists K large enough such that the solution is global. They use the dissipation in a crucial way, whereas our model does not have any (i.e. we study the case where $\nu = 0$). On the other hand, the rate of dispersive decay is not central to this argument and our improvement of the dispersive estimate allows for lowering the constant K for which one obtains a global solution.

On a separate note we remark that the dispersive SQG equation might be viewed as a natural generalization of the (not dispersive) Burgers-Hilbert Equation studied by Ifrim and Tataru in [16]. Their use of a new, modified energy method is based on the existence of a normal form. However, for the dispersive SQG equation such a transformation does not seem to be available.

1.2. The Inviscid Boussinesq System. We consider an incompressible, inviscid fluid $v : \mathbb{R}^+ \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ of variable density $\rho : \mathbb{R}^+ \times \mathbb{R}^3 \rightarrow \mathbb{R}$ under the influence of an external gravity force proportional to ρ in direction $\mathbf{k} \in \mathbb{R}^3$, described by Euler's equation coupled to a continuity equation and an equation of state:

$$\begin{cases} \rho(\partial_t v + v \cdot \nabla v) = -\nabla p + g\rho\mathbf{k}, \\ \partial_t \rho + v \cdot \nabla \rho = 0, \\ \operatorname{div} v = 0, \end{cases}$$

where $p : \mathbb{R}^+ \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is the fluid pressure and g is a gravitational constant.

Here we will further investigate the *Boussinesq approximation* of these equations, where the density variation is assumed to be small in comparison with the effects of gravity. For $2d$ flows – choosing $\mathbf{k} = e_2$ and $g = 1$ – these equations can then be reduced to a system of scalar vorticity equations (see [12], [15]), the so-called *inviscid Boussinesq system*:

$$(1.1) \quad \begin{cases} \partial_t \omega + u \cdot \nabla \omega = \partial_x \rho, \\ \partial_t \rho + u \cdot \nabla \rho = 0, \\ u = \nabla^\perp (-\Delta)^{-1} \omega, \end{cases}$$

where $\omega, \rho : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}$.

The question of global well-posedness for this equation remains unresolved. From a hyperbolic systems point of view these equations bear a lot of similarity to the SQG equation. In particular, using energy methods and Beale-Kato-Majda type blow-up criteria, local existence results in various spaces have been obtained by several authors ([7], [6], [19], [23]).

In the present article we are concerned with the stability of certain stationary solutions of the Boussinesq system (1.1) – in one sentence, we are going to study the

opposite of the Rayleigh-Bernard instability, namely the setup where warm fluid is on top of cooler fluid. We will prove long-time stability *in the absence of viscosity*. Here the temperature and density are assumed to be proportionally related, so that the cooler fluid is more dense. The gravitational force is thus expected to stabilize such a density (or temperature) distribution.

The phenomenon known as Rayleigh-Bernard convection has been studied by a number of authors for many years. The idea is simple: take a container filled with water which is at rest. Now heat the bottom of the container and cool the top of the container. It has been observed experimentally and mathematically that if the temperature difference between the top and the bottom goes beyond a certain critical value, the water will begin to move and convective rolls will begin to form. This effect is called Rayleigh-Bernard instability ([13],[11]).

Now, in the other case, when one cools the bottom and heats the top, it is expected that the system remains stable. In the presence of viscosity it is not difficult to prove this fact (see [11]). However, without the effects of viscosity (or temperature dissipation), it is conceivable for such a configuration to be unstable. Indeed, in the absence of viscosity, non-linear stability is not known to be true for most fluid equations, except in very specific cases (see, for example, [20], [3], [14]).

In this work, we prove that a particular stationary configuration is *non-linearly stable* on an interval of time which is proportional to the 4/3-power of the inverse of the size of the perturbation. Prior to this work, the long-time existence of non-trivial solutions was unknown. We emphasize that in the viscous case, stability of the stationary solution we will study is trivial. However, in the inviscid case, even *linear* stability seems to have been unknown prior to this work.

1.3. Plan of the Article. We start by considering the dispersive properties of the linear evolution of our problems in section 2. Given the lack of isotropy in the equation one might expect a slower rate of decay than for other dispersive $2d$ models. Proposition 2.1 shows that this is indeed the case – more precisely, solutions to the linear equation decay at a rate of $t^{-1/2}$ in L^∞ . With the help of special functions we can then show that this decay is sharp.

In section 3 we demonstrate how to exploit the dispersion to obtain an increased time of existence of solutions of the SQG equation. This is the content of Theorem 3.2.

Finally we show how the same semigroup as for the SQG equation comes up in the question of stability of solutions to the Boussinesq equation (section 4) and directly obtain a corresponding stability result (Theorem 4.2).

Notation. On \mathbb{R}^2 we shall denote by R_j the Riesz transform in the variable x_j , $1 \leq j \leq 2$, given by

$$R_j f(x) = - \int_{\mathbb{R}^2} i \frac{\xi_j}{|\xi|} \hat{f}(\xi) e^{ix \cdot \xi} d\xi.$$

For $s \in \mathbb{R}$ and $1 \leq p \leq \infty$ we denote by $W^{s,p}$ the inhomogeneous Sobolev space with derivatives up to order s in L^p , and write $H^s := W^{s,2}$, while \dot{H}^s stands for the homogeneous Sobolev space of derivatives of order s in L^2 . By $\dot{B}_{b,c}^a$ we shall denote

the homogeneous Besov space of a -th derivatives of the frequency localized pieces in L^b , summed with respect to ℓ^c .

2. DISPERSION

In this section we consider the dispersive properties of the linear equation

$$(2.1) \quad \partial_t \omega - R_1 \omega = 0, \quad \omega : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}.$$

Its semigroup is given by

$$e^{R_1 t} f(x) = \int_{\mathbb{R}^2} \hat{f}(\xi) e^{ix \cdot \xi - it \frac{\xi_1}{|\xi|}} d\xi = \int_{\mathbb{R}^2} \hat{f}(\xi) e^{it\phi(\xi)} d\xi,$$

where we denote by ϕ the phase¹

$$\phi(\xi) := \frac{x}{t} \xi - \frac{\xi_1}{|\xi|}.$$

A direct computation gives the determinant of the Hessian (in ξ) of ϕ as $-\frac{\xi_2^2}{|\xi|^6}$. The degeneracy of this along $\{\xi_2 = 0\}$ prevents us from applying stationary phase techniques directly to obtain a decay estimate for $e^{R_1 t} f$. Instead we use a non-singular integration by parts formula (see (2.6)), which gives a slower rate of decay of $t^{-1/2}$ (Proposition 2.1). As we will show further below, this decay is actually sharp (Proposition 2.5).

2.1. Dispersive Estimate.

Proposition 2.1. *Let $f \in C_c^\infty(\mathbb{R}^2)$. Then*

$$(2.2) \quad \|e^{R_1 t} f\|_{L^\infty} \leq C t^{-1/2} \|f\|_{\dot{B}_{1,1}^2}$$

for some constant $C > 0$.

Proof. We will localize f in frequency (using Littlewood-Paley theory), estimate the different pieces separately and sum in the end.

1. *Localization:* For $j \in \mathbb{Z}$ let P_j be the Littlewood-Paley projectors associated to a smooth bump function $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}$ with support near $|\xi| \sim 1$, i.e. $\widehat{P_j f}(\xi) =$

¹We omit the variables x, t from the notation since we regard them as fixed (but arbitrary).

$\varphi(2^{-j}\xi)\hat{f}(\xi)$. We note that by Young's convolution inequality

$$\begin{aligned}
\|e^{R_1 t} f\|_{L^\infty} &= \left\| e^{R_1 t} \left(\sum_{j \in \mathbb{Z}} P_j f \right) \right\|_{L^\infty} \leq \sum_{j \in \mathbb{Z}} \|e^{R_1 t} P_j f\|_{L^\infty} \\
&= \sum_{j \in \mathbb{Z}} \left\| \mathcal{F}^{-1} \left(e^{-it \frac{\xi_1}{|\xi|}} \varphi(2^{-j}\xi) \hat{f}(\xi) \right) \right\|_{L^\infty} \\
&= \sum_{j \in \mathbb{Z}} \left\| \mathcal{F}^{-1} \left(e^{-it \frac{\xi_1}{|\xi|}} \varphi(2^{-j}\xi) \right) * \mathcal{F}^{-1} \left(\tilde{\varphi}(2^{-j}\xi) \hat{f}(\xi) \right) \right\|_{L^\infty} \\
&\leq \sum_{j \in \mathbb{Z}} \left\| \mathcal{F}^{-1} \left(e^{-it \frac{\xi_1}{|\xi|}} \varphi(2^{-j}\xi) \right) \right\|_{L^\infty} \left\| \mathcal{F}^{-1} \left(\tilde{\varphi}(2^{-j}\xi) \hat{f}(\xi) \right) \right\|_{L^1} \\
&= \sum_{j \in \mathbb{Z}} \left\| \mathcal{F}^{-1} \left(e^{-it \frac{\xi_1}{|\xi|}} \varphi(2^{-j}\xi) \right) \right\|_{L^\infty} \|Q_j f\|_{L^1},
\end{aligned}$$

where Q_j is a Littlewood-Paley projector associated to a slightly ‘‘fattened’’ bump function $\tilde{\varphi}$ (which equals 1 on the support of φ).

Hence it suffices to estimate the semigroup on the frequency localizers φ . To this end we notice that by a change of variables

$$\begin{aligned}
(e^{R_1 t} P_j)(x) &= \int_{\mathbb{R}^2} \varphi(2^{-j}\xi) e^{ix \cdot \xi - it \frac{\xi_1}{|\xi|}} d\xi = 2^{2j} \int_{\mathbb{R}^2} \varphi(\xi) e^{i(2^j x) \cdot \xi - it \frac{\xi_1}{|\xi|}} d\xi \\
&= 2^{2j} (e^{R_1 t} P_0)(2^j x).
\end{aligned}$$

Since we will estimate this in L^∞ we can further reduce to estimating

$$(2.3) \quad \|e^{R_1 t} P_0\|_{L^\infty}$$

and our final estimate will be

$$\begin{aligned}
\|e^{R_1 t} f\|_{L^\infty} &\lesssim \sum_{j \in \mathbb{Z}} 2^{2j} \|e^{R_1 t} P_0\|_{L^\infty} \|Q_j f\|_{L^1} = \|e^{R_1 t} P_0\|_{L^\infty} \sum_{j \in \mathbb{Z}} 2^{2j} \|Q_j f\|_{L^1} \\
(2.4) \quad &= \|e^{R_1 t} P_0\|_{L^\infty} \|f\|_{\dot{B}_{1,1}^2}.
\end{aligned}$$

2. *Estimate of (2.3):* We want to estimate in L^∞

$$(2.5) \quad e^{R_1 t} P_0(x) = \int_{|\xi| \sim 1} \varphi(\xi) e^{ix \cdot \xi - it \frac{\xi_1}{|\xi|}} d\xi.$$

The standard idea is to use integration by parts. Letting

$$\Phi(\xi) := \nabla_\xi \phi(\xi) = \frac{x}{t} + \frac{\xi_2}{|\xi|^3} \xi^\perp, \quad \xi^\perp = (-\xi_2, \xi_1)^\top,$$

we consider the differential operator

$$\mathcal{L} := \frac{\text{Id} - i\Phi(\xi, x) \cdot \nabla_\xi}{1 + t |\Phi(\xi, x)|^2}$$

which leaves the exponential in (2.5) invariant:

$$(2.6) \quad \mathcal{L}e^{it\phi(\xi)} = e^{it\phi(\xi)}.$$

Integrating by parts gives

$$\begin{aligned} \int_{|\xi| \sim 1} \varphi(\xi) e^{ix \cdot \xi - it \frac{\xi_1}{|\xi|}} d\xi &= \int_{|\xi| \sim 1} \varphi(\xi) \mathcal{L}(e^{ix \cdot \xi - it \frac{\xi_1}{|\xi|}}) d\xi \\ &= \int_{|\xi| \sim 1} \mathcal{L}^\top(\varphi)(\xi) e^{ix \cdot \xi - it \frac{\xi_1}{|\xi|}} d\xi \end{aligned}$$

where

$$(2.7) \quad \mathcal{L}^\top(\varphi) = \frac{\varphi}{1 + t |\Phi(\xi, x)|^2} + \operatorname{div}_\xi \left(\frac{i\Phi}{1 + t |\Phi(\xi, x)|^2} \right) \varphi + \frac{i\Phi}{1 + t |\Phi(\xi, x)|^2} \nabla_\xi \varphi.$$

Hence

$$\|e^{R_1 t} P_0\|_{L_x^\infty} \leq \int_{|\xi| \sim 1} \left\| \mathcal{L}^\top \varphi(\xi) \right\|_{L_x^\infty} d\xi.$$

Remark 2.2. In estimating $\|\mathcal{L}^\top \varphi(\xi)\|_{L_x^\infty}$ two simple points seem noteworthy:

- (1) The L^∞ norm of φ and its derivatives pose no problems and can just be absorbed into the constant of our estimate.
- (2) The derivatives of Φ in ξ are independent of x and t , and uniformly bounded near $|\xi| \sim 1$, so they can equally be absorbed into the constant.

We estimate the three pieces in (2.7) separately.

In the second term we have

$$\operatorname{div}_\xi \left(\frac{i\Phi}{1 + t |\Phi(\xi, x)|^2} \right) = \frac{\operatorname{div}_\xi \Phi}{1 + t |\Phi|^2} - \frac{2t\Phi \cdot \nabla_\xi (|\Phi|^2)}{(1 + t |\Phi|^2)^2},$$

so this can be bounded by $\frac{C}{1 + t |\Phi(\xi, x)|^2}$. All in all we see that we only have to bound the integrals

$$\int_{|\xi| \sim 1} \frac{1}{1 + t |\Phi(\xi, x)|^2} d\xi \quad \text{and} \quad \int_{|\xi| \sim 1} \frac{\Phi}{1 + t |\Phi(\xi, x)|^2} d\xi.$$

The second term is bounded by the simple observation that by Cauchy-Schwarz

$$|\Phi| = \frac{1}{\sqrt{t}} \sqrt{t} |\Phi| \leq \frac{1}{\sqrt{t}} (1 + t |\Phi|^2),$$

so that

$$\int_{|\xi| \sim 1} \frac{|\Phi|}{1 + t |\Phi(\xi, x)|^2} d\xi \leq \frac{1}{\sqrt{t}} \int_{|\xi| \sim 1} 1 d\xi \leq \frac{C}{\sqrt{t}}.$$

Regarding the first term we claim (and will prove below):

Lemma 2.3.

$$(2.8) \quad \int_{|\xi| \sim 1} \frac{1}{1 + t |\Phi(\xi, x)|^2} d\xi \leq \frac{C}{\sqrt{t}},$$

with a constant $C > 0$ that is independent of x .

Putting together all the pieces we see that we have proved that

$$\|e^{R_1 t} P_0\|_{L^\infty} \leq \frac{C}{\sqrt{t}}.$$

In conjunction with (2.4) this shows the claim. \square

Proof of Lemma 2.3. Note: For $t \leq T$ for some fixed T this is trivial, so we need only concern ourselves with large t .

Straight from the definition of Φ we have

$$|\Phi|^2 \geq \left(\frac{|x|}{t} - \frac{|\xi_2|}{|\xi|^2} \right)^2,$$

so that we can estimate the integrand in Lemma 2.3 by

$$\begin{aligned} \int_{|\xi| \sim 1} [1 + t |\Phi|^2]^{-1} d\xi &\leq \int_{|\xi| \sim 1} \left[1 + t \left(\frac{|x|}{t} - \frac{|\xi_2|}{|\xi|^2} \right)^2 \right]^{-1} d\xi \\ (2.9) \quad &\leq \int_{r \sim 1} \int_0^{2\pi} \frac{1}{1 + t \left(\frac{|x|}{t} - \frac{|\sin \theta|}{r} \right)^2} d\theta r dr \\ &= t \int_{\rho \sim \frac{1}{\sqrt{t}}} \int_0^{2\pi} \frac{\rho^2}{\rho^2 + \left(\frac{|x|}{\sqrt{t}} \rho - |\sin(\theta)| \right)^2} d\theta \rho d\rho \end{aligned}$$

after changing variables to $\rho := \frac{r}{\sqrt{t}}$.

We let $b := \frac{|x|}{\sqrt{t}}$ and note that

$$\int_0^{2\pi} \frac{\rho^2}{\rho^2 + (b\rho - |\sin(\theta)|)^2} d\theta = 4 \int_0^{\frac{\pi}{2}} \frac{\rho^2}{\rho^2 + (b\rho - |\sin(\theta)|)^2} d\theta.$$

The delicate part of this integral is where $(b\rho - |\sin(\theta)|)^2$ is small. We thus distinguish three main cases depending on the size of the key parameter $b\rho = \frac{|x|}{\sqrt{t}}\rho \sim \frac{|x|}{t}$:

Case 1: $\frac{|x|}{t} \geq 1 + \epsilon$: We can then bound the integrand by

$$\int_0^{\frac{\pi}{2}} \frac{\rho^2}{\rho^2 + (b\rho - |\sin(\theta)|)^2} d\theta \lesssim \epsilon^{-2} \rho^2.$$

Case 2: $1 - \epsilon \leq \frac{|x|}{t} \leq 1 + \epsilon$: In this case $(b\rho - |\sin(\theta)|)^2$ is small on a set of size ϵ^2 , and away from it the integrand can be bounded by $\epsilon^{-2} \rho^2$, i.e.

$$\int_0^{\frac{\pi}{2}} \frac{\rho^2}{\rho^2 + (b\rho - |\sin(\theta)|)^2} d\theta \leq \epsilon^2 + \epsilon^{-2} \rho^2.$$

Case 3: $\frac{|x|}{t} \ll 1$: Here we have

$$\int_0^{\frac{\pi}{2}} \frac{\rho^2}{\rho^2 + (b\rho - |\sin(\theta)|)^2} d\theta \leq C\rho^2 + \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{\rho^2}{\rho^2 + (b\rho - |\sin(\theta)|)^2} d\theta.$$

This last integral we now estimate using a dyadic decomposition. We localize as follows: for the negative integers $j \leq 0$ we see that the set where $b\rho - |\sin(\theta)| \sim 2^j$ has size $\sim 2^j$, so that we have the bound

$$\begin{aligned}
 \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{\rho^2}{\rho^2 + (b\rho - |\sin(\theta)|)^2} d\theta &\leq \sum_{j=-\infty}^0 2^j \frac{\rho^2}{\rho^2 + 2^{2j}} \\
 &= \sum_{2^j < \rho} 2^j \frac{\rho^2}{\rho^2 + 2^{2j}} + \sum_{1 \geq 2^j \geq \rho} 2^j \frac{\rho^2}{\rho^2 + 2^{2j}} \\
 &\leq \sum_{2^j < \rho} 2^j + \sum_{1 \geq 2^j \geq \rho} 2^j \frac{\rho^2}{2^{2j}} \\
 &= \sum_{j=-\infty}^{\log_2(\rho)} 2^j + \rho^2 \sum_{j=\log_2(\rho)}^0 2^{-j} \\
 &= \sum_{j=-\log_2(\rho)}^{\infty} 2^{-j} + \rho^2 \sum_{j=0}^{-\log_2(\rho)} 2^j \\
 &\sim \rho \sum_{j=0}^{\infty} 2^{-j} + \rho^2(\rho^{-1} - 1) \\
 &= \rho + (\rho - \rho^2).
 \end{aligned}$$

Altogether we thus have the bound

$$\int_0^{\frac{\pi}{2}} \frac{\rho^2}{\rho^2 + \left(\frac{|x|}{\sqrt{t}}\rho - |\sin(\theta)|\right)^2} d\theta \leq \max\{\epsilon^{-2}\rho^2, \epsilon^2 + \epsilon^{-2}\rho^2, \rho\} \leq \rho,$$

once we choose $\epsilon \sim \sqrt{\rho}$.

Inserting this into (2.9) we have

$$\begin{aligned}
 \int_{|\xi| \sim 1} [1 + t|\Phi|^2]^{-1} d\xi &\leq t \int_{\rho \sim \frac{1}{\sqrt{t}}} \rho^2 d\rho \\
 &\sim t \left(\frac{1}{\sqrt{t}}\right)^3 = \frac{1}{\sqrt{t}},
 \end{aligned}$$

thus finishing the proof of the lemma. \square

Remark 2.4 (Dispersion for models with $1 < \alpha \leq 2$). The SQG equation can be viewed as the special case $\alpha = 1$ of the broader class of models

$$\partial_t \theta + u \cdot \nabla \theta = \frac{\partial_1}{|\nabla|^\alpha} \theta, \quad 1 \leq \alpha \leq 2,$$

with $u = \nabla^\perp (-\Delta)^{-\alpha/2} \theta = (-\frac{\partial_2}{|\nabla|^\alpha} \theta, \frac{\partial_1}{|\nabla|^\alpha} \theta)$. We note that for $\alpha = 2$ this yields the well-known *β -plane equation* (see e.g. [22]).

It turns out that for the cases $\alpha > 1$ the dispersion of the linear semigroups can be computed using standard stationary phase methods. More precisely, we have:

$$e^{\frac{\partial_1}{|\nabla|^\alpha} t} f(x) = \int_{\mathbb{R}^2} \hat{f}(\xi) e^{ix \cdot \xi - it \frac{\xi_1}{|\xi|^\alpha}} d\xi = \int_{\mathbb{R}^2} \hat{f}(\xi) e^{it\phi_\alpha(\xi)} d\xi,$$

where

$$\phi_\alpha(\xi) := \frac{x}{t} \xi - \frac{\xi_1}{|\xi|^\alpha}.$$

Then

$$\nabla_\xi \phi_\alpha = \frac{x}{t} + |\xi|^{-\alpha-2} ((\alpha-1)\xi_1^2 + \xi_2^2, \alpha\xi_1\xi_2),$$

so that for every $(t, x) \in \mathbb{R} \times \mathbb{R}^2$ there are at most two stationary points. A direct computation then gives the determinant of the Hessian of ϕ_α as

$$\det H_{\phi_\alpha} = \alpha^2 \frac{(\alpha-1)\xi_1^2 + \xi_2^2}{|\xi|^{4+2\alpha}},$$

which is non-degenerate away from the origin.

Rescaling as in the proof of Proposition 2.1 now yields

$$\begin{aligned} e^{\frac{\partial_1}{|\nabla|^\alpha} t} P_j(x) &= \int_{\mathbb{R}^2} \varphi(2^{-j}\xi) e^{ix \cdot \xi - it \frac{\xi_1}{|\xi|^\alpha}} d\xi \\ &= 2^{2j} \int_{\mathbb{R}^2} \varphi(\xi) e^{i2^j x \cdot \xi - it 2^{j(1-\alpha)} \frac{\xi_1}{|\xi|^\alpha}} d\xi \\ &= 2^{2j} e^{\frac{\partial_1}{|\nabla|^\alpha} 2^{j(1-\alpha)} t} P_0(2^j x), \end{aligned}$$

so that we obtain

$$\left\| e^{\frac{\partial_1}{|\nabla|^\alpha} t} P_j \right\|_{L^\infty} = 2^{2j} \left\| e^{\frac{\partial_1}{|\nabla|^\alpha} 2^{j(1-\alpha)} t} P_0 \right\|_{L^\infty} \lesssim 2^{2j} (2^{j(1-\alpha)} t)^{-1} = 2^{j(1+\alpha)} t^{-1}$$

from the stationary phase lemma. The full dispersive estimate thus reads

$$(2.10) \quad \left\| e^{\frac{\partial_1}{|\nabla|^\alpha} t} f \right\|_{L^\infty} \lesssim t^{-1} \|f\|_{\dot{B}_{1,1}^{1+\alpha}}, \quad 1 < \alpha \leq 2.$$

As announced we now turn to the question of the sharpness of the dispersive estimate.

2.2. Sharpness of the Dispersive Estimate. It turns out that for all smooth, radial L^1 functions f with $f(0) \neq 0$, the linear evolution $e^{Rt} f$ decays no faster than $\frac{1}{\sqrt{t}}$ in L^∞ :

Proposition 2.5. *Let f be a smooth, radial L^1 function with $f(0) \neq 0$.*

Then

$$e^{Rt} f(0) \sim f(0) \sqrt{\frac{2}{\pi t}} \cos\left(t - \frac{\pi}{4}\right) \quad \text{as } t \rightarrow \infty.$$

Proof. Let f be a smooth, radial function. Then \hat{f} is radial, so in particular

$$\begin{aligned} e^{Rt} f(0) &= \int_{\mathbb{R}^2} e^{-i\frac{\xi_1}{|\xi|}t} \hat{f}(\xi) d\xi = \int_0^\infty \hat{f}(r)r dr \int_0^{2\pi} e^{-it \cos(\vartheta)} d\vartheta \\ &= f(0) \int_0^{2\pi} e^{-it \cos(\vartheta)} d\vartheta. \end{aligned}$$

It turns out that

$$\int_0^{2\pi} e^{-it \cos(\vartheta)} d\vartheta = J_0(t),$$

where J_0 is the zeroth order Bessel function of the first kind. The asymptotics of this function are well known:

Lemma 2.6 (see page 364 in [1]). *For $t \gg 1$*

$$J_0(t) \sim \sqrt{\frac{2}{\pi t}} \cos\left(t - \frac{\pi}{4}\right).$$

This concludes the proof of the proposition. \square

3. INCREASED TIME OF EXISTENCE OF SOLUTIONS TO THE SQG EQUATION

Consider the equation

$$(3.1) \quad \partial_t \theta + u \cdot \nabla \theta = R_1 \theta$$

for $\theta : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}$ and

$$u = \nabla^\perp (-\Delta)^{-1/2} \theta = (-R_2 \theta, R_1 \theta),$$

where R_j denotes the Riesz transform in x_j , $j = 1, 2$.

Using the energy method and the Gagliardo-Nirenberg inequality one can prove the following blow-up criterion:

Lemma 3.1. *Let θ be a solution of the dispersive SQG equation (3.1) defined on a time interval containing $[0, T]$. Then for any $s > 0$ we have the bound*

$$(3.2) \quad \|\theta(T)\|_{H^s} \leq \|\theta_0\|_{H^s} \exp\left(c \int_0^T \|\nabla u(t)\|_{L^\infty} + \|\nabla \theta(t)\|_{L^\infty} dt\right)$$

for some universal constant $c > 0$.

Proof. We note that for the operator $J^s := (1 - \Delta)^{s/2}$ we have $\|J^s \theta\|_{L^2} \sim \|\theta\|_{H^s}$, so it suffices to bound $J^s \theta$ in L^2 . Applying J^s to the equation we see that

$$\partial_t J^s \theta + J^s (u \cdot \nabla \theta) = J^s R_1 \theta = R_1 J^s \theta.$$

Next we multiply by $J^s \theta$ and integrate over \mathbb{R}^2 to obtain

$$(3.3) \quad \partial_t \|J^s \theta\|_{L^2}^2 + \langle J^s (u \cdot \nabla \theta), J^s \theta \rangle_{L^2} = 0,$$

since for any $f \in L^2$ we have $\int f R_1 f = 0$.

We note that u is divergence-free, so

$$\langle u \cdot \nabla (J^s \theta), J^s \theta \rangle_{L^2} = 0,$$

and thus by the Gagliardo-Nirenberg inequality

$$|\langle J^s(u \cdot \nabla \theta), J^s \theta \rangle_{L^2}| \lesssim \|J^s \theta\|_{L^2}^2 (\|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}).$$

Combining this with (3.3) yields

$$\partial_t \|J^s \theta\|_{L^2}^2 \lesssim (\|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) \|J^s \theta\|_{L^2}^2,$$

so we need only appeal to Grönwall's Lemma to finish the proof. \square

We are now in the position to prove one of the main theorems of this article, thus demonstrating that solutions to the dispersive surface quasi-geostrophic equation (3.1) exist for a longer period of time than what standard hyperbolic estimates predict:

Theorem 3.2. *Let $\delta > 0$, $\mu > 0$. If $\|\theta_0\|_{H^{4+\delta}}, \|\theta_0\|_{W^{3+\mu,1}} \leq \epsilon$ there exists a unique solution $\theta \in C^0([0, T]; H^{4+\delta})$ of the initial value problem*

$$(3.4) \quad \begin{cases} \partial_t \theta + u \cdot \nabla \theta = R_1 \theta, \\ u = \nabla^\perp (-\Delta)^{-1/2} \theta, \\ \theta(0) = \theta_0 \in H^{4+\delta}, \end{cases}$$

with $T \sim \epsilon^{-4/3}$.

Moreover, for $t \in [0, T]$ we have $\|\theta(t)\|_{H^{4+\delta}} \lesssim \epsilon$.

The proof proceeds by bounding $\|\nabla u\|_{L^\infty}$ and $\|\nabla \theta\|_{L^\infty}$ using the dispersion and then invoking our previous Lemma 3.1.

Proof. By Duhamel's Principle we have

$$\theta(t) = e^{R_1 t} \theta_0 + \int_0^t e^{R_1(t-s)} (u(s) \cdot \nabla \theta(s)) ds.$$

For notational simplicity we introduce the two first-order pseudo-differential operators $L_1 := \nabla \nabla^\perp (-\Delta)^{-1/2}$ and $L_2 := \nabla$, for which $L_1 \theta = \nabla u$ and $L_2 \theta = \nabla \theta$.

Upon applying L_i to (3.1) and noting that $e^{R_1 \cdot}$ commutes with L_i we get

$$L_i \theta(t) = e^{R_1 t} L_i \theta_0 + \int_0^t e^{R_1(t-s)} L_i (u \cdot \nabla \theta)(s) ds.$$

Thus for any $\delta, \mu > 0$ the decay estimate (2.2) implies

$$\begin{aligned} \|L_i \theta(t)\|_{L^\infty} &\lesssim \frac{1}{\sqrt{t+1}} \|L_i \theta_0\|_{\dot{B}_{1,1}^2} + \int_0^t \frac{1}{\sqrt{t-s+1}} \|L_i (u \cdot \nabla \theta)\|_{\dot{B}_{1,1}^2} ds \\ &\lesssim \frac{1}{\sqrt{t+1}} \|\theta_0\|_{\dot{B}_{1,1}^3} + \int_0^t \frac{1}{\sqrt{t-s+1}} \|u \cdot \nabla \theta\|_{\dot{B}_{1,1}^3} ds \\ &\lesssim \frac{1}{\sqrt{t+1}} \|\theta_0\|_{W^{3+\mu,1}} + \int_0^t \frac{1}{\sqrt{t-s+1}} \|u \cdot \nabla \theta\|_{W^{3+\delta,1}} ds \\ &\lesssim \frac{1}{\sqrt{t+1}} \|\theta_0\|_{W^{3+\mu,1}} + \int_0^t \frac{1}{\sqrt{t-s+1}} \|\theta\|_{H^{4+\delta}}^2 ds. \end{aligned}$$

Now suppose that $\|\theta\|_{H^{4+\delta}} \leq 2\epsilon$ on a time interval $[0, T]$.² Then for $t \leq T$

$$\|L_i\theta(t)\|_{L^\infty} \leq c\epsilon \frac{1}{t^{1/2}} + \int_0^t \frac{c}{(t-s+1)^{1/2}} \epsilon^2 ds \leq c\left(\frac{\epsilon}{t^{1/2}} + \epsilon^2 t^{1/2}\right)$$

for a universal constant $c > 0$.

Now, using the energy inequality (3.2) from Lemma 3.1 we obtain

$$\|\theta(t)\|_{H^s} \leq \|\theta_0\|_{H^s} e^{c \int_0^t \frac{\epsilon}{s^{1/2}} + c^2 t^{1/2} dt} = \|\theta_0\|_{H^s} e^{c(\epsilon T^{1/2} + \epsilon^2 T^{3/2})}.$$

Choosing $s = 4 + \delta$ we see that $\|\theta\|_{H^{4+\delta}} \leq 2\epsilon$ so long as

$$\epsilon T^{1/2} + \epsilon^2 T^{3/2} \lesssim 1,$$

i.e. so long as $T \lesssim \epsilon^{-4/3}$. This completes the proof of the theorem. \square

Remark 3.3 (Longer Time of Existence for models with $1 < \alpha \leq 2$). For the related models discussed in Remark 2.4 (with parameter $1 < \alpha \leq 2$) one may go through the analogous steps of the above proof to obtain the condition

$$T(\log(T) - 1) \lesssim \epsilon^{-2},$$

thanks to the higher rate of dispersion (2.10). Hence we obtain an existence time of $T \sim \frac{\epsilon^{-2}}{|\log(\epsilon)|}$ for solutions in $H^{3+\alpha+\delta}$ with small initial data in $W^{2+\alpha+\mu,1}$ and $H^{3+\alpha+\delta}$ (for any given $\mu, \delta > 0$).

4. A STABILITY RESULT FOR THE INVISCID BOUSSINESQ SYSTEM

Recall from the introduction (page 3) the inviscid Boussinesq system

$$(4.1) \quad \begin{cases} \partial_t \omega + u \cdot \nabla \omega = \partial_x \rho, \\ \partial_t \rho + u \cdot \nabla \rho = 0, \\ u = \nabla^\perp (-\Delta)^{-1} \omega, \end{cases}$$

where $\omega, \rho : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}$.

In this section we draw the connection to the linear decay estimate of section 2 by linearizing the equations. We then show how this can provide a stability result for a certain stationary solution to the Boussinesq system (4.1).

To begin we notice that $\rho(t, x, y) = f(y)$ and $u \equiv \omega \equiv 0$ is a stationary solution of the Boussinesq system for any continuously differentiable function $f \in C^1$. Here we want to study the stability of the special stratified solution with $f(y) = -y$.

²We know that the maximal time interval $[0, T_{max}]$ where this is true has $T_{max} \geq c/\epsilon$.

4.1. Linearized Equations. The linearized equations around $(\rho, u, \omega) = (-y, 0, 0)$ read

$$(4.2) \quad \begin{cases} \partial_t \omega - \partial_x \rho = 0, \\ \partial_t \rho - u_2 = 0, \\ u_2 = \partial_x (-\Delta)^{-1} \omega. \end{cases}$$

Thus

$$\partial_{tt} \omega = \partial_t \partial_x \rho = \partial_x u_2 = \partial_{xx} (-\Delta)^{-1} \omega.$$

Upon taking the Fourier transform of both sides of this equation we get

$$\partial_{tt} \hat{\omega}(\xi, t) = -\frac{\xi_1^2}{|\xi|^2} \hat{\omega}(\xi, t),$$

the solutions of which are of the form

$$\hat{\omega}(\xi, t) = A(\xi) e^{-i \frac{\xi_1}{|\xi|} t} + B(\xi) e^{i \frac{\xi_1}{|\xi|} t},$$

from which one computes $u = \nabla^\perp (-\Delta)^{-1} \omega$ and $\hat{\rho}(\xi, t) = -|\xi|^{-1} A(\xi) e^{-i \frac{\xi_1}{|\xi|} t} + |\xi|^{-1} B(\xi) e^{i \frac{\xi_1}{|\xi|} t}$.

By Proposition 2.1 this implies that solutions of the linearized problem near $(\rho, u, \omega) = (-y, 0, 0)$ decay at the rate of $t^{-\frac{1}{2}}$ in L^∞ .

Remark 4.1. We note that the linearization around $(\rho, u, \omega) = (y, 0, 0)$ gives $\partial_{tt} \omega = -\partial_{xx} (-\Delta)^{-1} \omega$, so that the stationary solution $\rho = y$, $u \equiv \omega \equiv 0$ is *linearly unstable*. In the presence of viscosity this instability has been studied before (see for example [12, Chapter 8]).

4.2. Nonlinear Stability. Next we will show how this decay of the linear solutions can be used to establish the stability of the stationary solution $\rho = -y$, $u \equiv \omega \equiv 0$.

When perturbing around $(\rho, u, \omega) = (-y, 0, 0)$ we get the following system:

$$(4.3) \quad \begin{cases} \partial_t \omega + u \cdot \nabla \omega = \partial_x \rho, \\ \partial_t \rho + u \cdot \nabla \rho = u_2, \\ u = \nabla^\perp (-\Delta)^{-1} \omega. \end{cases}$$

We may write this as a perturbation of the linear system (4.2), namely

$$\begin{cases} \partial_t \omega - \partial_x \rho = F(u, \omega), \\ \partial_t \rho - u_2 = G(u, \rho), \\ u = \nabla^\perp (-\Delta)^{-1} \omega, \end{cases}$$

where $F = -u \cdot \nabla \omega$ and $G = -u \cdot \nabla \rho$ are quadratic.

We observe that these equations are of the same form as those in section 3, so the theory developed there can be carried over directly and yields the following

Theorem 4.2 (Increased time of stability of $(\rho, u, \omega) = (-y, 0, 0)$). *Let $\delta > 0$, $\mu > 0$, $\gamma > 0$. If $\|\omega_0\|_{H^{4+\delta}}$, $\|\omega_0\|_{H^{-1}}$, $\|\omega_0\|_{W^{3+\mu,1}}$, $\|\rho_0\|_{H^{5+\gamma}}$, $\|\rho_0\|_{W^{4+\mu,1}} \leq \epsilon$, then*

there exist unique functions $\rho \in C^0([0, T]; H^{5+\gamma})$, $\omega \in C^0([0, T]; H^{4+\delta} \cap H^{-1})$ such that the initial value problem for the Boussinesq system (4.1),

$$\begin{cases} \partial_t \omega + u \cdot \nabla \omega = \partial_x \rho, \\ \partial_t \rho + u \cdot \nabla \rho = 0, \\ u = \nabla^\perp (-\Delta)^{-1} \omega, \\ \omega(0) = \omega_0 \in H^{4+\delta} \cap H^{-1}, \\ \rho(0) = \rho_0 \in H^{5+\gamma}, \end{cases}$$

is solved by $(\rho - y, u, \omega)$. Here, $T \sim \epsilon^{-4/3}$.

In particular, small perturbations in $H^{4+\delta} \cap H^{-1}$ and $H^{5+\gamma}$ for ω and ρ (respectively) of the special solution $(\rho, u, \omega) = (-y, 0, 0)$ of (4.1) remain small for times $\lesssim \epsilon^{-4/3}$.

Remark 4.3. Given that ω is the vorticity of the fluid under consideration the requirement that $\|\omega_0\|_{H^{-1}} \leq \epsilon$ is physically sensible: It corresponds to the fluid having small kinetic energy.

Proof. Essentially the proof is the same as that of Theorem 3.2, so we only go over the two key ingredients: The dispersive estimate and the blow-up criterion.

Dispersive Estimate. As observed above, writing the equations for $(\rho - y, u, \omega)$ gives the system (4.3), the linear part (4.2) of which is

$$(4.4) \quad \begin{cases} \partial_t \omega - \partial_x \rho = 0, \\ \partial_t \rho - \partial_x (-\Delta)^{-1} \omega = 0, \end{cases}$$

and shares the decay properties of the SQG equation, hence the analogous estimates

$$\|(\nabla \rho, \omega)(t)\|_{L^\infty} \lesssim t^{-1/2} \|(\nabla \rho_0, \omega_0)\|_{\dot{B}_{1,1}^2}.$$

Blow-Up Criterion. We proceed in analogy with the SQG case: For $s \geq -1$ we take s derivatives of the first, and $s+1$ derivatives of the second equation in (4.4), multiply by $|\nabla|^s \omega$ and $|\nabla|^{s+1} \rho$ respectively, and integrate over \mathbb{R}^2 to obtain

$$\begin{aligned} \frac{1}{2} \partial_t \| |\nabla|^s \omega \|_{L^2} &= \langle |\nabla|^s \partial_x \rho, |\nabla|^s \omega \rangle_{L^2} - \langle |\nabla|^s (u \cdot \nabla \omega), |\nabla|^s \omega \rangle_{L^2}, \\ \frac{1}{2} \partial_t \| |\nabla|^{s+1} \rho \|_{L^2} &= \langle |\nabla|^{s+1} \partial_x (-\Delta)^{-1} \omega, |\nabla|^{s+1} \rho \rangle_{L^2} - \langle |\nabla|^{s+1} (u \cdot \nabla \rho), |\nabla|^{s+1} \rho \rangle_{L^2}. \end{aligned}$$

We add these two equalities to get

$$\frac{1}{2} \partial_t (\|\omega\|_{\dot{H}^s} + \|\rho\|_{\dot{H}^{s+1}}) = -\langle |\nabla|^s (u \cdot \nabla \omega), |\nabla|^s \omega \rangle_{L^2} - \langle |\nabla|^{s+1} (u \cdot \nabla \rho), |\nabla|^{s+1} \rho \rangle_{L^2},$$

since by integration by parts

$$\langle |\nabla|^s \partial_x \rho, |\nabla|^s \omega \rangle_{L^2} + \langle |\nabla|^{s+1} \partial_x (-\Delta)^{-1} \omega, |\nabla|^{s+1} \rho \rangle_{L^2} = 0.$$

But from this it follows by Gagliardo-Nirenberg estimates and Grönwall's Lemma (as in the proof of Lemma 3.1) that

$$\|\omega(T)\|_{\dot{H}^s} + \|\rho(T)\|_{\dot{H}^{s+1}} \lesssim (\|\omega_0\|_{\dot{H}^s} + \|\rho_0\|_{\dot{H}^{s+1}}) \times \exp\left(c \int_0^T \|\nabla u(t)\|_{L^\infty} + \|\nabla \theta(t)\|_{L^\infty} + \|\nabla \rho(t)\|_{L^\infty} dt\right).$$

Summation now gives the bound in inhomogeneous Sobolev spaces. \square

Acknowledgements. *The authors would like to thank Pierre Germain for his helpful comments.*

REFERENCES

- [1] Milton Abramowitz and Irene A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] A. Babin, A. Mahalov, and B. Nicolaenko. Global regularity of 3D rotating Navier-Stokes equations for resonant domains. *Appl. Math. Lett.*, 13(4):51–57, 2000.
- [3] J. Bedrossian and N. Masmoudi. Inviscid damping and the asymptotic stability of planar shear flows in the 2d Euler equations. *ArXiv e-prints*, June 2013.
- [4] Luis A. Caffarelli and Alexis Vasseur. Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation. *Ann. of Math. (2)*, 171(3):1903–1930, 2010.
- [5] Marco Cannone, Changxing Miao, and Liutang Xue. Global regularity for the supercritical dissipative quasi-geostrophic equation with large dispersive forcing. *Proc. Lond. Math. Soc. (3)*, 106(3):650–674, 2013.
- [6] Dongho Chae, Sung-Ki Kim, and Hee-Seok Nam. Local existence and blow-up criterion of Hölder continuous solutions of the Boussinesq equations. *Nagoya Math. J.*, 155:55–80, 1999.
- [7] Dongho Chae and Hee-Seok Nam. Local existence and blow-up criterion for the Boussinesq equations. *Proc. Roy. Soc. Edinburgh Sect. A*, 127(5):935–946, 1997.
- [8] J.-Y. Chemin, B. Desjardins, I. Gallagher, and E. Grenier. *Mathematical Geophysics*, volume 32 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press, Oxford University Press, Oxford, 2006.
- [9] Peter Constantin, Andrew J. Majda, and Esteban Tabak. Formation of strong fronts in the 2-D quasigeostrophic thermal active scalar. *Nonlinearity*, 7(6):1495–1533, 1994.
- [10] Peter Constantin and Vlad Vicol. Nonlinear maximum principles for dissipative linear nonlocal operators and applications. *Geom. Funct. Anal.*, 22(5):1289–1321, 2012.
- [11] Charles R. Doering and J. D. Gibbon. *Applied Analysis of the Navier-Stokes Equations*. Cambridge University Press, 1995. Cambridge Books Online.
- [12] P. G. Drazin. *Introduction to hydrodynamic stability*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2002.
- [13] P. G. Drazin and W. H. Reid. *Hydrodynamic Stability*. Cambridge University Press, second edition, 2004. Cambridge Books Online.
- [14] Tarek M. Elgindi. Inviscid damping for the incompressible porous medium equation, I, in preparation.
- [15] Z. Hassainia and T. Hmidi. On the inviscid Boussinesq system with rough initial data. *ArXiv e-prints*, February 2014.
- [16] J. K. Hunter, M. Ifrim, D. Tataru, and T. K. Wong. Long time Solutions for a Burgers-Hilbert Equation via a Modified Energy Method. *ArXiv e-prints*, January 2013.
- [17] A. Kiselev, F. Nazarov, and A. Volberg. Global well-posedness for the critical 2D dissipative quasi-geostrophic equation. *Invent. Math.*, 167(3):445–453, 2007.

- [18] Alexander Kiselev and Fedor Nazarov. Global regularity for the critical dispersive dissipative surface quasi-geostrophic equation. *Nonlinearity*, 23(3):549–554, 2010.
- [19] Xiaofeng Liu, Meng Wang, and Zhifei Zhang. Local well-posedness and blowup criterion of the Boussinesq equations in critical Besov spaces. *J. Math. Fluid Mech.*, 12(2):280–292, 2010.
- [20] Clément Mouhot and Cédric Villani. On Landau damping. *Acta Mathematica*, 207(1):29–201, 2011.
- [21] Serge Resnick. Dynamical problems in nonlinear advective partial differential equations. *PhD Thesis, University of Chicago*, 1995.
- [22] Jai Sukhatme and Leslie M. Smith. Local and nonlocal dispersive turbulence. *Physics of Fluids (1994-present)*, 21(5):–, 2009.
- [23] Zhaoyin Xiang and Wei Yan. On the well-posedness of the Boussinesq equation in the Triebel-Lizorkin-Lorentz spaces. *Abstr. Appl. Anal.*, 2012.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544, USA
E-mail address: tme2@math.princeton.edu

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, 251 MERCER STREET, NEW YORK 10012 NY, USA
E-mail address: klaus@cims.nyu.edu