

Dark Energy, Inflation, CMB Anisotropy and Polarization from Quantum Metric Fluctuations

Leonid Marochnik and Daniel Usikov

Physics Department, East-West Space Science Center, University of Maryland, College Park, MD 20742

Abstract

We propose a model of cosmological evolution of the early and late Universe which is consistent with observational data and naturally explains the origin of inflation and dark energy (DE). We show that the de Sitter accelerated expansion of the FLRW space with no matter fields (hereinafter “empty space”) is its natural state, and the model does not require either a scalar field or cosmological constant or any other hypotheses. Mathematically, this is due to the fact that the de Sitter state is an exact solution of the rigorous, mathematically consistent equations of one-loop quantum gravity for the empty FLRW space that are finite off the mass shell. Physically, this is due to the fact that the natural quantum metric fluctuations have the backreaction effect on the FLRW background, forming a self-polarized de Sitter graviton-ghost condensate which describes the condensation of gravitons on the horizon scale of the non-stationary Universe leading to its exponential expansion. The energy required to maintain the accelerated expansion is drawn from the graviton vacuum. At the start and the end of cosmological evolution, the Universe is assumed to be empty, which means that de Sitter expansion is a natural state of the Universe at the start and end of its cosmological evolution. The emptiness of the Universe at the start of its cosmological evolution automatically generates inflation. With an increase in the energy density of a new-born matter during inflation, the expansion begins to deviate from the de Sitter law, and eventually stops when the energy density of new-born matter reaches the first threshold which is the energy density of instantons. After that, the standard Big Bang cosmology begins. With a decrease in the energy density of matter by the end of the cosmological evolution of the Universe, the expansion first passes through the second threshold which again is the energy density of instantons. Here the quasi-de Sitter expansion (i.e. DE) is born, and then as the Universe empties, approaches the de Sitter law and asymptotically becomes such when the Universe becomes completely empty. This is observed as the DE effect. This scenario seems consistent with observational data. Existence of the first threshold explains the reason why the inflation is stopped. Existence of the second threshold explains why the DE acceleration is happening during the contemporary epoch of matter domination (coincidence problem). The Universe starts and ends with de Sitter expansion but the evolutionary process runs in these cases in opposite directions. It leads to the prediction that the signs of the parameter $1 + w$ should be opposite in both cases, and this fact is consistent with observations. The fluctuations of the number of gravitons lead to fluctuations of their energy density which in turn leads to the observed CMB temperature anisotropy of the order of $\sim 10^{-5}$ and CMB polarization. In the frame of this scenario, it is not a hypothetical scalar field that generates inflation and relic gravitational waves but on the contrary, the gravitational waves (gravitons) generate DE, inflation, CMB anisotropy and polarization.

1. Introduction

As is known, the idea of a possible key role of de Sitter solution for the early stages of evolution of the Universe goes back to de Sitter, Eddington and Lemaitre. Gliner [1] was the first to show that the equation of state $p = -\varepsilon$, generating de Sitter expansion of the Friedmannian Universe is the equation of state of anti-gravitating physical vacuum. On the basis of this idea, Zeldovich [2] showed that the cosmological constant producing the de Sitter expansion of the Friedmannian Universe is of the vacuum nature. In the pioneering work of Starobinsky [3], it was shown by direct computation that the quantum corrections to Einstein's equations that come from conformal anomalies lead to the nonsingular de Sitter solution instead of a standard Big Bang singularity (in some works, this solution is considered as an alternative to scalar field inflation, see e.g. [4]). The possibility of de Sitter expansion at the beginning of cosmological evolution was also found in works [5, 6]. Later on, it was turned out that the appearance of the de Sitter equation of state in the empty early Universe is not accidental because any quantum corrections to the Einstein equations lead to the energy density term of the form $\varepsilon = C \cdot \hbar H^4$ where C is a dimensionless constant, H is Hubble constant and \hbar is Planck constant. Such a form of the energy density term is simply a consequence of its dimension [7, section V]. It is usual to call space without matter fields empty. Hereinafter, the term "empty space" is used in this sense. In fact, space without matter fields is not empty, as it always has the natural quantum fluctuations of the metric, i.e. gravitons. Therefore, the empty (in this sense) space is filled with gravitons, which have the backreaction effect on its evolution over time. The quantum state of gravitons is determined by their interaction with a macroscopic field, and the macroscopic (background) geometry, in turn, depends on the state of gravitons. The background metric and the graviton operator appearing in the self-consistent theory are extracted from the unified gravitational field, which initially satisfies exact equations of quantum gravity. The classical component of the unified field is by definition a function of coordinates and time. The quantum component of the same unified field is described by a tensor operator function, which also depends on coordinates and time. Under such formulation of the problem, the original exact equations should be the operator equations of quantum theory of gravity in the Heisenberg representation (see below). The derivation of these equations and their relation to existing references can be found in [8]. For the first time, these equations (and exact solutions) were given by [9]. In work [9], the exact equations of one-loop quantum gravity for the empty space with FLRW geometry were obtained in the form of Bogoliubov-Born-Green-Kirkwood-Yvon hierarchy or BBGKY-chain. One of the exact solutions to the equations of the chain to the surprise of the authors turned out to be the de Sitter solution. The chain is invariant with respect to the Wick rotation $t = i\tau$, so there was a question as to how the real or imaginary time relates to this exact solution. It allows interpretation in terms of virtual gravitons [9, 7] as well as in terms of instanton fluctuations [7, 10]. In [7, 10], the arguments in favor of the exact de Sitter solution refer to the imaginary time. This means that in imaginary time (Euclidean space) quantum metric fluctuations form a self-consistent De Sitter gravitational instanton that can be thought of as describing the tunneling to the Lorentzian space of real time from "nothing" (see [10] and references therein). The topological non-equivalence between manifolds plays a role of a classically impenetrable barrier, quantum tunneling across which can create a flat inflationary Universe. In work [10], it was shown also that in the presence of matter, such tunneling is able to give birth to dark energy after the density of matter drops below a critical level, and the Universe becomes quite empty again. It is important to note

that the first rigorous mathematically consistent equations of one-loop quantum gravity which were obtained in works [7], [8] and [9] *are finite off the graviton mass shell*¹. In the empty space, they are the exact equations of self-consistent theory of gravitons in the Heisenberg representation with the ghost sector automatically providing one-loop finiteness off the mass shell. These equations are the only mathematically consistent of all in the available literature [8]. Because of conformal non-invariance and zero rest mass of gravitons, no conditions exist in the Universe to place gravitons on the mass shell precisely. Therefore, in the absence of one-loop finiteness, divergences arise in observables. To eliminate them, the Lagrangian of Einstein's theory must be modified, by amending the definition of gravitons. In other words, in the absence of one-loop finiteness, gravitons generate divergences, contrary to their own definition [7, 8]. Such a situation does not make any sense, so the one-loop finiteness off the mass shell is a prerequisite for internal consistency of the theory. As already noted, one of exact solutions to these equations of one-loop quantum gravity that are finite off the graviton mass shell is the de Sitter solution. Physically, it is the self-polarized condensate in the de Sitter space which describes the condensation of gravitons on the horizon scale of the non-stationary Universe. The theory of this macroscopic quantum effect is based only on the most general properties of quantum gravitational field in the non-stationary Universe, which are Bose-Einstein statistics, zero rest mass of gravitons, their conformal non-invariance and one-loop finiteness of quantum gravity off the graviton mass shell. The effect of condensation is clearly demonstrated by the fact that the state vectors have the form of quantum coherent superposition of vectors corresponding to different occupation numbers of gravitons and ghosts having the same wavelength of the order of the radius of 4-curvature. The solution describing the self-polarized condensate in the de Sitter space is presented in terms of field operators and state vectors. The analysis of this de Sitter solution is given in works [7, section VII] and [10]².

Thus, there is no need for the cosmological constant, hypothetical scalar fields and/or other hypotheses to generate the de Sitter exponential expansion of the empty FLWR Universe because such an expansion is its natural state.

2. Where does the Energy come from?

The quantum nature of this de Sitter solution sheds also a light on the nature of acceleration of the space expansion. The second Friedmannien equation reads

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\varepsilon + 3p)$$

Where $a(t)$ is a scale factor, dots are time derivatives and G is the gravitational constant. The accelerated expansion ($\ddot{a} > 0$) can take place only if $p < -\varepsilon/3$. In accordance with thermodynamics, the energy conservation law is $dE = -pdV$. The negative pressure leads to the fact that the total energy E of the Universe is increasing with the increasing of space volume V in the process of expansion. In de Sitter case, $p = -\varepsilon = \text{const}$ so that $E = \varepsilon V \sim a(t)^3$, i.e. in the process of expansion

¹ The one-loop quantum gravity with no matter fields is finite on the graviton mass shell [11]

² Technically, in both works [7] and [10] the solution to the equations of one-loop quantum gravity was obtained in the Euclidean space of imaginary time, and then was analytically continued (tunneled) into the Lorentz space of real-time. However, the process of transition to a real-time in these works is different, which leads to a difference in physical interpretation of the solutions. In this paper, we are following the procedure used in the work [10].

the energy of the Universe is increased indefinitely. To keep the energy density constant during de Sitter accelerated expansion, one needs a source of energy which is external to the space of the Universe. The question is where does such energy come from? The generally accepted interpretation of this fact (in case of DE, for instance) is that this is a feature of space itself. Our answer to this question is that the energy comes from the graviton vacuum. One can demonstrate this fact by simple qualitative consideration [9]. For the sake of clarity, we give it below. Let us to consider the balance of energy that is emerging in the space due to graviton creation and vanishing due to graviton annihilation. The characteristic energy of gravitons in these processes is $\hbar\omega \sim \hbar H$. Total probabilities of graviton creation and annihilation processes (normalized to the unity volume) w_{cr} and w_{ann} are proportional to the phase volume of one graviton $\omega^3/3\pi^2 \sim H^3/3\pi^2$. The exponent of the background–graviton coupling constant is unity if $\omega \sim H$. Thus, we obtain for w_{cr} and w_{ann} the following estimates

$$w_{cr} = \frac{\gamma}{3\pi^2} H^3 (\bar{N}_{\mathbf{k}} + 1)(\bar{N}_{-\mathbf{k}} + 1),$$

$$w_{ann} = \frac{\gamma}{3\pi^2} H^3 \bar{N}_{\mathbf{k}} \bar{N}_{-\mathbf{k}}.$$

Here $\gamma = O(1)$, $\bar{N}_{\pm\mathbf{k}} \sim N_g/2$ is the average number of gravitons with wavelengths that are near characteristic value $\omega \sim H$. Finally, we get the balance equation in the form

$$\begin{aligned} \varepsilon_g &= \hbar\omega(w_{cr} - w_{ann}) = \\ &= \frac{\gamma}{3\pi^2} \hbar H^4 (\bar{N}_{\mathbf{k}} + \bar{N}_{-\mathbf{k}} + 1) \simeq \frac{\gamma}{3\pi^2} N_g \hbar H^4 \end{aligned} \quad (1)$$

This estimate with accuracy of a numerical factor of the order of unity coincides with Eq. (8) which is obtained by exact computation [7]. Virtual gravitons with wavelength of the order of the horizon must appear and disappear in the graviton vacuum because of massless and conformal non-invariance of the graviton field. A non-zero balance of energy is due to the pure quantum process of spontaneous graviton creation, in other words, due to the uncertainty relation. The permanent creation and annihilation of virtual gravitons is not in exact balance because of the expansion of the Universe. The excess energy comes from the spontaneous process of graviton creation and is trapped by the background. This result is not unexpected. As is known, the vacuum Einstein equations with the cosmological constant Λ also produce the de Sitter solution in the FLRW metric corresponding to the equation of state $p_\Lambda = -\varepsilon_\Lambda$. A comprehensive review of the evolution of ideas that led to the understanding of Λ as a vacuum energy density can be found in work [12]. The important fact is that the physical vacuum with such an equation of state is an external source of energy for the empty FLRW space-time, and this is the reason why Λ is able to produce the accelerated expansion as well as de Sitter gravitational instanton [10].

3. Thresholds

At the first time, the Universe was empty (presumably) at the beginning of its cosmological evolution. The second time, it is going to become empty by the end of its cosmological evolution when the energy density of non-relativistic matter is approaching zero. Thus, one has to expect to find de Sitter expansion at the beginning and at the end of cosmological evolution of the Universe. At the beginning, it generates inflation, so that there is no need for hypothetical scalar fields. By the end, it generates the observed cosmological acceleration which is known as dark energy effect, and there is no need in the cosmological

constant or other hypotheses. If so, both the inflation and dark energy are one and the same macroscopic quantum effect of graviton condensation on the horizon scale of the non-stationary Universe. The instanton nature of this solution leads to the appearance of thresholds in both cases.

As was already mentioned, the exact de Sitter solution was obtained in imaginary time for the empty space and then analytically continued into real time. In the presence of matter, we have to follow the same procedure because there should exist a continuous transition to the case of empty space. In the Euclidean space of imaginary time $\tau = -it$, the first Friedmann equation reads [10]

$$3H_\tau^2 = 8\pi G(|\varepsilon_{inst}| - \varepsilon_m) \quad (2)$$

Where the energy density of instantons is ε_{inst} , the energy density of matter is ε_m and $H_\tau \equiv d \ln a / d\tau$ [10]. To be tunneled into the Lorentzian space of the Universe, solutions to (2) must exist. They do exist if the energy density of matter is below the threshold

$$\varepsilon_m \leq |\varepsilon_{inst}| \quad (3)$$

In the dark energy case, the energy density of non-relativistic matter is initially higher than this threshold, i.e. $\varepsilon_m > |\varepsilon_{inst}|$. So, when ε_m drops to the threshold (3) (and below) it marks the birth of dark energy which is a possible explanation to the fact that the birth of dark energy is taking place during the contemporary epoch of cosmological evolution (“coincidence problem”) [10]. In the case of inflation, the situation is reversed in the following sense. Presumably, there was no matter initially, so the Universe started with the natural de Sitter expansion. With time, the new-born matter began to appear; so that de Sitter expansion was changed to quasi-de Sitter, and finally, the inflation stops when the energy density of a new-born matter ε_M has increased up to the threshold $\varepsilon_M = |\varepsilon_{inst}|$. After that the standard Big Bang cosmology begins. It is convenient to characterize the difference between the cases of inflation and dark energy by the parameter $\delta = -\dot{H} / H^2$ (so-called “slow-roll” parameter in the scalar field theory) where the sign of δ indicates the direction of change in H . If the Hubble function H is increasing with time then $\delta < 0$ and vice versa. Note that $\delta = \delta_\tau$ where $\delta_\tau = -\dot{H}_\tau / H_\tau^2$ (here the dot is derivative over τ). In the case of DE, one has to expect an increase in H_τ with time because $\varepsilon_m \sim a^{-3}$ is decreasing (see Eq. (2)), and hence $\delta_{DE} < 0$. In the case of inflation, the energy density of a new-born matter ε_M is increasing with time (until it stops after it reaches the threshold (3)), so one has to expect $\delta_{inf} > 0$ in this case. Such behavior of parameter δ is consistent with observations (see section 5).

The following example is to illustrate the inflation case. The energy density of particles into which gravitons decayed satisfies a conservation equation [13]

$$\dot{\varepsilon}_M + 3H(\varepsilon_M + p_M) = \Gamma \varepsilon_g e^{-\Gamma(t-t_s)} \quad (4)$$

The right-hand side of (4) takes into account the flow of energy from gravitons. Here ε_g is energy density of gravitons forming de Sitter solution, Γ is the rate of decay of gravitons into other particles, and $t_{start} \equiv t_s$ is taken at the beginning of the process of filling the empty space with the new-born matter.

Assuming that products of decay of gravitons are highly relativistic, i.e. $\varepsilon_M = 3p_M$, we find the following solution to the Eq. (4)

$$\varepsilon_M = a^{-4} \int_{t_s}^t \Gamma \varepsilon_g a^4 e^{-\Gamma(t'-t_s)} dt' \quad (5)$$

The energy density of new-born matter (5) starts equal to zero at $t = t_s$ then first rises up to some maximum value and finally falls as the density is attenuated by the expansion of the Universe. The inflation is supposed to be stopped when the energy density of the new-born matter ε_M reaches the threshold $\varepsilon_M = |\varepsilon_{inst}|$. We do not know how primordial gravitons decay into other particles but we can consider some hypothetical example for pure illustrative purposes. In accordance with the inflation paradigm, at a sufficiently early stage of cosmological evolution all modes of interest must be deep inside the horizon which means that we deal with short wave gravitons. As is known [14], in this case³

$$\varepsilon_g = \varepsilon_g^0 (a_s / a)^4 \quad (6)$$

Taking into account Eq. (6), we get from (5)

$$\varepsilon_M = \varepsilon_g^0 \left(\frac{a_s}{a}\right)^4 [1 - e^{-\Gamma(t-t_s)}] \quad (7)$$

One can see from (7) that at $t = t_s$ the energy density of the new-born matter $\varepsilon_M(t_s) = 0$, then it increases to the maximum $\varepsilon_M^{\max} / \varepsilon_g^{\max} = (1 + 4H / \Gamma)^{-1}$ where $\varepsilon_g^{\max} = \varepsilon_g^0 (a_s / a_{\max})^4$, and after that the expansion continues with the usual Big Bang factor $\varepsilon_M \sim a^{-4}$. The inflation continues until the energy density of a new-born matter is increasing up to the threshold $\varepsilon_M = |\varepsilon_{inst}|$. After that it stops, and the standard Big Bang cosmology begins. This means that for the above example the condition $\varepsilon_M^{\max} \leq |\varepsilon_{inst}|$ must be satisfied. Existence of threshold in the DE case allows a natural explanation of the ‘‘coincidence problem’’ [10]. Solutions to Eq. (2) can exist only when the density of non-relativistic matter drops below the threshold (3) which corresponds to the redshift $1 + z_T = (\Omega_{DE} / \Omega_m)^{1/3}$. This threshold is consistent with observational data (see [10] for the discussion of DE case).

4. CMB anisotropy and polarization from fluctuations of the number of gravitons

³ Eq. (6) is valid for the gravitons of short wavelengths, i.e. for the condition $k\eta \gg 1$ where the conformal time is $\eta = \int dt / a$. In the process of expanding, the wavelengths of gravitons are stretched out, and after crossing the horizon they satisfy the reverse condition $k\eta \ll 1$ which produces the long wave modes for which instead of (6) we get $\varepsilon \sim a^{-2}$ and $\varepsilon \sim a^{-6}$ [15] (see also [7], section IV B).

The equation of state of graviton condensate forming de Sitter solution described above reads [7, 9] (cf. (1))

$$p = -\varepsilon = -\frac{3\hbar NH^4}{8\pi^2} \quad (8)$$

Here N is a dimensionless functional of parameters of state vector which is of the order of the number of gravitons in the Universe expanding with the Hubble velocity H . According to [7] (section VII), in such a condensate the phases in the graviton and ghost sectors correlate similarly, and the non-zero effect appears due to the difference of average occupation numbers for graviton's and ghost's instantons. In the Universe, the graviton condensate produced by quantum tunneling provided by instantons. The equation of state (8) is superficially similar to that which comes from quantum conformal anomalies [3]. Quantum corrections to the Einstein equations due to zero oscillations can provide a self-consistent De Sitter solution in the vicinity of Planck's value of curvature ([3], [16], [17]). In such a case, the equation of state is $\varepsilon \sim n\hbar H^4$ where the number of types of elementary particles n is of the order of ≤ 100 [18]. Conformal anomalies that arise due to regularization and renormalization procedures do not apply to the graviton condensate which is the solution of the equations of quantum gravity that are finite in the one-loop approximation. In the finite one-loop quantum gravity, the effect of conformal anomalies is exactly zero, and the De Sitter solution can be formed only by graviton-ghost instantons ([7], section X). In contrast to the conformal anomaly parameter n , the parameter N is arbitrary and can be a huge number. In our case, it is of the order of the number of gravitons in the Universe N_g , i.e. $N = \alpha N_g$ where $\alpha = O(1)$. In the contemporary Universe, we have $N_g \sim 10^{122}$ [9, 10].

The equation of state (8) was obtained under the assumption that the spectrum is flat and typical occupation numbers in the ensemble are large, so that squares of modules of probability amplitudes are likely to be described by Poisson distribution [7]. If so, from Eq. (8) it follows that in the energy units $\hbar = c = 1$ H reads

$$H^2 = 8\pi^2 \frac{M_{pl}^2}{\alpha N_g} \quad (9)$$

Where M_{pl} is the reduced Planck mass $M_{pl} = (\hbar c / 8\pi G)^{1/2} = 2.4 \cdot 10^{18} \text{ GeV} / c^2$. From (9) it follows that the fluctuations of the number of gravitons N_g that are presumably described by Gaussian distribution are $\langle (\Delta N_g)^2 \rangle / \langle N_g \rangle^2 = \langle N_g \rangle^{-1}$, so that

$$\frac{\langle (\Delta N_g)^2 \rangle}{\langle N_g \rangle^2} = \frac{\alpha}{8\pi^2} \cdot \frac{H^2}{M_{pl}^2} \quad (10)$$

Here $\langle (\Delta N_g)^2 \rangle = \langle N_g^2 \rangle - \langle N_g \rangle^2$. Due to Eq. (8), one gets

$$\frac{\langle (\Delta N_g)^2 \rangle}{\langle N_g \rangle^2} = \frac{\langle (\Delta \varepsilon)^2 \rangle}{\langle \varepsilon \rangle^2} \quad (11)$$

Thus, fluctuations of the number of gravitons produce fluctuations of energy density. They play the same role as scalar perturbations (density fluctuations) which are responsible for the anisotropy of CMB in the models of inflation using scalar fields. In other words, if the graviton condensate is responsible for the inflation then fluctuations of number of gravitons are the cause of the anisotropy of CMB. For the typical energy scale of inflation $H \approx 10^{15} \text{ GeV}$ and $\alpha = 1$ one gets from (10) and (11)

$$\left(\frac{\langle (\Delta \varepsilon)^2 \rangle}{\langle \varepsilon \rangle^2} \right)^{1/2} \approx 1.5 \cdot 10^{-5} \quad (12)$$

As is known, temperature fluctuations $\Delta T / T$ are of the same order of magnitude as the metric and density perturbations which contribute directly to $\Delta T / T$ via the Sachs-Wolfe effect. Thus, the fluctuations of the number of gravitons in the Universe are able to produce CMB anisotropy $\Delta T / T \sim 10^{-5}$ due to fluctuations of gravitational potential which in turn are of the order of fluctuations of energy density. Any mechanism generating the temperature anisotropy inevitably generates the CMB polarization as well, which is an order of magnitude below the temperature fluctuations ([19] and references therein). The B-mode of polarization of CMD is produced by gravitational waves. Over the de Sitter background, the mode functions of gravitons are (see, e.g. [9])

$$f(k\eta) = \left(1 - \frac{i}{k\eta}\right) e^{-ik\eta} \quad f^+(k\eta) = \left(1 + \frac{i}{k\eta}\right) e^{ik\eta} \quad (13)$$

The Bunch-Davies initial conditions select $f(k\eta)$ as a unique mode function in theories of inflation based on the scalar field hypothesis. Gravitons of instanton origin select the same $f(k\eta)$ as a unique mode function but because of a different reason⁴, so that there is no difference between power spectra of tensor perturbations produced by scalar fields and gravitons produced by instantons. Thus, both temperature fluctuations and B-mode of polarization of CMB can be of the same origin. Both of them can be produced by primordial gravitons.

5. Spectrum of metric fluctuations

The de Sitter solution considered above is produced by the flat spectrum of metric fluctuations [7, 10]. The spectrum slightly deviating from the flat spectrum produces the quasi-de Sitter solution. The observed tilt $n_s - 1$ of power spectrum k^{n_s-1} deviates slightly from the scale-invariant form corresponding to $n_s = 1$. The observed value is $n_s \approx 0.96$ [20]. This means that in reality we deal with a quasi-de Sitter expansion. To consider this case, we use Eq. (15) from [10] which can be presented in the following form

$$H^2 = \frac{6\pi^2 M_{pl}^2}{\int_0^\infty N_k [-\xi^2 + (1 + \xi)^2] e^{-2\xi} \xi d\xi} \quad (14)$$

⁴ Instanton solutions are sought in imaginary time by transition $\eta \rightarrow -i\eta$. Requirement of finiteness eliminates the $f^+(k\eta)$ solution (see [7] section VII and [11]), so only $f(k\eta)$ is the a unique mode function in this case

In the case of a flat spectrum, $N_k = N = \alpha N_g = \text{const}$, and (14) leads to (9) (after calculation of the integral in the denominator). Assuming that $N_k = N_0 (k / k_0)^\beta$ where k_0 is a pivot scale, we get from (14)

$$H^2 = \frac{6\pi^2 M_{pl}^2 (k_0 \eta)^\beta}{N_0 \int_0^\infty [-\xi^2 + (1 + \xi)^2] e^{-2\xi} \xi^{1+\beta} d\xi} \quad (15)$$

The straightforward calculation δ from (15) leads to $\delta = \beta / 2$. From Friedmann equations it follows that $\delta = 3(1 + w) / 2$ (where $w = p / \varepsilon$ is the equation of state parameter), so we get

$$\beta = 3(1 + w) \quad (16)$$

In the present approach, the deviations from de Sitter solution are due to any kind of matter that makes the space is non-empty. So, the parameter β is the one that characterizes the degree of by how much the space is filled by matter. In case of DE, the Planck data [21] obtained by combination of Planck+WP+BAO give, e.g. $w = -1.13_{-0.25}^{+0.24}$, i.e. $\beta_{DE} = 2\delta_{DE} \approx -0.39 < 0$, and the energy density of non-relativistic matter which is close to $\Omega_m \approx 0.3$. One can say that $\beta_{DE} \approx -0.39$ corresponds to the Universe, ~ 70 percent of which is already empty. Note the negative sign of β_{DE} as was expected for the dark energy case. Assuming homogeneous and isotropic Gaussian processes, we get the dimensionless variance for the two-point correlation function of fluctuations of the number of gravitons

$$\frac{\langle \delta N_k \delta N_{k'} \rangle}{N_k^2} = \frac{k^3}{2\pi^2} \frac{1}{N_k} = \frac{k^3}{2\pi^2} \frac{1}{N_0} \left(\frac{k}{k_0}\right)^{-\beta} \quad (17)$$

From (17) it follows that $n_s - 1 = -\beta$, and for $n_s \approx 0.96$, we get for the inflation case

$\beta_{\text{inf}} = 2\delta_{\text{inf}} \approx 0.04 > 0$. Note the positive sign of β_{inf} as was expected for the inflation case. Thus, the signs of parameter $\beta = 2\delta$ are consistent with the instanton origin of DE and inflation.

6. Conclusion

Based on the above consideration, the following scenario of cosmological evolution of the early and late Universe can be proposed. The de Sitter expansion is a natural state of empty FLRW space, so there is no need in scalar fields, cosmological constant and other hypotheses to generate such an expansion of the empty space. After the (quantum) birth of the Universe, it was presumably empty and the natural de Sitter expansion generates the inflation. In the process of filling the Universe with a new-born matter, de Sitter expansion becomes first quasi-de Sitter and then it stops when the energy density of the new-born matter reaches the threshold (energy density of instantons). After that the standard Big Bang cosmology begins. Nearing the end of the evolution, the Universe empties, and its expansion is approaching again the de Sitter law what is observed as the dark energy effect. The graviton vacuum is the external source of energy needed to maintain a constant energy density during the accelerated de Sitter expansion. This scenario is consistent with observational data. Fluctuations of the number of gravitons generate the CMB

anisotropy of the order of $\Delta T / T \sim 10^{-5}$ and inevitably they must generate the CMB polarization; the difference in signs of “slow-roll” parameter δ in cases of inflation and dark energy is consistent with the prediction of instanton theory; the threshold after which the dark energy appears is also consistent with observations (coincidence problem) [10].

Acknowledgment

We are grateful to Arthur Chernin for graciously agreeing to read our manuscript and make comments on our approach. One of us (LM) is grateful to Natalia Ptitsyna for stimulating discussions. We would like to express our deep appreciation and special thanks to Walter Sadowski for invaluable advice and help in the preparation of the manuscript.

References

1. E. B. Gliner , Soviet Physics JETP, 22, 378, 1966
2. Ya. B. Zeldovich, Soviet Physics –Uspekhi, 11,381, 1968
3. A.Starobinsky , Soviet Physics JETP Lett., 30, 682, 1979
4. G. Hinshaw et al. arXiv:1212.5226 v3, 2012
5. D. Kazanas, Astrophys. J., 241, L59,1980
6. K. Sato, Mon.Not.Roy.Astron.Soc., 195, 467,1981
7. L. Marochnik, D. Usikov and G. Vereshkov, J. Mod.Phys., 4, 48, 2013; arXiv:0811.4484, 2008
8. G. Vereshkov and L. Marochnik, J. Mod. Phys., 4, 285, 2013; arXiv: 1108.4256, 2011
9. L. Marochnik, D. Usikov and G. Vereshkov, Found. Phys., 38, 546, 2008; arXiv: 0709.2537, 2007
10. L. Marochnik, Grav. Cosmology, 19, 178, 2013; arXiv: 1306.6172, 2013
11. G. ‘t Hooft and M. Veltman , Ann. Inst. Henry Poincare, 20, 69, 1974
12. [P. J. E. Peebles](#) and [Bharat Ratra](#), Rev.Mod.Phys.75:559-606,2003; [arXiv:astro-ph/0207347](#)
13. S. Weinberg, Cosmology (section 4.2), Oxford, 2008
14. L. Marochnik, N. Pelikhov and G. Vereshkov, Astrophys. Space Sci., 34, 265, 1975
15. L.R.W. Abramo, R.H. Brandenberger and V.F. Mukhanov, Phys. Rev. D, 56, 3248, 1997
16. A. Starobinsky, Phys. Lett., B 91, 99, 1980

17. Ya. B. Zeldovich and A. Starobinsky, Sov. Phys. JETP 34, 1159, 1972
18. Ya. B. Zeldovich, Soviet Physics-Uspekhi, 24, 216, 1981
19. M. Kamionkowski, A. Kosowsky and A. Stebbins, Phys. Rev. D, 55, 7368, 1997, arXiv: astro-ph/9611125, v1, 1996
20. E. Komatsu et al., Astrophys. J. Suppl. 180,330, 2009; arXiv: 0803.0547v2, 2008
21. P.A.R. Ade et al., arXiv: 1303.5076, 2013