

IN WHAT CONDITION CAN THE 125 GEV HIGGS BOSON DECAY TO A PAIR OF ON-SHELL W-BOSONS?

Rasmiyya E. Gasimova^{1,2,3}, Vali A. Huseynov^{2,3,4}

¹Shamakhy Astrophysical Observatory National Academy of Sciences of Azerbaijan,
Y. Mammadaliyev Settlement, Shamakhy District, AZ5618, Azerbaijan;

²Department of Theoretical Physics, Baku State University, Z. Khalilov 23, AZ 1148, Baku,
Azerbaijan;

³Department of Physics, Qafqaz University, Baku-Sumgayit Road, 16 km., Khirdalan, Baku,
AZ0101, Azerbaijan;

⁴Department of General and Theoretical Physics, Nakhchivan State University, University Campus,
AZ 7012, Nakhchivan, Azerbaijan;

E-mail: gasimovar@yahoo.co.uk

E-mail: vgusseinov@yahoo.com

ABSTRACT

We have determined that the decay of the neutral boson at a mass around 125 GeV into an on-shell W^-W^+ -pair in a uniform magnetic field becomes, in principle, possible and the new decay channel of this boson in a magnetic field is allowed by the energy and total angular momentum conservation laws. The required magnetic field strength for observation of the measurable effect is $\sim 10^{23}$ G (or in Teslas $\sim 10^{19}$ T). The existence of the other neutral boson with the spin $J = 0$ and with the other mass is not either excluded in the mass range below $2m_W \cong 160.77$ GeV .

Recently a new neutral boson (NB) at a mass around 125 GeV [1, 2] with properties compatible with the Standard Model Higgs boson was discovered in the ATLAS and CMS experiments. This boson is described with $J^{PC} = 0^{++}$ where P is the parity, C is the charge conjugation, J is the spin. One of the decay channels of the discovered Higgs-like boson is $H \rightarrow WW^*$. One of these W^\mp -bosons is on-shell, the other one (W^*) is off-shell. According to the energy conservation law the decay of this NB into the on-shell W^-W^+ -boson pair is impossible because of $m_H < 2m_W$, where $m_W \cong 80.385$ GeV [3] is the W -boson mass. Therefore, this NB decays into one on-shell W -boson and one off-shell W^* -boson. The following natural questions arise. In what condition can the 125 GeV Higgs boson decay to a pair of on-shell W -bosons? How is realistic and promising the decay of the NB at a mass around 125 GeV into the two on-shell W -bosons in a magnetic field (MF)? Search for the answers to these questions determines the motivation for the presented investigation. We investigate the decay of the NB at a mass around 125 GeV observed at the LHC into a pair of on-shell W -bosons in a MF. The main purpose of this work is to determine the condition in what the NB at a mass around 125 GeV can decay to a pair of on-shell W -bosons?

The energy of a W^\mp -boson in a homogenous MF is given by the formula [4, 5]

$$E_{W^\mp}^2 = p_{\mp z}^2 + (2n_{\mp} + 1 - 2q_{\mp}s_{\mp z})eB + m_W^2, \quad (1)$$

where $B = |\vec{B}|$ is the strength of a MF whose intensity vector \vec{B} is directed along the Oz -axis, $p_{\mp z}$ ($s_{\mp z}$) are the third component of the momentum (spin) of a W^\mp -boson, $n_{\mp} = 0, 1, 2, \dots$ enumerates the Landau energy levels of a W^\mp -boson, $q_{\mp} = -1$ ($q_{\mp} = +1$) is the sign of the electric charge of a $W^-(W^+)$ -boson. A W^\mp -boson has three polarization states: $|W^\mp(s_{\mp} = 1, s_{\mp z} = +1)\rangle = |1, +1\rangle$, $|W^\mp(s_{\mp} = 1, s_{\mp z} = 0)\rangle = |1, 0\rangle$, $|W^\mp(s_{\mp} = 1, s_{\mp z} = -1)\rangle = |1, -1\rangle$, where s_{\mp} is the spin of a W^\mp -boson. Hereafter we will consider the case $n_{\mp} = 0$, $p_{\mp z} = 0$ and $q_{\mp}s_{\mp z} = +1$ that corresponds to the ground Landau level of the $W^-(W^+)$ -boson. When $s_{-z} = -1$ and $s_{+z} = +1$, we only have dealings with the states $|W^-(s_{-} = 1, s_{-z} = -1)\rangle = |1, -1\rangle$ and $|W^+(s_{+} = 1, s_{+z} = +1)\rangle = |1, +1\rangle$. In this case the W^\mp -boson energy satisfies the inequality $E_{W^\mp} = \sqrt{m_W^2 - eB} < m_W$ for an arbitrary B taken from the range $0 < B < B_{0W}$ where $B_{0W} = m_W^2/e$.

One of the main decay modes of the NB at a mass around 125 GeV is the $H \rightarrow WW^* \rightarrow l\nu l\nu$ channel [6]. According to the energy conservation law the decay of this NB into the on-shell W^-W^+ -boson pair is impossible. Therefore, this NB decays into one on-shell W -boson and one off-shell W^* -boson. However, if we place this NB in a uniform MF, the MF will affect on the W^- - and W^+ -bosons. If we consider the NB at a mass around 125 GeV in the rest frame and take into account the relation $E_{W^\mp} = \sqrt{m_W^2 - eB} < m_W$ for the $W^-(W^+)$ -boson with $s_{-z} = -1$ ($s_{+z} = +1$) in the energy conservation law $m_H = E_H = E_{W^-} + E_{W^+}$, we can see that in a sufficiently strong MF with the strength B_H the equality $m_H = 2\sqrt{m_W^2 - eB_H}$ is satisfied and the decay of the NB with the mass around 125 GeV into the two on-shell W^\mp -bosons becomes, in principle, energetically possible. So, as a result of the decay of 125 GeV NB in a MF we have the final diboson system W^-W^+ that consists of the on-shell W^- - and W^+ -bosons situating in a MF. The quantum states of the system W^-W^+ can have the spin equal to 0, 1, 2. Using the two polarization states $|W^-(s_{-} = 1, s_{-z} = -1)\rangle = |1, -1\rangle$ and $|W^+(s_{+} = 1, s_{+z} = +1)\rangle = |1, +1\rangle$ of W^\mp -bosons and the addition rule of spins we obtain three polarization states of the system W^-W^+ : $|0, 0\rangle$, $|1, 0\rangle$, $|2, 0\rangle$. It should be noted that in case of the longitudinal polarization of the spin of the W^\mp -boson we have $s_{\mp z} = 0$ and from the formula (1) we derive $E =$

$\sqrt{m_W^2 + eB} > m_W$ for an arbitrary B taken from the range $0 < B < B_{0W}$ and obtain for the mass of the decaying NB the relation $m = 2\sqrt{m_W^2 + eB} > 2m_W$ that contradicts to the condition $m_H \cong 125 \text{ GeV} < 2m_W$. So, in the mass range $0 < m < 2m_W$ there is no sense to investigate the case of the longitudinal polarization of the spin of the on-shell $W^-(W^+)$ -boson in a MF. According to the energy conservation law the energy of the final W^-W^+ -system $E_{W^-W^+}$ is to be in the range $0 < E_{W^-W^+} < 2m_W$. The polarization states $|W^-(s_- = 1, s_{-z} = -1)\rangle = |1, -1\rangle$ and $|W^+(s_+ = 1, s_{+z} = +1)\rangle = |1, +1\rangle$ that form the three possible final states $|0, 0\rangle, |1, 0\rangle, |2, 0\rangle$ only satisfy the condition $0 < E_{W^-W^+} < 2m_W$. The energy conservation law is as $m = m_H = 2\sqrt{m_W^2 - eB_H}$ when $W^-(s_- = 1, s_{-z} = -1)$ and $W^+(s_+ = 1, s_{+z} = +1)$ are produced on the ground Landau level. Since we consider a NB in the rest frame, the mass of the decaying NB can not be zero: $m \neq 0$. Taking into account the condition $m \neq 0$ we obtain from the formula $m = m_H = 2\sqrt{m_W^2 - eB_H}$ that $B_H \neq B_{0W}$ and $B_H < B_{0W}$. So, the mass of the NB at a mass around 125 GeV satisfies the condition $0 < m_H < 2m_W$ ($0 < m_H < 160.77 \text{ GeV}$) for an arbitrary B taken from the range $0 < B_H < B_{0W}$.

We can determine the charge conjugation $C_{W^-W^+}$, the parity $P_{W^-W^+}$ and the total angular momentum J of the W^-W^+ -system by the formulae:

$$C_{W^-W^+} = (-1)^{L_{W^-W^+} + S_{W^-W^+}}, \quad (2)$$

$$P_{W^-W^+} = (-1)^{L_{W^-W^+}} P_{W^-} P_{W^+} = (-1)^{L_{W^-W^+}}, \quad (3)$$

$$J = L_{W^-W^+} + S_{W^-W^+}, L_{W^-W^+} + S_{W^-W^+} - 1, \dots, |L_{W^-W^+} - S_{W^-W^+}| \quad (4)$$

where $P_{W^-}(P_{W^+})$ is the intrinsic parity for the $W^-(W^+)$ -boson, $L_{W^-W^+}$ ($S_{W^-W^+}$) is the orbital quantum number (the total spin) for the W^-W^+ -system. We take into account that the NB at a mass around 125 GeV also decays into the two photons. Therefore its spin J can not be 1 according to the Landau-Yang theorem [7, 8] and the charge conjugation C of this NB is $C = C_H = 1$. The decay $H \rightarrow W^-W^+$ is a weak process and C is not conserved in this process. It means that if the charge conjugation of the initial neutral H -boson is $C_H = 1$ before the reaction $H \rightarrow W^-W^+$, the charge conjugation $C_{W^-W^+}$ might be $+1$ or -1 after the reaction, or it might also go to a state that is not a $C_{W^-W^+}$ eigenstate. Here we assume that $C_{W^-W^+}$ (and $P_{W^-W^+}$) is either $+1$ or -1 after the reaction. The following combinations of $C_{W^-W^+}$ and $P_{W^-W^+}$ for the W^-W^+ -system are possible: A) $C_{W^-W^+} = +1, P_{W^-W^+} = +1$; B) $C_{W^-W^+} = +1, P_{W^-W^+} = -1$; C) $C_{W^-W^+} =$

$-1, P_{W^-W^+} = +1$; D) $C_{W^-W^+} = -1, P_{W^-W^+} = -1$. We have obtained that only in the cases A and B the NB with the spin $J = 0$ can exist in the mass range $0 \leq m_H \leq 2m_W$ and it can decay into the on-shell $W^-(s_- = 1, s_{-z} = -1)$ - and $W^+(s_+ = 1, s_{+z} = +1)$ -bosons in a MF:

$$\text{case A: if } C_{W^-W^+} = +1 \text{ and } P_{W^-W^+} = +1, J = 0, 2 \text{ (} S_{W^-W^+} = 0, 2; L_{W^-W^+} = 0 \text{),} \quad (10)$$

$$\text{case B: if } C_{W^-W^+} = +1 \text{ and } P_{W^-W^+} = -1, J = 0, 2 \text{ (} S_{W^-W^+} = 1; L_{W^-W^+} = 1 \text{),} \quad (11)$$

We have obtained $J = 0, 2$ for the spin of the NB at a mass around 125 GeV if $C_{W^-W^+} = +1$. One NB with the spin $J = 0$ has already been observed in the mass range $0 < m_H \leq 2m_W$ by the ATLAS and CMS Collaborations [1, 2]. However, the existence of the other NB with the spin $J = 0$ and with the other mass is not excluded in the mass range $0 < m_H \leq 2m_W$. The analysis of the above considered cases A, B, C and D show that the existence of the NB at a mass around 125 GeV with the spin $J = 0$ is allowed in the cases A and B. When W^-W^+ -pair are produced on the ground Landau level, the orbital quantum number $L_{W^-W^+}$ should be minimal. $L_{W^-W^+}$ is minimal only in the case A. So, the case A is more suitable for the particle with the spin $J = 0$. The MF strength required for the decay of the NB at a mass around 125 GeV into the on-shell $W^-(s_- = 1, s_{-z} = -1)$ - and $W^+(s_+ = 1, s_{+z} = +1)$ -bosons is calculated by the formula $B_H = B_{0W}[1 - (m_H/2m_W)^2]$. Let us perform the simple numerical estimations for the strength of the MF required for the decay of the NB at a mass around 125 GeV into the on-shell $W^-(s_- = 1, s_{-z} = -1)$ - and $W^+(s_+ = 1, s_{+z} = +1)$ -bosons in a MF. When $m \cong 125 \text{ GeV}$, the required MF strength is $\sim 10^{23} \text{ G}$ (or in Teslas $\sim 10^{19} \text{ T}$). The strongest (pulsed) MF ever obtained in a laboratory is 28 MG [9] that is much less than $\sim 10^{22} \text{ G}$. The maximum strength of the produced strong MF in noncentral heavy-ion collisions in the direction perpendicular to the reaction plane is estimated to be $\sim 10^{17} \text{ G}$ at the RHIC and $\sim 10^{18} \text{ G}$ at the LHC [10-16]. In lead-lead collisions at the LHC, the strength of the generated MF may reach $\sim 10^{20} \text{ G}$ [11, 12]. We hope that in the future collider experiments, when the strength of the produced strong MF reaches the magnitude $\sim 10^{22+23} \text{ G}$, the decay of the NB at a mass around 125 GeV into the on shell W^- - and W^+ -bosons can be observed experimentally. The decay of the NB at a mass around 125 GeV into the on shell W^- - and W^+ -bosons, probably, could be realized in the early stages after the Big Bang when sufficiently strong MFs existed.

We have determined that the decay of the NB at a mass around 125 GeV into an on-shell W^-W^+ -pair in a uniform MF becomes, in principle, possible and the new decay channel of this boson in a MF is allowed by the energy and total angular momentum conservation laws. The required MF strength for observation of the measurable effect is $\sim 10^{23} G$ (or in Teslas $\sim 10^{19} T$). The existence of the other NB with the spin $J = 0$ and with the other mass is not either excluded in the mass range below $2m_W \cong 16077 \text{ GeV}$. We hope that the possibility of the decay of the NB at a mass around 125 GeV into a pair of on-shell W -bosons in a uniform MF will be of interest of cosmologists and particle astrophysicists and it will also attract the experimental physicists' attention in future collider experiments.

ACKNOWLEDGEMENTS

R. G. and V. H. are very grateful to the Organizing Committee of the LHCP2014 Conference for the kind invitation to attend this conference.

References

1. G. Aad *et al.* (ATLAS Collaboration), Phys. Lett. B **716**, 1 (2012).
2. S. Chatrchyan *et al.* (CMS Collaboration), Phys. Lett. B **716**, 30 (2012).
3. J. Beringer *et al.*, Particle Data Group, PR D **86**, 010001 (2012).
4. T. Goldman, W. Tsai, and A. Yildiz, Phys. Rev. D **8**, 1926 (1972).
5. A. Salam, J. Strathdee, Nucl. Phys. B **90**, 203 (1975).
6. ATLAS Collaboration. ATLAS-CONF-2012-098; ATLAS-COM-CONF-2012-138.
7. L. D. Landau, Dokl. Akad. Nauk USSR **60**, 207 (1948).
8. C. N. Yang, Phys. Rev. **77**, 242 (1950).
9. B. A. Boyko *et al.*, in 12th IEEE Intern. Pulsed Power Conf., Monterey, Calif., USA, 1999, Digest of Technical Papers (Eds C. Stallings, H. Kirbie) (New York: IEEE, 1999).
10. V. Skokov, A. Y. Illarionov, V. Toneev, Int. J. Mod. Phys. A **24**, 5925 (2009).
11. W. -T. Deng and X. -G. Huang, Phys. Rev. C **85**, 044907 (2012).
12. A. Bzdak and V. Skokov, Phys. Lett. B **710**, 171 (2012).
13. D. E. Kharzeev, L. D. McLerran and H. J. Warringa, Nucl. Phys. A **803**, 227 (2008).
14. M. N. Chernodub, Phys. Rev. D **82**, 085011 (2010).
15. M. N. Chernodub, Phys. Rev. Lett. **106**, 142003 (2011).
16. M. N. Chernodub, Lect. Notes Phys. **871**, 143 (2013).