

When Higgs Meets Starobinsky in the Early Universe

Mahdi Torabian*

School of Particles and Accelerators, Institute for Research in Fundamental Sciences (IPM), 19395-5531, Tehran, Iran

The measurement of the Higgs mass has confirmed that the Standard Model electroweak vacuum is a shallow local minimum and is not absolutely stable. In addition to a probable unacceptably fast tunneling to the deep true minimum, it is not clear how the observable present-day vacuum could be reached from the early Universe particularly following inflation. In this note it is shown that these problems can be alleviated if the Higgs field is non-minimally coupled to a higher-curvature theory of gravity which is effective in deriving inflation a la Starobinsky.

PACS numbers: 14.80.Bn, 12.60.Fr, 98.80.Cq

Introduction

The great achievement of the LHC was the discovery of the Higgs particle of mass $m_h = 125.15 \pm 0.24$ GeV [1, 2]. It completed the scalar sector of the Standard Model of particle physics with all parameters determined. In particular, the Higgs quartic self-coupling parameter is deduced at the electroweak scale to be $\lambda(m_{EW}) \approx 0.13$. This parameter is the only parameter of the Standard Model which is not multiplicatively renormalized. Given the central value of top quark mass $m_t = 173.2 \pm 0.9$ GeV [3], its beta-function β_λ at low energy is chiefly dominated by the top Yukawa coupling and so is negative. The Standard Model is a renormalizable theory and thus one could in principle extrapolate it to an arbitrary high energy and make predictions. If no new physics intervenes, the effective Higgs potential can be computed at desired loop orders. The quartic coupling is monotonically decreasing, vanishes at an intermediate energy of about 10^{10} GeV and subsequently turns negative [4]. At high energy the gauge interactions take over and make the beta function positive. Then, the quartic coupling is increasing as it develops a new minimum which will be the global one. The potential is plotted in Figure 1.




FIG. 1. The 2-loop effective potential for the Higgs $V_{\text{eff}}(\varphi) = \frac{1}{2}m^2(\mu)\varphi^2 + \frac{1}{4}\lambda(\mu)\varphi^4$ extrapolated to an arbitrary high scale. Magnified parts of the potential are shown as an ordered series of insets. The stability is disfavored at 98% C.L.

The global minimum, located at tens of m_{Pl} , is deeply inside the quantum gravity regime. This picture could radically be altered if one sensibly includes quantum gravity effects at high energies. Not understanding that thoroughly, it is reasonable to abandon the naive extrapolation to an arbitrary high energy and limit to the Planck scale. Thus, the potential would basically be seen ill as unbounded from below. The electroweak vacuum is a local one and barrier separating it from the deep well is extremely small. If one naively computes the tunneling rate between the vacua, one finds that life-time of the present-day electroweak vacuum is greater than the age of the Universe and thus the vacuum is *metastable* [5]. However, if one includes higher dimensional Planck-suppressed operators, the conclusion will be drastically changed and the electroweak vacuum turns *unstable* and quickly decays to true vacuum [6]. Consequently, due to a huge negative cosmological constant, it leads to a catastrophic gravitational collapse.

Furthermore, this picture of the Higgs potential raises problems in relation to the early Universe. In order to end up in the energetically-disfavored present-day electroweak vacuum and prevent the Higgs from rolling down to the true minimum, a fine-tuning at level of *one part in a hundred million* in the Higgs value is needed [7, 8]. Moreover, it being granted that the Higgs is initially near the desired local minimum, it will not stick to that in the presence of Hubble-size quantum fluctuations during an intermediate to high scale inflation [9, 10].

New physics beyond the Standard Model could possibly stabilize the Higgs potential. However, excellent agreement of the Standard Model predictions with the experimental results puts tight constraints on new physics as it must have marginal effect on the electroweak fit. Moreover, generically new physics would inevitably introduce a naturalness problem to the scalar sector.

Here it is shown that the the stability problem can be alleviated without introducing unnaturalness if the Higgs field is non-minimally coupled to higher-curvature theory of gravity which account for the dynamics in the early Universe. Therefore, basically the new ingredient is not new particles or interactions, but is to properly take into account the gravitational interaction at high energy.

The Higgs Sector Action

The dynamics of the Higgs field which is non-minimally coupled to a higher-curvature theory of gravity is given by the following action

$$S = \int d^4x (-g)^{1/2} \left(\frac{1}{2} m_{\text{Pl}}^2 f(\varphi, R) - \frac{1}{2} g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi - V(\varphi) \right), \quad (1)$$

where $\varphi^2 = 2H^\dagger H$ and

$$f(\varphi, R) = (1 - \xi m_{\text{Pl}}^{-2} \varphi^2) R + \alpha m_{\text{Pl}}^{-2} R^2, \quad (2)$$

$$V(\varphi) = \frac{1}{2} m^2 \varphi^2 + \frac{1}{4} \lambda \varphi^4. \quad (3)$$

The non-minimal parameter ξ could be zero at tree-level. However, being related to a dimension 4 operator, it is unstable to quantum fluctuations and would be generated at loop-level in a curved spacetime [11]. Moreover, the term quadratic in the Ricci scalar is the simplest generalization to General Relativity regarding higher-order gravity. Although being a higher-derivative theory of gravity, it is free from Ostrogradski classical instability or the presence of spin-2 ghost (and also spin-0 ghost for positive α) in the spectrum [12]. Furthermore, this terms is needed for the renormalization of the effective potential. Namely, the presence of higher derivative terms is needed if one demands renormalizability.

Through a Weyl transformation of the metric

$$g_{\mu\nu}^E = (1 - \xi m_{\text{Pl}}^{-2} \varphi^2 + 2\alpha m_{\text{Pl}}^{-2} R) g_{\mu\nu} \equiv e^{\tilde{\chi}} g_{\mu\nu}, \quad (4)$$

one can move to the Einstein frame with the following action

$$S_E = \int d^4x (-g_E)^{1/2} \left(\frac{1}{2} m_{\text{Pl}}^2 R_E - \frac{1}{2} g_E^{\mu\nu} \partial_\mu \chi \partial_\nu \chi - \frac{1}{2} e^{-\tilde{\chi}} g_E^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi - V_E(\varphi, \chi) \right). \quad (5)$$

The term quadratic in the Ricci scalar introduces a new propagating scalar, *a.k.a.* Weyl scalar, in the Einstein frame. The canonically normalized dimensionful Weyl scalar field is $\chi = (3/2)^{1/2} m_{\text{Pl}} \tilde{\chi}$. Note that the Weyl scalar has a canonical kinetic term while the Higgs scalar has a non-canonical one. In fact, the Weyl and the Higgs fields interact both via the kinetic term and the scalar potential. Finally, the mass parameter in the Jordan and the Planck mass in Einstein frame, which is shown symbolically the same, might have slight different value depending on the value of ξ .

Stability Condition

The scalar potential in the Einstein frame reads as

$$\begin{aligned} V_E(\varphi, \chi) &= e^{-2\tilde{\chi}} V(\varphi) + \frac{1}{8} \alpha^{-1} m_{\text{Pl}}^4 (1 - e^{-\tilde{\chi}} (1 - \xi m_{\text{Pl}}^{-2} \varphi^2))^2 \\ &= \frac{1}{8\alpha} (1 - e^{-\tilde{\chi}})^2 m_{\text{Pl}}^4 \\ &+ \frac{1}{4} e^{-2\tilde{\chi}} \left[2m^2 + (e^{\tilde{\chi}} - 1) m_{\text{Pl}}^2 \xi / \alpha + (\lambda + \xi^2 / 2\alpha) \varphi^2 \right] \varphi^2, \end{aligned} \quad (6)$$

The second line in equation (6) gives the pure Weyl potential which is flat for large χ values. When the potential is mainly dominated by that, it derives a Starobinsky-like R^2 -inflation in the early Universe [13]. The Planck results on the CMB anisotropy ($A_s = 2.23 \pm 0.16 \times 10^{-9}$ 68% C.L.) and the primordial gravitational waves ($r < 0.11$ 95% C.L.) [9] constraint the free parameter α as

$$\alpha = (12\pi^2 r A_s)^{-1} \gtrsim 3.4 \times 10^7. \quad (7)$$

The simplest manifestation of the Starobinsky inflation predicts $r \approx 2.5 \times 10^{-3}$ and therefore $\alpha \approx 1.5 \times 10^9$. However, modifications to the model predict larger r and so smaller α works as well (see [14]). For completeness, although not quite relevant to the present study, there is also an upper bound on the value of α from gravitational experiments measuring Yukawa correction to the Newtonian potential as $\alpha \lesssim 10^{61}$ [15, 16].

From the last line in equation (6) one finds that depending on the sign and the absolute value of ξ , different stable solutions can be obtained. If the value of ξ parameter is large enough such that at the Planck scale the following constraint is satisfied

$$\lambda(m_{\text{Pl}}) + \xi(m_{\text{Pl}})^2 / 2\alpha \geq 0, \quad (8)$$

then the Higgs quartic coupling is positive-definite. Given the value of the Higgs quartic coupling at the Planck scale $\lambda(m_{\text{Pl}}) \approx -0.01$ the above constraint implies that (for $\alpha \approx 10^9$)

$$\xi(m_{\text{Pl}}) \gtrsim 4500 \quad \text{or} \quad \xi(m_{\text{Pl}}) \lesssim -4500. \quad (9)$$

If $\xi > 0$, then the Higgs potential is convex for any value of the scalar fields. If $\xi < 0$, then the stability can be reached only if the Higgs quadratic term is suppressed through the following constraint on the field values

$$\tilde{\chi}_0 \lesssim 2 \ln \tilde{\varphi}_0 + \ln(-\xi/2). \quad (10)$$

Furthermore, an absolutely stable potential exists for smaller values of ξ when condition (8) is relaxed. In fact, the Weyl field needs to take large value so that the Higgs quadratic term takes over the quartic term and makes the potential convex. Thus, stable solution exists provided that $\xi > 0$ and the following condition is met

$$\tilde{\chi}_0 \gtrsim 16.1 + 2 \ln \tilde{\varphi}_0 - \ln \xi, \quad (11)$$

so that the Higgs field receives a huge effective mass.

The parameter ξ basically normalizes (defines) the 4-dimensional Planck mass in the Einstein frame. An upper limit exists only when it is negative as $\xi \lesssim 10^{32}$. Later, using collider bounds, it will be argued that a tighter constraint on its absolute value can be assumed.

Fields Evolution

The evolution of the Higgs and the Weyl field in the early Universe in a homogenous/isotropic background

$$ds^2 = -dt^2 + a(t) dx, \quad \varphi = \varphi(t), \quad \chi = \chi(t), \quad (12)$$




FIG. 2. Time evolution of the Higgs (up) and the Weyl (down) fields. The initial conditions are mentioned on the top.

is governed by the following equations of motion

$$\ddot{\chi} + 3H\dot{\chi} + 6^{-1/2}e^{-\tilde{\chi}}\dot{\varphi}^2 + V_{\chi}^E = 0, \quad (13)$$

$$\ddot{\varphi} + 3H\dot{\varphi} - (2/3)^{1/2}m_{\text{Pl}}^{-1}\dot{\chi}\dot{\varphi} + V_{\varphi}^E = 0, \quad (14)$$

subject to the following metric equations

$$3H^2m_{\text{Pl}}^2 = \frac{1}{2}\dot{\chi}^2 + \frac{1}{2}e^{-\tilde{\chi}}\dot{\varphi}^2 + V_E, \quad (15)$$

$$-2\dot{H}m_{\text{Pl}}^2 = \dot{\chi}^2 + e^{-\tilde{\chi}}\dot{\varphi}^2. \quad (16)$$

These are coupled second-order differential equations that can be solved by numerical methods. The solutions as scalar fields time evolution, in Planck units for field values and time, are plotted in Figure 2. The Higgs field initial value is of order one in Planck mass. Its initial velocity could also be chosen order one, however, it is found that it has insignificant qualitative effect on the solutions. On the other hand, the Weyl field initial conditions are chosen such that the Universe experiences at least 60 e-folds of exponential expansion.

The solutions are interpreted as follows. The Higgs and the Weyl fields are initially frozen for tens of Planck time until they commence harmonic oscillations about their local minima. Then, the Higgs field oscillates with large




FIG. 3. solutions for different ξ parameter: $\xi=1000$ (black), 5000 (green), 10000 (red), 15000 (blue), 20000 (yellow)

amplitude which is rapidly decreasing in time. Meanwhile, after a tiny jump in the value, the Weyl field oscillates with rather small amplitude which is slowly decreasing in time. After about a hundred thousand Planck time the Weyl field slowly rolls down the potential from its initial value to smaller values while the Higgs, whose value is reduced to $10^{-8}m_{\text{Pl}}$, is still rapidly oscillating. This epoch lasts for about 60 Hubble times after which both fields oscillate about their minima and the Universe is filled by the Bose condensates of Higgs and Weyl particles. After many damped oscillations fields settle down in their minima near the origin. Finally, they decay and reheat the Universe through the Higgs decay products.

Needless to say, fields have slightly different evolution, although qualitatively the same, depending on the value of the non-minimal coupling parameter. This change of behavior is summarized in Figure 3.

The time evolution of the Hubble parameter is given in Figure 4. It is fairly constant in two time periods which can be interpreted as two inflationary epochs. The former (when $H \approx 8.5 \times 10^{-3}m_{\text{Pl}}$) could last for only few e-folds and the observable latter one (when $H \approx 6.5 \times 10^{-6}m_{\text{Pl}}$) lasts for about 60 e-folds. It is important to emphasize that by this time the Higgs field amplitude is less than $10^{-8}m_{\text{Pl}}$ so it later evolves to the electroweak vacuum.




FIG. 4. The time evolution of the Hubble parameter.

The Electroweak Vacuum

At late times, the Higgs is closed enough to the origin of the potential and, through the symmetry breaking mechanism, can be attracted to the local minima. It receives a non-zero vacuum expectation value and spontaneously breaks the electroweak symmetry. The expectation value and the curvature about the minimum are

$$\varphi_0^2 \approx -m^2/\lambda, \quad (17)$$

$$m_\varphi^2 \approx 2\varphi_0^2(\lambda + \xi^2/2\alpha). \quad (18)$$

Experimental data determines the free parameters as

$$-m(m_{\text{EW}})^2/\lambda(m_{\text{EW}}) \approx (246 \text{ GeV})^2, \quad (19)$$

$$\lambda(m_{\text{EW}}) + \xi(m_{\text{EW}})^2/2\alpha \approx 0.13. \quad (20)$$

Furthermore, the Weyl field receives a non-zero vacuum expectation value and mass as follows

$$\chi_0 \approx -\xi m_{\text{Pl}}^{-1} \varphi_0^2 \approx -10^{-5} \xi \text{ eV}, \quad (21)$$

$$m_\chi \approx (6\alpha)^{-1/2} m_{\text{Pl}} \approx 10^{13} \text{ GeV}. \quad (22)$$

Basically, the non-minimal coupling of the Higgs field which itself develops a non-zero vacuum expectation value through other mechanism induces a vacuum expectation value for the Weyl field. It has an important implication; as can be seen from the action (5), the kinetic term of the Higgs field is not canonically normalized. One needs to apply a field redefinition to make it

canonical which could be interpreted as the Higgs wavefunction renormalization

$$h^c \equiv e^{-\bar{\chi}_0/2} \varphi \approx (1 + \xi \varphi_0^2/m_{\text{Pl}}^2) \varphi. \quad (23)$$

Thereby, the Higgs coupling to the Standard Model particles would be modified. This modification has an observable effect at colliders by suppressing or enhancing the decay modes of the Higgs particle. The combined analysis of the ATLAS and CMS excludes $|\xi| \gtrsim 10^{32}$ at 95% C.L. This can be compared to a much tighter constraint one would obtain for the Higgs non-minimally coupled to the Einstein gravity [17].

On the other hand, equation (20) implies that the non-minimal coupling and higher-curvature gravity normalize the Higgs self-coupling parameter. If ξ is small the effect is insignificant. However, when ξ is big as equation (8) demands, then the effect could be substantial and in principle observable at colliders. In this regard, two extreme cases can be studied; either the first or the second term dominates the equation (20). If λ gives most of contribution at the electroweak scale, then $\xi/2\alpha$ is only needed to stabilize the potential at the Planck scale. Thus, ξ should be around the bounds given in equation (8). By computing the running of the non-minimal coupling β_ξ , one understands that ξ at the electroweak scale is about 10% less than ξ at the Planck scale. Altogether, $\xi(m_{\text{EW}})/2\alpha$ makes $\lambda(m_{\text{EW}})$ about 8% less than what would expect in its absence. In the other extreme limit, when $\xi/2\alpha$ contributes the most to equation (20) at the electroweak scale, it gives an upper bound $|\xi_{\text{EW}}| \lesssim 22000$.

Conclusion

In this note it has been argued that non-minimal coupling of the Higgs field to a higher-curvature theory of gravity could alleviate problems associated with metastability or instability of the electroweak vacuum. Essentially, pure gravitational effects, also operative in deriving inflation, could be applied to account for the present-day electroweak minimum without a need for fine-tuning in the initial Higgs value in the early Universe.

Acknowledgments I would like to thank ICTP, Trieste for hospitality during the completion of this work.

* Electronic address: mahdi@ipm.ir

-
- [1] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716** (2012) 1 [arXiv:1207.7214 [hep-ex]].
 [2] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716** (2012) 30 [arXiv:1207.7235 [hep-ex]].
 [3] [ATLAS and CDF and CMS and D0 Collaborations], arXiv:1403.4427 [hep-ex].
 [4] G. Degrassi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori and A. Strumia, JHEP **1208**

- (2012) 098 [arXiv:1205.6497 [hep-ph]].
 [5] M. Sher, Phys. Rept. **179** (1989) 273.
 [6] V. Branchina and E. Messina, Phys. Rev. Lett. **111** (2013) 241801 [arXiv:1307.5193 [hep-ph]].
 [7] O. Lebedev and A. Westphal, Phys. Lett. B **719** (2013) 415 [arXiv:1210.6987 [hep-ph]].
 [8] I. Bars, P. J. Steinhardt and N. Turok, Phys. Lett. B **726** (2013) 50 [arXiv:1307.8106].

- [9] P. A. R. Ade *et al.* [Planck Collaboration], *Astron. Astrophys.* (2014) [arXiv:1303.5076 [astro-ph.CO]].
- [10] P. A. R. Ade *et al.* [BICEP2 Collaboration], *Phys. Rev. Lett.* **112**, 241101 (2014) [arXiv:1403.3985 [astro-ph.CO]].
- [11] C. G. Callan, Jr., S. R. Coleman and R. Jackiw, *Annals Phys.* **59**, 42 (1970).
- [12] K. S. Stelle, *Phys. Rev. D* **16**, 953 (1977).
- [13] A. A. Starobinsky, *Phys. Lett. B* **91** (1980) 99.
- [14] I. Ben-Dayan, S. Jing, M. Torabian, A. Westphal and L. Zarate, *JCAP* **09** (2014)005 arXiv:1404.7349 [hep-th].
- [15] C. D. Hoyle, D. J. Kapner, B. R. Heckel, E. G. Adelberger, J. H. Gundlach, U. Schmidt and H. E. Swanson, *Phys. Rev. D* **70**, 042004 (2004) [hep-ph/0405262].
- [16] X. Calmet, S. D. H. Hsu and D. Reeb, *Phys. Rev. D* **77**, 125015 (2008) [arXiv:0803.1836 [hep-th]].
- [17] M. Atkins and X. Calmet, *Phys. Rev. Lett.* **110** (2013) 051301 [arXiv:1211.0281 [hep-ph]].