

# Dynamical generation of the weak and Dark Matter scales from strong interactions

Oleg Antipin<sup>a</sup>, Michele Redi<sup>a</sup>, Alessandro Strumia<sup>b</sup>

<sup>a</sup>*INFN, Sezione di Firenze, Via G. Sansone, 1; I-50019 Sesto Fiorentino, Italy*

<sup>b</sup>*Dipartimento di Fisica dell'Università di Pisa and INFN, Italy  
and National Institute of Chemical Physics and Biophysics, Tallinn, Estonia*

## Abstract

Assuming that mass scales arise in nature only via dimensional transmutation, we extend the dimension-less Standard Model by adding vector-like fermions charged under a new strong gauge interaction. Their non-perturbative dynamics generates a mass scale that is transmitted to the elementary Higgs boson by electro-weak gauge interactions. In its minimal version the model has the same number of parameters as the Standard Model, predicts that the electro-weak symmetry gets broken, predicts new-physics in the multi-TeV region and is compatible with all existing bounds, provides two Dark Matter candidates stable thanks to accidental symmetries: a composite scalar in the adjoint of  $SU(2)_L$  and a composite singlet fermion; their thermal relic abundance is predicted to be comparable to the measured cosmological DM abundance. Some models of this type allow for extra Yukawa couplings; DM candidates remain even if explicit masses are added.

## Contents

1	Introduction	2
2	Higgs Mass	4
3	Dark Matter	11
4	Conclusions	17
A	Effective potential	19

# 1 Introduction

The idea that the weak scale could be dynamically generated from strong interactions has a long history. Originally, technicolor models were developed as an alternative to the Higgs: the weak interactions of the techniquarks  $Q$  were chosen so that their condensates would break the SM electro-weak group and the weak scale was the technicolor scale. This scenario was disfavoured by flavour and precision data even before the first LHC run, where the Higgs and no new physics was observed.

Later, strong dynamics was invoked to generate a composite or partially-composite Higgs, although realising complete models is so complicated that model-building is usually substituted by postulating effective Lagrangians with the needed properties.

Recently, models where new strong dynamics does not break the electro-weak symmetry nor provide a composite Higgs have been considered in the literature, just because they are simple, phenomenologically viable and lead to interesting LHC phenomenology [1]. With abuse of language we use the old name ‘technicolor’. In this paper we show that these models

1. provide Dark Matter candidates;
2. provide a dynamical origin for the electro-weak scale, if we adopt the scenario of ‘finite naturalness’ [2, 3, 4].

Point 2 amounts to assuming that quadratically divergent corrections to the Higgs mass have no physical meaning and can be ignored, possibly because the fundamental theory does not contain any mass term [4]. In this context, dynamical generation of the weak scale via dimensional transmutation has been realised with weakly-coupled dynamics, in models where an extra scalar  $S$  has interactions that drive its quartic  $\lambda_S|S|^4$  negative around or above the weak scale:  $S$  acquires a vev at this scale, and its interaction  $\lambda_{HS}|H|^2|S|^2$  effectively becomes a Higgs mass term,  $m^2 = \lambda_{HS}\langle S \rangle^2$  [5]. A related possibility is that the scalar  $S$  is interacting with techniquarks [6] or charged under a technicolor gauge group [7] and again  $S$  acquires a vev or forms a condensate. In all these models  $\langle S \rangle$  can be pushed arbitrarily above the weak scale by making  $\lambda_{HS}$  arbitrarily small, leaving no observable signals.

We here consider simple models without any extra scalar  $S$  beside the Higgs doublet  $H$ . The SM is extended by adding a gauge group  $G_{TC}$  (for example  $SU(N)$ ) and techniquarks  $Q_L$  charged under the SM, as well as the corresponding  $Q_R$  in the conjugated representations of the gauge group  $G_{SM} \otimes G_{TC}$ , so that  $Q_L \oplus Q_R$  is vectorial. As a consequence the condensate  $\langle Q_L Q_R \rangle$  transforms as a singlet of  $G_{SM}$  and does not break it.

The techniquarks have no mass terms because of our assumption that only dimensionless couplings exist<sup>1</sup>; for certain assignments of their gauge quantum numbers, techniquarks

---

<sup>1</sup>Relaxing this hypothesis allows other interesting possibilities for Dark Matter that will be discussed in a separate publication [8].

can have Yukawa interactions  $y$  with the elementary SM Higgs doublet  $H$ . The scenario that we consider is described by the renormalizable Lagrangian

$$\mathcal{L} = \mathcal{L}_{\text{SM}}^{m=0} - \frac{1}{4} \mathcal{G}_{\mu\nu}^A + \bar{Q}_L^i i \not{D} Q_L^i + \bar{Q}_R^j i \not{D} Q_R^j + (y_{ij} H Q_L^i Q_R^j + \text{h.c.}) \quad (1)$$

where  $\mathcal{L}_{\text{SM}}^{m=0}$  describes the SM without the Higgs mass term, and  $\mathcal{G}_{\mu\nu}^A$  is the techni-color field strength. In models where Yukawa couplings  $y$  are not allowed (for example techni-quarks in the 3 of  $\text{SU}(2)_L$ ) the number of free parameters is the same as in the Standard Model: all new physics is univocally predicted. This new physics manifests as:

- Strong dynamics generates a dynamical scale  $\Lambda_{\text{TC}}$  that can be identified with the mass of the lightest vector meson resonance, the techni- $\rho$ , and spontaneously breaks accidental chiral symmetries conserved by the techni-strong interactions producing light pseudo-Goldstone bosons (GB). Using large  $N$  counting  $m_\rho = g_\rho f$  where  $f$  is the decay constant of the techni-pions and  $g_\rho \approx 4\pi/\sqrt{N}$ .
- In absence of techni-quark masses, the techni-pions  $\pi \sim Q_L Q_R$  acquire mass  $m_\pi^2 \approx \alpha_2 m_\rho^2 / 4\pi$  from the electro-weak gauge interactions that explicitly break the global techni-flavour accidental symmetries. Yukawa couplings also contribute to their masses; in absence of Yukawa couplings the lightest techni-pions could be a stable  $\text{SU}(2)_L$  triplet providing a viable DM candidate.
- The heaviest new particles are techni-baryons with mass  $m_B \approx N m_\rho$ . The lightest techni-baryon is stable and is a natural DM candidate; if it is a thermal relic, the observed DM abundance is reproduced for  $m_B \approx 100$  TeV [9].

The LHC phenomenology of techni-strong dynamics was discussed in [1]. The main new point of our work is the possible connection with the weak scale and implications for dark matter. Assuming that power divergences vanish [2, 4], the techni-strong interactions give a finite *negative* contribution to the Higgs squared mass term, such that the weak scale is dynamically generated. The Higgs physical mass arises as

$$M_h^2 \approx +\alpha_2^2 f^2 + y^2 \frac{m_\rho^2 f^2}{m_\pi^2} \quad (2)$$

so that *the techni-color scale is predicted to be*  $f \approx M_h/\alpha_2 \approx \text{few TeV}$ , or smaller in models where  $y$  is present and dominant in eq. (2). Unlike ordinary techni-color as a solution to the usual hierarchy problem, where the natural scale for new physics is the weak scale itself, in this scenario the natural mass scales are

$$m_\pi \sim 2 \text{ TeV}, \quad m_\rho \sim 20 \text{ TeV}, \quad m_B \sim 50 \text{ TeV}. \quad (3)$$

New physics effects in accelerator searches and precision experiments are well below the present sensitivity. In particular no new effects are generated in flavor physics. Techni-pions [11] and techni-baryons [12], stable due to accidental symmetries of the renormalizable Lagrangian, can provide a thermal Dark Matter candidate.

This work is organised as follows. In section 2 we consider the Higgs mass generated by the SM electro-weak gauge couplings, by the SM strong coupling, and by the Yukawa couplings of the Higgs with the techni-quarks, allowed in some models. Dark Matter is discussed in section 3. We conclude in section 4. In the appendix we present the technical details of the computation of the potential induced by Yukawa interactions.

## 2 Higgs Mass

We write the tree-level potential of the SM Higgs doublet  $H$  as

$$V = m^2|H|^2 + \lambda|H|^4. \quad (4)$$

If  $m^2 \equiv -M_h^2/2$  is negative, the Higgs doublet  $H$  develops the vacuum expectation value  $v = M_h/\sqrt{2\lambda} \approx 246.2$  GeV: expanding the potential  $V$  around its minimum as  $H = (0, (v+h)/\sqrt{2})$  shows that  $M_h \approx 125$  GeV is the tree-level mass of the physical Higgs boson  $h$ .

Under our assumptions, the only mass scale of the theory is set by the dynamical scale of the techni-color sector. Through loop corrections it induces other scales and in particular the Higgs mass parameter. Electro-weak interactions of the techni-quarks induce a 2-loop contribution, computed in section 2.1, and color charges give a 3 loop contribution to the Higgs mass, computed in section 2.2. If the Higgs couples to the techni-quarks through Yukawa interactions (for example if techni-quarks contain doublets and singlets under the electro-weak interactions) a contribution to the Higgs mass is also generated at 1-loop, computed in section 2.3.

### 2.1 Electro-weak interactions

Electro-weak gauge interactions give a minimal, quasi-model-independent, contribution to the Higgs mass, described by the non-perturbative techni-color multi-loop dressing of the two-loop Feynman diagram in fig. 1a (plus the associated seagull diagram): the Higgs interacts with the electro-weak vectors, that interact with the techni-quarks.

To leading order in the SM interaction, and to all orders in the techni-strong interactions, the techni-strong dynamics corrects the SM electro-weak gauge bosons propagator as

$$D_{\mu\nu}^{YY}(q) = -i\frac{\eta_{\mu\nu}}{q^2}(1 + g_Y^2\Pi_{YY}(q^2)) + i\xi_Y\frac{q_\mu q_\nu}{q^2} \quad (5)$$

$$D_{\mu\nu}^{ab}(q) = -i\frac{\eta_{\mu\nu}}{q^2}(1 + g_2^2\Pi_{WW}(q^2))\delta^{ab} + i\xi_W\frac{q_\mu q_\nu}{q^2}\delta^{ab} \quad (6)$$

where  $\xi_V$  are gauge-fixing parameters. Techni-strong dynamics is encoded in the  $\Pi_{VV}(q^2)$  functions. From the point of view of the techni-strong dynamics, they are the renormalised two-point functions of the currents  $J_\mu^a = \sum_i \bar{Q}^i \gamma_\mu T_{Q_i}^a Q^i$  (where  $Q^i = (Q_L^i, \bar{Q}_R^i)$ ) is a Dirac




Figure 1: The two loop contribution to the Higgs mass coming from the electro-weak gauge interactions of: a) a techni-quark, to be dressed with non-perturbative techni-interactions, approximated as: b) the techni-gluon condensate; c) the techni- $\rho$ . The extra seagull diagram is not explicitly plotted.

spinor and  $T^a$  are the SM gauge generators) corresponding to the unbroken part of the accidental global techni-flavour symmetry, partially gauged by electro-weak interactions:

$$i \int d^4x e^{iq \cdot x} \langle 0 | T J_\mu^V(x) J_\nu^{V'}(0) | 0 \rangle \equiv \delta^{VV'} (q^2 g_{\mu\nu} - q_\mu q_\nu) \Pi_{VV}(q^2). \quad (7)$$

The correction to the Higgs mass is

$$\Delta m^2 = -\frac{3}{4i} \int \frac{d^4q}{(2\pi)^4} \frac{3g_2^4 \Pi_{WW}(q^2) + g_Y^4 \Pi_{YY}(q^2)}{q^2}, \quad (8)$$

and, performing the Wick rotation to the Euclidian  $Q^2 = -q^2 > 0$ ,

$$\Delta m^2 = \frac{3}{4(4\pi)^2} \int dQ^2 \left[ 3g_2^4 \Pi_{WW}(-Q^2) + g_Y^4 \Pi_{YY}(-Q^2) \right]. \quad (9)$$

In general the integral above is UV-divergent, quadratically and logarithmically. In the case at hand, the unphysical power divergences are ignored because of our assumption of finite naturalness, and logarithmic divergences (that describe the RGE running of  $m^2$ ) are absent, because of our assumption that the only mass scale,  $\Lambda_{TC}$ , is generated dynamically. Thereby the generated squared Higgs mass term is finite and scheme independent.

We next show that the electro-weak interactions induce a calculable negative Higgs mass so that the electro-weak symmetry is spontaneously broken. We proceed in 3 steps: dispersion relations in section 2.1.1 show in general that  $\Delta m^2 < 0$ , Operator Product Expansion in section 2.1.2 shows that  $\Delta m^2$  is ultra-violet finite, vector meson dominance and/or large  $N$  in section 2.1.3 allow to give the estimate  $\Delta m^2 \approx -\alpha_2^2 f^2$ .

### 2.1.1 Dispersion relation

Under our assumptions, quadratically divergent terms are zero and we are interested in the dependence on the physical scales of the theory. To extract this we consider the variation of the Higgs mass with respect to the dynamical scale of the theory  $\Lambda_{\text{TC}}$ ,

$$\frac{\partial \Delta m^2}{\partial \Lambda_{\text{TC}}^2} = \frac{3g_2^4}{4(4\pi)^2} \int dQ^2 \left[ 3g_2^4 \frac{\partial \Pi_{WW}}{\partial \Lambda_{\text{TC}}^2} + g_Y^4 \frac{\partial \Pi_{YY}}{\partial \Lambda_{\text{TC}}^2} \right]. \quad (10)$$

The sign of the gauge correction  $\Delta m^2$  can be determined using the dispersion relation [16]

$$\frac{\partial \Pi_{VV}(q^2)}{\partial q^2} = \frac{1}{\pi} \int_0^\infty ds \frac{\text{Im} \Pi_{VV}(s)}{(s - q^2 - i\epsilon)^2}. \quad (11)$$

where we use the conventions of [17]. The optical theorem relates the cross-sections  $\sigma(s)$  to  $\text{Im} \Pi_{VV}(s)$ , allowing to show in general that  $\text{Im} \Pi_{VV} \leq 0$ .<sup>2</sup> For dimensional reasons, the dimension-less  $\Pi_{VV}$  can only depend on  $Q^2/\Lambda_{\text{TC}}^2$ . Thereby

$$\frac{\partial \Pi_{VV}}{\partial \Lambda_{\text{TC}}^2} = -\frac{Q^2}{\Lambda_{\text{TC}}^2} \frac{\partial \Pi_{VV}}{\partial Q^2} = \frac{Q^2}{\Lambda_{\text{TC}}^2} \frac{1}{\pi} \int \frac{\text{Im} \Pi_{VV}(s)}{(s + Q^2)^2} ds < 0 \quad (13)$$

where in the last step we used the dispersion relation. A similar relation holds for the hypercharge contribution. The integrand in (10) is negative definite corresponding to a negative  $\Delta m^2$  given the boundary condition  $\Delta m^2 = 0$  for  $\Lambda_{\text{TC}} = 0$ .

### 2.1.2 The ultra-violet tail

In a theory with a dynamical scale  $\Lambda_{\text{TC}}$ , arguments based on Operator Product Expansion allow to show that  $\partial \Delta m^2 / \partial \Lambda_{\text{TC}}^2$  is ultra-violet convergent as expected and to compute the high-energy tail of  $\Pi_{VV}(q^2)$ .  $\Pi_{VV}$  can be expanded as

$$\Pi_{VV}(q^2) \stackrel{q^2 \gg \Lambda_{\text{TC}}^2}{\simeq} c_1(q^2) + c_2(q^2) \langle 0 | m_Q \mathcal{Q}_L \mathcal{Q}_R | 0 \rangle + c_3(q^2) \langle 0 | \frac{\alpha_{\text{TC}}}{4\pi} \mathcal{G}_{\mu\nu}^2 | 0 \rangle + \dots \quad (14)$$

The first term (unity operator) does not contribute to (10). Indeed, at leading order it describes the diagram in fig. 1a with techni-quarks but neglecting their techni-color interactions, such that

$$c_1 = \frac{C}{12\pi^2} \ln(-q^2) + \dots \quad (15)$$

<sup>2</sup>As a check, replacing techni-color with a perturbative one-loop correction of fermions with explicit mass  $m_Q$ , one would obtain

$$\frac{\partial \Pi_{VV}(-Q^2)}{\partial m_Q^2} = -\frac{g^2}{2\pi^2} \frac{Q^2}{m_Q^2} \int_0^1 \frac{x^2(1-x)^2}{m_Q^2 + x(1-x)Q^2}. \quad (12)$$

Inserting this into eq. (10) the integrand is negative definite but the integral is logarithmically divergent. This corresponds to a contribution proportional to  $g^2 m_Q^2$  in the RG equation for the Higgs mass  $m^2$ . No such UV-divergent RGE effect is present in a techni-color theory that generates dynamically a mass scale  $\Lambda_{\text{TC}}$  from a dimension-less coupling  $g_{\text{TC}}$ , given that, in any mass-independent scheme such as Minimal Subtraction, only  $g_{\text{TC}}$  can appear in the RGE.

where  $C > 0$  is a model-dependent group theory factor given by  $C = \text{Tr} T^a T^a$  in terms of the  $SU(2)_L$  techni-quark generators (with a similar expression factor for the  $U(1)_Y$  generators). This high energy tail does not contain any mass scale, so that the associated quadratically divergent no-scale integral in eq. (9) vanishes, under our assumptions. The second term also vanishes, because it is proportional to the techni-quark masses  $m_Q$  that vanish under our assumption that the theory does not contain any mass scale.

The third term in eq. (14) is represented by the Feynman diagram in fig. 1b, which gives  $c_3 = -C'/q^4$  [16], where  $C' > 0$  is another order one model-dependent group theory factor. The techni-gluons form a positive condensate (the condensate is positive-defined in the Euclidian path-integral [16], in agreement with QCD lattice computations)

$$\langle 0 | \frac{\alpha_{\text{TC}}}{4\pi} \mathcal{G}_{\mu\nu}^A{}^2 | 0 \rangle = \kappa \Lambda_{\text{TC}}^4. \quad (16)$$

where  $\kappa > 0$  is an order-one coefficient. This allows to show that the UV contribution to the squared Higgs mass term is negative as expected:

$$\Delta m^2|_{\text{UV}} \simeq -\frac{3C'g_2^4}{4(4\pi)^2} \kappa \Lambda_{\text{TC}}^4 \int_{Q_{\text{min}}^2}^{\infty} \frac{dQ^2}{Q^4} \approx -\alpha_2^2 \frac{\kappa \Lambda_{\text{TC}}^4}{Q_{\text{min}}^2}. \quad (17)$$

The  $1/Q_{\text{min}}^2$  dependence on the artificial infra-red cut-off  $Q_{\text{min}} \sim \Lambda_{\text{TC}}$  shows that the dominant effects comes from virtual momenta  $Q^2$  around the techni-meson masses.

### 2.1.3 The infra-red and resonance region

The dominant contribution to the Higgs mass comes from the  $Q^2$  region densely populated by the techni-meson resonances. A variety of methods have been proposed to approximatively describe such region: vector meson dominance, Weinberg sum rules, large  $N$ , holographic models... As long as the techni-quarks are charged under the electro-weak group, they form, among the various mesons, spin-1 resonances that mix with the SM electro-weak vectors  $V_\mu$ . This is described by the effective Lagrangian

$$\mathcal{L}_{\text{eff}} = -\frac{1}{4g_0^2} V_{\mu\nu}^a V^{a\mu\nu} - \frac{1}{4g_\rho^2} \rho_{\mu\nu}^a \rho^{a\mu\nu} + \frac{f^2}{2} (V_\mu^a - \rho_\mu^a)^2 \quad (18)$$

such that the massless eigenstate has gauge coupling  $1/g_2^2 = 1/g_0^2 + 1/g_\rho^2$  and the orthogonal heavy state has mass  $m_\rho^2 = f^2(g_0^2 + g_\rho^2)$ . Integrating out the  $\rho$  at tree-level one finds:

$$\Pi_{VV}(q^2) = \frac{m_\rho^2}{g_\rho^2(q^2 - m_\rho^2 + i\epsilon)}. \quad (19)$$

Plugging eq. (19) into eq. (9) we obtain a logarithmically divergent infra-red correction to the squared Higgs mass term:

$$\Delta m^2 \approx -\frac{9g_2^4}{4(4\pi)^2} \int dQ^2 \frac{m_\rho^2}{g_\rho^2(Q^2 + m_\rho^2)} \sim -\frac{g_2^4 m_\rho^2}{(4\pi)^2 g_\rho^2} \log \frac{\Lambda^2}{m_\rho^2} \sim -\alpha_2^2 f^2. \quad (20)$$




Figure 2: *The three loop contribution to the Higgs mass coming from techni-quarks  $Q$  that only have color interactions. Similar diagrams can be drawn for graviton contributions.*

The integrand is negative definite and its size agrees with the naive expectation based on the Feynman diagram plotted in fig. 1c, including the  $1/g_\rho^2$  suppression of vector mixing. The logarithmic UV divergence here arises because this is only an approximate description, where an explicit mass term  $m_\rho$  substitutes the dynamical mechanism of mass generation. An infinite number of states would be needed to properly describe the non-perturbative dynamics.

In theories with large  $N$  this can be made more rigorous:  $\Pi_{VV}$  can be represented exactly as an infinite sum of poles corresponding to the physical quasi-stable techni-mesons of the theory:

$$\Pi_{VV}(q^2) = \frac{N}{16\pi^2} m_\rho^2 \sum_i \frac{c_i^2}{q^2 - m_i^2 + i\epsilon}. \quad (21)$$

where  $c_i$  are adimensional coefficients. The infinite number of resonances allows to reproduce the logarithmic divergence, that does not contribute to the Higgs mass zero under our assumption of finite naturalness.

These considerations offer an intuitive argument to understand the sign of  $\Delta m^2$ . The net effect of non-perturbative dynamics is creating a mass gap that stops the techni-quark contribution to the RGE running of  $g_2, g_Y$  below  $\Lambda_{TC}$ , effectively making  $g_2, g_Y$  smaller with respect to the perturbative case. As a consequence the unphysical power divergence present in the SM,  $\Delta m^2 \sim +g_{2,Y}^2 \Lambda^2$ , gets replaced by a finite physical effect  $\Delta m^2 \sim -g_{2,Y}^4 \Lambda_{TC}^2$ .

## 2.2 Color interactions

We next consider techni-color models where the techni-quarks have SM color interactions. For example, techni-quarks could be a color octet of  $SU(3)_c$ , charged also under the techni-color gauge group. Then techni-quarks cannot have any Yukawa coupling to the SM Higgs: both the Yukawa contribution of section 2.3 and the electro-weak contribution of section 2.1



are absent.

In these models, the Higgs mass is dominantly generated at three loops: the Higgs interacts with the top quark, that interacts with the gluons, that interact with the techni-quarks, as plotted in fig. 2. The computation can be performed along the lines of section 2.1 by defining  $\Pi_{GG}(q^2)$ , the techni-color correction to the gluon propagator. Summing the two diagrams of fig. 2, the result is ultraviolet-convergent:

$$\Delta m^2 = -\frac{64y_t^2 g_3^4}{(4\pi)^4} \int dQ^2 \Pi_{GG}(-Q^2) \sim \frac{y_t^2 g_3^4}{4\pi^4} f^2. \quad (22)$$

The computation of the sign is analogous to what described in the previous section (with  $\Pi_{WW}$  replaced by  $\Pi_{GG}$ ): in the present case we find a positive  $\Delta m^2$ , such that this contribution does not induce electro-weak symmetry breaking. The sign of the effect also corresponds to the intuitive reasoning presented at the end of the previous section: the sign is opposite to the known negative sign of the naive quadratic divergence associated with  $y_t$ , because  $g_3$  and thereby  $y_t$  are reduced by techni-strong dynamics.

We mention a final possibility. The techni-quarks could be completely neutral under the whole SM gauge group. In this situation only gravity mediates a contribution to the Higgs mass, proportional to the two-point function of the energy momentum tensor. Furthermore, a super-Planckian techni-color condensate would dynamically generate the Planck mass itself, within a dimensionless extension of Einstein gravity such as agravity [4]. The problem is that techni-color dynamics, dominated by a single non-perturbative coupling, has no free parameters and would also generate a large negative cosmological constant, which is at odd with observations.

### 2.3 Yukawa interactions

Finally, we consider the case where the gauge quantum numbers of the techni-quarks allow for Yukawa couplings to the elementary Higgs. This choice implies the existence of a technipion  $\pi_2$  with the same quantum numbers of the Higgs doublet  $H$ , that can then mix with  $H$ .

The left panel of fig. 3 shows the one-loop corrections to the squared Higgs mass generated by a weakly coupled techni-quark with Yukawa interactions to the Higgs. At strong coupling the physical degrees of freedom become bound state techni-hadrons that can be described using effective Lagrangian techniques. The techni-quark loop can be matched to an effective chiral Lagrangian, so that such diagrams collapses to a tree level diagram (right-handed panel of fig. 3) dominated by the lightest techni-mesons, the techni-pions  $\pi \approx Q_L Q_R$ . For simplicity we here consider Yukawa couplings that preserve the  $Q_L \leftrightarrow Q_R$  parity of the techni-strong interactions; a more general discussion can be found in the appendix. Similarly to quark masses in QCD, the Yukawa interactions produce the following term in the chiral Lagrangian,

$$y m_\rho f^2 \text{Tr}[HU] + \text{h.c.} \quad (23)$$



Figure 3: Correction to the Higgs mass coming from the Yukawa coupling with: a) a weakly coupled massive fermion; b) a massless strongly interacting fermion.

where  $U = \exp(i\pi^{\hat{a}}T^{\hat{a}}/f)$  is the Goldstone boson matrix. As we discuss in detail in the appendix, upon minimisation of the potential this term induces a mass mixing  $\approx ym_{\rho}fH\pi^*$  between the techni-pion and the elementary Higgs. This term also explicitly breaks accidental symmetries respected by gauge interactions.

What emerges is a two-Higgs doublet system where the extra Higgs doublet  $\pi_2$  is a heavy composite doublet with negligible vev. In order to compute the mass eigenstates, we need to compute the mass matrix. Including effects at tree and one-loop level in the SM couplings  $g_2$  and  $y$ , the mass matrix has the structure

$$\begin{array}{c} \pi_2 \\ H \end{array} \begin{pmatrix} \mathcal{O}(g_2^2) \pm \mathcal{O}(y^2)/(4\pi)^2 & \mathcal{O}(y)\sqrt{N}/(4\pi) \\ \mathcal{O}(y)\sqrt{N}/(4\pi) & -\mathcal{O}(y^2)N/(4\pi)^2 \end{pmatrix} m_{\rho}^2 \quad (24)$$

where we used the fact that the one-loop contribution of weak gauge interactions to  $m_{\pi}^2 \approx g_2^2 m_{\rho}^2/(4\pi)^2$  is positive (as known from the SM analogous computation of the  $\pi^+/\pi^0$  mass difference [13]), and added the one-loop Yukawa contribution (absent in the SM<sup>3</sup>). The  $HH^*$  entry describes the contribution of composite scalar resonances that can also mix with the Higgs giving a negative sub-leading contribution to its mass squared, see appendix for more details.

We see that the phenomenologically acceptable regime is the one where the Yukawa coupling is small,  $y \ll g$ , such that: 1) the loop contribution coming from the Yukawa coupling can be ignored; 2) the heaviest eigenstate is the techni-pion with squared mass  $m_{\pi}^2 > 0$ ; 3)

<sup>3</sup>The literature on composite Higgs models explored linear couplings of SM quarks to composite fermionic states, finding that they can give a negative contribution to the Higgs mass term. Simple UV completions require extra scalars as in the supersymmetric realisation of [14]. Here instead we compute the techni-pion potential induced by a bi-linear  $HQ_L Q_R$  Yukawa coupling, involving techni-quarks  $Q$  and a scalar  $H$  without techni-strong interactions.

the determinant of the mass matrix is dominated by the off-diagonal terms and is negative: the lightest eigenstate is the elementary Higgs, that acquires a *negative* squared mass term dominated by the mass mixing term in eq. (24) and given by a see-saw-like formula:

$$\Delta m^2 \sim -\frac{y^2}{(4\pi)^2} \frac{m_\rho^4 N}{m_\pi^2} \sim -y^2 \frac{m_\rho^2 f^2}{m_\pi^2}. \quad (25)$$

### 3 Dark Matter

The models described in this paper contain two Dark Matter (DM) candidates: techni-baryons and techni-pions. Their stability is guaranteed by accidental symmetries of the renormalizable Lagrangian, techni-baryon number and (possibly)  $G$ -parity [11].

In fact the presence of stable states is a generic prediction of the framework that implies restrictions on the representations of the techni-quarks under the SM gauge group, such that the stable states are viable DM candidates. In table 1 we summarise the simplest allowed charge assignments under the electro-weak group and the resulting DM candidates. Introducing techni-quark masses allows several other possibilities [8].

The new matter modifies the running of SM gauge couplings. Adding  $n_2$  weak doublets and  $n_3$  weak triplets in the  $N \oplus \bar{N}$  of  $SU(N)_{\text{TC}}$  the beta-function of  $SU(2)_L$  becomes

$$b_2 = -\frac{19}{6} + \frac{2N}{3}(n_2 + 4n_3) \quad (26)$$

such that the  $SU(2)_L$  gauge coupling does not develop a Landau pole below the Planck scale ( $b_2 \lesssim 5$ ) and possibly remains asymptotically free ( $b_2 < 0$ ) for small enough  $n_2, n_3, N$ . Higher  $SU(2)_L$  lead to Landau poles instead. The trans-Planckian Landau pole for hypercharge can be naturally avoided in models where hypercharge is embedded in  $SU(2)_R$  below a few TeV [10]; a technicolor sector could be used to dynamically break the extended gauge group.

#### 3.1 Techni-pions

If techni-quarks fill  $N_F$  fundamentals and anti-fundamentals of the  $SU(N)_{\text{TC}}$  gauge group with  $N \geq 3$ , the spontaneous symmetry breaking  $SU(N_F)_L \otimes SU(N_F)_R / SU(N_F)$  of the accidental global techni-flavor symmetry produces  $N_F^2 - 1$  Goldstone bosons in the adjoint of the unbroken  $SU(N_F)$ . These scalars acquire mass from effects that explicitly break the global symmetries. Within finite naturalness the only contribution to their masses is due to SM gauge interactions, and possibly to the techni-quark Yukawa couplings.

If Yukawa couplings are forbidden by the fermions quantum numbers, then the model is extremely predictive: it only has one free parameter — the techni-color scale — which is fixed by the Higgs mass under the hypothesis of finite naturalness. All the rest is univocally predicted: techni-pion masses, Dark Matter and its thermal relic abundance.

number of techni-flavors	Yukawa	$N = 3$		$N = 4$		
		TCb	TC $\pi$	TCb	TC $\pi$	
$N_F = 2$		2	3	1	3	under TC-flavor SU(2)
model 1: $\mathcal{Q} = 2_{Y=0}$	0	charged	3	1	3	DM, under SU(2) $_L$
$N_F = 3$		8	8	$\bar{6}$	8	under TC-flavor SU(3)
model 1: $\mathcal{Q} = 1_Y + 2_{Y'}$	1	1	no	1	no	DM, under SU(2) $_L$
model 2: $\mathcal{Q} = 3_{Y=0}$	0	3	3	1	3	DM, under SU(2) $_L$
$N_F = 4$		$\bar{20}$	15	$20'$	15	under TC-flavor SU(4)
model 1: $\mathcal{Q} = 4_{Y=0}$	0	charged	3	1	3	DM, under SU(2) $_L$
$N_F = 5$		$\bar{40}$	24	$\bar{50}$	24	under TC-flavor SU(5)
model 1: $\mathcal{Q} = 2_Y + 3_{Y'}$	1	1	no	charged	no	DM, under SU(2) $_L$
model 2: $\mathcal{Q} = 5_{Y=0}$	0	3	3	1	3	DM, under SU(2) $_L$

Table 1: *Dimension-less techni-color models that give viable techni-baryon (TCb) and/or technipion (TC $\pi$ ) Dark Matter candidates with  $Q = Y = 0$ . We consider models with SU( $N$ ) gauge group for  $N = \{3, 4\}$  and  $N_F = \{2, 3, 4, 5\}$  flavours of techni-quarks in its fundamental plus anti-fundamental. The darker rows give the techni-flavour content of the lightest TCb and TC $\pi$  considering only masses induced by techni-color interactions. The lighter rows consider models with viable assignments of electro-weak interactions and show, after including the mass splitting due to unbroken electro-weak interactions, the SU(2) $_L$  content of the lighter DM candidates.*

The SM gauge interactions give positive squared masses to the gauge-charged technipions, while SM singlets remain exact massless Goldstone bosons. If the  $N_F$  techni-quark flavors are composed of  $k$  irreducible (real or pseudo-real) representations of  $G_{\text{SM}}$ , then the technipions decompose under  $G_{\text{SM}}$  as

$$\text{Adj}_{\text{SU}(N_F)} = \left[ \sum_{i=1}^k r_i \right] \otimes \left[ \sum_{i=1}^k \bar{r}_i \right] \ominus 1 \quad (27)$$

so that  $k - 1$  technipions are neutral gauge singlets (the extra scalar singlet analog of the  $\eta'$  in QCD acquires mass from anomalies with techni-interactions and will not play a role in what follows).

One combination of singlets corresponds to a global symmetry anomalous under SU(2) $_L$ , so that the corresponding Goldstone boson acquires an axion-like couplings to SM vectors: an almost massless axion with a decay constant  $f \sim \text{TeV}$  would be grossly excluded by star cooling and other bounds. In absence of techni-quark Yukawa interactions, these bounds significantly reduce the space of models favouring the simplest models with  $k = 1$ . The techni-quarks should belong to a single irreducible representation  $j = (N_F - 1)/2$  of SU(2) $_L$  and, in order to obtain a neutral lightest techni-baryon, the techni-quark hypercharge should vanish. Then the  $N_F^2 - 1$  technipions lie in the following irreducible representations  $J$  of

$SU(2)_L$ :

$$\text{Adj}_{SU(N_F)} = \sum_{J=1}^{N_F-1} J. \quad (28)$$

Models of this kind were studied in [11], where it was pointed out that a discrete symmetry, “ $G$ -parity” exists in these theories (for zero hypercharge) due to the fact that  $SU(2)_L$  representations are real or pseudo-real.  $G$  parity acts on techni-quarks as  $Q \rightarrow \exp(i\pi T^2)Q^c$ , replacing any  $SU(2)_L$  representation with its conjugate representation, which is equivalent to the original representation. SM fields are neutral. On techni-pions  $G$  parity becomes the  $(-1)^J Z_2$  symmetry, so that techni-pions with even (odd) isospin ( $J$ ) are even (odd). Summarizing:

- Techni-pion singlets under  $SU(2)_L$  are  $G$ -even, do not acquire masses from SM gauge interactions and can have anomalous couplings to  $SU(2)_L$  vectors: they are excluded in our framework unless Yukawa couplings make them massive. They are absent if techni-quarks fill a single irreducible representation of  $SU(2)_L$ .
- Techni-pions in the 3 of  $SU(2)_L$  are  $G$ -odd and could be the lightest stable DM candidates. The simplest models are listed in table 1.
- Techni-pions in the 5 of  $SU(2)_L$  are  $G$ -even and are heavier,  $m_{\pi_5} \approx \sqrt{3}m_{\pi_3}$ : they undergo anomalous decays into electro-weak vectors,  $\pi_5 \rightarrow WW$ .
- Techni-pions in higher representations of  $SU(2)_L$ , if present, decay into lighter techni-pions respecting  $G$ -parity by emitting two  $SU(2)_L$  vectors, e.g.  $\pi_7 \rightarrow \pi_3 WW$ .

The situation is different in models where Yukawa couplings  $y$  of techni-quarks to the elementary Higgs are present. The Yukawa couplings break explicitly  $G$ -parity and accidental global symmetries so that the SM singlet techni-pions  $\eta$  receive non-zero masses given by eq. (56),  $M_\eta \sim |y|vm_\rho/m_{\pi_2}$  and star cooling bounds are easily avoided. Furthermore, techni-pions can now decay through the Higgs, so that only techni-baryons remain as dark matter candidates.

Models with techni-color gauge group  $SU(2) \sim Sp(2)$  are special: its fundamental representation is pseudo-real,  $2 \sim 2^*$ , so that the techni-flavour symmetry is enhanced becoming  $SU(2N_F)/Sp(2N_F)$ . The extra techni-pions are  $QQ$  scalars and there are no stable techni-baryons. Dangerous light techni-pions neutral under  $SU(2)_L$  are again absent if techni-quarks lie in a single representation of  $SU(2)_L$  with dimension  $2N_F$ . Within our assumptions however these models do not provide DM candidates because techni-pions are  $G$ -even.

### 3.2 Techni-baryons

Techni-baryons are technicolor singlet states constructed with  $N$  techniquarks. The stability of the lightest techni-baryon follows from the accidental techni-baryon number global symmetry.

Using the non-relativistic quark model, group theory allows to compute the electro-weak quantum numbers of the techni-baryons: their wave-function must be anti-symmetric in the techniquarks. The wave function is assumed to be antisymmetric in technicolor, and so must be symmetric in spin and flavour for the lightest techni-baryons that have no orbital angular momentum. Different techni-baryons are split by spin-spin interactions that prefer, as lightest techni-baryon, the one with smallest spin. As a consequence, the lightest techni-baryons have spin 0 (1/2) for even (odd)  $N \geq 2$ .

In general the  $SU(N_F)$  techniflavour representation of the lightest techni-baryon corresponds to a Young diagram with 2 rows having  $N/2$  boxes each (for  $N$  even) and to a Young diagram with 2 rows having  $(N + 1)/2$  and  $(N - 1)/2$  boxes respectively (for  $N$  odd). In particular, they are

$$\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \end{array} \quad \text{for } N = 3 \quad \text{and} \quad \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \quad \text{for } N = 4. \quad (29)$$

This is the end of the story, as long as technicolor interactions are involved.

Next, the components of a techni-baryon multiplet are split by SM gauge interactions, and possibly by techniquark Yukawa interactions. The lightest components are those with smallest  $G_{\text{SM}}$  charge.

Furthermore, electro-weak symmetry breaking induces extra splitting within the components of any electro-weak multiplet, with the result that the component with smallest electric charge is the lightest stable state [18]. Since DM direct detection constraints demand that DM does not couple at tree level to the  $Z$ , the DM hypercharge should be zero, which is possible for integer isospin.

### 3.3 Direct detection of Dark Matter

The previous discussion is summarised in table 1, which tells that the simplest TC models lead to the following viable stable DM candidates:

- Techni-baryons, fermions for odd  $N$  and scalars for even  $N$ . Their annihilation cross section is estimated to be  $\sigma v \sim g_{\text{TC}}^4/4\pi M^2$ , around the unitarity bound [9]. By performing a naive rescaling of the QCD non-relativistic  $p\bar{p}$  cross section,  $\sigma_{p\bar{p}}v \sim 100/m_p^2$ , we estimate that the cosmological thermal relic abundance of a techni-baryon equals the total DM abundance if its mass is loosely around  $m_B \sim 200$  TeV. A cosmological techni-baryon asymmetry can leave a higher abundance, allowing for a lighter  $m_B$ .




Figure 4: *The signals in Dark Matter direct detection produced by a DM techni-baryon with magnetic or electric dipole moment (line) or from a Minimal-Dark-Matter-like techni-pion with thermal abundance (star), compared to the present experimental LUX bound [20] and to the background due to neutrinos.*

- Scalar techni-pions, that fill a  $SU(2)_L$  triplet with hypercharge  $Y = 0$ . Techni-pions have small residual technicolor interactions (as well as small quartic couplings) and thereby behave as Minimal Dark Matter [18]. Their cosmological thermal relic abundance equals the total DM abundance if their mass is around 2.5 TeV [18]. Their spin-independent cross section for direct detection is  $\sigma_{SI} \approx 0.12 \cdot 10^{-46} \text{ cm}^2$  [19, 2], as plotted in fig. 4.

As already discussed, both mass scales suggested by the DM cosmological abundance arise naturally within the context of finite naturalness.

Techni-baryons have distinctive features in direct detection experiments: if DM is a neutral composite particle made of charged techniquarks, direct detection can be mediated by the photon [21]. Any such DM particle can have a non trivial form factor, dominated at low energy by the ‘charge radius’ interaction. For a scalar DM  $S$  this is the only interaction and can be written as

$$\frac{e}{\Lambda_{TC}^2} (S^* i \partial_\alpha S) \partial_\mu F^{\mu\alpha}. \quad (30)$$

The resulting cross section for direct detection is suppressed by four powers of the TC scale, and is negligible for  $\Lambda_{TC} \sim \text{few TeV}$ .



The situation is more promising if DM is a fermionic techni-baryon  $B$ , which generically has magnetic (and possibly electric) dipole moments,  $\mu$  and  $d$ . They are described by the effective operator

$$\bar{B}\sigma_{\mu\nu}\frac{\mu + id\gamma_5}{2}B F_{\mu\nu}. \quad (31)$$

Electro-magnetic dipoles give sizeable direct detection signals with a characteristic testable enhancement at low recoil-energy  $E_R$ , given that the DM/matter scattering is mediated by the massless photon. Furthermore, in the relevant non-relativistic limit, the cross-section induced by the magnetic dipole  $\mu$  is suppressed by two extra power of the relative DM/matter velocity  $v$  with respect to the cross section induced by the more speculative electric dipole  $d$  [21]

$$\frac{d\sigma}{dE_R} \approx \frac{e^2 Z^2}{4\pi E_R} \left( \mu^2 + \frac{d^2}{v^2} \right). \quad (32)$$

For simplicity, we here assumed a nucleus with  $Z \gg 1$ , a recoil energy  $E_R \ll m_N v^2$  and approximated the nuclear charge form factor with unity.

The magnetic  $g$ -factor, defined by  $\mu = ge/2m_B$ , is expected to be of order one for a strongly-coupled particle (while it is loop suppressed for an elementary particle). We also define the electric  $g$ -factor as  $d = g_E e/2m_B$ . In terms of such  $g$ -factors we find that the present direct detection bound is

$$g^2 + 1.2 \cdot 10^7 g_E^2 < \left( \frac{m_B}{5.1 \text{ TeV}} \right)^3 \quad (33)$$

dominated by LUX data [20, 22]. This bound assumes that techni-baryons constitute all galactic DM, and must be rescaled otherwise. Fig. 4 shows the resulting prediction in the usual plane  $(M_{\text{DM}}, \sigma_{\text{SI}})$  used to describe spin-independent direct detection of Dark Matter.

An electric dipole moment needs CP-violation. In our context, techni-quarks are strictly massless, such that the CP-violating techni-strong  $\theta$  term is not physical. A small  $g_E$  could be generated if techni-quark masses are included.

### 3.4 A worked example

More quantitative predictions can be given in the QCD-like scenario with  $N = N_F = 3$  [12]. In this case the spectrum can be obtained by rescaling known QCD results,

$$\frac{m_B}{m_\rho} \approx 1.3 \quad \frac{m_\pi}{m_\rho} \approx 0.1 \sqrt{J(J+1)} \quad (34)$$

where  $m_\rho$  is the mass of the lightest techni-vector resonance and techni-pions  $\pi$  lie in the  $J$  representation of  $\text{SU}(2)_L$ . The second estimate is obtained from the electro-magnetic splitting of QCD pions, see the appendix.



The lightest techni-baryons are an octet of flavour  $SU(3)$  and table 1 lists the two possible viable assignments for techni-quarks under  $SU(2)_L \otimes U(1)_Y$ :

$$\mathcal{Q} = \begin{cases} 1_{\mp 1/3} \oplus 2_{\pm 1/6} \\ 3_0 \end{cases} . \quad (35)$$

The hypercharges are determined by requiring that the lightest techni-baryon is neutral; in the first case their overall normalisation is determined by requiring that techni-quarks can have a Yukawa interaction with the Higgs. For this choice of quantum numbers the lightest techni-baryon is an electro-weak singlet with  $Y = Q = 0$ , avoiding direct detection constraints.

The lightest technibaryons decompose under  $SU(2)_L$  as

$$\mathbf{8} = \begin{cases} \mathbf{2}(p, n) \oplus \mathbf{3}(\Sigma^{\pm, 0}) \oplus \mathbf{2}(\Xi^0, \Xi^-) \oplus \mathbf{1}(\Lambda_0) & \text{for } \mathcal{Q} = 1 \oplus 2 \\ \mathbf{3} \oplus \mathbf{5} & \text{for } \mathcal{Q} = 3 \end{cases} . \quad (36)$$

In the  $\mathcal{Q} = 1 \oplus 2$  model we used the familiar names of the QCD octet. The lightest techni-baryon is  $\Lambda_0$ , that is analogous to the QCD state  $\Lambda_0^{\text{QCD}} \sim s(ud - du)$ . Its magnetic dipole moment can be estimated from QCD data:  $\mu_{\Lambda_0^{\text{QCD}}} = 0.61e/2m_p$  [23]. Inserting  $g = -0.61$  in eq. (33) we obtain the bound  $m_{\Lambda^0} > 3.7$  TeV. The previous QCD-based estimate of the DM annihilation cross section becomes exact, such that the cosmological DM density is reproduced for  $m_{\Lambda^0} \approx 200$  TeV. In this model there are no stable techni-pions.

In the  $\mathcal{Q} = 3_Y$  model the lightest technibaryon is a triplet  $\mathbf{3}_{3Y}$  of  $SU(2)_L$ , so that neutral DM is obtained for  $Y = 0$  and  $Y = \pm 1/3$ : the first possibility is allowed by direct detection constraints. Due to the absence of Yukawa and hypercharge interactions, the neutral member of the techni-pion triplet is the DM candidate, stable thanks to the accidental G-parity discussed in section 3.1. Its mass must be smaller than 2.5 TeV in order to avoid a thermal relic density bigger than the observed DM density. This implies that in this model the thermal relic density of the technibaryon dark matter is subdominant.

## 4 Conclusions

In conclusion, we presented a new class of models where the Standard Model is made dimension-less by dropping the mass term of the elementary Higgs and extended by adding techni-quarks with techni-color interactions arranged in such a way that they do *not* break the electro-weak gauge group nor generate a composite Higgs. Within the context of finite naturalness — the assumption that a QFT with no mass parameters nor power divergences might provide a revised concept of weak-scale naturalness and of the origin of mass scales — the simplest models of this type dynamically generate a mass term for the Higgs.

The elementary Higgs acquires a squared mass term  $m^2$  entirely determined in terms of weak interactions of the techni-quarks and of the techni-color scale. Using various approximation techniques that allow to control the techni-color dynamics, in section 2.1 we found

that the sign of  $m^2$  is negative, such that  $SU(2)_L \otimes U(1)_Y$  gets broken, and that the observed weak scale is obtained for a technicolor scale  $m_\rho \approx 4\pi M_h/\alpha_2 \approx 10 - 20$  TeV. This is large enough that such models do not pose any phenomenological problem. Techni-pions are lighter, as determined by their electro-weak interactions, and could give observable signals at LHC; in particular techni-pions  $\pi_5$  in the 5 of  $SU(2)_L$  undergo anomalous decays into pairs of electro-weak vectors,  $\pi_5 \rightarrow WW$ . Such models can have the same number of free parameters as the Standard Model: *all new physics is univocally predicted*, up to theoretical uncertainties in the technicolor dynamics, that could be reduced with respect to our estimates by performing dedicated lattice computations.<sup>4</sup>

Independently of the assumption of finite naturalness, the models studied in this paper contain two Dark Matter candidates: the lightest technibaryon  $B$  with mass  $m_B \sim 50$  TeV (section 3.2) and, in some models, the lightest technipion  $\pi_3$ , a triplet under  $SU(2)_L$  with mass  $m_{\pi_3} \sim 0.1m_\rho \sim 1 - 2$  TeV (section 3.1). Their thermal relic abundance is also univocally predicted, with the result that the observed cosmological Dark Matter abundance is naturally reproduced in the technipion case, while the technibaryon seems more likely to be a subdominant Dark Matter component, if a naive rescaling of the QCD  $p\bar{p}$  cross-section holds, and ignoring possible technibaryon asymmetries. The direct detection cross section of such DM candidates is predicted to be 2 – 3 orders of magnitude below present bounds. Magnetic moment interactions of technibaryons would lead to recoil events with a distinctive energy spectrum (section 3.3).

Table 1 offers a panoramic of models that lead to DM candidates. In some models the quantum numbers allow for Yukawa interactions between techniquarks and the elementary Higgs. Such Yukawas give extra negative contributions to the squared Higgs mass term (section 2.3), so that the technicolor scale needed to reproduce the weak scale gets lighter; in such models a singlet technipion is especially light. Models where techniquarks only have QCD interactions or gravitational interactions do not seem to lead to a promising phenomenology, as discussed in section 2.2.

## Acknowledgments

We wish to thank Roberto Franceschini and Giovanni Villadoro for discussions and collaboration at the early stages of this work. We thank Roberto Contino, Riccardo Rattazzi, Slava Rychkov, Riccardo Torre for advice about strong dynamics and the authors of [22] (in particular Eugenio del Nobile) for help about using their code. The work of OA and MR is supported by the MIUR-FIRB grant RBFR12H1MW.

---

<sup>4</sup>Technicolor dynamics generates a negative vacuum energy of order  $-\Lambda_{TC}^4$ . It can be canceled, compatibly with the scenario of dynamical mass generation in the SM sector, by adding another sector negligibly coupled to SM particles; this kind of sector is anyhow needed to account for the Planck mass. This cancellation is the usual huge fine-tuning associated with the cosmological constant problem, on which we have nothing to say.

# A Effective potential

The effective potential for the elementary Higgs and the techni-pions receives contributions at tree level in the Yukawa couplings and at loop level in the gauge and Yukawa couplings. It can be computed using the techniques reviewed in [15]: the relevant ingredients are the correlation functions of the composite operators of the theory. There are three main contributions: from SM gauge interactions at loop level (section A.1); from the possible Yukawa couplings at tree level (section A.2) and at loop level (section A.3). Summing these contributions, the full potential is studied in section A.4.

## A.1 Gauge contribution at one loop level

At 1-loop the SM gauge interactions induce a techni-pion mass that can be computed in terms of correlators of the vector ( $J_\mu^a = \sum_Q \bar{Q} \gamma_\mu T_Q^a Q$ ) and axial ( $J_\mu^a = \sum_Q \bar{Q} \gamma_\mu T_Q^a \gamma_5 Q$ ) symmetry currents. On general grounds these have the form,

$$\begin{aligned} i \int d^4x e^{iq \cdot x} \langle 0 | T J_\mu^V(x) J_\nu^{V'}(0) | 0 \rangle &\equiv \delta^{VV'} (q^2 g_{\mu\nu} - q_\mu q_\nu) \Pi_{VV}(q^2), \\ i \int d^4x e^{iq \cdot x} \langle 0 | T J_\mu^A(x) J_\nu^{A'}(0) | 0 \rangle &\equiv \delta^{AA'} (q^2 g_{\mu\nu} - q_\mu q_\nu) \Pi_{AA}(q^2). \end{aligned} \quad (37)$$

The one-loop techni-pion potential reads [15]:

$$V_{g1} \approx \frac{3}{2(4\pi)^2} \sum_i g_i^2 \text{Tr}[U T^i U^\dagger T^i] \int_0^\infty Q^2 dQ^2 [\Pi_{AA}(-Q^2) - \Pi_{VV}(-Q^2)] \quad (38)$$

where  $U = e^{i\pi^{\hat{a}} T^{\hat{a}}/f}$  is the Goldstone boson matrix,  $g_i$  are the SM couplings and  $T_i$  their generators. Gauge-charged techni-pions acquire positive squared masses, that, for the  $SU(2)_L$  interactions, are estimated as

$$m_\pi^2 \approx \frac{3g_2^2}{(4\pi)^2} J(J+1)m_\rho^2 \quad (39)$$

where  $J$  is the weak isospin of the techni-pion representation.

## A.2 Yukawa contribution at tree level

We now consider the potential generated by the Yukawa interactions. For concreteness we here focus on the case where techni-quarks  $Q = 2 \oplus 1$  fill one doublet and one singlet of  $SU(2)_L$  with hypercharges as in section 3.4. The 8 techni-pions decompose under  $SU(2)_L \otimes U(1)_Y$  as

$$8 = 2_{\pm 1/2} + 3_0 + 1_0. \quad (40)$$

In general there are two Yukawa couplings:

$$y H Q_L^1 Q_R^2 + y' H^\dagger Q_R^1 Q_L^2 + \text{h.c.} = H \bar{Q}_2 \left( \frac{y + y'^*}{2} + \gamma_5 \frac{-y + y'^*}{2} \right) Q_1 + \text{h.c.} \quad (41)$$

where on the right hand side we used Dirac spinors  $Q_i = (Q_L^i, \bar{Q}_R^i)$ . The phases of  $y$  and  $y'$  are not physical and can be chosen for convenience, for example real and positive. The terms above generate the tree level effective potential

$$V_{y0} = a_0 \text{Tr}[MU] + \text{h.c.} \quad (42)$$

where  $a_0 \approx -m_\rho f^2$  and

$$M = \begin{matrix} & \mathcal{Q}_R^1 & \mathcal{Q}_R^2 \\ \mathcal{Q}_L^1 & \begin{pmatrix} 0 & yH \\ y'H^\dagger & 0 \end{pmatrix} \end{matrix}. \quad (43)$$

The explicit result for the potential of the doublet ( $\pi_2$ ) and singlet ( $\eta$ ) techni-pions is

$$V_{y0} = -8\sqrt{2}a_0 \operatorname{Im} e^{-\frac{i\eta}{4\sqrt{3}f}} \frac{\sin \Delta/f}{\Delta} [yH^\dagger \pi_2 + y'\pi_2^\dagger H], \quad \Delta = \frac{1}{4}\sqrt{3\eta^2 + 8\pi_2^\dagger \pi_2}. \quad (44)$$

### A.3 Yukawa contribution at one loop level

To compute the one-loop Yukawa correction to the effective potential we proceed similarly to the gauge interactions. We formally introduce sources  $S_{ij}$  for the techni-quark bilinears  $\mathcal{Q}_L^i \mathcal{Q}_R^j(x)$  (such that, in the real theory of interest, it contains  $yH$  in some of its components) and write the effective Lagrangian that describes the Higgs/techni-pion system after having integrated out the heavier techni-strong dynamics. For simplicity we consider vectorial couplings as in these case fewer invariants exist. In a constant techni-pion configuration and to quadratic order in the sources  $S$ , the effective action has the following structure determined by the symmetries,

$$\mathcal{L}_{\text{eff}}^{QQ} = a_0 \delta^4(q) (\operatorname{Tr}[SU] + \text{h.c.}) + \Pi_0^{QQ}(q^2) \operatorname{Tr}[SS^\dagger] + \Pi_1^{QQ}(q^2) |\operatorname{Tr}[SU]|^2. \quad (45)$$

The first term linear in  $S$  describes the  $\mathcal{Q}_L \mathcal{Q}_R$  condensate. The form factors can be obtained integrating over the strong dynamics including techni-pion fluctuations. By construction they encode the two point functions of the techni-quark bilinears,

$$\langle 0 | \bar{\mathcal{Q}}^i \mathcal{Q}^j(q) \bar{\mathcal{Q}}^k \mathcal{Q}^l(-q) | 0 \rangle = i G_{\text{Adj}}^{QQ}(q^2) \left( \delta^{ik} \delta^{lj} - \frac{1}{3} \delta^{ij} \delta^{lk} \right) + i G_S^{QQ}(q^2) \delta^{ij} \delta^{lk} \quad (46)$$

where  $G_S^{QQ}$  and  $G_{\text{Adj}}^{QQ}$  correspond to the singlet and adjoint channels (namely, the octet for  $N_F = 3$ ). Matching eq.s (45) and (46) (for example choosing  $U = 1$ ) one finds

$$\Pi_0^{QQ} = G_{\text{Adj}}^{QQ}, \quad \Pi_1^{QQ} = G_S^{QQ} - \frac{1}{3} G_{\text{Adj}}^{QQ}. \quad (47)$$

At large  $N$  one has

$$G_{\text{Adj}}^{QQ}(q^2) = \frac{N}{16\pi^2} m_\rho^4 \sum_n \frac{c_{\text{Adj}_n}^2}{q^2 - m_{\text{Adj}_n}^2 + i\epsilon}, \quad G_S^{QQ}(q^2) = \frac{N}{16\pi^2} m_\rho^4 \sum_n \frac{c_{S_n}^2}{q^2 - m_{S_n}^2 + i\epsilon}, \quad (48)$$

where the coefficients  $c$  are of order 1 and the sum is over the scalar resonances in the theory. The sum does not include techni-pions because we only consider vectorial Yukawa couplings that do not generate 1 techni-pion states.

To obtain the effective action for the scalars we just need to set to zero the non dynamical components of  $S$  and add kinetic terms for the components of  $S$  associated to the Higgs. This produces

$$\mathcal{L}_{\text{eff}}^H = a_0 \delta^4(q) (\operatorname{Tr}[MU] + \text{h.c.}) + (q^2 + y^2 \Pi_0^{QQ}(q^2)) H^\dagger H + \Pi_1^{QQ}(q^2) |\operatorname{Tr}[MU]|^2. \quad (49)$$

The first term describes the tree level contribution discussed above. The second term encodes the tree level effect of mixing with heavy scalar resonances that gives the  $HH^*$  entry of the mass matrix in eq. (24),

$$m_H^2 = y^2 \Pi_0^{QQ}(0) \sim -\frac{y^2 N}{(4\pi)^2} m_\rho^2. \quad (50)$$

Performing the path integral with respect to  $H$  we obtain the one-loop Yukawa contribution to the techni-pion potential,

$$\begin{aligned} V_{y1} &= \frac{1}{2} \int \frac{d^4 Q}{(2\pi)^4} \ln \left[ Q^2 - y^2 \Pi_0^{QQ}(-Q^2) - y^2 \Pi_1^{QQ}(-Q^2) \sum_a \text{Tr}[T^a U] \text{Tr}[T^a U^\dagger] \right] \\ &\approx v_0^4 - \frac{y^2}{2(4\pi)^2} \sum_a \text{Tr}[T^a U] \text{Tr}[T^a U^\dagger] \int_0^\infty dQ^2 \Pi_1^{QQ}(-Q^2) \end{aligned} \quad (51)$$

where  $v_0$  is the contribution to the the vacuum energy and  $T^a$  are  $SU(3)$  matrices derived from (43).

One can prove that, similarly to the gauge contribution, the loop integral in (51) is finite: since  $\Pi_1^{QQ}$  is sensitive to the chiral symmetry breaking, the Operator Product Expansion demands that

$$\Pi_1^{QQ}(q^2) \stackrel{q^2 \gg \Lambda_{\text{TC}}^2}{\simeq} \frac{\langle 0 | (\bar{Q}_L \Gamma_1 Q_R) (\bar{Q}_R \Gamma_2 Q_L) | 0 \rangle}{q^4} \quad (52)$$

where  $\Gamma_{1,2}$  are appropriate matrices in techni-color and flavour space, see [16].

Contrary to the gauge contribution we are not aware of any theorem that guarantees the sign of this contribution. As an estimate the contribution above gives

$$\delta m_\pi^2 \sim \frac{y^2 m_\rho^2}{(4\pi)^2}. \quad (53)$$

Summing up all the contributions we obtain a mass matrix with the structure of eq. (24).

## A.4 Minimization of the potential

The vacuum is determined through the minimization of the potential

$$V_{\text{eff}}(\pi, \eta, H) = V_{g1} + V_{y0} + V_{y1} + m^2 |H|^2 + \lambda |H|^4 \quad (54)$$

where  $m^2 < 0$  is induced by gauge loops (section 2.1). The gauge-charged techni-pions  $\pi$  acquire a large mass from gauge loops and can be integrated out, leaving an effective potential for the lighter scalars: the elementary Higgs doublet  $H$  and the gauge-neutral techni-pion  $\eta$ . In the parameter range of interest for us,  $g \gg y$ , one has  $V_{y1} \approx 0$  and  $V_{g1} \approx \frac{1}{2} m_\pi^2 \pi^2 (1 - \eta^2/16f^2)$ , where, for simplicity, we expanded at second order in  $\eta/f$  sufficient to compute the mass of the singlet. We can freely redefine the phases of the Yukawa couplings  $y$  and  $y'$  so that  $yy'$  is real and negative. With this choice  $\eta = 0$  indeed is a local minimum of the effective potential

$$V_{\text{eff}}(\eta \ll f, H) \approx |H|^2 \left[ m^2 - 32 \frac{m_\rho^2 f^2}{m_\pi^2} \left( (|y| + |y'|)^2 - |yy'| \frac{\eta^2}{12f^2} \right) \right] + \lambda |H|^4. \quad (55)$$

Around the minimum  $\eta$  acquires a positive squared mass

$$M_\eta \sim |y| \frac{m_\rho v}{m_\pi} \quad (56)$$

without mixing with the Higgs, that receives a negative contribution to its  $m^2$  parameter.

## References

- [1] C. Kilic, T. Okui and R. Sundrum, JHEP 1002, 018 (2010) [[arXiv:0906.0577](#)].
- [2] M. Farina, D. Pappadopulo and A. Strumia, JHEP 1308, 022 (2013) [[arXiv:1303.7244](#)]. A. de Gouvea, D. Hernandez and T. M. P. Tait, Phys. Rev. D 89 (2014) 115005 [[arXiv:1402.2658](#)].
- [3] F. Englert, C. Truffin and R. Gastmans, Nucl. Phys. B 117 (1976) 407. W Bardeen, FERMILAB-CONF-95-391-T. C. T. Hill, [arXiv:hep-th/0510177](#).
- [4] A. Salvio and A. Strumia, JHEP 1406 (2014) 080 [[arXiv:1403.4226](#)].
- [5] R. Hempfling, Phys. Lett. B 379 (1996) 153 [[arXiv:hep-ph/9604278](#)]. J. P. Fatelo, J. M. Gerard, T. Hambye and J. Weyers, Phys. Rev. Lett. 74 (1995) 492. T. Hambye, Phys. Lett. B 371 (1996) 87 [[arXiv:hep-ph/9510266](#)]. W. -F. Chang, J. N. Ng and J. M. S. Wu, Phys. Rev. D 75 (2007) 115016 [[arXiv:hep-ph/0701254](#)]. R. Foot, A. Kobakhidze and R. R. Volkas, Phys. Lett. B 655 (2007) 156 [[arXiv:0704.1165](#)]. R. Foot, A. Kobakhidze, K. L. McDonald, R. R. Volkas, Phys. Rev. D 77 (2008) 035006 [[arXiv:0709.2750](#)]. S. Iso, N. Okada, Y. Orikasa, Phys. Rev. D 80 (2009) 115007 [[arXiv:0909.0128](#)]. S. Iso and Y. Orikasa, PTEP 2013 (2013) 023B08 [[arXiv:1210.2848](#)]. C. Englert, J. Jaeckel, V. V. Khoze and M. Spannowsky, [arXiv:1301.4224](#). E.J. Chun, S. Jung, H.M. Lee, [arXiv:1304.5815](#). T. Hambye and A. Strumia, Phys. Rev. D 88 (2013) 055022 [[arXiv:1306.2329](#)]. C. D. Carone and R. Ramos, Phys. Rev. D 88 (2013) 055020 [[arXiv:1307.8428](#)]. R. Foot, A. Kobakhidze, K. L. McDonald and R. R. Volkas, Phys. Rev. D 89 (2014) 115018 [[arXiv:1310.0223](#)]. A. Farzinnia, H. J. He and J. Ren, Phys. Lett. B 727 (2013) 141 [[arXiv:1308.0295](#)]. C. T. Hill, Phys. Rev. D 89 (2014) 073003 [[arXiv:1401.4185](#)]. J. Guo and Z. Kang, [arXiv:1401.5609](#). S. Benic and B. Radovic, Phys. Lett. B 732 (2014) 91 [[arXiv:1401.8183](#)]. H. Davoudiasl, I.M. Lewis, Phys. Rev. D 90, 033003 (2014) [[arXiv:1404.6260](#)]. K. Allison, C. T. Hill and G. G. Ross, [arXiv:1404.6268](#). G. M. Pelaggi, [arXiv:1406.4104](#). W. Altmannshofer, W. A. Bardeen, M. Bauer, M. Carena and J. D. Lykken, [arXiv:1408.3429](#).
- [6] T. Hur and P. Ko, Phys. Rev. Lett. 106, 141802 (2011) [[arXiv:1103.2571](#)]. M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann and K. Tuominen, Mod. Phys. Lett. A 29 (2014) 1450077 [[arXiv:1304.7006](#)].
- [7] T. Hambye and M. H. G. Tytgat, Phys. Lett. B 683 (2010) 39 [[arXiv:0907.1007](#)]. M. Holthausen, J. Kubo, K. S. Lim and M. Lindner, JHEP 1312 (2013) 076 [[arXiv:1310.4423](#)]. See also [[24](#)].
- [8] Work in progress.
- [9] K. Griest and M. Kamionkowski, Phys. Rev. Lett. 64 (1990) 615. B. von Harling and K. Petraki, [arXiv:1407.7874](#).
- [10] G. F. Giudice, G. Isidori, A. Salvio and A. Strumia, [arXiv:1412.2769](#).
- [11] Y. Bai and R. J. Hill, Phys. Rev. D 82, 111701 (2010) [[arXiv:1005.0008](#)].
- [12] R. Pasechnik, V. Beylin, V. Kuksa and G. Vereshkov, Eur. Phys. J. C 74 (2014) 2728 [[arXiv:1308.6625](#)].
- [13] E. Witten, Phys. Rev. Lett. 51, 2351 (1983).
- [14] D. Marzocca, A. Parolini and M. Serone, JHEP 1403 (2014) 099 [[arXiv:1312.5664](#)].
- [15] R. Contino, [arXiv:1005.4269](#).
- [16] M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, Nucl. Phys. B147 (1979) 385.
- [17] See e.g. M. Peskin, D.V. Schroeder, “An Introduction to Quantum Field Theory”, pag. 618. Our  $\Pi_{VV}$  is defined with the same sign as  $\Pi$  in this book.
- [18] M. Cirelli, N. Fornengo and A. Strumia, Nucl. Phys. B 753 (2006) 178 [[arXiv:hep-ph/0512090](#)]. M. Cirelli, A. Strumia and M. Tamburini, Nucl. Phys. B 787 (2007) 152 [[arXiv:0706.4071](#)].

- [19] J. Hisano, K. Ishiwata and N. Nagata, Phys. Rev. D 82 (2010) 115007 [[arXiv:1007.2601](#)]. R. J. Hill and M. P. Solon, Phys. Lett. B 707 (2012) 539 [[arXiv:1111.0016](#)].
- [20] LUX Collaboration, [arXiv:1405.5906](#).
- [21] See e.g. V. Barger, W. Y. Keung and D. Marfatia, Phys. Lett. B 696 (2011) 74 [[arXiv:1007.4345](#)].
- [22] E. Del Nobile, M. Cirelli and P. Panci, [arXiv:1307.5955](#).
- [23] Particle Data Group Collaboration, Chin. Phys. C 38 (2014) 090001.
- [24] J. Kubo, K. S. Lim and M. Lindner, [arXiv:1403.4262](#). In the past other authors considered a similar idea, but without an elementary Higgs: W. J. Marciano, Phys. Rev. D21 (1980) 2425. D. Lust, E. Papantonopoulos, K. Streng, and G. Zoupanos, Nucl. Phys. B268 (1986) 49.