

Ross Recovery with Recurrent and Transient Processes^{*†}

Hyungbin Park[‡]

Courant Institute of Mathematical Sciences,
New York University, New York, NY, USA

23 September 2015

Abstract

Recently, Ross [20] argued that it is possible to recover an objective measure from a risk-neutral measure. His model assumes that there is a finite-state Markov process X_t that drives the economy in discrete time $t \in \mathbb{N}$. Many authors extended his model to a continuous-time setting with a Markov diffusion process X_t with state space \mathbb{R} . Unfortunately, the continuous-time model fails to recover an objective measure from a risk-neutral measure in general. We determine under which information recovery is possible in the continuous-time model. It was proven that if X_t is *recurrent* under the objective measure, then recovery is possible. In this article, when X_t is *transient* under the objective measure, we investigate what information is sufficient to recover.

Keywords: Ross recovery, Markovian pricing operators, recurrence, transience

1 Introduction

Quantitative finance theory involves two related probability measures: a risk-neutral measure and an objective measure. The risk-neutral measure determines the prices of assets and options in a financial market. The risk-neutral measure is distinct from the objective measure, which describes the actual stochastic dynamics of markets. The conventional belief is that one cannot determine an objective measure by observing a risk-neutral measure. The best known example capturing this belief is the Black-Scholes model, which says that the drift of a stock under a risk-neutral measure is independent of the drift of the stock under an objective measure.

Recently, Ross [20] questioned this belief and argued that it is possible to recover an objective measure from a risk-neutral measure under some circumstances. His model assumes that there is an underlying process X_t that drives the entire economy with a finite number of states on discrete time $t \in \mathbb{N}$. This result can be of great interest to finance researchers and investors, and thus it

^{*}The author is grateful to Jonathan Goodman and Srinivasa Varadhan for technical insights. The author also appreciates two unknown referees for their constructive feedback.

[†]First version: 19 September 2014

[‡]hyungbin@cims.nyu.edu, hyungbin2015@gmail.com

is highly valuable to extend the Ross model to a continuous-time setting, which is practical and useful in finance.

In this paper, we investigate the possibility of recovering in a continuous-time setting $t \in \mathbb{R}$ with a time-homogeneous Markov diffusion process X_t with state space \mathbb{R} . In this setting, the risk-neutral measure contains some information about an objective measure. In general, however, the model unfortunately fails to recover an objective measure from a risk-neutral measure.

A key idea of recovery theory is that the reciprocal of the pricing kernel is expressed in the form $e^{\beta t} \phi(X_t)$ for some constant β and positive function $\phi(\cdot)$. For example, in the *consumption-based capital asset model* in [4] and [13], the pricing kernel is expressed in the above form. The basis of recovery theory is finding β and $\phi(\cdot)$. Thus, we obtain the pricing kernel and the relationship between the objective measure and the risk-neutral measure.

We will see that β and $\phi(\cdot)$ satisfy the second-order differential equation

$$\frac{1}{2}\sigma^2(x)\phi''(x) + k(x)\phi'(x) - r(x)\phi(x) = -\beta\phi(x). \quad (1.1)$$

Thus, recovery theory is transformed into a problem of finding a particular solution pair (β, ϕ) of this particular differential equation with $\phi(\cdot) > 0$. If such a solution pair were unique, then we could successfully recover the objective measure. Unfortunately, this approach categorically fails to achieve recovery because such a solution pair is never unique.

Many authors have extended the Ross model to a continuous-time setting and have also confronted the non-uniqueness problem. To overcome the non-uniqueness problem, all authors assumed more conditions onto their models so that the differential equation (1.1) has a unique solution pair satisfying the conditions.

Carr and Yu [5] introduced the notion of Long's discovery of the numeraire portfolio to extend the Ross model to a continuous-time setting. They assumed Long's portfolio depends on time t and the underlying process X_t , and then they derived the above differential equation (1.1). Carr and Yu also assumed that the process X_t is a time-homogeneous Markov diffusion on a *bounded* interval with regular boundaries at both endpoints. They also implicitly assumed that $\phi(\cdot)$ is in $L^2(w)$ for some measure w to apply the regular Sturm-Liouville theory, thereby obtaining a unique solution pair satisfying these conditions. Dubynskiy and Goldstein [7] explored Markov diffusion models with reflecting boundary conditions.

Walden [21] extended the results of Carr and Yu to the case that X_t is an *unbounded* process. Walden proved that recovery is possible if the process X_t is *recurrent* under the objective measure. In addition, he showed that when recovery is possible in the unbounded case, approximate recovery is possible from observing option prices on a bounded subinterval.

Qin and Linetsky [17] proved that recovery is possible if X_t is recurrent and the pricing kernel admits a Hansen-Scheinkman decomposition. They also showed that the Ross recovery has a close connection with Roger's potential approach to the pricing kernel. Borovicka, Hansen and Scheinkman [2] showed that the recovery is possible if the process X_t is *stochastically stable* under the objective measure. They also discussed applications of the recovery theory to finance and economics.

The papers of Borovicka, Hansen and Scheinkman [2], Qin and Linetsky [17] and Walden [21] assumed a common condition on X_t . Specifically, X_t is *recurrent* under the objective measure. The mathematical rationale for this condition is to overcome the non-uniqueness problem of the differential equation (1.1). Indeed, if existent, there is a unique solution pair (β, ϕ) of the equation (1.1) satisfying this condition and we will review this condition in Section 5.1.

In this article, we investigate the possibility of recovery when the process X_t is *transient* under the objective measure. We explore in this case what information is sufficient to recover. One of the main contributions is that if β is known and if X_t is *non-attracted* to the left (or right) boundary under the objective measure, then recovery is possible. To achieve this, we establish a graphical understanding of recovery theory. This topic is discussed in Section 4 and 5. In Section 6, two examples of recovery theory are explored: the Cox-Ingersoll-Ross (CIR) interest rate model and the Black-Scholes stock model. Section 7 summarizes this article.

2 Markovian pricing operators

A financial market is defined as a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ having a one-dimensional Brownian motion B_t with the filtration $\mathcal{F} = (\mathcal{F}_t)_{t=0}^\infty$ generated by B_t . All the processes in this article are assumed to be adapted to the filtration \mathcal{F} . \mathbb{P} is the objective measure of this market. We assume that there are a state variable X_t and a positive numeraire G_t in the market.

Let \mathbb{Q} be an equivalent measure on the market $(\Omega, \mathcal{F}, \mathbb{P})$ such that each risky asset discounted by the numeraire G_t is a martingale under measure \mathbb{Q} . It is customary that this measure \mathbb{Q} is referred to as a risk-neutral measure when G_t is a money market account. In this article, however, for any given positive numeraire G_t , we say \mathbb{Q} is a risk-neutral measure with respect to G_t . Set the Radon-Nikodym derivative by $\Sigma_t = \frac{d\mathbb{Q}}{d\mathbb{P}}|_{\mathcal{F}_t}$, which is known to be a martingale process on $(\Omega, \mathcal{F}, \mathbb{P})$ for $0 < t < T$. Using the martingale representation theorem, we can write in the stochastic differential equation form $d\Sigma_t = -\rho_t \Sigma_t dB_t$ for some ρ_t . It is well-known that W_t defined by

$$dW_t := \rho_t dt + dB_t \tag{2.1}$$

is a Brownian motion under \mathbb{Q} . We define *the reciprocal of the pricing kernel* by $L_t = G_t/\Sigma_t$.

Assumption 1. The state variable X_t is a time-homogeneous Markov diffusion process satisfying

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = \xi.$$

The range of X_t is an open interval $I = (c, d)$ with $-\infty \leq c < d \leq \infty$. $b(\cdot)$ and $\sigma(\cdot)$ are continuously differentiable on I and that $\sigma(x) > 0$ for $x \in (c, d)$.

It is implicitly assumed that both endpoints are unattainable because the range of the process is an open interval.

Assumption 2. The dynamics of the numeraire G_t is determined by X_t . More precisely, G_t follows

$$\frac{dG_t}{G_t} = (r(X_t) + v^2(X_t)) dt + v(X_t) dW_t, \quad G_0 = 1.$$

Assume that r and v are continuously differentiable on I and

$$\exp\left(-\frac{1}{2} \int_0^t v^2(X_s) ds - \int_0^t v(X_s) dW_s\right)$$

is a martingale.

We assume that we can extract these four functions $b(\cdot), \sigma(\cdot), r(\cdot)$ and $v(\cdot)$ from market prices data, thus they are assumed to be known ex ante. The above martingale assumption is to define a new measure by using the Girsanov theorem, for example, in the proof of Theorem E.1. It is noteworthy that if there is a money market account with interest rate, denoted by r_t , in the market, then r_t is equal to $r(X_t)$ because $e^{\int_0^t r_s ds} \cdot G_t^{-1}$ is a martingale under \mathbb{Q} .

Assumption 3. Assume that (the reciprocal of) the pricing kernel L_t is *transition independent* in the sense that there are a positive function $\phi \in C^2(I)$ and a real number β such that

$$L_t = e^{\beta t} \phi(X_t) \phi^{-1}(\xi) . \quad (2.2)$$

In this case, we say (β, ϕ) is a *principal pair* of the market.

The basis of recovery theory is finding the principal pair (β, ϕ) and then obtaining the objective measure \mathbb{P} by setting the Radon-Nikodym derivative

$$\left. \frac{d\mathbb{P}}{d\mathbb{Q}} \right|_{\mathcal{F}_t} = \Sigma_t^{-1} = e^{\beta t} \phi(X_t) \phi^{-1}(\xi) G_t^{-1} .$$

One important aspect in implementing the recovery approach is to decide how to choose state variables X_t . Many processes can serve as a state variable and the choice of a state variable depends on the purpose of use. One way is the short interest rate r_t . Investors interested in the price of bonds want to find the dynamics of r_t under an objective measure. Plenty of examples with interest rate state variables can be found in [18]. Another way is a stock market index process such as the Dow Jones Industrial Average and Standard & Poor's (S&P) 500. Refer to [1] for an empirical analysis of recovery theory with the state variable S&P 500.

3 Transformed measures

We investigate how recovery theory is transformed into a problem of a differential equation. Applying the Ito formula to the definition of L_t , we know

$$dL_t = (r(X_t) + v^2(X_t) + v(X_t)\rho_t) L_t dt + (v(X_t) + \rho_t)L_t dW_t .$$

From (2.2), we also have

$$dL_t = \left(\beta + \frac{1}{2}(\sigma^2 \phi'' \phi^{-1})(X_t) + (b\phi' \phi^{-1})(X_t) \right) L_t dt + (\sigma \phi' \phi^{-1})(X_t) L_t dW_t .$$

By comparing these two equations, we obtain

$$\begin{cases} \frac{1}{2}\sigma^2 \phi'' + (b - v\sigma)\phi' - r\phi = -\beta\phi , \\ \rho_t = (\sigma\phi' \phi^{-1} - v)(X_t) . \end{cases} \quad (3.1)$$

For convenience, set $k(x) := (b - v\sigma)(x)$. Using notation \mathcal{L} defined by $\mathcal{L}\phi(x) = \frac{1}{2}\sigma^2(x)\phi''(x) + k(x)\phi'(x) - r(x)\phi(x)$, we have the following theorem.

Theorem 3.1. Let (β, ϕ) be the principal pair of the market. Then (β, ϕ) satisfies $\mathcal{L}\phi = -\beta\phi$.

In other words, if (λ, h) is a solution pair of $\mathcal{L}h = -\lambda h$ with $h > 0$, then (λ, h) is a candidate pair for the principal pair of X_t .

We are interested in a solution pair (λ, h) of $\mathcal{L}h = -\lambda h$ with positive function h . There are two possibilities.

- (i) there is no positive solution h for any $\lambda \in \mathbb{R}$, or

- (ii) there exists a number $\bar{\beta}$ such that it has two linearly independent positive solutions for $\lambda < \bar{\beta}$, has no positive solution for $\lambda > \bar{\beta}$ and has one or two linearly independent solutions for $\lambda = \bar{\beta}$.

Refer to page 146 and 149 in [16]. In this article, we implicitly assumed the second case by Assumption 3.

It is easily checked that

$$e^{\lambda t} h(X_t) h^{-1}(\xi) G_t^{-1}$$

is a local martingale under \mathbb{Q} . When this is a martingale, one can *attempt* to recover the objective measure \mathbb{P} by setting this as a Radon-Nikodym derivative.

Definition 1. Let (λ, h) be a solution pair of $\mathcal{L}h = -\lambda h$ with positive function h . Suppose that $e^{\lambda t} h(X_t) h^{-1}(\xi) G_t^{-1}$ is a martingale. A measure obtained from the risk-neutral measure \mathbb{Q} by the Radon-Nikodym derivative

$$\left. \frac{d\cdot}{d\mathbb{Q}} \right|_{\mathcal{F}_t} = e^{\lambda t} h(X_t) h^{-1}(\xi) G_t^{-1}$$

is called *the transformed measure* with respect to the pair (λ, h) .

Clearly, the transformed measure with respect to the principal pair is the objective measure \mathbb{P} . We have the following proposition by (2.1) and (3.1).

Proposition 3.2. A process B_t^h defined by $dB_t^h = -(\sigma h' h^{-1} - v)(X_t) dt + dW_t$ is a Brownian motion under the transformed measure with respect to (λ, h) . Furthermore, X_t follows

$$\begin{aligned} dX_t &= (b - v\sigma + \sigma^2 h' h^{-1})(X_t) dt + \sigma(X_t) dB_t^h \\ &= (k + \sigma^2 h' h^{-1})(X_t) dt + \sigma(X_t) dB_t^h . \end{aligned} \tag{3.2}$$

Occasionally, we use the notation B_t instead of B_t^h without ambiguity.

Even when $e^{\lambda t} h(X_t) h^{-1}(\xi) G_t^{-1}$ is not a martingale, we can consider the diffusion process corresponding to (3.2).

Definition 2. The diffusion process X_t defined by

$$dX_t = (k + \sigma^2 h' h^{-1})(X_t) dt + \sigma(X_t) dB_t$$

is called the diffusion process *induced by* (λ, h) .

4 Recurrence and transience

4.1 Mathematical preliminaries

We establish the mathematical preliminaries for recurrent and transient processes. The contents of this section are indebted to [8], [12] and [14]. Consider the diffusion process induced by (λ, h) :

$$dX_t = \mu(X_t) dt + \sigma(X_t) dB_t, \quad X_0 = \xi,$$

where $\mu(\cdot) = (k + \sigma^2 h' h^{-1})(\cdot)$. A measure S defined by

$$dS(x) := e^{-\int_{\xi}^x \frac{2\mu(y)}{\sigma^2(y)} dy} dx = \frac{h^2(\xi)}{h^2(x)} e^{-\int_{\xi}^x \frac{2k(y)}{\sigma^2(y)} dy} dx$$

is called *the scale measure* of the process with respect to the pair (λ, h) .

Definition 3. The left boundary c is *attracting* if $S((c, \xi]) < \infty$ and *non-attracting* otherwise. Similarly, we say the right boundary d is *attracting* if $S([\xi, d)) < \infty$ and *non-attracting* otherwise.

Define a stopping time τ by the following way. Let $(c_n)_{n=1}^\infty, (d_n)_{n=1}^\infty$ be strictly monotone sequences with limits c and d , respectively. Set $\tau_n := \inf \{ t > 0 \mid X_t \notin (c_n, d_n) \}$ and $\tau := \lim_{n \rightarrow \infty} \tau_n$.

Proposition 4.1. The left boundary c is non-attracting if and only if

$$\text{Prob} \left(\lim_{t \rightarrow \tau} X_t = c \right) = 0 .$$

It is similar to the right boundary d .

Proposition 4.2. X_t is recurrent if and only if both boundary points c and d are non-attracting.

4.2 Graphical understanding

We establish a graphical understanding of recovery theory. Recall that $X_0 = \xi$ in Assumption 1. The purpose of this section is to understand the graphs shown in Figure 1 and 2. A solution h of a second-order differential equation is uniquely determined by the initial value and the initial velocity. By normalizing, we may assume $h(\xi) = 1$ such that a solution is determined by $h'(\xi)$. In this section, we describe the relationship between $h'(\xi)$ and the recovery of X_t with respect to h . Occasionally, we use the terminology without ambiguity: the transformed measure with respect to *tuple* $(\lambda, h'(\xi))$ means the transformed measure with respect to *pair* (λ, h) . The two terms tuple and pair will be used to distinguish between these meanings.

Definition 4. We say $(\lambda, h'(\xi)) \in \mathbb{R}^2$ is a candidate tuple or we say (λ, h) is a candidate pair if (λ, h) is a solution pair of $\mathcal{L}h = -\lambda h$ with $h(\cdot) > 0$ and $h(\xi) = 1$. Denote the set of the candidate tuples by \mathcal{C} .

$$\mathcal{C} := \{ (\lambda, h'(\xi)) \in \mathbb{R}^2 \mid \mathcal{L}h = -\lambda h, h > 0, h(\xi) = 1 \} .$$

We investigate graphical properties of \mathcal{C} . Let $\bar{\beta}$ be the maximum value of the first coordinate of elements of \mathcal{C} , that is, $\bar{\beta} := \max \{ \lambda \mid (\lambda, z) \in \mathcal{C} \}$. The maximum $\bar{\beta}$ is achieved as we discussed in Section 3. For any λ with $\lambda \leq \bar{\beta}$, we set

$$M_\lambda := \sup_{(\lambda, z) \in \mathcal{C}} z, \quad m_\lambda := \inf_{(\lambda, z) \in \mathcal{C}} z .$$

Proposition 4.3. Let $\lambda \leq \bar{\beta}$. For any z with $m_\lambda \leq z \leq M_\lambda$, the tuple (λ, z) is in \mathcal{C} .

Therefore, the supremum and infimum are, in fact, the maximum and minimum, respectively. Furthermore, the λ -slide of \mathcal{C} is a connected and compact set. See Appendix A for proof.

Theorem 4.4. The diffusion process induced by tuple (λ, M_λ) is non-attracted to the left boundary. For z with $m_\lambda \leq z < M_\lambda$, the diffusion process induced by tuple (λ, z) , is attracted to the left boundary.

Similarly, the diffusion process induced by tuple (λ, m_λ) is non-attracted to the right boundary. For z with $m_\lambda < z \leq M_\lambda$, the diffusion process induced by tuple (λ, z) is attracted to the right boundary. Therefore, for z with $m_\lambda < z < M_\lambda$, the diffusion process induced by tuple (λ, z) is attracted to both boundaries. For proof, see Appendix B.

Proposition 4.5. M_λ is a strictly decreasing function of λ and m_λ is a strictly increasing function of λ for $\lambda \leq \bar{\beta}$.

See Appendix C for proof.

Corollary 4.6. For $\bar{\beta}$, there are two possibilities:

- (i) there is a unique number \bar{z} such that $(\bar{\beta}, \bar{z})$ is in \mathcal{C} . In this case, the tuple $(\bar{\beta}, \bar{z})$ is the unique tuple in \mathcal{C} such that the induced diffusion process is recurrent.
- (ii) there is an infinite number of z 's such that $(\bar{\beta}, z)$ is in \mathcal{C} . In this case, for any such tuple $(\bar{\beta}, z)$ in \mathcal{C} , the induced diffusion process X_t is transient.

Refer to Section 6 for examples of (i) and also see Appendix F for an example of (ii). In Figure 1, the left graph is the case of (i) and the right graph is the case of (ii).




Figure 1: Candidate sets

We now explore a particular subset of \mathcal{C} . In general, for a candidate pair (λ, h) , $e^{\lambda t} h(X_t) G_t^{-1}$ is a local martingale. We are interested in pairs that induce martingales.

Definition 5. Let (λ, h) be a candidate pair. We say $(\lambda, h'(\xi))$ is an admissible tuple or we say (λ, h) is an admissible pair if $e^{\lambda t} h(X_t) G_t^{-1}$ is a martingale. Denote the set of the admissible tuples by \mathcal{A} .

$$\mathcal{A} := \{(\lambda, h'(\xi)) \in \mathbb{R}^2 \mid (\lambda, h) \text{ is a candidate pair and } e^{\lambda t} h(X_t) G_t^{-1} \text{ is a martingale}\}.$$

One of the main purposes of this section is to investigate the graphical properties of \mathcal{A} with the notion of recurrence and transience. It is noteworthy that \mathcal{C} and \mathcal{A} depend on ξ .

Proposition 4.7. Let $\lambda \leq \bar{\beta}$ and let $m_\lambda < z < M_\lambda$. If two tuples (λ, m_λ) and (λ, M_λ) are in \mathcal{A} , then (λ, z) is in \mathcal{A} . If at least one of (λ, m_λ) and (λ, M_λ) is not in \mathcal{A} , then (λ, z) is not in \mathcal{A} .

See Appendix D for proof.

Proposition 4.8. Let $\delta < \lambda \leq \bar{\beta}$. If (δ, M_δ) is in \mathcal{A} , then (λ, M_λ) is also in \mathcal{A} . Similarly, if (δ, m_δ) is in \mathcal{A} , then (λ, m_λ) is also in \mathcal{A} .

Refer to Appendix E for proof.

Proposition 4.9. Consider the case of (i) in Corollary 4.6. For $\bar{\beta}$, suppose that there is a unique number \bar{z} such that $(\bar{\beta}, \bar{z})$ is in \mathcal{C} . Then $(\bar{\beta}, \bar{z})$ is in \mathcal{A} .

For proof, see [16]. Figure 2 displays two examples of admissible set and summarizes this section.




Figure 2: Admissible sets

5 Recurrent and transient recovery

5.1 Recurrent recovery

We review recovery theory with the assumption that X_t is *recurrent* under the objective measure. With this assumption, we can successfully recover the objective measure from the risk-neutral measure. This theory is especially useful when the state variable X_t is an interest rate process because the actual dynamics of an interest rate is usually recurrent or mean-reverting in the actual real-world measure.

Proposition 5.1. If it exists, there is a unique admissible pair $(\bar{\beta}, \bar{\phi})$ of $\mathcal{L}\bar{\phi} = -\bar{\beta}\bar{\phi}$ with $\bar{\phi} > 0$ such that X_t is recurrent under the transformed measure with respect to the pair $(\bar{\beta}, \bar{\phi})$. In this case, we have $\bar{\beta} = \max\{\lambda \mid (\lambda, z) \in \mathcal{A}\}$.

This proposition is easily obtained from Section 4.2 and gives the following theorem.

Theorem 5.2. (Recurrent recovery) Suppose X_t is recurrent under the objective measure \mathbb{P} . We can then recover the objective measure \mathbb{P} from the risk-neutral measure \mathbb{Q} .

5.2 Transient recovery

We encounter several conditions under which we can recover the objective measure when X_t is transient under the objective measure. The state variable X_t is always transient under the transformed measure with respect to an admissible pair (λ, h) for any $\lambda < \bar{\beta}$. Therefore, without further information, recovery is impossible when the process X_t is transient under the objective measure.

We assume that we know the value β . Despite knowing this value β , we cannot achieve recovery in general. However, the following theorem says that recovery is possible under some circumstances. The proof is direct from Proposition 4.7.

Theorem 5.3. (Transient recovery) Suppose we know the value β . If only one of (β, m_β) and (β, M_β) is an admissible tuple, then we can recover the objective measure \mathbb{P} from the risk-neutral measure \mathbb{Q} .

When both of (β, m_β) and (β, M_β) are admissible tuples, we cannot uniquely determine the objective measure \mathbb{P} because there is an infinite number of admissible pairs.

We investigate another way for recovery. We will see that there is only one way to recover the objective measure such that X_t is non-attracted to the left (or right) boundary.

Proposition 5.4. For any fixed λ with $\lambda \leq \bar{\beta}$, if it exists, there is a unique admissible pair (λ, h) such that X_t is non-attracted to the left boundary under the corresponding transformed measure. In this case, $h'(\xi) = M_\lambda$.

Similarly, if it exists, there is a unique admissible pair (λ, h) such that X_t is non-attracted to the *right* boundary under the corresponding transformed measure. In this case, $h'(\xi) = m_\lambda$. This proposition is easily obtained from Section 4.2 and gives the following theorem.

Theorem 5.5. (Transient recovery) Suppose we know the value β . If X_t is non-attracted to the left (or right) boundary under the objective measure \mathbb{P} , then we can recover the objective measure \mathbb{P} from the risk-neutral measure \mathbb{Q} .

If X_t is attracted to both boundaries under the objective measure \mathbb{P} , then both (β, m_β) and (β, M_β) are admissible tuples. Thus there is an infinite number of admissible tuples such that X_t is attracted to both boundaries under the corresponding transformed measure.

We now shift our attention to the choice of β . When X_t is non-attracted to the left (right) boundary, to recover the objective measure, we confront a problem of determining the value β . How can we choose the value? One way is to use the long-term yield of bonds, which is defined by

$$\lim_{t \rightarrow \infty} \left(-\frac{1}{t} \cdot \log \mathbb{E}^{\mathbb{Q}} [G_t^{-1}] \right) .$$

See [15] and [18] as a reference.

6 Applications

6.1 Interest rates

We investigate the recovery theorem when the state variable X_t is an interest rate process r_t and the numeraire is the money market account:

$$X_t = r_t, \quad G_t = e^{\int_0^t r_s ds} .$$

Assume that r_t follows $dr_t = k(r_t) dt + \sigma(r_t) dW_t$. Then

$$\frac{1}{2} \sigma^2(r) h''(r) + k(r) h'(r) - r h(r) = -\lambda h .$$

is the corresponding second-order differential equation.

Example 6.1. We explore the CIR model:

$$dr_t = a(\theta - r_t) dt + \sigma\sqrt{r_t} dW_t$$

with the Feller condition $2a\theta \geq \sigma^2$. Consider

$$\frac{1}{2}\sigma^2 r h''(r) + a(\theta - r)h'(r) - r h(r) = \lambda h$$

with $h(r_0) = 1$. Set $k := \frac{\sqrt{a^2 + 2\sigma^2} - a}{\sigma^2}$. It can be shown that the maximum value $\bar{\beta}$ is equal to $ka\theta$. For any fixed λ with $\lambda \leq ka\theta$, denote the functions corresponding to tuples (λ, M_λ) and (λ, m_λ) by $h_\lambda(\cdot)$ and $g_\lambda(\cdot)$, respectively. Then we have $h_\lambda(x) = \psi_\lambda(x)/\psi_\lambda(r_0)$ where

$$\psi_\lambda(x) := e^{-kx} K\left(\frac{\lambda - ka\theta}{\sqrt{a^2 + 2\sigma^2}}, \frac{2a\theta}{\sigma^2}, \frac{2x\sqrt{a^2 + 2\sigma^2}}{\sigma^2}\right).$$

Here, $K(\cdot, \cdot, \cdot)$ is the Kummer confluent hypergeometric function. Refer to [17] for more details about g_λ . It can be shown that (λ, M_λ) is an admissible tuple, but (λ, m_λ) is not. Thus, the admissible set \mathcal{A} is described by

$$\mathcal{A} = \{(\lambda, h'_\lambda(r_0)) \in \mathbb{R}^2 \mid \lambda \leq ka\theta\}.$$

See [17] for more details.

Interest rates are usually recurrent in the real world; therefore, we assume that r_t is recurrent under the objective measure. $(ka\theta, e^{-k(r-r_0)})$ is the only admissible pair that induces the recurrent X_t under the corresponding transformed measure and under which the X_t is expressed by

$$dr_t = \sqrt{a^2 + 2\sigma^2} \left(\frac{a\theta}{\sqrt{a^2 + 2\sigma^2}} - r_t \right) dt + \sigma\sqrt{r_t} dB_t.$$

□

6.2 Stock prices

We investigate the recovery theorem when the state variable X_t is a stock price S_t and the dividends of the stock are paid out continuously with rate $\delta(S_t) dt$. Here $\delta(\cdot)$ is a deterministic function and the case of $\delta(\cdot) = 0$ is not excluded. Let the numeraire G_t be the wealth process defined by

$$G_t = S_t e^{\int_0^t \delta(S_u) du}.$$

Assume that the dynamics of S_t is given by

$$dS_t = (r(S_t) - \delta(S_t) + \sigma^2(S_t))S_t dt + \sigma(S_t)S_t dW_t.$$

Then it follows that

$$\frac{dG_t}{G_t} = (r(S_t) + \sigma^2(S_t)) dt + \sigma(S_t) dW_t.$$

The corresponding second-order differential equation is

$$\frac{1}{2}\sigma^2(s)s^2 h''(s) + (r(s) - \delta(s))s h'(s) - r(s)h(s) = -\lambda h(s).$$

It is noteworthy that that if there is a money market account with a constant interest rate r in the market, then $r(S_t) = r$.

Example 6.2. Assume that the state variable is a stock price S_t and the dividends of the stock are paid out continuously with rate δdt . Suppose S_t follows a geometric Brownian motion

$$dS_t = (r - \delta + \sigma^2)S_t dt + \sigma S_t dW_t, \quad S_0 = 1$$

and the numeraire is $G_t = S_t e^{\delta t}$. Consider

$$\frac{1}{2}\sigma^2 s^2 h''(s) + (r - \delta)sh'(s) - rh(s) = -\lambda h(s).$$

We want to find the candidate set and the admissible set. For $\lambda < \frac{1}{2}\left(\frac{\sigma}{2} - \frac{r-\delta}{\sigma}\right)^2 + r$, these solutions are given by $h(s) = c s^{l_1} + (1 - c) s^{l_2}$ for $0 \leq c \leq 1$ where

$$l_1 = \frac{1}{2} - \frac{r - \delta}{\sigma^2} - \sqrt{\left(\frac{1}{2} - \frac{r - \delta}{\sigma^2}\right)^2 + \frac{2(r - \lambda)}{\sigma^2}},$$

$$l_2 = \frac{1}{2} - \frac{r - \delta}{\sigma^2} + \sqrt{\left(\frac{1}{2} - \frac{r - \delta}{\sigma^2}\right)^2 + \frac{2(r - \lambda)}{\sigma^2}},$$

thus we have $m_\lambda = l_1$, $M_\lambda = l_2$. For $\lambda = \frac{1}{2}\left(\frac{\sigma}{2} - \frac{r-\delta}{\sigma}\right)^2 + r$, the only positive solution h with $h(1) = 1$ is $h(x) = x^{\frac{1}{2} - \frac{r-\delta}{\sigma^2}}$ and $m_\lambda = M_\lambda = \frac{1}{2} - \frac{r-\delta}{\sigma^2}$. For $\lambda > \frac{1}{2}\left(\frac{\sigma}{2} - \frac{r-\delta}{\sigma}\right)^2 + r$, there are no positive solutions. Thus, the candidate set is obtained. It can be easily shown that the admissible set is equal to the candidate set.

If S_t is recurrent under the objective measure, then the transformed measure with respect to tuple $\left(\frac{1}{2}\left(\frac{\sigma}{2} - \frac{r-\delta}{\sigma}\right)^2 + r, \frac{1}{2} - \frac{r-\delta}{\sigma^2}\right)$ is the objective measure. We used Theorem 5.2. Under this measure, S_t follows

$$dS_t = \frac{1}{2}\sigma^2 S_t dt + \sigma S_t dB_t.$$

We now assume that S_t is non-attracted to 0 and the value $\beta \left(\leq \frac{1}{2}\left(\frac{\sigma}{2} - \frac{r-\delta}{\sigma}\right)^2 + r\right)$ is known. Then, by Theorem 5.5, the transformed measure with respect to tuple

$$(\beta, M_\beta) = \left(\beta, \frac{1}{2} - \frac{r - \delta}{\sigma^2} + \sqrt{\left(\frac{1}{2} - \frac{r - \delta}{\sigma^2}\right)^2 + \frac{2(r - \beta)}{\sigma^2}}\right)$$

is the objective measure, under which S_t follows

$$dS_t = \left(\frac{\sigma^2}{2} + \sqrt{\left(\frac{\sigma^2}{2} - (r - \delta)\right)^2 + 2\sigma^2(r - \beta)}\right) S_t dt + \sigma S_t dB_t.$$

As a particular case, if β is equal to the long-term yield of bonds

$$\lim_{t \rightarrow \infty} \left(-\frac{1}{t} \cdot \log \mathbb{E}^{\mathbb{Q}} [G_t^{-1}]\right) = r,$$

then S_t follows

$$dS_t = \left(\frac{\sigma^2}{2} + \left|r - \delta - \frac{\sigma^2}{2}\right|\right) S_t dt + \sigma S_t dB_t$$

under the objective measure. The numeraire G_t satisfies $dG_t = \mu G_t dt + \sigma G_t dB_t$ with

$$\mu := \frac{\sigma^2}{2} + \delta + \left| r - \delta - \frac{\sigma^2}{2} \right|.$$

Thus, the Sharpe ratio of G_t is

$$\frac{\mu - r}{\sigma} = \begin{cases} \sigma + \frac{2(\delta - r)}{\sigma} & \text{if } r < \delta + \frac{\sigma^2}{2} \\ 0 & \text{if } r \geq \delta + \frac{\sigma^2}{2}. \end{cases}$$

□

7 Conclusion

This article extended the Ross model to a continuous-time setting model. In a continuous-time setting, the risk-neutral measure contains some information about the objective measure. Unfortunately, the model fails to recover an objective measure from a risk-neutral measure. We discussed several conditions under which the recovery of the objective measure from the risk-neutral measure is possible in a continuous-time model. When the state variable X_t is recurrent under the objective measure, recovery is possible.

We also determined the type of information that is sufficient for recovery when X_t is transient. It was shown that when X_t is transient, recovery is possible if the value β is known and only one of (β, m_β) and (β, M_β) is an admissible tuple. We can also recover the objective measure if the value β is known and X_t is non-attracted to the left (or right) boundary.

The following extensions for future research are suggested. First, it would be interesting to find sufficient conditions under which recovery is possible when X_t is attracted to both boundaries. We could not offer such conditions in this article. Second, it would be valuable to find financially and economically reasonable ways to determine β . Finally, much work remains to be conducted on the implementation and empirical testing of recovery theory in future research.

A Proof of Proposition 4.3

Proposition A.1. A solution h of $\mathcal{L}h = -\lambda h$ can be expressed by $h = uq$, where $q(x) = e^{-\int_0^x \frac{k(y)}{\sigma^2(y)} dy}$ and u is a solution of

$$u''(x) + \left(-\frac{d}{dx} \left(\frac{k(x)}{\sigma^2(x)} \right) - \frac{k^2(x)}{\sigma^4(x)} + \frac{2(-r(x) + \lambda)}{\sigma^2(x)} \right) u(x) = 0.$$

This can be shown by direct calculation. For more details, refer to [21], page 36.

We now prove Proposition 4.3.

Proof. For convenience, we may assume that $\xi = 0$, the left boundary is $-\infty$ and the right boundary is ∞ . Let h be a positive solution of $\mathcal{L}h = -\lambda h$ with $h(0) = 1$. Another solution, which is independent of h , is

$$h(x) \int_0^x \frac{1}{h^2(y)} e^{-\int_0^y \frac{2k(z)}{\sigma^2(z)} dz} dy.$$

(It can be obtained by direct calculation). Let S be the scale measure with respect to the pair (λ, h) , then

$$h(x) \int_0^x \frac{1}{h^2(y)} e^{-\int_0^y \frac{2k(z)}{\sigma^2(z)} dz} dy = h(x) \cdot S([0, x]).$$

The general solutions of $\mathcal{L}h = -\lambda h$ are expressed by $h(x)(c_1 + c_2 \cdot S([0, x]))$. Assume that at least one of $S((-\infty, 0])$ and $S([0, \infty))$ is finite. (Otherwise, h is the unique positive solution of $\mathcal{L}h = -\lambda h$, so $m_\beta = z = M_\beta$, in which case we have nothing to prove). We may assume $S((-\infty, 0]) < \infty$. Set $B := S((-\infty, 0])$. The general (normalized to $h_c(0) = 1$) solution is expressed by

$$h_c(x) := h(x) \left((1 - c) + \frac{c}{B} \cdot S((-\infty, x]) \right),$$

which is a positive function for and only for

$$\begin{cases} 0 \leq c \leq 1 & \text{if } S([0, \infty)) = \infty \\ -\frac{B}{D} \leq c \leq 1 & \text{if } D := S([0, \infty)) < \infty \end{cases}$$

Using $h'_c(0) = h'(0) + \frac{c}{B}$, we have that

$$\begin{cases} M_\lambda = h'(0) + \frac{1}{B}, & m_\lambda = h'(0) & \text{if } S([0, \infty)) = \infty \\ M_\lambda = h'(0) + \frac{1}{B}, & m_\lambda = h'(0) - \frac{1}{D} & \text{if } D = S([0, \infty)) < \infty \end{cases}$$

Furthermore, $h'_c(0)$ can be any value in $[m_\lambda, M_\lambda]$. Hence, for any z with $m_\lambda \leq z \leq M_\lambda$, the tuple (λ, z) is in \mathcal{A} . This completes the proof. \square

B Proof of Theorem 4.4

Proof. For convenience, we may assume that $\xi = 0$, the left boundary is $-\infty$ and the right boundary is ∞ . Suppose that the diffusion process induced by tuple (λ, M_λ) is attracted to the left boundary. Denote the scale measure with respect to tuple (λ, M_λ) by S . We have $B := S((-\infty, 0]) < \infty$. By the proof of Proposition 4.3, we know that $(\lambda, h'(0) + \frac{1}{B}) = (\lambda, M_\lambda + \frac{1}{B})$ is in \mathcal{C} . This is a contradiction.

We now show that for z with $m_\lambda \leq z < M_\lambda$, the diffusion process induced by tuple (λ, z) is attracted to the left boundary. Let h and g be the functions corresponding to tuple (λ, M_λ) and (λ, z) , respectively. Recall $q(x) = e^{-\int_0^x \frac{k(y)}{\sigma^2(y)} dy}$ in Proposition A.1. Write $h = uq$ and $g = vq$. It can be easily checked that $u(0) = v(0) = 1$ and $u'(0) > v'(0)$. Set $\Gamma := \frac{u'}{u} - \frac{v'}{v}$. By direct calculation, we have $\Gamma' = -\Gamma^2 - \frac{2v'}{v}\Gamma$. Because $\Gamma(0) > 0$ and $\Gamma = 0$ is an equilibrium point, we know that $\Gamma(x) \geq 0$ for all x . By differentiating $\Gamma(x) e^{\int_0^x (\Gamma(y) + \frac{2v'(y)}{v(y)}) dy}$, we have

$$\left(\Gamma'(x) + \Gamma^2(x) + \frac{2v'(x)}{v(x)}\Gamma(x) \right) e^{\int_0^x (\Gamma(y) + \frac{2v'(y)}{v(y)}) dy} = 0.$$

Thus, $\Gamma(x) e^{\int_0^x (\Gamma(y) + \frac{2v'(y)}{v(y)}) dy} = \Gamma(0)$, which yields

$$v^{-2}(x) = e^{-\int_0^x \frac{2v'}{v} dy} = \frac{\Gamma(x)}{\Gamma(0)} e^{\int_0^x \Gamma(y) dy}.$$

We obtain that

$$\begin{aligned}
\int_{-\infty}^0 \frac{1}{g^2(y)} e^{-\int_0^y \frac{2k(z)}{\sigma^2(z)} dz} dy &= \int_{-\infty}^0 v^{-2}(y) dy \\
&= \frac{1}{\Gamma(0)} \int_{-\infty}^0 \Gamma(y) e^{\int_0^y \Gamma(z) dz} dy \\
&= \frac{1}{\Gamma(0)} \left(1 - e^{-\int_{-\infty}^0 \Gamma(y) dy}\right) \leq \frac{1}{\Gamma(0)} < \infty.
\end{aligned}$$

Therefore, the diffusion process induced by (λ, z) is attracted to $-\infty$. \square

C Proof of Proposition 4.5

Proof. Let $\delta < \lambda$. Let g be a positive function of $\mathcal{L}g = -\delta g$ with $g(0) = 1$ and $g'(0) = M_\delta$. We show that there does not exist a positive function h of $\mathcal{L}h = -\lambda h$ with $h(0) = 1$ and $h'(0) = M_\delta$. This implies that $M_\delta > M_\lambda$. Suppose h is a positive solution of $\mathcal{L}h = -\lambda h$ with $h(0) = 1$ and $h'(0) = M_\delta$. Recall $q(x) = e^{-\int_0^x \frac{k(y)}{\sigma^2(y)} dy}$ in Proposition A.1. Write $h = uq$ and $g = vq$. Define $\Gamma = \frac{u'}{u} - \frac{v'}{v}$. Then

$$\Gamma' = -\Gamma^2 - \frac{2v'}{v}\Gamma - \frac{2(\lambda - \delta)}{\sigma^2}.$$

From $\Gamma(0) = 0$, we have that $\Gamma(x) > 0$ for $x < 0$ because if Γ ever gets close to 0, then term $-\frac{2(\lambda - \delta)}{\sigma^2}$ dominates the right hand side of the equation. Choose x_0 with $x_0 < 0$. For $x < x_0$, we have

$$-\frac{2v'(x)}{v(x)} = \frac{\Gamma'(x)}{\Gamma(x)} + \Gamma(x) + \frac{2(\lambda - \delta)}{\sigma^2(x)} \cdot \frac{1}{\Gamma(x)}.$$

Integrating from x_0 to x ,

$$-2 \ln \frac{v(x)}{v(x_0)} = \ln \frac{\Gamma(x)}{\Gamma(x_0)} + \int_{x_0}^x \Gamma(y) dy + \int_{x_0}^x \frac{2(\lambda - \delta)}{\sigma^2(y)} \cdot \frac{1}{\Gamma(y)} dy$$

which leads to

$$\frac{v^2(x_0)}{v^2(x)} \leq \frac{\Gamma(x)}{\Gamma(x_0)} e^{\int_{x_0}^x \Gamma(y) dy}$$

for $x < x_0$. Thus,

$$\begin{aligned}
\int_{-\infty}^{x_0} \frac{1}{g^2(y)} e^{-\int_0^y \frac{2k(z)}{\sigma^2(z)} dz} dy &= \int_{-\infty}^{x_0} \frac{1}{v^2(y)} dy \\
&\leq (\text{constant}) \cdot \int_{-\infty}^{x_0} \Gamma(y) e^{\int_{x_0}^y \Gamma(w) dw} dy \\
&= (\text{constant}) \cdot \left(1 - e^{-\int_{-\infty}^{x_0} \Gamma(w) dw}\right) \\
&\leq (\text{constant}) \\
&< \infty.
\end{aligned}$$

This implies that the diffusion process induced by the tuple (δ, M_δ) is attracted to the left boundary, which is a contradiction. \square

D Proof of Proposition 4.7

Suppose that two tuples (λ, m_λ) and (λ, M_λ) are in \mathcal{A} . Let h_m and h_M be the functions corresponding to (λ, m_λ) and (λ, M_λ) , respectively. Then $e^{\lambda t} h_m(X_t) G_t^{-1}$ and $e^{\lambda t} h_M(X_t) G_t^{-1}$ are martingales. For z with $m_\lambda \leq z \leq M_\lambda$, let h be the function corresponding to (λ, z) . Because h can be expressed as a linear combination of h_m and h_M , $e^{\lambda t} h(X_t) G_t^{-1}$ is also a martingale. Thus, (λ, z) is in \mathcal{A} .

We now show that $e^{\lambda t} h(X_t) G_t^{-1}$ is not a martingale if at least one of (λ, m_λ) and (λ, M_λ) is not in \mathcal{A} . Write $h = c h_m + (1 - c) h_M$ for some constant $0 \leq c \leq 1$. Because $e^{\lambda t} h_m(X_t) G_t^{-1}$ and $e^{\lambda t} h_M(X_t) G_t^{-1}$ are positive local martingales, they are supermartingales. We have

$$\begin{aligned} h(\xi) &= c h_m(\xi) + (1 - c) h_M(\xi) \\ &\geq c \mathbb{E}[e^{\lambda t} h_m(X_t) G_t^{-1}] + (1 - c) \mathbb{E}[e^{\lambda t} h_M(X_t) G_t^{-1}] \\ &= \mathbb{E}[e^{\lambda t} h(X_t) G_t^{-1}]. \end{aligned}$$

If at least one of (λ, m_λ) and (λ, M_λ) is not in \mathcal{A} , the above inequality is strict. This completes the proof.

E Proof of Proposition 4.8

Theorem E.1. Let (β, ϕ) be a solution pair of $\mathcal{L}\phi = -\beta\phi$ with positive function ϕ . Then

$$e^{\beta t} \phi(X_t) \phi^{-1}(\xi) G_t^{-1}$$

is a martingale if and only if the diffusion induced by (β, ϕ)

$$dX_t = (k + \sigma^2 \phi' \phi^{-1})(X_t) dt + \sigma(X_t) dB_t$$

does not explode.

Proof. Set

$$H_t := \exp\left(-\frac{1}{2} \int_0^t v^2(X_s) ds - \int_0^t v(X_s) dW_s\right).$$

Define a new measure \mathbb{L} by setting the Radon-Nikodym derivative $d\mathbb{L} = H_t d\mathbb{Q}$. By the Girsanov theorem, we know that a process Z_t defined by

$$dZ_t = v(X_t) dt + dW_t$$

is a Brownian motion under \mathbb{L} . It follows that

$$\begin{aligned} dX_t &= (b - v\sigma)(X_t) dt + \sigma(X_t) dZ_t \\ &= k(X_t) dt + \sigma(X_t) dZ_t. \end{aligned}$$

We used that that $k(x) = (b - v\sigma)(x)$. We show that $e^{\beta t} \phi(X_t) G_t^{-1}$ is a martingale under \mathbb{Q} if and only if $e^{\beta t - \int_0^t r(X_s) ds} \phi(X_t)$ is a martingale under \mathbb{L} . It is because

$$\begin{aligned} \mathbb{E}^{\mathbb{L}}[e^{\beta t - \int_0^t r(X_s) ds} \phi(X_t)] &= \mathbb{E}^{\mathbb{Q}}[e^{\beta t - \int_0^t r(X_s) ds} \phi(X_t) H_t] \\ &= \mathbb{E}^{\mathbb{Q}}[e^{\beta t} \phi(X_t) G_t^{-1}]. \end{aligned}$$

Since both are supermartingales, this computation gives the desired result.

From the contents of [16] on page 212 and 215, we know that $e^{\beta t - \int_0^t r(X_s) ds} \phi(X_t)$ is a martingale under \mathbb{L} if and only if

$$dX_t = (k + \sigma^2 \phi' \phi^{-1})(X_t) dt + \sigma(X_t) dZ_t$$

does not explode. This completes the proof. \square

Lemma E.2. Assume $\delta < \lambda \leq \bar{\beta}$. Let g and h be the functions corresponding to tuple (δ, M_δ) and (λ, M_λ) , respectively. Then we have $g'g^{-1} > h'h^{-1}$.

Proof. For convenience, we may assume that $\xi = 0$, the left boundary is $-\infty$ and the right boundary is ∞ . Recall $q(x) = e^{-\int_0^x \frac{k(y)}{\sigma^2(y)} dy}$ in Proposition A.1. Write $h = uq$ and $g = vq$. Define $\Gamma = \frac{h'}{h} - \frac{g'}{g} = \frac{u'}{u} - \frac{v'}{v}$. Then

$$\Gamma' = -\Gamma^2 - \frac{2v'}{v}\Gamma - \frac{2(\lambda - \delta)}{\sigma^2}.$$

It suffices to show that $\Gamma(x) < 0$ for all x . First, we show this for $x > 0$. We know $\Gamma(0) = M_\lambda - M_\delta < 0$. We have that $\Gamma(x) < 0$ for all $x > 0$ because Γ ever gets close to 0, then the term $-\frac{2(\lambda - \delta)}{\sigma^2}$ dominates the right hand side of the equation above.

We now show that $\Gamma(x) < 0$ for all $x < 0$. Suppose there exists $x_1 < 0$ such that $\Gamma(x_1) \geq 0$. Then for all $z < x_1$, it is obtained that $\Gamma(z) > 0$ since if Γ ever gets close to 0, then term $-\frac{2(\beta - \lambda)}{\sigma^2}$ dominates the right hand side of the equation. Fix a point x_0 such that $x_0 < x_1$, thus $\Gamma(x_0) > 0$. For $z < x_0$, we have

$$-\frac{2v'(z)}{v(z)} = \frac{\Gamma'(z)}{\Gamma(z)} + \Gamma(z) + \frac{2(\beta - \lambda)}{\sigma^2(z)} \cdot \frac{1}{\Gamma(z)}.$$

By the same argument in Appendix B, we have

$$\int_{-\infty}^{x_0} \frac{1}{v^2(y)} dy < \infty.$$

This implies that the diffusion process induced by tuple (δ, M_δ) is attracted to the left boundary, which is a contradiction. \square

We now prove Proposition 4.8.

Proof. Assume $\delta < \lambda \leq \bar{\beta}$. Let g and h be the functions corresponding to tuple (δ, M_δ) and (λ, M_λ) , respectively. Suppose that $e^{\delta t} g(X_t) G_t^{-1}$ is a martingale. Then by Proposition 3.2 and Theorem E.1, we know that the process X_t under the transformed measure with respect to (δ, M_δ) satisfies

$$dX_t = (k + \sigma^2 g' g^{-1})(X_t) dt + \sigma(X_t) dB_t$$

and does not explode to ∞ . The above lemma says $g'g^{-1} > h'h^{-1}$. By the comparison theorem, we know that a process Y_t with

$$dY_t = (k + \sigma^2 h' h^{-1})(Y_t) dt + \sigma(Y_t) dB_t$$

satisfies $Y_t \leq X_t$ almost everywhere, thus Y_t does not explode to ∞ . On the other hand, it is clear that Y_t does not explode to $-\infty$ because it is non-attracted to the left boundary. By Theorem E.1, we conclude that $e^{\lambda t} h(X_t) G_t^{-1}$ is a martingale. \square

F An example for Corollary 4.6

Consider the equation $\mathcal{L}h = -\lambda h$. Recall that $\bar{\beta} := \max\{\lambda \mid (\lambda, z) \in \mathcal{A}\}$. That is, $\bar{\beta}$ is the maximum value among all the λ 's of the solution pair (λ, h) with $h(\xi) = 1$ and $h(\cdot) > 0$. In this section, we explore an example such that $\mathcal{L}h = -\bar{\beta}h$ has two linearly independent positive solutions.

Let \mathcal{L} be such that

$$\mathcal{L}h(x) := h''(x) + \frac{x}{(1+x^2)^{3/4}} h'(x) \quad \text{for } x \in \mathbb{R}.$$

First, $\mathcal{L}h(x) = 0$ has two linearly independent positive solutions: $h_1(x) \equiv 1$ and $h_2(x) = \int_{-\infty}^x e^{-\int_0^y u(z) dz} dy$ where $u(z) = \frac{z}{(1+z^2)^{3/4}}$.

It is enough to show that $\bar{\beta} = 0$. That is, for any fixed $\lambda > 0$, the equation $\mathcal{L}h = -\lambda h$ has no positive solutions. Suppose there exists such a positive solution h . Define a sequence of functions by $g_n(x) := \frac{h(x+n)}{h(n)}$ for $n \in \mathbb{N}$. By direct calculation, g_n satisfies the following equation:

$$g_n''(x) + \frac{x+n}{(1+(x+n)^2)^{3/4}} g_n'(x) = -\lambda g_n(x). \quad (\text{F.1})$$

By the Harnack inequality stated below, we have that $(g_n)_{n=1}^\infty$ is equicontinuous on each compact set on \mathbb{R} ; thus we can obtain a subsequence $(g_{n_k})_{k=1}^\infty$ such that the subsequence converges on \mathbb{R} , say the limit function g . Since g_n is positive, the limit function g is nonnegative and g is a nonzero function because $g(0) = \lim_{k \rightarrow \infty} g_{n_k}(0) = 1$. On the other hand, it can be easily shown that the limit function g satisfies $g''(x) = -\lambda g(x)$ by taking limit $n \rightarrow \infty$ in equation F.1. Clearly there does not exist a nonzero nonnegative solution of this equation when $\lambda > 0$. This is a contradiction. The author appreciates Srinivasa Varadhan for this example.

Theorem F.1. (Harnack inequality)

Let $h : \mathbb{R} \rightarrow \mathbb{R}$ be a positive solution of

$$a(x)h''(x) + b(x)h'(x) + c(x)h(x) = 0.$$

Assume that $a(x)$ is bounded away from zero; that is, there is a positive number l such that $a(x) \geq l > 0$. Suppose that $a(x), |b(x)|$ and $|c(x)|$ are bounded by a constant K . Then for any $z > 0$, there exists a positive number $M = M(z, K)$ (depending on z and K , but on neither $a(\cdot), b(\cdot), c(\cdot)$ nor $h(\cdot)$) such that $\frac{h(x)}{h(y)} \leq M$ whenever $|x - y| \leq z$.

G Reference Functions

In this section, we focus on the function ϕ rather than the value β . We assume that we roughly know the behavior of ϕ ; for example, we know a function f such that $\phi^{-1}f$ is bounded below and above or such that $\mathbb{E}_\xi^\mathbb{P}[(\phi^{-1}f)(X_t)]$ converges to a nonzero constant. Knowing f means that we have information about ϕ near the area where the process X_t lies with high probability under the objective measure. Such a function f is called a *reference function* of ϕ . More generally and more formally, we define a reference function in the following way.

Definition 6. A positive function f is called a *reference function* of ϕ if

$$\lim_{t \rightarrow \infty} \frac{1}{t} \cdot \log \mathbb{E}_\xi^\mathbb{P}[(\phi^{-1}f)(X_t)] = 0.$$

Proposition G.1. Knowing a reference function is equivalent to knowing the value β .

Proof. Suppose we know a reference function f of ϕ . From

$$\mathbb{E}_\xi^{\mathbb{Q}}[G_t^{-1}f(X_t)] = \mathbb{E}_\xi^{\mathbb{P}}[(\phi^{-1}f)(X_t)] \phi(\xi) e^{-\beta t},$$

and by the definition of the reference function, we have that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \cdot \log \mathbb{E}_\xi^{\mathbb{Q}}[G_t^{-1}f(X_t)] = -\beta.$$

Hence, we know the value β . Conversely, suppose we know the value β . We show that for any admissible pair (β, f) of $\mathcal{L}f = -\beta f$, f is a reference function. We have

$$\mathbb{E}_\xi^{\mathbb{Q}}[G_t^{-1}f(X_t)] = e^{-\beta t} f(\xi)$$

and thus

$$\mathbb{E}_\xi^{\mathbb{P}}[(\phi^{-1}f)(X_t)] = \mathbb{E}_\xi^{\mathbb{Q}}[G_t^{-1}f(X_t)] e^{\beta t} \phi^{-1}(\xi) = (\phi^{-1}f)(\xi).$$

This completes the proof. □

References

- [1] Audrino, F., R. Huitema and M. Ludwig. *An Empirical Analysis of the Ross Recovery Theorem*. (2014) Working paper.
- [2] Borovicka, J., L.P. Hansen and J.A. Scheinkman. *Misspecified Recovery*. (2014) Working Paper.
- [3] Borovicka, J., L.P. Hansen, M. Hendricks and J.A. Scheinkman. *Risk Price Dynamics*. (2011) Journal of Financial Econometrics, vol 9, 3-65.
- [4] Breeden, D.T. *An Intertemporal Asset Pricing Model with Stochastic Consumption and Investment Opportunities* (1979) Journal of Financial Economics, vol 7(3), 265-296.
- [5] Carr, P. and J. Yu. *Risk, Return, and Ross Recovery*. (2012) The Journal of Derivatives, vol. 20, no. 1, 38-59.
- [6] Davydov, D. and V. Linetsky. *Pricing Options on Scalar Diffusions: An Eigenfunction Expansion Approach*. (2003) Operations Research, 185-290.
- [7] Dubynskiy, S and R. Goldstein *Recovering Drifts and Preference Parameters from Financial Derivatives* (2013)
- [8] Durrett, R. *Stochastic Calculus: A Practical Introduction*. (1996) CRC Press.
- [9] Goodman, J. and H. Park *Do Prices Determine Objective Measures?* (2014) Working Paper.
- [10] Hansen, L.P. and J.A. Scheinkman. *Long Term Risk: An Operator Approach*. (2009) Econometrica, vol 77, 177-234.
- [11] Hull, J.C. *Options, Futures, and Other Derivatives*. (1997) Pearson Prentice Hall.

- [12] Karatzas, I. and S.E. Shreve. *Brownian Motion and Stochastic Calculus*. (1991) Springer-Verlag. New York.
- [13] Karatzas, I. and S.E. Shreve. *Methods of Mathematical Finance*. (1998) Springer-Verlag. New York.
- [14] Karlin, S. and H. Taylor. *A Second Course in Stochastic Process*. (1981) Academic Press.
- [15] Martin, I. and S. Ross. *The Long Bond*. (2013) Working paper.
- [16] Pinsky, R.G. *Positive Harmonic Function and Diffusion*. (1995) Cambridge University Press. New York
- [17] Qin, L. and V. Linetsky. *Long Term Risk: A Martingale Approach*. (2014a) Working paper.
- [18] Qin, L. and V. Linetsky. *Positive Eigenfunctions of Markovian Pricing Operators: Hansen-Scheinkman Factorization and Ross Recovery*. (2014b) Working paper.
- [19] Romer, D. *Advanced Macroeconomics*. (2012) McGraw-Hill.
- [20] Ross, S.A. *The Recovery Theorem*. (2015) Journal of Finance, vol 70, no 2, 615–648.
- [21] Walden, J. *Recovery with Unbounded Diffusion Processes*. (2013) Working paper.
- [22] Zettl, A. *Sturm-Liouville Theory*. (2005) American Mathematical Society, vol 121.