# PRISM: Recovery of the primordial spectrum from Planck data

F. Lanusse<sup>1</sup><sup>\*</sup>, P. Paykari<sup>1</sup><sup>\*\*</sup>, J.-L. Starck<sup>1</sup>, F. Sureau<sup>1</sup>, and J. Bobin<sup>1</sup>

<sup>1</sup> Laboratoire AIM, UMR CEA-CNRS-Paris 7, Irfu, SAp, CEA Saclay, F-91191 Gif sur Yvette cedex, France.

June 19, 2022

## ABSTRACT

## Context.

*Aims.* The primordial power spectrum describes the initial perturbations that seeded the large-scale structure we observe today. It provides an indirect probe of inflation or other structure-formation mechanisms. In this letter, using our recently published **PRISM** algorithm, we recover the primordial power spectrum from Planck PR1 dataset.

*Methods.* PRISM is a sparsity-based inversion method, which aims at recovering features in the primordial power spectrum from the empirical power spectrum of the Cosmic Microwave Background (CMB). The ill-posed inverse problem involved is regularised using a sparsity prior on features in the primordial power spectrum in a wavelet dictionary. Although this non-parametric method does not assume a strong prior on the shape of the primordial power spectrum, it is able to recover both its global shape and localised features. As a results, this approach presents a robust way of detecting deviations from the currently favoured scale-invariant spectrum.

*Results.* We apply PRISM to 100 Planck simulated data to investigate its performance on Planck-like data. We then apply PRISM to the Planck PR1 power spectrum to recover the primordial power spectrum. We further test on simulations the algorithm's ability to recover a small localised feature at  $k \sim 0.125$  Mpc<sup>-1</sup>, causing a large dip at  $\ell \sim 1800$  in the angular power spectrum.

Conclusions. We find no significant departures from the fiducial Planck PR1 near scale-invariant primordial power spectrum with the spectral index  $A_s = 2.215 \times 10^{-9}$  and  $n_s = 0.9624$ .

Key words. Cosmology : Primordial Power Spectrum, Methods : Data Analysis, Methods : Statistical

## 1. Introduction

The primordial power spectrum describes the initial perturbations curvature perturbations which over time evolved to form the large scale structure we observe today. Because the physics of the early Universe are encoded in the primordial power spectrum, it represents an invaluable probe into primordial cosmology and its measurement is a crucial research area in modern cosmology. The currently favoured model describing the physics of the early Universe, inflation (Guth 1981; Linde 1982), produces initial perturbations from quantum fluctuations during an epoch of accelerated exponential expansion. This inflation process produces a power spectrum of specific shape and can leave characteristic features. For the simplest models of inflation, the power spectrum, generated by almost purely adiabatic perturbations, is predicted to be nearly scale invariant. Hence, it is often expressed in terms of an amplitude  $A_s$ , a spectral index  $n_s$  with an optional 'running'  $\alpha_s$ :

$$P(k) = A_s \left(\frac{k}{k_p}\right)^{n_s - 1 + \frac{1}{2}\alpha_s \ln(k/k_p)} , \qquad (1)$$

where  $k_p$  is a pivot scale. Exact scale invariance, known as the Harrison-Zeldovich (HZ) model which sets  $n_s = 1$  (and  $\alpha_s = 0$ ) (Harrison 1970; Zeldovich 1972), has been ruled out by different datasets. Instead, the near scale-invariant spectrum with  $n_s < 1$  fits the current observations very well (e.g., Planck Collaboration et al. 2013a). There exists more complex models generating deviations from scale invariance. For a lightning review of inflation you can refer to Planck Collaboration et al. (2013b).

The recent Planck mission temperature anisotropy data, combined with the WMAP large-scale polarisation, constrain the spectral index to  $n_s = 0.9603 \pm 0.0073$  (Planck Collaboration et al. 2013a), ruling out exact scale invariance at over  $5\sigma$ . Also, Planck does not find a statistically significant running of the scalar spectral index, obtaining  $\alpha_s = -0.0134 \pm 0.0090$ . On the other hand, high-resolution CMB experiments, such as the South Pole Telescope (SPT)<sup>1</sup>, report a small running of the spectral index;  $-0.046 < \alpha_s < -0.003$  at 95% confidence (Hou et al. 2012). However, in general, any such detections have been small and consistent with zero.

In Planck Collaboration et al. (2013a) an extensive investigation is performed to search for features in the primordial power spectrum. A penalised likelihood approach suggests a feature near the highest wavenumbers probed by Planck at an estimated significance of  $\sim 3\sigma$ . This nominally statistically significant feature is detected around  $k \sim 0.13 \text{ Mpc}^{-1}$ . It has been confirmed that the large dip at  $\ell \sim 1800$  in the CMB power spectrum, which is associated with residual electromagnetic interference generated by the drive electronics of the 4 K cooler, is in fact responsible for the features detected at these high wavenumbers.

With the recent release of BICEP2 B-mode polarisation data there has been a range of papers investigating

<sup>\*</sup> francois.lanusse@cea.fr

<sup>\*\*</sup> paniez.paykari@cea.fr

<sup>&</sup>lt;sup>1</sup> http://pole.uchicago.edu/spt/index.php

the shape of both the scalar and the tensor primordial power spectrum in light of this new data. Authors of Hazra et al. (2014) have combined Planck CMB temperature and BICEP2 B-mode polarisation data (BICEP2 Collaboration et al. 2014a,b) to show that, assuming inflationary consistency relation, the power-law form of the scalar primordial spectrum is ruled out at more than  $3\sigma$  confidence level. In fact, a break or step at large scales in the primordial scalar perturbation spectrum is more favourable.

Determining the shape of the primordial spectrum generally consists of two approaches, one by parametrisation and the second by reconstruction. Non-parametric methods suffer from the non-invertibility of the transfer function that describes the *transfer* from P(k) to the CMB power spectrum. The dependence on the transfer function has the form

$$C_{\ell}^{\rm th} = 4\pi \int_0^\infty d\ln k \Delta_{\ell}^2(k) P(k) , \qquad (2)$$

where  $\ell$  is the angular wavenumber that corresponds to an angular scale via  $\ell \sim 180^{\circ}/\theta$  and  $\Delta_{\ell}(k)$  is the radiation transfer function holding the cosmological parameters responsible for the evolution of the Universe. Due to both the singularity of the transfer function and the limitations on the data due to effects such as projection, cosmic variance, instrumental noise, point sources, and etc., a robust algorithm is necessary for an accurate reconstruction of the primordial power spectrum from CMB data.

In this letter, we use our recently published PRISM algorithm (Paykari et al. 2014), to reconstruct this spectrum from the LGMCA Planck PR1 data (Bobin et al. 2014).

# 2. PRISM Algorithm

A CMB experiment, such as Planck, measures the anisotropies in the CMB temperature  $\Theta(p)$  in direction p, which is described as  $T(p) = T_{\text{CMB}}[1 + \Theta(p)]$ . This anisotropy field can be expanded in terms of spherical harmonic functions  $Y_{\ell m}$  as  $\Theta(\boldsymbol{p}) = \sum_{\ell} \sum_{m} a_{\ell m} Y_{\ell m}(\boldsymbol{p})$ , where  $a_{\ell m}$  are the spherical harmonic coefficients, that have a Gaussian distribution with zero mean,  $\langle a_{\ell m} \rangle = 0$ , and variance  $\langle a_{\ell m} a^*_{\ell' m'} \rangle = \delta_{\ell \ell'} \delta_{m m'} C^{\text{th}}_{\ell}$ . In practice, we only observe a realisation of this underlying power spectrum on our sky, meaning we are restricted by cosmic variance, especially on large scales. In addition, data is contaminated with additive instrumental noise on small scales and, due to different Galactic foregrounds, some areas of the observed CMB map need to be masked which induces correlations between modes. Taking these effects into account and following the MASTER method from Hivon et al. (2002), the pseudo power spectrum  $\widetilde{C}_{\ell}$  and the empirical power spectrum  $\widehat{C}_{\ell}^{\text{th}}$ , which is defined as  $\widehat{C}_{\ell}^{\text{th}} = 1/(2\ell+1)\sum_{m} |a_{\ell m}|^2$ , can be related through their ensemble averages

$$\langle \widetilde{C}_{\ell} \rangle = \sum_{\ell'} M_{\ell\ell'} \langle \widehat{C}_{\ell'}^{\text{th}} \rangle + \langle \widetilde{N}_{\ell} \rangle , \qquad (3)$$

where  $M_{\ell\ell'}$  describes the mode-mode coupling between modes  $\ell$  and  $\ell'$  resulting from computing the transform on the masked sky. We note that in this expression  $\langle \widehat{C}_{\ell'}^{\text{th}} \rangle = C_{\ell'}^{\text{th}}$ , and we set  $C_{\ell} = \langle \widetilde{C}_{\ell} \rangle$  and  $N_{\ell} = \langle \widetilde{N}_{\ell} \rangle$ , where  $C_{\ell}$  and  $N_{\ell}$ refer to the CMB and the noise power spectra of the masked maps, respectively. We have assumed that the pseudo power spectrum  $\widetilde{C}_{\ell}$  follows a  $\chi^2$  distribution with  $2\ell + 1$  degrees of freedom and hence can be modelled as

$$\widetilde{C}_{\ell} = \left(\sum_{\ell'} M_{\ell\ell'} C_{\ell'}^{\text{th}} + N_{\ell}\right) Z_{\ell} , \qquad (4)$$

where  $Z_{\ell}$  is a random variable distributed according to  $(2\ell + 1)Z_{\ell} \sim \chi^2_{2\ell+1}$ .

# 2.1. Formulation of the inverse problem

The relation between the discretised primordial power spectrum  $P_k$  and the measured pseudo power spectrum  $\tilde{C}_{\ell}$ , computed on a masked noisy map of the sky, can be condensed in the following form

$$\widetilde{C}_{\ell} = \left(\sum_{\ell'k} M_{\ell\ell'} T_{\ell'k} P_k + N_{\ell}\right) Z_{\ell} , \qquad (5)$$

with matrix elements  $T_{\ell k} = 4\pi\Delta \ln k \Delta_{\ell k}^2$ , where  $\Delta \ln k$  is the logarithmic k interval for the discrete sampling chosen in the integration of the system of equations. Because of the non-invertibility of the **T** operator, recovering the primordial power spectrum  $P_k$  from the true CMB power spectrum  $C_{\ell}^{\text{th}}$  already constitutes an ill-posed inverse problem, made even more difficult by the mask and cosmic variance affecting the observed  $\tilde{C}_{\ell}$ . In PRISM, we address both the inversion problem and the control of the noise on the CMB spectrum due to sample variance in the framework of sparse recovery. The inversion problem in Equation 5 can be regularised in a robust way by using the sparse nature of the reconstructed signal as a prior.

#### 2.2. Pk sparse recovery formulation

If the signal to recover,  $P_k$  in our case, can be sparsely represented in an adapted dictionary  $\Phi$ , then this problem, known as the basis pursuit denoising BPDN, can be recast as an optimisation problem. In our case, the optimisation problem can be formulated as

$$\min_{X} \frac{1}{2} \parallel C_{\ell} - (\mathbf{MT}X + N_{\ell}) \parallel_{2}^{2} + \lambda \parallel \mathbf{\Phi}^{t}X \parallel_{0}, \quad (6)$$

where X is the reconstructed estimate for the primordial power spectrum  $P_k$ . The first term in Eq. (6) imposes a  $\ell_2$ fidelity constraint to the data while the second term promotes the sparsity of the solution in dictionary  $\Phi$ . The parameter  $\lambda$  tunes the sparsity constraint. Although the  $\ell_0$  optimisation problem stated in Equation (6) cannot be solved directly, its solution can be estimated by solving a sequence of weighted  $\ell_1$  minimisation problems (Candes et al. 2008) of the form:

$$\min_{X} \frac{1}{2} \parallel \frac{1}{\sigma_{\ell}} \overline{R}_{\ell}(X) \parallel_{2}^{2} + K \sum_{i} \lambda_{i} |[\mathbf{W} \boldsymbol{\Phi}^{t} X]_{i}|, \qquad (7)$$

where **W** is a diagonal matrix applying a different weight for each wavelet coefficient,  $\overline{R}_{\ell}(X)$  is an estimate of the residual  $C_{\ell} - (\mathbf{MT}X + N_{\ell})$  and  $\sigma_{\ell}$  is its standard deviation. The parameters  $\lambda_i$  are set with respect to the expected standard variation  $\sigma(w_i)$  of each wavelet coefficient such that  $\lambda_i = K\sigma(w_i)$  where K is a global regularisation parameter (usually set to K = 3 or K = 4) which translates into a significance level threshold for the detection of features. The weighted  $\ell_1$  problem (7) is solved several times using the popular iterative soft-thresholding algorithm (ISTA) and updating each time the weights **W** based on the solution of the previous iteration. This procedure is fully described in Paykari et al. (2014).

# 3. Results

Before applying the PRISM algorithm to the Planck data processed with the  $LGMCA^2$  pipeline, we investigate the performance of the algorithm on a set of simulations. The Planck team provides 100 of their 1000 CMB and instrumental noise simulated maps<sup>3</sup>. The cosmology used to set up these CMB simulations is different from the PR1 best-fit cosmology. As we want to have the exact same pipeline applied to both the data and the simulations (hence the same radiation transfer function), we create our own 100 CMB simulations based on PR1 fiducial cosmology. However, we use the simulated noise maps provided by the Planck team. In order to have a thorough comparison between the simulations and the Planck PR1 data, we run the simulations through the same pipeline as the Planck data, i.e. the LGMCA pipeline. For each simulation, we process the nine frequency channels through LGMCA, with the precomputed set of parameters (Bobin et al. 2014). Full sky 5 arcmin resolution noisy maps are obtained, which are then masked using a Galactic and point sources mask with  $f_{\rm sky} = 0.76$ . The pseudo power spectra are obtained by applying the empirical power spectrum estimator to the masked maps. We also build an estimate of the instrumental noise power spectrum  $N_{\ell}$  by processing the noise maps through the same pipeline in LGMCA, mask the output noise maps and apply the pseudo spectrum estimator to these masked maps. We set our estimate of  $N_{\ell}$  to the average of the 100 noise pseudo spectra.

To apply PRISM to the simulated data, we build a transfer function  $\mathbf{T}'$  adapted to the simulations, so that it includes the effects of the beam  $b_\ell$  and the HEALPix window  $h_{\ell}$  represented by two diagonal matrices  $\mathbf{B} = \text{diag}(b_{\ell}^2)$ and  $\mathbf{H} = \operatorname{diag}(h_{\ell}^2); \mathbf{T}' = \mathbf{HBT}$ . The radiation transfer function  $\mathbf{T}$  is computed for the PR1 fiducial cosmology using CAMB (http://camb.info). The lensing contribution to the CMB temperature power spectrum, also computed with CAMB for the PR1 fiducial cosmology, is taken into account as an additional contribution to  $N_{\ell}$ . Contrary to Paykari et al. (2014) where the algorithm was initialised to a scale invariant power spectrum, in this work we initialise the PRISM algorithm to the best fit Planck PR1 primordial power spectrum as we are looking for small deviations from the best fit power law which already fits the data. With this choice of initialisation, in the absence of evidence from the data, the reconstruction will not depart from the best fit power law. For the reconstruction of the simulations and the Planck data, we fix the regularisation parameter K to K = 4 which robustly removes the noise due to sampling variance and we use bi-orthogonal Battle-Lemarié wavelets of order 1 with 9 dyadic scales. This choice of wavelet dictionary is rather generic and not specifically tuned to recover physically motivated features but these



Fig. 1: Reconstructions for the primordial power spectra from 100 simulations and Planck PR1 data. The  $1\sigma$  and  $2\sigma$  dispersion of the reconstructed spectra from the simulations are shown in blue bands around the mean of the reconstructions (blue line). We note that these bands do not include the errors due to point sources and beam uncertainties. The Planck fiducial power spectrum used for the simulations, with  $A_s = 2.215 \times 10^{-9}$  and  $n_s = 0.9626$  is shown in yellow dashed line. The inset shows a close up of the main figure.

wavelets exhibit limited oscillations and have 2 vanishing moments which makes them well suited to recover a near scale invariant power spectrum in logarithmic scale.

We note that the reconstruction of the primordial power spectrum is limited by different effects on different scales. On very large scales, the recovery is limited by cosmic variance and geometrical projection of the modes and on small scales we are limited by effects such as instrumental noise and point sources. These limitations leave us with a window through which the primordial power spectrum can be constrained with a good accuracy and the detection of features is possible. For the Planck data, we expect this window to be in the range  $k \sim 0.005 - 0.20$  Mpc<sup>-1</sup>.

In Figure 1 we show the reconstructed spectra from the simulations and the data. The blue contours show the  $1\sigma$  and  $2\sigma$  dispersion of the 100 reconstructed spectra for the CMB simulations based on the PR1 fiducial cosmology. Note that theses bands do not include the errors due to point sources and beams uncertainties, which are especially important in small scales. The mean reconstructed power spectrum shown, in blue, perfectly fits the input PR1 best fit power law, in dashed yellow, in the entire reconstructed range. Of course it does not mean that the algorithm is able to perfectly reconstruct an unknown power spectrum over this entire range but that with the regularisation level used for these reconstructions, no significant departures from the best fit power law have been detected.

We find that the reconstructed spectrum from the LGMCA PR1 power spectrum, shown in red on Figure 1, remains within the  $1\sigma$  bar of the reconstructed spectra from the PR1 best fit power law. Thus, we find no significant departure from the PR1 best-fit near scale-invariant spectrum.

<sup>&</sup>lt;sup>2</sup> LGMCA codes and Planck PR1 data are available at http: //www.cosmostat.org/planck\_pr1.html

<sup>&</sup>lt;sup>3</sup> http://wiki.cosmos.esa.int/planckpla

As a complementary test of PRISM on Planck-like data we assessed the algorithm's ability to recover a small local departure from the best fit PR1 power law. We created a set of CMB simulations from a fiducial primordial power spectrum with a small localised test feature causing a dip in the angular power spectrum around  $\ell \sim 1800$ . The aim of this set of simulations is to mimic the feature suggested in Planck Collaboration et al. (2013a) to account for the large dip in the angular power spectrum, which was later confirmed as being caused by residual electromagnetic interferences. Our test primordial power spectrum is built from the best fit PR1 power law with an added feature around k = 0.125, shown in red on the top panel of Figure 2. The presence of this feature causes a dip in the angular power spectrum as can be seen on the residuals  $\Delta C_{\ell}$  between the fiducial angular power spectrum and the PR1 best fit  $C_{\ell}$ , shown in red on the bottom panel of Figure 2.

From this test primordial power spectrum, we generate a set of 100 CMB simulations using the exact same procedure as previously mentioned and we apply PRISM to the measured angular pseudo-power spectra with the exact same parameters. The mean of the reconstructed power spectra is shown in yellow on the top panel of Figure 2 and the  $1\sigma$  and  $2\sigma$  dispersion of the reconstruction is shown as green contours. One can see that the feature is successfully detected and the reconstruction show little bias in position and amplitude. The bottom panel of Figure 2 shows in yellow the residual  $\Delta C_{\ell}$  between the fiducial angular power spectrum and the mean of the angular power spectrum derived from the PRISM reconstructed primordial power spectra. As can be seen, using the primordial power spectrum reconstructed with PRISM allows a much better fit to the data than a power law and the reconstructed angular power spectra fall inside the  $1\sigma$  region due to cosmic variance, shown as a blue region. If such a feature existed in the LGMCA processed Planck PR1 data, PRISM would therefore have been able to detect it.

# 4. Conclusions

The simplest models of inflation are currently the most favoured by the data and predict a near scale-invariant power spectrum with a possible small running. The CMB spectrum provides a possibility to measure the primordial spectrum. However, the singular nature of the radiation transfer function, the joint estimation of the cosmological parameters and the primordial power spectrum, along with the different sources of noise impede the full recovery of the primordial spectrum. Therefore, devising a robust technique able to detect deviations from scale invariance is important.

In this paper we have applied our recently published algorithm, PRISM Paykari et al. (2014), to the Planck PR1 data to recover the primordial power spectrum. PRISM is a sparse recovery method, making use of the sparsity of the primordial power spectrum as well as an adapted modelling for the noise of the CMB power spectrum. This algorithm assumes no prior shape for the primordial spectrum and does not require a coarse binning of the power spectrum, making it sensitive to both global smooth features (e.g. running of the spectral index) as well as local sharp features (e.g. a bump or an oscillatory feature). We have reconstructed the primordial power spectrum from the LGMCA PR1 Planck dataset in the range  $k \sim 0.005 - 0.20 \text{ Mpc}^{-1}$ .



Fig. 2: Top panel shows a fiducial primordial power spectrum with a feature around  $k = 0.125 \text{ Mpc}^{-1}$  in red and in green contours the  $1\sigma$  and  $2\sigma$  dispersion of 100 reconstructions from simulated CMB spectra. The mean of reconstructions is shown in yellow. Bottom panel shows the residuals  $\Delta C_{\ell}$  between the  $C_{\ell}$  for the fiducial primordial power spectrum with a feature and the  $C_{\ell}$  for the best fit Planck power law in red and for the mean reconstructed primordial power spectrum in yellow. The green bands indicate the  $1\sigma$  and  $2\sigma$  bands for the  $\Delta C_{\ell}$  from the simulations and the dashed blue lines show the  $1\sigma$  region due to cosmic variance.

We have not detected any significant deviations from a scale-invariant power spectrum, whether local or global such as a running of the spectral index. However, we have tested that PRISM would have been able to recover a small localised feature around  $k \sim 0.125$ , similar to the one suggested in Planck Collaboration et al. (2013a) to account for a dip in the angular power spectrum at around  $\ell \sim 1800$ , if it were present on the LGMCA Planck PR1 data.

## Acknowledgments

This work is supported by the European Research Council grant SparseAstro (ERC-228261)

## References

- BICEP2 Collaboration, Ade, P. A. R., Aikin, R. W., et al. 2014a, ArXiv e-prints
- BICEP2 Collaboration, Ade, P. A. R., Aikin, R. W., et al. 2014b, ArXiv e-prints
- Bobin, J., Sureau, F., Starck, J.-L., Rassat, A., & Paykari, P. 2014, A&A, 563, A105
- Candes, E. J., Wakin, M. B., & Boyd, S. P. 2008, Journal of Fourier Analysis and Applications, 14(5):877
- Guth, A. H. 1981, Phys. Rev. D, 23, 347
- Harrison, E. R. 1970, Phys. Rev. D, 1, 2726
- Hazra, D. K., Shafieloo, A., Smoot, G. F., & Starobinsky, A. A. 2014, ArXiv e-prints
- Hivon, E., Górski, K. M., Netterfield, C. B., et al. 2002, ApJ, 567, 2 Hou, Z., Reichardt, C. L., Story, K. T., et al. 2012, ArXiv e-prints
- Linde, A. D. 1982, Physics Letters B, 108, 389
- Paykari, P., Lanusse, F., Starck, J.-L., Sureau, F., & Bobin, J. 2014, A&A, 566, A77
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2013a, ArXiv e-prints
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2013b, ArXiv e-prints
- Zeldovich, Y. B. 1972, MNRAS, 160, 1P